Binding and sequestration of poison frog alkaloids by a plasma globulin

Aurora Alvarez-Buylla[1]*, Maria Dolores Moya-Garzon[2,3], Alexandra E. Rangel[4], Elicio E. Tapia[5], Julia
Tanzo[2,3] , H. Tom Soh[4,6,7], Luis A. Coloma[5], Jonathan Z. Long[2,3,8,9], Lauren A. O'Connell[1,3,10]*

[1]Department of Biology, Stanford University, Stanford, CA, USA
[2]Department of Pathology, Stanford University, Stanford, CA, USA
[3]Sarafan ChEM-H, Stanford University, Stanford, CA, USA
[4]Department of Radiology, Stanford University, Stanford, CA, USA
[5]Centro Jambatu de Investigación y Conservación de Anfibios, Fundación Jambatu, San Rafael, Quito,
Ecuador
[6]Department of Electrical Engineering, Stanford University, Stanford, CA, USA
[7]Chan Zuckerberg Biohub, San Francisco, CA, USA.
[8]Stanford Diabetes Research Center, Stanford University, Stanford, CA, USA
[9]Wu Tsai Human Performance Alliance, Stanford University, Stanford, CA, USA
[10]Wu Tsai Institute for Neuroscience, Stanford University, Stanford, CA, USA

*corresponding authors: auroraab@stanford.edu, loconnel@stanford.edu

Key words: plasma globulin, alkaloid binding, serpin, protein evolution
Running title: Alkaloid-Binding Globulin (ABG)

ABSTRACT

Alkaloids are important bioactive molecules throughout the natural world, and in many animals they serve
as a source of chemical defense against predation. Dendrobatid poison frogs bioaccumulate alkaloids
from their diet to make themselves toxic or unpalatable to predators. Despite the proposed roles of
plasma proteins as mediators of alkaloid trafficking and bioavailability, the responsible proteins have not
been identified. We use chemical approaches to show that a ~50 kDa plasma protein is the principal
alkaloid binding molecule in blood from poison frogs. Proteomic and biochemical studies establish this
plasma protein to be liver-derived alkaloid-binding globulin (ABG) that is a member of the serine-protease
inhibitor (serpin) family. In addition to alkaloid binding activity, ABG sequesters and regulates the
bioavailability of "free" plasma alkaloids *in vitro*. Unexpectedly, ABG is not related to saxiphilin, albumin,
or other known vitamin carriers, but instead exhibits sequence and structural homology to mammalian
hormone carriers and amphibian biliverdin binding proteins. Alkaloid-binding globulin (ABG) represents
a new small molecule binding functionality in serpin proteins, a novel mechanism of plasma alkaloid
transport in poison frogs, and more broadly points towards serpins acting as tunable scaffolds for small
molecule binding and transport across different organisms.

33
34

1. INTRODUCTION

Alkaloids are nitrogenous small molecules that play important ecological and physiological roles throughout nature, one of which is mediating predator-prey interactions. Species across many taxa, including plants, insects, marine invertebrates, and vertebrates, have co opted alkaloids as chemical defenses, methods for hunting, and pheromone signals. Some of the most potent alkaloid toxins, including batrachotoxin (BTX), saxitoxin (STX), and tetrodotoxin (TTX), act specifically by affecting voltage gated sodium (NaV), potassium, and calcium channels, leading to disruption of nerve and muscle cells [1–3]. In the blue-ringed octopus, *Hapalochlaena lunulata*, TTX is used to paralyze prey [4], while in the pufferfish *Takifugu niphobles* it also acts as a pheromone [5], and in the California newt *Taricha torosa* it is a defense against predation [6]. Other less-potent alkaloids also play important roles in predator-prey interactions. For example, Lepidoptera insects (butterflies and moths) and Coleoptera beetles sequester pyrrolizidine alkaloids from plants for predation, defense, and production of pheromones [7,8]. Although the identities of these alkaloids are well documented, less is known about the physiological mechanisms that allow animals to produce, sequester, and autoresist these small molecules. Identifying and characterizing proteins that interact with alkaloids in these ecological contexts allows us to better understand how animal physiology has coevolved with alkaloids.

Despite the important ecological and physiological roles of alkaloids in animals, the molecular mechanisms involved in alkaloid production, transport, and resistance have been more elusive and typically focused on a single alkaloid or specific structural class of alkaloids. In grasshoppers and moths, passive absorption of pyrrolizidine alkaloids is followed by conversion into a non-toxic form by haemolymph flavin-dependent monooxygenase, allowing the insects to avoid autointoxication [9]. In some beetle species, ATP-binding cassette (ABC) transporters actively pump pyrrolizidine alkaloids into reservoir defensive glands [10]. In vertebrates, the proteins that allow for the sequestration of alkaloids without autotoxicity are unclear with the exception of previous work with TTX and STX. The pufferfish saxitoxin and tetrodotoxin binding protein (PSTBP) was originally identified in the plasma of *Fugu pardalis* [11], and is thought to play a role in the transport of TTX and STX to the site of bioaccumulation in the liver and ovaries in many pufferfish species [12]. The soluble protein saxiphilin has been proposed as a toxin sponge for STX in various species of amphibians [13], although it remains unclear whether these species come into contact with STX in nature and whether saxiphilin may act as the predominant STX transporter in the plasma of frogs. While these insights have advanced our understanding of toxin physiology, studies in vertebrates have been narrowly focused on a few potent alkaloids with high-specificity binding proteins. However, other animals carry a wide diversity of alkaloids for chemical defense and the physiological mechanisms that allow accumulation of structurally diverse alkaloids are unknown.

Some species of frogs sequester a remarkable diversity of dietary alkaloids onto their skin as a chemical defense. This trait has independently evolved in several frog families, including Dendrobatidae in Central and South American and Mantellidae in Madagascar. Over 500 compounds have been found on the skin of *Dendrobatidae* frogs, with some alkaloids sourced from ants, mites, millipedes and beetles [14–16]. Within dendrobatids, alkaloid-based chemical defenses have evolved independently at least three times [17,18], where non-toxic species do not uptake alkaloids onto their skin even when they are present in their diet [19–22]. Well-studied poison frog alkaloids include pumiliotoxins (PTX), whose documented targets include sodium and potassium gated ion channels [23,24], and decahydroquinolines (DHQ), which affect nicotinic acetylcholine receptors [25]. Epibatidine was first identified in the genus *Epipedobates* and specifically binds certain nicotinic receptors, leading it to be proposed as a therapeutic

analgesic alternative to morphine [26]. Although there is limited research into the mechanisms of
sequestration and autoresistance of alkaloids in poison frogs [13,27–30], it is likely this process involves
alkaloid transport through circulation for these dietary compounds to end up in skin storage glands. Based
on the extensive work on plasma small molecule transport in mammals, one might expect that proteins
like albumin, which is an abundant and promiscuous small molecule binder in the blood [31–33], or
vitamin transporters [34–36], which are able to interact with diet derived molecules, might be involved in
alkaloid sequestration in poison frogs. In this study, we tested the hypothesis that poison frogs have an
alkaloid-binding protein in the plasma and aimed to uncover its functional role and evolutionary
significance. We predicted this protein would bind a range of poison frog alkaloids and would be present
in frogs that are chemically defended in nature, but not in undefended species.
89
2. RESULTS
An alkaloid-like photoprobe identifies a binding protein in poison frog plasma
We used a biochemical strategy to directly test for alkaloid binding in the plasma of different
species of poison frogs. To do this, we obtained a UV crosslinking probe with an indolizidine functional
group that shares structural similarity to the poison frog alkaloid pumiliotoxin **251D** (PTX) (Figure 1A,
functional group highlighted in blue). Upon UV irradiation, the diazirine (green) enables protein
crosslinking and the subsequent probe-protein complex can be conjugated to a fluorophore for gel-based
visual analysis or biotin for streptavidin enrichment. Application of this photocrosslinking approach
outside of mammalian systems has been remarkably limited, and in frogs has been limited to studying
neuromuscular receptors [37,38]. We found the PTX-like photoprobe shows binding activity within the
plasma of three species of dendrobatid poison frogs, *Oophaga sylvatica*, *Dendrobates tinctorius*, and
Epipedobates tricolor (Figure 1B). In these species, this binding activity was largely restricted to a few
bands in the 50-60 kDa range, and was a similar size across different species. Plasma from a non-toxic
dendrobatid poison frog (*Allobates femoralis)*, a mantellid poison frog (*Mantella aurantiaca)*, the cane
toad (*Rhinella marina)*, and humans showed no binding activity with the photoprobe (Figure 1B). We
further tested whether the presence of alkaloids would compete off photoprobe binding. In *O. sylvatica*,
photoprobe binding resulted in two bands that showed competition by the addition of PTX,
decahydroquinoline (DHQ), or epibatidine (epi), but not with nicotine (Figure 1C). In *D. tinctorius*, the
photoprobe showed a two-band binding pattern, where the bottom band was competed by PTX and there
was slight competition of both bands with DHQ and epibatidine, but no competition with nicotine (Figure
1D). In both *O. sylvatica* and *D. tinctorius*, competition occurred when PTX was 10-fold higher in
concentration than the photoprobe (Figure S1A,C). In *E. tricolor* plasma from some individuals two bands
were observed while in others only one band was seen, and these were more faint in the presence of
DHQ or epibatidine but not PTX or nicotine (Figure 1E). We conclude from these photocrosslinking
experiments that plasma binding of alkaloids in three species of chemically defended dendrobatid poison
frogs is mediated by a ~50-60 kDa plasma protein.

116
Figure 1: Alkaloid-like photocrosslinking probes show binding and competition in poison frog plasma. (A) Structures of
alkaloid-like photocrosslinking probe and alkaloids tested, with the functional group in blue, the diazirine group in green, and the
terminal alkyne in yellow. **(B)** Plasma from different species (*Oophaga sylvatica, Dendrobates tinctorius, Epipedobates tricolor,*
Allobates femoralis, Rhinella marina, and humans, from left to right) show different plasma photoprobe binding activity and
competition. Orange lines on phylogeny indicate independent evolutionary origins of chemical defense in Dendrobatidae and
Mantellidae. **(C)** *Oophaga sylvatica* plasma shows crosslinking, and competition with pumiliotoxin (PTX), decahydroquinoline
(DHQ), and epibatidine (epi), but not nicotine (nic). **(D)** *Dendrobates tinctorius* plasma shows crosslinking, and competition with
PTX, but not DHQ, epi, or nic. **(E)** *Epipedobates tricolor* plasma shows crosslinking, and competition with DHQ and slightly with
epi, but not PTX or nic.

Proteomic analysis identifies an alkaloid-binding globulin

To identify the alkaloid-binding protein found in the plasma assays, we performed a pull-down
paired with gel-punch proteomics on three conditions: no photoprobe (negative control), photoprobe only
(positive control), and photoprobe with PTX competitor (Figure 2A). A biotin handle, instead of the
fluorophore used above, was chemically added to the photoprobe for the enrichment of proteins using
streptavidin beads. We used an untargeted proteomics approach to quantify and compare these enriched
fractions using a proteome reference created from the *O. sylvatica* genome. On average, 3876 unique
peptides were found per sample, mapping to 433 *O. sylvatica* proteins (Figure 2B). The most highly
abundant protein in the photoprobe condition had an average number of peptide spectral counts of
1224.5 and was competed off in the photoprobe with PTX condition by 64% (Figure 2B), resembling

background levels (Figure 2C). This protein is annotated as serine protease inhibitor A1 (serpin-A1),
which encodes for the protein alpha-1-antitrypsin (A1AT). As our subsequent experiments demonstrate
this protein functions as an alkaloid binding and sequestration protein, we refer to it hereafter as "alkaloid-
binding globulin" (ABG). Mapping the 72 unique peptides onto the protein sequence of ABG showed full
coverage across the protein, excluding the signal peptide, which in other serpins is cleaved during
secretion (Figure S1D). We conclude that ABG functions as a major alkaloid binding protein in poison
frog plasma.
144

Figure 2: Proteomics identifies serpinA1 as the main pumiliotoxin binding protein in *Oophaga sylvatica* plasma. (A)
Streptavidin blot of the proteins pulled down from *O. sylvatica* plasma across the three conditions: no photoprobe, photoprobe,
and photoprobe plus competitor pumiliotoxin (PTX). **(B)** Quantitative proteomics output in terms of percent competition defined
as 100% - average spectral counts in the photoprobe + ptx condition divided by average spectral counts in the photoprobe only
condition. Average was taken across two replicates. **(C)** The number of spectral counts across conditions for the serpinA1
protein, each replicate is shown individually.
152

Structural predictions of ABG show binding pocket similarities to mammalian hormone carriers
The identification of ABG as the principal alkaloid binding protein in plasma was unexpected as
plasma binding of small molecules is commonly mediated by albumin. Nevertheless, in mammals,
members of the serpinA family function as carriers of lipophilic hormones, providing plausibility to the
hypothesis that frog serpin family members may also bind small molecules. Therefore, we sought further
structural insights into ABG using protein structure predictions and molecular docking simulations to
examine if this protein has a predicted binding pocket for small molecules. Using AlphaFold to predict the
structure of the full protein sequence without the signal peptide resulted in a high confidence structure
(Figure 3A). We then compared it to the structures of serpinA6/corticosteroid-binding globulin (CBG,
Figure 3B) [39], biliverdin-binding serpin (BBS, Figure 3C) [40], serpinA1/alpha-1-antitrypsin (A1AT,
Figure S2A) [41], and serpinA7/thyroxine binding globulin (TBG, Figure S2B) [42]. The AlphaFold
prediction for *O. sylvatica* ABG (*Os*ABG) demonstrated a conserved structural element of three alpha
helices backed by a set of beta sheets, which is the small molecule binding pocket in CBG, BBS, and
TBG (black boxes, Figure 3A-C, Figure S2B), and also exists in the non-small molecule binding A1AT

(black box, Figure S2A). When a molecular docking simulation was run with the whole *Os*ABG protein
as the search space and PTX as the ligand, the highest affinity binding site was in the same binding
pocket defined by this structural motif (Figure 3D). Although the overall structural components of the
binding pockets show similarities across *Os*ABG, CBG, BBS, and TBG, the individual amino acids
coordinating the small molecule binding differ across proteins (Figure 3D-F, Figure S2D-E). These results
offer a structural explanation for PTX binding by ABG and highlight the homology between ABG and other
small molecule binding globulins.

174
Figure 3: Predicted ABG structure and binding pocket resembles that of other small molecule binding serpins. (A)
AlphaFold structure predicted with the protein sequence of the *Oophaga sylvatica* ABG, with color representing model
confidence and predicted binding pocket based on molecular docking simulation indicated with a black box. **(B)** Crystal structure
for rat SerpinA6/corticosteroid-binding globulin, with the cortisol molecule shown in orange (PDB# 2V95). **(C)** Crystal structure
for tree frog *Boana punctata* biliverdin-binding serpin (BBS), with biliverdin shown in orange (PDB# 7RBW). **(D)** Close-up of
predicted binding pocket of PTX in *O. sylvatica* ABG, with residues proximal to PTX highlighted in magenta. The structure of
PTX is indicated on the top right. **(E)** Close-up of cortisol binding in CBG (PDB# 2V95), with proximal residues highlighted in
magenta. Cortisol structure is displayed on the top right. **(F)** Close-up of biliverdin binding in BBS (PDB# 7RBW), with some
proximal residues highlighted in magenta. Biliverdin structure is shown on the top right.
184

Recombinant expression recapitulates binding activity of different ABG proteins

To confirm ABG binding activity *in vitro* and compare across different species, we recombinantly
expressed and purified *O. sylvatica* ABG (*Os*ABG), and its closest homolog from the *D. tinctorius* and *E.*
tricolor transcriptomes (*Dt*ABG and *Et*ABG, respectively). The resulting purified protein doublet (Figure
S4B) is likely due to differences in post-translational glycosylation, as *Os*ABG has two predicted N-
glycosylation sites [43]. As expected, purified *Os*ABG recapitulated the binding and competition seen
with the plasma, where the photoprobe was most fully competed by the presence of PTX, and also
competed off by DHQ and epibatidine, but not nicotine (Figure 4A). The competition activity with the
purified protein was noticeable at a ratio of one to one photoprobe to PTX (Figure S1B). Purified *Dt*ABG
and *Et*ABG required higher concentrations of protein to see a signal and showed much weaker
photoprobe binding, which was competed off by the presence of PTX and DHQ in the case of *Dt*ABG
(Figure 4B), and DHQ and epibatidine in the case of *Et*ABG (Figure 4C). Together these results confirm
the plasma findings that ABG is a multi-alkaloid binding protein with different specificities and affinities
across poison frog species, and that *Os*ABG alone is sufficient to recapitulate the crosslinking activity
observed in the plasma.
Given the predicted binding pocket from the molecular docking simulations, and the differences
in binding activity of the ABG proteins in different poison frog species, we used a sequence (Figure 4D)
and predicted structure (Figure 4E) informed approach to mutate residues that might coordinate alkaloid
binding in the hypothesized pocket. Using the molecular docking simulation, we identified 6 residues with
proximity to the docked PTX molecule that might have important binding activity: Y36, S268, D273, W276,
S374, D383 (Figure 4E). Mutating different sets of these binding residues in the *Os*ABG sequence led to
a disruption of binding and competition. The combined mutations of Y36A, W276A, S374A, D383A and
Y36A, S268A, D273A, D383A disrupted binding to the photoprobe completely (Figure 4E). The single
point mutation of D383A weakened photoprobe binding significantly, to the point of being nearly
undetectable (Figure 4E). These results demonstrate that mutating residues in the binding pocket
identified through molecular docking disrupts binding activity of *Os*ABG, providing biochemical evidence
that the structurally predicted binding pocket of ABG indeed is the relevant binding site for PTX.
Furthermore, we have identified a set of residues that are necessary for PTX binding with high affinity,
showing that the plasma binding activity is coordinated by specific amino acids in *Os*ABG.
213

215
Figure 4: Recombinant expression and binding pocket mutants confirm plasma binding activity and binding pocket
predictions. (A) Photoprobe crosslinking and competition with different compounds of 10 µg recombinantly expressed and
purified OsABG recapitulates the binding activity seen in the plasma (Figure 1C). **(B)** Photoprobe crosslinking with 35 µg
recombinantly expressed *Dendrobates tinctorius* ABG shows crosslinking, and competition with PTX and DHQ. **(C)** Photoprobe
crosslinking with 60 µg recombinantly expressed *Epipedobates tricolor* ABG shows crosslinking, and competition with DHQ and
epibatidine. **(D)** Alignment of protein sequence of proteins homologous to *Os*ABG across species shows conservation of certain
amino acids. Coloring of amino acids is based on the RasMol "amino" coloring scheme, which highlights amino acid properties.
(E) Potential binding residues were identified from the molecular docking simulation. Three different mutants were made based
on specific amino acids in the binding pocket, with either a combination of four different alanine substitutions (m1 - yellow and
teal residues, and m2 - yellow and green residues) or a single alanine substitution at D383 (m3). PTX is shown in magenta.
Oxygen atoms on the molecules are highlighted in red, nitrogen in blue. **(F)** Binding pocket mutants lose binding activity of the
photoprobe in the original reaction conditions.

228
OsABG sequesters free PTX in solution with high affinity

Previous work has described the binding affinities of small molecule binding serpins and their
important role in regulating the pool of free versus bound ligands in circulation [44–47]. We hypothesized
that *Os*ABG might play a similar role for alkaloids in the poison frog plasma. To test this, we examined

both the binding affinity of *Os*ABG for PTX and its ability to regulate the pool of bioavailable alkaloids in
solution. Using microscale thermophoresis (MST) we found a dissociation constant (K_D) of 1.58 µM
(Figure 5A). To test the ability of *Os*ABG to sequester alkaloids *in vitro*, we used a 3 kDa molecular weight
cutoff centrifuge filter to separate the "bound" and "free" PTX (Figure 5B), which we then quantified by
liquid chromatography-mass spectrometry. We found that in the presence of *Os*ABG, the amount of "free"
PTX is dramatically reduced, while that of nicotine is not (Figure 5C). These results show that *Os*ABG is
able to bind PTX in solution with high affinity, and therefore may regulate the amount of free PTX in
solution. This regulation of bioavailable pools of PTX in circulation may have downstream consequences
on sequestration, transcription, and signaling throughout the organism.

242
243 **Figure 5: OsABG sequesters free PTX in solution (A)** Microscale thermophoresis (MST) of labeled *Os*ABG with PTX finds a
244 dissociation constant (Kd) of 1.58 uM. **(B)** A 3 kDA molecular weight cut off (MWCO) centrifuge filter was used to separate "free"
245 versus "bound" alkaloids in solutions with and without *Os*ABG present, to later be quantified with LC-MS. **(C)** The percent of
246 "free" PTX **251D** (green) dropped when *Os*ABG was present, however the amount of "free" nicotine" remained unchanged by
247 the presence of *Os*ABG.

OsABG is highly expressed in wild frogs and binds ecological toxins

We next sought to better understand the functional role of OsABG in a context relevant to the
ecology and physiology of poison frogs. *Oophaga sylvatica* frogs were collected across three different
locations in Ecuador (Figure 6A). Tissue RNA sequencing revealed that *Os*ABG mRNA is expressed
very highly in the liver compared to other tissues (Figure 6B). However a reanalysis of previously
published proteomics data shows that OsABG is detectable in the liver, skin, and gut, consistent with
movement through the plasma (Figure 6C). Hierarchical clustering of all unique serpinA genes in the
genome shows that OsABG is most closely related to two other serpinA1 genes, Os4677 and Os4682
(Figure 6D). The liver expression of OsABG is higher than all other serpinA genes, and is higher than the
expression of albumin in the liver (Figure 6D). Field-collected *O. sylvatica* frog skin contains alkaloids
from many different classes, with 33% of the summed alkaloid load being histrionicotoxins, followed by
22% in 5,8-indolizidines, 15% in 3,5-indolizidines, 13% in 5,6,8-indolizidines, and 10% in
decahydroquinolines (Figure 6E). Further crosslinking experiments with purified *Os*ABG found that it also
binds a histrionicotoxin-like base ring structure (HTX), indolizidine (indol), and shows slight competition
by a toxin mixture created from wild frog skin extracts (Figure 6F). Together, these data characterize the
expression profile and distribution of OsABG and show that it is capable of binding other alkaloid classes
that are found in wild frogs.

Figure 6: OsABG is expressed in the liver and binds ecologically relevant alkaloids **(A)** Wild *Oophaga sylvatica* were collected across three locations in Ecuador, n=10 per location. **(B)** Comparison of mRNA expression levels across tissues found high expression in the liver, and low to no expression in the skin and gut. **(C)** Reanalysis of proteomics from Caty et al., 2022 found that OsABG protein is present in the gut, liver, and skin. **(D)** The liver expression level of OsABG was higher than that of other members of the serpinA family found in the genome, and of albumin. **(E)** Dorsal skin alkaloids fell into 9 different classes, with the size of the circle representing the averaged percent of total skin alkaloid load. **(F)** Photoprobe binding was competed by PTX, DHQ, epi, a histrionicotoxin-like compound (HTX), and indolizidine ring without R groups (indol), and slightly by a mixture of skin toxins from the wild specimens (TM). Photoprobe binding was not competed by nicotine or cortisol (cort).

3. DISCUSSION

Alkaloid-binding globulin (ABG) represents a new small molecule binding functionality for a member of the serpin family, with a structurally conserved binding pocket similar to mammalian hormone carriers and biliverdin-binding serpin. The broader implication of this finding is that it provides evidence for convergent evolution of serpin proteins for the binding and transport of small molecules across taxa and for different physiological requirements. Most serpin proteins are known for the anti-protease inhibitory activity, however there are members of the serpin superfamily that have been instead characterized as non-inhibitory small molecule binding globulins. In tree frogs, biliverdin-binding serpin (BBS) has been identified as the primary protein that binds biliverdin [48], the heme metabolite that is responsible for green coloration in the lineage, although it remains unclear whether BBS is also responsible for the transport of biliverdin across tissues. In mammals, cortisol-binding globulin (CBG, serpinA6) and thyroxine-binding globulin (TBG, serpinA7) play important roles in the transport and

regulation of plasma hormone concentrations [49,50]. However, serpins are not the only carriers of
lipophilic small molecules in mammalian plasma, with many other hormones and vitamins moving through
plasma on albumin [31,32], Vitamin-D binding protein [51,52], and sex-hormone binding globulin [53,54],
which have no homology to serpin proteins. That CBG and TBG have evolved small molecule transport
from a protease inhibitor background [55] is surprising, and the discovery of ABG shows that this activity
may have evolved either independently or from hormone transport in poison frogs for a vastly different
purpose. BBS, CBG, and TBG all bind their respective ligands in a similar structural pocket [40], however
the identity of the residues that coordinate binding vary across the proteins. Mutations of key binding site
residues in *Os*ABG were able to disrupt binding, confirming the binding pocket predictions and the
functional homology to BBS, CBG, and TBG. A more detailed mutational scanning would be necessary
to fully understand which binding pocket residues coordinate the binding of different alkaloids. It would
be interesting to expand binding experiments to more classes of alkaloids, although we are limited by the
commercial availability of many of these compounds. The similarities between ABG and mammalian
hormone carriers raises the possibility that alkaloids may also act as signaling molecules, which would
require further characterization of other alkaloid targets in poison frogs and more mechanistic information
about ABG alkaloid transport. The identification of this novel alkaloid carrier opens up new avenues of
research in the evolution of small molecule transporters and their relevant physiological functions,
especially in amphibians. The data presented suggest that OsABG is using the binding site already
present in serpin family proteins, and has encoded remarkable specificity for certain alkaloids within this
site.
Alkaloid Binding Globulin (ABG) has different binding specificity across poison frog species and
independent evolutionary origins of acquired chemical defense, suggesting that there exist conserved
mechanisms of alkaloid sequestration across the poison frog clade. Of the species tested, the photoprobe
showed binding activity only in dendrobatid species that can acquire alkaloid chemical defenses from
their diet, namely *O. sylvatica*, *D. tinctorius*, and *E. tricolor*, which represent two independent origins of
chemical defense within the poison frog lineage [17]. This suggests that ABG has evolved in dendrobatid
frogs that are capable of acquired chemical defense, as binding was not seen in the dendrobatid without
chemical defenses in nature (*A. femoralis*), the Malagasy poison frog (*M. aurantiaca*), cane toads, or
human plasma. Malagasy poison frogs represent a convergent evolution of many of the notable
phenotypes in the dendrobatid lineage, and have been found to contain pumiliotoxins and other
dendrobatid alkaloids [56–58]. The lack of binding activity in the plasma may indicate that they have
evolved different molecular mechanisms for alkaloid transport and autoresistance. Using competition of
the photoprobe by presence of excess alkaloid as a proxy for binding activity, we found differences in the
competition activity of different alkaloids across species. This different plasma binding suggests that *O.*
sylvatica has a more promiscuous alkaloid binding pocket, which may be related to the high diversity of
alkaloids found in wild *O. sylvatica* frogs [27,59,60]. Future characterization of plasma binding activity
with more species and alkaloid compounds would be critical to fully trace the evolution of ABG binding
across the poison frog lineage. The identification of these three ABG proteins opens up the possibility of
further discoveries of ABGs in other toxic species, and a more comprehensive understanding of the
biochemical parameters involved in binding structurally different ligands. Overall, the diversity in plasma
binding seen across phylogenetically close species may reflect the diversity in environmental pressures
that have led to species-specific adaptations at the molecular scale.
*Os*ABG is expressed at high levels and binds additional alkaloids found in high abundance on the
skin of field-collected *O. sylvatica*, indicating its physiological and ecological relevance for this poison

frog species. *Os*ABG is produced in the liver as is the case with most members of the serpin family in
humans [61], cows [62], rats [63], baboons [64], and macaques [62]. The liver expression level of *Os*ABG
was much higher than other serpinA proteins and albumin, which in most vertebrates is the most
abundant plasma protein [32,65,66], further highlighting the importance of this protein for poison frog
physiology. Taken together, this supports a model where *Os*ABG is produced in the liver, binds alkaloids
present in the blood or recently absorbed by the intestines, and then may transport alkaloids to the skin
for bioaccumulation. The micromolar K_D value we identified for *Os*ABG with PTX is higher than the
previously reported value for CBG [44,46], and TBG [67], as well as that reported for the "toxin sponge"
protein saxiphilin [68], which are all in the nanomolar range. The lower affinity of *Os*ABG provides further
support to the hypothesis that *Os*ABG may be acting as a transporter protein to other tissues, and would
be in line with the hypothesis that there may be other mechanisms involved in autoresistance to
circulating alkaloids not bound by protein [69]. It is also possible that *Os*ABG has affinities in the
nanomolar range for other ligands which were not tested in this study. Previous work with CBG and TBG
in mammals show that they bind 70-90% of their respective ligands in the plasma, thus regulating the
concentration of free hormone in the blood [50,70]. Furthermore, CBG and TBG have transport activity,
as the release of the ligand is induced by proteolytic cleavage of the reactive center loop which creates
a conformational change of the protein [49,71]. ABG may be acting in a similar way to mediate the
bioaccumulation of dietary alkaloids onto the skin. The reduction of "free" PTX when *Os*ABG is present
in vitro further supports this, given that *Os*ABG would be regulating the pool of unbound alkaloids
available in the plasma. Further characterization using the respective protease from *O. sylvatica* would
be necessary to validate the alkaloid release through proteolytic cleavage of the reactive center loop as
well as genetic knockout frogs to fully characterize the organismal role of *Os*ABG. Given these findings
we conclude that ABG may play this important transport role in poison frog physiology and chemical
defense, which greatly broadens our understanding of this previously elusive mechanism in poison frogs.
Furthermore, given its striking similarities with mammalian hormone carriers and biliverdin binding serpin,
our findings open up additional questions regarding the functional convergence of serpins as small
molecule transporters in distantly related taxa.
360

Summary
This study presents the first evidence of an alkaloid binding protein in the poison frog plasma with a
suggested role as a transporter molecule. We found a novel alkaloid binding function for a member of
the serpin family, contributing to a mounting body of evidence suggesting that the small molecule binding
activity of certain serpins has evolved multiple times from their protease inhibitor precursors with a
structurally conserved binding pocket. Our alkaloid comparisons with *Os*ABG show that this serpin
binding pocket has been fine-tuned for specificity for certain small molecule substrates, and may play an
important ecological and physiological role in the evolution of chemical defense in poison frogs. The
discovery of a serpin that can bind a number of different ligands sets up the possibility of future efforts to
understand and engineer ABG's binding pocket for new ligands once an in-depth understanding of the
binding specificity is achieved.
372

4. MATERIALS AND METHODS
Animal usage
All animal procedures were approved by the Institutional Animal Care and Use Committee at
Stanford (protocol #34153). Topical benzocaine was used for anesthesia prior to the euthanasia of all

animals. Laboratory-bred animals were purchased from Understory Enterprises (Ontario, Canada) or
Josh's Frogs (Michigan, USA) depending on the species. Animals were either euthanized for plasma
collection upon arrival, or housed in 18^3 inch glass terraria, and fed a diet of non-toxic *Drosophila*
melanogaster until euthanasia. Plasma and tissues from a total of 62 animals were used for this study,
consisting of 32 lab-bred animals and 30 field-collected animals that are described below. Sample size
of field-collected animals was determined based on variability seen in previous studies, and for laboratory
experiments based on experimental needs in terms of volume of plasma.
384
Plasma collections
Lab-bred, and therefore non-toxic, poison frogs were anesthetized with topical application of 10%
benzocaine on the ventral skin, and euthanized with cervical translocation. Blood was collected directly
from the cervical cut using a heparinized capillary tube (22-362-566, Fisher Scientific, Waltham, MA) and
deposited into a lithium heparin coated microvette tube (20.1282.100, Sarstedt, Nümbrecht, Germany).
Blood was spun down 10 minutes at 5000 rpm at 4°C on a benchtop centrifuge, and the top layer which
contains the plasma was removed and pipetted into a microcentrifuge tube. This was stored at -80°C
until it was used for experiments.
393
UV crosslinking and competition using alkaloid-like photoprobe
Photocrosslinking methods follow methods outlined in Kim *et al.*, 2020 [72]. Plasma or purified
protein was thawed on ice. The total reaction volume was 50 µL and all experiments were performed in
a clear 96 well plate. For plasma, 5 µL of undiluted plasma was mixed with 40 µL of PBS for each reaction.
For purified *O. sylvatica* protein, 10 µg of protein was diluted into PBS per reaction to a volume of 45 µL.
For *E. tricolor* and *D. tinctorius* protein, 60 µg and 35 µg were used, respectively; a higher amount of
protein was used because no photocrosslinking was detected at 10 µg for these proteins. To this, either
2.5 µL of DMSO was added as vehicle control, or 2.5 µL of competitor compound dissolved in DMSO
was added at a final concentration of 100 µM for plasma competition experiments unless indicated
otherwise, or 1 mM for purified protein experiments unless indicated otherwise. The competitor
compounds were: custom synthesized pumiliotoxin **251D** (PTX, PepTech, Burlington, MA),
decahydroquinoline (DHQ, 125741, Sigma-Aldrich, St. Louis, MO), epibatidine (epi, E1145, Sigma-
Aldrich), a histrionicotoxin-like compound (HTX, ENAH2C55884A-50MG, Sigma-Aldrich), indolizidine
(indol, ATE24584802-100MG, Sigma-Aldrich), nicotine (nic, N3876-100ML, Sigma-Aldrich), cortisol (cort,
H0888-1G, Sigma-Aldrich). The "toxin mixture" (TM) used as a competitor in Figure 6 was made by taking
20µL of each of the skin alkaloid extracts from wild frogs described below, evaporating it under gentle
nitrogen gas flow, and resuspending in 100µL of DMSO. This was followed by addition of 2.5 µL of
photoprobe (Z2866906198, Enamine, Kyiv, Ukraine) dissolved in DMSO on ice, for a final photoprobe
concentration of 5 µM in plasma experiments and 100 µM in purified protein experiments. This was
incubated on ice for 10 minutes, and then UV crosslinked (Stratalinker UV 1800 Crosslinker, Stratagene,
La Jolla, CA) for 5 minutes on ice. TAMRA visualization of crosslinked proteins was done by adding 3 µl
TBTA (stock solution: 1.7 mM in 4:1 v/v DMSO:tert-Butanol; H66485-03, Fisher), 1 µl Copper (II) Sulfate
(stock solution: 50 mM in water; BP346-500, Fisher), 1 µl Tris (2-carboxyethyl) phosphine hydrochloride
(freshly prepared, stock solution: 50 mM; J60316-06, Fisher), and 1 µl TAMRA-N3 (stock solution: 1.25
mM in DMSO; T10182, Fisher), incubating at room temperature for 1 hour, and quenching the reaction
by boiling with 4x SDS loading buffer. This was run on a Nupage 4-12% Bis-Tris protein gel (NP0323BOX,
Invitrogen, Waltham, MA) and the in-gel fluorescence of the gel was visualized using a LI-COR Odyssey

imaging system (LI-COR Biosciences, Lincoln, Nebraska) at 600 nm for an exposure time of 30 seconds.
After imaging the TAMRA signal, the same gel was coomassie stained (InstantBlue, ISB1L, Abcam,
Cambridge, UK) and visualized the same way at 700 nm.
For proteomic identification of competed proteins, plasma samples were pooled from five different
individuals and were crosslinked with either no photoprobe and equivalent amounts of DMSO, 5 μM
photoprobe and DMSO, or 5 μM photoprobe and 100 μM PTX as described above. Each condition was
set up as 24 individual reactions and pooled after crosslinking. To attach a biotin handle, 3 μl TBTA, 1 ul
CuSO4, 1 μl TCEP, and 1.14 μl Biotin-N3 (stock solution: 9.67 mM in DMSO; 1265, Click Chemistry
Tools, Scottsdale, AZ), were added for each reaction and this was incubated at room temperature for 1
hour, rotating. After incubation, each condition was run through a 3kDa MWCO centrifuge filter twice
(UFC800324, Amicon, Millipore-Sigma, Burlington, MA) to dilute excess Biotin-N3 until reaching a 900x
dilution. Pulldown of biotinylated photoprobe-protein complexes was achieved with a magnetic bead strep
pulldown following the protocol outlined in Wei *et al*., 2021[73]. The pre and post-pulldown samples were
run on a gel for a streptavidin blot (Figure 2A) and silver stain. After verifying pulldown efficacy, samples
were run on SDS-PAGE gels in two replicates (one lane each) for each condition. Gels were fixed in
50:50 water:MeOH with 10% Acetic Acid for 1-2 hours. For the first replicate, the gel was run for a short
period, and an approximately one centimeter squared portion containing the whole lane for each condition
was excised and fixed. For the second replicate, the gel was run completely and the proteins between
39 kDa and 64 kDa were excised and fixed using the ladder as a size reference. Sections were chopped
into 1 mm pieces under sterile conditions and stored at 4°C in 100μL of water with 1% acetic acid until
processed for proteomics.
442
Proteomic identification of pulled down proteins across conditions
For proteomics analyses, SDS-PAGE gel slices approximately 1 cm in length were prepared for
proteolytic digestion. Each fixed gel slice was diced into 1 mm cubes under sterile conditions, and then
rinsed with 50 mM ammonium bicarbonate to remove residual acidification from the fixing process.
Following rinsing, the gels were incubated in 80% acetonitrile in water for five minutes; the solvent was
removed and then the gel pieces were incubated with 10 mM DTT dissolved in water at room temperature
for 20 minutes. Following reduction, alkylation was performed using 30 mM acrylamide for 30 minutes at
room temperature to cap free reduced cysteines. Proteolysis was performed using trypsin/lysC
(Promega, Madison, WI) in 50 mM ammonium bicarbonate overnight at 37°C. Resulting samples were
spun to pellet gel fragments prior to extraction of the peptides present in the supernatant. The resulting
peptides were dried by speed vac before dissolution in a reconstitution buffer (2% acetonitrile with 0.1%
formic acid), with an estimated 1 μg on-column used for subsequent LC-MS/MS analysis.
The liquid chromatography mass spectrometry experiment was performed using an Orbitrap
Eclipse Tribrid mass spectrometer RRID:022212 (Thermo Scientific, San Jose, CA) with liquid
chromatography using an Acquity M-Class UPLC (Waters Corporation, Milford, MA). A flow rate of 300
nL/min was used, where mobile phase A was 0.2% formic acid in water and mobile phase B was 0.2%
formic acid in acetonitrile. Analytical columns were prepared in-house with an I.D. of 100 microns pulled
to a nanospray emitter using a P2000 laser puller (Sutter Instrument, Novato, CA). The column was
packed using C18 reprosil Pur 1.8 micron stationary phase (Dr. Maisch) to an approximate length of 25
cm. Peptides were directly injected onto the analytical column using a 80 minute gradient (2%–45% B,
followed by a high-B wash). The mass spectrometer was operated in a data dependent fashion using
CID fragmentation in the ion trap for MS/MS spectra generation.

465  For data analysis, the .RAW data files were processed using Byonic v4.1.5 (Protein Metrics,
466 Cupertino, CA) to identify peptides and infer proteins based on a proteomic reference created with the
467 *O. sylvatica* genome annotation. Proteolysis with Trypsin/LysC was assumed to be specific with up to 2
468 missed proteolytic cleavages. Precursor mass accuracies were held within 12 ppm, and 0.4 Da for
469 MS/MS fragments in the ion trap. Cysteine modified with propionamide were set as fixed modifications
470 in the search, and other common modifications (e.g. oxidation of methionine) were also included. Proteins
471 were held to a false discovery rate of 1%, using standard reverse-decoy technique [74].
472

473 **Identification of ABG proteins in different species and sequence confirmation**
474  To identify potential ABG proteins in other species, we used the *Os*ABG protein sequence
475 identified in the proteomics as the query and searched against blast databases created from the *Allobates*
476 *femoralis* genome, and *Epipedobates tricolor*, *Dendrobates tinctorius*, and *Mantella aurantiaca*
477 transcriptomes. The top hit from each blast search was used as the most probable ABG gene from those
478 species. To ensure that the sequences did not contain sequencing or alignment errors, the gene from *O.*
479 *sylvatica, D. tinctorius, and E. tricolor* was amplified using PCR and sequence confirmed with sanger
480 sequencing. Total RNA was extracted from flash frozen liver tissue from three lab-bred, non-toxic,
481 individuals from each species using the Monarch total RNA Miniprep Kit (T2010S, New England Biolabs,
482 Ipswich, MA) following manufacturer instructions. This was used to create cDNA using the SuperScript
483 III First-Strand Synthesis kit (18080-400, Invitrogen), following manufacturer instructions with an
484 oligo(dT)20 primer. This was used for a PCR using Phusion High Fidelity DNA polymerase (F-530,
485 Thermo Scientific) and the primers and cycling conditions described below. PCRs were analyzed using
486 a 1% agarose gel for presence of a single band, cleaned up (NucleoSpin Gel and PCR cleanup,
487 740609.50, Takara Bio, Shiga, Japan), and transformed into pENTR vectors using a D-TOPO kit (45-
488 0218, Invitrogen). Plasmids containing the ABG sequences from each individual were then mini prepped
489 (27106X4, Qiagen, Hilden, Germany) and sanger sequenced with M13F and M13R primers (Azenta Life
490 Sciences, South San Francisco, California). Sequences were aligned using Benchling (Benchling Inc,
491 San Francisco, California) software, with MAFFT used for DNA alignments and Clustal Omega used for
492 protein alignments.
493

species	fwd primer	rev primer	Tm
O. sylvatica	CACCATGAAACTTTTCGTCTACCT GTGTTTCAGC	CTATTTTGTTGGGTCTACTATTCTT CCGCTG	68°C
D. tinctorius	CACCATGAAGCTTTTCGTCTTCCT ATGTTTCAGCC	CTATTTTGTTGGGTTTATTATTTTTC CATTCAAAATATCG	66°C
E. tricolor	CACCATGAAGCTTTTCATCTTCCT GTGTTTGAGCC	CTATTTTGTTGGGTCTATTATTCTT CCGGAGAAAAC	68°C

494 Cycling conditions: 98°C for 30 seconds, [98°C for 10 seconds, Tm for 30 seconds, 72°C for 2 minutes]
495 x 34 cycles, 72°C for 10 minutes
496

497 **Protein structure prediction and molecular docking analyses**
498  The *Os*ABG protein folding was predicted using the amino acid sequence, edited for point
499 mutations found across all three individuals used for sequence verification, and the AlphaFold google

colab   notebook
(https://colab.research.google.com/github/deepmind/alphafold/blob/main/notebooks/AlphaFold.ipynb)[7
5]. The predicted structure is provided in the supplementary information. The default AlphaFold
parameters were used. Molecular docking was performed using the UCSF Chimera software
(https://www.cgl.ucsf.edu/chimera/)[76], using AutoDock Vina [77,78] with the three dimensional
structure of PTX **251D** (Pubchem CID 6440480). The whole protein was used as the search space with
the default search parameters (5 binding modes, exhaustiveness of search of 8, and a maximum energy
difference of 3 kcal/mol). The docking result with the highest predicted affinity was used and is included
in the supplementary files. Protein structures and docking were visualized using PyMol for publication
quality images.

Recombinant protein expression and binding assays

Recombinant ABG proteins were expressed by Kemp Proteins (Maryland, USA) through their
custom insect cell protein expression and purification services. The reagents and vectors used are
proprietary to Kemp Proteins, however the general expression and purification details are as follows. The
verified protein sequences described above or the point mutations (Figure 4) were codon-optimized for
SF9 insect expression, and a 10xHIS tag was added to the C-terminal end. For *Os*ABG, a 1 L expression
was performed, for all other sequences (other species and mutants) a 50 mL expression was used. For
the 1 L expression, a multiplicity of infection of one was used for the p1 baculovirus and the supernatant
was collected after 72 hours. To this, 5 mL of Qiagen Ni-NTA resin washed and equilibrated in Buffer A
(20 mM Sodium Phosphate, 300 mM NaCl, pH=7.8) was added and it was mixed overnight at 4°C.
Afterwards, it was packed in a 5 mL Bio-Scale column and washed with 3 column volumes (CV) of Buffer
A, followed by washing with 5% Buffer B (20 mM Sodium Phosphate, 300 mM NaCl, 500 mM Imidazole,
pH=7.8) for 5 CV. Protein was eluted with a linear gradient from 5-60% over 25 CV, and 6 mL fractions
were collected throughout. All of the fractions containing protein were pooled and concentrated to 1
mg/mL using an Amicon centrifugal filter with a 10 kDa MWCO, the buffer was exchanged to PBS, it was
filtered through a 0.2 um filter, aliquoted, and frozen at -80°C. Protein expression and purification resulted
in a clear band by western blot (Figure S4A) and a clean doublet pattern by coomassie (Figure S4B)
closely resembling that seen in the plasma crosslinking results (Figure 1C) in both reduced and non-
reduced conditions. For the 50mL expression of *Dt*ABG, *Et*ABG, and mutant *Os*ABG proteins, a 10%
ratio of p1 virus to media was used and the supernatant was collected after 72 hours, to which 1 mL of
Qiagen Ni-NTA resin washed and equilibrated in Buffer A (20 mM Sodium Phosphate, 300 mM NaCl,
pH=7.4) was added. This was mixed overnight at 4°C and then packed into a 1 mL Bio-Scale column,
washed with 3 CV of Buffer A, washed with 5 CV of 5% Buffer B (20 mM Sodium Phosphate, 300 mM
NaCl, 500 mM imidazole, pH=7.4), and eluted with 5 CV of 50% Buffer B. Fractions containing protein
were buffer exchanged into PBS, and the final concentrations were approximately 0.2 mg/mL, with
varying final volumes. Protein expression and purification resulted in a clear band by western blot (Figure
S4C,E,G,I,K), and a clean doublet by coomassie (Figure S4D,F,H,J,L) in both reduced and non-reduced
conditions.

Determination of dissociation constant using Microscale Thermophoresis (MST)

To determine the binding affinity of *Os*ABG for PTX, we used Microscale Thermophoresis (MST)
to determine the dissociation constant (K_D). To do this, we used the Monolith system (Nanotemper
Technologies, München, Germany). Purified *Os*ABG protein was labeled using the protein labeling kit

Red-NHS 2nd generation (MO-L011, Nanotemper) which dyes primary lysine residues in the protein. The
kit was used following manufacturer instructions, however a 1.5x excess of dye was used instead of 3x
as this was found to better achieve a degree of labeling of ~0.5. To remove aggregates during the assay,
PBS-Tween was used for protein labeling and all dilutions. The labeled protein was centrifuged for 10
minutes at 20,000g on a benchtop centrifuge, and the supernatant was taken to further remove any
aggregation. The concentration was measured prior to calculating and setting up dilution series. A final
concentration of 10nM $OsABG$ was used, and a 16 tube 2x serial dilution series of PTX 251D was made
with the highest concentration being 1000uM. The concentration of DMSO was maintained consistent
across the dilution series. The labeled $OsABG$ was incubated for 10 minutes prior to loading into
capillaries, and three biological replicates were pipetted and run separately. The Monolith premium
capillaries (MO-K025, Nanotemper) were used, the MST power was set to Medium, and the excitation
power was set to auto-detect. The three replicates were compiled and plotted together using GraphPad
Prism (GraphPad Software, San Diego, California), and a dissociation model was fit to the data to
calculate the K_D. The raw data is included in the supplementary information (will be included with full
submission).
559
Determination of free versus bound alkaloids
Solutions with 4 uM of $OsABG$ protein, 4 uM of either PTX **251D** or nicotine, and a final volume
of 100 uL were made and incubated for one hour at room temperature. This was transferred to a 3 kDa
MWCO centrifugal filter (UFC500396, Amicon) and spun at max speed on a benchtop centrifuge at 4C
for 45 minutes. The top and bottom fractions were brought up to 100 uL with ultrapure water and
transferred to new tubes, where 300 uL of 2:1 Acetonitrile:Methanol was added, after which they were
vortexed and centrifuged at max speed on a benchtop centrifuge at 4C for 10 minutes. The supernatant
was transferred to autosampler vials for quantitation of the amount of alkaloid in each fraction with mass
spectrometry. Each condition was run in triplicate. Samples were analyzed using an Agilent Quadrupole
time-of-flight LC-MS instrument, with MS analysis performed by electrospray ionization (ESI) in positive
mode. Metabolites were separated with an Eclipse Plus C18 column (Agilent 959961-902) with normal
phase chromatography. Mobile phases were: buffer A (water with 0.1% formic acid) and buffer B (90%
acetonitrile, 10% water with 0.1% formic acid). The flow rate was maintained constant at 0.7 mL/min
throughout the LC protocol. The LC gradient elution was set as follows: starting at 5% B held till 0.51
minutes, linear gradient from 5 to 25% B in 1.5 minutes, linear gradient from 25 to 50% B in 23 minutes,
linear gradient from 50 to 95% B in 30 seconds, 95% B held for 2 minutes, linear gradient from 95 to 5%
B in 1 minute, and 5% B held for 1.5 minutes to equilibrate the column to the initial conditions. The total
run time was 30 minutes and the injection volume was 10 uL. Data was analyzed using the Agilent
MassHunter software; the extracted ion chromatograms for PTX were searched using the exact mass
M+1 of 252.2333, and nicotine was searched using the exact mass M+1 of 163.123, with a tolerance of
a symmetric +/- 100 ppm. Extracted ion chromatograms were smoothed once before automatically
integrating to get the abundance values. Abundance values were used to calculate the fractions above
and below the filter for each replicate, which were then plotted with GraphPad. All raw data is provided
as mzXML files through DataDryad (accessing information will be added for full submission)..
584
Field collections of *Oophaga sylvatica*
The frog samples used in this paper are the same as those used for the project described in
Moskowitz *et al*., 2022 [79]. For each location, 10 *O. sylvatica* individuals were collected under collection

permit 0013-18 IC-FAU-DNB/MA issued by the Ministerio del Ambiente de Ecuador, between the hours
of 7:00-18:00 during early May to early June 2019. All individuals were euthanized the same day as
collection. Prior to euthanizing, frogs were sexed, weighed, and the snout-vent length was measured.
Orajel (10% benzocaine) was used as an anesthetic prior to cervical dislocation. Once euthanized, frogs
were immediately dissected and the liver, intestines, and half of the dorsal skin were stored in RNAlater
in cryotubes at room temperature. The other half of the dorsal skin was placed in methanol in glass tubes
at room temperature. Once back in the lab, all tissues were stored at -20°C until further processing. All
tissues were transported to the United States under CITES permits 19EC000036/VS, 19EC000037/VS,
19EC000038/VS.
597
Alkaloid extraction, detection, and analysis
All following steps were performed under a hood. Skins were taken out of methanol with forceps
and weighed. From the methanol that the skin was stored in, 1 mL was taken and syringe filtered through
a 0.45µ PTFE filter (44504-NP, Thermo Scientific) into the new glass vial with a PTFE cap (60940A-2,
Fisher) filled with 25 µL of 1 µg/µL (-)-Nicotine (N3876-100ML, Sigma Aldrich), for a total of 25 µg of
added nicotine. Tubes were capped and vortexed, and stored at -80°C celsius for 24 hours, during which
proteins and lipids should precipitate. After 24 hours, tubes were taken out of the -80°C and quickly
syringe filtered through a 0.45µ PTFE filter again into a new glass vial. A 100 µL aliquot was added to an
GC-MS autosampler vial, and remaining solution was stored in the original capped vial at -80°C.
GC-MS analysis was performed on a Shimadzu GCMS-QP2020 instrument with a Shimadzu 30m
x 0.25 mmID SH-Rxi-5Sil MS column closely following the protocol outlined in Saporito *et al*., 2010 [80].
In brief, GC separation of alkaloids was achieved using a temperature program from 100 to 280°C at a
rate of 10°C per minute with He as the carrier gas (flow rate: 1 mL/min). This was followed by a 2 minute
hold and additional ramp to 320°C at a rate of 10°C/minute for column protection reasons, and no
alkaloids appeared during this part of the method. Compounds were analyzed with electron impact-mass
spectrometry (EI-MS). The GC-MS data files were exported as CDF files and the Global Natural Products
Social Network (GNPS) was used to perform the deconvolution and library searching against the AMDIS
(NIST) database to identify all compounds (https://gnps.ucsd.edu) [81]. For deconvolution (identification
of peaks and abundance estimates) the default parameters were used, for the library search the precursor
ion mass tolerance was set to 20000 Da and the MS/MS fragment ion tolerance to 0.5 Da. The resulting
dataset was filtered to keep only compounds that matched to our spiked-in nicotine standard, alkaloids
previously found in poison frogs from the Daly 2005 database [15], or compounds with the same base
ring structure and R groups as the classes defined in Daly 2005. All GC-MS data as CDF files are
available through the GNPS public data repository (accessing information will be added for full
submission).
Once the feature table from the GNPS deconvolution was filtered to only include only poison frog
alkaloids and nicotine, the abundances values were normalized by dividing by the nicotine standard and
skin weight. This filtered and normalized feature table was used for all further analyses and visualizations.
All steps were carried out with R version 4.0.4, and code is included in supplementary data (will be
included with full submission).
628
RNA extraction and library preparation
RNA extraction followed the Trizol (15596018, Thermo Fisher) RNA isolation protocol outlined in
Caty *et al.* 2019 [27] according to the manufacturer's instructions, and with sample randomization to avoid

batch effects. RNA quality was measured on a Agilent Tapestation RNA screentape (Agilent, Santa Clara,
CA), and quantified using a Qubit Broad Range RNA kit (Q10210, Invitrogen). In the liver and intestines,
samples with RIN scores greater than 5 were kept, RNA was normalized to the same Qubit concentration,
and mRNA was isolated and library prepped using the NEB Directional RNA sequencing kit (E7765L,
New England Biolabs) with the PolyA purification bundle (E7490L, New England Biolabs) and 96 Unique
Dual Indices (E7765L, New England Biolabs). The skin RIN scores were much lower, signaling potential
RNA degradation, ribosomal degradation was instead used to isolate mRNA. Following normalization
within all skin RNA samples to the same Qubit concentration, we used the Zymo RiboFree Total RNA
Library Prep kit (R3003-B, Zymo Research, Irvine, CA) following manufacturer instructions. After library
prep for all tissues was complete library size was quantified with the Agilent Tapestation D1000
screentape, and concentration was measured with the Qubit dsDNA high sensitivity kit (Q33231,
Invitrogen). All libraries within a tissue type were pooled to equimolar amounts and sequenced on two
lanes of an Illumina NovaSeq (Illumina, San Diego, CA) machine to obtain 150 bp paired-end reads.
645
RNA expression analysis and identification of *O. sylvatica* serpinA genes
Analysis of RNA expression levels followed the protocol outlined by Payne *et al*., 2022 [82]. The
Trim-galore! wrapper tool [83] was used to trim adapter sequences with cutadapt [84] and quality filter
the reads (trim_galore --paired --phred33 --length 36 -q 30 --stringency 1 -e 0.001). All trimmed reads are
available through the NCBI BioProject (accessing information will be added for full submission). *Kallisto*
[85] was used to pseudoalign the reads to a reference created with the coding sequence of the annotated
O. sylvatica genome. These abundances were combined into a matrix, and the trimmed-mean of M-
values (TMM) normalized counts were used for all further analyses. Additional serpinA genes were found
in the genome by searching for all genes annotated with "serpina" in the header, and by blasting the
*Os*ABG protein sequence against the genome (e-value < 1e-60) and including any additional genes not
annotated with "serpina." Four sequences were removed because they were exact matches of the full
gene   (OopSylGTT00000004683),   the   N-terminal   end   (OopSylGTT00000004650,
OopSylGTT00000004685), or the C-terminal (OopSylGTT00000004676) end sequence of another
serpina gene, and therefore could be potential duplications caused by annotation or assembly errors. To
create the protein tree (Figure 6D), ClustalW was used to align the sequences, a distance matrix was
created using identity, and neighbor joining was used to construct the tree. The albumin gene was
determined by blasting the protein sequences of *Xenopus laevis* albumin A (Uniprot #P08759), *X. laevis*
albumin B (Uniprot #P14872), and the Asian toad *Bombina maxima* albumin (Uniprot #Q3T478) against
the *O. sylvatica* genome. In all three cases, the top hit was the same (OopSylGTT00000003067),
therefore this was assumed to be the most likely albumin candidate in the genome and was used to plot
the TMM expression for comparison. All plots were created in R version 4.0.4, and all analysis and plotting
code is available in the supplementary files (will be included with full submission).
668

5. ACKNOWLEDGEMENTS
The authors acknowledge that this research was conducted on the ancestral lands of the Muwekma
Ohlone people at Stanford; we understand the implications of the historical and present colonialism
experienced by the Ohlone people and celebrate their continued stewardship of their lands. We thank
Andrea Terán Valdéz for their assistance coordinating field work and their kindness. We would also like
to thank the Laboratory of Organismal Biology, the Long Lab, the Soh lab, Joel Francis, and Cheyenne

Payne for helpful discussions and guidance throughout this project. Centro Jambatu researchers thank
the Saint Louis Zoo for their commitment and sustained support for amphibian research.
677

6. FUNDING
This work was supported by the National Science Foundation (IOS-1822025) and the New York Stem
Cell Foundation (LAO). This work was also supported by the Vincent Coates Foundation Mass
Spectrometry Laboratory, Stanford University Mass Spectrometry (RRID:SCR_017801) utilizing the
Thermo Orbitrap Eclipse nanoLC/MS system (RRID:SCR_022212) that was purchased with funding from
National Institutes of Health Shared Instrumentation grant 1S10OD030473. This work was supported in
part by NIH P30 CA124435 utilizing the Stanford Cancer Institute Proteomics/Mass Spectrometry Shared
Resource. AAB is supported by a NSF Graduate Research Fellowship (DGE-1656518) and an HHMI
Gilliam Fellowship (www.hhmi.org, GT13330). MDMG was supported by the Fundacion Alfonso Martin
Escudero postdoctoral fellowship. LAO is a New York Stem Cell Foundation – Robertson Investigator.

7. DATA AVAILABILITY
All raw data, analysis scripts, and intermediate data analysis files will be publicly available at the time of
publication.
692

693 **8. COMPETING INTERESTS**
694 The authors have filed a provisional patent application based on the idea of using alkaloid-binding
695 globulins as anti-toxin sponges.
696

9. AUTHOR CONTRIBUTIONS
AAB - Conceptualization, Methodology, Validation, Formal Analysis, Investigation, Data Curation, Writing
- Original Draft, Writing - Review and Editing, Visualization
MDMG - Methodology, Validation, Formal Analysis, Investigation
AER - Methodology, Validation, Formal Analysis, Investigation
ET - Methodology, Validation, Investigation
JT - Methodology
LAC - Methodology, Resources
HTS - Methodology, Resources
JZL - Conceptualization, Methodology, Resources, Writing - Review and Editing, Supervision
LAO - Conceptualization, Resources, Writing - Review and Editing, Supervision, Funding Acquisition
708

709 **10. SUPPLEMENTARY MATERIAL**

710
Figure S1: Dose response of photoprobe competition (A) Plasma from *O. sylvatica* shows crosslinking to the
photoprobe, and competition by pumiliotoxin **251D** (PTX) occurs when there is 10:1 PTX:photoprobe concentration

in the reaction. **(B)** Purified O. sylvatica Alkaloid Binding Globulin (ABG) shows crosslinking to the photoprobe and
competition by PTX when there is a 1:1 PTX:photoprobe concentration. **(C)** Plasma from *D. tinctorius* shows
crosslinking to the photoprobe, and competition by PTX occurs when there is 10:1 PTX:photoprobe concentration
in the reaction.
717

718
719   **Figure S2: Peptide coverage over ABG protein sequence** All unique peptides from one replicate of the
720   proteomics were mapped onto the protein sequence of *O. sylvatica* ABG, showing that there were peptides covering
721   the full length of the protein.
722

723
Figure S3: Structure and binding pockets of closely related serpinA proteins (A) Crystal structure for human
SerpinA1/alpha-1-antitrypsin (PDB# 1HP7) contains the structural elements of other serpin binding pockets (black
box), however is not documented to have small-molecule binding capabilities. **(B)** Close-up of A1AT (PDB# 1HP7),
with pocket residues highlighted in magenta. **(C)** Crystal structure for human SerpinA7/thyroxine-binding globulin
(PDB# 2RIW) binds thyroxine (orange) in the same structural pocket as other serpins. **(D)** Close-up of thyroxine
binding pocket of TBG (PDB# 2RIW), with proximal residues highlighted in magenta. Thyroxine structure is
displayed on the top right. **(E)** Alignment of proximal residues (within 5 angstroms of small molecule) across small
molecule binding serpins *Os*ABG, biliverdin binding serpin (BBS, PDB# 7RBW), corticosteroid binding globulin
(CBG, PDB# 2V95), and thyroxine binding globulin (TBG, PDB# 2RIW) shows that most residues that may be
involved in coordinating small molecule binding are not conserved, despite the structural conservation of the binding

pocket. Percentages indicate the total percent identity of the protein sequences, the small number above each
residue indicates the position of that amino acid in the protein sequence. Only proximal residues are shown, blank
spaces are not representative of any specific sequence or spacing.

Figure S4: Recombinant expression and purification of Alkaloid Binding Globulin (ABG) proteins
in insect cells. HIS blot **(A)** and coomassie **(B)** of the recombinantly expressed and purified *O. sylvatica*
alkaloid-binding globulin (*Os*ABG) show a clear doublet pattern in both reduced (R) and non-reduced
(NR) conditions. HIS blot **(C)** and coomassie **(D)** of recombinantly expressed *D. tinctorius* ABG, identified
by the most-homologous protein to OsABG in the *D. tinctorius* transcriptome, show a clear purification.
HIS blot **(E)** and coomassie **(F)** of recombinantly expressed *E. tricolor* ABG, identified by the most-
homologous protein to OsABG in the *E. tricolor* transcriptome, show a clear purification. HIS blot **(G)** and
coomassie **(H)** of recombinantly expressed *Os*ABG mutant 1 (Y36A, W276A, S374A, D383A) shows a
clear purification. HIS blot **(I)** and coomassie **(J)** of recombinantly expressed *Os*ABG mutant 2 (Y36A,

S268A, D273A, D383A) shows a clear purification. HIS blot **(K)** and coomassie **(L)** of recombinantly
expressed *Os*ABG mutant 3 (D383A) shows a clear purification.
750

11. REFERENCES

1. Wright SN. Irreversible block of human heart (hH1) sodium channels by the plant alkaloid
lappaconitine. Mol Pharmacol. 2001;59: 183–192. doi:10.1124/mol.59.2.183

2. Wang S-Y, Tikhonov DB, Mitchell J, Zhorov BS, Wang GK. Irreversible block of cardiac mutant
Na+ channels by batrachotoxin. Channels . 2007;1: 179–188. doi:10.4161/chan.4437

3. Efimova SS, Zakharova AA, Ostroumova OS. Alkaloids Modulate the Functioning of Ion Channels
Produced by Antimicrobial Agents via an Influence on the Lipid Host. Front Cell Dev Biol. 2020;8:
537. doi:10.3389/fcell.2020.00537

4. Asakawa M, Matsumoto T, Umezaki K, Kaneko K, Yu X, Gomez-Delan G, et al. Toxicity and Toxin
Composition of the Greater Blue-Ringed Octopus from Ishigaki Island, Okinawa Prefecture, Japan.
Toxins . 2019;11. doi:10.3390/toxins11050245

5. Matsumura K. Tetrodotoxin as a pheromone. Nature. 1995;378: 563–564. doi:10.1038/378563b0

6. Bucciarelli GM, Li A, Zimmer RK, Kats LB, Green DB. Quantifying tetrodotoxin levels in the
California newt using a non-destructive sampling method. Toxicon. 2014;80: 87–93.
doi:10.1016/j.toxicon.2014.01.009

7. Hartmann T, Theuring C, Schmidt J, Rahier M, Pasteels JM. Biochemical strategy of sequestration
of pyrrolizidine alkaloids by adults and larvae of chrysomelid leaf beetles. J Insect Physiol.
1999;45: 1085–1095. doi:10.1016/s0022-1910(99)00093-1

8. Cogni R, Trigo JR, Futuyma DJ. A free lunch? No cost for acquiring defensive plant pyrrolizidine
alkaloids in a specialist arctiid moth (Utetheisa ornatrix). Mol Ecol. 2012;21: 6152–6162.
doi:10.1111/mec.12086

9. Wang L, Beuerle T, Timbilla J, Ober D. Independent recruitment of a flavin-dependent
monooxygenase for safe accumulation of sequestered pyrrolizidine alkaloids in grasshoppers and
moths. PLoS One. 2012;7: e31796. doi:10.1371/journal.pone.0031796

10. Strauss AS, Peters S, Boland W, Burse A. ABC transporter functions as a pacemaker for
sequestration of plant glucosides in leaf beetles. Elife. 2013;2: e01096. doi:10.7554/eLife.01096

11. Yotsu-Yamashita M, Sugimoto A, Terakawa T, Shoji Y, Miyazawa T, Yasumoto T. Purification,
characterization, and cDNA cloning of a novel soluble saxitoxin and tetrodotoxin binding protein
from plasma of the puffer fish, Fugu pardalis. Eur J Biochem. 2001;268: 5937–5946.
doi:10.1046/j.0014-2956.2001.02547.x

12. Yotsu-Yamashita M, Nagaoka Y, Muramoto K, Cho Y, Konoki K. Pufferfish Saxitoxin and
Tetrodotoxin Binding Protein (PSTBP) Analogues in the Blood Plasma of the Pufferfish , , , and.
Mar Drugs. 2018;16. doi:10.3390/md16070224

13. Abderemane-Ali F, Rossen ND, Kobiela ME, Craig RA, Garrison CE, Chen Z, et al. Evidence that
toxin resistance in poison birds and frogs is not rooted in sodium channel mutations and may rely
on "toxin sponge" proteins. J Gen Physiol. 2021;153. doi:10.1085/jgp.202112872

14. Daly JW, Kaneko T, Wilham J, Garraffo HM, Spande TF, Espinosa A, et al. Bioactive alkaloids of
frog skin: Combinatorial bioprospecting reveals that pumiliotoxins have an arthropod source.
Proceedings of the National Academy of Sciences. 2002;99: 13996–14001.
doi:10.1073/pnas.222551599 PMID - 12381780

15. Daly JW, Spande TF, Garraffo HM. Alkaloids from Amphibian Skin: A Tabulation of Over Eight-
Hundred Compounds. J Nat Prod. 2005;68: 1556–1575. doi:10.1021/np0580560 PMID - 16252926

16. Saporito RA, Spande TF, Martin Garraffo H, Donnelly MA. Arthropod Alkaloids in Poison Frogs: A
Review of the "Dietary Hypothesis." HETEROCYCLES. 2009. p. 277. doi:10.3987/rev-08-sr(d)11

17. Santos JC, Coloma LA, Cannatella DC. Multiple, recurring origins of aposematism and diet
specialization in poison frogs. Proceedings of the National Academy of Sciences. 2003;100:
12792–12797. doi:10.1073/pnas.2133521100 PMID - 14555763

18. Summers K. Convergent evolution of bright coloration and toxicity in frogs. Proceedings of the
National Academy of Sciences. 2003;100: 12533–12534. doi:10.1073/pnas.2335928100 PMID -
14569014

19. Daly JW, Secunda SI, Garraffo HM, Spande TF, Wisnieski A, Cover JF. An uptake system for
dietary alkaloids in poison frogs (Dendrobatidae). Toxicon. 1994;32: 657–663. doi:10.1016/0041-
0101(94)90335-2 PMID - 7940573

20. Caldwell JP. The evolution of myrmecophagy and its correlates in poison frogs (Family
Dendrobatidae). J Zool. 1996;240: 75–101. doi:10.1111/j.1469-7998.1996.tb05487.x

21. Darst CR, Menéndez-Guerrero PA, Coloma LA, Cannatella DC. Evolution of dietary specialization
and chemical defense in poison frogs (Dendrobatidae): a comparative analysis. Am Nat. 2004;165:
56–69. doi:10.1086/426599 PMID - 15729640

22. Toft CA. Feeding ecology of thirteen syntopic species of anurans in a seasonal tropical
environment. Oecologia. 1980;45: 131–141. doi:10.1007/bf00346717 PMID - 28310947

23. Vandendriessche T, Abdel-Mottaleb Y, Maertens C, Cuypers E, Sudau A, Nubbemeyer U, et al.
Modulation of voltage-gated Na+ and K+ channels by pumiliotoxin 251D: A "joint venture" alkaloid
from arthropods and amphibians. Toxicon. 2008;51: 334–344. doi:10.1016/j.toxicon.2007.10.011
PMID - 18061227

24. Daly JW, Gusovsky F, McNeal ET, Secunda S, Bell M, Creveling CR, et al. Pumiliotoxin alkaloids:
A new class of sodium channel agents. Biochem Pharmacol. 1990;40: 315–326. doi:10.1016/0006-
2952(90)90694-g PMID - 2165404

25. Okada T, Wu N, Takashima K, Ishimura J, Morita H, Ito T, et al. Total Synthesis of
Decahydroquinoline Poison Frog Alkaloids ent-cis-195A and cis-211A. Molecules. 2021. p. 7529.
doi:10.3390/molecules26247529

26. Spande TF, Garraffo HM, Edwards MW, Yeh HJC, Pannell L, Daly JW. Epibatidine: a novel
(chloropyridyl)azabicycloheptane with potent analgesic activity from an Ecuadoran poison frog.
Journal of the American Chemical Society. 1992. pp. 3475–3478. doi:10.1021/ja00035a048

27. Caty SN, Alvarez-Buylla A, Byrd GD, Vidoudez C, Roland AB, Tapia EE, et al. Molecular
physiology of chemical defenses in a poison frog. J Exp Biol. 2019;222: jeb204149.

826   doi:10.1242/jeb.204149 PMID - 31138640

827 28. Alvarez-Buylla A, Payne CY, Vidoudez C, Trauger SA, O'Connell LA. Molecular physiology of
828   pumiliotoxin sequestration in a poison frog. PLoS One. 2022;17: e0264540.
829   doi:10.1371/journal.pone.0264540

830 29. O'Connell LA, LS50: Integrated Science Laboratory Course, O'Connell JD, Paulo JA, Trauger SA,
831   Gygi SP, et al. Rapid toxin sequestration modifies poison frog physiology. J Exp Biol. 2021;224.
832   doi:10.1242/jeb.230342

833 30. Tarvin RD, Borghese CM, Sachs W, Santos JC, Lu Y, O'Connell LA, et al. Interacting amino acid
834   replacements allow poison frogs to evolve epibatidine resistance. Science. 2017;357: 1261–1266.
835   doi:10.1126/science.aan5061 PMID - 28935799

836 31. Peters T, Jr. All About Albumin: Biochemistry, Genetics, and Medical Applications. Academic
837   Press; 1995. Available: https://play.google.com/store/books/details?id=i1DC3KlTAB8C

838 32. Baker ME. Albumin, steroid hormones and the origin of vertebrates. J Endocrinol. 2002;175: 121–
839   127. doi:10.1677/joe.0.1750121

840 33. Czub MP, Handing KB, Venkataramany BS, Cooper DR, Shabalin IG, Minor W. Albumin-Based
841   Transport of Nonsteroidal Anti-Inflammatory Drugs in Mammalian Blood Plasma. J Med Chem.
842   2020;63: 6847–6862. doi:10.1021/acs.jmedchem.0c00225

843 34. Haddad JG, Matsuoka LY, Hollis BW, Hu YZ, Wortsman J. Human plasma transport of vitamin D
844   after its endogenous synthesis. J Clin Invest. 1993;91: 2552–2555. doi:10.1172/JCI116492

845 35. Hall CA. Transcobalamins I and II as natural transport proteins of vitamin B12. J Clin Invest.
846   1975;56: 1125–1131. doi:10.1172/JCI108187

847 36. Kanai M, Raz A, Goodman DS. Retinol-binding protein: the transport protein for vitamin A in
848   human plasma. J Clin Invest. 1968;47: 2025–2044. doi:10.1172/JCI105889

849 37. Kiefer H, Lindstrom J, Lennox ES, Singer SJ. Photo-affinity labeling of specific acetylcholine-
850   binding sites on membranes. Proc Natl Acad Sci U S A. 1970;67: 1688–1694.
851   doi:10.1073/pnas.67.4.1688

852 38. Shinozawa T, Sokabe M, Terada S, Matsusaka H, Yoshizawa T. Detection of cyclic GMP binding
853   protein and ion channel activity in frog rod outer segments. J Biochem. 1987;102: 281–290.
854   doi:10.1093/oxfordjournals.jbchem.a122052

855 39. Klieber MA, Underhill C, Hammond GL, Muller YA. Corticosteroid-binding globulin, a structural
856   basis for steroid transport and proteinase-triggered release. J Biol Chem. 2007;282: 29594–29603.
857   doi:10.1074/jbc.M705014200

858 40. Manoilov KY, Ghosh A, Almo SC, Verkhusha VV. Structural and Functional Characterization of a
859   Biliverdin-Binding Near-Infrared Fluorescent Protein From the Serpin Superfamily. J Mol Biol.
860   2022;434: 167359. doi:10.1016/j.jmb.2021.167359

861 41. Kim S, Woo J, Seo EJ, Yu M, Ryu S. A 2.1 A resolution structure of an uncleaved alpha(1)-
862   antitrypsin shows variability of the reactive center and other loops. J Mol Biol. 2001;306: 109–119.
863   doi:10.1006/jmbi.2000.4357

42. Qi X, Loiseau F, Chan WL, Yan Y, Wei Z, Milroy L-G, et al. Allosteric modulation of hormone
release from thyroxine and corticosteroid-binding globulins. J Biol Chem. 2011;286: 16163–16173.
doi:10.1074/jbc.M110.171082

43. Gupta R, Brunak S. Prediction of glycosylation across the human proteome and the correlation to
protein function. Pac Symp Biocomput. 2002; 310–322. Available:
https://www.ncbi.nlm.nih.gov/pubmed/11928486

44. Chan WL, Carrell RW, Zhou A, Read RJ. How changes in affinity of corticosteroid-binding globulin
modulate free cortisol concentration. J Clin Endocrinol Metab. 2013;98: 3315–3322.
doi:10.1210/jc.2012-4280

45. Pemberton PA, Stein PE, Pepys MB, Potter JM, Carrell RW. Hormone binding globulins undergo
serpin conformational change in inflammation. Nature. 1988;336: 257–258. doi:10.1038/336257a0

46. Gardill BR, Vogl MR, Lin H-Y, Hammond GL, Muller YA. Corticosteroid-binding globulin: structure-
function implications from species differences. PLoS One. 2012;7: e52759.
doi:10.1371/journal.pone.0052759

47. Lewis JG, Bagley CJ, Elder PA, Bachmann AW, Torpy DJ. Plasma free cortisol fraction reflects
levels of functioning corticosteroid-binding globulin. Clin Chim Acta. 2005;359: 189–194.
doi:10.1016/j.cccn.2005.03.044

48. Taboada C, Brunetti AE, Lyra ML, Fitak RR, Faigón Soverna A, Ron SR, et al. Multiple origins of
green coloration in frogs mediated by a novel biliverdin-binding serpin. Proc Natl Acad Sci U S A.
2020;117: 18574–18581. doi:10.1073/pnas.2006771117

49. Robbins J. Thyroxine transport and the free hormone hypothesis. Endocrinology. 1992;131: 546–
547. doi:10.1210/endo.131.2.1639006

50. Siiteri PK, Murai JT, Raymoure WJ, Kuhn RW, Hammond GL, Nisker JA. The Serum Transport of
Steroid Hormones. Proceedings of the 1981 Laurentian Hormone Conference. 1982. pp. 457–510.
doi:10.1016/b978-0-12-571138-8.50016-0

51. Bouillon R, Schuit F, Antonio L, Rastinejad F. Vitamin D Binding Protein: A Historic Overview.
Front Endocrinol . 2019;10: 910. doi:10.3389/fendo.2019.00910

52. Chun RF. New perspectives on the vitamin D binding protein. Cell Biochem Funct. 2012;30: 445–
456. doi:10.1002/cbf.2835

53. Fortunati N. Sex Hormone-Binding Globulin: Not only a transport protein. What news is around the
corner? Journal of Endocrinological Investigation. 1999. pp. 223–234. doi:10.1007/bf03343547

54. Round P, Das S, Wu T-S, Wähälä K, Van Petegem F, Hammond GL. Molecular interactions
between sex hormone–binding globulin and nonsteroidal ligands that enhance androgen activity.
Journal of Biological Chemistry. 2020. pp. 1202–1211. doi:10.1016/s0021-9258(17)49880-2

55. Spence MA, Mortimer MD, Buckle AM, Minh BQ, Jackson CJ. A Comprehensive Phylogenetic
Analysis of the Serpin Superfamily. Mol Biol Evol. 2021;38: 2915–2929.
doi:10.1093/molbev/msab081

56. Garraffo HM, Caceres J, Daly JW, Spande TF, Andriamaharavo NR, Andriantsiferana M. Alkaloids
in Madagascan frogs (Mantella): pumiliotoxins, indolizidines, quinolizidines, and pyrrolizidines. J

Nat Prod. 1993;56: 1016–1038. doi:10.1021/np50097a005

57. Fischer EK, Roland AB, Moskowitz NA, Vidoudez C, Ranaivorazo N, Tapia EE, et al. Mechanisms
of Convergent Egg Provisioning in Poison Frogs. Curr Biol. 2019;29: 4145–4151.e3.
doi:10.1016/j.cub.2019.10.032

58. Clark VC, Raxworthy CJ, Rakotomalala V, Sierwald P, Fisher BL. Convergent evolution of
chemical defense in poison frogs and arthropod prey between Madagascar and the Neotropics.
Proc Natl Acad Sci U S A. 2005;102: 11617–11622. doi:10.1073/pnas.0503502102

59. McGugan JR, Byrd GD, Roland AB, Caty SN, Kabir N, Tapia EE, et al. Ant and Mite Diversity
Drives Toxin Variation in the Little Devil Poison Frog. J Chem Ecol. 2016;42: 537–551.
doi:10.1007/s10886-016-0715-x PMID - 27318689

60. Santos JC, Tarvin RD, O'Connell LA. A Review of Chemical Defense in Poison Frogs
(Dendrobatidae): Ecology, Pharmacokinetics, and Autoresistance. Chemical Signals in Vertebrates
13. 2016. pp. 305–337. doi:10.1007/978-3-319-22026-0_21

61. Fagerberg L, Hallström BM, Oksvold P, Kampf C, Djureinovic D, Odeberg J, et al. Analysis of the
human tissue-specific expression by genome-wide integration of transcriptomics and antibody-
based proteomics. Mol Cell Proteomics. 2014;13: 397–406. doi:10.1074/mcp.M113.035600

62. Merkin J, Russell C, Chen P, Burge CB. Evolutionary dynamics of gene and isoform regulation in
Mammalian tissues. Science. 2012;338: 1593–1599. doi:10.1126/science.1228186

63. Yu Y, Fuscoe JC, Zhao C, Guo C, Jia M, Qing T, et al. A rat RNA-Seq transcriptomic BodyMap
across 11 organs and 4 developmental stages. Nat Commun. 2014;5: 3230.
doi:10.1038/ncomms4230

64. Pipes L, Li S, Bozinoski M, Palermo R, Peng X, Blood P, et al. The non-human primate reference
transcriptome resource (NHPRTR) for comparative functional genomics. Nucleic Acids Res.
2013;41: D906–14. doi:10.1093/nar/gks1268

65. Doweiko JP, Nompleggi DJ. Reviews: The Role of Albumin in Human Physiology and
Pathophysiology, Part III: Albumin and Disease States. Journal of Parenteral and Enteral Nutrition.
1991. pp. 476–483. doi:10.1177/0148607191015004476

66. Merlot AM, Kalinowski DS, Richardson DR. Unraveling the mysteries of serum albumin-more than
just a serum protein. Front Physiol. 2014;5: 299. doi:10.3389/fphys.2014.00299

67. Nonneman D, Rohrer GA, Wise TH, Lunstra DD, Ford JJ. A variant of porcine thyroxine-binding
globulin has reduced affinity for thyroxine and is associated with testis size. Biol Reprod. 2005;72:
214–220. doi:10.1095/biolreprod.104.031922

68. Chen Z, Zakrzewska S, Hajare HS, Alvarez-Buylla A, Abderemane-Ali F, Bogan M, et al. Definition
of a saxitoxin (STX) binding code enables discovery and characterization of the anuran saxiphilin
family. Proc Natl Acad Sci U S A. 2022;119: e2210114119. doi:10.1073/pnas.2210114119

69. Tarvin RD, Santos JC, O'Connell LA, Zakon HH, Cannatella DC. Convergent Substitutions in a
Sodium Channel Suggest Multiple Origins of Toxin Resistance in Poison Frogs. Mol Biol Evol.
2016;33: 1068–1081. doi:10.1093/molbev/msv350 PMID - 26782998

70. Refetoff S. Thyroid Hormone Transport Proteins: Thyroxine-Binding Globulin, Transthyretin, and

942   Albumin. Encyclopedia of Hormones. 2003. pp. 483–490. doi:10.1016/b0-12-341103-3/00287-4

943 71. Hammond GL, Smith CL, Paterson NA, Sibbald WJ. A role for corticosteroid-binding globulin in
944   delivery of cortisol to activated neutrophils. J Clin Endocrinol Metab. 1990;71: 34–39.
945   doi:10.1210/jcem-71-1-34

946 72. Kim JT, Jedrychowski MP, Wei W, Fernandez D, Fischer CR, Banik SM, et al. A Plasma Protein
947   Network Regulates PM20D1 and N-Acyl Amino Acid Bioactivity. Cell Chem Biol. 2020;27: 1130–
948   1139.e4. doi:10.1016/j.chembiol.2020.04.009

949 73. Wei W, Riley NM, Lyu X, Bertozzi CR, Long JZ. Protocol for cell type-specific labeling, enrichment,
950   and proteomic profiling of plasma proteins in mice. STAR Protoc. 2021;2: 101014.
951   doi:10.1016/j.xpro.2021.101014

952 74. Elias JE, Gygi SP. Target-decoy search strategy for increased confidence in large-scale protein
953   identifications by mass spectrometry. Nat Methods. 2007;4: 207–214. doi:10.1038/nmeth1019

954 75. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein
955   structure prediction with AlphaFold. Nature. 2021;596: 583–589. doi:10.1038/s41586-021-03819-2

956 76. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF
957   Chimera--a visualization system for exploratory research and analysis. J Comput Chem. 2004;25:
958   1605–1612. doi:10.1002/jcc.20084

959 77. Trott O, Olson AJ. AutoDock Vina: improving the speed and accuracy of docking with a new
960   scoring function, efficient optimization, and multithreading. J Comput Chem. 2010;31: 455–461.
961   doi:10.1002/jcc.21334

962 78. Eberhardt J, Santos-Martins D, Tillack AF, Forli S. AutoDock Vina 1.2.0: New Docking Methods,
963   Expanded Force Field, and Python Bindings. J Chem Inf Model. 2021;61: 3891–3898.
964   doi:10.1021/acs.jcim.1c00203

965 79. Moskowitz NA, Aurora-Alvarez-Buylla, Morrison CR, Chamba A, Rentería J, Tapia EE, et al.
966   Poison frog diet and chemical defense are influenced by availability and selectivity for ants.
967   doi:10.1101/2022.06.14.495949

968 80. Saporito RA, Donnelly MA, Madden AA, Garraffo HM, Spande TF. Sex-related differences in
969   alkaloid chemical defenses of the dendrobatid frog Oophaga pumilio from Cayo Nancy, Bocas del
970   Toro, Panama. J Nat Prod. 2010;73: 317–321. doi:10.1021/np900702d

971 81. Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, et al. Sharing and community
972   curation of mass spectrometry data with Global Natural Products Social Molecular Networking. Nat
973   Biotechnol. 2016;34: 828–837. doi:10.1038/nbt.3597

974 82. Payne C, Bovio R, Powell DL, Gunn TR, Banerjee SM, Grant V, et al. Genomic insights into
975   variation in thermotolerance between hybridizing swordtail fishes. Mol Ecol. 2022.
976   doi:10.1111/mec.16489

977 83. Krueger F, James F, Ewels P, Afyounian E, Schuster-Boeckler BF. TrimGalore: v0. 6.7-DOI via
978   Zenodo. Zenodo; 2021.

979 84. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
980   EMBnet.journal. 2011. p. 10. doi:10.14806/ej.17.1.200

981   85. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nat
982   Biotechnol. 2016;34: 525–527. doi:10.1038/nbt.3519

983