Higher-order dissimilarity in biodiversity:

Identifying dissimilarities of spatial or temporal dissimilarity structures

Ryosuke Nakadai[1,2,3], Keita Fukasawa[1], Taku Kadoya[1,4], Fumiko Ishihama[1]

[1] Biodiversity division, National Institute for Environmental Studies, Onogawa 16-2, Tsukuba,

Ibaraki 305-8506, Japan

[2] Faculty of Environment and Information Sciences, Yokohama National University,

Tokiwadai 79-7, Hodogaya-ku, Yokohama, Kanagawa 240-8501, Japan

[3] Institute for Multidisciplinary Sciences, Yokohama National University, Tokiwadai 79-5,

Hodogaya-ku, Yokohama, Kanagawa 240-8501, Japan

[4] Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba,

Ibaraki 305-8571, Japan

* Correspondence: Ryosuke Nakadai

r.nakadai66@gmail.com; nakadai-ryosuke-pt@ynu.ac.jp

Running title: Higher-order dissimilarity

ORCID

Ryosuke Nakadai 0000-0002-9512-8511

Keita Fukasawa 0000-0002-9563-457X

Taku Kadoya 0000-0002-7816-5484

Fumiko Ishihama 0000-0001-8515-5914

Acknowledgements

We thank Dr. Satoshi Aoki for his advice on choosing a coalescent simulator and Prof. Pierre

Legendre for his insightful comments on the manuscript, drawing from his profound and

expert knowledge.

Data availability

34 Simulated data and codes will be uploaded after acceptance of the manuscript to Figshare.

Conflict of interest

37 The authors declare no conflict of interests.

Author contributions

40 RN, FI, KF, and TK designed the study; RN conceived the idea of higher-order dissimilarity,

41 analysed the datasets, and wrote the first draft of the manuscript, with significant inputs from

42 KF, FI, and TK; KF and RN developed the methodology to quantitatively evaluate the idea

43 introduced in the present study; and FI conceived the basic concept of the methodology. All

44 authors have contributed substantially to the final version of this manuscript.

Abstract

1. Elucidating biodiversity patterns and their background processes is critical in biodiversity

science. Dissimilarity, which is calculated based on multivariate biological quantities, is a

major component of biodiversity. As spatial and temporal biodiversity information

availability increases, the scope of dissimilarity studies has been expanded to cover

various levels and types of spatio-temporal biodiversity facets (e.g., gene, community,

and ecosystem function), and diverse pairwise dissimilarity indices have been developed.

However, further development of the dissimilarity concept is required in comparative

studies on spatio-temporal structures of biodiversity compositional patterns, such as those

exploring commonalities of biogeographical boundaries among taxa, compared to the

conventional ones to consider higher dimensions of dissimilarity: dissimilarity of

dissimilarity structures.

2. This study proposes a novel and general concept, high-order dissimilarity (HOD), for

quantitatively evaluating the dissimilarities of spatial or temporal dissimilarity structures

among different datasets, proposes specific implementations of HOD as operational

indices, and illustrates the potential resolution of scientific and practical questions

through HOD.

3. We further demonstrate the advantages of the HOD concept by applying it to actual

patterns, such as long-term and/or large-spatial hypothetical monitoring datasets.

4. Our conceptual framework on HOD extends the existing framework of biodiversity

science and is versatile, with many potential applications in acquiring more valuable

information from ever-increasing biodiversity data.

Keywords:

autocorrelation, beta diversity, dissimilarity matrix, genetic differentiation, multivariate

analysis, spatio-temporal structure

Introduction

Nature's patterns are ubiquitous regardless of the level of biological organisation, and the elucidation of their determinants has been a long-standing and fundamental issue in biodiversity science. Dissimilarity is a key element of such patterns and has been studied in terms of spatial and temporal differences and changes in various biological entities, mainly at the same level of biological organisation (Anderson et al. 2011). This encompasses differences and changes in genetic alleles, species composition, interaction networks, and ecosystem functions (Whittaker 1960, 1972; Raymond & Rousset 1995; Mori et al. 2015; Poisot et al. 2012). The scope of dissimilarity is continuously expanding, covering various levels and types of biodiversity facets, which has stimulated the development of conceptual and analytical frameworks for the analysis of the determinants of dissimilarity patterns using various dissimilarity indices (Koleff et al. 2003; Lozupone & Knight 2005; Baselga 2010; Rocchini et al. 2018; Mammola & Cardoso 2020). Notably, despite the varieties, all these indices have a standard form in that one dissimilarity value is calculated from two vectors of multivariate biological quantities (i.e., pairwise dissimilarity indices), and the calculation of dissimilarity values for all pairs of vectors yields a dissimilarity matrix (Anderson et al. 2011).

Some of the spatio-temporal structures of biodiversity composition patterns, however, are beyond the scope of pairwise dissimilarity and identifying, quantifying, and understanding them necessitates higher-order considerations. In a spatial context, higher-order considerations are useful for the analysis of categorically different dissimilarity matrices, such as among assemblages of different taxonomic groups and among species (i.e., comparative biogeography and phylogeography). For example, biogeographical researchers have attempted to identify common biogeographical borders among taxa (Wallace 1894; Holt et al. 2013; Whittaker et al. 2013; Komaki 2021); similarities or dissimilarities in spatial composition dissimilarities among multiple taxa need to be evaluated. In the temporal context, higher-order consideration would provide a framework for quantitatively evaluating dissimilarity in compositional changes (i.e., temporal asynchrony of compositional changes), such as among sites and different trophic groups. For example, temporal changes in community composition (i.e., temporal beta diversity: Legendre 2019; Nakadai 2022) have

recently been expanded to multiple sites at large spatial scales (Magurran et al. 2019;

Nakadai 2020; Gotelli et al. 2022), thus necessitating quantitative evaluation of inter-site

dissimilarity between temporal dynamics of community composition measured using

pairwise comparison. Furthermore, loss and biotic homogenisation in species composition

and intraspecific haplotypes have been studied in-depth since identifying human-induced

impacts on biodiversity within the context of conservation (Olden 2006; Valtonen et al. 2017).

Moreover, quantifying the temporal change of spatial dissimilarity structure from a baseline

at a time point is essential to assessing the impacts. Despite the apparent importance and

scientific need for higher-order consideration in dissimilarities, general concepts and formal

statistical methods for quantitative comparisons across conventional pairwise dissimilarities

have not been fully developed.

To address this limitation, we introduce a general framework for considering

dissimilarities at a higher order (hereafter referred to as higher-order dissimilarity; HOD)

than that of conventional pairwise dissimilarities. HOD is a natural extension of the pairwise

dissimilarity indices and considers differences between dissimilarity matrices. In addition, it

can be a novel building block in the studies of the dissimilarities of spatio-temporal structures

of biodiversity compositional patterns. First, in the present study, we formally define the

novel and general concept of HOD to quantitatively evaluate dissimilarities of spatial or

temporal dissimilarity structures among different datasets (Fig. 1a). Second, we designed the

concept of HOD and proposed its implementation as a general statistical method. Third, we

demonstrated the advantages of the HOD concept by applying it to actual patterns, such as

long-term and/or large-spatial hypothetical monitoring datasets. Finally, we discuss the

potential impacts of the concept of "higher-order dissimilarity" and the developed analytical

framework on a wide range of research fields, including understanding complex spatio-

temporal structures in biodiversity and future methodological challenges.

Figure 1

Figure 1 Schematic of higher-order dissimilarity (HOD). A spatial case of HOD is shown in

(a). Although conventional dissimilarity has focused on dissimilarity among the same

dimension, higher-order dissimilarity can compare dissimilarity structures without regard to

whether they are of the same dimension. For visualisation, only three compositional types are

hypothesised, corresponding to three levels of darkness in each colour (e.g., blue, green, and

red). Specifically, we hypothesised that dissimilarities between sites (or time points) of the

same level of darkness were zero (i.e., identical composition), those between sites of one

darker or lighter difference were 0.5, and those between the darkest and lightest sites were

1.0. The grey arrows indicate conventional dissimilarity (i.e., first-order dissimilarity) and the

black arrows indicate HOD. Among compared datasets, black and white stars indicate

identical sites or time points. The potential analytical framework for HOD based on spatial

and temporal comparisons (i.e., spatial and temporal HOD) is shown in (b). The upper panel

shows examples of spatial HOD, and the lower panel shows examples of temporal HOD. In

both cases, the left-hand side shows examples of community structure comparisons between

different guilds and intraspecific genetic structure between species, while the right-hand side

shows examples of changes in spatial HOD or temporal HOD along a temporal or spatial axis.

Higher-order dissimilarity

In biodiversity science, one dissimilarity value is generally calculated from two vectors of

multivariate biological quantities (e.g., community composition or allele frequency).

Specifically, a community dissimilarity index (e.g., Bray–Curtis dissimilarity; Odum 1950)

and a genetic distance index (e.g., F_{st}; Wright 1969) were calculated based on vectors of

species abundance in two target communities and vectors of genetic allele frequency in two

target populations, respectively. In addition, multiple vectors are summarised into a matrix

called a site-species (or time-species) matrix (e.g., compositional vectors for sites or time

points). The calculation of dissimilarity values for all pairs of vectors yields a dissimilarity

matrix, which is a conventional pairwise dissimilarity; we call this first-order dissimilarity

(Fig. 1a). Here, we extend the first-order dissimilarity to formally define HOD as

dissimilarities of spatial or temporal dissimilarity structures (Fig. 1a). We also define 'order'

as nested times for calculating dissimilarity, e.g., conventional pairwise dissimilarity is the

first order; and dissimilarity of dissimilarity matrices the second order. By definition, the

concept can be extended to a higher order where appropriate, such as to the third order (i.e.,

dissimilarity of second-order dissimilarity matrices).

The concept of HOD is a generalisation of the traditional analytical approach to

identify species groups sharing the same trend and boundary in space and time (Holt et al.

2013; Gotelli et al. 2022). In addition, HOD targets the spatial structure, which has been the

primary subject of biogeography and temporal structure. Specifically, to spatially and

temporally distinguish HODs, we use the terms' spatial HOD' and 'temporal HOD' (Fig. 1b).

Given the recent rapid growth of spatio-temporal monitoring data, HODs regarding temporal

variation in spatial structure (i.e. temporal change in spatial HOD, Fig. 1b) or spatial

variation in temporal structure (i.e. spatial change in temporal HOD, Fig. 1b) will be

applicable in future analyses. The subsequent sections mainly focus on second-order

dissimilarities similar to spatial and temporal HODs to elucidate the concept.

Implementation of the higher-order dissimilarity

The most basic calculation of the HOD index (Fig. 2) is based on the RV coefficient

(Escoufier 1973; Robert & Escoufier 1976), which is the multivariate generalisation of the

squared Pearson correlation that correlates two matrices with corresponding sites (Blanchet et

al. 2014). Before comparing two multivariate data matrices (i.e., conventional dissimilarity

matrices), one may employ a dissimilarity index as a transformation technique for each data

matrix. Specifically, principal coordinate analysis (PCoA) should be applied to each

conventional dissimilarity matrix. This process yields rectangular data matrices, ensuring

each dissimilarity matrix adheres to Euclidean properties. Then, the RV coefficient can be

calculated through a comparison of two PCoA matrices (i.e., X and X' in Fig. 2). The RV

coefficient yields values ranging from 0 (indicating no correlation) to 1 (representing perfect

correlation). Therefore, we compute the HOD index as one minus the RV coefficient to

transform it into a dissimilarity measure, which we refer to as the contrast RV coefficient.

Figure 2

Figure 2 Schematic representation of specific analytical procedures for higher-order

dissimilarity (HOD). For visualisation, only three compositional types are hypothesised,

corresponding to three levels of darkness in each colour (e.g., blue, green, and red).

Specifically, we hypothesised that dissimilarities between sites (or time points) of the same

level of darkness were zero (i.e., identical composition), between sites of one darker or

lighter difference were 0.5, and between darkest and lightest sites were 1.0. The asterisk

notes the necessity of the Kriging process if the specific sampling points are not identical.

203   When sampling points comprising two biodiversity datasets are obtained at the same

spatial or temporal location and correspond precisely one-to-one, their dissimilarity matrix

elements also correspond. However, this is not always true. For example, when comparing

two phylogeographic patterns, the sample points of each species are usually in different

locations. In such a case, the elements of dissimilarity matrices do not correspond one-to-one

and temporal or spatial proximity information is required for comparison. To calculate the

HOD index for such an 'irregular dataset' (i.e., a dataset where the two compositional vectors

to be compared were observed at different spatial and temporal points and did not correspond

on a one-to-one basis), kriging, which is an interpolation method based on variograms

(Wagner 2004; Legendre & Legendre 2012; Borcard et al. 2023), helps estimate the values of

one data set at the locations of the other before the calculation. After interpolating the

missing values, the methods for calculating the HOD index (i.e., contrast RV coefficient)

remain the same as described above. However, spatio-temporal interpolation by kriging is not

always effective and must be used cautiously when there is either no spatial/temporal

autocorrelation or weak autocorrelation, or when there are disconnected patterns. One

possible approach, based on spatial or temporal weight matrices such as Moran's I

(Wartenberg 1985; Lee 2001; Eckardt and Mateu 2021), could be effective except in the

absence of complete spatial or temporal autocorrelation.

To check the properties of the HOD indices, we summarised the specific values of spatial

HOD for pairs of simple sample cases in Table 1. While the degrees of spatial (or temporal)

autocorrelation are not directly relevant to this issue in the case of complete datasets, they are

included as reference information. When interpolating using kriging, the degree of

autocorrelation would potentially affect the accuracy of the calculation. Here, spatial

autocorrelation was obtained through regression analysis of distance-based RDA to the

PCNM axes of spatial distances (Legendre et al. 2015). To calculate both the RV coefficient

and the p-values using the function "coeffRV" of the *FactorMineR* package (Lê et al. 2008)

in R (R Core Team, 2022).

Table 1 Simple sample cases to show properties of HOD indices. Patterns X and Y represent

pairs for calculating HODs. *Adjusted R²* (X) and (Y) indicate the degree of spatial

autocorrelation for each pattern X and Y, obtained through regression analysis of distance-

based RDA to the PCNM axes of spatial distances. The HOD indices calculated as contrast

RV coefficient (i.e., 1 – RV coefficient) are shown. For visualisation purposes, only three

compositional types are hypothesised, corresponding to three levels of darkness in each

colour (e.g., blue and green). Specifically, we hypothesised that dissimilarities between sites

of the same level of darkness were zero (i.e., identical composition), between sites of one

darker or lighter difference were 0.5, and between the darkest and lightest sites were 1.0.

Association	Pattern		Regression-PCNM (Space)		HOD	
	X	Y	*Adjusted R²* (X)	*Adjusted R²* (Y)	Contrast RV coefficient	*P*-value
A-A			0.8242	0.8242	0.0000	0.0000
B-B			0.7501	0.7501	0.0000	0.0000
C-C			–0.1388	–0.1388	0.0000	0.0000
A-B			0.8242	0.7501	0.7223	0.0010
B-C			0.7501	–0.1388	0.9892	0.5542
C-A			–0.1388	0.8242	0.9317	0.1177
A-A'			0.8242	0.8242	0.7776	0.0034
B-B'			0.7501	0.7501	0.9317	0.1177
C-C'			–0.1388	–0.1388	1.0000	0.9971
A-C'			0.8242	–0.1388	0.9971	0.7369

Resolvable questions through the higher-order dissimilarity approaches

Theoretically, HOD concepts apply to any dissimilarity dataset if the relative position (i.e.,

sites or time points) can be determined based on the absolute distance between pairs of

datasets. For example, the monitoring of the genetic structure is one of the most urgent

targets of biodiversity monitoring (Hoban et al. 2021; O'Brien et al. 2022), although no

effective monitoring framework has been developed. In this context, the HOD provides an

ideal framework for the quantitative evaluation of genetic structure temporally (Fig. 3). As

frequently probable examples, we show two hypothetical scenarios of genetic structure

changes: one is a stable scenario (orange line and dots), and the other is a genetically

homogenised scenario (blue line and dots) in Fig. 3 (see Supplementary Text 1 for detailed

information about construction for visualisation). Fig. 3(a) indicates temporal changes in the

values of the HOD index (i.e., contrast RV coefficient) to time 0 (starting point), and the

lower Fig. 3(b) represents mapped overviews of changes in genetic structure over time, from

which we calculated the HOD values. The scarcity of such spatially coordinated data on

genetic structures over a wide area will soon be addressed, particularly following the

development of environmental DNA technology (Tsuji et al. 2023), and our framework will

be an essential contribution to genetic diversity monitoring.

Figure 3

Figure 3 Sample application of higher-order dissimilarity (HOD) to genetic monitoring using

two simulated datasets based on two types of scenarios from time 0 to time 100. In both

scenarios, the stable conditions are hypothesised by time 30 (shared section in light green).

After time 30, the genetic structure is stable in one scenario (upper section in orange) but

homogenised in the other (lower section in blue). The change of the calculated HOD index

(i.e., contrast RV coefficient) compared to the value at time 0 is shown in (a). The genetic

structure is visualised in (b), and the colours were determined using the two axes of NMDS

(details see Supplementary text 1).

To show the advantages of HOD more comprehensively, we summarised the potential

applications of HOD in Table 2, including five types of datasets: community composition,

genetic frequency, ecosystem function, interaction network, and phylogeny. Most cases are

straightforward extensions of those shown in Fig. 1b. One particular case is that HOD

approaches are applicable to phylogenetic patterns despite the difficulty in extracting single-

dimensional values, for example, the output of non-metric multidimensional scaling based on

multiple traits. HOD approaches can be applied to various community ecology, biogeography,

phylogeography, and macroevolution datasets. Particularly in the case of examples covering

macroevolution (i.e., phylogenetic structures), a previous study had a similar, though not

identical, perspective (Duarte 2011). This fact further emphasises the importance of

developing a field of study for HOD.

Table 2 Potential applications of the higher-order dissimilarity (HOD) approach

Target axis of HOD	Targets of comparison	Question
(a) community composition		
Time	Space	How do temporal changes in community composition vary in space?
		Do closer sites show more similar community compositional change across time?
Time	Taxa	Which taxa share similar changes in community composition in time?
Space	Time	How does spatial composition change over time?
		When did a large change in community compositional structure happen?
Space	Taxa	Which taxa share a similar community composition structure in space?
(b) population genetics		
Time	Space	How does temporal genetic frequency change in space?
		Do closer sites show more similar changes in genetic frequency across time?
Time	Taxa	Which taxa share similar changes in genetic frequency over time?
Space	Time	How does the spatial genetic structure change over time?
		When did a large change in genetic structure occur?
Space	Taxa	Which taxa share similar changes in genetic frequency in space?
(c) ecosystem function		
Time	Space	How does temporal ecosystem function change in space?
		Do closer sites show more similar changes in ecosystem function across time?
Space	Time	How does spatial genetic structure change over time?
		When did a large change in the genetic structure occur?

(d) interaction network

Time	Space	How does the temporal network structure change in space? Do closer sites show more similar changes in network structure across time?
Space	Time	How does spatial network structure change in time? When did large change in network structure happen?

(e) macroevolution

Phylogeny	Trait	Which traits share similar evolutionary changes across macroevolutionary history?

The main focus has been on comparisons between dissimilarity matrices with similar

properties, but this is not necessarily true. Examples include comparisons among

communities of different trophic levels and comparisons between community structures and

genetic structures in community members. In traditional approaches, revealing the relative

importance of the driving processes of local communities, such as abiotic vs. biotic factors

and/or top-down vs. bottom-up processes as drivers of community structures, has been a

major concern (Smith et al. 2010; HilleRisLambers et al. 2012). The HOD approach is a

different perspective that directly evaluates the similarity of patterns without explicitly

considering the driving processes. Low values of HOD among data from different trophic

levels of communities and different hierarchies of genetic and community data allowed us to

generate novel hypotheses with more direct evidence, as the processes shaping them are

considered common among the compared targets (taxonomic differences; Fig. 1b).

Conversely, high HOD values suggest that the process of constructing the overall system is

complex and contingent. From an application perspective, high values of spatial HOD also

suggest that achieving conservation through establishing protected areas, setting focal species

for conservation, or implementing a single conservation policy would be difficult, as the

driving processes differ among taxa and levels of biological entities.

Future challenges

In the HOD approaches proposed here, we cannot distinguish between parts of continuous

patterns and, indeed, disjunctive patterns in dissimilarities, for example, either simple

distance-decay patterns or patterns due to isolation by geographic barriers. Therefore, where

the commonality of spatial disconnections is the main concern, combining HOD approaches

with existing approaches would be more effective in identifying disjunct barriers (e.g., Soltis

et al. 2006). In temporal structures, approaches separating trend components have been used

(Cowpertwait & Metcalife 2009), and potentially combined with such methods, it may be

possible to evaluate continuous and disjunctive dissimilarities separately. Furthermore,

spatially- or temporally-constrained clustering approaches (Guénard & Legendre 2022) based

on the HOD matrix can depict disjunctive patterns of higher-order structures in community

compositions. However, further testing is required to clarify how much complex hierarchy is

allowed for a sufficiently accurate evaluation for calculating the HOD index.

Although we mainly focused on the second-order dissimilarities in the present study, the

applicability of third- and higher-order dissimilarities would increase when the amount and

dimension of biodiversity information continue to grow. To illustrate the benefit of HOD, we

showed temporal changes in genetic structure (Fig. 3). The figure shows a pattern in a single

species and, thus, second-order dissimilarity (i.e., space × time). If this dataset is available for

multiple species, the similarity and dissimilarity of temporal changes in spatial structure in

population genetics can be quantitatively evaluated among species using third-order

dissimilarities (i.e., space×time×species). In such multidimensional situations, a hierarchical

multiple-factor analysis (Le Dien & Pagès 2003) can be a useful exploratory tool for grasping

the structure of communities that affect the higher-order HOD. This simple example tells us

that the breadth of areas that the HOD concept can adopt is rapidly expanding, aligning with

the forthcoming explosive increase in available dissimilarity information.

Conclusions

The rapid increase in the availability of spatial and temporal biodiversity information

necessitates comparative studies on spatio-temporal structures of biodiversity compositional

patterns, which require further development of the dissimilarity concept over conventional

ones to consider higher dimensions of dissimilarity: dissimilarity of dissimilarity structures.

Our study introduced the concept of higher-order dissimilarity (HOD), which can account for

dissimilarities of spatial or temporal dissimilarity structures. For example, this framework can

be applied to various types of broad-scale biodiversity monitoring datasets and enables the

evaluation of temporal changes in the spatial HOD to a baseline. Furthermore, the concept is

applicable even if the compared datasets have entirely different origins (e.g., different

taxonomic groups and hierarchies of biological organisation, Fig. 1b) as long as they are

summarised into dissimilarity information at their level. We mainly focused on the

biodiversity dataset in the present study; the concept of HOD applies to all kinds of

dissimilarity matrices, not just those based on biodiversity. For example, various

contemporary issues arise in the social-ecological system, and this HOD will be useful in

examining the relationships between different layers, such as the economy, culture, and

biodiversity. Therefore, the HOD concept and related approaches pave the way for novel

dimensional approaches that deal with large amounts of dissimilarity information.

References

Anderson, M.J., Crist, T.O., Chase, J.M., Vellend, M., Inouye, B.D., Freestone, A.L. et al. (2011) Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. *Ecology Letters,* 14, 19–28.

Baselga, A. (2010) Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography,* 19, 134–143.

Borcard, D., Gillet, F., & Legendre, P. (2023) Numerical ecology with R (R による数値□態学), Japanese edition. Kyoritsu Shuppan Co. Ltd, Tokyo.

Cowpertwait, P.S., & Metcalfe, A.V. (2009) Introductory time series with R. Springer Science & Business Media.

Duarte, L.S. (2011) Phylogenetic habitat filtering influences forest nucleation in grasslands. Oikos, 120, 208–215.

Gotelli, N.J., Moyes, F., Antão, L.H., Blowes, S.A., Dornelas, M., McGill, B.J. et al. (2022) Long□term changes in temperate marine fish assemblages are driven by a small subset of species. *Global Change Biology,* 28, 46–53.

Guénard, G., & Legendre, P. (2022) Hierarchical clustering with contiguity constraint in R. *Journal of Statistical Software*, 103, 1–26.

Eckardt, M., & Mateu, J. (2021) Partial and semi□partial statistics of spatial associations for multivariate areal data. *Geographical. Analysis*, 53, 818–835.

HilleRisLambers, J., Adler, P.B., Harpole, W.S., Levine, J.M., & Mayfield, M.M. (2012)

Rethinking community assembly through the lens of coexistence theory. *Annual Review*

of Ecology, Evolution and Systematics, 43, 2012.

Hoban, S., Campbell, C.D., da Silva, J. M., Ekblom, R., Funk, W.C., Garner, B.A., et al.

(2021) Genetic diversity is considered important but interpreted narrowly in country

reports to the Convention on Biological Diversity: Current actions and indicators are

insufficient. *Biological Conservation*, 261, 109233.

Holt, B.G., Lessard, J.P., Borregaard, M.K., Fritz, S.A., Araújo, M.B., Dimitrov, D. (2013)

An update of Wallace's zoogeographic regions of the world. *Science,* 339, 74–78.

Koleff, P., Gaston, K.J., & Lennon, J.J. (2003) Measuring beta diversity for presence–

absence data. *Journal of Animal Ecology,* 72, 367–382.

Komaki, S. (2021) Widespread misperception about a major East Asian biogeographic

boundary exposed through bibliographic survey and biogeographic meta‐analysis.

Journal of Biogeography 48, 2375–2386.

Le Dien, S., & Pagès, J. (2003) Hierarchical multiple factor analysis: application to the

comparison of sensory profiles. *Food Quality and Preference*, 14, 397–403.

Lee, S. I. (2001) Developing a bivariate spatial association measure: an integration of

Pearson's r and Moran's I. *Journal of Geographical Systems,* 3, 369–385.

Legendre, P. (2019) A temporal beta-diversity index to identify sites that have changed in

exceptional ways in space–time surveys. *Ecology and Evolution,* 9, 3500–3514

Legendre, P., Fortin, M. J., & Borcard, D. (2015) Should the Mantel test be used in spatial

analysis? *Methods in Ecology and Evolution*, 6, 1239–1247.

Legendre, P. & Legendre, L. (2012) Numerical ecology, 3rd edn. *Elsevier*. Amsterdam

Lozupone, C., & Knight, R. (2005) UniFrac: a new phylogenetic method for comparing

microbial communities. *Applied and Environmental Microbiology,* 71, 8228–8235.

Magurran, A.E., Dornelas, M., Moyes, F., & Henderson, P.A. (2019) Temporal β diversity—

A macroecological perspective. *Global Ecology and Biogeography,* 28, 1949–1960.

Mammola, S., & Cardoso, P. (2020) Functional diversity metrics using kernel density

n‐dimensional hypervolumes. *Methods in Ecology and Evolution,* 11, 986–995.

Mori, A. S., Isbell, F., Fujii, S., Makoto, K., Matsuoka, S., & Osono, T. (2016) Low

multifunctional redundancy of soil fungal diversity at multiple scales. *Ecology Letters*,

19, 249–259.

Nakadai, R. (2020) Degrees of compositional shift in tree communities vary along a gradient

of temperature change rates over one decade: Application of an individual‐based

temporal beta‐diversity concept. *Ecology and Evolution* 10, 13613–13623.

Nakadai, R. (2022) Development of novel temporal beta-diversity indices for assessing

community compositional shifts accounting for changes in the properties of individuals.

Ecological Indicators, 144, 109427.

O'Brien, D., Laikre, L., Hoban, S., Bruford, M.W., Ekblom, R., Fischer, M.C. et al. (2022)

Bringing together approaches to reporting on within species genetic diversity. *Journal of*

Applied Ecology, 59, 2227–2233.

Odum, E.P. (1950) Bird populations of the Highlands (North Carolina) Plateau in relation to

plant succession and avian invasion. *Ecology* 31, 587–605.

Olden, J.D. (2006). Biotic homogenization: a new research agenda for conservation

biogeography. *Journal of Biogeography.* 33, 2027–2039.

Poisot, T., Canard, E., Mouillot, D., Mouquet, N., & Gravel, D. (2012) The dissimilarity of

species interaction networks. *Ecology Letters*, 15, 1353–1361.

Raymond, M., & Rousset, F. (1995). An exact test for population differentiation. *Evolution*

49, 1280–1283.

Rocchini, D., Luque, S., Pettorelli, N., Bastin, L., Doktor, D., Faedi, N. et al. (2018)

Measuring β‐diversity by remote sensing: A challenge for biodiversity monitoring.

Methods in Ecology and Evolution, 9, 1787–1798.

Smith, J.E., Hunter, C.L., & Smith, C. M. (2010) The effects of top–down versus bottom–up

control on benthic coral reef community structure. *Oecologia,* 163, 497–507.

Soltis, D.E., Morris, A.B., McLachlan, J.S., Manos, P.S., & Soltis, P.S. (2006) Comparative

phylogeography of unglaciated eastern North America. *Molecular Ecology,* 15, 4261–

4293.

Tsuji, S., Shibata, N., Inui, R., Nakao, R., Akamatsu, Y., & Watanabe, K. (2023)

Environmental DNA phylogeography: Successful reconstruction of phylogeographic

patterns of multiple fish species from cups of water. *Molecular Ecology Resources*, 23,

1050–1065.

Valtonen, A., Hirka, A., Szőcs, L., Ayres, M. P., Roininen, H., & Csóka, G. (2017).

Long☐term species loss and homogenization of moth communities in Central Europe.

Journal of Animal Ecology, 86, 730–738.

Wartenberg, D. (1985) Multivariate spatial correlation: a method for exploratory

geographical analysis. *Geographical Analysis,* 17, 263–283.

Wallace, A.R. (1894) What are zoological regions? *Nature,* 49, 610–613.

Whittaker R.H. (1960) Vegetation of the Siskiyou mountains, Oregon and California.

Ecological Monographs 30, 279–338

Whittaker R.H. (1972) Evolution and measurement of species diversity. *Taxon,* 21, 213–251

Whittaker, R.J., Riddle, B.R., Hawkins, B.A., & Ladle, RJ (2013) The geographical

distribution of life and the problem of regionalization: 100 years after Alfred Russel

Wallace. *Journal of Biogeography,* 40, 2209–2214.

Wright, S. (1969) Genetics and the evolution of populations. Vol. 2: The theory of gene

frequencies.

Appendices

Figure S1 Hexagon grids used in simulation.

Supplementary text 1 Detailed information on simulated genetic structure across four major

islands in Japan.

(a)

Space

First-order dissimilarity
(Conventional dissimilarity)

Dissimilarity between
a pair of
dissimilarity structures

Higher-order dissimilarity (HOD)

	⬡	⬡	⬡
⬡	0		
⬡	0.5	0	
⬡	1.0	0.5	0

First-order dissimilarities
among hexagon-grids
within each color

(b)

Space

Spatial HOD among species group or taxa

Example cases
Communites   Genes

Top predators   *Species 1*

Primary consumers   *Species 2*

Primary producers   *Species 3*

Temporal changes of spatial HOD

Time 1   *Time 2*   *Time 3*

Time

Temporal HOD among species group or taxa

Example cases
Communites   Genes

Top predators   *Species 1*

Primary consumers   *Species 2*

Primary producers   *Species 3*

Spatial changes of temporal HOD

Site 1

Site 2

Site 3

Spatial information

Euclidian distance

Dissimilarity matrix 1

Dissimilarity matrix 2

Spatial distance matrix

PCoA

PCoA

Kriging*

1 − RV coefficient

*If the sampling points were mismatched, kriging compliment the incomplete information.

HOD index