A new means of energy supply driven by terahertz photons recovers related neural activity

Xiaoxuan Tan[1,2,*]

[1] Astronaut Center of China, Beijing 100084, China.

[2] Innovation Laboratory of Terahertz Biophysics, National Innovation Institute of Defense Technology, Beijing 100071, China.

*Corresponding author.

Email: t15822082707@163.com (X.T.).

Abstract

Continuous and efficient energy capture represents a long-sought dream of mankind. The brain is a major energy consuming organ, an adult brain accounts for about 2% of the body weight but consumes about 20% of the body's energy. However, it is still unclear how the brain achieve efficient use of energy. Here, using nerve cells as test subjects, we found that THz photons with a specific frequency can effectively restore the reduced frequency of action potentials caused by inadequate adenosine 5'-triphosphate (ATP) supply, which has been demonstrated as a novel mode of energy supply, present photons emission at a particular frequency from the breaking of the ATP phosphate bond. This energy supply mechanism may play a key biophysical basis for explaining how the body efficiently obtain energy, because the quantized chemical reactions could have a high energy efficiency and ultrahigh selectivity compared with the traditional thermochemistry and photochemistry.

Introduction

As a universal energy-storing molecule, adenosine 5'-triphosphate (ATP) is used by virtually all living organisms, and plays an important role in both physiological and pathophysiological processes[1-3]. In addition, ATP also regulates many brain functions such as synaptic transmission[4], glutamate release[5], pain sensation[6] and have complex interactions with other signaling such as reactive oxygen species (ROS) or calcium[7-9]. Despite ATP is important for both health and disease, the detailed mechanisms underlying the efficient energy supply of ATP are poorly understood, especially *in vivo*.

It has become increasingly popular for people to take some foods or drugs as functional beverage that are expected to obtain energy to achieve the purpose of maintaining the efficient work of the human body for a long time. However, functional energy foods have very limited energy supply, and stimulants are banned in most cases because of their side effects[10,11]. Early observations have proposed that human high intelligence is involved

in spectral redshift of biophotonic activities in the brain because biophoton spectral redshift

could be a more economical and effective measure of biophotonic signal communications

and information processing in the brain[12,13], The improvement in cognitive function is

driven by the need for more effective communication that requires less energy, but, so far,

there is no direct experimental evidence for the role of photons in brain information

processing and energy supply.

A recent study proposed the concept for the quantized chemical reaction resonantly

driven by multiple FIR/MIR photons[14-16]. Theoretic studies revealed that THz photons

released by ATP phosphate bond breakage could drive life activities. Nevertheless, the

quantum mechanisms underlying the THz bioeffects remain elusive, and the experimental

evidences are still lacking. In this work, we employed nerve cells as test object to study the

effects of THz photons on energy supply. We surprisingly found that without introducing

any exogeneous substances and implanting electrodes, THz photons could effectively

recover reduced action potential emission frequency due to inadequate ATP supply, and

we ruled out the possibility of restoring ATP production for energy supply (Fig.1A). The

phenomenon can clearly demonstrate that THz photons can regulate brain information

processing and energy metabolism directly.

Results and discussion

Photons with a specific frequency can participate in multiple life activities such as ion

channel dynamics and DNA unwinding through changing vibration levels of a confined

molecule excited (Fig.1B)[17-20]. Previous studies have proposed that the principle of

terahertz wave energy supply is the release of 34 ± 2 THz photons by phosphate bond

break, then photons acts as a massless reagent to supply energy[15,16]. We therefore

calculated whether the IR vibration spectra of ATP have strong formants at 34 ± 2 THz by

density functional theory (DFT). As Fig.1C shown, the ATP shows significant THz

absorption in this range. In particular, ATP supplies energy by breaking high-energy

phosphate bonds and the THz frequency (34 ± 2 THz) indeed could resonate with the

vibration of phosphate bond. These data agree with the previous results and suggest that

34±2 THz may well be the direct energy source for life activities such as biosynthesis.

Next, we designed and fabricated terahertz lasers that can be used to irradiate biological

tissues. The λ ~ 8.6 μm quantum cascade laser (QCL) wafer was grown on an n-doped

(Si, 2×10^{17} cm^{-3}) InP substrate by solid source molecular beam epitaxy (MBE) based on

a slightly-diagonal bound-to-continuum design, as is identical to that in Ref. 21 (for detailed

protocol, see Supplemental Methods). As Fig.2 shown, the power of QCL has good

portability and easy integration. The laser output by QCL is coupled into the fiber by the

coupler for collimation. The fiber (PIR600/700-100-FC/PC-FT-SP30) with a core diameter

of 200 μm and a numerical aperture (NA) of 0.35±0.05 we used is flexible, so there is good

flexibility to change the stimulation target. We delivered the THz laser with the following

parameters: average irradiation power 30 ± 2 mW (measurements were made before and

after each experiment to ensure stable output)，pulse width 2 us, and repetition rate 200

kHz (for detailed protocol, see Supplemental Methods).

To determine whether 34 ± 2 THz photons have the function of energy supply, we first

asked whether 34 ± 2 THz photons could restore the diminished life activities caused by

insufficient energy supply. To answer this question, we took neurons as the test object

(layer-5 pyramidal cells in acute slices of mouse prefrontal cortex) and performed patch-

clamp experiment to quantify the firing frequency of neurons' action potentials. We first

inhibited mitochondrial ATP generation and electron transfer pathways by using

mitochondrial ATP production inhibitor (mito-I) composed of rotenone (mitochondrial

complexⅠblocker, applied to inhibit cyclophilin D to reduce mPTP activity in cells, 10 μM),

anti-mycin A (to inhibit electron transfer between Cyt b and Cyt c of complexⅢ, 4 μM) and

oligomycin (to inhibit ADP phosphorylation and electron transport, 10 μM)[22]. The

experimental results show that mito-I we selected can effectively block the energy supply

of neurons. As Fig.3A shown，about 2 minutes after the addition of mito-I to artificial

cerebrospinal fluid (ACSF), the neurons were injected with 400 pA currents and the firing

frequency of action potential (AP) was significantly lower than that before the addition of

mito-I (from 20.67 ± 1.33 to 13.67 ± 2.09 Hz, n = 6, t_5 = 4.134 , P = 0.0090 , paired Student's

t test; Fig. 3B). When THz applied, the AP frequency increased significantly from 13.67 ±

2.09 to 17.00 ± 1.84 Hz (n = 6, t_5 = 5.000, P = 0.0041, paired Student's *t* test; Fig. 3B),

basically returned to the level before adding mito-I (from 20.67 ± 1.33 to 17.00 ± 1.84 Hz,

n = 6, t_5 = 2.314, P = 0.0686, paired Student's *t* test; Fig. 3B). Although slightly reduced,

mito-I had no significant effect on the AP amplitude (from 80.92 ± 3.54 to 74.94 ± 6.33 mV,

n = 6, t_5 = 2.089, P = 0.0910, paired Student's *t* test; Fig. 3B) and halfwidth (from 1.71 ±

0.17 to 1.61 ± 0.32 ms, n = 6, t_5 = 0.6535, P = 0.5423, paired Student's *t* test; Fig. 3B).

Meanwhile, although both mito-I and THz illumination led to a slight decrease in the AP

area (from 142.4 ± 5.65 to 118.0 ± 8.45 mV·ms, n = 6, t_5 = 3.937 , P = 0.0110 and from

118.0 ± 8.45 to 78.03 ± 13.79 mV·ms, n = 6, t_5 = 2.716 , P = 0.0420 ,respectively, paired

Student's *t* test; Fig. 3B), their kinetic properties such as rise slope and decay slope did not

change significantly (Fig. S1).

The effect of mito-I accumulated over time, about 8 minutes after the addition of mito-I

to ACSF, the neurons no longer generated APs when 400 pA currents were injected.

Therefore, we injected a much stronger 600 pA current (Fig. 3C). Interestingly, at this point

the frequency of AP decreased from 7.33 ± 1.23 to 4.67 ± 0.67 Hz during THz illumination

(Fig. 3D), the AP amplitude and halfwidth were also dramatically reduced from 76.26 ± 2.30

to 64.04 ± 3.17 mV during THz illumination (Fig. 3D) and from 1.46 ± 0.03 to 0.89 ± 0.05

ms during THz illumination (Fig. 3D), which resulted in the area decreased from 115.9 ±

5.49 to 60.74 ± 6.48 mV·ms (Fig. 3D). Meanwhile, dynamic properties such as rise slope

and decay slope were significantly enhanced (from 154.2 ± 5.89 to 174.3 ± 6.26 mV/ms, n

= 6; from -50.01 ± 1.64 to -76.04 ± 2.39 mV/ms, respectively; Fig. 3D). We speculate that

this is because the AP is affected by ATP and other factors, such as Na^+ - K^+ pump activity,

ion concentration inside and outside the membrane, enzyme activity, membrane properties

such as membrane resistance, etc[23-26]. Long periods of inadequate energy supply leads

to irreversible changes or damage to a variety of influencing factors (Fig. S3), under the

complex and comprehensive action of many factors, THz illumination at this time enhanced

the dynamic properties of AP but did not restore the AP frequency. Besides, Our previous

study found that THz radiation can change the kinetic properties of ion channels and

different cell states lead to different excitability changes[19,20], so we speculate that THz

may enhance the kinetic properties of Na+ channels in this cellular state, leading to an

increase in the rising slope of the AP, which also accelerate the repolarization process,

leading to the decrease in the AP amplitude.

Does THz illumination change the amount of ATP in nerve tissue? We next employed

two-photon Ca^{2+} imaging to visualize the changes of ATP during THz illumination *in vivo*.

We measured ATP in neurons and astrocytes respectively in mouse sensory cortex. The

sensory cortex was injected with an ATP-sensitive probe (rAAV-hSyn-GRAB (ATP1.0),

target neurons or rAAV-GfaABC1D-GRAB (ATP1.0), target astrocytes). We performed

real-time (40 Hz) two-photon imaging over the entire course of THz illumination (2 mins

before, 4 mins THz illumination, 4 mins after; Fig. 4A). Since the two fluorescence probes

we used have just been developed and no relevant reports have been reported[27], we

first tested the effectiveness of the two probes. As the Fig. S1 shown, after the addition of

mito-I (the drug composition was the same as that of patch clamp experiment, and the

concentration was 5 times that of brain slice experiment), the fluorescence intensity

decreased significantly and after the addition of ATP (100 µM), the fluorescence intensity

increased, which support the effectiveness of the fluorescent probes we used.

Next, we used the above two probes to test the direct effects of THz illumination on

ATP in neurons and astrocytes *in vivo* via two-photon imaging. An example neuron (Fig.

4B, C) showed that THz illumination reliably induced ATP in neurons during the illumination

time (2-6 min) and astrocytes changed in the same way as neurons (Fig. 4D, E,H,I). The

changes of intracellular ATP in HEK293T cells is consistent with the extracellular ATP in

vivo (Fig. S3)[28,29]. We speculate that the reason is that the body may have a

"compensatory" response, that is, the THz photons are energy-supplying during the

illumination, so the cells no longer need to synthesize large amounts of ATP, resulting in

reduced ATP release. We also

Previous studies have reported that ATP could play the role of peroxidase in

mitochondria to remove excess reactive oxygen species (ROS) and ruled out the possibility

that ATP promoted the decomposition of ROS by releasing energy[30]. We therefore tested

whether and how THz illumination in the nerve cells affect the ROS in mitochondria. We

used a mito-targeted reporter (MitoSOX Red, 5 μM) to measure mitochondrial superoxide

production, imaging time was 10 min for each group (2 mins before, 4 mins THz illumination,

4 mins after)[31]. A typical example of changes of ROS in mitochondria is shown in Fig.

4F&G, in which there is obvious increase of ROS during THz illumination (Fig. 4J). These

results together suggest that THz illumination results in a decrease of ATP in nerve cells

and affects the biochemical reactions involved in ATP such as the breakdown of ROS.

Conclusion

In this study, we report that THz photons can effectively recover reduced action potential

emission frequency due to inadequate ATP supply and change the amount of ATP and

ROS in nerve cells without introducing any exogeneous substances. Besides, this control

technology has good targeting and direction of brain tissue *in vivo*, which is fundamentally

different from some well-known and widely-deployed human function regulation technology

such as optogenetics and electrode implantation. Our results illustrate that biophotons may

play a key role in neural information processing and encoding and may be involved in

quantum brain mechanism.

We hope that our findings can bring a new viewpoint that neural communication or

physiological signal transduction is likely to be essentially the transmission of endogenous

terahertz waves and other high-frequency electromagnetic signals and provide an

explanation of how energy can be used efficiently. However, due to the complex coherence

of biological quantum information[32-36], its related exploration still has a long way to go.

Overall, if the generation, transmission, and coupling of neural signals are in the form of

THz photons, then exogenous THz photons input can change and even regulate various

life activities, which can not only explain the efficient and low-consumption information

processing of human beings, but also provide new ideas for exploring the mysteries of the

brain and the development of artificial intelligence. At the same time, the relevant technical

means have great application prospects in the fields of disease treatment and human

function regulation.

Acknowledgements

This work was supported by the National Natural Science Foundation of China
(No.12225511 and T2241002), National Defense Science and Technology Innovation
Special Zone, and the National Supercomputer Center in Tianjin. We thank the support
from Prof. Yulong Li and Dr Zhaofa Wu of Peking University. C.C. acknowledges the
support from the XPLORER Prize. The authors are grateful to Dr. Bo Song for inspiring our
work.

Competing interests

192   The authors declare no competing interests.

References

[1]. Y. M. Abbas, D. Wu, S. A. Bueler, C. V. Robinson, J. L. Rubinstein, Structure of V-ATPase
from the mammalian brain. Science 367, 1240-1246 (2020).

[2]. R. C. Coll, J. R. Hill, C. J. Day, A. Zamoshnikova, D. Boucher, N. L. Massey, J. L. Chitty,
J. A. Fraser, M. P. Jennings, A. A. B. Robertson, K. Schroder, MCC950 directly targets the
NLRP3 ATP-hydrolysis motif for inflammasome inhibition. Nat. Chem. Biol. 15, 556-559
(2019).

[3]. F. Di Virgilio, A. C. Sarti, S. Falzoni, E. De Marchi, E. Adinolfi, Extracellular ATP and P2
purinergic signalling in the tumour microenvironment. Nature reviews. Cancer 18, 601-618
(2018).

[4]. G. Burnstock, Historical review: ATP as a neurotransmitter. Trends Pharmacol. Sci. 27,
166-176 (2006).

[5]. P. Illes, G. Burnstock, Y. Tang, Astroglia-Derived ATP Modulates CNS Neuronal Circuits.
Trends Neurosci. 42, 885-898 (2019).

[6]. G. Burnstock, A unifying purinergic hypothesis for the initiation of pain. The Lancet 347,
1604-1605 (1996).

[7]. G. S. Shadel, T. L. Horvath, Mitochondrial ROS signaling in organismal homeostasis. Cell
163, 560-569 (2015).

[8]. D. Nolfi-Donegan, A. Braganza, S. Shiva, Mitochondrial electron transport chain: Oxidative
phosphorylation, oxidant production, and methods of measurement. Redox Biology 37,
101674 (2020).

[9]. E. Bertero, C. Maack, Calcium Signaling and Reactive Oxygen Species in Mitochondria.
Circ. Res. 122, 1460-1478 (2018).

[10]. N. S. Guest, T. A. Vandusseldorp, M. T. Nelson, J. Grgic, B. J. Schoenfeld, N. D. M.
Jenkins, S. M. Arent, J. Antonio, J. R. Stout, E. T. Trexler, A. E. Smith-Ryan, E. R. Goldstein,
D. S. Kalman, B. I. Campbell, International society of sports nutrition position stand:
caffeine and exercise performance. J. Int. Soc. Sport. Nutr. 18 (2021).

[11]. B. Nourbakhsh, N. Revirajan, B. Morris, C. Cordano, J. Creasman, M. Manguinao, K.
Krysko, A. Rutatangwa, C. Auvray, S. Aljarallah, C. Jin, E. Mowry, C. Mcculloch, E.
Waubant, Safety and efficacy of amantadine, modafinil, and methylphenidate for fatigue in
multiple sclerosis: a randomised, placebo-controlled, crossover, double-blind trial. Lancet
Neurol. 20, 38-48 (2021).

[12]. Z. Wang, N. Wang, Z. Li, F. Xiao, J. Dai, Human high intelligence is involved in spectral
redshift of biophotonic activities in the brain. Proc. Natl. Acad. Sci. U.S.A 113, 8753-8758
(2016).

[13]. R. Tang, J. Dai, Biophoton signal transmission and processing in the brain. J.Photochem
139, 71-75 (2014).

[14]. K. Wu, C. Qi, Z. Zhu, C. Wang, B. Song, C. Chang, Terahertz Wave Accelerates DNA
Unwinding: A Molecular Dynamics Simulation Study. J. Phys. Chem. Lett 11, 7002-7008
(2020).

[15]. F. Zhang, B. Song, L. Jiang, The quantized chemical reaction resonantly driven by
multiple MIR-photons: From nature to the artificial. Nano Res. 14, 4367-4369 (2021).

[16]. N. Li, D. Peng, X. Zhang, Y. Shu, F. Zhang, L. Jiang, B. Song, Demonstration of
biophoton-driven DNA replication via gold nanoparticle-distance modulated yield oscillation.
Nano Res. 14, 40-45 (2021).

[17]. Z. Zhu, C. Chen, C. Chang, B. Song, Terahertz-Light Induced Structural Transition and
Superpermeation of Confined Monolayer Water. ACS Photonics 8, 781-786 (2021).

[18]. C. Zhang, Y. Yuan, K. Wu, Y. Wang, S. Zhu, J. Shi, L. Wang, Q. Li, X. Zuo, C. Fan, C.
Chang, J. Li, Driving DNA Origami Assembly with a Terahertz Wave. Nano Lett. 22, 468-
475 (2022).

[19].T. Xiao, K. Wu, Y. Ding, X. Yang, C. Chang, Y. Yang, Reversible and gain modulation of
neuronal responses and sensorimotor behavior by mid-infrared stimulation.
bioRxiv. https://doi.org/10.1101/2022.04.27.489693

[20]. X. Liu, Z. Qiao, Y. Chai, Z. Zhu, K. Wu, W. Ji, D. Li, Y. Xiao, L. Mao, C. Chang, Q. Wen,
B. Song, Y. Shu, Nonthermal and reversible control of neuronal signaling and behavior by
midinfrared stimulation. Proc. Natl. Acad. Sci. U.S.A. 118 (2021).

[21]. J. C. Zhang, F. Q. Liu, S. Tan, D. Y. Yao, L. J. Wang, L. Li, J. Q. Liu, Z. G. Wang, High-
performance uncooled distributed-feedback quantum cascade laser without lateral
regrowth. Appl. Phys. Lett. 100, 112105 (2012).

[22]. A. Agarwal, P. Wu, E. G. Hughes, M. Fukaya, M. A. Tischfield, A. J. Langseth, D. Wirtz,
D. E. Bergles, Transient Opening of the Mitochondrial Permeability Transition Pore Induces
Microdomain Calcium Transients in Astrocyte Processes. Neuron 93, 587-605 (2017).

[23]. A. Gidon, T. A. Zolnik, P. Fidzinski, F. Bolduan, A. Papoutsi, P. Poirazi, M. Holtkamp, I.
Vida, M. E. Larkum, Dendritic action potentials and computation in human layer 2/3 cortical
neurons. Science 367, 83-87 (2020).

[24]. P. S. Kaeser, W. G. Regehr, Molecular mechanisms for synchronous, asynchronous, and
spontaneous neurotransmitter release. Annu. Rev. Physiol. 76, 333-363 (2014).

[25]. W. Wang, R. Mackinnon, Cryo-EM Structure of the Open Human Ether-à-go-go -Related
K + Channel hERG. Cell 169, 422-430 (2017).

[26]. E. T. Kavalali, The mechanisms and functions of spontaneous neurotransmitter release.
Nat. Rev. Neurosci. 16, 5-16 (2015).

[27]. Z. Wu, K. He, Y. Chen, H. Li, S. Pan, B. Li, T. Liu, F. Xi, F. Deng, H. Wang, J. Du, M. Jing,
Y. Li, A sensitive GRAB sensor for detecting extracellular ATP in vitro and in vivo. Neuron
110, 770-782 (2022).

[28].S. Arai, R. Kriszt, K. Harada, et al. Liang-Sheng Looi. RGB-color intensiometric indicators
visualize spatiotemporal dynamics of ATP in single cells.   Angew Chem Int Ed 57, 10873-
10878 (2018). .

[29].Lobas, M., Tao, R., Nagai, J. et al. A genetically encoded single-wavelength sensor for
imaging cytosolic and cell surface ATP. Nat Commun 10, 711 (2019)

[30]. Y. Shi, M. Tang, C. Sun, Y. Pan, L. Liu, Y. Long, H. Zheng, ATP Mimics pH-Dependent
Dual Peroxidase-Catalase Activities Driving H_2O_2 Decomposition. CCS Chemistry 1, 373-
383 (2019).

[31]. K. M. Robinson, M. S. Janes, J. S. Beckman, The selective detection of mitochondrial
superoxide by live cell imaging. Nat. Protoc. 3, 941-947 (2008).

[32]. S. Tarantini, M. N. Valcarcel-Ares, P. Toth, A. Yabluchanskiy, Z. Tucsek, T. Kiss, P.
Hertelendy, M. Kinter, P. Ballabh, Z. Süle, E. Farkas, J. A. Baur, D. A. Sinclair, A. Csiszar,
Z. Ungvari, Nicotinamide mononucleotide (NMN) supplementation rescues
cerebromicrovascular endothelial function and neurovascular coupling responses and
improves cognitive function in aged mice. Redox Biology 24, 101192 (2019).

[33].M. Tegmark, Importance of quantum decoherence in brain processes.
Phys. rev., E Stat. phys. plasmas fluids relat. interdiscip. topics 61, 4194-4206 (2000).

[34]. G. Liu, C. Chang, Z. Qiao, K. Wu, Z. Zhu, G. Cui, W. Peng, Y. Tang, J. Li, C. Fan, Myelin
Sheath as a Dielectric Waveguide for Signal Propagation in the Mid-Infrared to Terahertz
Spectral Range. Adv. Funct. Mater. 29, 1807862 (2019).

[35]. Z. Xiang, C. Tang, C. Chang, G. Liu, A new viewpoint and model of neural signal
generation and transmission: Signal transmission on unmyelinated neurons. Nano Res. 14,
590-600 (2021).

[36]. Z. X. Xiang, C. X. Tang, C. Chang, G. Z. Liu, A primary model of THz and far-infrared
signal generation and conduction in neuron systems based on the hypothesis of the
ordered phase of water molecules on the neuron surface I: signal characteristics. Sci. Bull.
65, 308-317 (2020).

Fig.1. Mechanistic study of the energy supply driven by THz photons. (**A**) Schematic indicating the process of THz photons release and it acted on nerve cells. (**B**) Theoretic absorption spectra of the bulk water model (orange curve) and the THz photons absorption spectra of biomolecules (blue curve). (**C**) Theoretic absorption spectra of the ATP. Inset, schematic illustration of the P-O stretches around 34±2 THz.

306
307

308 **Fig.2. Schematic drawing of the THz illumination device.** The QCL was described in

309 detail in a previous report[21]. The biological tissue such as brain slice is incubated in a

310 chamber containing perfusion solution (also see SI methods).

311
312
313

Fig.3. terahertz photons recovers related neural activity. (A) Representative AP spiking

before adding mito-I, after adding mito-I and THz illumination after adding mito-I (2-3 min).

(B) Comparison of the changes in frequency, amplitude, halfwidth and area, the three

groups of data of each figure correspond to the three states of the Fig.3A . **(C)**

Representative AP spiking before and during THz illumination (8-9 min). **(D)** Comparison

of the changes in frequency, amplitude, halfwidth and area, the two groups of data of each

figure correspond to the two states of the Fig.3C . *, **, *** and **** represent P < 0.05,

0.01, 0.001 and 0.0001, respectively. Repeated measurement ANOVA. Error bars

represent s.e.m.

Fig.4. THz illumination leads to a decrease in ATP and an increase in ROS. (A) Schematic diagram depicting the experimental protocol in which an AAV encoding promoter is injected into the mouse sensor cortex, followed by two-photon imaging. (B and C) Exemplar fluorescence images, pseudocolor images, and individual traces of the fluorescence response of ATP measured in neurons (THz illumination at 4-6 minutes). (D and E)

Exemplar fluorescence images, pseudocolor images, and individual traces of the

fluorescence response of ATP measured in astrocytes (THz illumination at 4-6 minutes).

(H) The summary data of ATP in neurons represent 15 ROIs from 5 mice. (I) The summary

data of ATP in astrocytes represent 15 ROIs from 5 mice. (J) The summary data of ROS

represent 15 ROIs from 3 coverslips. Scale bars represent 20 μm. *, **, *** and ****

represent $P < 0.05$, 0.01, 0.001 and 0.0001, respectively. Repeated measurement ANOVA.

Error bars represent s.e.m.

Supporting information

A new means of energy supply driven by terahertz photons

recovers related neural activity

Xiaoxuan Tan[1,2,*]

[1] Astronaut Center of China, Beijing 100084, China.

[2] Innovation Laboratory of Terahertz Biophysics, National Innovation Institute of Defense Technology, Beijing 100071, China.

*Corresponding author.

Email: t15822082707@163.com (X.T.).

Methods

Animals

C57BL/6J male mice (2-3 months old) were provided by the Laboratory Animal Center of the Qsinghua University. All experimental procedures were performed in accordance with institutional animal welfare guidelines with the approval of Qsinghua University Animal Care and Use Committee.

THz-QCL laser

We used a quantum cascade terahertz laser for this study, as describe in previous studies[1,2]. The quantum cascade laser (QCL) wafer was grown on an n-doped (Si, 2 × 10^{17} cm^{-3}) InP substrate by solid source molecular beam epitaxy (MBE) based on a slightly-diagonal bound-to-continuum design. The epi-wafer was etched into double channel geometry with a ridge width of 9 μm. After etching, Fe-doped semi-insulating layers were regrown by metal-organic chemical vapor deposition (MOCVD). Silicon oxide was grown by plasma-enhanced chemical vapor deposition (PECVD) as electrical isolation layer. We set a 3 μm wide window for electron injection on each ridge by photolithography and wet etching and ohmic contact was provided by a Ti/Au layer. Then we deposited a Ge/Au/Ni/Au metal contact layer on the backside of the sample after thinning the substrate down to 120 μm. Finally, the sample was high-reflectivity (HR) coated with Al$_2$O$_3$/Ti/Au/Al$_2$O$_3$ (200/10/100/120 nm) on the back facets by e-beam evaporation, and then cleaved into 4-mm-long devices. The beam of laser was collimated by an anti-reflectivity (AR) coated high-aperture collimation lens (CL) (LightPath 390037IR1, NA = 0.86, f = 1.87 mm).

Mouse neucortical slice preparation.

Coronal slices of prefrontal cortex were obtained from C57/B6 mice (postnatal ~2 months). We anesthetized the mice with sodium pentobarbital (10 mg/kg of body weight,

intraperitoneal injection) and sacrificed with decapitation. Brain tissues were then

immediately dissected out and immersed in ice-cold oxygenated (95% O_2 and 5% CO_2)

slicing solution in which 126 mM NaCl was substituted by 213 mM sucrose. Slices (250

µm in thickness) were cut with a microtome (VT-1200S, Leica, Germany) and incubated in

the aerated normal ACSF containing (in mM): 126 NaCl, 2.5 KCl, 2 $MgSO_4$, 2 $CaCl_2$, 26

$NaHCO_3$, 1.25 NaH_2PO4, and 25 dextrose (315 mOsm, pH 7.4) and maintained at 34.5 ℃

for 1.5 h. After incubation, slices were kept in the same solution at room temperature until

use.

Electrophysiological recordings from mouse neocortical slices.

Before recordings, the slices were transferred into a recording chamber and perfused with

a regular ACSF (0.9 ml/min). Cortical neurons were visualized with an upright infrared

differential interference contrast (IR-DIC) microscope (BX51WI, Olympus) equipped with a

water-immersed objective (40x, NA 0.8). The patch pipettes for somatic recordings had

impedance of 4-6 MΩ. Electrical signals were acquired using a Multiclamp 700B amplifier

(Molecular Devices), digitized and sampled by Micro 1401 mk II (Cambridge Electronic

Design) at 25 kHz using spike2 software.

Virus injection.

Prior to surgery, the mice were fixed in a stereotactic frame (RWD, Shenzhen, China) under

a-combination of xylazine (10mg/kg) anesthesia and ketamine (100mg/kg) analgesia. A

volume of ~150 nl virus was injected using calibrated glass microelectrodes connected to

an infusion pump (micro 4, WPL, USA) at a rate of 30 nl/min. The coordinates were defined

as dorso-ventral (DV) from the brain surface, anterior-posterior (AP) from bregma, and

medio-lateral (ML) from the midline (in mm).

The glass pipettes used to perform virus injection were beveled at 45° with a 10 to 15 µm

opening. After infusing solution, we kept the pipette in the brain for about 10 min before

pulling up the pipette to avoid backflow during withdrawal. About 150 nl volume of rAAV-

hSyn-GRAB (ATP1.0) or rAAV-GfaABC1D-GRAB (ATP1.0) solution (~2×10^{12} infectious

units/ml) was injected into sensory cortex slowly (AP: ~-1.5 mm, ML: `~2.0mm, DV:

~0.35mm) for imaging calcium activity with GCaMP6f.

Two-photon Ca^{2+} imaging. A commercial Nikon A1R two-photon microscope system was

used to perform two-photon calcium imaging. Two-photon excitation beam was emitted by

a mode-locked Ti: Sa laser (model "Mai-Tai Deep See", Spectra Physics). We utilized a

water-immersion objective (Nikon) with 25 ×/1.10NA to perform imaging. The excitation

wavelength was set to 910 nm for Gcamp6f calcium imaging experiments. For somatic

imaging, the dimension of field-of-view (FOV) was set 200 μm × 200 μm. We acquired

images of 512 × 512 pixels at 30-Hz frame rate. The average power reaching the cortical

surface ranged from 30 to 40 mW, depending on the expression efficiency of virus and

depth of imaging. Within an imaging time window of ~3 min (~1 min per time, three times

in total) for each imaging, no sign of photo-damage was observed. Before the experiment,

the mice were fixed under microscope objective frequently with the chamber to adapt to

the imaging state for real calcium activity.

Quantum-chemical calculations. Quantum-chemical calculations were performed to

obtain the infrared spectroscopy of ATP. The PBE0 exchange correlation functional and 6-

311G* basis set were employed in the DFT calculations. The polarized continuum model

(PCM) was used for modeling solvation effects together with water as the solvent. The

geometry optimization calculations were performed first to obtain the stable conformations

of molecules and the lowest energy conformation was used for calculating results. The

frequency of absorbing/emissive photon was identified by the vibrational mode. All

calculations were performed by the Gaussian 16 software package and Gaussview 6.0

visualization program[3-5].

Cell cultures

HEK293T cells were obtained from ATCC (CRL-3216) and verified based on their

morphology under the microscope and by their growth curve. HEK293T cells were cultured

at 37℃ in 5% CO2 in DMEM (Biological industries) supplemented with 10% (v/v) fetal

bovine serum (FBS, GIBCO) and 1% penicillin-streptomycin (Biological Industries).

Rat primary neurons were prepared from 0-day old (P0) pups (male and female, randomly

selected) in Ref.6-7. Cortical and hippocampal neurons were dissociated from the dissected brains in 0.25% Trypsin-EDTA (Gibco) and plated on 12-mm glass coverslips coated with poly-D-lysine (Sigma-Aldrich) in neurobasal medium (Gibco) containing 2% B-27 supplement (Gibco), 1% GlutaMAX (Gibco), and 1% penicillin-streptomycin (Gibco). Based on glial cell density, after approximately 3 days in culture (DIV 3), cytosine β-D-arabinofuranoside (Sigma) was added to the hippocampal cultures in a 50% growth media exchange at a final concentration of 2 uM. The neurons were cultured at 37°C in 5% CO_2.

Expression of MaLion in HEK293T cells

For confocal imaging, HEK293T cells were plated on 12-mm glass coverslips in 24-well plates and grown to 60-80% confluence for transfection. Cells were transfected using a mixture containing 1 μg DNA and 1 μg PEI for 4-6 hours and imaged 24-48 hours after transfection.

Measurement of mitochondrial ROS production

Mitochondrial production of ROS in cells was measured using MitoSOX Red (Invitrogen/Thermo Fisher Scientific), a mitochondrion-specific hydroethidine-derivative fluorescent dye[8]. In brief, cells were incubated with MitoSOX (5 uM; for 30 min, at 37°C, in the dark). The cells were then washed with PBS and MitoSOX fluorescence was measured by confocal imaging.

Statistical analysis

We used GraphPad Prism 5 and MATLAB to perform the statistical analysis. Except where indicated otherwise, summary data are presented as the mean ± SEM. All measurements were taken from distinct samples.

References

[1] C. Zhang, Y. Yuan, K. Wu, Y. Wang, S. Zhu, J. Shi, L. Wang, Q. Li, X. Zuo, C. Fan, C. Chang, J. Li, Driving DNA Origami Assembly with a Terahertz Wave. *Nano Lett.* 22, 468-475 (2022).

[2] J. C. Zhang, F. Q. Liu, S. Tan, D. Y. Yao, L. J. Wang, L. Li, J. Q. Liu, Z. G. Wang, High-performance uncooled distributed-feedback quantum cascade laser without lateral

regrowth. *Appl. Phys. Lett.* 100, 112105 (2012).

[3] R. Krishnan, J. S. Binkley, R. Seeger, J. A. Pople. Self-consistent molecular orbital

methods. XX. A basis set for correlated wave functions. *J. Chem. Phys.* 72, 650-654 (1980).

[4]C. Adamo , M. Cossi, V. Barone. An accurate density functional method for the study of

magnetic properties: the PBE0 model. *J. Mol. Struct,* 493(1-3):145-157 (1999).

[5]Gaussian 16, Revision B.01, Frisch, M. J. et al. Gaussian, Inc., Wallingford CT, 2016.

[6] Z. Wu, K. He, Y. Chen, H. Li, S. Pan, B. Li, T. Liu, F. Xi, F. Deng, H. Wang, J. Du, M.

Jing, Y. Li, A sensitive GRAB sensor for detecting extracellular ATP in vitro and in vivo.

Neuron. 110, 770-782 (2022).

[7] W.Peng, Z. Wu, K. Song, S. Zhang, Y. Li, M. Xu, Regulation of sleep homeostasis

mediator adenosine by basal forebrain glutamatergic neurons. *Science.* 369,1208 (2020).

[8] K. M. Robinson, M. S. Janes, J. S. Beckman, The selective detection of mitochondrial

superoxide by live cell imaging. Nat. Protoc. 3, 941-947 (2008).

493   **Fig.S1. The decay slope (A) and rise slope (B) of the action potential.**

499   **Fig. S2. The fluorescence response to mito-I and ATP measured in neurons.**

Fig. S1. The changes in membrane potential caused by mito-I at different times.

Fig. S4. The changes of intracellular ATP during THz irradiation. (A) Fluorescence

images of HEK293T expressing MaLionG and MaLionG-neg when the irradiation duration

511   is 0s (left) and 350s (right) respectively. **(B)** Fluorescence images of HEK293T expressing

512   MaLionR and MaLionR-neg when the irradiation duration is 0s (left) and 350s (right)

513   respectively. **(C)** and **(D)** Normalized F/F0 was measured in MaLion and MaLion-neg

514   expressing HEK293T cells in response to THz continuously applied (left: green; right: red);

515   n=25 cells from 3 cultures. Scale bars represent 100 μm.

517