Neuronal IL-17 controls *C. elegans* developmental diapause through CEP-1/p53

Abhishiktha Godthi[1,2,], Sehee Min[1,2,], Srijit Das[1], Johnny Cruz-Corchado[1,2], Andrew Deonarine[1],

Kara Misel-Wuchter[3], Priya D. Issuree[3] and Veena Prahlad[1,2] *

Affiliations: [1]Department of Cell Stress Biology

Roswell Park - Comprehensive Cancer Center

Elm and Carlton Streets

CGP-BLSC   L3-307

Buffalo, New York 14263

[2]Department of Biology,

338 Biology Building East,

143 Biology Building, Iowa City, IA 52242-1324.

[3]Department of Internal Medicine,

503 Eckstein Medical Research Building,

431 Newton Road, Iowa City, IA 52242

* Corresponding author email**:** : veena.prahlad@roswellpark.org

Abstract. How metazoan development is coordinated with nutrient availability remains unclear.

Here, we show that the activity of CEP-1, the *C. elegans* ortholog of the tumor suppressor p53,

switches organismal development between reproductive growth and dormancy by responding to

nutrient availability signaled by neuronal cytokine, ILC-17.1. Specifically, ILC-17.1 is released

from amphid neurons upon food availability and promotes glucose utilization and represses CEP-

1/p53: CEP-1/p53 repression promotes growth. In the absence of ILC-17.1, CEP-1/p53 is

activated, accumulates in post-embryonically dividing progenitor blast cells, upregulates cell-cycle

inhibitors, decreases cytochrome C levels, and causes larvae to arrest as stress-resistant,

quiescent dauers. We propose a model whereby ILC-17.1 signaling links nutrient availability and

energy metabolism to cell cycle progression and cell fate decisions during development through

CEP-1/p53. These findings also suggest that the developmental function of the p53-gene family

could have shaped their evolution and are relevant to our understanding of neuroimmune

mechanisms in cancer.

Main

During metazoan development, nutrient availability is coordinated with the division, growth and metabolic activity of individual cells through cell-cell communication. This is also the case in the invertebrate *C. elegans*, a free-living bacterivore, which displays a dramatic developmental plasticity to ensure that its growth and reproduction match available resources[1-10]. When *C. elegans* larvae hatch under optimal conditions (e.g., in the laboratory at 20°C, at low population densities, on abundant food provided by lawns of the *E. coli* strain, OP50) they continue development into reproducing adults. However, if they hatch under suboptimal conditions, such as in the paucity of food, at high population densities, or high ambient temperatures, larvae implement an alternative developmental program and arrest as quiescent, stress-resistant larvae called 'dauer' larvae. Dauer larvae display metabolic and organismal phenotypes specialized for dispersal and survival, and can remain for months in an arrested state to resume development into reproductive adults when conditions become favorable again[1-10]. Studies from several labs have identified molecular pathways that mediate the dauer decision, showing that growth promoting molecules, the best-characterized of which are the insulins, the transforming β growth factor (TGFβ/DAF-7) and lipid based dafachronic acid hormones, are released by sensory neurons and other cells to license continued growth and development; adverse environments inhibit these growth promoting signals and trigger dauer arrest[1-11]. In addition, a number of quantitative trait loci (QTL) also modulate dauer arrest[12]. Yet, how the dauer entry decision results in a coordinated change in cell fates across different tissues and is linked with the systemic shut-down of anabolic pathways still remains poorly understood.

An important group of proteins that mediate cell-cell communication and metabolism in metazoa are secreted proteins called cytokines[13,14]. The IL-17 cytokines are one such evolutionarily conserved family of proinflammatory cytokines that are released by specialized immune cells at the interface between an organism and the outside world to activate immune surveillance, barrier

function and wound healing, but in humans, also promote cancers and autoimmune disease such

as psoriasis[15,16]. More recently IL-17s have been shown to play key immunometabolic roles in

modulating mammalian physiology and disease[17]. Here, we show that the *C. elegans* IL-17

ortholog, ILC-17.1, is released from amphid neurons in response to food, and coordinates cell

division and cell fate decisions with metabolism by controlling the activity of the *C. elegans* tumor

suppressor p53 ortholog, CEP-1. Specifically, ILC-17.1 release that occurs in the presence of

food suppresses CEP-1/p53. CEP-1/p53 is activated upon the loss of ILC-17.1, and remarkably,

this switches whole organism development from continuous growth to dormancy. The p53-like

tumor suppressor genes are found in all multicellular animals where they prevent the transmission

of damaged DNA by activating a multifaceted program that controls cell cycle checkpoints,

mediates reversible growth arrest or apoptosis, and controls metabolic flux in cells [18-21]. Our

studies show that these functions of CEP-1/p53 also act to control developmental quiescence of

a whole organism, *C. elegans*, suggesting that the developmental function of the p53- gene family

could have shaped their evolution[22-24].

Results

ILC-17.1 is released by *C. elegans* amphid neurons in response to food and prevents dauer

arrest.

C. elegans express three IL-17 orthologs[25,26]. Of these, ILC-17.1, which has been shown to act

as a neuromodulator, is expressed by a subset of specialized neurons with epithelial properties

called amphid neurons that are exposed to the outside and sense information from the

environment[25-27]. We discovered that a deletion of *ilc-17.1,* syb5296*,* that removes almost all the

coding sequence (2173bp out of 2980 bp; Supplementary Fig. S1a) and abolishes mRNA

expression (Supplementary Fig. S1b) caused larvae to constitutively enter the dauer state even

under optimal growth conditions (Supplementary Fig. S1c-d)*.* Dauer larvae can be identified by

their growth arrest, distinct morphology that includes specialized cuticular structures called alae,

a constricted pharynx, and arrested germline, as well as decreased pharyngeal pumping rates

and detergent (1% SDS) resistance[3-9]. *ilc-17.1*(syb5296)X dauer larvae displayed all these

characteristic morphological and physiological features (Supplementary Fig. S1e-j). Under

optimal growth conditions at 20°C, *ilc-17.1* deleted larvae entered the dauer state transiently:

approximately 30% (31.2 ±5%) were SDS-resistant 48 hours post-hatching on OP50 bacteria,

and most larvae exited dauer and become reproductive adults by 72 hrs (Supplementary Fig.

S1c, d). Under the same conditions, none of the wildtype larvae were detergent resistant, nor did

any larvae arrest as dauers (Supplementary Fig. S1c, d). However, as with other mutations that

promote dauer entry[3,4,28], the dauer arrest of *ilc-17.1* deletion mutants was dramatically

accentuated at the slightly more stressful ambient temperature of 25°C, which still supported the

growth of all wildtype larvae into reproductive adults, but caused practically 100% of larvae lacking

ilc-17.1 to enter and persist in the arrested dauer state (Fig.1a-d).

Because the molecular basis for dauer regulation has been the target of extensive genetic

screens that are considered to be saturated[3,5,29-31], and IL-17.1 had not been previously implicated

in the regulation of dauer, we confirmed the role of *ilc-17.1* in dauer entry by (i) examining the

developmental phenotype of another independently generated CRISPR deletion in *ilc-17.1,*

syb5297 (loss of 2188 bp of 2980 bp; Supplementary Fig. S1a), (ii) by backcrossing *ilc-17.1,*

syb5297, and (iii) by downregulating *ilc-17.1* and its receptor[25,26] in wildtype (N2) larvae using

RNA interference (RNAi). Like *ilc-17.1* (syb5296), *ilc-17.1* (syb5297) created using a different

guide RNA (see Materials and Methods) also displayed a completely penetrant dauer arrest at

25°C (Fig. 1a,b). In *C. elegans* there have been no off-target effects observed using CRISPR

even after whole genome sequencing by several labs[32-34]. Nevertheless, due to reviewer's

skepticism, we considered whether a similarly large CRISPR-mediated deletion of any gene

could, in itself, predispose larvae to enter dauer, and examined whether of the deletion of a related

gene *F25D1.3,* syb7367(loss of 1475 bp of 1642 bp) also triggered dauer arrest. This did not

(Supplementary Fig. S1k). *ilc-17.1* (syb5296) larvae, however, continued to arrest as dauers even

after they were backcrossed 2X time into a wildtype background (Supplementary Fig. S1k). In

addition, RNAi-mediated downregulation of *ilc-17.1* in wildtype (N2) animals, but not control RNAi

treatment, also caused a small but significant number of larvae to arrest as SDS-resistant dauer

larvae at 25°C, (Supplementary Fig. S1l). ILC-17.1 signals though cytokine receptors which in *C.*

elegans consists of ILCR-1/ILCR-2[25,26]. The ILCR-2 subunit is widely expressed, as determined

using a bicistronic SL2 cassette to tag the *ilcr-2* receptor at its endogenous locus with GFP

(Supplementary Fig. S2a). Downregulation of *ilcr-2* in wildtype (N2) animals through RNAi caused

over one-third, 39.6% ± 4%, of the wildtype larvae to arrest as dauers and did not alter the dauer

entry of *ilc-17.1* deleted larvae (Fig. 1e). RNAi, being a partial downregulation of mRNA levels,

is generally insufficient to induce dauer arrest with even known dauer pathway genes[3]. Therefore,

the RNAi-induced dauer entry of larvae exposed to *ilc-17.1* or *ilcr-2* RNAi for less than 48 hrs.

post-hatching, (see Materials and Methods) was not only statistically significant, but also provided

additional support for the role of *ilc-17.1* in dauer entry. Thus, together, these observations

indicated that during normal development, ILC-17.1 signaling, likely through the ubiquitously

expressed ILCR-2 receptors, was required for growth of larvae to reproducing adults; in the

absence of ILC-17.1, or the ILCR-2 receptor, larvae arrested growth and entered the dauer state,

transiently, or for prolonged durations.

The dauer phenotype of *ilc-17.1* deletion mutants could be rescued by re-expressing *ilc-17.1*

under the control of its own promoter and 3' UTR regions (Fig.1b; see Supplementary Fig. S2b

for expression levels of rescue construct in comparison to the endogenous gene). The extent of

rescue varied based on the animal's diet (ILC-17.1 re-expression caused an approximately 32.8%

rescue on OP50 bacteria, Fig 1b, but 63.6% rescue on HT115 bacteria, Supplementary Fig. S2c;

Chi-squared = 121, df = 1, p-value <2e-16), suggesting that the role of ILC-17.1 signaling in dauer

might be related to food, a prominent modulator of the *C.elegans* dauer decision[3-9]. The systemic

effects of the *ilc-17.1* deletion and rescue were surprising because previous studies using

transgenic *ilc-17.1* reporter constructs had shown that *ilc-17.1* mRNA expression is restricted to

a small subset of neurons, a pair of amphid sensory neurons (ASE) and two pairs of interneurons

(AUA and RMG)[25]. The expression of the endogenous *ilc-17.1* gene was also restricted to a few

neurons: mCherry expressed as a bicistronic SL2 cassette along with *ilc-17.1* mRNA from the

endogenous *ilc-17.1* locus was also only visible in a few dye-filling amphid neurons

(Supplementary Fig. S2d).   Therefore, to understand how ILC-17.1 acts systemically, we re-

expressed *ilc-17.1* ectopically in *ilc-17.1* deleted larvae, remote from its normal site of expression,

using the muscle-specific *unc-54* promoter (Supplementary Fig. S2e), and asked whether ILC-

17.1 could act in a cell non-autonomous manner to promote development, or whether its only

functioned cell autonomously in amphid neurons. Our expectation was that if ILC-17.1 controlled

developmental pathways by acting in a cell-autonomous manner in the amphid neurons, ectopic

expression would not rescue the dauer arrest, whereas a cell non-autonomous or endocrine mode

of activity could result in a rescue. Consistent with a cell non-autonomous mode of action, and as

was the case for the previously reported effects of ILC-17.1 on behavior[25], ILC-17.1 expressed

ectopically from the *unc-54* promoter rescued dauer arrest of *ilc-17.1* deleted larvae (Fig.1b and

Supplementary Fig. S2c). The rescue from this ectopic expression did not vary with bacterial diet

(Fig 1b vs Supplementary Fig. S2c), perhaps because of its non-neuronal expression and/or

higher expression levels when compared to that from the endogenous promoter (Supplementary

Fig. S2b, f). RNAi mediated downregulation of *ilcr-2* reduced the percentage of *ilc-17.1* larvae that

were rescued from dauer arrest by re-expressing ILC-17.1 either under its own promoter, or from

the muscle promoter (Fig. 1e). This effect was significant, but only ~10%, reflecting our limited

understanding of the stoichiometry of ILC-17.1 ligand-receptor interactions as they relate to the

timing of dauer arrest that occurs in the absence of ILC-17.1. Nevertheless, these studies

indicated that ILC-17.1 signaling was required for continuous growth and suggested that ILC-17.1

might normally be secreted to act cell non-autonomously to promote development.

Therefore, to more directly examine whether ILC-17.1 is normally secreted from amphid neurons,

we immunolocalized ILC-17.1 protein in *C. elegans* larvae using strains expressing ILC-17.1

tagged at its endogenous locus using a HA-tag. Suspecting that food played a role in ILC-17.1

signaling, we conducted these studies on larvae that were hatched in the absence or presence of

food (OP50; Fig. 1f). While *ilc-17.1* mRNA localization was not altered by food availability, and

remained restricted to the amphid neurons both in the presence and absence of food as detected

by mCherry expressed from the endogenous *ilc-17.1* locus (Fig. 1f; left), ILC-17.1 protein was

present outside the amphid neurons, concentrated in the pharynx when larvae were exposed to

food (OP50), but not when larvae hatched in the absence of food (Fig.1f; right, quantification in

Supplementary Fig. S2g). This suggested that ILC-17.1 protein was secreted and either retained,

stabilized, or taken up by the pharynx in response to food signals [we could not reliably detect

ILC-17.1 protein in distal tissues such as the intestine or coelomocytes]. The secretion of ILC-

17.1 protein was also supported by immunolocalization studies on the muscle-expressed ILC-

17.1 protein which, although expressed under a well-characterized muscle-specific promoter (i.e.

unc-54 ; Supplementary Fig. S2e), could be constitutively detected outside muscle cells,

throughout the animal (Supplementary Fig.S2h). Thus, the presence of ILC-17.1 protein in regions

outside its cellular sites of mRNA synthesis provided strong support that ILC-17.1 is secreted,

and in the case of the endogenous ILC-17.1 protein, secretion occurs in response to food

availability.

Sequencing total RNA (RNA-seq) extracted from bleach synchronized larvae grown for 32-36

hours at 25°C when *ilc-17.1* mutant larvae were *en route* to dauer arrest, and wildtype larvae were

in their late L2/early L3 stage *en route* to developing into reproducing adults, showed that *ilc-17.1*

mutant larvae expressed the molecular signatures of dauer larvae[6,35-37] that were distinct from

wildtype larvae and from *ilc-17.1* deletion mutants that re-expressed ILC-17.1 (Supplementary

Figure S3a; Supplementary Table 1-3). Thus, anabolic processes such as DNA replication and

ribosome biogenesis were downregulated (Fig. 1g), and catabolic processes and stress

responses such as autophagy, fatty acid metabolism, glutathione metabolism, and xenobiotic

defense pathways were upregulated (Fig. 1h). In addition, *ilc-17.1* deleted larvae expressed

dauer-specific collagen genes that differ from the collagens of larvae undergoing continuous

development[38] (Fig. 1i; Supplementary Table 4). To evaluate the extent to which the *ilc-17.1*

deleted dauers were similar to previously described *C. elegans* dauers that accumulate in the

population upon starvation or crowding*,* we compared the global gene expression changes of *ilc-*

17.1 mutant larvae with previously published gene expression changes in larvae entering dauer

in response to starvation[37], or pheromone treatment[37] (Supplementary Fig. S3 b,c; Supplementary

Tables 5, 6). Consistent with a role for ILC-17.1 in signaling nutrient availability, the expression

profile of *ilc-17.1* deleted larvae was more closely correlated to dauer larvae generated by

starvation[37] (Supplementary Fig. S3b), compared to ascaroside induced dauers[36] (Supplementary

Fig. S3c). These differences were surprising and suggest that the trigger and route of dauer entry

determines the gene expression profile of *C. elegans* dauer larvae[39,40].

ILC-17.1 loss activates CEP-1/p53 to trigger dauer entry through the activation of DAF-

16/FOXO, DAF-3/SMAD-DAF-5/Ski complex and steroid hormone pathways.

To understand how the loss of ILC-17.1 was triggering dauer entry we first considered the

possibility that, through its role as a neuromodulator[25], the secretion of ILC-17.1 modulated the

animals' sensory perception of food or feeding behavior, and the lack of *ilc-17.1* prevented larvae

from finding or ingesting food. However, this did not seem to be the case (Supplementary Fig.

S4a): *ilc-17.1* deleted animals did not exhibit decreased chemotaxis towards lawns of OP50, or

towards organic molecules like lysine or diacetyl thought to mimic the presence of bacteria[41].

Animals lacking ILC-17.1 were also able to feed like wildtype animals, and the rate of pharyngeal

pumping, a measure that is typically correlated with food uptake, was the same in wildtype animals

and animals lacking ILC-17.1, both in the presence and absence of food (Supplementary Fig.

S4b). Also, prior to their dauer entry, *ilc-17.1* deletion mutants accumulated slightly, but not

significantly fewer numbers of red latex beads in their intestinal lumen that served as a proxy for

rates of bacterial ingestion (Supplementary Fig. S4c). These data suggested that *ilc-17.1* deficient

larvae arrested development and entered a dauer diapause state despite being able to find and

ingest nutrients.

Therefore, to answer how the loss of ILC-17.1 triggered dauer entry we next adopted a genetic

approach, leveraging the wealth of studies on *C. elegans* dauer formation[2-11,28]. In *C. elegans*,

insulins released in the presence of optimal food activate or inhibit the sole insulin receptor

ortholog, DAF-2, to modulate the function of the Forkhead transcription factor DAF-16. DAF-2

signaling supports continuous growth programs: loss of DAF-2 signaling leads to the activation of

DAF-16 and promotes dauer entry[2-11,28]. Likewise, TGFβ/DAF-7 ligands released under optimal

growth conditions antagonize the activity of the transcription factor complex, DAF-3/SMAD-DAF-

5/Ski, to permit continuous growth: inhibition of DAF-7 activates DAF-3/SMAD-DAF-5/Ski and

leads to dauer formation[2-11,28]. Both insulins and TGF-β result in the production of the sterol based

dafachronic acid hormones that act through DAF-12, a nuclear hormone receptor, to support the

development of adult reproductive physiology[2-11,28]. The absence of dafachronic acid pathway

components also triggers dauer arrest, and providing exogenous dafachronic acids to mutant

animals deficient in insulin or TGFβ signaling, or decreasing DAF-12 activity, can bypass their

dauer arrest[3,42]. Finally, a mutation in a transmembrane guanylyl cyclase *daf-11,* that causes

chemosensory defects also triggers dauer arrest[3,43]. *ilc-17.1* deletion mutants lacked obvious

chemosensory defects (Supplementary Fig. 4a-c), and continued to arrest as dauers on 5mM 8-

Bromo-cGMP, which is known to rescue *daf-11*-associated dauer formation[3,43] (Supplementary

Fig. 4d). Therefore, to identify signal transduction pathways through which the loss of ILC-17.1

might act to control the dauer decision, we examined whether the other dauer effectors, *daf-16,*

daf-5, or dafachronic acid signaling, modulated dauer formation of *ilc-17.1* mutant larvae.

A *daf-16* loss of function mutation, mu86[44], completely suppressed dauer formation in animals

lacking ILC-17.1, indicating that activation of DAF-16 was one mechanism responsible for their

dauer arrest (Fig. 2a). The activation of DAF-16 in animals lacking ILC-17.1 could be quantified

by an increase in the proportion of larvae with DAF-16::GFP[45] positive intestinal nuclei even at

20°C (Fig. 2b), and the upregulation of DAF-16 target genes[46,47] both at 20°C (Supplementary

Fig. S4e) and 25°C (Fig.2c). DAF-16 activation could be suppressed by re-expressing ILC-17.1

in *ilc-17.1* deletion mutants under its endogenous promoter, or ectopically, under the muscle-

specific promoter (Fig.2c). Similarly, a loss of function mutation in *daf-5*, e1386[48] also completely

suppressed the dauer arrest of *ilc-17.1* mutant larvae (Fig. 2d). Finally, and consistent with the

role of the DAF-12 pathway downstream of DAF-16 and DAF-5/DAF-3, downregulating *daf-12* by

RNAi significantly decreased the number of larvae that arrested as dauers (Supplementary Fig.

S5a), and exposing *ilc-17.1* deficient larvae to exogenous Δ7-dafachronic acid at concentrations

of 50 nM[42], partially but significantly suppressed their dauer arrest and only ~65% instead of

~100% of the larvae were dauers (Fig 2e). The activation of DAF-16/FOXO and DAF-3/SMAD-

DAF-5/Ski by the loss of ILC-17.1 was not simply because *ilc-17.1* deleted larvae were deficient

in the expression of insulin signaling and TGFβ/ DAF-7 signaling pathway components. The

expression levels of *daf-2* mRNA and the 28 insulin ligands[49,50] were not decreased in *ilc-17.1*

deletion mutants, (Supplementary Fig. S5b *daf-2* boxed; Supplementary Table 7), and the

expression of *ins-4* was even slightly higher (Supplementary Fig. S5c). The expression of insulin

peptides INS-4 and DAF-28[50] were also not altered in larvae deficient in *ilc-17.1*, as seen using

GFP translational reporters of these insulin ligands (Supplementary Fig. S5d, e). In addition, the

dauer arrest of a mutation in the insulin receptor, *daf-2* (e1370)[51], could not be rescued by ILC-

17.1 overexpression, suggesting that ILC-17.1 did not act genetically downstream of *daf-2*

(Supplementary Fig. S5f). Similarly, the mRNA expression levels of the main TGFβ components

were mostly unaltered in *ilc-17.1*, and *daf-5* levels were even downregulated (Supplementary

Fig. S5g; Supplementary Table 8). Thus, these data together indicated that ILC-17.1 acted

upstream of DAF-16/FOXO, DAF-3/SMAD-DAF-5/Ski, and the DAF-9/DAF-12 endocrine

pathways to modulate dauer entry.

In wildtype animals, DAF-16/FOXO and DAF-3/SMAD-DAF-5/Ski complex are thought to act

independently, and in parallel to modulate the dauer decision[3,5]. Therefore, to identify

mechanism(s) that led to the activation of both these pathways in *ilc-17.1* deleted larvae, we used

RNAi to downregulate genes known to interact with these dauer effectors[52-59] (Supplementary Fig.

S5a). Intriguingly, amongst the possible candidates, we found that the downregulation of *cep-1*,

the *C. elegans* ortholog of p53[60,61] (Supplementary Fig. S5a), or a deletion in *cep-1*, gk138,

completely rescued the dauer arrest of *ilc-17.1* deletion mutants and nearly all (98%) of *cep-1;ilc-*

17.1 double mutant larvae grew into reproductive adults (Fig 2f-h).

Although p53 is best studied as a tumor suppressor, p53-like proteins modulate vertebrate

development through as yet poorly understood mechanisms[18-24]. The *C. elegans* p53 ortholog,

cep-1 has not been previously implicated in *C. elegans* development. Nevertheless, as could be

predicted by the dauer rescue of *ilc-17.1* larvae upon *cep-1* deletion, the overexpression of CEP-

1 alone was sufficient to arrest growth and promote an almost completely penetrant dauer

phenotype in larvae that developed at 25°C (Fig. 3a, b; see Supplementary Fig. S6a for mRNA

levels). CEP-1 overexpression was achieved by expressing a functional, fluorescently tagged

CEP-1 under its own promoter, CEP-1::GFP, that was able to complement a *cep-1* deletion, *cep-*

1(lg12501),[62-64] as a genomically integrated multicopy array in a wildtype background. CEP-

1::GFP did not induce dauer arrest in the background of the *cep-1* deletion, *cep-1(lg12501)*,

(Figure 3c), indicating that it was the higher expression levels of *cep-1* (Supplementary Fig. S6a),

and not the GFP tag on CEP-1/p53, that was responsible for the dauer arrest of CEP-1::GFP

overexpressing larvae. CEP-1/p53 overexpression not only caused dauer arrest at 25°C, but also

prompted larvae to transiently enter dauer at 20°C, as seen by their SDS-resistance

(Supplementary Fig. S6b), phenocopying the *ilc-17.1* deletion. RNA-seq analysis showed that

genes differentially expressed in CEP-1/p53 overexpressing larvae *en route* to their dauer arrest

significantly overlapped with those expressed in larvae lacking ILC-17.1 (65% of the 15,443

differentially expressed genes overlapped, padj.< 0.05; Supplementary Fig. S6c; Supplementary

Tables 9, 10). CEP-1/p53 overexpressing larvae also upregulated dauer-specific collagens

(Supplementary Fig. S6d; Supplementary Table 4), downregulated genes involved in anabolism

(Supplementary Fig. S6e), and upregulated genes involved in catabolism (Supplementary Fig.

S6f). In addition, between 15 hours to 32 hours post-hatching at 25°C, CEP-1/p53 overexpressing

larvae and *ilc-17.1* deletion mutants showed a remarkably constant level of gene expression, and

only 340 and 34 genes respectively changed expression ($p<0.05$), compared to the > 10,000

genes that were either upregulated or downregulated in wildtype larvae and *cep-1;ilc-17.1* double

mutant larvae that were in the process of developing into reproductive adults (PCA analysis;

Supplementary Fig. S6g; Supplementary Table 11), suggesting that the dauer decision was

implemented by 15 hrs post-hatching.

Like in the ILC-17.1 deficient animals, the completely penetrant 25°C-dauer arrest of larvae

overexpressing CEP-1 was dependent on DAF-16, DAF-5 and the steroid hormone pathway.

Downregulating *daf-16* expression by RNAi (Fig. 3d; we could not generate CEP-1/p53

overexpressing and *daf-16* double mutant larvae, perhaps because of the site of integration of

CEP-1::GFP transgene into the genome) or crossing CEP-1/p53 overexpressing larvae into *daf-*

5, e1386, suppressed their dauer arrest (Fig. 3e). Treatment with 50nM exogenous Δ7-

dafachronic acid also allowed a significant proportion of CEP-1/p53 overexpressing larvae to

bypass dauer arrest (Fig. 3e). In accordance, DAF-16 target genes were also upregulated in

larvae overexpressing CEP-1/p53 at both 25°C (Fig 3f) and 20°C (Supplementary Fig. S7a). In

addition, and most convincing, the upregulation of *daf-16* target genes in *ilc-71.1* deletion mutant

larvae was *cep-1* dependent at both 25°C and 20°C (Fig 3f; Supplementary Fig. S7a). As we

showed for ILC-17.1, CEP-1/p53 induced dauer entry appeared to be somewhat specific to

reduced ILC-17.1 signaling, as the 25°C-dauer arrest of *daf-2*(e1370) III larvae that occurred due

to reduced insulin signaling did not depend on *cep-1* (Supplementary Fig. S7b). Moreover, in

agreement with previous reports[65,66], *cep-1* was not required for dauer induction at high

temperatures as *cep-1* (gk138) I larvae could enter dauer at 27°C just like wildtype animals

(Supplementary Fig. S7c).

To confirm that CEP-1/p53 was indeed activated in the absence of ILC-17.1, we tested whether

known targets of CEP-1/p53, the BH3-only proteins *egl-1* and *ced-13*[67-70], were upregulated in *ilc-*

17.1 deleted larvae in a CEP-1/p53 dependent manner. As could be expected, *egl-1* and *ced-13*

mRNA levels were upregulated upon CEP-1/p53 overexpression in larvae both at 25°C (Fig 3g)

and 20°C (Supplementary Fig. S7d). *egl-1* and *ced-13* were also upregulated in a CEP-1/p53-

dependent manner in *ilc-17.1* deletion mutants at both 25°C and 20°C, and could be suppressed

by rescuing *ilc-17.1* deletion mutants with ILC-17.1 re-expressed under its endogenous-, or

muscle-specific promoters (Fig 3g; Supplementary Fig. S7d). In addition, C̲hromatin

I̲mmunop̲recipitation followed by quantitative PCR (ChIP-qPCR) using animals carrying an

endogenous CRISPR/Cas9 FLAG tagged *cep-1* showed that more CEP-1/p53 was bound to the

promoter regions of *egl-1* and *ced-13* in larvae lacking *ilc-17.1* when compared to wildtype larvae,

and the increased occupancy was abrogated when *ilc-17.1* deletion mutants were rescued from

dauer arrest by re-expressing ILC-17.1 (Fig. 3h).

In mammalian cells, activation of p53 in response to stress increases p53 protein levels due to its

escape from constitutive degradation mediated by the E3 ubiquitin ligase MDM2[18,20]. *C. elegans*

genome lacks MDM2 orthologs, but CEP-1/p53 is controlled through translational and

posttranslational modifications[62,71], although the mechanisms of activation remain incompletely

understood. Western analysis of FLAG-tagged endogenous CEP-1/p53 in *C. elegans* showed

that CEP-1::FLAG expression was increased in larvae lacking *ilc-17.1* (Fig 3i). As a positive

control we used gamma irradiation at doses known to activate CEP-1/p53 in the germline, and

this also increased CEP-1::FLAG levels (Fig 3i)[67,69]. These results together showed that CEP-

1/p53 was activated upon loss of ILC-17.1 signaling and triggered the dauer arrest of *ilc-17.1*

deletion mutants through the DAF-16/FOXO and DAF-3/SMAD-DAF-5/Ski (and likely DAF-12)

pathways. Thus, it appeared that ILC-17.1-CEP-1/p53 could act as a node to integrate and

coordinate the diverse cellular, physiological and metabolic effects of the DAF-2/Insulin signaling-

and TGFβ/DAF-7-growth hormone signaling pathways that normally act in parallel to control dauer

arrest. However, as we showed for DAF-2, these transcription complexes also likely receive

independent information through the insulin receptor, DAF-2, and the TGFβ/DAF-7 receptors.

ILC-17.1 modulates CEP-1/p53 activity in ventral blast cells (P cells) of larvae and directly

or indirectly controls cell cycle progression and cell fate programs.

During *C. elegans* development, most cell divisions are completed during embryogenesis, but a

subset of somatic and germline cells divide post-embryonically during and beyond the L1 stage,

to generate adult tissues[72-77]. Previous studies have shown that upon dauer entry these cells

maintain or re-establish multipotency to be able to continue development when conditions

become favorable again[78,79], and this is mediated by the known dauer signaling pathways that act

to different extents, in different post-embryonically developing tissues, to modulate their cell cycle

G1/S transition, natural re-programming, and/or differentiation[80-87]. Because of its conserved

functions in promoting reversible cell cycle arrest and stem cell differentiation[19-21,88], we

speculated that CEP-1/p53 activated by the loss of ILC-17.1, was also restraining cell cycle

progression and suppressing adult fates in post-embryonically developing cells when nutrients

were scarce. To test this possibility, we (i) examined the localization of CEP-1/p53 in developing

ilc-17.1 deleted larvae and *cep-1* overexpressing larvae during their dauer entry, (ii) examined

whether CEP-1/p53 activity in *ilc-17.1* deletion mutants (and CEP-1/p53 overexpressing larvae)

altered cell fate or increased the expression of cell cycle inhibitors [89,90], and (iii) examined whether

decreasing the expression of the cell cycle inhibitors that may be upregulated could systemically

rescue dauer arrest.

Previous studies have shown that CEP-1/p53 is expressed in the adult germline, and in a subset

of larval pharyngeal muscle and neuronal cells[60,62]. This has been visualized by other groups

using the same functional, fluorescently tagged CEP-1, CEP-1::GFP which complements the *cep-*

1 deletion, *cep-1*(*lg12501*), where endogenous CEP-1 protein is not expressed[62]. We found that

in addition to these cells, CEP-1/p53 was also expressed in the ventral somatic progenitor blast

cells (P cells) early during larval development (15-20 hours post-hatching; Fig. 4a). The ventral P

blast cells are multipotent progenitor cells that divide post-embryonically during larval stage 1 (L1)

to generate neuroblasts, hypodermal blast cells and vulval precursor cells which later form adult

tissue [72,73,77,91].   These cells express the Paired-box gene *pax*-3, the sole *C. elegans* PAX-3

transcription factor homolog, during these early stages of development[92]. CEP-1::GFP colocalized

with approximately 50% of *pax*-3-driven mCherry that can be used to mark P cells, 15 hours post-

hatching at 20°C, and accumulated in more *pax-3* expressing cells under dauer inducing

conditions of 25°C (Fig. 4a, b). In addition, CEP-1::GFP expression levels in P blast cells

increased at 25°C in larvae overexpressing CEP-1/p53 or lacking ILC-17.1, when compared to

larvae at 20°C, or when compared to CEP-1::GFP expression in *cep-1* (lg12501) larvae that do

not arrest as dauers (Fig. 4c). Notably, CEP-1::GFP expression in blast cells of *cep-1* (lg12501)

larvae did not vary between 20°C and 25°C (Fig. 4c), indicating that increased CEP-1/p53 levels

in blast cells of larvae overexpressing CEP-1/p53, or lacking ILC-17.1, was likely related to their

dauer entry and not due to changes in temperature *per se*. Thus, during dauer arrest of *ilc-17*

deletion mutants, or CEP-1/p53 overexpressing larvae, CEP-1/p53 appeared to be activated in

the ventral P blast cells that would arrest lineage progression as dauers.

Larvae overexpressing CEP-1/p53 and *ilc-17.1* deleted larvae also upregulated cell cycle

inhibitors that are known to be direct or indirect targets of CEP-1/p53[90]: one of the two p21

homologs, *cki-1,* and *phg-1,* the *C. elegans* homologue of Growth arrest-specific 1 or Gas1 were

upregulated (Fig. 4d). *phg-1,* but not *cki-1,* was a direct target of CEP-1/p53, as *ilc-17.1*

deletion mutants displayed increased occupancy of CEP-1/p53 at the promoter of the *phg-1*

gene (Fig. 4e), but not the *cki-1* gene (Supplementary Fig. S7e). The increased expression of

these genes and occupancy at *phg-1* promoter was restored to wildtype levels upon re-

expressing ILC-17.1 (Fig. 4d, e). For technical reasons we could not localize *phg-1* or *cki-1*

mRNA in larvae during these early stages of development and therefore do not know whether

the increased expression occurred in the P blast cells. Nevetheless, the increased expression

of *phg-1* and *cki-1* played a causal role in dauer arrest: downregulation by RNAi was sufficient

to significantly, but modestly rescue dauer arrest in *ilc-17.1* deletion mutants and CEP-1/p53

overexpressing larvae, and promote their development into adults (Supplementary Fig. S7f).

Crossing *ilc-17.1* deletion mutants into *phg-1* deletion mutants, ok2741, resulted in a greater

rescue and ~40% escaped dauer arrest and grew into reproductive adults, but surprisingly, over

60% of the *ilc-17.1;phg-1* double mutants still remained arrested as dauers (Fig. 4f;

Supplementary Fig. S7g), suggesting that CEP-1/p53 activity might be acting in cooperation

with other cell cycle or dauer pathway genes. In addition, a proportion of *ilc-17.1* deletion

mutant and CEP-1/p53 overexpressing larvae that appeared morphologically dauer*,* displayed

modest but significant non-dauer physiological traits on *phg-1* and *cki-1* RNAi such as increased

rates of sporadic pumping (Supplementary Fig. S7h; wildtype larvae at larval stage 2, L2, were

used as controls), suggesting there there may also occur a wider, albeit incomplete, dauer

rescue in some tissues. The dauer arrest of *daf-2* mutant larvae which did not depend on *cep-*

1, was also not rescued upon downregulating *phg-1* or *cki-1*, nor were the pumping rates of *daf-*

2 dauers altered, and these remained even lower than in *ilc-17.1* deleted or CEP-1/p53

overexpressing dauers (Supplementary Fig. S7f, h). These data together showed that cell cycle

inhibitors that were CEP-1/p53-targets were upregulated upon loss of ILC-17.1 signaling or

overexpressing CEP-1/p53 and downregulating these inhibitors was sufficient, at least in part, to

cause a systemic rescue from dauer arrest.

Given these observations, we asked whether overexpressing CEP-1/p53 in P blast cells alone,

under the *pax-3* promoter, might drive larvae to arrest as dauers. This expectation turned out to

be too simplistic, and was not supported, as the majority (85.7 ± 5%; n=6 experiments, 122 larvae

total) of *pax-3p::CEP-1* expressing larvae grew into adults at 25°C, although 14.3 ± 4.5% did not

complete development and arrested at various early larval (<L4) stages. However, CEP-1/p53

expression in the P blast cells caused anteroposterior patterning defects in larvae and 11.8 ±2.9%

of larvae displayed dorsalization, vulvaless and dumpy phenotypes (representative example; Fig.

4g) suggesting that CEP-1/p53 expression could alter the cell fates of the P blast cell lineages[76,93-

95]. Consistent with these observations, in larvae lacking ILC-17.1 or overexpressing CEP-1/p53,

HOM-C homeobox genes required for anterio-posterior body patterning[76,93-95] were differentially

expressed compared to wildtype or *ilc-17.1; cep-1* double mutant larvae (Fig 4h; p< 0.05). This

included *lin*-39 (Scr/Dfd/Hox3–5) which maintains P cells competent to respond to later inductive

signals and prevents aberrant cell fusions, and *mab-5* (Antp/Hox6–8) and *egl*-5 (Abd-A/Abd-

B/Hox9–13), involved in mid- and posterior body patterning, which were maintained at modestly,

but significantly higher levels when compared to larvae continuously developing into adults

(compare expression levels between 15 hr and 32 hr larvae in Supplementary Table 12). Thus,

while further studies are required to better understand how CEP-1/p53 activity in *ilc-17.1* deleted

larvae triggers dauer arrest, these data together indicate that CEP-1/p53-induced increase in cell

cycle inhibitors such as *phg-1* (in P cells or other cells), and its ability to directly or indirectly

modulate cell fate in the ventral P cell lineage during development, where CEP-1/p53

accumulates during dauer arrest, may collaborate to pause growth programs of the animal.

ILC-17.1 links glucose metabolism to CEP-1/p53 activity in progenitor blast cells (P cells)

of larvae.

The release of ILC-17.1 from amphid neurons in response to food suggested the intriguing

possibility that ILC-17.1 signaling linked CEP-1/p53-mediated cell fate decisions to energy

metabolism, since limited nutrient availability, glucose starvation or energy deprivation are all

known to activate p53 in mammalian cells[88,96]. Indeed, the dauer arrest of *ilc-17.1* mutant larvae

appeared related to an impairment in their ability to utilize glucose normally available through their

diets: decreasing glucose availability by downregulating *fgt-1,* the main Glucose transporter

(GLUT) responsible for glucose absorption in *C. elegans*[97], decreased the extent of dauer-rescue

conferred by ILC-17.1 re-expression in *ilc-17.1* mutant larvae (Fig. 5a). Similarly, *fgt-1* RNAi

increased the numbers of *ilc-17.1* deletion mutant larvae that arrested as dauers under optimal

conditions at 20°C (Fig. 5b). Promoting glucose utilization and/or metabolism by providing extra

glucose[98] rescued the dauer arrest of *ilc-17.1* deficient larvae (Fig. 5c; Supplementary Fig. S8a).

This effect of glucose supplementation also depended on glucose utilization and metabolism.

Impairing glucose import through *fgt-1* RNAi decreased the percentage of larvae that were

rescued by the extra glucose (Supplementary Fig. S8b), and no rescue was observed on the non-

hydrolyzable glucose analog 2-Deoxy-d-glucose (2-DOG; Supplementary Fig. S8c) which cannot

undergo glycolysis and does not enter the metabolic pathway. We ruled out the involvement of

mitochondrial reactive oxygen species, ROS, known to be increased under glucose-rich diets[98]

by using mitochondria-targeted antioxidant, Mito-Tempo, that scavenges ROS (Supplementary

Fig. S8c). The rescue also did not appear to be due to the non-specific effect of nutritional

supplementation (e.g. L-glutamine, another important carbon source for metabolism, or skim milk

powder that can be ingested by *C.* elegans[99]; Supplementary Fig. S8d). Downregulating *fgt-1* in

wildtype larvae, while not sufficient to trigger dauer arrest, modestly increased the expression of

CEP-1/p53 target *egl-1* (but not *ced-13*; Supplementary Fig. S8e). Thus, these data supported

the possibility that the activation of CEP-1/p53 upon the loss of ILC-17.1 could, at least in part,

occur due an impairment in the *ilc-17.1* deleted larvae's ability to utilize glucose.

If glucose impairment was responsible for CEP-1/p53 activation, glucose supplementation should

also rescue the dauer arrest of larvae overexpressing CEP-1/p53. This was the case (Fig 5c;

Supplementary Fig. S8a), also arguing that decreased glucose intake in the absence of ILC-17.1

was not a consequence of CEP-1/p53 activation, but instead acted logically upstream of CEP-

1/p53 activation. In agreement, providing extra glucose decreased the levels of CEP-1::GFP that

accumulated in the ventral P blast cells (Fig. 5d), suppressed the increased expression of *phg-1*

the direct target of CEP-1/p53, which was in part responsible for dauer arrest of *ilc-17.1* deleted

animals (Fig. 5e), and decreased DAF-16/FOXO activation (Fig.5e). Surprisingly, not only did

CEP-1/p53 appear to be activated by the inability of *ilc-17.1* to utilize glucose normally, but CEP-

1/p53 activity, in turn decreased the expression levels of key metabolic enzymes involved in

glucose metabolism[100]: phosphofructokinase-1.2 (*pfk-1.2*), the rate limiting enzyme in glycolysis,

and cytochrome *c* (*cyc-2.2*), the subunit of complex IV of the electron transport chain responsible

for the final transfer of reducing equivalents to O_2. This was visible in the RNA-seq data

(Supplementary Figure S9 a-f; Supplementary Table 13) and confirmed by qRT-PCR

(Supplementary Figure S9g, Fig. 5f). In mammalian cells, p53 activity is known to decrease the

expression of enzymes required for glycolysis[88] and indeed, the decrease in mRNA expression

of *pfk-1.2* was a consequence of CEP-1/p53 activation in the *ilc-17.1* deleted larvae: the deletion

of *cep-1* rescued the low mRNA levels (Supplementary Figure S9g). Surprisingly, the decrease

in *cyc-2.2* expression was also dependent on CEP-1/p53*,* but also on DAF-16/FOXO, as the

deletion of *cep-1* or *daf-16* rescued the low *cyc-2.2* mRNA levels in *ilc-17.1* deleted larvae (Fig.

5f). Moreover, although the downregulation of *pfk-1.2* or *cyc-2.2* alone was not sufficient to induce

dauer in wildtype animals, RNAi induced downregulation of *cyc-2.2* in *ilc-17.1* deleted larvae

increased the percentage of larvae that arrested as dauers under optimal conditions at

temperatures of 20°C, suggesting that CEP-1/p53-dependent decrease of *cyc-2.2* could feedback

onto *ilc-17.1* loss to increase dauer propensity (Supplementary Figure S9h).   These data,

together, strongly suggested that the loss of ILC-17.1 resulted in an impairment in glucose

utilization and thus activated CEP-1/p53, which, upon activation, could further limit glucose

utilization by downregulating genes important for glycolysis and OXPHOS. Thus, ILC-17 secretion

from amphid neurons that occurred in response to food availability appeared to link cell fate

decisions with energy metabolism through CEP-1/p53.

ILC-17.1 suppresses CEP-1/p53 in *C. elegans* and human epithelial cells.

Since our data pointed towards a model whereby ILC-17.1 signaling that occurred constitutively,

under favorable conditions in the presence of food, suppressed CEP-1/p53 to promote continuous

development, we directly examined whether IL-17 suppressed CEP-1/p53 activity. As was shown

in B lymphoma cells[101], stimulation of untransformed human epithelial cells with human

recombinant IL-17A also induced a modest but significant downregulation of p53 expression

levels (Fig. 6a). This was similar to the decrease in CEP-1/p53 levels seen in *C. elegans* upon

overexpression of ILC-17.1 (Fig. 3i). For the experiments in mammalian cells, IL-17A stimulation

of epithelial cells was confirmed by measurement of secreted CXCL5, a key cytokine downstream

of IL-17A signaling[102] (Supplementary Fig. S10a).

In *C. elegans* too, increased ILC-17.1 levels suppressed CEP-1/p53 activation: overexpressing

ILC-17.1 in animals also overexpressing CEP-1/p53 inhibited their dauer arrest (Fig. 6b),

inhibited the increase in mRNA levels of CEP-1/p53 target genes, *egl-1*, *ced-13* in CEP-1/p53

overexpressing animals (Fig 3g; Supplementary Fig S7d), *cki-1* and *phg-1* (Fig. 4d), and

partially restored the decreased *cyc-2.2* mRNA levels that resulted from CEP-1/p53

overexpression (Fig 5f). In *C. elegans* the activity of CEP-1/p53 has been best studied in the

adult germline where CEP-1/p53 controls germline apoptosis in response to genotoxic

insults[60,62,67,68]. Therefore, we used gamma irradiation to cause DNA damage in day-one adult

animals, and measured the number of germline apoptotic corpses that result 24 hr. later, to

determine whether increased ILC-17.1 levels could also inhibit this canonical activity of CEP-

1/p53 (Fig 6c). As has been previously shown[68], and consistent with the role of p53 in

controlling DNA-damage induced-, but not physiological apoptosis, *cep-1 (gk138)* adults lacking

functional CEP-1/p53 displayed only a mild decrease in the rates of physiological apoptosis

when compared to wildtype animals (Fig. 6c). However, as also shown previously[68], in these

animals gamma irradiation did not increase apoptosis (Fig. 6c). CEP-1/p53 overexpression,

surprisingly, caused a significant increase in physiological apoptosis, and a marked increase in

irradiation-induced apoptosis, both of which were decreased upon also overexpressing ILC-17.1

in these animals (Fig 6c). Overexpressing *ilc-17.1* alone also decreased both physiological or

irradiation-induced apoptosis compared to that in wildtype animals, but this effect was mild and

did not reach significance (average nos. of cells undergoing physiological apoptosis in N2 v ILC-

17.1 overexpressing animals were 3.56 ± 1.3 and 2.53 ± 1.1 cells, and upon irradiation, these

increased to 7.1 ± 1.8 in N2, and 5.72 ± 2.2 in IL-17 overexpressing animals). Also, for reasons

we do not understand, *ilc-17.1* deletion mutants did not display changes in apoptosis when

compared to wildtype animals, suggesting perhaps that the functions of ILC-17.1 in adult tissue,

and/or germline cells may be more nuanced when compared to its developmental function.

Nevertheless, these data confirmed that, consistent with our working model (Fig. 6d), ILC-17.1

could inhibit CEP-1/p53. In addition, they suggested the possibility that the roles of *ilc-17.1*

signaling and CEP-1/p53 activity might diverge post-development.

Discussion

Here we show, for the first time, a role for the cytokine IL-17, and p53 in the development of an

invertebrate species, *C. elegans*. Specifically, we show that CEP-1/p53 repression during early

larval development ensures the continuous growth of the animal into reproductive adults, and this

repression is mediated by ILC-17.1 released by amphid neurons upon food availability. Reduced

ILC-17.1 signaling, expected to occur when food is scarce, activates CEP-1/p53, which then

triggers dauer arrest through the well characterized dauer activating pathways, DAF-16/FOXO,

DAF-3/SMAD-DAF-5/Ski and also, perhaps also through steroid hormone signaling. In its role in

promoting dauer arrest CEP-1/p53 accumulates in multipotent P blast cells that differentiate

postembryonically into adult tissues, restrains cell cycle progression, and alters cell fate decisions.

In addition, while promoting dauer arrest, CEP-1/p53 alters metabolic flux by directly or indirectly

controlling the levels of phosphofructokinase (*pfk*-1) and cytochrome C (*cyc-2.2*). Notably, these

are functions of p53 in its role as a tumor suppressor. We propose a working model whereby ILC-

17.1 secretion from amphid neurons that occurs in response to food availability links cell fate

decisions with energy metabolism through the conserved, multifaceted program of CEP-1/p53

(Fig. 6d). Our data also support the possibility that this ancient function of p53, conserved in all

multicellular animals and shown here to modulate an invertebrate developmental program, could

have been the driving force in the evolution of the p53 gene-family.

It is intriguing that during normal *C. elegans* development CEP-1/p53 appears to modulate the

fate of multipotent P blast cells. In mammals too, normal p53 activity is essential for maintaining

stem cell quiescence in the hematopoietic system and other organs during steady state tissue

homeostasis[103,104], and unrestrained p53 activity is associated with a developmental syndrome

resembling CHARGE (coloboma, heart defects, atresia choanae, growth retardation, genital

abnormalities, and ear abnormalities), through its effects on cell cycle progression, apoptosis

and the migration of neural crest cells[24]. Cytokines such as IL-17 modulate glucose metabolism

in mammalian models, and glucose deprivation can, in turn, activate p53[88,96]. Thus, as seen in

C. elegans, the ability of p53-proteins to link metabolic control with cell cycle progression and

stem cell fate could be a conserved mechanism leveraged by several signaling pathways similar

to IL-17s to coordinate metazoan development with nutrient availability.

The specific mechanisms by which ILC-17.1/CEP-1 axis promotes dauer entry remain to be better

explored.   In addition, we are yet to dissect the temporal requirements, sites of action, and

mechanisms and consequences of ILC-17.1 signaling required to control CEP-1/p53 activation,

as well as its effects on adult traits such as longevity and stress resistance--traits linked to other

dauer-activating pathways. While the discovery of new *C. elegans* dauer pathway genes is

surprising given the extensive genetic screens that have been conducted[3,5,29-31], genes that

require a complete deletion to confer a dauer phenotype, QTLs that interact in complex ways to

limit the discovery of dauer phenotypes, or genes that show functional redundancy and tolerance

to quantitative variation such as seen with CEP-1/p53 overexpression, could have been missed.

Indeed, a previously characterized *ilc-17.1* mutation *ilc-17.1*(tm5218), generated through EMS

mutagenesis and in the background of a linked mutation in *npr-1,* itself known to be a modifier of

the dauer phenotype, does not trigger dauer arrest under the same conditions (Supplementary

Figure S10b). Intriguingly, our RNA-seq data show that gene expression changes between 15 hr.

post-hatching and 32 hr. post-hatching are negligible in *ilc-17.1* deletion mutants and CEP-1/p53

overexpressing larvae, suggesting that this pathway is one of the very early checkpoints in the

larval decision between continuous growth and dauer arrest[3,4]. Thus, it is plausible that ILC-17.1

and CEP-1/p53 could even act as early as in the maternal germline, or during embryogenesis, to

modulate developmental decisions.

It is interesting that ILC-17.1 is secreted by *C. elegans* sensory neurons to control CEP-1/p53

activity. Neurons control dauer entry also though TGFβ, insulin and steroid hormone signaling. In

addition, other stress-responsive transcriptional programs such as the cellular response to protein

misfolding that are cell autonomous in individual cells are controlled through neuronal signaling

in a metazoan such as *C. elegans*, allowing the nervous system to coordinate tissue-specific

transcriptional and epigenetic responses with organismal physiology and behavior[105-108]. Nutrient

scarcity has shaped much of evolution, and the ability of cells and organisms to sense nutrient

availability and prepare in anticipation to pause cell cycle progression and growth and maintain a

quiescent state until resources are optimal, has unquestionable selective advantages. For *C.*

elegans, like for most organisms, the availability of food is not guaranteed. Neuronal ILC-17.1

signaling appears to be one mechanism which allows the animal to match the cell fate programs

in the developing larvae and metabolic requirements with resource availability. In mammals,

resident neurons and immune molecules in peripheral tissue profoundly influence cell fate

decisions to remodel tissue architecture during development, organogenesis, inflammatory

responses, and in diseases such as cancer[109,110]. It is therefore not far-fetched to imagine that

mammalian neurons control p53 activity, cell quiescence and stem cell fate in response to nutrient

availability through IL-17s, as seen in *C. elegans*, although, perhaps in mammals this occurs

locally through peripheral innervation and immune cells. Such a control mechanism could open

new avenues for therapeutic intervention in cancer and other IL-17-mediated diseases.

Materials and Methods

C. elegans **strains**

C. elegans strains used in this study are listed in Table 1. Strains were procured from

Caenorhabditis Genetics Center (CGC, Twin Cities, MN), generated in the laboratory or

generated by Suny Biotech (Suzhou, Jiangsu, China 215028). All strains will be available upon

request, after publication.

Strain Name	Gene name	Source	Additional information
N2, *C. elegans* var *Bristol*	N2, Wild-type	Caenorhabditis Genetics Center	
TJ1	*cep-1* (gk138) I	Caenorhabditis Genetics Center	
CF1038	*daf-16* (mu86) I	Caenorhabditis Genetics Center	
TG12	*cep-1*(lg12501) I*; unc-119* (ed4) III; gtIs1 [CEP-1::GFP + *unc-119* (+)]	Caenorhabditis Genetics Center	Pcep-1::CEP-1(let-585 3'UTR)::GFP; from construct pRH53
CB1370	*daf-2* (e1370) III	Caenorhabditis Genetics Center	
VEP024	praEx021 (*ilc-17.1*p::*ilc-17.1* cDNA::3XFLAG::*ilc-17.1* 3'UTR; p*myo-2*::mCherry::*unc-54* 3'UTR)	Prahlad Lab	ILC-17.1 overexpression under its own promoter and 3'UTR

VEP031	praEx022 (*unc-54*p::*ilc-17.1* cDNA::3XFLAG::*tbb-2* 3'UTR; p*dat-1*::GFP::*unc-54* 3'UTR)	Prahlad Lab	ILC-17.1 overexpression under the muscle promoter and 3'UTR
VEP032	*ilc-17.1* (syb5296) X	Prahlad Lab/ SunyBiotech	*ilc-17.1* deletion, 2173bp deletion, and the 15bp and 127bp sequences were left in the 5' and 3' deletion end, respectively of the 2315 bp *ilc-17.1* gene
VEP033	*ilc-17.1* (syb5296) X; praEx022 (*unc-54*p::*ilc-17.1* cDNA::3XFLAG ::*tbb-2* 3'UTR; p*dat-1*::GFP::*unc-54* 3'UTR)	Prahlad Lab	ILC-17.1 paracrine rescue of *ilc-17.1* deletion
VEP034	*ilc-17.1* (syb5296)X; *cep-1* (gk138) I	Prahlad Lab	
VEP035	*ilc-17.1* (syb5296); *daf-16* (mu86) I	Prahlad Lab	
VEP036	*unc-119* (ed4); gtIs1 [CEP-1::GFP + *unc-119* (+)]	Prahlad Lab	CEP-1 overexpression (ref: [60])
VEP037	*ilc-17.1* (syb5296); praEx022 (*unc-54*p::*ilc-17.1* cDNA:: 3XFLAG::*tbb-2* 3'UTR; p*dat-1*::GFP::*unc-54* 3'UTR);	Prahlad Lab	IL-17.1 overexpression;

	unc-119 (ed4); gtIs1 [CEP-1::GFP + *unc-119* (+)]		CEP-1/p53 overexpression
VEP038	*ilc-17.1* (syb5296); praEx022 (*unc-54*p::*ilc-17.1* cDNA:: 3XFLAG::*tbb-2* 3'UTR; p*dat-1*::GFP::*unc-54* 3'UTR); *daf-2* (e1370) III	Prahlad Lab	
VEP039	*daf-2* (e1370) III; *cep-1* (gk138) I	Prahlad Lab	
VEP040	*cep-1* ((syb6099) [*cep-1*::3XFLAG]) I	Prahlad Lab/ SunyBiotech	Endogenous *cep-1* 3X FLAG tagged at C-terminus
VEP041	*ilc-17.1* (syb5296)X ; *cep-1* (pra02 [*cep-1*::3XFLAG]) I	Prahlad Lab	
VEP042	praEx022 (*unc-54*p::*ilc-17.1* cDNA::3XFLAG::*tbb-2* 3'UTR; p*dat-1*::GFP::*unc-54* 3'UTR); *ilc-17.1* (syb5296) X ; *cep-1* (pra02 [cep-1::3XFLAG]) I	Prahlad Lab	
VEP043	*ilc-17.1* (pra03 [*ilc-17.1*::SL2::mCherry]) X	Prahlad Lab/ SunyBiotech	
VEP044	*ilc-17.1* (pra04 [*ilc-17.1*::3xHA]) X	Prahlad Lab/ SunyBiotech	
VEP045	*ilcr-2* (pra05 [*ilcr-2*::SL2::GFP]) II	Prahlad Lab/ SunyBiotech	
CB1386	daf-5(e1386) II	Caenorhabditis Genetics Center	

VEP061	*ilc-17.1* (syb5297)	Prahlad Lab/ SunyBiotech	*ilc-17.1* deletion, 2188bp deletion, and the 9bp and 118bp sequences were left in the 5' and 3' deletion end, respectively of the 2315 bp *ilc-17.1* gene
RW10645	stIs10645 [pax-3p::HIS-24::mCherry + *unc-119*(+); *unc-119* (ed3)	Caenorhabditis Genetics Center	
VEP062	stIs10645 [pax-3p::HIS-24::mCherry + *unc-119*(+); *unc-119* (ed3); *ilc-17.1*(syb5296)	Prahlad Lab	
VEP059	stIs10645 [pax-3p::HIS-24::mCherry + *unc-119*(+); *unc-119* (ed3); gtIs1[cep-1::GFP + *unc-119*(+)]	Prahlad Lab	
VEP063	praEx026(*pax-3*p::CEP-1A cDNA::3xFLAG::*unc-54* UTR; p*myo-2*::mCherry::*unc-54* UTR)	Prahlad Lab	
AX5690	npr-1(ad609)X; ilc-17.1(tm5218)X	Mario de Bono's lab/ C. elegans National Bioresource Project, Japan	

608 **Generation of CRISPR strains**

609 CRISPR/Cas9 was used to create following *C. elegans* strains:

Background Strain	Type of editing	Description/position of editing	Gene	Resulting strain
N2, *C. elegans* var Bristol	Deletion (syb5296)	2173bp deletion, and the 15bp and 127bp sequences were left in the 5' and 3' deletion end, respectively of the 2135 bp *ilc-17.1* gene. **sgRNA used - <u>CCA</u>AAATCACCACACAGGACA AA**	(*ilc-17.1*)X	VEP032
N2, *C. elegans* var Bristol	Deletion (syb5297)	2188bp deletion, and the 9bp and 118bp sequences were left in the 5' and 3' deletion end, respectively of the 2135 bp *ilc-17.1* gene. **sgRNA used- tagGCAAATGCGAATGCGGA<u>TG G</u>**	(*ilc-17.1*)X	VEP032
N2, *C. elegans* var Bristol	Insertion; *SL2::mCherry*	C'-terminus	(*ilc-17.1*)X	VEP043
N2, *C. elegans* var Bristol	Insertion; 3X HA	C'-terminus	(*ilc-17.1*)X	VEP044
N2, *C. elegans* var Bristol	Insertion; SL2::GFP	C'-terminus	(*ilcr-2*)II	VEP045

N2, *C. elegans* var Bristol	Insertion; 3X FLAG	C'-terminus	(*cep-1*)I	VEP040
N2, *C. elegans* var Bristol	Deletion (1475bp of 1692 bp)	Prahlad Lab/ SunyBiotech	F25D1.3 (syb7367) V	PHX7367

Generation of transgenic strains

i)   Backcrossing of *ilc-17.1* (syb5296) X

The *ilc-17.1 (syb5296)* X strain was crossed to wildtype (N2) males, and the the male F1 progeny which harbored the *ilc-17.1 (syb5296)* allele on their X chromosome were backcrossed to the *ilc-17.1 (syb5296)* X strain and homozygous F2s were selected by PCR. This procedure was conducted 2X times to generate the 2X backcrossed *ilc-17.1 (syb5296)* X line.

ii)   Generation of *cep-1* overexpression strain

TG12 (*cep-1*(lg12501) I; *unc-119* (ed4) III; [CEP-1::GFP + *unc-119* (+)]) is a strain that expresses a functional CEP-1 tagged with GFP integrated into its genome, as determined by the rescue of *cep-1*(lg12501) phenotype[62]. To overexpress CEP-1 we backcrossed the TG12 strain with wild-type (N2) worms and selected F2 progeny that were homozygous for the wild-type *cep-1* gene, lacked the *cep-1*(lg12501) I mutation (confirmed by PCR), and were homozygous for the CEP-1::GFP transgene (confirmed by 100% GFP expression amongst the F3 and F4 progeny). Overexpression of cep-1 mRNA was verified by qPCR (Supplementary Figure S6d). The CEP-1::GFP construct was PCR amplified from the final transgenic VEP037 *C. elegans* strain, and sequence verified.

iii)   Generation of ILC-17.1 overexpression strain under its own promoter and 3'UTR

To generate the strain VEP024 – praEx021 [(*ilc-17.1*p::*ilc-17.1* (cDNA)::3xFLAG::*ilc-17.1* 3'UTR); p*myo-2*::mCherry::*unc-54* 3'UTR], which overexpressed *ilc-17.1* under the endogenous *ilc-17.1* promoter and *ilc-17.1* 3'UTR, we first amplified 3 kb of genomic sequence upstream of

the translational start site (*ilc-17.1* promotor) and 1 kb of genomic sequence downstream from

the translational stop codon (*ilc-17.1* 3' UTR). These regions were cloned using Gateway

technology into pDONR221. *ilc-17.1* cDNA fused to C-terminal 3xFLAG sequence was

synthesized as a gBlock (Integrated DNA technologies). These three fragments were then

submitted for gene synthesis service through GenScript to generate the expression vector

pUC57(*ilc-17.1*p:: *ilc-17.1* (cDNA)::3xFLAG:: *ilc-17.1* 3'UTR). All plasmids were sequence

verified. The *ilc-17.1* expression vector was then injected at 97.5 ng/ul along with the co-

injection marker pCFJ90 (p*myo-2*::mCherry::*unc-54* 3'UTR) at 2.5 ng/ul by InVivo Biosystems

injection express service. Animals expressing mCherry were singled, lines transmitting the

extrachromosomal array were established, and mCherry positive progeny were PCR verified to

ensure they were transmitting *ilc-17.1*p:: *ilc-17.1* (cDNA)::3xFLAG:: *ilc-17.1* 3'UTR. These lines

were then harvested for Western blot to verify expression of protein.

iv)   **Generation of ILC-17.1 overexpression strain under the *unc-54* muscle promoter**

644   **and *tbb-2* 3'UTR**

To generate VEP031, we overexpressed ILC-17.1 under muscle promoter by fusing - [(*unc-*

*54*p:: *ilc-17.1* (cDNA)::3xFLAG] which was synthesized as a gBlock (Integrated DNA

technologies) and *tbb-2* 3' UTR which was amplified from genomic DNA. The two fragments

were cloned into pUC19 plasmid as the backbone using Gibson assembly[111] to create the

overexpression plasmid [(*unc-54*p:: *ilc-17.1* (cDNA)::3xFLAG::*tbb-2* 3'UTR); p*dat-1*::ssmito] and

sequence verified. The construct was injected at 100 ng/ul along with the co-injection marker

that expressed GFP in the dopaminergic neurons under a *dat-1* promoter [p*dat-1*], at 10 ng/ul

into wildtype (N2) worms. Animals expressing GFP in their dopaminergic neurons were singled,

lines were established, and GFP-positive progeny were PCR verified to ensure that they were

also transmitting harbored the *ilc-17.1* transgene. These lines were then harvested for Western

blot to verify expression of protein.

v)   **Generation of CEP-1 overexpression strain under the *pax-3* P-cell promoter and**

***unc-54* 3'UTR**

To generate the strain VEP063 – praEx026(*pax-3*p::CEP-1A cDNA::3xFLAG::*unc-54* UTR;

p*myo-2*::mCherry::*unc-54* UTR)], which overexpressed CEP-1 under the P-cell promotor *pax-3*

and *unc-54* 3'UTR, we first amplified 2655 bp of genomic sequence upstream of the

translational start site (*pax-3* promotor), 1932 bp of CEP-1A full length cDNA sequence without

a stop codon from N2 cDNA, and 851 bp of sequence was amplified from vector pDONRP2r-

P3(3xFLAG::*unc-54* 3'UTR). These three sequences were then cloned using a BP reaction into

the appropriate DONR vectors for the Multisite Gateway Pro cloning kit (Invitrogen) according to

manufacturer's guidelines. An expression vector was created by performing the multisite LR

reaction using pDEST R1-R2 with the three entry clones generated above to generate pDEST

R1-R2 (*pax-3*p::CEP-1A::3xFLAG::*unc-54* 3' UTR) according to manufacturer's instructions. All

plasmids were sequence verified. The CEP-1 expression vector was then injected at 120 ng/ul

along with the co-injection marker pCFJ90 (p*myo-2*::mCherry::*unc-54* 3'UTR) at 5 ng/ul into N2

wildtype animals. Animals expressing mCherry were singled, lines transmitting the

extrachromosomal array were established, and mCherry positive progeny were then harvested

for Western blot to verify expression of protein.

vi)   **Generation of crosses.**

Crosses between strains were generated according to standard procedures. The primers used to

genotype strains are included in the table below.

	Forward primer	Reverse primer
VEP032	TGAATGAGCAAATCGACCAG	AGTTTTGAGATTGAGGGCAAA
	CAGGTCAAGGGGCTAAAACA	CCCACATTAATACTCGCACCT
CF1038	CGGATCAGGAAGAACCAGAA	CATTCCCCGAGTACGAGCTA
	AAGCTGCTGCCTTCACTCTC	ATGGGACTCGAGTTGAGTGG

TJ1	CGGTCTTTTTGGACGATGAA	GTAAACATTGCCGCGTTTTT
	GCGATTCGAAATCCAAGAAG	AGGAACGCATCCATGAAGAC
TJ12	GCCTTCCTCTATTCACTTGAAAC	ACCGAGAGCTCCATGAAGAA
	TCACTCTTTGAATTGGTTCTTTTG	GCTTACGTTCGATTGGCTTC
	AAAAACGCGGCAATGTTTAC	GAGCTTTGTGGGAAGCAATC
RB2075	GACAGTTACGCTGTGCTTGG	CTCCCGGGGTATTCTGAAAT
	GGTGATTCTCCCGTTGGTAA	AAACGGAAATGGCAGAGAGA
CB1370	TAA ATC TCG GAA TTC TGG TGA A	GGA TGA GAC TGT CAA GAT TGG A
	AAG ATG AAC GAG CTG GAG GA	AGA TTG GAG ATT TCG GAA TGG
VEP040	AGA ATT CGA GAA ACA TAA TGA CAT CTT	AAA AAT GCG ATT ATA AAA AGG AAC A
CB1386	CGG CTT GAT CGG ATA GTG AT	AAG AGC AAG GAG TGC GAA AA
	Wild-type PCR product was digested by BstNI	

Antibodies used:Growth conditions for *C. elegans* strains

All strains were grown and maintained at 20°C unless otherwise mentioned. Animals were grown

in 20°C incubators (humidity controlled) on 60mm nematode growth media (NGM) plates by

passaging 8-15 L4s (depending on the strain) onto a fresh plates. Extra care was taken to ensure

equal worm densities across all strains. Animals were fed *Escherichia coli* OP50 obtained from

Caenorhabditis Genetics Center (CGC) that were seeded (OD$_{600}$=1.5 and this was strictly

maintained throughout the experiments) onto culture plates 2 days before use. The NGM plate

thickness was controlled by pouring 8.9ml of autoclaved liquid NGM per 60mm plate. Laboratory

temperature was maintained at 20°C and monitored throughout. For all experiments, age-

matched day-one hermaphrodites, or larvae timed to reach specific developmental stages as

mentioned in the figure legend, were used.

For all experiments with larvae, the timings were adjusted to account for the differences in growth

rates at different temperatures. Thus, at 25°C, larvae were harvested 30-36 hrs. post bleach hatch

or egg lay, and at 20°C, larvae were harvested 40 hrs. post bleach hatch or egg lay.

Obtaining synchronized embryos by 'bleach-hatching'

Bleach-hatching' was performed as previously described[105]. Populations of 250-300 gravid adults

were generated by passaging L4s, as described above. These plates were used for obtaining

synchronized embryos by bleach-induced solubilization of the adults. Specifically, animals were

washed off the plates with 1X PBS and pelleted by centrifuging at 2665Xg for 30s. The PBS was

removed carefully, and worms were gently vortexed in the presence of bleaching solution [250μl

1N NaOH, 200μl standard (regular) bleach and 550μl sterile water] until all the worm bodies had

dissolved (approximately 5-6 minutes), and only eggs were viable. The eggs were pelleted by

centrifugation (2665Xg for 45s), bleaching solution was carefully removed and then embryos were

washed with sterile water 3-4 times and counted under the microscope. The desired number of

embryos were seeded on fresh OP50 plates and allowed to grow at 20°C or 25°C for specific time

periods depending on the experimental need. If >5% eggs remained unhatched, these plates

were discarded.

Dauer Assay

Embryos were allowed to hatch and grow on OP50 plates at 20°C or 25°C for this assay. Embryos

were generated according to one of the following two methods: (i) Day-one gravid adults that had

grown under normal culture conditions (on OP50 at 20°C) were bleach dissolved as described

above, and ~50-100 embryos obtained from these gravid adults were seeded on fresh OP50

plates or (ii) day-one gravid adults were allowed to lay eggs on fresh OP50 (or RNAi) plates for

2-4 hours at 20°C, the adults were removed, and then embryos were allowed to develop at 20°C

or 25°C under humified condition for 48-72 hrs. The former, bleach-synchronization method was

used for RNA-seq method and experiments requiring large numbers of age-matched larvae. The

latter, timed-egg lay method was used for all other exeriments. For each of the strains and

experiments, when we had to use bleach synchronization, we first ensured that bleaching alone

did not alter the percentage of dauers or gene expression profiles by conducting a pilot experiment

using both methods side-by-side. This was the case for all experiments, except experiments

where glucose rescued the dauer phenotype. Here, although the trends were the same, and

glucose resulted in a significant rescue of ilc-17.1 deletion mutants and cep-1 OE larvae that were

generated either by bleach-synchronization or timed-egg lay, bleach synchronization caused a

lower percentage to be rescued from dauer arrest. To assess dauer entry, the number of larvae

that arrested as dauers (determined by phenotype and/or resistance to 1%SDS) and those that

developed into L4s or adults were counted and percentage of dauers was calculated. The total

number of embryos in each plate was ~50-100. Each experiment was performed in triplicate or

more.

For the *ilc-17.1* deletion mutants, in order to acertain that the dauer phenotype was not simply

due to the presence of a background mutation, we also conducted the dauer assay after

backcrossing the *ilc-17.1* (syb5296) X animals with wild-type Bristol N2 animals, 2X times. The

backcrossed strains also arrested as dauers under that same conditions and 100 ± 0% larvae

(n=3 independent repeats, 212 larvae scored), arrested as dauers at 25°C as determined by SDS

resistance compared to 0% of wildtype larvae.

SDS treatment

Larvae were washed off the dauer assay plates, washed 1X with PBS to remove bacteria, treated

with 1% sodium dodecyl sulfate (SDS) for 30 minutes, washed again with M9, and then transferred

onto a fresh plate. Larvae could be scored easily as dead/dissolved or as live dauer larvae (based

on their phenotype, and movement).

RNAi mediated downregulation of genes

All RNAi clones used were verified by sequencing, and plates were seeded with RNAi bacteria

for a maximum of two days before being used. Day-one gravid adults grown on OP50 were

allowed to lay eggs on RNAi-bacteria seeded plates for 2-4 hours at 20°C. A second generation

of RNAi knockdown was performed where necessary by transferring L4's on existing RNAi

plates to fresh plates for knockdown in their progeny. The adults were allowed lay eggs and

then removed, and the embryos were transferred to 25°C (or kept at 20°C) under humified

condition for 48-72 hrs. The number of larvae that arrested as dauers (determined by phenotype

and/or resistance to 1%SDS) and those that developed into L4s or adults were counted and

percentage of dauers was calculated.

Exposure to food [OP50] availability

Gravid day-one *ilc-17.1*(pra03 [*ilc-17.1*::SL2::mCherry]) X adults or *ilc-17.1*(pra04 [*ilc-*

17.1::3xHA]) X adults were bleach dissolved and embryos were placed either on empty plates

with no food or NGM plates seeded with OP50 and allowed to grow for 24-36 hours at 25°C. To

observe mRNA expression of *ilc-17.1, ilc-17.1*(pra03 [*ilc-17. 1:*SL2::mCherry]) X larvae were

anesthetized with 1mM levamisole, mounted on 1% agarose pads, and imaged using a

Confocal SPE microscope using optimal settings. mCherry expression served to mark cells that

expressed *ilc-17.1* mRNA. To identify where ILC-17.1 protein was expressed prior to food

exposure, and after food exposure, 24-36 hour *ilc-17.1*(pra04 [*ilc-17.1*::3xHA]) X larvae were

fixed and immunostained using Bouin's Tube Fixation method (see below).

Immunofluorescence

Nematodes were fixed and immunostained using a modification of the Bouin's Tube Fixation

method [112]. Worms were fixed for 30 min. at room temperature (RT) in 400 µl Bouin's fix (Sigma

Aldrich) + 600 µl methanol and 10 µl β-mercaptoethanol by tumbling, freeze cracked three times

in liquid nitrogen and again tumbled for 30 mins at RT. For permeabilization, the fixative was

removed and exchanged for borate-Triton-βmercaptoethanol (BTB: 1xBorate Buffer, 0.5% Triton

and 2% β-mercaptoethanol) solution. Worms were tumbled 3 times for 1 hour each in fresh BTB

solution at RT. Worms were washed with PBS-0.05% Tween and incubated in block-solution

(5% BSA). Staining with primary antibody was performed overnight at 4 °C, incubation with

secondary antibody (Donkey anti-Mouse Alexa Fluor 488, 1:2000) for 2-4 hours at RT. All

antibody dilutions were performed using antibody buffer containing 5% BSA. Samples were

mounted onto glass slides using VECTASHIELD antifade mounting medium (Vector

Laboratories, Burlingame, CA, USA). Primary antibodies used: (i) Monoclonal ANTI-FLAG M2,

Sigma Aldrich, 1:500, (ii) Mouse anti-HA antibody, Thermofisher; 2-2.2.14, at 1:1000), (iii) anti β-

Actin, Cell Signaling Technology, #4967, 1:1000.

Chemotaxis

Chemotaxis assay was performed on 9 cm petri dishes containing NGM. Two marks were made

on the back of the plate at opposite sides of the plate about 0.5 cm from the edge of the agar.

About 5 µl of attractant diluted in water was placed on the agar over one mark, and 5 µl of water

was placed as the control over the opposite mark. Attractants and concentrations used are

listed in drugs and metabolites section. 5 ul sodium azide with the concentration of 1M was also

placed at both the attractant source and the control source. This drug could anesthetize animals

within about a 0.5-cm radius of the attractant. Age synchronized day-one adult worms were

transferred to the middle of the NGM plates at a point equidistant from the middle of each

odorant. After 1 hr., the assay was quantified by counting the number of worms that had left the

center origin a chemotaxis index was calculated [#Odor - #Control] / [#Odor + #Control]. Each

repeat consisted of 50 worms, and experiments were repeated a minimum of three times.

Measuring Pharyngeal pumping

Synchronized day-one adults were singled onto NGM plates seeded with OP50, and onto plates

without OP50. Pharyngeal pumping rates were determined by recording the pharyngeal region of

animals by video using a Leica S9i digital stereo microscope at 5X magnification and slowing

down the video to manually count the number of 'pumps' in 10 seconds, three times per animal.

The mean of these pumps was determined and the number of pumps/minute calculated. One

complete cycle of synchronous contraction and relaxation of the corpus and the terminal bulb was

counted as a pump.

Feeding of fluorescent latex beads

Overnight 5ml OP50 culture LB was pelleted and resuspended in fresh 0.5ml LB to concentrate

bacteria. 1 µl of fluorescent beads of 0.5µm mean particle size that mimic size of *E. coli*. (Sigma

L3280, red fluorescence) was added to 1 ml of concentrated OP50 [1:1000 ratio (v/v)], and 100ul

of the mixture was seeded onto NGM plates [113]. Synchronized day-one adults were allowed to lay

eggs on the bead containing plates for 2 hours at 20°C and then removed. After 40 hours, larvae

were picked onto 1% agarose pads on glass slides and anesthetized using 10mM levamisole. Z-

stack images of the larvae were taken on Leica TCS SPE confocal microscope and number of

beads within a set area near the tail region per larvae was quantified.

Drug and metabolite treatment

812 The following drugs or metabolites were used:

Drug/metabolite	Source	Final concentration

Glucose	Research products International	50/100mM
2-Deoxy-d-glucose	R&D systems	50mM
Mitotempo	Millipore sigma	0.1mM
L-Glutamine	Sigma-Aldrich	0.26mM
Skim Milk powder	HyVee - instant nonfat dry milk	20mM
Diacetyl	Sigma-Aldrich	0.01% v/v
Lysine	Sigma-Aldrich	1M
(25S)-Δ7-Dafachronic Acid	Cayman Chemical	50nM
8-Bromo-cGMP	Cayman Chemical	5mM

The drug solution (0.5 ml) was spread onto NGM agar plates containing an OP50 lawn and left

for 1 hr. to dry or used for chemotaxis assays as described. Age synchronized day-one adult

worms were transferred to the middle of the NGM plates for 4 hrs. and dauer assay were

performed as described. Sterile water was used as control.

8-Bromo-cGMP solution was first added to standard NGM plates and allowed to dry for 1 hour.

1.5 OD op50 was added on top and allowed to grow for 1 day before starting the dauer assay as

described.

Plates containing D-glucose were prepared as described in published methods [98], except

previously except that we seeded live OP50 onto Glocose plates. Stock solutions of 0.1 and 1M

Sterile D-glucose was prepared and added into NGM agar to obtain final concentration of 5mM

and 50mM, respectively. For control plate, equivalent amount of sterile water was added to

NGM agar. The plates were left to dry for 2 days at RT before seeding 1.5 OD OP50 onto them.

After drying at RT for another 1day, dauer assays were performed as described.

Fluorescence Image analysis

ImageJ (Fiji) v1.53i was used for measuring fluorescence intensity as follows: single planes [,

DAF-16::GFP, CEP-1::GFP, TG12] or projections of z-planes [DAF-28, INS-4, fluorescent latex

beads,pax-3p::mCherry] were used for measuring fluorescence intensity. The region of interest

was circled using the circle selection tool. The mean fluorescence intensity of each circled area

was recorded using the measurement tool. When appropriate, the background fluorescence

intensity was measured and subtracted. Quantification of measurements was done in Microsoft

excel/GraphPad.

RNA-sequencing and Data analysis

a) **RNA isolation, library preparation and sequencing**

838   For the Total RNA-seq samples: Day-one adult worms were bleach-hatched and ~3200

839   eggs/genotype (~800 eggs/plate and 4 plates/genotype) were seeded on fresh OP50

840   plates and allowed to grow for 30-34 hrs. at 25°C. Worms were washed with sterile water

841   and total RNA was extracted from biological triplicates using the Direct-zol RNA Miniprep

842   Kits (catalog no. R2050, Zymo Research). Libraries were prepared using the Illumina

843   Stranded Total RNA Prep

844   with Ribo-Zero Plus, for rRNA depletion (catalog no. 20040525, Illumina). Samples

845   were sequenced in one lane of the Illumina NovaSeq 6000, generating 2x150bp paired-

846   end reads.

847   For the mRNA-seq samples : Day-one adult worms were bleach-hatched and ~3200

848   eggs/genotype (~800 eggs/plate and 4 plates/genotype) were seeded on fresh OP50

849   plates and allowed to grow for either 15 hours or 32 hours at 25°C. Worms were washed

850   with sterile water and total RNA was extracted from biological triplicates using the Direct-

851   zol RNA Miniprep. Libraries were prepared using Illumina TruSeq Stranded mRNA kit.

Samples were sequenced in one lane of the Illumina NovaSeq 6000, generating 2x100bp

paired-end reads.

b) RNA-seq analysis

The quality of the RNA sequences was assessed with FastQC. Adapters and sequence

reads with a quality lower than Q25 were trimmed by using Trimgalore [114](v0.67).

Ribosomal RNA (rRNA) contamination was filtered with sortmeRNA[115] (version 4.3.4).

Sequencing reads were aligned to the *C. elegans* genome (WBcel235[116]) using the Star

aligner[117] (v2.7.9a) and then the aligned reads were quantified with FeatureCounts

v2.0.1 from the R package Rsubread[118] . Differential expression analysis was performed

using DESeq2[119] and genes with FDR corrected p-value of < 0.05 were considered

significant. Pairwise distance analysis (sample-to-sample) was performed by using

normalized counts coupled with the variance stabilization transformation (VST) on the

complete set of genes and calculating the Euclidean distance between the replicates.

Principal Component Analysis (PCA) was done using the plotPCA function in DESeq2[119]

with the VST transformed data. The mRNA-seq was analyzed following a similar

protocol as described above, excluding the SortmeRNA step.

The microarray data for the *C. elegans* dauer stage was obtained from the

Supplementary Table 1 of the study by Wang, J. & Kim (2003)[37]. The expression values

of the dauer stage were defined as the dauers expression (fold-change) at 0hr relative to

the mixed staged reference RNA. Sequences IDs from the array were converted to

standard gene nomenclature using Simplemine in Wormbase[120]. The correlation

between the expression at the dauer stage and the fold change in expression in the *ilc-*

17.1 deletion mutant relative to the wild type, was determined by performing a Spearman

Rank Test.

The expression of C. elegans larvae (L1) treated with ascaroside cocktail was obtained

from Cohen, et al. [121]. The correlation between the Fold-Change in expression in the

larvae treated with ascaroside relative to N2 larvae and the fold change in expression in

the *ilc-17.1* deletion mutant relative to the wild type, was determined by performing a

Spearman Rank Test. The differentially expressed genes (DEGs) between N2 and daf-

1(m40) was obtained from. Hu, et al. [122] and were presented as Log_2(Fold-Change)

values. The overlap between differentially expressed genes (DEGs) in the cep-1

overexpression mutant and the *ilc-17.1* deletion mutant was tested using Fisher exact

test from the Gene Overlap package[123]. Venn diagram was done using the R package

VennDiagram[124].

c) Functional analysis

Gene ontology analysis (GO) and KEGG enrichment analysis was performed on the

differentially expressed genes by using ClusterProfiler 4.0[125]. Terms with an FDR

corrected p-value <0.05 were considered significant. The annotations were obtained

from R package org.Ce.eg.db[126] (version 3.14) and the KEGG database[127]. The

annotations of genes as part of the metabolic pathways were obtained from the

database Wormpaths[100] and heatmaps were done using the R packages

Complexheatmap and pheatmap[128,129].

Progenitor blast cell (P cells) imaging and quantification.

Synchronized day-one adults were allowed to lay eggs on OP50 plates for 2 hours at 20°C and

then removed. The plates were then placed at 20C or 25°C. After 15 hours, about 10 L1 larvae

were picked onto 1% agarose pads on glass slides and anesthetized using 10mM levamisole and

imaged on a Leica SP8. Cells expressing pax-3p::mCherry and CEP-1::GFP were scored, and

the percent pax-3p::mCherry cells that also expressed CEP-1::GFP was calculated for individual

worms.

DiO dye filling assay to identify amphid neurons

A stock dye solution containing 2mg/ml DiO (Molecular Probes, catalog # D-275) in dimethyl

formamide is made and maintained under dark conditions. Day-one adult worms were

transferred into an eppendorf tube with 1 ml of M9, spun down at 2000-3000 rpm and

supernatant removed. Worms were resuspended in 1 ml of M9 and 5 microliter DiO stock sol

(1:200 dilution) was added and incubated on a slow shaker for 2 hours at room temperature.

Worms were spun down and washed with M9 twice before transferring them onto agar pads

with 1mM levamisole to visualize by fluorescence using a Leica TCS SPE confocal microscope.

Dauer pharyngeal pumping

Synchronized day-one adults were allowed to lay eggs on OP50 plates for 2 hours at 20°C and

then removed. The plates were then placed at 25°C, and larvae were allowed to arrest as dauers

or grow into L4 larvae (RNAi treatment) . On day 3, larvae that remained arrested as dauers were

used to measure pharyngeal pumping rates using Zeiss Axio Observer.A1 at 40X magnification.

WT (N2) L2 stage larvae were used to measure pumping rates as controls. Pumping rates were

measured by counting the number of grinder movements in the terminal bulb per minute. The

average of three biological replicates consisting of 10 dauer larvae each was quantified.

Scoring apoptotic nuclei with Acridine Orange

Acridine orange (AO) staining was performed as previously described (Hefel et al. 2021). AO

staining to detect apoptotic germline cells was performed by picking 50 L4 animals to a fresh

plate and allowing them to grow into day-one adults for 24 hours at 20°C. Day 1 adults were

then either exposed to 75 Gy from a Cesium source or left unirradiated. 24 hours later, AO

staining was performed.. For each trial, 10 mg/ml AO stain was freshly prepared and then

diluted 1:400 with M9 buffer. Worms were then picked to a tube of the diluted AO stain, wrapped

tightly with foil, and rotated at room temperature for 1 hour. After mixing, worms were

transferred to a fresh OP50 plate and allowed to crawl away from residual AO stain for 1 hour

before being picked to a droplet of 10 mM levamisole on a 1% agarose pad. A cover slip was

added, and worms were imaged immediately using the Leica fluorescence microscope on 60x

magnification. It was important to visualize AO within 10 mins after mounting animals on the

pad. The number of AO stain positive cells found in 12 individual gonad arms per experiment

were recorded for each sample.

Gamma irradiating *C. elegans* to activate CEP-1/p53

100 L4 larvae were picked onto an OP50 seeded plate, allowed to mature into day-one adults

and exposed to 75 Gy gamma irradiation, and subsequently harvested for Western blot

analysis. To determine the number of AO positive apoptotic nuclei, at least 50 L4 larvae were

exposed to 75 Gy gamma irradiation. The number of AO positive cells were scored 24 hrs. later,

when animals had matured into day-one adults.

Western blot analysis

i.   **Western blot analysis of *C. elegans***

Western blot analysis was performed with 30–100 adult day-one animals, or approximately 400

30-36 hr. larvae as indicated. Animals were harvested in 15 µl of 1X PBS (pH 7.4), and 4X

Laemmli sample buffer (catalog no. 1610737, Bio-Rad) supplemented with 10% β-

mercaptoethanol was added and samples were boiled for 30 min. Whole-worm lysates were

resolved on 12% SDS-PAGE gels and transferred onto nitrocellulose membrane (catalog no.

1620115, Bio-Rad). Membranes were blocked with Odyssey Blocking Buffer (part no. 927–50000,

LI-COR). Immunoblots were imaged using LI-COR Odyssey Infrared Imaging System (LI-COR

Biotechnology, Lincoln, NE). Mouse anti-FLAG M2 antibody (catalog no. F1804,

RRID:AB_262044,_Sigma Aldrich) was used at 1:500 to detect CEP-1::FLAG. Mouse anti-HA

antibody (Thermofisher; 2-2.2.14), was used at 1:1000 to detect ILC-17.1::HA. Mouse anti-α-

tubulin primary antibody (AA4.3, RRID:AB_579793), developed by C. Walsh, was obtained from

the Developmental Studies Hybridoma Bank (DSHB), created by the National Institute of Child

Health and Human Development (NICHD) of the National Institute of Health (NIH), and

maintained at the Department of Biology, University of Iowa. The following secondary antibodies

were used: Donkey anti-mouse IgG (H and L) Antibody IRDye 800CW Conjugated (Licor) and

Alexa Fluor 680 goat anti-rabbit IgG (H+L) (Invitrogen). LI-COR Image Studio software

(RRID:SCR_015795) was used to quantify protein levels in different samples, relative to α-tubulin

levels. Fold change of protein levels was calculated relative to wildtype (N2)/ controls.

ii.   Western blot analysis of epithelial cell lines

A549 epithelial cells were lysed in a modified M2 buffer containing 20mM

tris[hydroxymethyl]aminomethane, 150mM NaCl at pH 7.4, 0.5% NP40, 3mM EDTA, 3mM

EGTA, 4mM PMSF, and a cOmplete Mini, EDTA-free protease inhibitor tablet according to the

manufacturer's instructions (Roche, #11836170001). Epithelial cells were collected by

mechanical dislodgment in lysis buffer at individual time points, incubated for 20 minutes on ice,

and spun at 14,000 x g for 5 minutes to remove insoluble material. 25ug protein was resolved

by NuPAGE™ 4-12% Bis-Tris gel (Invitrogen) and transferred to PVDF for immunoblotting. Anti-

p53 (Cell Signaling Technology, #9282) and β-actin (DSHB, 224-236-1) primary antibodies were

used with HRP-conjugated secondary antibodies to anti-rabbit IgG (Jackson ImmunoResearch,

#111-035-003) and anti-mouse IgG (Invitrogen, #62-6520). Densitometry was analyzed with

ImageJ software.

RNA extraction and real-time quantitative reverse-transcriptase PCR (RT-PCR)

Worms were bleach-hatched as described above and ~400-800 eggs/plate (2 plates/strain) were

seeded on fresh OP50 plates and were allowed to grow (i) for 30-34 hrs. at 25°C or (ii) for 36 hrs.

at 20°C and then harvested for RNA extraction. RNA was extracted as described earlier [130].

Briefly, plates were washed with sterile water and centrifuged. Water was carefully removed and

300 µl of Trizol (catalog no. 400753, Life Technologies) was added and snap-frozen immediately

in liquid nitrogen. Samples were thawed on ice and then lysed using a Precellys 24 homogenizer

(Bertin Corp.). RNA was then purified as detailed with appropriate volumes of reagents modified

to 300 µl of Trizol. The RNA pellet was dissolved in 17 µl of RNase-free water. The purified RNA

was then treated with deoxyribonuclease using the TURBO DNA-free kit (catalog no. AM1907,

Life Technologies) as per the manufacturer's protocol. cDNA was generated by using the iScript

cDNA Synthesis Kit (catalog no. 170–8891, Bio-Rad). qRT-PCR was performed using PowerUp

SYBR Green Master Mix (catalog no. A25742, Thermo Fisher Scientific) in QuantStudio 3 Real-

Time PCR System (Thermo Fisher Scientific) at a 10 µl sample volume, in a 96-well plate (catalog

no. 4346907, Thermo Fisher Scientific). The relative amounts of mRNA were determined using

the $\Delta\Delta C_t$ method for quantitation. We selected *pmp-3* as an appropriate internal control for gene

expression analysis in *C. elegans*.

All relative changes of mRNA were normalized to either that of the wild-type control or the control

for each genotype (specified in figure legends). Each experiment was repeated a minimum of

three times. For qPCR reactions, the amplification of a single product with no primer dimers was

confirmed by melt-curve analysis performed at the end of the reaction. Primers were designed

using Primer3 software and generated by Integrated DNA Technologies. The primers used for

the qRT-PCR analysis are listed below:

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
pfk-1.2	TTGCATCGAATCTGTGAAGC	TCGCTGCAGTCAAAGCTAGA
cyc-2.2	CGGTGGAGCTATTCCAGAAG	GCAACTTGTCCGGATTGTCT
egl-1	TCCAAGCTAGCAGCAATGTG	GCGAAAAGTCCAGAAGACG

ced-13	GCAACTCAAACACCGTTGAA	CAATGCTGGCATACGTCTTG
phg-1	CAGAGGAGCTTGTACCGACA	TCGTCTCTAGCAGTGCATGT
sod-3	CACTGCTTCAAAGCTTGTTCA	ATGGGAGATCTGGGAGAGTG
mtl-1	TGGATGTAAGGGAGACTGCAA	CATTTTAATGAGCCGCAGCA
lys-7	GCCGTCAAACTTGGCATCTT	GGGTTGTATGCACGAACGAA
pmp-3	TAGAGTCAAGGGTCGCAGTG	ATCGGCACCAAGGAAACTGG
cep-1(a+b isoform)	GCTCACTCTGTCGACTGCTGAGT	AACCCAAGTGTATCTGGGAACTTT
cep-1(a isoform)	GTTGTGCTCGACTCCCAAAAG	GGCACGCTTCTCAATTACAAGTT

1001

Chromatin immunoprecipitation (ChIP)

1002

Chromatin immunoprecipitation (ChIP) was performed as described earlier (Das et al., 2020).

Day-one adult worms were bleach-hatched as described above and ~1600 eggs were seeded

(~800 eggs/plate and 2 plates/genotype) on fresh OP5O plates. Plates were kept at 25°C for 30-

34 hrs. and larvae were washed with 1X PBS (pH 7.4) and cross-linked with freshly prepared 2%

formaldehyde (catalog no. 252549, Sigma Aldrich) at room temperature for 10 min followed by

addition of 250 mM Tris (pH 7.4) at room temperature for 10 min. Samples were then washed

three times in ice-cold 1X PBS supplemented with protease inhibitor cocktail and snap-frozen in

liquid nitrogen. The worm pellet was resuspended in FA buffer [50 mM HEPES (pH 7.4), 150 mM

NaCl, 50 mM EDTA, 1% Triton-X-100, 0.5% SDS and 0.1% sodium deoxycholate], supplemented

with 1 mM DTT and protease inhibitor cocktail. The suspended worm pellet was lysed using a

Precellys 24 homogenizer (Bertin Corp.), and then sonicated in a Bioruptor Pico Sonication

System (catalog no. B0106001, Diagenode) (15 cycles of 30 s on/off).

Endogenous CEP-1 was immunoprecipitated with anti-FLAG M2 magnetic bead (catalog no. M-8823, Sigma-Aldrich). Beads were first pre-cleared with salmon sperm DNA (catalog no. 15632–011, Invitrogen). Worm lysate was incubated at 4°C overnight with the pre-cleared FLAG beads. Beads were washed with low salt, high salt and LiCl wash buffers and then eluted in buffer containing EDTA, SDS and sodium bicarbonate. The elute was then de-crosslinked overnight in presence of Proteinase K. The DNA was purified by ChIP DNA purification kit (catalog no. D5205, Zymo Research). qPCR analysis of DNA was performed as described above using primer sets specific for different target genes. For all ChIP experiments, 10% of total lysate was used as 'input' and chromatin immunoprecipitated by different antibodies were expressed as % input values. The primers used for ChIP experiments, and the expected amplicon sizes are as follows:

Gene	Position	Forward primer (5'-3')	Reverse primer (5'-3')	Amplicon size
egl-1	TSS +4 to +123)	CTCACCTTTGCCTCAACCTC	CGAGGAGAAGTCCTGAGACG	120 bp
ced-13	Promoter (-428 to -310)	CTATTCTTGGCCGTGCTCAT	AGGCAATCTAGCATGCACCT	119 bp
phg-1	Promoter -428 to -294)	GCCAAACCTTCCAGATTTACA	TTCCTAGATAAGGGTTAGATGATGAGA	135 bp
phg-1	Intron 1 (+223 to +326)	AAGCTGAGCTCCGAAAACAA	TTTCCCGCTAAACGAGACAT	104 bp

cki-1	Promoter (-698 to -722)	TTT TCC ATA CTT CAC TAG TCA AAA CCT	GAC AGT GAG AAG CTT TCG TAT TGA	103 bp
cki-1	Promoter (-542 to -428)	TTC CTC ATA ATC ACG GAG CA	GGA ACC GAA GTG GTC AGA TG	114 bp
cki-1	Promoter (-323 to -217)	TCT CCG ACT GCT GAC CT	CGA GAA GGG GTG GAG TCA TA	106 bp

1025

1026 **Treatment of mouse epithelial cell lines with human IL-17 and Nutlin3A**

1027 A549 epithelial cells were seeded in a 6-well plate overnight and stimulated with 50-100ng/mL

1028 recombinant human-IL-17 (Peprotech, SKU 200-17) or 10uM Nutlin3A (Tocris, 675576-98-4),

1029 and supernatants and cell lysates were harvested 18 hours later.

1030

QUANTIFICATION AND STATISTICAL ANALYSIS

Each 'experiment' refers to a set of independent biological samples that undergo treatment

together. No statistical methods were used to predetermine sample size, and experiments were

not randomized, but some were blinded. For the assays where we scored dauers, for each

experiment, independent biological samples of the different strains were allowed to lay eggs and

the eggs were subjected to 'treatment' or 'control' conditions. The results of an experiment

generated counts categorized as either (a) dauers, or (b) non-dauers. The counts of dauers/non-

dauers for the different types of experiments were used to generate a contingency table, and a

chi-squared test was performed to test the independence (whether the results are related or not)

of the experiment. This Chi-square statistic, degrees of freedom (df) and p value for each

experiment is reported. Yates' continuity correction was used to prevent overestimation of

statistical significance for small data. For qPCR or other experiments, the data were analyzed by

using Student's t-test and/or one-way ANOVA with Tukey's correction (GraphPad Prism software)

as described in respective figure legends. P values are indicated as follows: *$p<0.05$; **$p< 0.01$;

***$p<0.001$, ns, not significant.

1046

1047   **DATA AVAILABILITY**

1048   The following data sets were generated

1049   Genes differentially expressed in the following *C. elegans* larvae bleach-hatched and allowed to

1050   grow for 30-36 hr. at 25°C: wild type (N2), *ilc-17.1*(syb5296) X and praEx022 (*unc-54*p::*ilc-17.1*

1051   cDNA::3XFLAG::*tbb-2* 3'UTR; p*dat-1*::GFP::*unc-54* 3'UTR).

1052

1053   Genes deferentially expressed in the following *C. elegans* larvae bleach-hatched and allowed to

1054   grow for 15 hr. and 32 hr. at 25°C: wild type (N2), *ilc-17.1*(syb5296) X, *cep-1* (gk138) I, *ilc-*

1055   *17.1*(syb5296) X; *cep-1* (gk138) I, and VEP036 (unc-119 (ed4); gtIs1 [CEP-1::GFP + unc-119

1056   (+)].

1057

1058   The data discussed in this publication have been deposited in NCBI's Gene Expression

1059   Omnibus (Edgar et al., 2002) and are accessible through GEO Series accession number

1060   GSE218596, , https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE218596 and

1061   GSE229132, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE229132

1062

Figure 1: ILC-17.1 is released by larval amphid neurons in response to food availability and reduced ILC-17.1 signaling causes dauer arrest.

a. Schematic of experiment to assess dauer arrest. For each experiment, independent samples of 10 day-one adults of each strain laid eggs for two hours at 20°C on 60 mm NGM plates with OP50 lawns. Adults were removed, and embryos were transferred to 25°C. Approximately 48 hr. later, larval phenotype was scored and categorized as either dauers, or larvae that continued development into stage 4 larvae (L4) and/or young adults. If unsure, SDS-resistance was used to confirm dauer arrest. Extent of crowding on plates was comparable. [Representative average numbers of embryos/plates from experiments in **b** are 64, 68, 56, 46 and 65 for wildtype (N2), *ilc-17.1*(syb5296) X, *ilc-17.1*(syb5297) X, *ilc-17.1*(syb5296) rescued with *ilc-17.1p::ilc-17.1*, and *ilc-17.1*(syb5296) rescued with *unc-54p::ilc-17.1* respectively]. Pearson's Chi-squared test with Yates' continuity correction was used to determine significance throughout the manuscript, unless otherwise stated. Chi-squared values were calculated for whole experiments, and for pair-wise comparisons. All legends show Chi-squared, df and p values for the whole experiment; p values for pair-wise comparisons are depicted on graphs using ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$ and ns, not significant.

b. Percentage larvae that arrest as dauers. ***$p<0.001$. (*n*=3-12 experiments; Chi-squared = 2359.6, df = 4, p-value < 2.2e-16.)

c, **d**. Representative micrographs 48 hours post-hatching at 25°C. **Top (c):** Wildtype (N2) grew to L4s. **Bottom (d):** *ilc-17.1*(syb5296) larvae arrested as dauers. Scale bar, 0.5mm.

e. Top: Schematic of experiment. **Bottom**: Percentage of wildtype (N2) and *ilc-17.1* larvae that arrest as dauers on control (L4440) and *ilcr-2* RNAi. ***$p < 0.001$, ns, not significant (n=3-10 experiments. Chi-squared = 1815.1, df = 7, p-value < 2.2e-16.)

1089

f. Top: Schematic of experiment to localize *ilc-17.1* mRNA and ILC-17.1 protein in the presence

and absence of food (OP50). Embryos were hatched at 20°C and imaged after 24-36 hours (L1

stage larvae). **Left panels:** Representative micrographs of z-section showing mCherry co-

expressed with *ilc-17.1* mRNA in amphid neurons. **Right panels**: projection of z-sections showing

immunostaining with anti-HA antibody to visualize CRISPR tagged endogenous ILC-17.1::HA

protein. Pharyngeal region of larvae **(top panel)** hatched in the absence of food, and **(middle**

panel) in the presence of food (OP50). B**ottom panels:** Controls (N2) that do not express HA or

mCherry used to assess specificity of the fluorescence signal. (n>3 experiments, 4-5 larvae each).

Scale bar, 5μm.

g. KEGG enrichments associated with differentially downregulated genes (p.adjust <0.05) in *ilc-*

17.1 mutant larvae (RNA-seq on 32-36 hr. larvae).

h. KEGG enrichments associated with differentially upregulated genes (p.adjust <0.05) in *ilc-17.1*

mutant larvae (RNA-seq on 32-36 hr. larvae).

 i. Heatmap depicting expression levels ($\log_2$ normalized counts) of the major dauer-specific

collagens in wildtype (N2) as L2-L3 larvae *en route* to development into reproductive adults, *ilc-*

17.1 deletion mutants *en route* to dauer entry and *ilc-17.1* deletion mutants rescued from dauer

entry with *unc-54p::ilc-17.1.*

Bars show mean ± S.E.M. Individual data points in the bar graphs in **b** and **e** represent the %

dauers/experiment, and the bar graph depicts mean of these percentages.

Figure 2: Reduced ILC-17.1 signaling acts through DAF-16/FOXO, DAF-3/SMAD-DAF-5/Ski complex, steroid hormone signaling and CEP-1/p53 to control dauer arrest.

a. Percent dauers in *ilc-17.1* deletion mutants and *ilc-17.1;daf-16* double mutant larvae. ***p < 0.001, (*n*=4-8 experiments, Chi-squared = 497.65, df = 1, p-value < 2.2e-16).

b. Left: Percent larvae showing DAF-16::GFP nuclear localization in intestinal cells. Larvae were grown at 20°C for 36 hrs**.** *n*=3 experiments with 10-15 larvae each. Each data point represents average no. of larvae with nuclear DAF-16::GFP/experiment. ***p < 0.001, (unpaired t-test). Note: experiments were conducted at 20°C, since DAF-16::GFP localized constitutively in intestinal nuclei even in wildtype larvae at 25°C. **Right:** representative micrographs showing intestinal DAF-16::GFP (boxed) in wildtype and *ilc-17.1* deletion mutant larvae. Scale bar, 5μm

c. Average *sod-3, mtl-1, and lys-7* mRNA levels in larvae 32-36 hr. post-hatching at 25°C. mRNA levels were determined relative to *pmp-3* and normalized to wildtype (N2) values. *n*=4-10 experiments. ***p < 0.001, **p < 0.01, ns=not significant, (unpaired t-test).

d. Percent dauers in *ilc-17.1* mutants and *ilc-17.1;daf-5* double mutants. ***p < 0.001, (*n*=3-5 experiments, Chi-squared = 787.3, df = 3, p-value < 2.2e-16).

e. Percent dauers in *ilc-17.1* mutants on control (H$_2$O) and exogenous 50nM Δ7-dafachronic acid. ***p < 0.001, (*n*=3-5 experiments, Chi-squared = 784.32, df = 3, p-value < 2.2e-16).

f-h. f, percent dauers in *ilc-17.1* mutants and *ilc-17.1;cep-1* double mutants. ***p < 0.001, (*n*=3-5 experiments, Chi-squared = 1029.2, df = 3, p-value < 2.2e-16). Representative micrographs of **g,** *ilc-17.1* deleted larvae arrested as dauers, and **h,** *ilc-17.1;cep-1* double mutants that grew to L4s, 48 hr. post-hatching at 25°C. Scale bar, 0.5mm

1140

1141   Bars show the mean ± S.E.M. Individual data points in bar graphs in **a, d, e, f** represent the

1142   average % dauers/experiment, Pearson's Chi-squaredd test with Yates' continuity correction.

1143

Figure 3: CEP-1/p53 acts downstream of reduced ILC-17.1 signaling and activates DAF-16/FOXO, DAF-3/SMAD-DAF-5/Ski complex to control dauer entry.

a-c. a, Percent dauers in wildtype (N2), *cep-1 (lg12501)*;CEP-1::GFP animals that do not overexpress CEP-1/p53, and CEP-1/p53 overexpressing larvae. ***$p < 0.001$, (n=3-4 experiments, Chi-squared = 569.35, df = 2, p-value < 2.2e-16). For *cep-1* mRNA expression levels see **Supplementary Fig. S6a**. Representative micrographs 48 hrs. post-hatching at 25°C, showing in **b,** CEP-1/p53 overexpressing larvae arrested as dauers and **c,** *cep-1 (lg12501)*; CEP-1::GFP as L4s. Scale bar, 0.5mm.

d. Percent dauers in wildtype (N2), and CEP-1/p53 overexpressing larvae on control (Ctrl; L4440) and *daf-16* RNAi. ***$p < 0.001$, **$p < 0.01$ (n=3-4 experiments, Chi-squared = 494.22, df = 3, p-value < 2.2e-16).

e. Percent dauers in wildtype (N2), *daf-5* (e1386), CEP-1/p53 overexpressing larvae, CEP-1/p53 overexpressing larvae crossed to *daf-5* (e1386), and N2 and CEP-1/p53 overexpressing larvae treated with 50nM Δ7-dafachronic acid (DA). ***$p < 0.001$, **$p < 0.01$, ns, not significant, (n=3-4 experiments, Chi-squared = 982.66, df = 5, p-value < 2.2e-16).

f. Average *sod-3, mtl-1, and lys-7* mRNA levels in larvae 32-36 hr. post-hatching at 25°C. mRNA levels were determined relative to *pmp-3* and normalized to wildtype (N2) values. n=4-10 experiments. ***$p < 0.001$, **$p < 0.01$, ns=not significant, (unpaired t-test).

g. Average *egl-1* and *ced-13* mRNA levels in 32-36 hr. larvae grown at 25°C. mRNA levels were determined relative to *pmp-3* and normalized to wildtype (N2) values. n=6-7 experiments. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns=not significant, (unpaired t-test).

h. CEP-1 occupancy (expressed as percent input) in 32-36 hr. larvae grown at 25°C, measured at the promoter proximal regions of *egl-1* and *ced-13* (**Top,** schematic)*.* CEP-1 occupancy was assayed in animals where endogenous *cep-1* was FLAG tagged at its C-terminus using CRISPR/Cas9, and immunoprecipitated with anti-FLAG antibody. Strains used: CEP-1::FLAG expressing animals in control wildtype background (Ctrl), in *ilc-17.1* background, and in *ilc-17.1* rescued from dauer arrest by *unc-54p::ilc-17.1* (*ilc-17.1;unc-54p::ilc-17.1* CDS). n=4 experiments. ***p < 0.001, **p < 0.01, *p < 0.05. (unpaired t-test).

i. Top: Representative Western blot showing CEP-1::FLAG and tubulin when crossed into *unc-54p::ilc-17.1,* and *ilc-17.1* deletion backgrounds. *cep-1* RNAi serves to show specificity. Samples constitute 32-36 hr. larvae grown at 25°C, except for positive control, irradiated sample (4; adults). **Bottom**: CEP-1 levels were quantified relative to tubulin and normalized to values from control animals (Ctrl; CEP-1::FLAG in wildtype background). n=4 experiments. Numbers in data bars correspond to representative bands on western. ***p < 0.001, **p < 0.01 (unpaired t-test).

Bars show the mean ± S.E.M. Individual points in bar graphs in **a, d, e** represent the % dauers/experiment, Pearson's Chi-squaredd test with Yates' continuity correction.

Figure 4. CEP-1/p53 accumulates in multipotent progenitor cells (P cells) during dauer entry, upregulates cell cycle inhibitors, and modulates cell fate.

a. Representative micrographs of 15 hr. L1 larvae at 25°C, showing *pax-3p::*mCherry marking ventral postembryonic blast cells (P cells; **top panel**; see schematic above), CEP-1::GFP expression (**middle panel**), and overlap (**bottom panel**) to show co-localization. Co-localized cells are marked with *. CEP-1::GFP expressing cells that do not co-localize with *pax-3p*::mCherry marked with #. Insets: zoom-in on the P blast cells. Scale bar=5um.

b. Percent co-localization between *pax-3p::*mCherry-expressing cells and CEP-1::GFP expressing cells at 20°C and 25°C. n=3 experiments, 10-15 larvae/experiment.***p < 0.001, (unpaired t-test). Each data point represents the percent of *pax-3p;mCherry*-expressing cells that colocalize with CEP-1::GFP in one animal.

c. Fluorescent intensity (arbitrary units; AU) measured in P cells of L1 larvae, 15 hr. post-hatching at 20°C and 25°C. Strains used: *cep-1* (lg12501); CEP-1::GFP animals that do not overexpress CEP-1/p53, CEP-1/p53 overexpressing larvae, and *ilc-17.1* deletion mutants expressing CEP-1::GFP. n=3 experiments, 5-6 larvae/experiment. ***p < 0.001, ns, not significant (unpaired t-test). Each data point represents the average fluorescence intensity value across five P cells in one animal (see **Material and Methods**).

d. Average *cki-1* and *phg-1* mRNA levels in 32-36 hr. larvae grown at 25°C. mRNA levels were determined relative to *pmp-3* and normalized to wildtype (N2) values. n=4-6 experiments. *p < 0.05. (unpaired t-test).

e. CEP-1 occupancy (expressed as percent input) in 32-36 hr. larvae grown at 25°C, measured at the promoter proximal region of *phg-1* (**Top,** schematic)*.* CEP-1 occupancy was assayed in

1215   animals where endogenous *cep-1* was FLAG tagged at its C-terminus using CRISPR/Cas9, and

1216   immunoprecipitated with anti-FLAG antibody. Stains used: CEP-1::FLAG expressing animals in

1217   control wildtype background (Ctrl), in *ilc-17.1* background, and in *ilc-17.1* rescued from dauer with

1218   *ilc-17.1; unc-54p::ilc-17.1* CDS. n=4 experiments. *p < 0.05. (unpaired t-test).

1219

1220   **f**. Percent dauers in *ilc-17.1* larvae and *ilc-17.1;phg-1* double mutants. **p < 0.05, ***p<0.001 (n=6

1221   experiments, Chi-squared = 826.95, df = 3, p-value< 2.2e-16). Individual points in the bar graph

1222   represent the % dauers/experiment, Pearson's Chi-squared test with Yates' continuity correction.

1223

1224   **g**. **Top:** Representative micrographs of wildtype (N2) animals and animals overexpressing CEP-

1225   1 driven by the *pax*-3 promoter or 25°C. Scale bar, 50µm. **Bottom:** Percent larvae that displayed

1226   body morphology defects. **p<0.01, (unpaired t-test)

1227

1228   **h**. **Top:** Schematic showing homeobox gene expression patterns during post-embryonic

1229   development. **Bottom:** Heatmap depicting expression levels ($\log_2$ normalized counts) of the

1230   homeobox genes on Chromosome III, 32 hrs post-hatching at 25°C, in *ilc-17.1* deletion mutants

1231   and CEP-1 overexpressing larvae that enter dauer, compared to *ilc-17.1; cep-1* double mutants

1232   and wildtype (N2) larvae that continue development. (Also see **Supplementary Table 12**).

1233

1234   Bars show the mean ± S.E.M.

1235

1236

Figure 5. Impaired glucose utilization in ILC-17.1 deficient larvae activates CEP-1/p53

and decreases the expression of Cytochrome C.

a. Left: Schematic of experimental design. **Right**: Percent dauers at 25°C amongst wildtype (N2),

ilc-17. deletion mutants, and *ilc-17.1* deletion mutants rescued by expressing *unc-54p::ilc-17.1* on

Control (L4440) and *fgt-1* RNAi. ***p < 0.001, **p < 0.01, ns, not significant. (n= 5 experiments,

Chi-squared = 1817.1, df = 5, p-value < 2.2e-16).

1243

1244 **b. Left**: Schematic of experimental design. **Right**: Percent SDS (1%)-resistant dauers that

1245 accumulate amongst wildtype (N2) and *ilc-17.1* deletion mutants growing on Control (L4440) and

1246 *fgt-1* RNAi under optimal growth conditions of 20°C. ***p < 0.001, **p < 0.01, ns, not significant.

1247 (n= 3 experiments, Chi-squared = 206.82, df = 7, p-value < 2.2e-16).

1248

1249 **c. Left**: Schematic of experimental design. **Right**: Percent dauers amongst wildtype (N2) larvae,

1250 *ilc-17.1* deletion mutants and CEP-1/p53 overexpressing animals on Control (H_2O) and 50mM

1251 glucose at 25°C. **p < 0.01, ns, not significant. (n= 3-6 experiments, Chi-squared = 2882.3, df =

1252 5, p-value < 2.2e-16).

1253

1254 **d**. Fluorescent intensity (arbitrary units; AU) measured in P cells in L1 larvae on Control (H_2O)

1255 and 50mM glucose, 15 hrs. post-hatching at 25°C. Strains: *cep-1* (lg12501); CEP-1::GFP and

1256 CEP-1/p53 overexpressing larvae. n=2-3 experiments, 3-5 larvae/experiment. ***p < 0.001,

1257 **p < 0.01, ns, not significant (unpaired t-test). Each data point indicates the average fluorescence

1258 intensity value across five P cells in one animal.

1259

1260 **e.** Average *phg-1, sod-3 and mtl-1* mRNA levels in 32-36 hr. larvae that arrest as dauers (on H_2O,

1261 control) and larvae that bypass dauer (50mM glucose) at 25°C. mRNA levels were determined

1262 relative to *pmp-3* and normalized to wildtype (N2) values. n=3 experiments. ***$p < 0.001$, **

1263 $p < 0.01$, ns, not significant (unpaired t-test).

1264

1265 **f**. Average *cyc-2.2* mRNA levels in 32-36 hr. larvae grown at 25°C. mRNA levels were determined

1266 relative to *pmp-3* and normalized to the wildtype (N2) values. n=4-6 experiments. ***$p < 0.001$,

1267 **$p < 0.01$, *$p < 0.05$. (unpaired t-test).

1268

1269 Bars show the mean ± S.E.M. Individual points in the bar graphs in **a, b, c** represent the %

1270 dauers/experiment, Pearson's Chi-squared test with Yates' continuity correction.

1271

Figure 6: ILC-17.1 suppresses CEP-1/p53 in *C. elegans* and mammalian epithelial cells.

a. Top: Representative Western blot showing p53 levels in A459 epithelial cells stimulated for 18hrs. with increasing doses of rIL-17. β-actin was used as a loading control and Nutlin was used as a positive control. **Bottom:** Quantification of p53 levels normalized to actin levels. n= 4 independent experiments. **p<0.01. (One-Way ANOVA with uncorrected Fisher's LSD; df=2).

b. Percent dauers upon overexpressing ILC-17.1 (expression under the *unc-54* promoter) in *cep-1* overexpressing larvae at 25°C. ***p < 0.001. (n=4 experiments, Chi-squared = 1245.4, df = 2, p-value < 2.2e-16).

c. Top: Schematic of germline apoptosis (apoptotic cells in red) scored with Acridine Orange. **Bottom:** Average numbers of apoptotic cells in day-two adult animals under control, non-irradiated conditions, and upon irradiation with 75 Gy (day-one adults were irradiated). n= 3 experiments, and 12 gonad arms/experiment. ***p < 0.001, **p < 0.01 (analysis of variance (ANOVA) with Tukey's correction, df=13). Note: p values between N2 vs *ilc-17.1* OE (*unc-54p*), non-irradiated and irradiated are p=0.0192 and p=0.0623, if compared by themselves (unpaired t-test), but do not rise to significance when corrected for the multiple comparisons.

d. Working model: ILC-17.1 secretion from larval amphid neurons that occurs in response to food availability links cell fate decisions with glucose metabolism and DAF-16, DAF-3/DAF-5 activation through CEP-1/p53.

Data in all graphs show mean ± S.E.M.

Supplementary Figure S1: ILC-17.1 modulates *C. elegans* developmental diapause.

a. Schematic of *ilc-17.1* gene depicting the syb5296 and syb5297 deletions made by CRISPR/Cas9 editing using two distinct guide RNAs. *ilc-17.1* (syb5296) X deletion mutants were used in all future experiments.

b. Average mRNA expression in *ilc-17.1* deletion mutants relative to wildtype (N2). n=4-8 experiments. ***p < 0.001 (unpaired t-test,).

c. Schematic of experiment leveraging the resistance of dauers to 1% SDS used to identify dauers developing under optimal growth conditions at 20°C.

d. Percent *ilc-17.1* larvae that transiently enter the dauer state as seen by SDS-resistance 48 hrs and 72 hrs. post-hatching under optimal conditions at 20°C. ***p < 0.001, **p < 0.01 (n=4 experiments, Chi-squared = 166.53, df = 3, p-value < 2.2e-16).

e-i. Representative micrographs showing DIC images of **e**, whole *ilc-17.1* deletion mutant dauer larvae (inset shows alae), **f,** pharyngeal region and **h,** arrested germline 72 hr. post-hatching at 25°C. Wildtype (N2) larval stage 2 (L2) animals used as comparisons: **g**, pharyngeal region and **i**, germline. Scale bar, a=50μm, f-i, 5 μm.

j. Average pharyngeal pumping rates/minute in *ilc-17.1* deletion mutants relative to wildtype (N2) larval stage 2 (L2) animals. n=4-8 experiments. ***p < 0.001, (unpaired t-test).

k. Percent F25D1.3*,* syb7367, and 2X backcrossed *ilc-17.1* larvae that arrest as dauers at 25°C. ***p < 0.001, (n=3 experiments; 50-100 larvae each).

1321

1322   **l. Top:** Schematic of experiment. **Bottom:** Percent wildtype (N2) larvae that arrest as dauers as

1323   determined by SDS-resistance, 48 hrs post-hatching on control (Ctrl; L4440) and *ilc-17.1* RNAi at

1324   25°C. *$p < 0.05$. (*n*=5 experiments, Chi-squared = 5.1712, df = 1, p-value = 0.02296).

1325   Bars show the mean ± S.E.M**.** Individual points in the bar graphs in **d, k** represent the %

1326   dauers/experiment, Pearson's Chi-squared test with Yates' continuity correction.

1327

Supplementary Figure S2: Characterization of ILC-17.1 signaling pathway components in

C. elegans.

a. Top: Schematic of CRISPR insertion of SL2::GFP into *ilcr-2* locus to identify sites of *ilcr-2*

expression. **Bottom, Upper panels:** Representative micrographs showing GFP expression in

cells of the pharynx (**left**), epidermis (**middle**; arrows) and gonad (**right**). **Bottom, lower panels:**

Control, wildtype (N2) non-transgenic animals not expressing GFP confirm specificity of GFP

expression. Scale bar=5μm.

1335

1336 **b.** Average expression in *ilc-17.1* mRNA in 32-26 hr. larvae of wildtype (N2), *ilc-17.1* deletion

1337 mutants rescued with *ilc-17.1p::ilc-17.1* and *unc-54p::ilc-17.1*. Values normalized to *pmp-3* and

1338 shown relative to wildtype (N2) values. *p<0.05 (unpaired t-test).

1339

c. Percentage of wildtype (N2) and *ilc-17.1* larvae that arrest as dauers on control HT115

(L4440) bacteria. ***p < 0.001 (n=10-20 experiments, Chi-squared = 3206, df = 3, p-value <

2.2e-16). Representative average numbers of embryos/plates of wildtype (N2), *ilc-17.1*, *ilc-17.1*

rescued with *ilc-17.1p::ilc-17.1*, and *ilc-17.1* rescued with *unc-54p::ilc-17.1* on HT115 (L4440)

bacteria are 59, 66, 66, 46 respectively. Individual points in the bar graphs represent the %

dauers/experiment, Pearson's Chi-squareddtest with Yates' continuity correction.

1346

d. Top: Schematic of CRISPR insertion of SL2::mCherry into *ilc-17.1* locus. **Bottom, upper**

panel: Representative z-section micrographs showing mCherry expression in amphid neurons

of *C. elegans* expressing mCherry as a bicistronic SL2 cassette along with the endogenous *ilc-*

17.1 gene to report on sites of *ilc-17.1* mRNA expression. Amphid neurons were identified by

DiO dye filling (green); mCherry overlapped with a subset of DiO filled neurons. **Bottom, lower**

panel: Control, wildtype (N2) animals not expressing mCherry confirm specificity of mCherry

expression. Scale bar=5μm.

1354

1355   **e. Top:** Schematic of *ilc-17.1::*flag driven by the *unc-54* promoter [*ilc-17.1* OE *(unc-54p)*] that

1356   was used to rescue *ilc-17.1* deletion mutants to assess ILC-17.1 protein localization. **Bottom:**

1357   Representative micrographs showing a z-section of these animals immunostained with α-FLAG

1358   antibody, and α-actin (to show expression in body wall muscle cells). Scale bar=5μm

1359

f. Top: Representative Western blots showing (**upper panel**) expression of ILC-17.1::FLAG

rescue constructs under the *ilc-17.1* promoter (1) and under the *unc-54* promoter (2) in *ilc-17.1*

deletion mutants. Wildtype (N2) were used to show specificity of α-FLAG antibody. (**lower**

panel), tubulin. **Bottom:** Quantification of ILC-17.1::FLAG levels in *ilc-17.1* deletion mutant

animals relative to tubulin, and normalized to FLAG levels in (1). n=3 experiments. ***$p < 0.001$

(unpaired t-test).

1366

1367   **g.** Quantification of fluorescent intensity in pharynx versus background (arbitrary units; AU), in

1368   larvae harboring CRISPR tagged endogenous ILC-17.1::HA**,** immunostained with anti-HA

1369   (**Figure 1f**). n=3 experiments. 5-6 larvae/experiment. ***$p < 0.001$, (unpaired t-test).

1370

1371   Representative micrographs showing z-sections through animals overexpressing ILC-17.1::FLAG

1372   in their body wall muscle cells as in **e**. Note: ILC-17.1::FLAG is detected at the amphid

1373   commissures near the pharynx (**top panel**; arrow) and in the epidermis (**bottom panel**; arrows)

1374   in addition to the body wall muscle cells (**e**), Scale bar=5μm

1375

1376   Bars show the mean ± S.E.M.

1377

1378

1379   **Supplementary Figure S3: RNA-seq analysis of ILC-17.1 deficient larvae.**

1380   **a.** Pair-wise distance matrix of RNA-seq samples shows the expected clustering of total RNA of

1381   the biological triplicates of each strain [Strains used: wildtype (N2), *ilc-17.1*, and *ilc-17.1* rescued

1382   with *unc-54p::ilc-17.1* CDS]

1383

1384   **b.** Spearman correlation between all genes in a microarray analysis of dauer larvae that were

1385   induced by starvation and *ilc-17.1* deleted larvae *en route* to dauer arrest grown for 32-36 hrs. at

1386   25°C.

1387

1388   **c**. Spearman correlation between all genes in L1 larvae exposed to ascarosides and *ilc-17.1*

1389   deleted larvae *en route* to dauer arrest grown for 32-36 hrs. at 25°C.

1390

Supplementary Figure S4: ILC-17.1 loss does not affect the animal's chemotaxis or

pumping and is associated with changes in *daf-16* target gene expression.

a. Top: Schematic of experimental design. **Bottom:** Average chemotaxis index towards OP50,

0.01% v/v diacetyl, and 1M lysine for *ilc-17.1* deletion mutants and wildtype (N2) day-one adults

at 20°C. n= 5-8 experiments. ns, not significant (unpaired t-test). Data points: average

chemotaxis/experiment.

1397

1398 **b.** Number of pharyngeal pumps/min in *ilc-17.1* deletion mutants and wildtype (N2) day-one adults

1399 at 20°C, on OP50 lawns, and on plates without food. n=3 experiments, with at least 5

1400 animals/experiment scored. Data points: pumping rates of individual animals. ns, not significant

1401 (unpaired t-test).

1402

1403 **c.** Average numbers of fluorescent latex beads in similar areas of distal gut lumen of 40-hour *ilc-*

1404 *17.1* deletion mutants and wildtype (N2) larvae at 20°C. n=3 experiments, with at least 10

1405 animals/experiment. Data points: the numbers of beads in individual animals. ns, not significant

1406 (unpaired t-test).

1407

1408 **d.** Percent dauers amongst wildtype (N2) and *ilc-17.1* deletion mutants raised on Control (H_2O)

1409 and 5mM 8-bromo-cGMP at 25°C. ***$p < 0.001$, Pearson's Chi-squared test with Yates

1410 correction, (n=3 experiments. Chi squared = 798, df = 3, p-value <2e-16.). Individual points in

1411 the bar graphs represent the % dauers/experiment,

1412

1413 **e.** Average *sod-3, mtl-1, and lys-7* mRNA levels in 40 hr. old larvae grown at 20°C. mRNA levels

1414 were determined relative to *pmp-3* and normalized to wildtype (N2) values. n=3 experiments.

1415 ***$p < 0.001$, **$p < 0.01$, ns, not significant (unpaired t-test).

1416

1417   Data in all graphs show mean ± S.E.M.

1418

Supplementary Figure S5: ILC-17.1 deficiency does not alter the expression levels of Insulin signaling- or TGFβ signaling- pathway components.

a. Percent dauers in *ilc-17.1* deletion mutant larvae following RNAi-mediated downregulation of several genes known to interact with DAF-16 and DAF3/DAF-5. *daf-16* RNAi was used as a positive control. ***p < 0.001, ns, not significant. (*n=3*-5 experiments, Chi-squared = 156.66, df = 3, p-value < 2.2e-16).

b. Heatmap depicting expression levels ($\log_2$ normalized counts) of *daf-2* (boxed) and insulin ligands in the *ilc-17.1* deletion mutants, *ilc-17.1* deletion mutants rescued by *unc-54p::ilc-17.1* and wildtype (N2). Arrows highlight lack of changes in *ins-1* and *ins-18* mRNA. RNA-seq data from larvae grown for 32-36 hrs. at 25°C.

c. Average mRNA levels (insulins) in 32-36 hr. larvae grown at 25°C. mRNA levels were determined relative to *pmp-3* and normalized to wildtype (N2) values. n=3 experiments. **p < 0.001, ns, not significant (unpaired t-test).

d. Left: Representative micrographs showing DAF-28::GFP expression in the last intestinal cell of 32-36hr. larvae at 25°C of *ilc-17.1* deletion mutants and wildtype (N2). Images are projections of confocal z-sections. Scale bar: 5μm. **Right:** Quantification of fluorescent intensity (arbitrary units, AU) of DAF-28::GFP. n=3 experiments. Data points: the fluorescence intensity of DAF-28::GFP in individual animals. ns, not significant (unpaired t-test).

e. Top: Representative micrographs showing INS-4::GFP expression in the pharynx and neurons of 32-36 hr. larvae at 25°C in *ilc-17.1* deletion mutants and wildtype (N2). Images are projections of confocal z-sections. Scale bar: 5μm. **Bottom:** Bars represent percent larvae that

1444   express GFP in the different tissues where INS-4 is known to be expressed. n=3 experiments of

1445   5-10 larvae of each strain. ns, not significant (unpaired t-test).

1446

1447   **f.** Overexpressing ILC-17.1 under *unc-54p::ilc-17.1* CDS [*ilc-17.1* OE (*unc-54p)*] in *daf-2*

1448   (e1370) III strains does not rescue their dauer arrest. ***p < 0.001, ns, not significant (n=3

1449   experiments, Chi-squared = 521, df = 3, p-value < 2.2e-16).

1450

1451   **g.** Heatmap depicting expression levels (log$_2$ normalized counts) of TGFβ signaling pathway

1452   components in the *ilc-17.1* deletion mutants, *ilc-17.1* deletion mutants rescued by *unc-54p::ilc-*

1453   *17.1*, and wildtype (N2).

1454

1455   Data in all graphs show mean ± S.E.M. Individual points in the bar graphs in **e**, **g**, represent the

1456   % dauers/experiment, Pearson's Chi-squared test with Yates' continuity correction.

1457

1458 **Supplementary Figure S6: CEP-1/p53 overexpression induces dauer arrest.**

1459 **a. Top:** Schematic of *cep-1* mRNA, isoform b. **Bottom:** Average *cep-1* isoform b mRNA levels in

1460 32-36 hr. larvae grown at 25°C. mRNA levels were determined relative to *pmp-3* and normalized

1461 to wildtype (N2) values. *n*=3 experiments. **p < 0.001, *p < 0.05 (unpaired t-test).

1462

1463 **b. Top:** Schematic of SDS treatment to assess dauer formation at 20°C. **Bottom:** Percent of

1464 wildtype (N2) and *cep-1* overexpressing larvae (*cep-1* OE) that enter a dauer state during

1465 development, 48 hrs. and 72 hrs. post-hatching at 20°C. ***p < 0.001, (n=3 experiments, Chi-

1466 squared = 414.65, df = 3, p-value < 2.2e-16). Individual points in the bar graphs represent the %

1467 dauers/experiment, Pearson's Chi-squared test with Yates' continuity correction.

1468

1469 **c**. Venn diagram depicting overlap between significantly differentially expressed genes (p<0.05)

1470 in *ilc-17.1* deletion mutants and CEP-1/p53 overexpressing larvae, 32-36 hr. post-hatching at

1471 25°C *en route* to dauer entry. p=0.0; hypergeometric test (Also see **Supplementary Tables 9,**

1472 **10**)

1473

1474 **d**. Heatmap comparing expression levels (log$_2$ normalized counts) of the major dauer-specific

1475 collagens in the *ilc-17.1* deletion mutants, CEP-1/p53 overexpressing larvae, *ilc-17.1; cep-1*

1476 double mutants, *ilc-17.1* deletion mutants rescued by *unc-54p::ilc-17.1,* and wildtype (N2).

1477

1478 **e.** KEGG enrichments associated with differentially downregulated genes (p.adjust <0.05) in CEP-

1479 1/p53 overexpressing larvae [RNA-seq: 32-36 hr. larvae grown at 25°C; **Supplementary Tables**

1480 **9**].

1481

1482   **f**. KEGG enrichments associated with differentially upregulated genes (padjust <0.05) in CEP-

1483   1/p53 overexpressing larvae [RNA-seq: 32-36 hr. larvae grown at 25°C; **Supplementary Tables**

1484   **9**].

1485

1486   **g.** Principle Component Analysis **(PCA)** of the observed variance in the triplicate RNA-seq

1487   samples collected during development, 15 hrs. and 32 hrs. post-hatching at 25°C, of *ilc-17.1*

1488   deletion mutants, CEP-1/p53 overexpressing larvae, *ilc-17.1; cep-1* double mutants and

1489   wildtype (N2) larvae (**Supplementary Tables 9).**.

1490

1491   Data in all graphs show mean ± S.E.M.

Supplementary Figure S7: CEP-1/p53 overexpression upregulates DAF-16/FOXO targets and cell cycle inhibitors to induce dauer arrest.

Average *sod-3, mtl-1, and lys-7* mRNA levels in 40 hr. larvae grown at 20°C. mRNA levels were determined relative to *pmp-3* and normalized to wildtype (N2) values. Strains used are on X-axis. n=6-10 experiments. ***p < 0.001, **p < 0.01, ns, not significant. (unpaired t-test).

b. Percent dauers in *daf-2* and *cep-1;daf-2* double mutants. Note: no rescue. n=3 experiments. ns, not significant. (Chi-squared = Not appropriate, df = 1, p-value = no significant)

c. Percent high temperature (27.5°C; HID phenotype) dauers in wildtype (N2) and *cep-1* deletion mutants. ns, not significant. (n=4 experiments, Chi-squared = 0.67295, df = 1, p-value = 0.412).

d. Average *egl-1* and *ced-13* mRNA levels in 40 hr. larvae grown at 20°C. mRNA levels were determined relative to *pmp-3* and normalized to wildtype (N2) values. n=6-7 experiments. ***p < 0.001, **p < 0.01, ns, not significant (unpaired t-test).

e. CEP-1 occupancy (expressed as percent input) in 32-36 hr. larvae grown at 25°C, measured at three promoter proximal regions of *cki-1* (Schematic on **top**). Strains used: CEP-1::FLAG expressing animals in wildtype background (Ctrl), in *ilc-17.1* deletion background. n=4 experiments. ns, not significant. (unpaired t-test).

f. Percent dauers in *ilc-17.1* larvae, CEP-1/p53 overexpressing larvae and *daf-2* larvae upon RNAi induced downregulation of *cki-1* and *phg-1.* ***p < 0.001, **p < 0.01, ns, not significant (n=6 experiments, Chi-squared = 7443.7, df = 14, p-value < 2.2e-16).

g. Representative micrograph of *ilc-17.1; phg-1* double mutants showing the dauer rescue of some individuals but not others (dauers marked with arrow).

h. Average pumps/minute in *ilc-17.1, cep-1 and daf-2* dauers when subjected to control (L4440; Ctrl), *cki-1* and *phg-1* RNAi. n= 4 experiments, and 10 dauers were scored per experiment. *$p < 0.05$, ns, not significant (unpaired t-test). Pumping rates in wildtype (N2), L2 larvae are shown for comparison.

Data in all graphs show mean ± S.E.M. Individual points in the bar graphs in **b**, **c**, **f**, represent the % dauers/experiment, Pearson's Chi-squared test with Yates' continuity correction.

Supplementary Figure S8: CEP-1/p53 activity can be repressed by glucose supplementation.

a. Percent dauers amongst *ilc-17.1* deletion mutants, CEP-1/p53 overexpressing larvae and wildtype (N2) larvae on 5 mM, 10mM and 50 mM glucose at 25°C. ***$p < 0.001$, **$p < 0.01$, ns, not significant. (*n*= 5-9 experiments, Chi-squared = 5263.6, df = 11, p-value < 2.2e-16)

b. Percent dauers amongst *ilc-17.1* deletion mutants and wildtype (N2) larvae on 50mM glucose on Control (Ctrl; L4440) and *fgt-1* RNAi. **$p < 0.01$, *$p < 0.05$, ns, not significant. (*n*=6 experiments, Chi-squared = 299.36, df = 3, p-value < 2.2e-16).

c. Percent dauers amongst *ilc-17.1* deletion mutants and wildtype (N2) larvae upon exposure to 50mM DOG, 50mM glucose, and 0.1mM MitoTEMPO at 25°C (see X-axis for different conditions). ***$p < 0.001$, ns, not significant, (n=3- experiments, Chi-squared = 1251.7, df = 9, p-value < 2.2e-16).

d. Percent dauers amongst *ilc-17.1* deletion mutants and wildtype (N2) larvae on 20mM Skim Milk powder and 0.26 mM L-Glutamine at 25°C. ns, not significant, (n=3 experiments, Chi-squared = 648.89, df = 5, p-value < 2.2e-16).

e. Average mRNA levels of *egl-1* and *ced-13* in wildtype (N2) larvae subjected to Control (Ctrl; L4440) and *fgt-1* RNAi. mRNA levels were determined relative to *pmp-3* and normalized to wildtype (N2) values. *n=3* experiments, *$p < 0.05$, ns, not significant (unpaired t-test).

Data in all graphs show mean ± S.E.M. Individual points in the bar graphs in **a-d** represent the % dauers/experiment, Pearson's Chi-squared test with Yates' continuity correction.

Supplementary Figure S9: ILC-17.1 deficiency and CEP-1/p53 overexpression modulate

key glucose metabolic enzymes.

a-f. Heatmap depicting expression levels ($\log_2$ normalized counts) of glycolysis and mitochondrial

OXPHOS genes in the *ilc-17.1* deletion mutants, CEP-1/p53 overexpressing animals, *ilc-17.1;*

cep-1 double mutants, *ilc-17.1* rescued with *unc-54p::ilc-17.1*, and wildtype (N2). Arrows:

highlight decreases in *pfk-1.2* and *cyc-2.2* mRNA levels in *ilc-17.1* deleted and CEP-1/p53

overexpressing larvae (Boxed). RNA-seq data was collected as previously described.

1561

1562 **g**. Average *pfk-1.2* mRNA levels in 32-36 hr. old larvae grown at 25°C. mRNA levels were

1563 determined relative to *pmp-3* and normalized to wildtype (N2) values. Strain names on X-axis.

1564 n=4-6 experiments. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$. (unpaired t-test).

1565

h. Top: Schematic of SDS treatment to assess dauer formation at 20°C. **Bottom:** Percent of

wildtype (N2) and *ilc-17.1* deletion mutant larvae that enter a dauer state at 20°C, 48 hr. post-

hatching, when subjected to control (Ctrl; L4440), *cyc-2.2* and *pfk-1.2* RNAi. **$p < 0.01$, ns, not

significant (*n*=3-4 experiments, Chi-squared = 213.38, df = 5, p-value < 2.2e-16) . Individual

points in the bar graphs represent the % dauers/experiment, Pearson's Chi-squared test with

Yates' continuity correction.

1572

1573 Data in all graphs show mean ± S.E.M.

Supplementary Figure S10: rIL-17 induces the expected cytokine response in human epithelial cells.

a. Soluble CXCL5 levels in the supernatant of A459 epithelial cells stimulated for 18hrs. with rIL-17 measured by ELISA. Data is a summary of 2 independent experiments, performed in triplicates. *p<0.05, (unpaired t-test). Data show mean ± S.E.M.

b. Percent wild-type (N2) and *npr-1*(ad609); *ilc-17.1*(tm5218) double mutant larvae that arrest as dauers at 25°C. ns, not significant (n=3 experiments; 50-100 larvae each).

1587   References

1588

1.   Golden, J.W. & Riddle, D.L. The Caenorhabditis elegans dauer larva: developmental effects of

pheromone, food, and temperature. *Dev Biol* **102**, 368-78 (1984).

2.   Swanson, M.M. & Riddle, D.L. Critical periods in the development of the Caenorhabditis elegans

dauer larva. *Dev Biol* **84**, 27-40 (1981).

3.   Hu, P.J. Dauer. *WormBook*, 1-19 (2007).

4.   Karp, X. Working with dauer larvae. *WormBook* **2018**, 1-19 (2018).

5.   Fielenbach, N. & Antebi, A. C. elegans dauer formation and the molecular basis of plasticity. *Genes

Dev* **22**, 2149-65 (2008).

6.   Burnell, A.M., Houthoofd, K., O'Hanlon, K. & Vanfleteren, J.R. Alternate metabolism during the

dauer stage of the nematode Caenorhabditis elegans. *Exp Gerontol* **40**, 850-6 (2005).

7.   Rashid, S., Wong, C. & Roy, R. Developmental plasticity and the response to nutrient stress in

Caenorhabditis elegans. *Dev Biol* **475**, 265-276 (2021).

8.   Cohen, S.M., Wrobel, C.J.J., Prakash, S.J., Schroeder, F.C. & Sternberg, P.W. Formation and

function of dauer ascarosides in the nematodes Caenorhabditis briggsae and Caenorhabditis

elegans. *G3 (Bethesda)* **12**(2022).

9.   Braendle, C., Milloz, J. & Felix, M.A. Mechanisms and evolution of environmental responses in

Caenorhabditis elegans. *Curr Top Dev Biol* **80**, 171-207 (2008).

10.   Diaz, S.A. & Viney, M. The evolution of plasticity of dauer larva developmental arrest in the

nematode Caenorhabditis elegans. *Ecol Evol* **5**, 1343-53 (2015).

11.   Ewald, C.Y., Landis, J.N., Porter Abate, J., Murphy, C.T. & Blackwell, T.K. Dauer-independent

insulin/IGF-1-signalling implicates collagen remodelling in longevity. *Nature* **519**, 97-101 (2015).

12.   Harvey, S.C., Shorto, A. & Viney, M.E. Quantitative genetic analysis of life-history traits of
1611   Caenorhabditis elegans in stressful environments. *BMC Evol Biol* **8**, 15 (2008).

1612   13.   Zhu, J. & Thompson, C.B. Metabolic regulation of cell growth and proliferation. *Nat Rev Mol Cell*
1613   *Biol* **20**, 436-450 (2019).

1614   14.   Altan-Bonnet, G. & Mukherjee, R. Cytokine-mediated communication: a quantitative appraisal of
1615   immune complexity. *Nat Rev Immunol* **19**, 205-217 (2019).

1616   15.   Mills, K.H.G. IL-17 and IL-17-producing cells in protection versus pathology. *Nat Rev Immunol*
1617   (2022).

1618   16.   Li, X., Bechara, R., Zhao, J., McGeachy, M.J. & Gaffen, S.L. IL-17 receptor-based signaling and
1619   implications for disease. *Nat Immunol* **20**, 1594-1602 (2019).

1620   17.   Douglas, A., Stevens, B. & Lynch, L. Interleukin-17 as a key player in neuroimmunometabolism.
1621   *Nat Metab* **5**, 1088-1100 (2023).

1622   18.   Hall, P.A. & Lane, D.P. Tumor suppressors: a developing role for p53? *Curr Biol* **7**, R144-7 (1997).

1623   19.   Jain, A.K. & Barton, M.C. p53: emerging roles in stem cells, development and beyond.
1624   *Development* **145**(2018).

1625   20.   Levine, A.J. p53: 800 million years of evolution and 40 years of discovery. *Nat Rev Cancer* **20**, 471-
1626   480 (2020).

1627   21.   Rutkowski, R., Hofmann, K. & Gartner, A. Phylogeny and function of the invertebrate p53
1628   superfamily. *Cold Spring Harb Perspect Biol* **2**, a001131 (2010).

1629   22.   Lu, W.J., Amatruda, J.F. & Abrams, J.M. p53 ancestry: gazing through an evolutionary lens. *Nat*
1630   *Rev Cancer* **9**, 758-62 (2009).

1631   23.   Bowen, M.E., Mulligan, A.S., Sorayya, A. & Attardi, L.D. Puma- and Caspase9-mediated apoptosis
1632   is dispensable for p53-driven neural crest-based developmental defects. *Cell Death Differ* **28**,
1633   2083-2094 (2021).

1634   24.   Van Nostrand, J.L. *et al.* Inappropriate p53 activation during development induces features of

1635   CHARGE syndrome. *Nature* **514**, 228-32 (2014).

1636   25.   Chen, C. *et al.* IL-17 is a neuromodulator of Caenorhabditis elegans sensory responses. *Nature*

1637   **542**, 43-48 (2017).

1638   26.   Flynn, S.M. *et al.* MALT-1 mediates IL-17 neural signaling to regulate C. elegans behavior,

1639   immunity and longevity. *Nat Commun* **11**, 2099 (2020).

1640   27.   Lillis, M., Zaccardi, N.J. & Heiman, M.G. Axon-dendrite and apical-basolateral sorting in a single

1641   neuron. *Genetics* **221**(2022).

1642   28.   Golden, J.W. & Riddle, D.L. A pheromone-induced developmental switch in Caenorhabditis

1643   elegans: Temperature-sensitive mutants reveal a wild-type temperature-dependent process. *Proc*

1644   *Natl Acad Sci U S A* **81**, 819-23 (1984).

1645   29.   Thomas, J.H., Birnby, D.A. & Vowels, J.J. Evidence for parallel processing of sensory information

1646   controlling dauer formation in Caenorhabditis elegans. *Genetics* **134**, 1105-17 (1993).

1647   30.   Vowels, J.J. & Thomas, J.H. Genetic analysis of chemosensory control of dauer formation in

1648   Caenorhabditis elegans. *Genetics* **130**, 105-23 (1992).

1649   31.   Ailion, M. & Thomas, J.H. Isolation and characterization of high-temperature-induced Dauer

1650   formation mutants in Caenorhabditis elegans. *Genetics* **165**, 127-44 (2003).

1651   32.   Li, W. & Ou, G. The application of somatic CRISPR-Cas9 to conditional genome editing in

1652   Caenorhabditis elegans. *Genesis* **54**, 170-81 (2016).

1653   33.   Paix, A. *et al.* Scalable and versatile genome editing using linear DNAs with microhomology to

1654   Cas9 Sites in Caenorhabditis elegans. *Genetics* **198**, 1347-56 (2014).

1655   34.   Xu, S. The application of CRISPR-Cas9 genome editing in Caenorhabditis elegans. *J Genet Genomics*

1656   **42**, 413-21 (2015).

35.   Holt, S.J. Staying alive in adversity: transcriptome dynamics in the stress-resistant dauer larva. *Funct Integr Genomics* **6**, 285-99 (2006).

36.   Cohen, S.M., Sun, J.J., Schroeder, F.C. & Sternberg, P.W. Transcriptional Response to a Dauer-Inducing Ascaroside Cocktail in Late L1 in C. elegans. *MicroPubl Biol* **2021**(2021).

37.   Wang, J. & Kim, S.K. Global analysis of dauer gene expression in Caenorhabditis elegans. *Development* **130**, 1621-34 (2003).

38.   Wirick, M.J. *et al.* daf-16/FOXO blocks adult cell fate in Caenorhabditis elegans dauer larvae via lin-41/TRIM71. *PLoS Genet* **17**, e1009881 (2021).

39.   Kyani-Rogers, T. *et al.* Developmental history modulates adult olfactory behavioral preferences via regulation of chemoreceptor expression in Caenorhabditiselegans. *Genetics* **222**(2022).

40.   Ow, M.C., Borziak, K., Nichitean, A.M., Dorus, S. & Hall, S.E. Early experiences mediate distinct adult gene expression and reproductive programs in Caenorhabditis elegans. *PLoS Genet* **14**, e1007219 (2018).

41.   Bargmann, C.I. Chemosensation in C. elegans. *WormBook*, 1-29 (2006).

42.   Motola, D.L. *et al.* Identification of ligands for DAF-12 that govern dauer formation and reproduction in C. elegans. *Cell* **124**, 1209-23 (2006).

43.   Vowels, J.J. & Thomas, J.H. Multiple chemosensory defects in daf-11 and daf-21 mutants of Caenorhabditis elegans. *Genetics* **138**, 303-16 (1994).

44.   Lin, K., Dorman, J.B., Rodan, A. & Kenyon, C. daf-16: An HNF-3/forkhead family member that can function to double the life-span of Caenorhabditis elegans. *Science* **278**, 1319-22 (1997).

45.   Libina, N., Berman, J.R. & Kenyon, C. Tissue-specific activities of C. elegans DAF-16 in the regulation of lifespan. *Cell* **115**, 489-502 (2003).

46.   Lee, S.S., Kennedy, S., Tolonen, A.C. & Ruvkun, G. DAF-16 target genes that control C. elegans life-span and metabolism. *Science* **300**, 644-7 (2003).

47.   Kenyon, C. & Murphy, C.T. Enrichment of regulatory motifs upstream of predicted DAF-16 targets.

1682   *Nat Genet* **38**, 397-8; author reply 398 (2006).

1683   48.   da Graca, L.S. *et al.* DAF-5 is a Ski oncoprotein homolog that functions in a neuronal TGF beta

1684   pathway to regulate C. elegans dauer development. *Development* **131**, 435-46 (2004).

1685   49.   Pierce, S.B. *et al.* Regulation of DAF-2 receptor signaling by human insulin and ins-1, a member of

1686   the unusually large and diverse C. elegans insulin gene family. *Genes Dev* **15**, 672-86 (2001).

1687   50.   Ritter, A.D. *et al.* Complex expression dynamics and robustness in C. elegans insulin networks.

1688   *Genome Res* **23**, 954-65 (2013).

1689   51.   Gems, D. *et al.* Two pleiotropic classes of daf-2 mutation affect larval arrest, adult behavior,

1690   reproduction and longevity in Caenorhabditis elegans. *Genetics* **150**, 129-55 (1998).

1691   52.   Zecic, A. & Braeckman, B.P. DAF-16/FoxO in Caenorhabditis elegans and Its Role in Metabolic

1692   Remodeling. *Cells* **9**(2020).

1693   53.   Oh, S.W. *et al.* JNK regulates lifespan in Caenorhabditis elegans by modulating nuclear

1694   translocation of forkhead transcription factor/DAF-16. *Proc Natl Acad Sci U S A* **102**, 4494-9

1695   (2005).

1696   54.   Greer, E.L. *et al.* An AMPK-FOXO pathway mediates longevity induced by a novel method of

1697   dietary restriction in C. elegans. *Curr Biol* **17**, 1646-56 (2007).

1698   55.   Leiser, S.F., Begun, A. & Kaeberlein, M. HIF-1 modulates longevity and healthspan in a

1699   temperature-dependent manner. *Aging Cell* **10**, 318-26 (2011).

1700   56.   Perrin, A.J. *et al.* Noncanonical control of C. elegans germline apoptosis by the insulin/IGF-1 and

1701   Ras/MAPK signaling pathways. *Cell Death Differ* **20**, 97-107 (2013).

1702   57.   Gurkar, A.U. *et al.* Dysregulation of DAF-16/FOXO3A-mediated stress responses accelerates

1703   oxidative DNA damage induced aging. *Redox Biol* **18**, 191-199 (2018).

1704   58.   Baruah, A. *et al.* CEP-1, the Caenorhabditis elegans p53 homolog, mediates opposing longevity

1705   outcomes in mitochondrial electron transport chain mutants. *PLoS Genet* **10**, e1004097 (2014).

1706   59.   Ewald, C.Y., Castillo-Quan, J.I. & Blackwell, T.K. Untangling Longevity, Dauer, and Healthspan in

1707   Caenorhabditis elegans Insulin/IGF-1-Signalling. *Gerontology* **64**, 96-104 (2018).

1708   60.   Derry, W.B., Putzke, A.P. & Rothman, J.H. Caenorhabditis elegans p53: role in apoptosis, meiosis,

1709   and stress resistance. *Science* **294**, 591-5 (2001).

1710   61.   Schumacher, B., Hofmann, K., Boulton, S. & Gartner, A. The C. elegans homolog of the p53 tumor

1711   suppressor is required for DNA damage-induced apoptosis. *Curr Biol* **11**, 1722-7 (2001).

1712   62.   Schumacher, B. *et al.* Translational repression of C. elegans p53 by GLD-1 regulates DNA damage-

1713   induced apoptosis. *Cell* **120**, 357-68 (2005).

1714   63.   Levi-Ferber, M. *et al.* It's all in your mind: determining germ cell fate by neuronal IRE-1 in C.

1715   elegans. *PLoS Genet* **10**, e1004747 (2014).

1716   64.   Hicks, T. *et al.* R-loop-induced irreparable DNA damage evades checkpoint detection in the C.

1717   elegans germline. *Nucleic Acids Res* **50**, 8041-8059 (2022).

1718   65.   Arum, O. & Johnson, T.E. Reduced expression of the Caenorhabditis elegans p53 ortholog cep-1

1719   results in increased longevity. *J Gerontol A Biol Sci Med Sci* **62**, 951-9 (2007).

1720   66.   Lee, C.C. *et al.* Mutation of a Nopp140 gene dao-5 alters rDNA transcription and increases germ

1721   cell apoptosis in C. elegans. *Cell Death Dis* **5**, e1158 (2014).

1722   67.   Gartner, A., Boag, P.R. & Blackwell, T.K. Germline survival and apoptosis. *WormBook*, 1-20 (2008).

1723   68.   Greiss, S., Schumacher, B., Grandien, K., Rothblatt, J. & Gartner, A. Transcriptional profiling in C.

1724   elegans suggests DNA damage dependent apoptosis as an ancient function of the p53 family. *BMC*

1725   *Genomics* **9**, 334 (2008).

1726   69.   Schumacher, B. *et al.* C. elegans ced-13 can promote apoptosis and is induced in response to DNA

1727   damage. *Cell Death Differ* **12**, 153-61 (2005).

70.   Schertel, C. & Conradt, B. C. elegans orthologs of components of the RB tumor suppressor

complex have distinct pro-apoptotic functions. *Development* **134**, 3691-701 (2007).

71.   Sendoel, A., Kohler, I., Fellmann, C., Lowe, S.W. & Hengartner, M.O. HIF-1 antagonizes p53-

mediated apoptosis through a secreted neuronal tyrosinase. *Nature* **465**, 577-83 (2010).

72.   Sulston, J.E. & Horvitz, H.R. Post-embryonic cell lineages of the nematode, Caenorhabditis

elegans. *Dev Biol* **56**, 110-56 (1977).

73.   Sternberg, P.W. Vulval development. *WormBook*, 1-28 (2005).

74.   Horvitz, H.R. & Sternberg, P.W. Nematode postembryonic cell lineages. *J Nematol* **14**, 240-8

(1982).

75.   Chisholm, A.D. & Hardin, J. Epidermal morphogenesis. *WormBook*, 1-22 (2005).

76.   Herman, M.A. Hermaphrodite cell-fate specification. *WormBook*, 1-16 (2006).

77.   Costa, M., Weir, M., Coulson, A., Sulston, J. & Kenyon, C. Posterior pattern formation in C. elegans

involves position-specific expression of a gene containing a homeobox. *Cell* **55**, 747-56 (1988).

78.   Euling, S. & Ambros, V. Reversal of cell fate determination in Caenorhabditis elegans vulval

development. *Development* **122**, 2507-15 (1996).

79.   Braendle, C. & Felix, M.A. Plasticity and errors of a robust developmental system in different

environments. *Dev Cell* **15**, 714-24 (2008).

80.   Karp, X. & Greenwald, I. Control of cell-fate plasticity and maintenance of multipotency by DAF-

16/FoxO in quiescent Caenorhabditis elegans. *Proc Natl Acad Sci U S A* **110**, 2181-6 (2013).

81.   Tenen, C.C. & Greenwald, I. Cell Non-autonomous Function of daf-18/PTEN in the Somatic Gonad

Coordinates Somatic Gonad and Germline Development in C. elegans Dauer Larvae. *Curr Biol* **29**,

1064-1072 e8 (2019).

82.   Hong, Y., Roy, R. & Ambros, V. Developmental regulation of a cyclin-dependent kinase inhibitor

controls postembryonic cell cycle progression in Caenorhabditis elegans. *Development* **125**, 3585-

97 (1998).

83.   Karp, X. & Ambros, V. Dauer larva quiescence alters the circuitry of microRNA pathways regulating

cell fate progression in C. elegans. *Development* **139**, 2177-86 (2012).

84.   O'Keeffe, C. & Greenwald, I. EGFR signal transduction is downregulated in C. elegans vulval

precursor cells during dauer diapause. *Development* **149**(2022).

85.   Colella, E., Li, S. & Roy, R. Developmental and Cell Cycle Quiescence Is Mediated by the Nuclear

Hormone Receptor Coregulator DIN-1S in the Caenorhabditis elegans Dauer Larva. *Genetics* **203**,

1763-76 (2016).

86.   Narbonne, P. & Roy, R. Inhibition of germline proliferation during C. elegans dauer development

requires PTEN, LKB1 and AMPK signalling. *Development* **133**, 611-9 (2006).

87.   Wong, C., Kadekar, P., Jurczak, E. & Roy, R. Germline stem cell integrity and quiescence are

controlled by an AMPK-dependent neuronal trafficking pathway. *PLoS Genet* **19**, e1010716

(2023).

88.   Vousden, K.H. & Ryan, K.M. p53 and metabolism. *Nat Rev Cancer* **9**, 691-700 (2009).

89.   van den Heuvel, S. Cell-cycle regulation. *WormBook*, 1-16 (2005).

90.   Derry, W.B. *et al.* Regulation of developmental rate and germ cell proliferation in Caenorhabditis

elegans by the p53 gene network. *Cell Death Differ* **14**, 662-70 (2007).

91.   Cameron, S., Clark, S.G., McDermott, J.B., Aamodt, E. & Horvitz, H.R. PAG-3, a Zn-finger

transcription factor, determines neuroblast fate in C. elegans. *Development* **129**, 1763-74 (2002).

92.   Thompson, K.W. *et al.* The Paired-box protein PAX-3 regulates the choice between lateral and

ventral epidermal cell fates in C. elegans. *Dev Biol* **412**, 191-207 (2016).

93.   Salser, S.J. & Kenyon, C. Patterning C. elegans: homeotic cluster genes, cell fates and cell

1774   migrations. *Trends Genet* **10**, 159-64 (1994).

1775   94.   Sternberg, P.W. Genetic control of cell type and pattern formation in Caenorhabditis elegans. *Adv*

1776   *Genet* **27**, 63-116 (1990).

1777   95.   Burglin, T.R. & Ruvkun, G. The Caenorhabditis elegans homeobox gene cluster. *Curr Opin Genet*

1778   *Dev* **3**, 615-20 (1993).

1779   96.   Jones, R.G. *et al.* AMP-activated protein kinase induces a p53-dependent metabolic checkpoint.

1780   *Mol Cell* **18**, 283-93 (2005).

1781   97.   Feng, Y., Williams, B.G., Koumanov, F., Wolstenholme, A.J. & Holman, G.D. FGT-1 is the major

1782   glucose transporter in C. elegans and is central to aging pathways. *Biochem J* **456**, 219-29 (2013).

1783   98.   Schulz, T.J. *et al.* Glucose restriction extends Caenorhabditis elegans life span by inducing

1784   mitochondrial respiration and increasing oxidative stress. *Cell Metab* **6**, 280-93 (2007).

1785   99.   Wu, Z. *et al.* Dietary Restriction Extends Lifespan through Metabolic Regulation of Innate

1786   Immunity. *Cell Metab* **29**, 1192-1205 e8 (2019).

1787   100.   Walker, M.D. *et al.* WormPaths: Caenorhabditis elegans metabolic pathway annotation and

1788   visualization. *Genetics* **219**(2021).

1789   101.   Li, Q. *et al.* IL-17 induces radiation resistance of B lymphoma cells by suppressing p53 expression

1790   and thereby inhibiting irradiation-triggered apoptosis. *Cell Mol Immunol* **12**, 366-72 (2015).

1791   102.   Chen, K. *et al.* IL-17 Receptor Signaling in the Lung Epithelium Is Required for Mucosal Chemokine

1792   Gradients and Pulmonary Host Defense against K. pneumoniae. *Cell Host Microbe* **20**, 596-605

1793   (2016).

1794   103.   Liu, Y. *et al.* p53 regulates hematopoietic stem cell quiescence. *Cell Stem Cell* **4**, 37-48 (2009).

1795   104.   Cheung, T.H. & Rando, T.A. Molecular regulation of stem cell quiescence. *Nat Rev Mol Cell Biol* **14**,

1796   329-40 (2013).

1797   105.   Das, S., Min, S. & Prahlad, V. Gene bookmarking by the heat shock transcription factor programs

1798   the insulin-like signaling pathway. *Mol Cell* **81**, 4843-4860 e8 (2021).

1799   106.   Das, S. *et al.* Serotonin signaling by maternal neurons upon stress ensures progeny survival. *Elife*

1800   **9**(2020).

1801   107.   Tatum, M.C. *et al.* Neuronal serotonin release triggers the heat shock response in C. elegans in

1802   the absence of temperature increase. *Curr Biol* **25**, 163-174 (2015).

1803   108.   Ooi, F.K. & Prahlad, V. Olfactory experience primes the heat shock transcription factor HSF-1 to

1804   enhance the expression of molecular chaperones in C. elegans. *Sci Signal* **10**(2017).

1805   109.   Venkatesh, H. & Monje, M. Neuronal Activity in Ontogeny and Oncology. *Trends Cancer* **3**, 89-112

1806   (2017).

1807   110.   Curry, R.N. & Glasgow, S.M. The Role of Neurodevelopmental Pathways in Brain Tumors. *Front*

1808   *Cell Dev Biol* **9**, 659055 (2021).

1809   111.   Gibson, D.G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat*

1810   *Methods* **6**, 343-5 (2009).

1811   112.   Nonet, M.L. *et al.* Caenorhabditis elegans rab-3 mutant synapses exhibit impaired function and

1812   are partially depleted of vesicles. *J Neurosci* **17**, 8061-73 (1997).

1813   113.   Nika, L., Gibson, T., Konkus, R. & Karp, X. Fluorescent Beads Are a Versatile Tool for Staging

1814   Caenorhabditis elegans in Different Life Histories. *G3 (Bethesda)* **6**, 1923-33 (2016).

1815   114.   Krueger, F. TrimGalore. (2021).

1816   115.   Kopylova, E., Noe, L. & Touzet, H. SortMeRNA: fast and accurate filtering of ribosomal RNAs in

1817   metatranscriptomic data. *Bioinformatics* **28**, 3211-7 (2012).

1818   116.   Yates, A.D. *et al.* Ensembl Genomes 2022: an expanding genome resource for non-vertebrates.

1819   *Nucleic Acids Res* **50**, D996-D1003 (2022).

1820   117.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

1821   118.   Liao, Y., Smyth, G.K. & Shi, W. The R package Rsubread is easier, faster, cheaper and better for

1822   alignment and quantification of RNA sequencing reads. *Nucleic Acids Research* **47**, e47-e47 (2019).

1823   119.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-

1824   seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1825   120.   Davis, P. *et al.* WormBase in 2022-data, processes, and tools for analyzing Caenorhabditis elegans.

1826   *Genetics* **220**(2022).

1827   121.   Cohen, S.M., Sun, J.J., Schroeder, F.C. & Sternberg, P.W. Transcriptional Response to a Dauer-

1828   Inducing Ascaroside Cocktail in Late L1 in C. elegans. (microPublication Biology, 2021).

1829   122.   Hu, M., Crossman, D., Prasain, J.K., Miller, M.A. & Serra, R.A. Transcriptomic Profiling of DAF-

1830   7/TGFβ Pathway Mutants in C. elegans. *Genes* **11**, 288 (2020).

1831   123.   Shen, L. & Sinai, M. GeneOverlap: Test and visualize gene overlaps. (2019).

1832   124.   Chen, H. VennDiagram: Generate High-Resolution Venn and Euler Plots. (2018).

1833   125.   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.

1834   *Innovation (N Y)* **2**, 100141 (2021).

1835   126.   Carlson, M. org.Ce.eg.db: Genome wide annotation for Worm. (2019).

1836   127.   Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M. KEGG for

1837   taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res* (2022).

1838   128.   Gu, Z. Complex heatmap visualization. *iMeta* **1**, e43 (2022).

1839   129.   Kolde, R. pheatmap: Pretty Heatmaps. (2019).

1840   130.   Chikka, M.R., Anbalagan, C., Dvorak, K., Dombeck, K. & Prahlad, V. The Mitochondria-Regulated

1841   Immune Pathway Activated in the C. elegans Intestine Is Neuroprotective. *Cell Rep* **16**, 2399-414

1842   (2016).

1843

1844

Acknowledgements

We thank the V.P. laboratory, Drs. Sarit Smolikove, Josep Comeron, Anna Malkova and Michael

Petrascheck for comments, Dr. Peter Ratcliffe, Oxford, for his generous gift of *CeHIF-1* antibody,

and Dr. Mario de Bono, Institute of Science and Technology Austria (ISTA) for *C. elegans* strains.

Nematode strains were provided by the Caenorhabditis Genetics Center (CGC) (funded by the

NIH Infrastructure Programs P40 OD010440). This work was supported by NIH R01 AG060616

(V.P.).

Author Contributions

All authors designed the study, performed experiments, analyzed data, and drafted the

manuscript. P.D.I coordinated the mammalian experiments; V.P. coordinated *C. elegans*

experiments and the project. All authors approved the final version of the manuscript.

1856

1857 **Declaration of Interests**

1858 The authors declare no competing interests

1859

Figure 1

Figure 2

a — 25°C, 48 hrs. post-egg lay

% dauers

***

0.0%

ilc-17.1   *ilc-17.1;*
              daf-16
              (mu86)

b

% larvae (nuclear DAF-16::GFP)

***

N2   *ilc-17.1*

20°C, 40 hrs post-egg lay

N2 — DAF-16::GFP — 5µm
ilc-17.1 — DAF-16::GFP — 5µm

c

Relative fold change (mRNA, 25°C)

□ *sod-3*
▨ *mtl-1*
▤ *lys-7*

*** ** *** *** *** *** ** ns

N2   *ilc-17.1*   *daf-16*   *ilc-17.1;*   *ilc-17.1*   *ilc-17.1*
                                daf-16   *rescued;*   *rescued;*
                                               (unc-54p)   *(ilc-17.1p)*

d — 25°C, 48 hrs. post-egg lay

% dauers

***   ***

0.0%   0.0%

N2   *daf-5*   *ilc-17.1*   *ilc-17.1;*
      (e1386)   *daf-5*

e — 25°C, 48 hrs. post-egg lay

% dauers

***   ***

0.0%

Control   Dafachronic   Control   Dafachronic
           Acid (50 nM)   Acid (50 nM)

|__ N2 __| |__ *ilc-17.1* __|

f — 25°C, 48 hrs. post-egg lay

% dauers

***   ***

0.0%   0.25%

N2   *cep-1*   *ilc-17.1*   *ilc-17.1;*
      (gk138)   *cep-1*

g — 0.5mm — *ilc-17.1*

h — 0.5mm — *ilc-17.1; cep-1*

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

a

ilc-17.1 (syb5296)   870bp

Deletion (2173 bp)

ilc-17.1 (syb5297)   792bp

Deletion (2188 bp)

b

Relative fold change (*ilc-17.1* mRNA, 20°C)

***

0.0014±0.00

N2   *ilc-17.1* (syb5296)

c

Egg lay on OP50 plate at 20ºC, 2 hrs. → Place embryos at **20°C** → SDS resistance after 48 and 72 hrs.

d

20°C, 48 and 72 hrs. post-egg lay

% SDS resistance

*** ** **

N2
ilc-17.1

48 hrs   72 hrs

e   50µm   dauer

alae

f   5µm   dauer

Buccal cavity   Metacorpus   Isthmus   Terminal bulb

g   5µm   L2 larva

Buccal cavity   Metacorpus   Isthmus   Terminal bulb

h   5µm   dauer

germline

i   5µm   L2 larva

germline

j

Average pumps/min

***

N2 (L2 larvae)   *ilc-17.1*

k

% dauers

*** ns

N2   *F25D1.3* (syb7367)   *ilc-17.1* (syb5296) 2X outcross

l

Egg lay on RNAi at 20ºC, 2 hrs. → Place embryos at **25°C** → SDS resistance after 48 hrs.

% SDS resistance

*

Ctrl RNAi   *ilc-17.1* RNAi
N2

Figure S2

Ilc-17.1 HA?

Figure S3

a

b

c

Figure S4

a Day-one adults placed at center of plate and control (H$_2$O), OP50, or odor placed at opposite end. Chemotaxis index, after 1 hour

Figure S5

a

% dauers vs RNAi: Ctrl, *daf-16*, *daf-12*, *cep-1*, *pha-4*, *jnk-1*, *aak-1*, *aak-2*, *par-4*, *strd-1*, *hsf-1*, *ncl-1*, *pmk-1*, *sta-2*, *daf-2*, *flp-8*, *daf-15*, *raga-1*, *rheb-1*, *rict-1*, *egl-9*, *vhl-1*, *skn-1*, *wdr-23*, *hif-1*

ilc-17.1; **25°C**

*** ns

b

Expression scale 0–15

Columns: *ilc-17.1*, *ilc-17.1* rescued, N2

Left: *daf-2*, *daf-28*, *ins-1*, *ins-10*, *ins-11*, *ins-12*, *ins-13*, *ins-14*, *ins-15*, *ins-16*, *ins-17*, *ins-18*, *ins-19*, *ins-2*, *ins-20*, *ins-21*, *ins-22*, *ins-23*, *ins-24*, *ins-25*, *ins-26*

Right: *ins-27*, *ins-28*, *ins-29*, *ins-3*, *ins-30*, *ins-31*, *ins-32*, *ins-33*, *ins-34*, *ins-35*, *ins-36*, *ins-37*, *ins-38*, *ins-39*, *ins-4*, *ins-5*, *ins-6*, *ins-7*, *ins-8*, *ins-9*

c

Relative fold change (mRNA, 25°C) — N2 (grey), *ilc-17.1* (red)

ins-3, *ins-4*, *ins-17*, *ins-19*, *ins-32*, *ins-37*

ns, **, ns

d

N2 / *ilc-17.1*, *daf-28p*::GFP, 5μm

Arbitrary Units (AU), ns — daf-28p::GFP, daf-28p::GFP; *ilc-17.1*

e

N2 / *ilc-17.1*, *ins-4p*::GFP, 5μm

% animals expressing ins-4p::GFP — ins-4p::GFP (grey), ins-4p::GFP; *ilc-17.1* (red)

neurons, muscle, epithelium, pharynx — ns

f

25°C, 48 hrs post-egg lay

% dauers — N2, *ilc-17.1* OE (*unc-54p*), *daf-2 (e1370)*, *ilc-17.1* OE (*unc-54p*); *daf-2*

*** ns

g

Expression scale 6–12

Columns: *ilc-17.1*, *ilc-17.1* rescued, N2

bra-1, *daf-1*, *daf-14*, *daf-3*, *daf-4*, *daf-5*, *daf-7*, *daf-8*

Figure S6

Figure S7

Figure S8

Figure S9

Figure S10

a

b