Apicomplexans predict thermal stress mortality in the Mediterranean coral

Paramuricea clavata

Anthony M Bonacolta[1,2], Jordi Miravall[3], Daniel Gómez-Gras[4,5], Jean-Baptiste Ledoux[6], Paula

López-Sendino[3], Joaquim Garrabou[3], Ramon Massana[3], Javier del Campo*[1,2]

[1]Department of Marine Biology and Ecology, Rosenstiel School of Marine, Atmospheric and

Earth Science, University of Miami, Miami, Florida, 33149 USA

[2]Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Barcelona, Catalonia, Spain

[3]Institut de Ciències del Mar – CSIC, Barcelona, Catalonia, Spain

[4]Hawaiʻi Institute of Marine Biology, University of Hawaiʻi at Mānoa, Kaneohe, HI, USA

[5]Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Institut de Recerca de la

Biodiversitat (IRBIO), Universitat de Barcelona, Barcelona, Spain

[6]CIIMAR/CIMAR, Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade

do Porto, Porto, Portugal.

*Corresponding Author: jdelcampo@ibe.upf-csic.es, Institut de Biologia Evolutiva (UPF-CSIC),

Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Catalonia, Spain

Abstract

The octocoral *Paramuricea clavata* is an ecosystem architect of the Mediterranean temperate reefs that is currently threatened by episodic mass mortality events related to global warming. Local average thermal regimes nor recent thermal history have been shown to play a significant role in population thermotolerance in this species. The microbiome, however, may play an active role in the thermal stress susceptibility of corals, potentially holding the answer as to why corals show differential sensitivity to heat-stress. To investigate this, the prokaryotic and eukaryotic microbiome of *P. clavata* collected from around the Mediterranean was characterized before experimental heat-stress to determine if its microbial composition influences the thermal response of the holobiont. We found that the prokaryotic community was not informative in predicting the thermal susceptibility of *P. clavata*. On the other hand, members of *P. clavata's* microeukaryotic community were significantly correlated with thermal stress sensitivity. Syndiniales from the *Dino-Group I Clade 1* were significantly enriched in thermally resistant corals, while the apicomplexan corallicolids were significantly enriched in thermally susceptible corals. Corallicolids are associated with 70% of coral genera around the world, yet the ecological role of this general anthozoan symbiont has yet to be determined. We hypothesize that *P. clavata* mortality following heat-stress may be caused by a shift from apparent commensalism to parasitism in the corallicolid-coral host relationship driven by the added stress. Our results show the potential importance of corallicolids and the rest of the microeukaryotic community of corals to understanding thermal stress response in corals and provides a useful tool to guide conservation efforts and future research into coral-associated microeukaryotes.

46 **Keywords:** Microbiome, 18S rRNA, Metabarcoding, Octocorallia, *P. clavata*, Apicomplexa,

47 Protists, Syndiniales, Cnidaria, Coral

Background

Paramuricea clavata is a slow-growing octocoral prevalent in the temperate Mediterranean Sea [1]. Gorgonians, such as *P. clavata*, are habitat-forming species of biodiversity-rich hard bottom benthic communities where they provide complex habitat for numerous organisms contributing significantly to the biodiversity of the region [2]. *P. clavata's* slow growth rate, long population turn-over time, high post-settlement mortality, and low genetic connectivity puts it at high-risk of extinction caused by natural and anthropogenic stressors [1]. *P. clavata* populations have been facing increased mortality due to episodic mass mortality events (MMEs) [3]. MMEs, which were once uncommon in temperate ecosystems such as the Mediterranean Sea, have been increasing since the turn of the century, with most of these occurring in the Western Mediterranean [4]. The increase in the frequency, duration, and intensity of marine heatwaves driven by global warming have been suggested to be the primary cause of the increase in MMEs in the Mediterranean. Thus, MMEs are leading to the current biodiversity crisis in the region [5,6]. *P. clavata* are heavily affected by MMEs with mortality sometimes reaching 80% in affected populations and recovery hindered by the recurrence of marine heatwaves. The loss of this habitat-forming species causes a functional shift in Mediterranean hard-bottom communities leading to simplified habitats with lower species diversity [7,8]. To prevent the significant decline of *P. clavata* and other habitat forming octocorals in the Mediterranean, the effect of thermal stress on the *P. clavata* holobiont must be better understood.

Thermal stress typically causes death in anthozoans by disrupting metabolic pathways, increasing physiological stress, decreasing host defense, causing energy constraints, and promoting the growth of opportunistic pathogens [9–11]. In *P. clavata,* exposure to higher

temperatures causes usually partial or full necrosis of colonies, where branches first change color

from red to grey before denudation of the skeleton [12]. This contrasts with "bleaching" which

takes place in zooxanthellate corals and is when the algal symbiont leaves the coral host during

periods of high heat-stress [13]. *P. clavata* is more sensitive to elevated seawater temperatures

than other gorgonians, marked by a significant loss in energy reserves, reduced feeding ability,

and partial mortality after heat-stress [14]. Furthermore, a companion study using the same *P.*

clavata from this research found no significant association between local average thermal

regimes nor recent thermal history and population thermotolerance; however, inter-individual

variation within populations was found [15]. A complete characterization of the *P. clavata*

microbiome before heat-stress can better elucidate the role of the microbiome to overall

holobiont health and fitness in the face of warming ocean temperatures.

Some octocoral-associated microorganisms are considered beneficial members of the

coral holobiont contributing to nitrogen fixation, sulfur-cycling, antibiotic production, and host

defense [16]. *P. clavata* has a consistent composition of the bacterial component of the

microbiome across regions, dominated (above ~90% relative abundance) by the common coral

symbiont *Endozoicomonas,* a gammaproteobacterium of the order Oceanospirillales [17,18].

Notably, *P. clavata*'s bacterial community differs significantly from the microbiome of the water

surrounding the organism and that of neighboring octocorals, such as *Eunicella* [19]. Other

bacteria commonly found within *P. clavata* include Alphaproteobacteria, Betaproteobacteria, and

Actinobacteria [17]. Despite the research into the bacterial community found in *P. clavata*

throughout the Mediterranean Sea, there has been no research into the microeukaryotic

community of this coral which constitutes a significant portion of the microbial diversity in other

corals and may contribute significantly to holobiont fitness in a changing climate [20].

Microeukaryotes (protists and unicellular fungi) are important components of many of

the Earth's ecosystems, including being the main contributors to oceanic primary production and

carbon mobilization in the microbial loop [21]. They may also play an important role in many

marine holobionts. The corallicolids, which are part of a lineage of parasitic alveolates known as

the Apicomplexa are associated with 70% of screened coral genera globally [22].

Apicomplexans also include such well-known human parasites like *Plasmodium*, the causative

agent of malaria, and *Toxoplasma*, a wide-spread mammal-infecting pathogen. While the

corallicolids, based on their distribution and presence in healthy individuals [23], seem to be

commensals in the coral microbiome, their evolutionary context suggests a more parasitic role

for this general anthozoan associate.

To better understand the microbiome of *P. clavata* and its ability to predict thermal stress

susceptibility we conducted a common garden heat-stress experiment using corals collected from

12 populations (148 total individuals) across a wide geographic range of *P. clavata* (See

Supplementary Figure 1, Additional File 1). We used an 18S rRNA and 16S rRNA gene

metabarcoding approach to get a holistic view of the microbes which may confer thermal

resiliency to *P. clavata.*

Methods

Sample Collection

Branch fragments (30 colonies per population) of healthy adult *P. clavata* were collected

from four localities (11 discrete populations) by scuba across the 18th-19th of September 2017

encompassing much of the known Mediterranean distribution of the octocoral. From Croatia, *P.*

clavata populations from Balun (Lat, Lon; 43.803889, 15.255000) and Mana (43.800273,

15.266389) were sampled. Three populations were sampled from Portofino (Liguria, Italy):

Altare (44.308918, 9.178645), Indiano (44.312476, 9.166889), and Lighthouse (44.298650,

9.219014). Three populations were sampled from Scandola (Corsica, France): Palazzinu

(42.379854, 8.549999), Gargallu (42.371839, 8.534394), and Palazzu (42.379873, 8.546163).

Three populations were sampled from Medes Islands (Catalonia, Spain): La Vaca (42.048047,

3.226322), Pota de Llop (42.0497, 3.2254), and Tascons (42.042189, 3.2269). Lastly, a

population thought to be *P. clavata* was sampled from an Atlantic locality from Sagres, Portugal

(37.012129, -8.924541). While underwater, samples were initially stashed in small plastic jars or

bags (no more than 10 per container) until being brought to the surface. Characteristic

information from each site can be found in Supplementary Table 1 (See Supplementary Table 1,

Additional File 2). Sampling was conducted as described in Gómez-Gras et al. 2022 [15]. In

short, 0.5 cm branch fragments were sampled for microbiome analyses at time of collection,

stored in a 2 mL cryovial containing 96% EtOH, flash-frozen using liquid nitrogen, and shipped

to the Institut de Ciències del Mar (CSIC) in Barcelona, Catalonia, Spain on ice. 10-17

individuals per population were included in microbiome analyses, depending on their

availability. 10 cm long coral specimens were maintained in a recirculating aquaria system

overnight, then transferred to 2 L bags containing water (10 coral fragments per bag). Bags were

placed into polystyrene boxes (3 bags per box) then shipped to Barcelona. For the coral

specimens from Medes and Scandola, the coral were transported in coolers immediately after

sampling, skipping the overnight aquaria step of the other populations. Transportation times

between the sampling location and CSIC ranged from several hours (Medes) to 36 hours

(Portofino).

Thermal Stress Experiment

The experiment was conducted in the Aquarium Experimental Zone of the Institut de Ciències del Mar in Barcelona (ICM-CSIC). Sets of 70 L aquarium tanks were divided between control and treatment groupings, each containing 3 replicate tanks per population with 10 coral fragments in each (30 individuals in total per population and experimental condition). To establish and maintain experimental conditions, each tank has individual heaters, temperature controllers, water pumps, and HOBO temperature data logs (recorded every 5 mins). Coral fragments were first acclimated to the aquarium environment for 2 months before experimentation began. Any coral showing signs of disease or necrosis during this period was promptly discarded before experimentation began. For the control tanks, water temperature was maintained at 16-18ºC for the entirety of the experiment. For the treatment tanks, the temperature increased step-wise from 18ºC to 25ºC over the course of three days. Water for both treatments was sourced from a buffer tank with 50 mm sand-filtered Mediterranean seawater pumped from 15 m depth. The coral fragments were fed three times per week until the end of the experiment. Feeding alternated between 3 mL of Bentos Nutrition Marine Active Supplement (Maim, Vic, Spain) and a tablet of frozen copepods (*Cyclops* spp.; Ocean nutrition, Antwerp, Belgium). Following 21 days of experimentation, the responses of individual coral fragments were assessed by measuring the percentage of tissue necrosis. Percent necrosis per fragment was assessed on days 14 and 21. Thermal response for each individual was categorized as follows by interpretation of the results of the two necrosis measurements: Super Resistant (0% necrosis), Resistant (< 25% necrosis), Tolerant (25-75% necrosis), Sensitive (>75% necrosis), and Hyper Sensitive (100% necrosis). Supplementary Table 1 (See Supplementary Table 1, Additional File 3) shows the thermal resiliency categories for each coral sample at the end of the experiment.

DNA Extraction and rRNA Gene Metabarcoding

P. clavata microbiome samples were thawed at room temperature and then processed using a modified DNeasy Mini Kit (QIAGEN) protocol. In short, samples were digested for 3 hours at 56ºC with slow agitation in 180 µl of tissue lysis buffer and 20 µl of proteinase K. During this time samples were mixed frequently by vortexing. Once the incubation ended 200 µl of Lysis buffer was added to the tissue lysates followed by 200 µl of 96-100% ethanol. Tissue lysates were then mixed by vortexing and the resulting mixtures were pipetted (discarding the central axis and the sclerites released from the tissues) into DNeasy Mini spin columns placed in 2 ml collection tubes. The columns were centrifuged for 1 minute at 8000 rpm and room temperature. The collection tubes with the mixture were discarded and the columns were placed in new collection tubes. The samples contained in the DNeasy Mini spin columns were then washed in two steps. First, 500 µl of Wash buffer 1 was added to the columns followed by a centrifugation of 1 minute at 8000 rpm and room temperature. Second, 500 µl of Wash buffer 2 was added and columns were centrifuged for 3 minutes at 14000 rpm and room temperature. Finally, to elute the DNA the columns were placed in 1.5ml Eppendorf tubes and 200 µl of Elution buffer was added to the center of the spin column membrane. Lastly, the columns were incubated for 1 minute at room temperature and then they were centrifuged for 1 minute at 8000 rpm at room temperature. DNA concentration from purified samples was measured using Qubit fluorometer. Concentrations that were higher than 1 ng/µl were considered acceptable for PCR. The V4 region of the prokaryotic 16S rRNA gene was amplified according to the Earth Microbiome Project protocol [53] and the V4 region of the eukaryotic 18S rRNA gene was amplified using a nested PCR approach validated in del Campo et al. 2019 [54]. Briefly, for the

18S rRNA metabarcoding, the samples were first amplified with UNonMet-PCR primers which

reduce the metazoan signal (18S-EUK581-F 5'-GTGCCAGCAGCCGCG-3', 18S-EUK1134-R

5'-TTTAAGTTTCAGCCTTGCG-3'), then reamplified with primers from Comeau et al. 2011

[55]. PCR mixture for one sample contained 4 µl 5X HS Buffer, 0.4 µl dNTP, 1 µl each of the

forward and reverse primers (diluted at 1:10), 0.6 µl DMSO, 0.2 µl Phusion DNA Polymerase, 3

µl of the sample DNA template, and reaction volumes made up to 20 µl with MilliQ water (9.8

µl). The PCR program of the thermal cycler was set up as follows: an initial denaturation step at

98ºC for 30 seconds, subsequent 35 cycles of denaturation at 98ºC for 10 seconds, annealing at

51.1ºC for 30 seconds and elongation at 72ºC for 1 minute, all of them followed by a final

elongation at 72ºC for 5 minutes. Gel electrophoresis was used to ensure PCR amplification was

successful. Electrophoresis gel was made with 1g of ultrapure agarose dissolved in 100 ml of 1X

Tris-acetate-EDTA buffer (TAE) and stained with 5 µl of SYBR safe DNA gel stain. The gelling

process has been carried out at room temperature for about 20 minutes and the agarose gel was

ran at 100 V for about 20 minutes at room temperature. The agarose gel was finally visualized

under UV light using a Bio-Rad Chemi XRS Gel Documentation system and Bio-Rad Quantity

One software. Those samples that didn't show a clear band on the agarose gel were amplified

again adding 0.2 µl MgCl2 50mM to the PCR mixture for one sample. The rest of the

components of the PCR mixture for one sample were maintained at the same volumes as before

and reaction volumes made up to 20 µl with MilliQ water (9.6 µl). The PCR program of the

thermal cycler was set up as before and PCR samples were checked again by an electrophoresis.

The PCR products were cleaned using QIAquick PCR Purification Kit (Qiagen). The buffer PB

of this purification kit was used with a pH indicator added to it. A volume of 8 µl of the PCR

sample was mixed with 40 µl of the Buffer PB and transferred to a QIAquick column. Columns

were centrifuged for 60 s to bind DNA to the column membrane. Next columns were washed by

adding 750 µl of Buffer PE and centrifuging them in two steps of 60 s each one. Finally, DNA

was eluted by adding 50 µl of Buffer PB and centrifuging the columns for 1 minute. DNA

concentrations from purified PCR samples were checked using Qubit fluorometer before being

sent to the Integrated Microbiome Resource facility at the Centre for Comparative Genomics and

Evolutionary Bioinformatics at Dalhousie University for further 18S rRNA amplification and

sequencing. 96 holes microplates were prepared in order to send them to Dalhousie University

for sequencing on a Illumina MiSeq using 300+300 bp paired-end V3 chemistry. Plates were

prepared following the receiver instructions. Plate holes were filled with 10 µl of each of the

DNA purified samples and the PCR purified samples and plates were sealed using an adhesive

plastic seal.

Microbiome Analysis

Raw sequencing data for this project has been deposited on NCBI (BioProject:

PRJNA928446). Primers were removed from reads using Cutadapt v3.1 [56]. The trimmed reads

were then processed in R using DADA2 [24]. For the 18S rRNA gene amplicons, forward reads

were truncated at 230 bp and reverse reads were truncated at 195 bp. Reads were then denoised

and joined into amplicon sequence variants (ASVs). Chimeras were removed using the

'removeBimeraDenovo' command with the "consensus" option. Taxonomy was assigned to 18S

ASVs in DADA2 using PR2 v4.13.0 [57]. For the 16S rRNA gene amplicons, forward reads

were truncated at 250 bp and reverse reads at 210 bp. Truncated reads were then denoised and

joined into ASVs before assigning taxonomy using SILVA v138 [58] in DADA2. Sequence

tables, taxonomy tables, and metadata for both the 16S and 18S rRNA gene amplicon datasets

were uploaded into a Phyloseq object in R for further filtering and analysis [59]. For the 16S

rRNA gene amplicons, ASVs corresponding to Chloroplast and Mitochondria were removed.

One sample was also removed for having under 250 reads. Additionally, ASVs were kept in the

dataset only if they were in the most abundant 99% of taxa in at least one of the samples.

Following this filtering 4 546 ASVs and 92 samples remained. The 18S amplicon were subjected

to the same filtering parameters except ASVs corresponding to Metazoa and Embryophyceae

were also removed from the dataset, resulting in 3181 ASVs across 138 samples. Bubble plots,

alpha-diversity, and beta-diversity figures were constructed using ggplot2 and tidyverse

packages [60]. Statistical analyses were conducted using the microbiomeMarker package and

vegan [61,62]. Analysis of similarities (ANOSIM) within vegan was used to test for differences

in beta-diversity between groups using 999 permutations. Linear Discriminant Analysis Effect

Size (LEfSe) [63] and Analysis of Compositions of Microbiomes with Bias Correction

(ANCOM-BC) [64] were used to find significantly enriched and differentially abundant taxa

between treatment groups. For the ANCOM-BC test, region of origin was treated as a fixed

variable and the p-values were adjusted using false discover rate ("fdr").

In order to place the corallicolid ASVs in a phylogenetic context, a multiple sequence

alignment (MSA) was created using publicly available full-length 18S rRNA sequences of

corallicolids and the program MAFFT. Abundant corallicolid ASVs (more than 50% of

corallicolid signal in at least one sample) were then extracted from the dataset. Hmmalign was

then used to map these ASVs to the full length 18S rRNA corallicolid MSA [65]. A maximum

likelihood phylogenentic tree was constructed with RAxML using this MSA (GTRCAT

substitution model, N = 1 000) [66]. The resulting tree was modified for publication using ggtree

[67]. For the dinoflagellate and ciliate trees, a similar approach was taken, except no publicly

available sequences were used in their construction and alignment.

To assess cross-domain positive-correlation in the occurrence patterns of ASVs, the 16S

and 18S rRNA gene amplicon tables were combined for samples present in both datasets. After

setting a filter prevalence threshold of 0.001, correlation patterns were inferred for the combined

dataset using SPARCC which considers the compositional nature of metabarcoding data [68]. A

network was then constructed using SpiecEasi with a p-value threshold of 0.05 (ref. 30). The top

families exhibiting positive co-occurrence relationships were plotted in a chordplot using the

MicroEco package [70].

RESULTS

The bacterial microbiome of *Paramuricea clavata*

For the 16S rRNA gene sequencing, 1332678 total reads were recovered after processing in DADA2 [24]. Following filtering, a total of 4540 Amplicon Sequence Variants (ASVs) across 92 samples were left for bacterial community analysis. The water and food microbiome samples were taxonomically distinct from the coral microbiome samples in this study (See Supplementary Figure 2, Additional File 4). The population originally sampled from Sagres, Portugal exhibited a distinct microbiome from the other *P. clavata* populations (Fig. 1) and was later determined to be a different species based on genomic work [25]**.** It was hence removed from further analysis. Oceanospirillales, mostly consisting of *Endozoicomonas*, constituted most of the bacterial community of *P. clavata* (Fig. 1). Other highly abundant bacterial orders include Alteromonadales, Rhodobacterales, and Flavobacteriales. A high number of unidentified bacterial ASVs were also present in the dataset. Subsequent BLAST analysis of these ASVs yielded no definitive taxonomic identification.

Figure 1. Relative abundance of microbial orders within *P. clavata* across its geographic distribution. According to 16S rRNA gene amplicons (A) and 18S rRNA gene amplicons (B) metabarcoding. *P. clavata* populations are listed from west to east, with major localities color coded. Sagres is denoted with a star as this population was excluded from further analysis since it is a different species. Microbial taxa are ordered taxonomically. Only those taxa with a median relative abundance above 1% are shown, the rest are listed as "other". IS is short for Incertae sedis, meaning there is

uncertainty in the taxonomic position within the broader taxonomic group. Relative abundance bubbles

are shown at 80% opacity and correspond to within domain (Prokaryotic or Eukaryotic) relative

abundance. The upper-bound of the standard deviation is shown by the opaquer outer bubble. The lower-

bound of the standard deviation is shown by a solid circle. Lower bounds with a negative value are not

shown.

Pair-wise analysis of the bacterial communities at the ASV-level between localities

showed significant differences in Shannon-Wiener alpha-diversity (Fig. 2). Additionally, the

beta-diversity of the bacterial community between localities was significantly different as well

(ANOSIM; $P = 0.001$; Fig. 3). Overall, the bacterial community of *P. clavata* clustered closely

based on geographic origin.

Figure 2. Shannon-Wiener Alpha Diversity of the prokaryome and eukaryome of *P. clavata* across sampled localities and thermal tolerance categories. Mean comparison p-values are added to significant group comparisons.

Figure 3. Aitchison distance PCAs of the prokaryome and eukaryome of *P. clavata* across sampled localities and thermal tolerance categories. Analysis of similarities (ANOSIM) within vegan was used to test for differences in beta-diversity between groups using 999 permutations. The beta-diversities of both the bacterial and microeukaryotic communities between localities were significant (ANOSIM; P = 0.001).

The dominance of *Endozoicomonas* (Proteobacteria; Oceanospirillales) in the bacterial community of *P. clavata* masked any fine-scale differences in the bacterial community composition across the range of thermal susceptibilities (Fig. 4). Analyzing the bacterial community with *Endozoicomonas* removed, did not reveal any noticeable pattern in taxon composition in relation to thermal stress susceptibility (See Supplementary Figure 3, Additional File 5). Additionally, an ANCOM-BC analysis of the Prokaryotic Orders corrected for the region of origin and false discovery rate did not identify any significant bacterial taxa between the thermal susceptibility categories.

Figure 4. Relative abundance of microbial orders within *P. clavata* across thermal susceptibility categories. According to 16S rRNA gene amplicons (A) and 18S rRNA gene amplicons (B) metabarcoding. The Sagres population was not included in this analysis (see Methods section). Hyper

Sensitive (n_{bac} = 29; n_{euk} = 39) corals were those which showed 100% necrosis during the heat experiment.

Sensitive (n_{bac} = 3; n_{euk} = 16) corals were those that showed < 75% necrosis. Tolerant (n_{bac} = 42; n_{euk} = 61)

corals were those that showed 25-75% necrosis. Resistant (n_{bac} = 3; n_{euk} = 7) corals showed < 25%

necrosis at the end of heat-stress. Super Resistant (n_{bac} = 4; n_{euk} = 5) corals showed 0% necrosis at the end

of heat-stress. Microbial taxa are ordered taxonomically. Only those taxa with a median relative

abundance above 1% are shown, the rest are listed as "other". IS is short for Incertae sedis, meaning there

is uncertainty in the taxonomic position within the broader taxonomic group. Relative abundance bubbles

are shown at 80% opacity and correspond to within domain relative abundance. The upper-bound of the

standard deviation is shown by the opaquer outer bubble. The lower-bound of the standard deviation is

shown by a solid circle. Lower bounds with a negative value are not shown.

The eukaryotic microbiome of *Paramuricea clavata*

For the 18S rRNA sequencing, 18955358 total reads were recovered, resulting in 3181

ASVs across 138 samples following filtering. This community is dominated by 3 major clades of

Alveolates: Apicomplexans, Ciliates, and Syndiniales (Fig. 1). One of the primary

microeukaryotic taxa present, Dino-Group-I (Dinoflagellata; Syndiniales), exhibited its highest

relative abundance in the more Eastern populations (Balun, Mana, & Lighthouse) (Fig. 1).

Corallicolids (Apicomplexa; Corallicolida) exhibited a high relative abundance across every

population sampled, excluding Lighthouse (Fig. 1). Additionally, three distinct genera of ciliates

(Ciliophora) were found across the surveyed geographic range of *P. clavata*, with *Licnophoria*

spp. being prevalent at Tascons, Lighthouse, and Altare (Fig. 1). *Hypotrichia* spp. were found

mainly in the Western populations (Tascons, Pota de Llop, La Vaca, Palazzu, Palazzinu, &

Gargallu) (Fig. 1). Lastly, *Suctoria* spp. were most prevalent within the *P. clavata* of the Medes

and Portofino localities (Fig. 1).

At the ASV-level, the locality-level microeukaryotic communities of *P. clavata* exhibited

higher Shannon-Wiener alpha-diversity compared to the bacterial communities (Fig. 2).

Additionally, the microeukaryotic community only showed a significant difference in the alpha-

diversity of *P. clavata* between Croatia vs Medes. Furthermore, the beta-diversity between

localities was significantly different for the microeukaryotic community (ANOSIM; P = 0.001;

Fig. 3).

Syndiniales were enriched in the super-resistant corals (LEfSe; P = 0.0016). Specifically,

Dino-Group I Clade 1 abundance in the holobiont was associated with the increased

thermotolerance in *P. clavata* and was enriched in the super-resistant individuals (LEfSe; P =

0.0027; Fig. 4). At the ASV level, ASV_20 and ASV_13 were the *Dino-Group I Clade 1* ASVs

which were highly abundant in the resistant groups relative to the other Alveolates (See

Supplementary Figure 4, Additional File 6). Concurrently, a high relative abundance of

corallicolids within the holobiont was associated with higher sensitivity to thermal stress (Fig. 4).

LEfSe analysis revealed that the *Corallicolidae* family was significantly enriched in the sensitive

corals (LEfSe; P = 0.015). ASV_6, within the COR3 clades of corallicolids, showed the highest

abundance amongst of the alveolates in the hyper-sensitive corals (See Supplementary Figure 4,

Additional File 6). ANCOM-BC analysis of the microeukaryotic Orders corrected for the region

of origin and false discovery rate also confirmed these associations, with corallicolids being

significantly more abundant (~2.3 natural log fold higher abundance; ANCOMBC, p_{adj} = 0.0003)

in the sensitive and hypersensitive corals compared to the resistant and hyper resistant ones,

while Dino-Groups I and II showed an opposite pattern (Fig. 5; ~1.4 natural log fold lower

abundance; ANCOMBC, p_{adj} = 0.04). On the other hand, ciliates showed a steady abundance

throughout thermal sensitivity groups, except for ASV_7 (*Ephelota mammillata*) which was

highly abundant in the sensitive and tolerant groupings (See Supplementary Figure 4, Additional

File 6). Alpha-diversity and beta-diversity analysis of both the bacterial community and

microeukaryotic community yielded no significant differences between any of the thermal

tolerance categories (Figs. 2 & 3).

Figure 5. ANCOMBC Significant Taxa. Significant differentially abundant microbial taxa between

sensitive (hyper sensitive + sensitive; n_{euk} = 55) vs resistant (super resistant + resistant; n_{euk} = 12) samples

according to ANCOMBC analysis while controlling for region of origin and false discovery rate. Natural

log fold differences in abundance for each significant taxa are shown with SD. Adjusted p-values are

shown. IS is short for Incertae sedis, meaning there is uncertainty in the taxonomic position within the

broader taxonomic group.

Phylogenetic placement of the microeukaryotes

Phylogenetic analysis of the most prevalent Alveolate ASVs found within *P. clavata*

revealed a diversity of clades associated with the octocoral across the surveyed geographic range.

ASV_6, which was the most abundant corallicolid ASV and found in most samples, was most

abundant in the Scandola populations (Fig. 6). An evolutionary placement algorithm based on

RAxML placed this ASV as closely related to MN022452_Corallicolidae_COR3_sp. (99.22%

Similarity), a sequence recovered from the deep-sea octocoral, *Callogorgia delta* in the Gulf of

Mexico. A second clade was identified that most likely belonged to the Corallicolidae COR2

group (Fig. 6). These ASVs (ASV_16, ASV_19, ASV_22) have a low relative abundance within

P. clavata but are found across most samples (Fig. 6). This group is most closely related to

MN022429_Corallicolidae_COR2_sp. (95.84% Similarity to ASV_16), a sequence recovered

from the deep-sea Antipatharian *Leiopathes glaberrima*, also from the Gulf of Mexico. Lastly, a

third clade of prevalent corallicolids was identified within *P. clavata* that shows close relation to

JX943883_Corallicolidae_COR1_sp. (93.04% Similarity to ASV_105), a sequence found in the

stony coral *Porites astreoides,* from the Bahamas.

Figure 6. RAxML maximum-likelihood tree of corallicolids. Primary corallicolid ASVs recovered

from *P. clavata* placed within the tree structure using EPA. Relative abundance of each ASV at each

respective population is shown in the heatmap. Populations are ordered from west to east, with major

localities color coded. Relative abundance (green) and sample prevalence (purple) of the ASV across the

entire dataset is shown in the barplots to the right.

403   Examination of the Dinoflagellate ASVs revealed distinct clades present within *P.*

clavata. *Dino-Group-I Clade 1* ASVs were most prevalent across the dataset with ASV_13 being

the most abundant across all localities except Scandola, where ASV_12 is most prevalent (See

Supplementary Figure 5, Additional File 7). The closest blast results for these sequences were to

uncultured microeukaryotes isolated from intertidal mudflat sediments and ocean surface water

from the East and South China Seas, respectively [26,27] (100% Similarity). One

Symbiodiniaceae ASV (ASV_43) was identified in some of the populations (Tascons, La Vaca,

Palazzinu, Lighthouse, Altare, Mana & Balun) despite *P. clavata* being azooxanthellate (See

Supplementary Figure 5, Additional File 7). The ASV from this study corresponds to

Symbiodinium Temperate Clade A which has been identified in Mediterranean corals before [28].

It's closest blast result from NCBI was to a *Symbiodinium sp. Temperate Clade A* sequence

(99.21% Similarity) recovered from the stony coral *Mussismilia hispida* [29].

415   The ciliate ASVs within the dataset revealed three primary clades found variably

abundant across the surveyed distribution of *P. clavata*. ASVs corresponding to *Holostichia*

diademata (Order: Hypotrichia) (ASV_8) and *Ephelota mammillata* (Order: Suctoria) (ASV_7)

were the most abundant and prevalent across the dataset overall (See Supplementary Figure 6,

Additional File 8). Interestingly, BLAST results for ASV_8, shows 100% similarity to

Holosticha pullaster (Order: Hypotrichia) isolated from a freshwater pond in the USA [30].

ASV_7 showed high similarity to *E. mammillata* recovered from *Tubularia* hydroids (98.51%

Similarity) [31]. *Licnophora macfarlandi* (Order: Licnophoria), was found across most

populations, with its highest abundance being in Mana, Croatia (See Supplementary Figure 6,

Additional File 8). Blast results show that these sequences were highly similar to *L. macfarlandi*

(92.66% Similarity to ASV_3) recovered from the fluid in the respiratory trees of the giant

California sea cucumber *Parastichopus californicus* [32].

Co-occurrence between members of the microbiome

A microbial co-occurrence network was constructed using SPARCC to analyze the

positive correlations between the major microbial families of the *P. clavata* holobiont.

Corallicolidae & *Dino-Group I Clade 1* had the strongest significant co-association between all

the major taxa (Fig. 7). This co-association is of note when you consider these two group's

antagonistic relative abundance patterns across the thermal susceptibility groups (Fig. 4). The

three ciliate families also showed some significant positive associations with the corallicolids

and *Dino-Group I Clade 1*. *Endozoicomonas*, *Pseudoalteromonadaceae*, *Vibrionaceaee*, and

Flavobacteriaceae were the primary bacterial families that showed significant co-occurrence

with the different microeukaryotic members of the *P. clavata* holobiont (Fig. 7).

Figure 7. Chord plot of *P. clavata* microbiome. Chord plot of significant positive co-association patterns between the top microbial families across both the 16S and 18S rRNA gene amplicon datasets for *P. clavata*. Interactions were constructed using SPARCC and SpiecEasi with a p-value threshold of 0.05 for significance. Self-interactions have been removed. Eukaryotic families are shown in black text, while Prokaryotic families are shown in red text. IS is short for Incertae sedis, meaning there is uncertainty in the taxonomic position within the broader taxonomic group.

DISCUSSION

The bacterial community of *P. clavata* showed significant differences in its make-up from location to location, most likely attributed to ASV-level diversity within the

Endozoicomonas which dominated the bacteriome of each population (Figs. 1, 2, & 4). This is

consistent with previous work into the bacterial component of the *P. clavata* holobiont [18,33].

Additionally, the prevalence of unidentified bacterial ASVs in the dataset points to the

underrepresentation of octocoral-associated bacteria in the SILVA reference database. These

results exemplify the need to identify more bacteria associated with a healthy symbiotic state in

coral holobionts.

The microeukaryotic community of *P. clavata* is dominated by three major groups of

alveolates: *Apicomplexa* (mostly corallicolids), *Dinoflagellata* (mostly *Syndiniales*), and

Ciliophora. This community shows variability in its taxonomic make-up at the order level from

location to location, while also showing significant differences in beta-diversity at the ASV level

(Figs. 1 & 3). Each of the alveolates found in *P. clavata* showed locality-specific differences,

highlighting the specificity and potential co-adaptations of these associations in coral holobionts

(Fig. 6, and see Supplementary Figures 5 & 6, Additional Files 7 & 8). Despite being within a

clade of obligate parasites, corallicolids are general anthozoan associates found world-wide in

healthy corals, even at depths as deep as 1400 m, and in some cases can be as abundant as

Symbiodiniaceae [34,35]. They tend to be the most abundant and prevalent microeukaryotic

coral symbiont in azooxanthellate corals. The ones recovered from this study showed close

relation to those found within deep-sea cnidarians, including octocorals and antipatharians, of the

Gulf of Mexico from Vohsen *et al.* [36] (Fig. 6). A third clade of corallicolids were closely

related to one recovered from the shallow water Caribbean coral *Porites astreoides* [37]. This

diversity of corallicolids present within *P. clavata,* alone, hints at the potential global diversity of

corallicolids in other anthozoan hosts.

Syndiniales, which have been found in high prevalence in reef environments before, also

constituted a major portion of the eukaryotic microbiome. They are an Order of parasites found

at the base of the dinoflagellates which can infect a variety of marine macro- and microscopic

organisms [38]. *Syndiniales* from *Dino-Group I* are one of the main microeukaryotic taxa found

in the hard coral *Pocillopora damicornis* [39], however this group is also ubiquitous in global

plankton communities [40]. Additionally, *Dino-Group 1 Syndiniales* with identical 18S rRNA

V4 sequences may associate with different hosts, suggesting a generalist host specificity for this

clade or the need for a different marker gene to decipher its true diversity [41]. *Syndiniales* have

the potential to disrupt nutrient cycling [42], and thus could likely play a role in (de)stabilizing

symbiotic relationships, which rely on nutrient dynamics, in a tight-knit symbiotic system such

as a coral holobiont. Their differential abundance within *P. clavata* across different thermal

susceptibility categories highlights the necessity of studying the role of parasitism in the

nutritional dynamics of marine holobionts, especially in the face of stressors. Hudson et al. 2006

noted that a healthy system may be one that is rich in parasitic species due to the impact of

parasite-mediated effects on population dynamics, interspecific competition, and energy flow

[43]. Applying this perspective to coral holobionts may help us better understand the higher

thermal tolerance of *P. clavata* rich in potential parasites such as the corallicolids or *Syndiniales*.

The high abundance of these microbes within the *P. clavata* holobiont may contribute to a

healthy and resilient system by influencing the microbial community dynamics when confronted

with heat-stress.

Within the ciliate lineages recovered from *P. clavata*, three genera (*Licnophora*,

Holosticha, & *Ephelota*) showed variable abundances across the surveyed distribution of the

coral host. *L. macfarlandi*, has been found as an endosymbiont in the respiratory trees of

holothuroids, but also in association with corals, likely as a secondary invader hunting other

microbes [44,45]. It is a noted predator of protozoans and is speculated to be attracted to other

ciliate species found at the interface of coral lesions [45]. *L. macfarlandi* within *P. damicornis*

also exhibited geographic variability in abundance [39], which may depend upon the presence of

other ciliates on which it preys. Another ciliate, *H. diademata,* has also been found within corals

often crawling slowly on the coral surface [45]. It is a noted bacterivore so it may feed upon the

bacteria of the coral, exerting top-down control upon that portion of the microbiome, and

therefore influencing the cycling of certain nutrients in the holobiont. The presence of

bacterivores, such as ciliates, may increase microbial activity and turnover within the ecosystem

[46]. While the previous two ciliates are predominately associated with diseased corals [47], the

third ciliate genera, *Ephelota,* has never previously been described in corals; however, suctorian

ciliates are typically commensal and have previously been recovered as epibionts of crustaceans,

mollusks, fishes, and even other protists [48]. Thus, it may be using the stability and structure

provided by the *P. clavata* host to stabilize itself in the water column and capture prey more

readily.

This study identified the key microbes associated with *P. clavata,* an important habitat-

forming species impacted by MMEs, and provides evidence of the microbiome's potential role in

the thermal susceptibility of the host. In *P. clavata*, the overwhelming abundance of

Endozoicomonas masked any major differences in bacterial composition. On the other hand, the

microeukaryotic community has distinct taxonomic composition associated with different

degrees of thermal tolerance. *Dino-Group I Clade 1* was significantly enriched in thermally

tolerant corals, while corallicolids were significantly enriched in thermally sensitive corals.

These results may explain inter-individual variation in thermotolerance within *P. clavata*

populations observed by Gómez-Gras et al. 2022 [15]. How the relative abundance of these

apicomplexans confers thermal susceptibility to the coral host is not yet understood. There's the

possibility that the symbiosis between corallicolids and their coral host shifts from an apparent

commensalism towards parasitism as added stressors, such as heat, breakdown the relationship.

The context-dependency of this relationship can explain the abundance of corallicolids in

apparently healthy coral. Briefly, the coral host may have evolved resistance to corallicolids at a

significant fitness cost leading to a state of commensalism without the presence of addition

stressors; however, heat-stress may cause these corallicolid-tolerant corals to exhibit higher

mortality as the apicomplexans shift towards a parasitc relationship with the host, similar to the

apparent commensalism of parasites idea proposed by Miller et al. 2006 [49]. This potential

interaction and the widespread distribution of the corallicolids would have significant

implications when interpreting threats to corals in the context of warming ocean temperatures.

Showing strong co-association with the corallicolids were the *Syndiniales*, specifically

those belonging to *Dino-Group I Clade 1* (Fig. 7). These parasites are known to infect a variety

of multicellular and microbial hosts and have previously been recovered from coral reef

sediments, plankton, and corals [50]. Members include microeukaryotic parasites such as

Duboscquella sp. which is known to parasitize ciliates [51]. This opens the possibility that they

may be infecting either the coral host, it's microeukaryotic members, or both. While significant

positive co-association between *Dino-Group I* and the ciliates recovered from *P. clavata* was

apparent in the network analysis, the association between *Dino-Group I* and the corallicolids was

most prevalent (Fig. 7). While this can simply be the result of fluctuations in two of the most

highly abundant microeukaryotes found in the dataset, it may also represent an interesting case

of parasitism of a coral endosymbiont by another member of the microeukaryotic community. Its

association with the corallicolids hints at a possible positive role of this parasite preventing the

overabundance of corallicolid growth in the coral holobiont which may be associated with

decreased thermal tolerance. *Syndiniales* have been proposed to have the capacity to control the

sizes of their host populations since a singular infection can produce hundreds of dinospores to

further infect new hosts [52]. The vast diversity of known microeukaryotic hosts of *Syndiniales*

combined with the larger cell size of corallicolids (~12 um in length [22]) supports the

possibility of this relationship. Possible explanations for this interaction could be top-down

control of the corallicolid population by the *Syndiniales*, niche competition, easier co-infection of

an already infected host, or excess dinospore production by *Syndiniales*. In either case, the

relationship between *Dino-Group I*, the corallicolids and the coral host needs to be investigated

further to decipher the ecological relevance of this potentially pivotal interaction within the coral

holobiont.

Conclusions

We have shown here that a deeper characterization of the, often overlooked,

microeukaryotic portion of the coral microbiome can have major implications on coral health in

a warming ocean. The recently described corallicolids are significantly correlated with thermally

susceptible *P. clavata*, yet their role within coral holobionts globally has yet to be uncovered.

Along with other genetic, environmental, and physiological mechanisms the microbiome,

especially the microeukaryotic portion, should be considered a vital component when assessing

coral thermal response. Future research should focus on identifying the mechanisms by which

these microbes may confer thermal resiliency to individual corals and how this knowledge can

best be used to manage fragile coral ecosystems.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Competing Interests

The authors declare that they have no competing interests.

Funding

This study has been supported by project PID2020-118836GA-I00 financed by MCIN / AEI /10.13039/501100011033 awarded to Javier del Campo and startup funds from the University of Miami,Rosenstiel School of Marine, Atmospheric and Earth Sciences This research was supported by the strategic funding UIDB/04423/2020, UIDP/04423/2020 and 2021.00855. CEECIND through national funds provided by FCT – Fundação para a Ciência e a Tecnologia. Joaquim Garrabou and Paula López-Sendino, Daniel Gómez Gras acknowledges the funding by the "Severo Ochoa Centre of Excellence" (CEX2019-000928-S), Interreg-Med

Authors' Contributions

JdC conceptualized the study. JdC, DG, JBL, PL, JG collected the samples and

performed the experiment. JdC and JM prepped the samples for sequencing. AMB conducted the

formal analysis, generated the figures, and wrote the manuscript. JdC, DG, JBL, PL, JG, and

R.M provided the laboratory resources for the experiment. All authors read and approved the

final manuscript.

Acknowledgments

We appreciate the contributions of the following scientists in helping with our sampling

effort: Bensoussan, N, Cerrano C, Ferretti E, Kipson S, Bakran-Petricioli T, Serrao EA, Paulo D,

Boavida J, Montero-Serra I, López-Sanz A, Milanese M, Linares C.

References

1. Linares C, Doak DF, Coma R, Díaz D, Zabala M. Life history and viability of a long-lived

marine invertebrate: The octocoral Paramuricea clavata. Ecology. 2007;88:918–28.

2. Ballesteros E. Mediterranean coralligenous assemblages: A synthesis of present knowledge.

Oceanogr Mar Biol. 2006;44:123–95.

3. Linares C, Doak DF. Forecasting the combined effects of disparate disturbances on the

persistence of long-lived gorgonians: A case study of Paramuricea clavata. Mar Ecol Prog Ser.

2010;402:59–68.

4. Garrabou J, Gómez-Gras D, Ledoux JB, Linares C, Bensoussan N, López-Sendino P, et al.

Collaborative Database to Track Mass Mortality Events in the Mediterranean Sea. Front Mar Sci.

2019;6.

5. Garrabou J, Gómez-Gras D, Medrano A, Cerrano C, Ponti M, Schlegel R, et al. Marine

heatwaves drive recurrent mass mortalities in the Mediterranean Sea. Glob Chang Biol.

2022;28:5708–25.

6. Garrabou J, Ledoux J-B, Bensoussan N, Gómez-Gras D, Linares C. Sliding Toward the of

Mediterranean Coastal Marine Rocky Ecosystems. 2021.

7. Verdura J, Linares C, Ballesteros E, Coma R, Uriz MJ, Bensoussan N, et al. Biodiversity loss

in a Mediterranean ecosystem due to an extreme warming event unveils the role of an

engineering gorgonian species. Sci Rep. 2019;9:1–11.

8. Gómez-Gras D, Linares C, Dornelas M, Madin JS, Brambilla V, Ledoux JB, et al. Climate

change transforms the functional identity of Mediterranean coralligenous assemblages. Ecol Lett.

2021;24:1038–51.

9. van de Water JAJM, Chaib De Mares M, Dixon GB, Raina JB, Willis BL, Bourne DG, et al.

Antimicrobial and stress responses to increased temperature and bacterial pathogen challenge in

the holobiont of a reef-building coral. Mol Ecol. 2018;27:1065–80.

10. Hou J, Xu T, Su D, Wu Y, Cheng L, Wang J, et al. RNA-seq reveals extensive transcriptional

response to heat stress in the stony coral Galaxea fascicularis. Front Genet. 2018;9:1–10.

11. Li J, Long L, Zou Y, Zhang S. Microbial community and transcriptional responses to

increased temperatures in coral Pocillopora damicornis holobiont. Environ Microbiol.

2021;23:826–43.

12. Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P, Cigliano M, et al. Mass

mortality in Northwestern Mediterranean rocky benthic communities: Effects of the 2003 heat

wave. Glob Chang Biol. 2009;15:1090–103.

13. Baker AC, Glynn PW, Riegl B. Climate change and coral reef bleaching: An ecological

assessment of long-term impacts, recovery trends and future outlook. Estuar Coast Shelf Sci.

Elsevier Ltd; 2008;80:435–71.

14. Tignat-Perrier R, van de Water JAJM, Guillemain D, Aurelle D, Allemand D, Ferrier-Pagès

C. The Effect of Thermal Stress on the Physiology and Bacterial Communities of Two Key

Mediterranean Gorgonians. Appl Environ Microbiol. American Society for Microbiology;

2022;88.

15. Gómez-Gras D, Bensoussan N, Ledoux JB, López-Sendino P, Cerrano C, Ferretti E, et al.

Exploring the response of a key Mediterranean gorgonian to heat stress across biological and

spatial scales. Sci Rep. Nature Publishing Group UK; 2022;12:1–14.

16. Van De Water JAJM, Allemand D, Ferrier-Pagès C. Host-microbe interactions in octocoral

holobionts - recent advances and perspectives. Microbiome. Microbiome; 2018;6:1–28.

17. Vezzulli L, Pezzati E, Huete-Stauffer C, Pruzzo C, Cerrano C. 16SrDNA Pyrosequencing of

the Mediterranean Gorgonian Paramuricea clavata Reveals a Link among Alterations in Bacterial

Holobiont Members, Anthropogenic Influence and Disease Outbreaks. PLoS One. 2013;8:1–10.

18. La Rivière M, Garrabou J, Bally M. SPATIAL AND TEMPORAL ANALYSIS OF

BACTERIAL DIVERSITY ASSOCIATED WITH THE MEDITERRANEAN GORGONIAN

PARAMURICEA CLAVATA. Rapp Comm int Mer medit. 2010;13007.

19. van de Water JAJM, Melkonian R, Voolstra CR, Junca H, Beraud E, Allemand D, et al.

Comparative assessment of Mediterranean gorgonian-associated microbial communities reveals

conserved core and locally variant bacteria. Microb Ecol. 2017;73:466–78.

20. Ainsworth TD, Fordyce AJ, Camp EF. The Other Microeukaryotes of the Coral Reef

Microbiome. Trends Microbiol. 2017. p. 980–91.

21. Corliss JO. Biodiversity and biocomplexity of the protists and an overview of their

significant roles in maintenance of our biosphere. Acta Protozool. 2002;41:199–219.

22. Kwong WK, Irwin NATT, Mathur V, Na I, Okamoto N, Vermeij MJAA, et al. Taxonomy of

the Apicomplexan Symbionts of Coral, including Corallicolida ord. nov., Reassignment of the

Genus Gemmocystis, and Description of New Species Corallicola aquarius gen. nov. sp. nov.

and Anthozoaphila gnarlus gen. nov. sp. nov. J Eukaryot Microbiol. 2021;68:1–9.

23. Kwong WK, Mathur V, Vermeij MJA, Keeling PJ. A widespread coral-infecting

apicomplexan contains a plastid encoding chlorophyll biosynthesis. 2018;

24. Ul-Hasan S, Bowers RM, Figueroa-Montiel A, Licea-Navarro AF, Michael Beman J, Woyke

T, et al. Community ecology across bacteria, archaea and microbial eukaryotes in the sediment

and seawater of coastal Puerto Nuevo, Baja California. PLoS One. 2019;14:1–7.

25. Campo J del, Pons MJ, Herranz M, Wakeman KC, Valle J del, Vermeij MJAA, et al.

Validation of a universal set of primers to study animal-associated microeukaryotic communities.

Environ Microbiol. John Wiley & Sons, Ltd; 2019;21:3855–61.

26. Comeau AM, Li WKW, Tremblay JÉ, Carmack EC, Lovejoy C. Arctic ocean microbial

community structure before and after the 2007 record sea ice minimum. PLoS One. 2011;6.

27. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal. EMBnet Stichting; 2011;17:10.

28. Callahan BJ, Mcmurdie PJ, Rosen MJ, Han AW, A AJ. DADA2: High resolution sample

inference from Illumina amplicon data. Nat Methods. 2016;13:581–3.

29. Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, et al. The Protist Ribosomal

Reference database (PR2): a catalog of unicellular eukaryote small sub-unit rRNA sequences

with curated taxonomy. Nucleic Acids Res. Oxford University Press; 2013;41:D597-604.

30. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal

RNA gene database project: Improved data processing and web-based tools. Nucleic Acids Res.

2013;41:590–6.

31. McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. PLoS One. 2013;8.

32. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the

Tidyverse. J Open Source Softw. 2019;4:1686.

33. Cao Y. Tools for microbiome marker identification • microbiomeMarker [Internet]. [cited

2022 Apr 25]. Available from:

https://yiluheihei.github.io/microbiomeMarker/articles/microbiomeMarker-vignette.html

34. Oksanen J, Blanchet F. Package "vegan." cran.ism.ac.jp. 2022;

35. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic

biomarker discovery and explanation. Genome Biol. 2011;12:R60.

36. Lin H, Peddada S Das. Analysis of compositions of microbiomes with bias correction. Nat

Commun. Springer US; 2020;11:1–11.

37. Zhang Z, Wood WI. A profile hidden Markov model for signal peptides generated by

HMMER. Bioinformatics. 2003;19:307–8.

38. Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics. 2014;30:1312–3.

39. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. Ggtree: an R Package for Visualization and

Annotation of Phylogenetic Trees With Their Covariates and Other Associated Data. Methods

Ecol Evol. 2017;8:28–36.

40. Friedman J, Alm EJ. Inferring Correlation Networks from Genomic Survey Data. PLoS

Comput Biol. 2012;8:1–11.

41. Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA. Sparse and

Compositionally Robust Inference of Microbial Ecological Networks. PLoS Comput Biol.

2015;11:1–25.

42. Liu C, Cui Y, Li X, Yao M. Microeco: An R package for data mining in microbial

community ecology. FEMS Microbiol Ecol. Oxford University Press; 2021;97:1–9.

43. Coelho M, Pearson G, Boavida J, Paulo D, Aurelle D, Daniel G, et al. Not out of the

Mediterranean□: Atlantic populations of the gorgonian Paramuricea clavata are a separate sister

species under further lineage diversification. Mol Ecol. 2023;1–18.

44. Wu W, Huang B, Liao Y, Sun P. Picoeukaryotic diversity and distribution in the subtropical-

tropical South China Sea. FEMS Microbiol Ecol. 2014;89:563–79.

45. Ma Y, Ouyang J, Wei N, Zhang S, Yan X, Xiong H. Microeukaryotic community structure

and dynamics during macrobenthos bioremediation in intertidal mudflat sediments of Sansha

Bay, China. Jundishapur J Microbiol. 2017;10.

46. Visram S, Wiedenmann J, Douglas AE. Molecular diversity of symbiotic algae of the genus

Symbiodinium (Zooxanthellae) in cnidarians of the Mediterranean Sea. J Mar Biol Assoc United

Kingdom. 2006;86:1281–3.

47. Picciani N, de Lossio e Seiblitz IG, de Paiva PC, e Castro CB, Zilberberg C. Geographic

patterns of Symbiodinium diversity associated with the coral Mussismilia hispida (Cnidaria,

Scleractinia) correlate with major reef regions in the Southwestern Atlantic Ocean. Mar Biol.

Springer Berlin Heidelberg; 2016;163:1–11.

48. Chen L, Dong J, Wu W, Xin Y, Warren A, Ning Y, et al. Morphology and molecular

phylogeny of a new hypotrich ciliate, Anteholosticha songi nov. spec., and an American

population of Holosticha pullaster (Müller, 1773) Foissner et al., 1991 (Ciliophora, Hypotrichia).

Eur J Protistol. Elsevier GmbH; 2020;72:125646.

49. Riley JL, Katz LA. Widespread distribution of extensive chromosomal fragmentation in

ciliates. Mol Biol Evol. 2001;18:1372–7.

50. Lynn DH, Strüder-Kypke M. Phylogenetic Position of Licnophora, Lechriopyla, and

Schizocaryum, Three Unusual Ciliates (Phylum Ciliophora) Endosymbiotic in Echinoderms

(Phylum Echinodermata). J Euk. 2002;49:460–8.

51. La Rivière M, Roumagnac M, Garrabou J, Bally M. Transient Shifts in Bacterial

Communities Associated with the Temperate Gorgonian Paramuricea clavata in the

Northwestern Mediterranean Sea. PLoS One. 2013;8.

52. Kwong WK, del Campo J, Mathur V, Vermeij MJA, Keeling PJ, del Campo J, et al. A

widespread coral-infecting apicomplexan with chlorophyll biosynthesis genes. Nature. Springer

US; 2019;568:103–7.

53. Vohsen SA, Anderson KE, Gade AM, Gruber-Vodicka HR, Dannenberg RP, Osman EO, et

al. Deep-sea corals provide new insight into the ecology, evolution, and the role of plastids in

widespread apicomplexan symbionts of anthozoans. Microbiome. Microbiome; 2020;8:1–15.

54. Vohsen SA, Anderson KE, Gade AM, Gruber-Vodicka HR, Dannenberg RP, Osman EO, et

al. Deep-sea corals provide new insight into the ecology, evolution, and the role of plastids in

widespread apicomplexan symbionts of anthozoans. Microbiome. Microbiome; 2020;8:1–15.

55. Kirk NL, Thornhill DJ, Kemp DW, Fitt WK, Santos SR. Ubiquitous associations and a peak

fall prevalence between apicomplexan symbionts and reef corals in Florida and the Bahamas.

Coral Reefs. 2013;32:847–58.

56. Guillou L, Viprey M, Chambouvet A, Welsh RM, Kirkham AR, Massana R, et al.

Widespread occurrence and genetic diversity of marine parasitoids belonging to Syndiniales

(Alveolata). Environ Microbiol. 2008;10:3349–65.

57. Clerissi C, Brunet S, Vidal-Dupiol J, Adjeroud M, Lepage P, Guillou L, et al. Protists within

corals: the hidden diversity. Front Microbiol. Frontiers; 2018;9:2043.

58. Engelen S, Hingamp P, Sieracki M, Vargas C, Audic S, Henry N, et al. Eukaryotic plankton

diversity in the sunlit ocean. Science (80-). 2015;348:1261605-1/11.

59. Clarke LJ, Bestley S, Bissett A, Deagle BE. A globally distributed Syndiniales parasite

dominates the Southern Ocean micro-eukaryote community near the sea-ice edge. ISME

Journalp. Springer US; 2019;13:734–7.

60. Chambouvet A, Morin P, Marie D, Guillou L. Control of toxic marine dinoflagellate blooms

by serial parasitic killers. Science (80-). 2008;322:1254–7.

61. Hudson PJ, Dobson AP, Lafferty KD. Is a healthy ecosystem one that is rich in parasites?

Trends Ecol Evol. 2006;21:381–5.

62. Lynn DH, Str M. Scuticociliate Endosymbionts of Echinoids (Phylum Echinodermata):

Phylogenetic Relationships among Species in the Genera Entodiscus , Plagiopyliella ,

Thyrophylax , and Entorhipidium (Phylum Ciliophora) Author (s): Denis H . Lynn and

Michaela Strüde. 2022;91:1190–9.

63. Sweet MJ, Séré MG. Ciliate communities consistently associated with coral diseases. J Sea

Res. Elsevier B.V.; 2016;113:119–31.

64. Trap J, Bonkowski M, Plassard C, Villenave C, Blanchart E. Ecological importance of soil

bacterivores for ecosystem functions. Plant Soil. 2016;398:1–24.

65. Ravindran C, Raveendran HP, Irudayarajan L. Ciliated protozoan occurrence and association

in the pathogenesis of coral disease. Microb Pathog. Elsevier Ltd; 2022;162:105211.

66. Dovgal I V. Evolution, phylogeny and classification of Suctorea (Ciliophora). Protistology.

2002;2.

67. Miller MR, White A, Boots M. THE EVOLUTION OF PARASITES IN RESPONSE TO

TOLERANCE IN THEIR HOSTS: THE GOOD, THE BAD, AND APPARENT

COMMENSALISM. Evolution (N Y). John Wiley & Sons, Ltd; 2006;60:945–56.

68. Clerissi C, Brunet S, Vidal-Dupiol J, Adjeroud M, Lepage P, Guillou L, et al. Protists within

corals: the hidden diversity. Front Microbiol. 2018;9:2043.

69. Harada A, Ohtsuka S, Horiguchi T. Species of the Parasitic Genus Duboscquella are

Members of the Enigmatic Marine Alveolate Group I. Protist. Urban & Fischer; 2007;158:337–

47.

70. Jephcott TG, Alves-de-Souza C, Gleason FH, van Ogtrop FF, Sime-Ngando T, Karpov SA,

et al. Ecological impacts of parasitic chytrids, syndiniales and perkinsids on populations of

marine photosynthetic dinoflagellates. Fungal Ecol. Elsevier; 2016;19:47–58.

Additional Information

Supplementary Figure 1, Additional File 1. Map showing locations of sampled localities of ***Paramuricea clavata***. Sagres is denoted by a star as this population was later determined to be a different species.

787
788
789

Supplementary Table 1, Additional File 2, Sample Metadata. Excel format (.xls). Information on each coral specimen analyzed in this study including thermal-tolerance categories, necrosis percentages, origin name and location, as well as environmental characteristics of population like depth and temperature at time of sampling.

793
794
795
796

Supplementary Table 2, Additional File 3, Thermal Categories System. Excel format (.xls). Information on how each coral's mortality was assessed and categorized at days 14 and 21 of the heat-stress experiment. This table also shows how these two measurements were interpreted for the final thermal tolerance category for the coral fragment.

798
799
800
801

804 **Supplementary Figure 2, Additional File 4. Stacked bar plots comparing the microbial**

805 **order composition of *P. clavata*.** The prokaryotic (A) and eukaryotic (B) communities of the

806 coral vs. the water it was sampled from is depicted. Only those taxa with a median relative

807 abundance above 1% are shown, the rest are listed as "Below Threshold".

Supplementary Figure 3, Additional File 5. Relative abundance of bacterial orders within

P. clavata* across thermal tolerance categories with the highly abundant *Endozoicomonas

removed. Hyper Sensitive corals were those which showed 100% necrosis after the heat-stress

experiment. Super Resistant corals showed 0% necrosis at the end of heat-stress. Microbial taxa

are ordered taxonomically. Only those taxa with a median relative abundance above 1% are

shown, the rest are listed as "other". IS is short for Incertae sedis, meaning there is uncertainty in

the taxonomic position within the broader taxonomic group. Relative abundance bubbles

correspond to within domain relative abundance. The high abundance of Bacteria IS can be

contributed to poor representation of octocoral bacteria in the SILVA reference database.

Supplementary Figure 4, Additional File 6. Maximum-likelihood tree of top alveolate ASVs

across the thermal tolerance categories of *P. clavata*. Heatmap abundance account for the

relative abundance of each specific ASV across the alveolates shown.

Supplementary Figure 5, Additional File 7. Maximum-likelihood tree of top dinoflagellate ASVs across the sampled populations of *P. clavata.* Populations are ordered from west to east, with major geographic localities color coded. ASV taxonomy is assigned to species level if sequences share above 98% similarity with their reference. ASV abundance is also shown across the whole data set (green bars) as well the ASV prevalence across all the samples (purple bars).

Supplementary Figure 6, Additional File 8. Maximum-likelihood tree of top ciliate ASVs

across the sampled populations of *P. clavata*. Populations are ordered from west to east, with

major geographic localities color coded [Medes = Green, Scandola = Red, Portofino = Yellow,

Croatia = Cyan]. ASV taxonomy is assigned to species level if sequences share above 98%

similarity with their reference. ASV abundance is also shown across the whole data set (green

bars) as well the ASV prevalence across all the samples (purple bars).