Apicomplexans predict thermal stress mortality in the Mediterranean coral

Paramuricea clavata

Anthony M Bonacolta[1,2], Jordi Miravall[3], Daniel Gómez-Gras[4,5], Jean-Baptiste Ledoux[6], Paula

López-Sendino[3], Joaquim Garrabou[3], Ramon Massana[3], Javier del Campo*[1,2]

[1]Department of Marine Biology and Ecology, Rosenstiel School of Marine, Atmospheric and Earth

Science, University of Miami, Miami, Florida, 33149 USA

[2]Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Barcelona, Catalonia, Spain

[3]Institut de Ciències del Mar – CSIC, Barcelona, Catalonia, Spain

[4]Hawaiʻi Institute of Marine Biology, University of Hawaiʻi at Mānoa, Kaneohe, HI, USA

[5]Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Institut de Recerca de la

Biodiversitat (IRBIO), Universitat de Barcelona, Barcelona, Spain

[6]CIIMAR/CIMAR, Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade

do Porto, Porto, Portugal.

*Corresponding Author: jdelcampo@ibe.upf-csic.es, Institut de Biologia Evolutiva (UPF-CSIC),

Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Catalonia, Spain

ABSTRACT

Paramuricea clavata is an azooxanthellate octocoral currently threatened by episodic mass

mortality events related to global warming. The microbiome may play an active role in the thermal

stress susceptibility of corals, potentially holding the answer as to why corals show differential

sensitivity to heat-stress. The microbiome of *P. clavata* collected from around the Mediterranean

was characterized right before experimental heat-stress using the 16S and 18S small subunits

(SSU) of the rRNA gene to determine if it influences the thermal response of the holobiont. *P.*

clavata's microeukaryotic community was significantly correlated with thermal stress sensitivity,

while bacterial assemblages were not. Syndiniales dinoflagellates Group I Clade 1 were

significantly enriched in thermally resistant corals, while the apicomplexan Corallicolida were

significantly enriched in thermally susceptible corals. Our results show the importance of the

microeukaryotic communities to understanding thermal stress response in corals and provides a

useful tool to guide conservation efforts.

Paramuricea clavata is a slow-growing octocoral prevalent in the temperate Mediterranean

Sea (1). Gorgonians, such as *P. clavata*, are habitat-forming species of biodiversity-rich hard

bottom benthic communities where they provide complex habitat for numerous organisms

contributing significantly to the biodiversity of the region (2). *P. clavata's* slow growth rate, long

turn-over time, high post-settlement mortality, and low genetic connectivity puts it at high-risk of

extinction caused by natural and anthropogenic stressors (1). *P. clavata* populations have been

facing increased mortality due to episodic mass mortality events (MMEs) (3). MMEs, which were

once uncommon in temperate ecosystems such as the Mediterranean Sea, have been increasing

since the turn of the century, with most of these occurring in the Western Mediterranean (4). The

increase in the frequency, extension, and intensity of marine heatwaves driven by global warming

have been suggested to be the primary cause of the increase in MMEs in the Mediterranean. Thus,

MMEs are leading to the current biodiversity crisis in the region (5,6). *P. clavata* are heavily

affected by MMEs with mortality sometimes reaching 80% in affected populations and recovery

hindered by the recurrence of marine heatwaves. The loss of this habitat-forming species causes a

functional shift in Mediterranean hard-bottom communities leading to simplified habitats with

lower species diversity (7,8). To prevent the significant decline of *P. clavata* and other habitat

forming octocorals in the Mediterranean, the effect of thermal stress on the *P. clavata* holobiont

must be better understood.

Thermal stress typically causes death in anthozoans by disrupting metabolic pathways,

increasing physiological stress, decreasing host defense, causing energy constraints, and

promoting the growth of opportunistic pathogens (9–11). In *P. clavata,* exposure to higher

temperatures causes usually partial or full necrosis of colonies, where branches first change color

from red to grey before denudation of the skeleton (12). This contrasts with "bleaching" which

takes place in zooxanthellate corals and is when the algal symbiont leaves the coral host during

periods of high heat-stress. *P. clavata* is more sensitive to elevated seawater temperatures than

other gorgonians, marked by a significant loss in energy reserves, reduced feeding ability, and

partial mortality after heat-stress (13). A complete characterization of the *P. clavata* microbiome

before heat-stress can better elucidate the role of the microbiome to overall holobiont health and

fitness in the face of warming ocean temperatures.

Octocoral-associated microorganisms are considered beneficial members of the coral

holobiont contributing to nitrogen fixation, sulfur-cycling, antibiotic production, and host defense

(14). *P. clavata* has a regionally stable bacteriome dominated (above ~90% relative abundance)

by the common coral symbiont *Endozoicomonas,* a Gammaproteobacteria of the order

Oceanospirillales (15,16). Notably, *P. clavata*'s bacterial community differs significantly from the

microbiome of the water surrounding the organism and that of neighboring octocorals, such as

Eunicella (17). Other bacteria commonly found within *P. clavata* include Alphaproteobacteria,

Betaproteobacteria, and Actinobacteria (15). Despite the research into the bacterial community

found in *P. clavata* throughout the Mediterranean Sea, there has been no research into the

microeukaryotic community which constitutes a significant portion of the microbial diversity and

can contribute significantly to holobiont fitness in a changing climate (18).

Microeukaryotes (protists and unicellular fungi) are important drivers of many of the

Earth's ecosystems including being the main contributors to oceanic primary production and

carbon mobilization in the microbial loop. They may influence the cycling of nutrients in the

holobiont as well as play a role in host defense. The corallicolids, which are part of a lineage of

parasitic alveolates known as the Apicomplexa are associated with 70% of corals globally (19).

Apicomplexans also include such well-known human parasites like *Plasmodium*, the causative

agent of malaria, and *Toxoplasma*, a wide-spread mammal-infecting pathogen. While the

corallicolids, based on their distribution and presence in healthy individuals (20), seem to be

commensals in the coral microbiome, their evolutionary context suggests a more parasitic role for

this general anthozoan associate.

To better understand the microbiome of *P. clavata* and its ability to predict thermal stress

susceptibility we conducted a common garden heat-stress experiment using corals collected from

12 populations (148 total individuals) across the geographic range of *P. clavata* (Extended Data

Fig. 1). We used an 18S rRNA and 16S rRNA gene metabarcoding approach to get a holistic view

of the microbes which may confer thermal resiliency to *P. clavata.*

RESULTS

The bacterial microbiome of *Paramuricea clavata*

For the 16S rRNA gene sequencing, 1 332 678 total reads were recovered after processing in DADA2. Following filtering, a total of 4 540 Amplicon Sequence Variants (ASVs) across 92 samples were left for bacterial community analysis. The water and food microbiome samples were taxonomically distinct from the coral microbiome samples in this study (Extended Data Fig. 2). The population originally sampled from Sagres, Portugal exhibited a distinct microbiome from the other *P. clavata* populations (Fig. 1) and was later determined to be a different species based on genomic work (21)**.** It was hence removed from further analysis. Oceanospirillales, mostly consisting of Endozoicomonas, constituted most of the bacterial community of *P. clavata* (Fig 1). Other highly abundant bacterial orders include Alteromonadales, Rhodobacterales, and Flavobacteriales. A high number of unidentified bacterial ASVs were also present in the dataset. Subsequent BLAST analysis of these ASVs yielded no definitive taxonomic identification.

Pair-wise analysis of the bacterial communities between localities showed significant differences in Shannon-Wiener alpha-diversity (Extended Data Fig. 3). Additionally, the beta-diversity of the bacterial community between localities was significantly different as well (ANOSIM; P = 0.001; Extended Data Fig. 4). Overall, the bacterial community of *P. clavata* clustered closely based on geographic origin.

The dominance of Endozoicomonas (Proteobacteria; Oceanospirillales) in the bacterial community of *P. clavata* masked any fine-scale differences in the bacterial community composition across the range of thermal susceptibilities (Fig. 2). Analyzing the bacterial community with Endozoicomonas removed, did not reveal any noticeable pattern in taxon composition in relation to thermal stress susceptibility (Extended Data Fig. 5); however, an

ANCOMBC comparison between the super-resistant and hyper-sensitive thermal susceptibility

categories showed that Pseudomonadales and Francisellales were significantly more abundant in

hyper-sensitive corals (Fig. 3).

The eukaryotic microbiome of *Paramuricea clavata*

For the 18S rRNA sequencing, 18 955 358 total reads were recovered, resulting in 3 181

ASVs across 138 samples following filtering. This community is dominated by 3 major clades of

Alveolates including: Apicomplexans, Ciliates, and Syndiniales (Fig. 1). One of the primary

microeukaryotic taxa present, Dino-Group-I (Dinoflagellata; Syndiniales), exhibited its highest

relative abundance in the more Eastern populations (Balun, Mana, & Lighthouse) (Fig. 1).

Corallicolids (Apicomplexa; Corallicolida) exhibited a high relative abundance across every

population sampled, excluding Lighthouse (Fig. 1). Additionally, three distinct genera of ciliates

(Ciliophora) were found across the surveyed geographic range of *P. clavata*, with *Licnophoria*

spp. being prevalent at Tascons, Lighthouse, and Altare (Fig. 1). *Hypotrichia* spp. were found

mainly in the Western populations (Tascons, Pota de Llop, La Vaca, Palazzu, Palazzinu, &

Gargallu) (Fig. 1). Lastly, *Suctoria* spp. were most prevalent within the *P. clavata* of the Medes

and Portofino localities (Fig. 1).

At the ASV-level, the locality-level microeukaryotic communities of *P. clavata* exhibited

higher Shannon-Wiener alpha-diversity compared to the bacterial communities (Extended Data

Fig. 3). Additionally, the microeukaryotic community only showed a significant difference in the

alpha-diversity of *P. clavata* between Croatia vs Medes. Furthermore, the beta-diversity between

localities was significantly different for the microeukaryotic community (ANOSIM; $P = 0.001$;

Extended Data Fig. 4).

Syndiniales were enriched in the super-resistant corals (LEfSe; P = 0.0016). Specifically,

Dino-Group I Clade 1 abundance in the holobiont was associated with the increased

thermotolerance in *P. clavata* and was enriched in the super-resistant individuals (LEfSe; P =

0.0027; Fig. 2). At the ASV level, ASV_20 and ASV_13 were the Dino-Group I Clade 1 ASVs

which were highly abundant in the resistant groups relative to the other Alveolates (Extended Data

Fig. 8). Concurrently, a high relative abundance of corallicolids within the holobiont was

associated with higher sensitivity to thermal stress (Fig. 2). LEfSe analysis revealed that the

Corallicolidae family was significantly enriched in the sensitive corals (LEfSe; P = 0.015). ASV_6,

within the COR3 clades of corallicolids, showed the highest abundance amongst of the alveolates

in the hyper-sensitive corals (Fig. 9). ANCOMBC analysis also confirmed these associations, with

corallicolids being significantly more abundant (averaging ~2.5 natural log fold higher abundance)

in the sensitive corals compared to the resistant ones, while Dino-Groups I and II showed an

opposite pattern (Fig. 3). On the other hand, ciliates showed a stable abundance throughout thermal

sensitivity groups, except for ASV_7 (*Ephelota mammillata*) which was highly abundant in the

sensitive and tolerant groupings (Extended Data Fig. 8). Alpha-diversity and beta-diversity

analysis of both the bacterial community and microeukaryotic community yielded no significance

differences between any of the thermal tolerance categories (Extended Data Figs. 3 & 4).

Phylogenetic placement of the microeukaryotic microbiome

Phylogenetic analysis of the most prevalent Alveolate ASVs found within *P. clavata*

revealed a diversity of clades associated with the Octocoral across the surveyed geographic range.

ASV_6, which was the most abundant corallicolid ASV and found in most samples, was most

abundant in the Scandola populations (Fig. 4). An evolutionary placement algorithm placed this

ASV as closely related to MN022452_Corallicolidae_COR3_sp. (99.22% Similarity), a sequence

recovered from the deep-sea Octocoral, *Callogorgia delta* in the Gulf of Mexico. A second clade

was identified that most likely belonged to the Corallicolidae COR2 group (Fig. 4). These ASVs

(ASV_16, ASV_19, ASV_22) have a low relative abundance within *P. clavata* but are found

across most samples (Fig. 4). This group is most closely related to

MN022429_Corallicolidae_COR2_sp. (95.84% Similarity to ASV_16), a sequence recovered

from the deep-sea Antipatharian *Leiopathes glaberrima*, also from the Gulf of Mexico. Lastly, a

third clade of prevalent corallicolids was identified within *P. clavata* that shows close relation to

JX943883_Corallicolidae_COR1_sp. (93.04% Similarity to ASV_105), a sequence found in the

stony coral *Porites astreoides,* from the Bahamas.

Examination of the Dinoflagellate ASVs revealed distinct clades present within *P. clavata*.

Dino-Group-I Clade 1 ASVs were most prevalent across the dataset with ASV_13 being the most

abundant across all localities except Scandola, where ASV_12 is most prevalent (Extended Data

Fig. 6). The closest blast results for these sequences were to uncultured microeukaryotes isolated

from intertidal mudflat sediments and ocean surface water from the China Sea (22,23) (100%

Similarity). One Symbiodiniaceae ASV (ASV_43) was identified in some of the populations

(Tascons, La Vaca, Palazzinu, Lighthouse, Altare, Mana & Balun) despite *P. clavata* being

azooxanthellate (Extended Data Fig. 6). The ASV from this study corresponds to Symbiodinium

Temperate Clade A which has been identified in Mediterranean corals before (24). It's closest blast

result from NCBI was to a *Symbiodinium sp.* Temperate Clade A sequence (99.21% Similarity)

recovered from the stony coral *Mussismilia hispida* (25).

The ciliate ASVs within the dataset revealed three primary clades found variably abundant

across the surveyed distribution of *P. clavata*. ASVs corresponding to *Holostichia diademata*

(Order: Hypotrichia) (ASV_8) and *Ephelota mammillata* (Order: Suctoria) (ASV_7) were the most

abundant and prevalent across the dataset overall (Extended Data Fig. 7). Interestingly, BLAST

results for ASV_8, shows 100% similarity to *Holosticha pullaster* (Order: Hypotrichia) (100%

Similarity) isolated from a freshwater pond in the USA (26). ASV_7 showed high similarity to *E.*

mammillata recovered from *Tubularia* hydroids (98.51% Similarity) (27). *Licnophora*

macfarlandi (Order: Licnophoria), was found across most populations, with its highest abundance

being in Mana, Croatia (Extended Data Fig. 7). Blast results show that these sequences were highly

similar to *L. macfarlandi* (92.66% Similarity to ASV_3) recovered from the fluid in the respiratory

trees of the giant California sea cucumber *Parastichopus californicus* (28).

Co-occurrence between members of the microbiome

A microbial co-occurrence network was constructed using SPARCC to analyze the positive

correlations between the major microbial families of the *P. clavata* holobiont. Corallicolidae &

Dino-Group I Clade 1 had the strongest significant co-association between all the major taxa (Fig.

5). This co-association is of note when you consider these two group's antagonistic relative

abundance patterns across the thermal susceptibility groups (Fig. 2). The three ciliate families

showed consistent associations with the corallicolids and Dino-Group I Clade 1. Endozoicomonas,

Pseudoalteromonadaceae, Vibrionaceaee, and Flavobacteriaceae were the primary bacterial

families that showed significant co-occurrence with the different microeukaryotic members of the

P. clavata holobiont (Fig. 5).

DISCUSSION

The bacterial community of *P. clavata* showed very little differences in its taxonomic

make-up from location to location (Fig. 1), which is consistent with previous work into the

bacterial component of the *P. clavata* holobiont (16,29). The differences in beta-diversity and

alpha-diversity between the different localities where *P. clavata* was sampled can be contributed

to the vast diversity of Endozoicomonas found across the surveyed distribution. The prevalence of

unidentified bacterial ASVs in the dataset points to the underrepresentation of octocoral-associated

bacteria in the SILVA reference database. These results exemplify the need to identify more

bacteria associated with a healthy symbiotic state in coral holobionts.

The microeukaryotic community of *P. clavata* is dominated by three major groups of

alveolates: Apicomplexa (mostly corallicolids), Dinoflagellata (mostly Syndiniales), and

Ciliophora. This community shows more variability in its taxonomic make-up from location to

location, while also showing significant differences in beta-diversity (Fig. 1 & Extended Data Fig.

4). Each of the alveolates found in *P. clavata* showed locality-specific differences, highlighting

the specificity and potential co-adaptations of these associations in coral holobionts (Fig. 4,

Extended Data Figs. 6 & 7). Corallicolids are general anthozoan associates found world-wide,

even at depths as deep as 1400 m, and in some cases can be as abundant as Symbiodiniaceae

(30,31). They tend to be the most abundant and prevalent microeukaryotic coral symbiont in

azooxanthellate corals. The ones recovered from this study showed close relation to those found

within deep-sea cnidarians, including octocorals and antipatharians, of the Gulf of Mexico from

Vohsen *et al.* (32) (Fig. 4). A third clade of corallicolids were closely related to one recovered

from the shallow water Caribbean coral *Porites astreoides* (33). This diversity of corallicolids

present within *P. clavata,* alone, hints at the potential global diversity and ubiquity of corallicolids

in anthozoan hosts.

Syndiniales, which have been found in high prevalence in reef environments before, also

constituted a major portion of the eukaryotic microbiome. They are an order of parasites found at

the base of the dinoflagellates which can infect a variety of marine macro- and microscopic

organisms (34). Syndiniales from Dino-Group I are one of the main microeukaryotic taxa found

in the hard coral *Pocillopora damicornis* (35), however this group is also ubiquitous in global

plankton communities (36). Additionally, Dino-Group 1 Syndiniales with identical 18S rRNA V4

sequences may associate with different hosts, suggesting a generalist host specificity for this clade

or the need for a different marker gene to decipher its true diversity (37). Syndiniales have the

potential to disrupt nutrient cycling, and thus would likely play a role in (de)stabilizing symbiotic

relationships, which rely on nutrient dynamics, in a tight-knit symbiotic system such as a coral

holobiont. Their differential abundance within *P. clavata* across different thermal susceptibility

categories highlights the necessity of studying the role of parasitism in the nutritional dynamics of

marine holobionts, especially in the face of stressors. Hudson et al. 2006 noted that a healthy

system may be one that is rich in parasitic species due to the impact of parasite-mediated effects

on population dynamics, interspecific competition, and energy flow (38). Applying this

perspective to coral holobionts may help us better understand the higher thermal tolerance of *P.*

clavata rich in Syndiniales. The high abundance of these parasites within the *P. clavata* holobiont

may contribute to a healthy and resilient system by influencing the microbial community dynamics

when confronted with heat-stress.

Within the ciliate lineages recovered from *P. clavata*, three genera (*Licnophora*,

Holosticha, & *Ephelota*) showed variable abundances across the surveyed distribution of the coral

host. *L. macfarlandi*, has been found as an endosymbiont in the respiratory trees of holothuroids,

but also in association with corals, likely as a secondary invader hunting other microbes (39,40).

It is a noted predator of protozoans and is speculated to be attracted to other ciliate species found

at the interface of coral lesions (40). *L. macfarlandi* within *P. damicornis* also exhibited geographic

variability in abundance (35), which may depend upon the presence of other ciliates on which it

preys. Another ciliate, *H. diademata,* has also been found within corals often crawling slowly on

the coral surface (40). It is a noted bacterivore so it may feed upon the bacteria of the coral, exerting

top-down control upon that portion of the microbiome, and therefore influencing the cycling of

certain nutrients in the holobiont. The presence of bacterivores may increase microbial activity

and turnover within the ecosystem (41). While the previous two ciliates are predominately

associated with diseased corals (42), the third ciliate genera, *Ephelota,* has never previously been

described in corals, however Suctorian ciliates are typically commensal and have previously been

recovered as epibionts of crustaceans, mollusks, fishes, and even other protists (43). Thus, it may

be using the stability and structure provided by *P. clavata* to stabilize itself in the water column

and capture prey more readily.

This study identified the key microbes associated with *P. clavata,* an important habitat-

forming species impacted by MMEs, and provides evidence of the microbiome's potential role in

the thermal susceptibility of the host. One of the bacterial taxa found to be associated with

thermally resilient coral was Pseudomonadales, which may be beneficial to a healthy coral

holobiont as it can prevent the growth of harmful Flavobacteria which may lead to disease (44).

However, in *P. clavata*, the overwhelming abundance of Endozoicomonas didn't allow for any

other fine scale differences in bacterial composition to be detected. On the other hand, the

microeukaryotic community has distinct taxonomic composition associated with different degrees

of thermal tolerance. Dino-Group I Clade 1 was significantly enriched in thermally tolerant corals,

while corallicolids were significantly enriched in thermally sensitive corals. How the relative

abundance of these apicomplexans confers thermal susceptibility to the coral host is not yet

understood. There's the possibility that the symbiosis between corallicolids and their coral host

shifts from an apparent commensalism towards parasitism as added stressors, such as heat,

breakdown the relationship. The context-dependency of this relationship can explain the

abundance of corallicolids in apparently healthy coral. The coral host may have evolved resistance

to corallicolids at a significant fitness cost leading to a state of commensalism without the presence

of addition stressors; however, heat-stress may cause the corallicolid-tolerant corals to exhibit

higher mortality as the apicomplexans shift towards parasitism, similar to the hypothesis proposed

by Miller et al. 2006 (45). This potential interaction and the widespread distribution of the

corallicolids would have significant implications when interpreting threats to corals in the context

of warming ocean temperatures.

Showing strong co-association with the corallicolids were the Syndiniales, specifically

those belonging to Dino-Group I Clade 1 (Fig. 5). These parasites are known to infect a variety of

multicellular and microbial hosts and have previously been recovered from coral reef sediments,

plankton, and corals (46). Members include microeukaryotic parasites such as *Duboscquella sp.*

which is known to parasitize ciliates (47). This opens the possibility that they may be infecting

either the coral host, it's microeukaryotic members, or both. While significant positive co-

association between Dino-Group I and the ciliates recovered from *P. clavata* was apparent in the

network analysis, the association between Dino-Group I and the corallicolids was most prevalent

(Fig. 5). While this can simply be the result of fluctuations in two of the most highly abundant

microeukaryotes found in the dataset, it may also represent an interesting case of parasitism of a

coral endosymbiont by another member of the microeukaryotic community. Its association with

the corallicolids hints at a possible positive role of this parasite preventing the overabundance of

corallicolid growth in the coral holobiont which may be associated with decreased thermal

tolerance. Syndiniales have been proposed to have the capacity to control the sizes of their host

populations since a singular infection can produce hundreds of dinospores to further infect new

hosts (48). The vast diversity of known microeukaryotic hosts of syndiniales combined with the

larger cell size of corallicolids (~12 um in length (19) supports the possibility of this relationship.

Possible explanations for this interaction could be top-down control of the corallicolid population

by the Syndiniales, niche competition, easier co-infection of an already infected host, or excess

dinospore production by Syndiniales. In either case, the relationship between Dino-Group I, the

corallicolids and the coral host needs to be investigated further to decipher the ecological relevance

of this potentially pivotal interaction within the coral holobiont.

We have shown here that a deeper characterization of the often-overlooked

microeukaryotic portion of the coral microbiome can have major implications of coral health in a

warming ocean. Along with other genetic, environmental, and physiological mechanisms the

microbiome should be considered a vital component when assessing coral thermal response. Future

research should focus on identify the mechanisms by which these microbes confer thermal

resiliency to individual corals and how this knowledge can best be used to manage fragile coral

ecosystems.

Conflict of Interest Statement

The authors declare no conflicts of interest.

Acknowledgments

We appreciate the contributions of the following scientists in helping with our sampling effort:

Bensoussan, N, Cerrano C, Ferretti E, Kipson S, Bakran-Petricioli T, Serrao EA, Paulo D, Boavida

J, Montero-Serra I, López-Sanz A, Milanese M, Linares C. This research was supported by the

strategic funding UIDB/04423/2020, UIDP/04423/2020 and 2021.00855.CEECIND through

national funds provided by FCT – Fundação para a Ciência e a Tecnologia. Joaquim Garrabou and

Paula López-Sendino, Daniel Gómez Gras acknowledges the funding by the "Severo Ochoa Centre

of Excellence" (CEX2019-000928-S), Interreg-Med Programme MPA-Engage

(1MED15_3.2_M2_337), the European Union Horizon 2020 research and innovation programme

(Futuremares SEP-210597628) H2020 MERCES project.

ONLINE METHODS

Sample Collection

Branch fragments (30 colonies per population) of healthy adult *P. clavata* were collected

from four localities (11 discrete populations) encompassing much of the known Mediterranean

distribution of the octocoral. 10-17 individuals per population were included in microbiome

analyses, depending on their availability. From Croatia, *P. clavata* populations from Balun (Lat,

Long; 43.803889, 15.255000) and Mana (43.800273, 15.266389) were sampled. Three

populations were sampled from Portofino (Liguria, Italy): Altare (44.308918, 9.178645), Indiano

(44.312476, 9.166889), and Lighthouse (44.298650, 9.219014). Three populations were sampled

from Scandola (Corsica, France): Palazzinu (42.379854, 8.549999), Gargallu (42.371839,

8.534394), and Palazzu (42.379873, 8.546163). Three populations were sampled from Medes

Islands (Catalonia, Spain): La Vaca (42.048047, 3.226322), Pota de Llop (42.0497   ,

3.2254), and Tascons (42.042189, 3.2269). Lastly, a population thought to be *P. clavata* was

sampled from an Atlantic locality from Sagres, Portugal (37.012129, -8.924541).

Thermal Stress Experiment

All branch fragments were transported to the Institut de Ciències del Mar (CSIC) in

Barcelona, Catalonia, Spain where they were housed in aquaria containing locally sourced

seawater. Colony fragments were distributed in different tanks, according to their sampling sites

and the two temperature treatments. Thermal treatments consisted of one control at which water

temperature was maintained at 18ºC, and another heated treatment at which water temperature was

maintained at 25ºC. The gorgonians were fed with copepods (*Cyclops* spp.) until the end of the

experiment. Following 21 days of incubation, the response of the coral fragment was assessed by

measuring the percentage of tissue necrosis. Thermal response was categorized as follows: Super-

resistant (0% necrosis), Resistant (< 25% necrosis), Tolerant (25-75% necrosis), Sensitive (>75%

necrosis), and Hyper-sensitive (100% necrosis). Extended Data Table 1 shows the thermal

resiliency categories for each coral sample at the end of the experiment. For each sample, 0.5 cm

length fragments were collected into a 2 mL cryovial containing 96% EtOH and stored at -80ºC.

DNA Extraction and rRNA Gene Metabarcoding

P. clavata samples were thawed at room temperature and then processed using a modified

DNeasy Mini Kit (QIAGEN) protocol. DNA concentration from purified samples was measured

using Qubit fluorometer. Concentrations that were higher than 1 ng/µl were considered acceptable

for PCR. The V4 region of the prokaryotic 16S rRNA gene was amplified according to the Earth

Microbiome Project protocol (49) and the V4 region of the eukaryotic 18S rRNA gene was

amplified using a nested PCR approach validated in del Campo et al. 2019 (50). Briefly, for the

18S rRNA metabarcoding, the samples were first amplified with UNonMet-PCR primers which

reduce the metazoan signal (18S-EUK581-F 5'-GTGCCAGCAGCCGCG-3′, 18S-EUK1134-R

5'-TTTAAGTTTCAGCCTTGCG-3′), then reamplified with primers from Comeau et al. 2011

(51). The PCR products were cleaned using QIAquick PCR Purification Kit (Qiagen). DNA

concentrations from purified PCR samples were checked using Qubit fluorometer before being

sent to the Integrated Microbiome Resource facility at the Centre for Comparative Genomics and

Evolutionary Bioinformatics at Dalhousie University for further 18S rRNA amplification and

sequencing.

Microbiome Analysis

Raw sequencing data for this project has been deposited on NCBI (XXXXXX). Primers

were removed from reads using Cutadapt v3.1 (52). The trimmed reads were then processed in R

using DADA2 (53). For the 18S rRNA gene amplicons, forward reads were truncated at 230 bp

and reverse reads were truncated at 195 bp. Reads were then denoised and joined into amplicon

sequence variants (ASVs). Chimeras were removed using the 'removeBimeraDenovo' command

with the "consensus" option. Taxonomy was assigned to ASVs using PR2 v4.13.0 (54). For the

16S rRNA gene amplicons, forward reads were truncated at 250 bp and reverse reads at 210 bp.

Truncated reads were then denoised and joined into ASVs before assigning taxonomy using

SILVA v138 (55). Sequence tables, taxonomy tables, and metadata for both the 16S and 18S

amplicon datasets were uploaded into a Phyloseq object in R for further filtering and analysis (56).

For the 16S rRNA gene amplicons, ASVs corresponding to Chloroplast and Mitochondria were

removed. One sample was also removed for having under 250 reads. Additionally, in each sample

ASVs that represent 1% or less of the reads were discarded. Following this filtering 4 546 ASVs

and 92 samples remained. The 18S amplicon were subjected to the same filtering parameters

except ASVs corresponding to Metazoa and Embryophyceae were also removed from the dataset,

resulting in 3 181 ASVs across 138 samples. Bubble plots, alpha-diversity, and beta-diversity

figures were constructed using ggplot2 and tidyverse packages (57). Statistical analyses were

conducted using the microbiomeMarker package and vegan (58,59). Analysis of similarities

(ANOSIM) within vegan was used to test for differences in beta-diversity between groups using

999 permutations. Linear Discriminant Analysis Effect Size (LEfSe) and Analysis of

Compositions of Microbiomes with Bias Correction (ANCOMBC) were used to find significantly

enriched and differentially abundant taxa between treatment groups.

In order to place the corallicolid ASVs in a phylogenetic context, a multiple sequence

alignment (MSA) was created using publicly available full-length 18S rRNA sequences of

corallicolids and the program MAFFT. Abundant corallicolid ASVs (more than 50% of

corallicolid signal in at least one sample) were then extracted from the dataset. Hmmalign was

then used to map these ASVs to the full length 18S rRNA corallicolid MSA (60). A maximum

likelihood phylogenentic tree was constructed with RAxML using this MSA (GTRCAT

substitution model, N = 1 000) (61). The resulting tree was modified for publication using ggtree

(62). For the dinoflagellate and ciliate trees, a similar approach was taken, except no publicly

available sequences were used in their construction and alignment.

To assess cross-domain positive-correlation in the occurrence patterns of ASVs, the 16S

and 18S amplicon tables were combined for samples present in both datasets. After setting a filter

prevalence threshold of 0.001, correlation patterns were inferred for the combined dataset using

SPARCC which considers the compositional nature of metabarcoding data (63). A network was

then constructed using SpiecEasi with a p-value threshold of 0.05 (ref. 30). The top families

exhibiting positive co-occurrence relationships were plotted in a chordplot using the MicroEco

package (65). All bioinformatic scripts used in this study can be found at

https://github.com/Abonacolta/Bonacolta_2022_Pcla.

References:

1. Linares C, Doak DF, Coma R, Díaz D, Zabala M. Life history and viability of a long-lived marine invertebrate: The octocoral Paramuricea clavata. Ecology. 2007;88(4):918–28.

2. Ballesteros E. Mediterranean coralligenous assemblages: A synthesis of present knowledge. Oceanogr Mar Biol. 2006;44:123–95.

3. Linares C, Doak DF. Forecasting the combined effects of disparate disturbances on the persistence of long-lived gorgonians: A case study of Paramuricea clavata. Mar Ecol Prog Ser. 2010;402:59–68.

4. Garrabou J, Gómez-Gras D, Ledoux JB, Linares C, Bensoussan N, López-Sendino P, et al. Collaborative Database to Track Mass Mortality Events in the Mediterranean Sea. Front Mar Sci. 2019;6(November).

5. Garrabou J, Gómez-Gras D, Medrano A, Cerrano C, Ponti M, Schlegel R, et al. Marine heatwaves drive recurrent mass mortalities in the Mediterranean Sea. Glob Chang Biol. 2022;28(19):5708–25.

6. Garrabou J, Ledoux J-B, Bensoussan N, Gómez-Gras D, Linares C. Sliding Toward the of Mediterranean Coastal Marine Rocky Ecosystems. 2021. 291–324 p.

7. Verdura J, Linares C, Ballesteros E, Coma R, Uriz MJ, Bensoussan N, et al. Biodiversity loss in a Mediterranean ecosystem due to an extreme warming event unveils the role of an engineering gorgonian species. Sci Rep. 2019;9(1):1–11.

8. Gómez-Gras D, Linares C, Dornelas M, Madin JS, Brambilla V, Ledoux JB, et al. Climate change transforms the functional identity of Mediterranean coralligenous assemblages. Ecol Lett. 2021;24(5):1038–51.

9. van de Water JAJM, Chaib De Mares M, Dixon GB, Raina JB, Willis BL, Bourne DG, et

al. Antimicrobial and stress responses to increased temperature and bacterial pathogen

challenge in the holobiont of a reef-building coral. Mol Ecol. 2018;27(4):1065–80.

10.   Hou J, Xu T, Su D, Wu Y, Cheng L, Wang J, et al. RNA-seq reveals extensive

transcriptional response to heat stress in the stony coral Galaxea fascicularis. Front Genet.

2018;9(FEB):1–10.

11.   Li J, Long L, Zou Y, Zhang S. Microbial community and transcriptional responses to

increased temperatures in coral Pocillopora damicornis holobiont. Environ Microbiol.

2021;23(2):826–43.

12.   Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P, Cigliano M, et al. Mass

mortality in Northwestern Mediterranean rocky benthic communities: Effects of the 2003

heat wave. Glob Chang Biol. 2009;15(5):1090–103.

13.   Tignat-Perrier R, van de Water JAJM, Guillemain D, Aurelle D, Allemand D, Ferrier-

Pagès C. The Effect of Thermal Stress on the Physiology and Bacterial Communities of

Two Key Mediterranean Gorgonians. Appl Environ Microbiol. 2022;88(6).

14.   Van De Water JAJM, Allemand D, Ferrier-Pagès C. Host-microbe interactions in

octocoral holobionts - recent advances and perspectives. Microbiome. 2018;6(1):1–28.

15.   Vezzulli L, Pezzati E, Huete-Stauffer C, Pruzzo C, Cerrano C. 16SrDNA Pyrosequencing

of the Mediterranean Gorgonian Paramuricea clavata Reveals a Link among Alterations in

Bacterial Holobiont Members, Anthropogenic Influence and Disease Outbreaks. PLoS

One. 2013;8(6):1–10.

16.   La Rivière M, Garrabou J, Bally M. Spatial and temporal analysis of bacterial diversity

associated with the mediterranean gorgonian Paramuricea clavata. Rapp Comm int Mer

medit. 2010;13007.

17.   van de Water JAJM, Melkonian R, Voolstra CR, Junca H, Beraud E, Allemand D, et al.

Comparative assessment of Mediterranean gorgonian-associated microbial communities

reveals conserved core and locally variant bacteria. Microb Ecol. 2017;73(2):466–78.

18.   Ainsworth TD, Fordyce AJ, Camp EF. The Other Microeukaryotes of the Coral Reef

Microbiome. Vol. 25, Trends in Microbiology. 2017. p. 980–91.

19.   Kwong WK, Irwin NATT, Mathur V, Na I, Okamoto N, Vermeij MJAA, et al. Taxonomy

of the Apicomplexan Symbionts of Coral, including Corallicolida ord. nov., Reassignment

of the Genus Gemmocystis, and Description of New Species Corallicola aquarius gen.

nov. sp. nov. and Anthozoaphila gnarlus gen. nov. sp. nov. J Eukaryot Microbiol.

2021;68(4):1–9.

20.   Kwong WK, Mathur V, Vermeij MJA, Keeling PJ. A widespread coral-infecting

apicomplexan contains a plastid encoding chlorophyll biosynthesis. 2018;

21.   Coelho M, Pearson G, Boavida J, Paulo D, Aurelle D, Daniel G, et al. Not out of the

Mediterranean : Atlantic populations of the gorgonian Paramuricea clavata are a separate

sister species under further lineage diversification. Mol Ecol. 2023;1–18.

22.   Wu W, Huang B, Liao Y, Sun P. Picoeukaryotic diversity and distribution in the

subtropical-tropical South China Sea. FEMS Microbiol Ecol. 2014;89(3):563–79.

23.   Ma Y, Ouyang J, Wei N, Zhang S, Yan X, Xiong H. Microeukaryotic community

structure and dynamics during macrobenthos bioremediation in intertidal mudflat

sediments of Sansha Bay, China. Jundishapur J Microbiol. 2017;10(8).

24.   Visram S, Wiedenmann J, Douglas AE. Molecular diversity of symbiotic algae of the

genus Symbiodinium (Zooxanthellae) in cnidarians of the Mediterranean Sea. J Mar Biol

Assoc United Kingdom. 2006;86(6):1281–3.

25.   Picciani N, de Lossio e Seiblitz IG, de Paiva PC, e Castro CB, Zilberberg C. Geographic

487   patterns of Symbiodinium diversity associated with the coral Mussismilia hispida

488   (Cnidaria, Scleractinia) correlate with major reef regions in the Southwestern Atlantic

489   Ocean. Mar Biol. 2016;163(11):1–11.

490   26.   Chen L, Dong J, Wu W, Xin Y, Warren A, Ning Y, et al. Morphology and molecular

491   phylogeny of a new hypotrich ciliate, Anteholosticha songi nov. spec., and an American

492   population of Holosticha pullaster (Müller, 1773) Foissner et al., 1991 (Ciliophora,

493   Hypotrichia). Eur J Protistol. 2020;72:125646.

494   27.   Riley JL, Katz LA. Widespread distribution of extensive chromosomal fragmentation in

495   ciliates. Mol Biol Evol. 2001;18(7):1372–7.

496   28.   Lynn DH, Strüder-Kypke M. Phylogenetic Position of Licnophora, Lechriopyla, and

497   Schizocaryum, Three Unusual Ciliates (Phylum Ciliophora) Endosymbiotic in

498   Echinoderms (Phylum Echinodermata). J Euk. 2002;49(6):460–8.

499   29.   La Rivière M, Roumagnac M, Garrabou J, Bally M. Transient Shifts in Bacterial

500   Communities Associated with the Temperate Gorgonian Paramuricea clavata in the

501   Northwestern Mediterranean Sea. PLoS One. 2013;8(2).

502   30.   Kwong WK, del Campo J, Mathur V, Vermeij MJA, Keeling PJ, del Campo J, et al. A

503   widespread coral-infecting apicomplexan with chlorophyll biosynthesis genes. Nature.

504   2019 Apr 3;568(7750):103–7.

505   31.   Vohsen SA, Anderson KE, Gade AM, Gruber-Vodicka HR, Dannenberg RP, Osman EO,

506   et al. Deep-sea corals provide new insight into the ecology, evolution, and the role of

507   plastids in widespread apicomplexan symbionts of anthozoans. Microbiome. 2020;8(1):1–

508   15.

32.   Vohsen SA, Anderson KE, Gade AM, Gruber-Vodicka HR, Dannenberg RP, Osman EO,

et al. Deep-sea corals provide new insight into the ecology, evolution, and the role of

plastids in widespread apicomplexan symbionts of anthozoans. Microbiome. 2020;8(1):1–

15.

33.   Kirk NL, Thornhill DJ, Kemp DW, Fitt WK, Santos SR. Ubiquitous associations and a

peak fall prevalence between apicomplexan symbionts and reef corals in Florida and the

Bahamas. Coral Reefs. 2013;32(3):847–58.

34.   Guillou L, Viprey M, Chambouvet A, Welsh RM, Kirkham AR, Massana R, et al.

Widespread occurrence and genetic diversity of marine parasitoids belonging to

Syndiniales (Alveolata). Environ Microbiol. 2008;10(12):3349–65.

35.   Clerissi C, Brunet S, Vidal-Dupiol J, Adjeroud M, Lepage P, Guillou L, et al. Protists

within corals: the hidden diversity. Front Microbiol. 2018 Aug 31 [cited 2021 Sep

29];9(August):2043.

36.   Engelen S, Hingamp P, Sieracki M, Vargas C, Audic S, Henry N, et al. Eukaryotic

plankton diversity in the sunlit ocean. Science (80-). 2015;348(6237):1261605-1/11.

37.   Clarke LJ, Bestley S, Bissett A, Deagle BE. A globally distributed Syndiniales parasite

dominates the Southern Ocean micro-eukaryote community near the sea-ice edge. ISME

Journalp. 2019;13(3):734–7.

38.   Hudson PJ, Dobson AP, Lafferty KD. Is a healthy ecosystem one that is rich in parasites?

Trends Ecol Evol. 2006;21(7):381–5.

39.   Lynn DH, Str M. Scuticociliate Endosymbionts of Echinoids (Phylum Echinodermata):

Phylogenetic Relationships among Species in the Genera Entodiscus , Plagiopyliella ,

Thyrophylax , and Entorhipidium (Phylum Ciliophora) Author (s): Denis H . Lynn and

Michaela Strüde. 2022;91(5):1190–9.

40.   Sweet MJ, Séré MG. Ciliate communities consistently associated with coral diseases. J

Sea Res. 2016;113:119–31.

41.   Trap J, Bonkowski M, Plassard C, Villenave C, Blanchart E. Ecological importance of soil

bacterivores for ecosystem functions. Plant Soil. 2016;398(1–2):1–24.

42.   Ravindran C, Raveendran HP, Irudayarajan L. Ciliated protozoan occurrence and

association in the pathogenesis of coral disease. Microb Pathog. 2022;162(April

2021):105211.

43.   Dovgal I V. Evolution, phylogeny and classification of Suctorea (Ciliophora).

Protistology. 2002;2(4).

44.   Gignoux-Wolfsohn SA, Vollmer S V. Identification of Candidate Coral Pathogens on

White Band Disease-Infected Staghorn Coral. PLoS One. 2015 Aug 4 [cited 2022 May

26];10(8).

45.   Miller MR, White A, Boots M. The evolution of parasites in response to tolerance in their

hosts: the good, the bad, and apparent commensalism. Evolution (N Y). 2006 May 1 [cited

2022 May 26];60(5):945–56.

46.   Clerissi C, Brunet S, Vidal-Dupiol J, Adjeroud M, Lepage P, Guillou L, et al. Protists

within corals: the hidden diversity. Front Microbiol. 2018;9(August):2043.

47.   Harada A, Ohtsuka S, Horiguchi T. Species of the Parasitic Genus Duboscquella are

Members of the Enigmatic Marine Alveolate Group I. Protist. 2007 Jul 18;158(3):337–47.

48.   Jephcott TG, Alves-de-Souza C, Gleason FH, van Ogtrop FF, Sime-Ngando T, Karpov

SA, et al. Ecological impacts of parasitic chytrids, syndiniales and perkinsids on

populations of marine photosynthetic dinoflagellates. Fungal Ecol. 2016 Feb 1;19:47–58.

49.   Ul-Hasan S, Bowers RM, Figueroa-Montiel A, Licea-Navarro AF, Michael Beman J,

556   Woyke T, et al. Community ecology across bacteria, archaea and microbial eukaryotes in

557   the sediment and seawater of coastal Puerto Nuevo, Baja California. PLoS One.

558   2019;14(2):1–7.

559   50.   Campo J del, Pons MJ, Herranz M, Wakeman KC, Valle J del, Vermeij MJAA, et al.

560   Validation of a universal set of primers to study animal-associated microeukaryotic

561   communities. Environ Microbiol. 2019 Oct 1 [cited 2021 Sep 29];21(10):3855–61.

562   51.   Comeau AM, Li WKW, Tremblay JÉ, Carmack EC, Lovejoy C. Arctic ocean microbial

563   community structure before and after the 2007 record sea ice minimum. PLoS One.

564   2011;6(11).

565   52.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

566   EMBnet.journal. 2011 May 2 [cited 2020 Aug 14];17(1):10.

567   53.   Callahan BJ, Mcmurdie PJ, Rosen MJ, Han AW, A AJ. DADA2: High resolution sample

568   inference from Illumina amplicon data. Nat Methods. 2016;13(7):581–3.

569   54.   Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, et al. The Protist Ribosomal

570   Reference database (PR2): a catalog of unicellular eukaryote small sub-unit rRNA

571   sequences with curated taxonomy. Nucleic Acids Res. 2013 Jan 26 [cited 2014 May

572   1];41(Database issue):D597-604.

573   55.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal

574   RNA gene database project: Improved data processing and web-based tools. Nucleic

575   Acids Res. 2013;41(D1):590–6.

576   56.   McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis

577   and Graphics of Microbiome Census Data. PLoS One. 2013;8(4).

57.   Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to

the Tidyverse. J Open Source Softw. 2019;4(43):1686.

58.   Cao Y. Tools for microbiome marker identification • microbiomeMarker. [cited 2022 Apr

25]. Available from:

https://yiluheihei.github.io/microbiomeMarker/articles/microbiomeMarker-vignette.html

59.   Oksanen J, Blanchet F. Package "vegan." cran.ism.ac.jp. 2022 [cited 2022 May 5];

Available from: http://cran.ism.ac.jp/web/packages/vegan/vegan.pdf

60.   Zhang Z, Wood WI. A profile hidden Markov model for signal peptides generated by

HMMER. Bioinformatics. 2003;19(2):307–8.

61.   Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 2014;30(9):1312–3.

62.   Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. Ggtree: an R Package for Visualization and

Annotation of Phylogenetic Trees With Their Covariates and Other Associated Data.

Methods Ecol Evol. 2017;8(1):28–36.

63.   Friedman J, Alm EJ. Inferring Correlation Networks from Genomic Survey Data. PLoS

Comput Biol. 2012;8(9):1–11.

64.   Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA. Sparse and

Compositionally Robust Inference of Microbial Ecological Networks. PLoS Comput Biol.

2015;11(5):1–25.

65.   Liu C, Cui Y, Li X, Yao M. Microeco: An R package for data mining in microbial

community ecology. FEMS Microbiol Ecol. 2021;97(2):1–9.

Figure Legends

Figure 1. Relative abundance of microbial orders within *P. clavata* across its geographic distribution according to 16S rRNA gene amplicons (A) and 18S rRNA gene amplicons (B) metabarcoding. *P. clavata* populations are listed from west to east, with major localities color coded. Sagres is denoted with a star as this population was excluded from further analysis since it is a different species. Microbial taxa are ordered taxonomically. Only those taxa with a median

607   relative abundance above 1% are shown, the rest are listed as "other". IS is short for Incertae sedis,

608   meaning there is uncertainty in the taxonomic position within the broader taxonomic group.

609   Relative abundance bubbles correspond to within domain (Prokaryotic or Eukaryotic) relative

610   abundance. The Sagres population was removed from further analysis after being determined to

611   be a different species.

Figure 2. Relative abundance of microbial orders within *P. clavata* across thermal susceptibility

categories according to 16S rRNA gene amplicons (A) and 18S rRNA gene amplicons (B)

metabarcoding. The Sagres population was not included in this analysis (see Methods section).

Hyper Sensitive (n_{bac} = 29; n_{euk} = 39) corals were those which showed 100% necrosis during the

heat experiment. Sensitive (n_{bac} = 3; n_{euk} = 16) corals were those that showed < 75% necrosis.

Tolerant (n_{bac} = 42; n_{euk} = 61) corals were those that showed 25-75% necrosis. Resistant (n_{bac} = 3;

n_{euk} = 7) corals showed < 25% necrosis at the end of heat-stress. Super Resistant (n_{bac} = 4; n_{euk} =

5) corals showed 0% necrosis at the end of heat-stress. Microbial taxa are ordered taxonomically.

Only those taxa with a median relative abundance above 1% are shown, the rest are listed as

"other". IS is short for Incertae sedis, meaning there is uncertainty in the taxonomic position within

the broader taxonomic group. Relative abundance bubbles correspond to within domain relative

abundance.

Figure 3. Significant differentially abundant microbial taxa between sensitive (hyper sensitive +

sensitive; n_{bac} = 32; n_{euk} = 55) vs resistant (super resistant + resistant; n_{bac} = 7; n_{euk} = 12) samples

according to ANCOMBC analysis. Natural log fold differences in abundance for each significant

taxa are shown with SD. IS is short for Incertae sedis, meaning there is uncertainty in the

taxonomic position within the broader taxonomic group. Panel A shows the bacterial taxa, while

panel B shows the eukaryotic taxa.

Figure 4. RAxML maximum-likelihood tree of Corallicolids with the primary Corallicolid ASVs

recovered from *P. clavata* placed within the tree structure using EPA. Relative abundance of each

ASV at each respective population is shown in the heatmap. Populations are ordered from west to

east, with major localities color coded. Relative abundance (green) and sample prevalence (purple)

of the ASV across the entire dataset is shown in the barplots to the right.

Figure 5. Chord plot of significant positive co-association patterns between the top microbial

families across both the 16S and 18S rRNA gene amplicon datasets for *P. clavata*. Interactions

were constructed using SPARCC and SpiecEasi with a p-value threshold of 0.05 for significance.

Self-interactions have been removed. Eukaryotic families are shown in black text, while

Prokaryotic families are shown in red text. IS is short for Incertae sedis, meaning there is

uncertainty in the taxonomic position within the broader taxonomic group.