Effect of depth on crowding investigated with a multi-depth plane display

Samuel P. Smithers[1*], Yulong Shao[1], James Altham[1], and Peter J. Bex[1*]

[1] Department of Psychology, Northeastern University, Boston, MA, USA

[*] Corresponding authors: SPS (s.smithers@northeastern.edu) and PJB (p.bex@northeastern.edu)

Abstract

Crowding occurs when highly visible objects are rendered unrecognizable in the presence of nearby features. Several studies have demonstrated that crowding can be reduced by depth differences between target and flanking features, which challenges the assumption of real-world significance because real objects typically occur at differing depths. However, most previous studies tested only small differences in depth and/or used standard stereo displays in which disparity, accommodation, and defocus blur are inconsistent with the real world. For this study we developed a multi-depth plane display consisting of three screens at different distances from the observer to test how crowding is affected by large, real differences in target-flanker depth. Overall, perceptual error decreased with increasing target-flanker spacing, but increased when the target or flankers were displayed at a different depth from fixation and one another. Our findings demonstrate that depth can increase crowding when depth differences are large. Analysis of the perceived appearance of the stimuli suggest that outside Panum's Fusional Area, crowding is increased by suppression to maintain single vision rather than by an increase in clutter because of diplopia. These findings suggest that crowding may be a more significant problem in real scenes than is estimated with 2D displays.

Introduction

Visual crowding is a phenomenon in which a peripherally viewed object, that is easily identifiable in isolation, becomes harder to recognize or identify when it is surrounded by other objects, despite still being clearly visible (Bouma, 1970; Herzog et al., 2015; Levi, 2008; Pelli, 2008; Whitney and Levi, 2011). There has been a tremendous amount of research on crowding and the topic has been extensively reviewed over the years (e.g., Herzog et al., 2015; Manassi and Whitney, 2018; Pelli, 2008; Strasburger et al., 2011; Whitney and Levi, 2011). This high level of interest is not surprising given that the natural environment is usually cluttered and mostly viewed in the peripheral visual field. Consequently, it is assumed that crowding has a large impact on our daily lives with implications for many everyday visual tasks including driving (Xia et al., 2020), reading (Legge, 2007) and face recognition (Kalpadakis-Smith et al., 2018; Louie et al., 2007). Crowding also has important clinical implications for people with central vision impairment (e.g., macular degeneration), amblyopia and dyslexia (Whitney and Levi, 2011). Through this body of research, we have been able to build a good understanding of what factors mediate and moderate crowding and develop unifying theories (Balas et al., 2009; Harrison and Bex, 2015; Keshvari and Rosenholtz, 2016).

However, despite the large body of literature, one area that remains surprisingly understudied is how crowding is affected by depth. This is important because we live in a three-dimensional environment and so to understand the significance of crowding in the real world it is essential to understand the role of depth information.

Kooi et al. (1994) were the first to investigate how depth influenced crowding. The

authors found that compared to when a peripheral target and flankers were at the same depth,

there was a release from crowding when the target and flankers were stereoscopically presented

in crossed (front) and uncrossed (behind) disparities (Kooi et al., 1994). Sayim et al. (2008)

found that when flankers were at fixation depth, there was less foveal crowding when the target

was stereoscopically presented in front or behind the flankers. Felisberti et al. (2005) also found

an effect of depth on crowding in the parafovea, but to a more limited extent. They reported a

release from crowding in two out of three subjects when the target appeared behind the flankers,

but only in one of the three subjects when the target was in front of the flankers (Felisberti et al.,

2005). Astle et al. (2014) extended on previous research by testing the effect of depth on

crowding over a range of stereoscopically produced target-flanker disparities (±800 arcsec). In

line with previous studies, Astle et al. (2014) found that when the target was at fixation depth,

increasing target-flanker disparity systematically reduced crowding. They also found that the

release from crowding was asymmetric whereby crowding was greater when the flankers were in

front of the target compared to behind it (Astle et al., 2014).

Taken together these studies appear to paint the picture that a difference in depth between

a target and flankers reduces crowding. This challenges the assumption that crowding is a

significant problem in three dimensional natural environments. However, the generalizability of

these studies to the real world is arguably limited by the fact depth cues were generated

stereoscopically. One potential problem with stereo displays is that disparity information often

does not match with accommodation or defocus blur which can detrimentally effect depth

perception (Hoffman et al., 2008; Watt et al., 2005). Furthermore, all the above studies tested

only small depth differences between target and flankers.

To our knowledge only three studies have investigated the effect of depth on crowding

using a real depth display. All three studies used two orthogonally positioned screens viewed

through a half-transparent mirror allowing stimuli to be displayed on two different depth planes

simultaneously (Eberhardt and Huckauf, 2019, 2017, 2020). Unlike previous studies that

investigated the effect of depth on crowding by varying the distance in depth between the target

and flankers, Eberhardt and Huckauf (2017) kept the target and flankers at the same depth and

presented them 0.06 diopters in front, behind or at the same depth as fixation. The authors found

that this small difference in depth between the stimuli and fixation planes had no effect on the

amount of crowding compared to when everything was presented at the fixation plane (Eberhardt

and Huckauf, 2017). Eberhardt and Huckauf (2019) repeated and expanded on this work with the

addition of a 'far' ($\pm$ 0.1 diopter) depth difference between fixation and the stimuli in addition to

the 'near' ($\pm$ 0.06 diopter) condition. The results showed that the amount of crowding was

greater when the stimuli were far from fixation compared to when they were near to fixation,

regardless of whether the stimuli were in front or behind the fixation plane (Eberhardt and

Huckauf, 2019). Most recently, Eberhardt and Huckauf (2020) used the same approach to

investigate the effect of depth on crowding when the target and flankers were separated in depth.

When flanker depth was varied while the target was at fixation depth, crowding increased with

increasing target-flanker depth difference (Eberhardt and Huckauf, 2020). This result is the

opposite to the findings of other studies in which small differences in depth between the target

and flankers were introduced with stereoscopic disparity (Astle et al., 2014; Felisberti et al.,

2005; Kooi et al., 1994; Sayim et al., 2008). Furthermore, the authors also found that when

flanker depth was varied while the target was at fixation depth, crowding was greater when the

flankers were behind the target and fixation compared to in front of it (Eberhardt and Huckauf,

2020). When target depth was varied while the flankers were at fixation, there was more

crowding when the target was in front of the flankers and fixation compared to when it was

behind them (Eberhardt and Huckauf, 2020).

In light of the above discrepancies, there is a clear need for further research to improve

our understanding of how depth information affects crowding. There is a particular need for

studies that use real depth and that test large depth differences that reflect those experienced in

the real world. Therefore, the aim of this study was to investigate the effect of real depth on

crowding using a multi-depth plane display consisting of three screens at 0.4 m, 1.26 m, and 4 m.

Methods

Subject recruitment

A total of 110 naïve subjects were recruited from the Northeastern undergraduate population in

exchange for course credit as compensation for their time. Of these, seven were excluded

immediately following screening because they did not meet the inclusion criteria of normal or

corrected-to-normal vision. An additional five subjects were excluded because of issues with the

experimental code (N=3), calibration issues (N=1) or because they did not follow the task

instructions (N=1). Thus, a total of 98 subjects (32 male, 66 female) contributed usable data

across five separate experiments (22 in experiment 1, 19 in experiment 2, 21 in experiments 3

and 4, and 15 in experiment 5). All subjects read and signed an informed consent form approved

by the University Ethics Board before the experiment began. The experimental procedure was

approved by the institutional review board at Northeastern University, and the experiment was

performed in accordance with the tenets of the Declaration of Helsinki.

Setup

We developed a multi-depth plane display consisting of three screens at 0.4 m (near screen), 1.26

m (mid screen) and 4 m (far screen) from the observer (Figure 1). These distances were chosen

because 4 m is close to optical infinity while 0.4 m is the clinical standard for near distance, with

1.26 m being in the middle of these two extremes in log units. For the far screen we used an

ultra-short throw projector (Bomaker Polaris 4K Laser Projector) projected onto a 240 cm x 250

cm white wall at the end of the room. The mid and near screens were made from 101.5 cm x 76

cm and 46 cm x 36.5 cm clear 5mm acrylic respectively held in position by a frame constructed

from extruded aluminum rails. The transparent acrylic was covered on one side with ClearBright

transparent display film (Lux Labs Inc., Boston, MA, USA). This nanoparticle film enabled us to

display images from two projectors (InFocus IN3116) onto the near and mid screens while

preserving their transparency (Hsu et al., 2014), thus allowing observers to simultaneously view

adjacent stimuli presented on all three depths planes at the same time. The screens were

temporally synchronized and spatially aligned to ensure display timing and position were

consistent across the three depth planes.

Figure 1: Illustration of the multi-depth plane display setup used for this study.

Spatial calibration

Each observer performed a subjective spatial calibration at the start of Experiments 1-4 to ensure

that the angular positions of the stimuli and fixation point relative to primary gaze were the same

(or as close as realistically possible) across all three depth planes. The center of the far screen

(which was at primary gaze when the observers head was positioned in the chin and head rest)

was used as the fixed reference against which all three screens were calibrated. With their head

in the chin and head rest observers were instructed to sequentially position a series of targets so

that they were in the same position as a fixed reference target that was displayed on the reference

screen. The calibration was performed under monocular conditions using the observer's

dominant eye (determined by the Miles test). The calibration was performed monocularly

because during a preliminary experiment subjects reported finding the task too difficult to

complete under binocular conditions (e.g., because of diplopia, an issue that is considered further

in the Discussion section). For Experiment 5, the spatial calibration was performed by the

experimenter prior to the experiment using a webcam positioned at the observer's cyclopean

point. Observers were asked to maintain the same head position within the chin and head rest

throughout the experiment.

Stimuli and experimental conditions

The target and flankers were inspired by those used in previous studies (Harrison and Bex, 2017,

2015). The target was a white Landolt-C displayed against a black background at 10˚ eccentricity

to the right or left of the fixation point at primary gaze. Following the structure of a Landolt-C

optotype, the target was 1˚ in dimeter with a 0.25˚ line width and a 0.25˚ opening. The opening

orientation of the target was random each trial, drawn from a uniform 360˚ distribution. The

target was surrounded by a white flanker ring formed of four circular arcs (Figure 2). In each

depth combination described below, the outer edge of the target was separated from the inner

edge of the flanker ring by 5˚, 2.5˚, 1.25˚ or 0.625˚. These distances of separation represented

1/2, 1/4, 1/8 and 1/16 of the target eccentricity respectively. The line width and size of the four

gaps in the flanker ring were kept constant (0.25˚). For each target depth, a control condition

with no flanker ring was also included. The fixation point consisted of a 1˚ white circle with a

black cross (Figure 2). This design was used because it has been reported to elicit more stable

fixation compared to a plain circle or cross hair alone (Thaler et al., 2013). The background on

all screens was always set to the lowest RGB (i.e. R=G=B=0) value which was transparent on

the near and mid screens, and black on the far screen. Since the mid screen was dimmest

(R=G=B=255, white=13.56 cd/m^2), the luminance of stimuli displayed on the near and far

screens were calibrated to match the luminance of the mid screen using a Photo Research PR-

650 spectroradiometer (Photo Research, Chatsworth, CA, USA). The room lights were switched

off for the duration of the experiment.

Figure 2: Schematic illustration of experimental stimuli and fixation marker. In all conditions, except the

no-flanker condition, the randomly orientated target (a white Landolt-C) was surrounded by a white

flanker ring at one of four possible target-flanker distances. The stimuli was viewed against a black

background.

A summary of the stimulus conditions tested in each experiment is provide in Table 1. Briefly, in

Experiments 1 and 2 the target was always presented at fixation depth while the depth of the

flanker was varied. In Experiment 1 the target and fixation point were always on the mid screen

while each flanker was displayed in front, behind or at fixation depth. In Experiment 2 the target

and fixation point were presented on the near or far screen while the flanker was presented at one

of the three depths across trials for each fixation distance. Experiments 3 and 4 were the same as

Experiments 1 and 2 except that the flanker ring was presented at fixation depth while the target

depth was varied. In Experiment 3, the flanker and fixation point were always on the mid screen

while the target was displayed on one of the three depth planes across trials. In Experiment 4, the

flankers and fixation point were presented on the near or far screen while the target was

presented at each of the three depths across trials for each fixation distance. For Experiment 5,

the target, flanker ring and fixation were presented together on the same screen at three different

fixation depths. This was to investigate whether there was an effect of absolute viewing distance

on crowding when there was no depth difference between the target and flanker ring.

Table 1: Summary of the stimulus conditions in each experiment. The same color scheme is used to represent the near (0.4 m), mid (1.26 m) and far (4 m) screens throughout the manuscript. Note that for target-flanker spacing, 'infinity' = no flanker ring.

Experiment	Fixation depth (m)	Target depth (m)	Flanker depth (m)	Target-flanker spacing (degrees)	No. of stimulus conditions	Total no. presentations	Sample size
Exp 1	Mid (1.26)	Mid (1.26)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°	13	130 (10 reps per stimulus)	N = 22
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Mid (1.26)	Mid (1.26)	Far (4)	0.625°, 1.25°, 2.5°, 5°			
Exp 2	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	26	260 (10 reps per stimulus)	N = 19
	Near (0.4)	Near (0.4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°			
	Near (0.4)	Near (0.4)	Far (4)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 3	Mid (1.26)	Near (0.4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity	15	150 (10 reps per stimulus)	N = 21
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Mid (1.26)	Far (4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 4	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	30	240 (8 reps per stimulus)	N = 21
	Near (0.4)	Mid (1.26)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Near (0.4)	Far (4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Near (0.4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Mid (1.26)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 5	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	15	150 (10 reps per stimulus)	N = 15
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			

Each condition was repeated 10 times in Experiments 1, 2, 3 and 5 and eight times in

Experiment 4, with equal number of repeats on either side of the fixation point. The order of

testing was randomized.

An example trial is illustrated in Figure 3. Briefly, at the start of each trial, observers

were instructed to focus on the central fixation point. The fixation point appeared on the fixation

screen 1 sec before stimulus onset in Experiments 1, 3 and 5, and 1.5 sec before stimulus onset in

Experiments 2 and 4. The target and flanker ring (if present) were displayed for 0.25 sec. The

fixation point remained on the screen for 1 sec before being replaced by a randomly oriented

Landolt-C that was the same size as the target Landolt-C. The observer used the mouse to rotate

this Landolt-C until it matched the perceived orientation of the target and then clicked to report

the orientation. In all experiments except Experiment 5, the observer then reported the perceived

location of the target relative to the position of the flanker ring. There were 5 options to choose

from, "target in center of ring", "target inside ring but not in the center", "ring obstructs target",

"target outside ring", and "unsure or no ring". Observers were instructed to select the option that

was the closest match to how they perceived the stimulus. If an observer missed the stimulus or

they were unsure of the targets' location relative to the flanker ring (e.g., because of diplopia)

they were instructed to select the "unsure or no ring" option. Following this selection, the next

trial started immediately. Before starting an experiment, all participants underwent training

during which they were encouraged to complete as many practice trials as they needed to

familiarize themselves with the task. These practice trials did not contribute to the collected data.

Figure 3: A step-by-step example of a single trial. Note that step five was not

included in Experiment 5.

Data processing and statistical analysis

<u>Perceptual error</u>

To quantify crowding for each stimulus condition we first calculated the report error for each

presentation, i.e., the angular difference between the reported gap orientation of the target and

the actual gap orientation. For each subject, this generated 10 report errors per stimulus condition

in Experiments 1, 2, 3 and 5, and eight per stimulus condition in Experiment 4. We then used

these report errors to calculate the circular standard deviation for each stimulus condition using

the Circular Statistics Toolbox (Berens, 2009) in MATLAB R2022a (MathWorks Inc., Natick,

MA, USA). We refer to this measure as perceptual error, which is reported in degrees.

The effect of target/flanker depth and target-flanker spacing on perceptual error was

analyzed using mixed effects multiple linear regression, fitted using the lme4 package (Bates et

al., 2015) in R (R Core Team, 2022). Perceptual error was used as the response (dependent)

variable after being natural log transformed to correct for positive skew. For all five experiments,

we included target-flanker spacing as a fixed effect. We also included flanker depth for

Experiments 1 and 2, target depth for Experiments 3 and 4 or target-flanker-fixation depth for

Experiment 5. Target-fixation depth or flanker-fixation depth were also included as a third fixed

effect for Experiment 2 and 4 respectively. All possible pairwise interactions of fixed effects

were fitted in the full model. Subject identification (ID) was included as a random effect in all

experiments to control for repeated measures. We first used model simplification, whereby

models were compared with one another using a likelihood ratio test (LRT), to remove non-

significant interactions sequentially. This enabled us to test the effect of individual fixed effects

that were not involved in an interaction. We then used the minimum model (i.e., model

containing all fixed effects of interest plus any significant two-way interactions) to report the

significance of each main effect or important interaction(s) by using a LRT to compare the

minimum model with the same model but with the effect/interaction of interest removed. Note

that data for the control condition (i.e., no flanker ring) were excluded from statistical analysis.

This was to ensure that a) the fixed effects remained independent and b) the effect of having no

flanker ring did not mask any of the effects we were most interested in (see Supplementary

Material for a more detailed explanation).

<u>Perceived target position relative to flanker ring</u>

For statistical analysis, the results for the perceived target position relative to the flanker ring

were converted to a binary response. A value of 1 meant that the observer perceived the target as

being inside the flanker ring (i.e., they chose either "target in center of ring" or "target inside ring

but not in the center") while a value of 0 meant the subject did not perceive the target as being

inside the flanker ring (i.e., they chose "ring obstructs target", "target outside ring" or "unsure or

no ring").

These data were then analyzed using mixed effects binary logistic regression, fitted using

the lme4 package (Bates et al., 2015) in R (R Core Team, 2022). We used the DHARMa package

(Hartig, 2022) to run residual diagnostics that are often overlooked in logistic regression.

Perceived target position was used as the binary response variable (i.e., 1 = target inside ring and

0 = target not inside ring). The fixed effect, and pairwise interaction for each experiment were

the same as those described above for the analysis of perceptual error. Subject ID was included

as a random effect in all experiments to control for repeated measures. Model simplification (to

remove non-significant interactions) and significance testing was the same as described above

for the analysis of perceptual error. Data for the control condition (i.e., no flanker ring) were

excluded from statistical analysis for the same reasons as those outlined above.

The separate results showing the number of times observers chose each of the five

choices for perceived target position relative to the flanker ring can be found in the

Supplementary Material (Figures S1-S4).

Results

Perceptual error

<u>Target always at fixation depth with variable flanker depth</u>

Experiment 1: Target and fixation at 1.26 m with flanker ring presented at each depth (Figure 4A). Perceptual error was higher when the flanker ring was displayed in front of, or behind, the target and fixation plane (mixed effects multiple linear regression models compared via LRT: $\chi^2_{(2)} = 20.67$, p <0.001). As expected, perceptual error decreased as target-flanker spacing increased ($\chi^2_{(1)} = 31.71$, p <0.001).

Experiment 2: Target and fixation at 0.4 m or 4 m with flanker ring presented at each depth (Figure 4B). Perceptual error tended to increase with increasing flanker depth when the flanker ring was behind the target and fixation plane, however, the effect of flanker depth was much weaker when the flanker ring was in front of the target and fixation (interaction between target-fixation depth and flanker depth: $\chi^2_{(2)} = 17.18$, p<0.001). Perceptual error decreased with increasing target-flanker spacing, in a manner corresponding to target-fixation depth (interaction between target-flanker spacing and target-fixation depth: $\chi^2_{(1)} = 9.2$, p=0.002).

Figure 4: Perceptual error results for Experiments 1 (A) and 2 (B) in which flanker depth was varied while the target was always at fixation depth. Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below each graph shows the depth of the flanker ring (based on the same color key as the box plot) relative to the depth of the fixation point and target (shown in black) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker spacing due to limited space within the figure.

Flanker ring always at fixation depth with variable target depth

Experiment 3: Flanker ring and fixation at 1.26 m with target presented at each depth (Figure 5A). Overall, perceptual error was higher when the target was displayed in front of, or behind, the flanker ring and fixation ($\chi^2_{(2)} = 36.93$, p <0.001), with the greatest perceptual error occurring when the target was behind the flanker ring and fixation. Perceptual error decreased as target-flanker spacing increased ($\chi^2_{(1)} = 45.99$, p <0.001).

Experiment 4: Flanker ring and fixation at 0.4 m or 4 m with target presented at each depth (Figure 5B). Perceptual error was greatest when the target was displayed behind the flanker ring and fixation, while the effect of depth was less prominent when the target was displayed in front of the flanker ring and fixation (interaction between flanker-fixation depth and target depth: $\chi^2_{(2)} = 61.87$, p<0.001). Perceptual error decreased as target-flanker spacing increased ($\chi^2_{(1)} = 29.92$, p<0.001).

Figure 5: **Perceptual error results for Experiments 3 (A) and 4 (B) in which target depth was varied while the flanker ring was always at fixation depth.** Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below each graph shows the depth of the target (based on the same color key as the box plot) relative to the depth of the fixation point and flanker ring (shown in black) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker spacing due to limited space within the figure.

Target and flanker ring at fixation depth

Experiment 5: Target and flanker ring always at fixation depth (Figure 6). Perceptual error decreased as target-flanker spacing increased ($\chi^2_{(1)}$ = 64.8, p<0.001). There was also a significant effect of depth ($\chi^2_{(2)}$ = 8.75, p=0.013) resulting from a higher perceptual error for stimuli displayed at 4 m.

Figure 6: **Perceptual error results for Experiment 5 in which the target and flanker were always**

presented at the same depth as the fixation point. Box plots show medians plus the interquartile range

(IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and

lower quartiles. Points show the perceptual error for each individual observer. The visual aid below the

graph shows the depth of the target, flanker, and fixation point (based on the same color key as the box

plot) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target,

and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker

spacing.

Perceived target position relative to the flanker ring

Target always at fixation depth with variable flanker depth

Experiment 1: Target and fixation at 1.26 m with flanker ring presented at each depth

(Figure 7A and supplementary Figure S1). In Experiment 1, most observers consistently

reported seeing the target inside the flanker ring. However, when the flanker ring was away from

target-fixation depth there was a reduction in the proportion of observers who consistently

reported seeing the target within the flanker ring when the target-flanker spacing was small

(mixed effects binary logistic regression models compared via LRT; interaction between target-

flanker spacing and flanker depth: $\chi^2_{(2)} = 36.12$, p <0.001). This effect was greatest when the

flanker ring was displayed behind the target and fixation.

Note that in all four experiments, the response to the control (i.e., no flanker condition)

indicates the probability of an observer making an inaccurate report (e.g., they selected the

wrong option by mistake or missed the stimulus and reported a guess rather than selecting the

"Unsure" option).

Experiment 2: Target and fixation at 0.4 m or 4 m with flanker ring presented at each

depth (Figure 7B and supplementary Figure S2). As with the previous experiments, when the

flanker ring was away from target-fixation depth, there was a reduction in the overall proportion

of trials in which observers reported seeing the target within the flanker ring (interaction between

target-fixation depth and flanker depth: $\chi^2_{(2)} = 374.26$, p < 0.001). The greatest reduction

occurred when target-flanker spacing was smaller ($\chi^2_{(1)} = 178.65$, p < 0.001).

Figure 7: **Proportion of trials in which observers reported seeing the target inside the flanker ring**

for Experiments 1 (A) and 2 (B). In both experiments flanker depth was varied while the target was

always at fixation depth**.** Box plots show medians plus the interquartile range (IQR), whiskers are the

lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points

show the proportion of trials for each individual observer. The visual aid below each graph shows the

depth of the flanker ring (based on the same color key as the box plot) relative to the depth of the fixation

point and target (shown in black) for each stimulus condition. The relative sizes of, and distances

between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown

for the 0.625˚ target-flanker spacing due to limited space within the figure.

<u>Flanker ring always at fixation depth with variable target depth</u>

Experiment 3: Flanker ring and fixation at 1.26 m with target presented at each depth

(Figure 8A and supplementary Figure S3). When the target was away from flanker-fixation

depth, observers were less likely to report seeing the target within the flanker ring. This effect

was greatest when target-flanker spacing was smaller and when the target was displayed in front

of the flanker ring and fixation (interaction between target-flanker spacing and target depth: $\chi^2_{(2)}$

$= 100.38$, p <0.001).

Experiment 4: Flanker ring and fixation at 0.4 m or 4 m with target presented at each

depth (Figure 8B and supplementary Figure S4). As with the previous experiment, the greater

the difference between target depth and flanker-fixation depth, the less likely observers were to

report seeing the target within the flanker ring (interaction between flanker-fixation depth and

target depth: $\chi^2_{(2)} = 779.49$, p < 0.001), particularly when target-flanker spacing was small

(interaction between target-flanker spacing and target depth: $\chi^2_{(2)} = 23.27$, p < 0.001). However,

target-flanker spacing had a weaker effect on the proportion of trials in which observers reported

seeing the target within the flanker ring when the flanker ring and fixation were near compared

to far (interaction between target-flanker spacing and flanker-fixation depth: $\chi^2_{(1)} = 56.23$, p $<$

0.001).

Figure 8: **Proportion of trials in which observers reported seeing the target inside the flanker ring for Experiments 3 (A) and 4 (B).** In both experiments target depth was varied while the flanker ring was always at fixation depth**.** Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the proportion of trials for each individual observer. The visual aid below each graph shows the depth of the target (based on the same color key as the box plot) relative to the depth of the fixation point and flanker ring (shown in black) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker spacing due to limited space within the figure.

Discussion

In all experiments, perceptual error decreased as the target-flanker spacing increased, which is in line with previous crowding research (Bouma, 1970; Pelli, 2008). Overall, perceptual error was higher when the target or flanker ring were presented at a different depth from the other and from fixation. The greatest differences in perceptual error among the different depth conditions occurred when the target or flanker ring were presented behind fixation depth. In contrast, when the target or flanker ring were presented in front of fixation depth, the differences in perceptual error among depth conditions were notably smaller. The results also show that

when the target, flanker ring and fixation were all at the same depth plane, crowding tended to

increase with the absolute distance of the stimuli from the observer.

The main finding that a depth difference between the flankers and the target led to an

increase, rather than a decrease, in perceptual error is different from the findings of several

previous studies that investigated the effect of depth on crowding using a stereoscopic

presentation (Astle et al., 2014; Felisberti et al., 2005; Kooi et al., 1994; Sayim et al., 2008). For

example, Astle et al (2014) found that when the target was presented at fixation depth

(comparable to Experiments 1 and 2 in our study), increasing the disparity between the target and

flankers resulted in less crowding, and ultimately a complete release from crowding. The authors

also reported that crowding was greater when flankers were presented in crossed disparity (in

front of the target and fixation), compared to uncrossed disparity (behind the target and fixation)

(Astle et al., 2014). Similarly, when the flankers were presented at fixation depth (comparable to

Experiments 3 and 4 in our study), Sayim et al (2008) found that foveal crowding was lower

when the target was presented in front of or behind the flanker and fixation.

Several explanations for crowding have been proposed in which crowding occurs because

target and flanking features are inappropriately grouped (Banks et al., 1979; Pachai et al., 2016).

Consistent with this explanation, crowding can be alleviated when the flankers are grouped

together and segmented from the target (Herzog and Manassi, 2015). One piece of evidence in

support of this account is the reduction of crowding that can occur when flankers are presented at

a different depth from the target (Astle et al., 2014; Felisberti et al., 2005; Kooi et al., 1994;

Sayim et al., 2008). Our results challenge this view of crowding because it would predict that

increasing the depth difference between a target and flankers would decrease the probability of

the flankers being grouped and increase the probability of their being segmented from the target,

thereby decreasing crowding, the opposite of our finding. Other models of crowding, such as

texture processing (Balas et al., 2009; Freeman and Simoncelli, 2011; Parkes et al., 2001; Wallis

and Bex, 2012), sampling errors (Ester et al., 2015, 2014; Harrison and Bex, 2017, 2016, 2015),

attentional resolution limits (He et al., 1996) or saccade properties (Nandy and Tjan, 2012) are

agnostic concerning the influence of depth differences among targets and flankers. Therefore,

each of these models would require modification to explain the increase in crowding that we

observe with large real depth differences between a target and its flankers.

In contrast with studies that controlled depth with small stereoscopic disparities, our

results do agree with the findings of Eberhardt and Huckauf (2020) who also used a real-depth

display and tested larger differences in depth between the target and flankers than previous

studies had examined. When the target was at fixation depth (comparable to Experiments 1 and 2

in our study), Eberhardt and Huckauf (2020) found that crowding increased with increasing

target-flanker depth difference. The authors also found that crowding was greater when the

flankers were behind the target and fixation compared to in front of it (Eberhardt and Huckauf,

2020). Our findings from Experiments 1 and 2 are consistent with these results. However, when

the flankers were at fixation depth (comparable to Experiments 3 and 4 in our study), Eberhardt

and Huckauf (2020) found that crowding was higher when the target was in front of the flankers

and fixation compared to when it was behind them. In contrast, in Experiments 3 and 4 we found

that perceptual error was greatest when the target was presented behind the flanker ring and

fixation rather than in front of it. This difference may be due to a combination of defocus blur

when the target was behind fixation combined with the finding from Experiment 5 which found

that in our study the absolute distance from the observer resulted in higher perceptual error.

Why the inconsistencies across studies? It is important to consider that most of the

previous studies used very small depth differences, the largest being 0.1 diopters in Eberhardt

and Huckauf (2020) and Eberhardt and Huckauf (2019). In comparison, the depth difference in

our study ranged from 0.54 diopters (between the mid and far screens) to 2.25 diopters (between

the near and far screens). This difference has several implications that may help to explain the

results. It is possible that small differences in depth between the target and flankers reduce

crowding (Astle et al., 2014; Felisberti et al., 2005; Kooi et al., 1994; Sayim et al., 2008), while

larger differences in depth increase it (Eberhardt and Huckauf, 2020; and results from our study)

and models of crowding will simply have to accommodate this peculiar property. Although this

pattern is quite surprising, there is precedent for non-linear effects of depth on perceptual

processing outside the crowding literature. In a multiple object tracking task, when the targets

and distractors were distributed equally on two depth planes, tracking accuracy was better than

the accuracy during trials in which targets and distractors appeared on a single depth plane when

the depth difference between planes was small (~6 cm) (Ur Rehman et al., 2015; Viswanathan

and Mingolla, 2002), but worse when the depth difference between planes was large (50 cm) (Ur

Rehman et al., 2015).

How could depth affect perception in this nonlinear way? The geometry of the eyes and

the 3D environment create interocular disparity differences between the images in each eye.

When corresponding features in the images in each eye can be identified, the images can be

fused to generate a binocular perceptual experience of depth (for review see Nityananda and

Read, 2017)). There is a horizontal and vertical spatial limit to the interocular disparity

difference over which the visual system attempts to identify corresponding retinal points in order

to fuse different 2D interocular images as a single 3D object (Westheimer, 1979). This 2D spatial

correspondence limit creates a 3D volume along the horopter known as Panum's Fusional Area

(Panum, 1858). Outside Panum's Fusional Area, there is a loss of stereoscopic depth perception

and either double vision (diplopia) may be experienced, or single vision may occur if the image

from one eye is suppressed (Georgeson and Wallis, 2014; Ono et al., 1977). The spatial scale of

Stereoscopic Panum's Fusional Area depends on the spatial frequency of the stimuli (Maiello et

al., 2020; Pulliam, 1981; Reynaud and Hess, 2017; Wilcox and Allison, 2009; Yang and Blake,

1991) and increases with eccentricity (Hampton and Kertesz, 1983; Qin et al., 2006). It is

possible that differences between fusion and depth perception within Panum's Fusional Area and

diplopia or suppression outside of it, may explain the patterns of results observed between

studies. Small stereoscopic disparity differences between target and flankers create differences in

perceived depth that can alleviate the effects of crowding, potentially through grouping (Herzog

and Manassi, 2015) or by independent processing of target and flankers in different depth-

selective channels (Norcia et al., 1985; Pulliam, 1981; Reynaud and Hess, 2017; Tyler et al.,

1994; Wilcox and Allison, 2009; Yang and Blake, 1991) by any of the computational models of

crowding based on texture processing (Balas et al., 2009; Freeman and Simoncelli, 2011; Parkes

et al., 2001; Wallis and Bex, 2012), sampling errors (Ester et al., 2015, 2014; Harrison and Bex,

2017, 2016, 2015) attentional resolution limits (He et al., 1996) or saccade properties (Nandy and

Tjan, 2012). Large interocular disparity differences may induce diplopia or suppression that

could increase perceptual error by increasing the number of features (diplopia) or removing the

benefit of binocular summation (suppression) that can help decrease the effects of crowding

(Siman-Tov et al., 2021).

In this study, when asked, only a few observers reported experiencing diplopia, and those

that did, reported only experiencing it in a minority of presentations for which they selected the

'unsure or no ring' option when reporting the perceived location of the target. As mentioned in

the Methods sections, observers did report experiencing diplopia when trying to perform the

spatial calibration task under binocular conditions during a pilot experiment. However, it should

be noted that the spatial calibration required observers to simultaneously concentrate on stimuli

at two different depths for an extended length of time. The calibration is therefore a

fundamentally different task from our crowding experiments in which stimuli were only briefly

presented in the observer's peripheral visual field, which could explain why few observers

reported diplopia in the main experiments. If the image from one eye were partly suppressed to

prevent diplopia, the reported location of the target relative to the flanker ring offers us potential

clues as to which eye was likely being suppressed. For instance, assuming no head movement

following the spatial calibration performed using the dominant eye, for presentations in which

the observer reported that the target was inside the flanker ring, it's likely that it was the image

from the non-calibrated eye that was suppressed. Conversely, if the observer reported seeing the

target outside the flanker ring, or that the target was obscured by the edge of the flanker ring,

then this would instead imply it was the image from the calibrated eye that was being

suppressed. However, as well as suppression of the calibrated eye, a perceived misalignment of

the target and flanker ring could also mean that the observer didn't maintain the same head

position as they had during the calibration. Although a chin and head rest were used to ensure the

observer maintained a steady head position throughout the experiment, it remains possible that

small head movements did occur. Furthermore, the larger the depth difference between the target

and flanker ring, and the smaller the target-flanker spacing, the more sensitive the spatial

calibration would have been to changes in head position. Indeed, Figures 7 and 8 show that when

either the target or flanker ring were away from fixation depth, participants reported seeing the

target in the center of the flanker ring on fewer trials when the target-flanker spacing was small.

Given that observers were less likely to perceive the target inside the flanker ring when

the target and flanker ring were separated by depth, one explanation for the higher perceptual

error could be perceptual occlusion of the target by the flanker ring. To examine how the trials in

which observers reported that the target was not inside the flanker ring (i.e., target obstructed

ring, target was outside ring, or they were unsure) affected the overall results, we recalculated

perceptual error using only trials in which the observer reported seeing the target inside the

flanker ring (supplementary Figures S5 and S6) and reanalyzed these data the same way as for

perceptual error calculated from the full data set. Apart from Experiment 2, the effect of flanker

and/or target depth remained statistically significant (see supplementary material for analysis

results) meaning that across the majority of experiments misalignment or occlusion of the target

by the flanker ring is not enough to explain the overall results. This can be seen visually by

comparing Figures 4 and 5 with their corresponding figures in the supplementary material

(Figures S5 and S6 respectively).

Our findings do not support the widely held view that depth differences between targets

and flankers always reduces crowding, instead the effect of depth on crowding appears to be

more complex than originally assumed. Our results, alongside those of previous studies, suggest

that small differences in depth between targets and flankers do not have the same effect on

crowding as larger differences in depth in which either the target or flankers are presented

outside the limits of Panum's Fusional Area. We suggest that large differences in depth between

target and flankers can increase crowding because of suppression to maintain single vision as this

removes the benefits of binocular summation. This property confirms the importance of

crowding in natural scenes.

Acknowledgements

561 We thank members of the Bex lab and Eskew lab at Northeastern University for valuable

562 feedback during pilot experiments. This work was supported by NIH grant R01EY032162

563 (P.J.B.). The Northeastern University IRB approved procedures, and all observers consented to

564 participation.

Author Contributions

567 S.P.S. and P.J.B. conceived the study and the idea of the multi-depth plane display and designed

568 the experiments. S.P.S. designed and built the multi-depth plane display, wrote all code, and

569 performed analyses. S.P.S., Y.S., and J.A. conducted the experiments. S.P.S. and P.J.B. wrote the

570 manuscript with input from all authors.

Data Availability

573 All data and code required to replicate the analyses in this paper will be made available on an

574 online repository at the stage of publication. It will also be made available to reviewers during

575 peer review.

References

Astle AT, McGovern DP, McGraw P V. 2014. Characterizing the role of disparity information in alleviating visual crowding. *J Vis* **14**:1–14. doi:10.1167/14.6.8

Balas B, Nakano L, Rosenholtz R. 2009. A summary-statistic representation in peripheral vision explains visual crowding. *J Vis* **9**:1–18. doi:10.1167/9.12.1

Banks WP, Larson DW, Prinzmetal W. 1979. Asymmetry of visual interference. *Percept Psychophys* **25**:447–456. doi:10.3758/BF03213822

Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *J Stat Softw* **67**:1–48. doi:10.18637/jss.v067.i01

Berens P. 2009. CircStat: a MATLAB toolbox for circular statistics. *J Stat Softw* **31**:1–29.

Bouma H. 1970. Interaction effects in parafoveal letter recognition. *Nature* **226**:177–178.

Eberhardt LV, Huckauf A. 2019. Crowding in depth for binocular and monocular observation. *Attention, Perception, Psychophys* **81**:1951–1961.

Eberhardt LV, Huckauf A. 2017. Examining crowding using a real three-dimensional experimental setup. *Proc Latv Acad Sci* **71**:313–319. doi:10.1515/prolas-2017-0055

Eberhardt L V, Huckauf A. 2020. Crowding effects across depth are fixation-centered for defocused flankers and observer-centered for defocused targets. *Brian Sci* **10**:596. doi:10.3390/brainsci10090596

Ester EF, Klee D, Awh E. 2014. Visual crowding cannot be wholly explained by feature pooling. *J Exp Psychol Hum Percept Perform* **40**:1022–1033. doi:10.1037/a0035377

Ester EF, Zilber E, Serences JT. 2015. Substitution and pooling in visual crowding induced by similar and dissimilar distractors. *J Vis* **15**:1–12. doi:10.1167/15.1.4

Felisberti FM, Solomon JA, Morgan MJ. 2005. The role of target salience in crowding.

Perception **34**:823–833. doi:10.1068/p5206

Freeman J, Simoncelli EP. 2011. Metamers of the ventral stream. *Nat Neurosci* **14**:1195–1204.

doi:10.1038/nn.2889

Georgeson MA, Wallis SA. 2014. Binocular fusion, suppression and diplopia for blurred edges.

Ophthalmic Physiol Opt **34**:163–185. doi:10.1111/opo.12108

Hampton DR, Kertesz AE. 1983. The extent of Panum's area and the human cortical

magnification factor. *Perception* **12**:161–165. doi:10.1068/p120161

Harrison WJ, Bex PJ. 2017. Visual crowding is a combination of an increase of positional

uncertainty, source confusion, and featural averaging. *Sci Rep* **7**:45551.

doi:10.1038/srep45551

Harrison WJ, Bex PJ. 2016. Ungrouping overcrowded explanations of visual crowding: reply to

Pachai et al. *Curr Biol* **26**:R353–R354. doi:10.1016/j.cub.2016.03.024

Harrison WJ, Bex PJ. 2015. A unifying model of orientation crowding in peripheral vision. *Curr*

Biol **25**:3213–3219. doi:10.1016/j.cub.2015.10.052

Hartig F. 2022. DHARMa: Residual diagnostics for hierarchical (multi-level/mixed) regression

models. R package version 0.4.6. http://florianhartig.github.io/DHARMa/

He S, Cavanagh P, Lntriligator J. 1996. Attentional resolution and the locus of visual awareness.

Nature **383**:334–337. doi:10.1038/258748a0

Herzog MH, Manassi M. 2015. Uncorking the bottleneck of crowding: a fresh look at object

recognition. *Curr Opin Behav Sci* **1**:86–93. doi:10.1016/j.cobeha.2014.10.006

Herzog MH, Sayim B, Chicherov V, Manassi M. 2015. Crowding, grouping, and object

recognition: a matter of appearance. *J Vis* **15**:1–18. doi:10.1167/15.6.5

Hoffman DM, Girshick AR, Akeley K, Banks MS. 2008. Vergence-accommodation conflicts

hinder visual performance and cause visual fatigue. *J Vis* **8**:1–30. doi:10.1167/8.3.33

Hsu CW, Zhen B, Qiu W, Shapira O, DeLacy BG, Joannopoulos JD, Soljačić M. 2014.

Transparent displays enabled by resonant nanoparticle scattering. *Nat Commun* **5**:1–6.

doi:10.1038/ncomms4152

Kalpadakis-Smith A V., Goffaux V, Greenwood JA. 2018. Crowding for faces is determined by

visual (not holistic) similarity: Evidence from judgements of eye position. *Sci Rep* **8**:12556.

doi:10.1038/s41598-018-30900-0

Keshvari S, Rosenholtz R. 2016. Pooling of continuous features provides a unifying account of

crowding. *J Vis* **16**:1–15. doi:10.1167/16.3.39

Kooi FL, Toet A, Tripathy SP, Levi DM. 1994. The effect of similarity and duration on spatial

interaction in peripheral vision. *Spat Vis* **8**:255–279. doi:10.1163/156856894X00350

Legge GE. 2007. Psychophysics of reading in normal and low vision. Mahwah, NJ: Lawrence

Erlbaum Associates Publishers.

Levi DM. 2008. Crowding - an essential bottleneck for object recognition: a mini-review. *Vision*

Res **48**:635–654. doi:10.1016/j.visres.2007.12.009

Louie EG, Bressler DW, Whitney D. 2007. Holistic crowding: selective interference between

configural representations of faces in crowded scenes. *J Vis* **7**:1–11. doi:10.1167/7.2.24

Maiello G, Chessa M, Bex PJ, Solari F. 2020. Near-optimal combination of disparity across a

log-polar scaled visual field. *PLoS Comput Biol* **16**:1–28. doi:10.1371/journal.pcbi.1007699

Manassi M, Whitney D. 2018. Multi-level crowding and the paradox of object recognition in

clutter. *Curr Biol* **28**:R127–R133. doi:10.1016/j.cub.2017.12.051

Nandy AS, Tjan BS. 2012. Saccade-confounded image statistics explain visual crowding. *Nat*

Neurosci **15**:463–469. doi:10.1038/nn.3021

Nityananda V, Read JCAA. 2017. Stereopsis in animals: evolution, function and mechanisms. *J*

Exp Biol **220**:2502–2512. doi:10.1242/jeb.143883

Norcia AM, Suiter EE, Tyler CW. 1985. Electrophysiological evidence for the existence of

coarse and fine disparity mechanisms in human. *Vision Res* **25**:1603–1611.

doi:10.1016/0042-6989(85)90130-0

Ono H, Angus R, Gregor P. 1977. Binocular single vision achieved by fusion and suppression.

Percept Psychophys **21**:513–521. doi:10.3758/BF03198731

Pachai M V., Doerig AC, Herzog MH. 2016. How best to unify crowding? *Curr Biol* **26**:R352–

R353. doi:10.1016/j.cub.2016.03.003

Panum PL. 1858. Physiologische untersuchungen über das sehen mit zwei augen. Kiel:

Schwerssche Buchhandlung.

Parkes L, Lund J, Angelucci A, Solomon JA, Morgan M. 2001. Compulsory averaging of

crowded orientation signals in human vision. *Nat Neurosci* **4**:739–744. doi:10.1038/89532

Pelli DG. 2008. Crowding: a cortical constraint on object recognition. *Curr Opin Neurobiol*

18:445–451. doi:10.1016/j.conb.2008.09.008

Pulliam K. 1981. Spatial frequency analysis of three-dimensional vision. *Proc SPIE Vis Simul*

Image Realis II **303**:71–77. doi:https://doi.org/10.1117/12.932660

Qin D, Takamatsu M, Nakashima Y. 2006. Disparity limit for binocular fusion in fovea. *Opt Rev*

13:34–38. doi:10.1007/s10043-006-0034-5

R Core Team. 2022. R: a language and environment for statistical computing.

Reynaud A, Hess RF. 2017. Characterization of spatial frequency channels underlying disparity

sensitivity by factor analysis of population data. *Front Comput Neurosci* **11**:63.

doi:10.3389/fncom.2017.00063

Sayim B, Westheimer G, Herzog MH. 2008. Contrast polarity, chromaticity, and stereoscopic

depth modulate contextual interactions in vernier acuity. *J Vis* **8**:1–9. doi:10.1167/8.8.12

Siman-Tov Z, Lev M, Polat U. 2021. Binocular summation is affected by crowding and tagging.

Sci Rep **11**:4843. doi:10.1038/s41598-021-83510-8

Strasburger H, Rentschler I, Jüttner M. 2011. Peripheral vision and pattern recognition: a review.

J Vis **11**:1–82. doi:10.1167/11.5.13.

Thaler L, Schütz AC, Goodale MA, Gegenfurtner KR. 2013. What is the best fixation target?

The effect of target shape on stability of fixational eye movements. *Vision Res* **76**:31–42.

doi:10.1016/j.visres.2012.10.012

Tyler CW, Barghout L, Kontsevich LL. 1994. Computational reconstruction of the mechanisms

of human stereopsis. *Proc SPIE Comput Vis Based Neurobiol* **2054**:52–68.

Ur Rehman A, Kihara K, Matsumoto A, Ohtsuka S. 2015. Attentive tracking of moving objects

in real 3D space. *Vision Res* **109**:1–10. doi:10.1016/j.visres.2015.02.004

Viswanathan L, Mingolla E. 2002. Dynamics of attention in depth: Evidence from multi-element

tracking. *Perception* **31**:1415–1437. doi:10.1068/p3432

Wallis TSA, Bex PJ. 2012. Image correlates of crowding in natural scenes. *J Vis* **12**:1–19.

doi:10.1167/12.7.6

Watt SJ, Akeley K, Ernst MO, Banks MS. 2005. Focus cues affect perceived depth. *J Vis* **5**:834–

862. doi:10.1167/5.10.7

Westheimer G. 1979. Cooperative neural processes involved in stereoscopic acuity. *Exp Brain*

Res **36**:585–597. doi:10.1007/BF00238525

Whitney D, Levi DM. 2011. Visual Crowding: a fundamental limit on conscious perception and

object recognition. *Trends Cogn Sci* **15**:160–168. doi:10.1016/j.tics.2011.02.005.Visual

Wilcox LM, Allison RS. 2009. Coarse-fine dichotomies in human stereopsis. *Vision Res*

49:2653–2665. doi:10.1016/j.visres.2009.06.004

Xia Y, Manassi M, Nakayama K, Zipser K, Whitney D. 2020. Visual crowding in driving. *J Vis*

20:1–7. doi:10.1167/jov.20.6.1

Yang Y, Blake R. 1991. Spatial frequency tuning of human stereopsis. *Vision Res* **31**:1176–1189.

doi:10.1016/0042-6989(91)90043-5