Large depth differences between target and flankers can increase crowding: Evidence from a multi-depth plane display

Samuel P. Smithers[1*], Yulong Shao[1], James Altham[1], and Peter J. Bex[1*]

[1] Department of Psychology, Northeastern University, Boston, MA, USA

[*] Corresponding authors: SPS (s.smithers@northeastern.edu) and PJB (p.bex@northeastern.edu)

Abstract

Crowding occurs when the presence of nearby features causes highly visible objects to become unrecognizable. Although crowding has implications for many everyday tasks and the tremendous amounts of research reflect its importance, surprisingly little is known about how depth affects crowding. Most available studies show that stereoscopic disparity reduces crowding, indicating that crowding may be relatively unimportant in three-dimensional environments. However, most previous studies tested only small stereoscopic differences in depth in which disparity, defocus blur, and accommodation are inconsistent with the real world. Using a novel multi-depth plane display, this study investigated how large (0.54 - 2.25 diopters), real differences in target-flanker depth, representative of those experienced between many objects in the real world, affect crowding. Our findings show that large differences in target-flanker depth increased crowding in the majority of observers, contrary to previous work showing reduced crowding in the presence of small depth differences. Furthermore, when the target was at fixation depth, crowding was generally more pronounced when the flankers were behind the target as opposed to in front of it. However, when the flankers were at fixation depth, crowding was generally more pronounced when the target was behind the flankers. These

23 findings suggest that crowding from clutter outside the limits of binocular fusion can still have a

24 significant impact on object recognition and visual perception in the peripheral field.

Introduction

Visual crowding is a phenomenon in which a peripherally viewed object, that is easily

identifiable in isolation, becomes harder to recognize or identify when it is surrounded by other

objects, despite still being clearly visible (Bouma, 1970; Herzog et al., 2015; Levi, 2008; Pelli,

2008; Whitney and Levi, 2011). The importance of crowding is reflected by the tremendous

amount of research on the topic and it has been extensively reviewed over the years (e.g., Coates

et al., 2021; Herzog et al., 2015; Manassi and Whitney, 2018; Pelli, 2008; Strasburger et al.,

2011; Whitney and Levi, 2011). This high level of interest is not surprising given that the natural

environment is usually cluttered and mostly viewed in the peripheral visual field. Consequently,

it is assumed that crowding has a large impact on our daily lives with implications for many

everyday visual tasks including driving (Xia et al., 2020), reading (Legge, 2007) and face

recognition (Kalpadakis-Smith et al., 2018; Louie et al., 2007). Crowding also has important

clinical implications for people with glaucoma (Ogata et al., 2019; Shamsi et al., 2022), as well

as for other disorders including central vision impairment (e.g., macular degeneration),

amblyopia and dyslexia (Whitney and Levi, 2011). Through this body of research, we have been

able to build a good understanding of the factors that mediate and moderate crowding and

develop unifying theories (e.g., Balas et al., 2009; Harrison and Bex, 2015; Keshvari and

Rosenholtz, 2016; van den Berg et al., 2012, 2010). However, despite the large body of

literature, one area that remains surprisingly understudied is how depth affects crowding. This is

important because we live in a three-dimensional environment and so to understand the

significance of crowding in the real world it is essential to understand the role of depth

information.

Kooi et al. (1994) were the first to investigate how depth influences crowding. The

authors found that compared to when a peripheral target and flankers were at the same depth,

there was a release from crowding when the target and flankers were stereoscopically presented

in crossed (front) and uncrossed (behind) disparities (Kooi et al., 1994). Sayim et al. (2008)

found that when flankers were at fixation depth, there was less foveal crowding when the target

was stereoscopically presented in front or behind the flankers. Felisberti et al. (2005) also found

an effect of depth on crowding in the parafovea, but to a more limited extent. They reported a

release from crowding in two out of three subjects when the target appeared behind the flankers,

but only in one of the three subjects when the target was in front of the flankers (Felisberti et al.,

2005). Astle et al. (2014) extended on previous research by testing the effect of depth on

crowding over a range of stereoscopically produced target-flanker disparities (±800 arcsec). In

line with previous studies, Astle et al. (2014) found that when the target was at fixation depth,

increasing target-flanker disparity systematically reduced crowding. They also found that the

release from crowding was asymmetric whereby crowding was greater when the flankers were in

front of the target compared to behind it (Astle et al., 2014).

Taken together these studies appear to paint the picture that a difference in depth between

a target and flankers reduces crowding. This could challenge the concept that crowding is

ubiquitous in three-dimensional natural environments (Whitney and Levi, 2011) because it

implies that depth differences between natural objects could prevent crowding from being a

problem (Figure 1). However, all the above studies tested only small depth differences between

target and flankers that do not reflect the often-large differences in depth between objects in the

real world. Furthermore, the generalizability of these studies to real world depth is also limited

by the fact that depth cues were generated stereoscopically. One potential problem with stereo

displays is that disparity information often does not match with accommodation or defocus blur

which can detrimentally affect depth perception (Hoffman et al., 2008; Watt et al., 2005).

Figure 1: Example of crowding in a simulated everyday scene. (A) When fixating on the red dot above

the horizon, it is relatively easy to spot and recognize the person crossing on the right. However, it is

more difficult, or impossible, to recognize the person about to cross the road on the left because of the

presence of nearby signs. Images like this one are often used to illustrate how crowding is ubiquitous in

everyday natural scenes. **(B)** However, in real life the two signs and the person on the left may be several

meters apart and so each would appear at a different depth from the observer. Currently, little is known

about how large, real differences in depth between objects affect crowding. See acknowledgments for

image attributions.

To our knowledge, four studies have investigated the effect of depth on crowding using a

real depth display. All four studies used two orthogonally positioned screens viewed through a

half-transparent mirror allowing stimuli to be displayed on two different depth planes

simultaneously (Eberhardt et al., 2021; Eberhardt and Huckauf, 2019, 2017, 2020). Eberhardt

and Huckauf (2017) kept a target and flankers at the same depth and presented them 0.06

diopters (i.e., 1/focal length in meters), in front, behind or at the same depth as fixation. The

authors found that this small difference in depth between the stimuli and fixation planes had no

effect on the amount of crowding compared to when everything was presented at the fixation

plane (Eberhardt and Huckauf, 2017). Eberhardt and Huckauf (2019) repeated and expanded on

this work with the addition of a $\pm\,0.1$ diopter depth difference between fixation and the stimuli in

addition to the $\pm\,0.06$ diopter condition. The results showed that crowding was greater when the

stimuli were furthest from fixation compared to when they were closer to fixation, regardless of

whether the stimuli were in front or behind the fixation plane (Eberhardt and Huckauf, 2019).

More recently, Eberhardt and Huckauf (2020) used the same approach to investigate the effect of

depth on crowding when the target and flankers were separated in depth. When flanker depth

was varied while the target was at fixation depth, crowding increased with increasing target-

flanker depth difference (Eberhardt and Huckauf, 2020). This result is the opposite to the

findings of other studies in which small differences in depth between the target and flankers were

introduced with stereoscopic disparity (Astle et al., 2014; Felisberti et al., 2005; Kooi et al.,

1994; Sayim et al., 2008). Furthermore, the authors also found that when flanker depth was

varied while the target was at fixation depth, crowding was greater when the flankers were

behind the target and fixation compared to in front of it (Eberhardt and Huckauf, 2020). When

target depth was varied while the flankers were at fixation, there was more crowding when the

target was in front of the flankers and fixation compared to when it was behind them (Eberhardt

and Huckauf, 2020).

In light of the above discrepancies, and the prevalence of depth information in real world

scenes, there is a pressing need for a better understanding of how visual crowding is affected by

depth differences representative of those experienced between objects in the real world.

Therefore, using a novel multi-depth plane display consisting of screens at 0.4 m (near screen),

1.26 m (mid screen), and 4 m (far screen) viewing distances (Figure 2), we investigate how

crowding is affected by large, real differences in target-flanker depth. These viewing distances

were chosen because 4 m is close to optical infinity while 0.4 m is the clinical standard for near

distance, with 1.26 m being in the middle of these two extremes in log units. Given in diopters,

the near, mid, and far screens were at 2.5 diopters, 0.79 diopters and 0.25 diopters receptively.

Figure 2: Illustration of the multi-depth plane display setup used for this study. The near and mid screens

consisted of transparent acrylic covered with ClearBright display film (see Methods for details) that

enabled projected images to be displayed on to the screens while preserving their transparency (Hsu et al.,

2014). This enabled observers to simultaneously view adjacent stimuli presented at different depth planes

at the same time. The screens were temporally synchronized and spatially aligned (see Methods) to ensure

display timing and position were consistent across the three depth planes.

Results

For the main study, naïve observers were tasked with reporting the orientation of a peripherally

viewed target (Figure 3) in a series of five experiments (see Methods for details and Figure 4 for

example trial). The target was surrounded by a flanker ring with a target-flanker spacing of 5˚,

2.5°, 1.25°, 0.625°, or infinity, i.e., no flanker ring (Figure 3). In Experiments 1 and 2 the target

was always presented at fixation depth while the depth of the flanker ring was varied.

Conversely, in Experiments 3 and 4 the flanker ring was presented at fixation depth while target

depth was varied. Finally, for Experiment 5, the target, flanker ring and fixation were presented

together on the same screen at three different depths. To quantify crowding for each stimulus

condition we used the reported target orientation across repeated trials to calculate perceptual

error (see Methods), which is reported in degrees.

Due to the nature of the experimental set up, and the large distances being tested, it is

possible that head movements, diplopia, or local suppression (see Discussion) could have

resulted in differences in the perceived position of the target relative to the flanker ring.

Therefore, in all experiments except Experiment 5, observers were also asked to report the

perceived location of the target relative to the position of the flanker ring (Figure 4). This

enabled us to examine how perceptual error was affected by factors such as target misalignment

and possible diplopia (see Discussion).

Figure 3: Schematic illustration of experimental stimuli and fixation marker. In all conditions, except the

no-flanker condition, the randomly orientated target (a white Landolt-C) was surrounded by a white

flanker ring at one of four possible target-flanker distances. The stimuli were viewed against a black

background.

Figure 4: A step-by-step example of a single trial. Note that in the actual experiments a text description

of each of the five options in step five was also included within the option boxes. The text descriptions for

the different options were: "Target in center of ring" (top left), "Target inside ring but NOT in the center"

(top right), "Ring obstructs target" (middle left), "Target outside ring" (middle right), and "Unsure or no

ring" (bottom center). Step five was not included in Experiment 5 as all parts of the stimulus were always

presented at the same depth.

<u>Perceptual error</u>

<u>Target always at fixation depth with variable flanker depth</u>

Experiment 1: Target and fixation at 1.26 m with flanker ring presented at each depth

(Figure 5A and Figure 5-figure supplement 1A). Perceptual error was higher when the flanker

ring was displayed in front of, or behind, the target and fixation plane (mixed effects multiple

linear regression models compared via LRT for effect of flanker depth: $\chi^2_{(2)} = 20.67$, p <0.001).

As expected, perceptual error decreased as target-flanker spacing increased (effect of target-

flanker spacing: $\chi^2_{(1)} = 31.71$, p <0.001). There was no significant interaction between flanker

depth and target-flanker spacing ($\chi^2_{(2)} = 5.56$, p=0.062).

Experiment 2: Target and fixation at 0.4 m or 4 m with flanker ring presented at each

depth (Figure 5B and Figure 5-figure supplement 1B). Perceptual error tended to increase

with increasing flanker depth when the flanker ring was behind the target and fixation plane,

however, the effect of flanker depth was much weaker when the flanker ring was in front of the

target and fixation (interaction between target-fixation depth and flanker depth: $\chi^2_{(2)} = 17.18$,

p<0.001). Perceptual error decreased with increasing target-flanker spacing, in a manner

corresponding to target-fixation depth (interaction between target-flanker spacing and target-

fixation depth: $\chi^2_{(1)} = 9.2$, p=0.002). There was no significant interaction between target-flanker

spacing and flanker screen and ($\chi^2_{(2)} = 4.52$, p=0.104).

Figure 5: Perceptual error results for Experiments 1 (A) and 2 (B) in which flanker depth was

varied while the target was always at fixation depth. Box plots show medians plus the interquartile

range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper

and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below

each graph shows the depth of the flanker ring (based on the same color key as the box plot) relative to

the depth of the fixation point and target (shown in black) for each stimulus condition. The relative sizes

of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation

point is only shown for the 0.625˚ target-flanker spacing due to limited space within the figure.

<u>Flanker ring always at fixation depth with variable target depth</u>

Experiment 3: Flanker ring and fixation at 1.26 m with target presented at each depth

(Figure 6A and Figure 6-figure supplement 1A). Overall, perceptual error was higher when

the target was displayed in front of, or behind, the flanker ring and fixation (effect of target

depth: $\chi^2_{(2)} = 36.93$, p <0.001), with the greatest perceptual error occurring when the target was

behind the flanker ring and fixation. Perceptual error decreased as target-flanker spacing

increased (effect of target-flanker spacing: $\chi^2_{(1)} = 45.99$, p <0.001). There was no significant

interaction between target depth and target-flanker spacing ($\chi^2_{(2)} = 3.55$, p=0.17).

Experiment 4: Flanker ring and fixation at 0.4 m or 4 m with target presented at each

depth (Figure 6B and Figure 6-figure supplement 1B). Perceptual error was greatest when the

target was displayed behind the flanker ring and fixation, while the effect of depth was less

prominent when the target was displayed in front of the flanker ring and fixation (interaction

between flanker-fixation depth and target depth: $\chi^2_{(2)} = 61.87$, p<0.001). Perceptual error

decreased as target-flanker spacing increased (effect of target-flanker spacing: $\chi^2_{(1)} = 29.92$,

p<0.001). There was no significant interaction between flanker-fixation depth and target-flanker

spacing ($\chi^2_{(1)} = 0.14$, p=0.71) nor between target depth and target-flanker spacing ($\chi^2_{(2)} = 1.43$,

p=0.488).

Figure 6: **Perceptual error results for Experiments 3 (A) and 4 (B) in which target depth was varied while the flanker ring was always at fixation depth.** Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below each graph shows the depth of the target (based on the same color key as the box plot) relative to the depth of the fixation point and flanker ring (shown in black) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker spacing due to limited space within the figure.

Target and flanker ring at fixation depth

Experiment 5: Target and flanker ring always at fixation depth (Figure 7). Perceptual error decreased as target-flanker spacing increased (effect of target-flanker spacing: $\chi^2_{(1)} = 64.8$, p<0.001). There was also a significant effect of target-flanker-fixation depth ($\chi^2_{(2)} = 8.75$, p=0.013) resulting from a higher perceptual error for stimuli displayed at 4 m. There was no significant interaction between target-flanker-fixation depth and target-flanker spacing ($\chi^2_{(2)} = 1.72$, p=0.424).

Figure 7: **Perceptual error results for Experiment 5 in which the target and flanker were always presented at the same depth as the fixation point.** Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below the graph shows the depth of the target, flanker, and fixation point (based on the same color key as the box plot) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker spacing.

Perceived target position relative to the flanker ring

Target always at fixation depth with variable flanker depth

Experiment 1: Target and fixation at 1.26 m with flanker ring presented at each depth (Figure 8A and Figure 8-figure supplement 1). In Experiment 1, most observers consistently reported seeing the target inside the flanker ring. However, when the flanker ring was away from target-fixation depth there was a reduction in the proportion of observers who consistently reported seeing the target within the flanker ring when the target-flanker spacing was small (mixed effects binary logistic regression models compared via LRT; interaction between target-flanker spacing and flanker depth: $\chi^2_{(2)} = 36.12$, p <0.001). This effect was greatest when the flanker ring was displayed behind the target and fixation.

Note that in all four experiments, the response to the control (i.e., no flanker condition)

indicates the probability of an observer making an inaccurate report (e.g., they selected the

wrong option by mistake or missed the stimulus and reported a guess rather than selecting the

"Unsure" option).

Experiment 2: Target and fixation at 0.4 m or 4 m with flanker ring presented at each

depth (Figure 8B and Figure 8-figure supplement 2). As with the previous experiments, when

the flanker ring was away from target-fixation depth, there was a reduction in the overall

proportion of trials in which observers reported seeing the target within the flanker ring

(interaction between target-fixation depth and flanker depth: $\chi^2_{(2)} = 374.26$, p < 0.001). The

greatest reduction occurred when target-flanker spacing was smaller (effect of target-flanker

spacing: $\chi^2_{(1)} = 178.65$, p < 0.001). There was no significant interaction between flanker depth

and target-flanker spacing ($\chi^2_{(2)} = 0.24$, p=0.886) nor between target-fixation depth and target-

flanker spacing ($\chi^2_{(1)} = 1.64$, p=0.2)

Figure 8: **Proportion of trials in which observers reported seeing the target inside the flanker ring**

for Experiments 1 (A) and 2 (B). In both experiments flanker depth was varied while the target was

always at fixation depth**.** Box plots show medians plus the interquartile range (IQR), whiskers are the

lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points

show the proportion of trials for each individual observer. The visual aid below each graph shows the

depth of the flanker ring (based on the same color key as the box plot) relative to the depth of the fixation

point and target (shown in black) for each stimulus condition. The relative sizes of, and distances

between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown

for the 0.625˚ target-flanker spacing due to limited space within the figure.

<u>Flanker ring always at fixation depth with variable target depth</u>

Experiment 3: Flanker ring and fixation at 1.26 m with target presented at each depth

(Figure 9A and Figure 9-figure supplement 1). When the target was away from flanker-

fixation depth, observers were less likely to report seeing the target within the flanker ring. This

effect was greatest when target-flanker spacing was smaller and when the target was displayed in

front of the flanker ring and fixation (interaction between target-flanker spacing and target depth:

$\chi^2_{(2)} = 100.38$, p <0.001).

Experiment 4: Flanker ring and fixation at 0.4 m or 4 m with target presented at each

depth (Figure 9B and Figure 9-figure supplement 2). As with the previous experiment, the

greater the difference between target depth and flanker-fixation depth, the less likely observers

were to report seeing the target within the flanker ring (interaction between flanker-fixation

depth and target depth: $\chi^2_{(2)} = 779.49$, p < 0.001), particularly when target-flanker spacing was

small (interaction between target-flanker spacing and target depth: $\chi^2_{(2)} = 23.27$, p < 0.001).

However, target-flanker spacing had a weaker effect on the proportion of trials in which

observers reported seeing the target within the flanker ring when the flanker ring and fixation

were near compared to far (interaction between target-flanker spacing and flanker-fixation depth:

$\chi^2_{(1)} = 56.23$, p < 0.001).

Figure 9: **Proportion of trials in which observers reported seeing the target inside the flanker ring**

for Experiments 3 (A) and 4 (B). In both experiments, target depth was varied while the flanker ring was

always at fixation depth**.** Box plots show medians plus the interquartile range (IQR), whiskers are the

lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points

show the proportion of trials for each individual observer. The visual aid below each graph shows the

depth of the target (based on the same color key as the box plot) relative to the depth of the fixation point

and flanker ring (shown in black) for each stimulus condition. The relative sizes of, and distances

between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown

for the 0.625˚ target-flanker spacing due to limited space within the figure.

Individual differences in perceptual error and perceived target position in experienced

observers

While we were able to detect overall statistically significant trends in the results, it is apparent

from Figures 5-9 that there is a large amount of variation within the data. This raises questions

about whether the variation is because the effects of depth are highly variable between some

observers, or because the naïve observers were not very stable at indicating their perception. The

latter could be due to the naïve observers being relatively inexperienced at the task, despite

receiving training since each observer only participated in a single experiment. Therefore, to

investigate the effects of depth at the individual level we repeated the study with a small number

of highly trained, experienced observers who performed all five experiments (see Methods for

details).

<u>Target always at fixation depth with variable flanker depth</u>

The small sample size means direct comparisons between these results and the results reported

above for the main study should be avoided. Nevertheless, the overall effects reported above are

also apparent in most of the experienced observers (Figure 10). However, Figure 10 shows that

even among highly trained, experienced observers there is a large amount of variation both

between and within subjects. For instance, when the target and fixation are on the near screen

(0.4 m) and target-flanker spacing is 0.625˚, ES3 and ES4 both exhibit a similar increase in

perceptual error with increasing flanker depth, which is consistent with the overall effect shown

in figure 5B. In contrast, ES2 shows the opposite trend for the same stimulus condition, a

reduction in perceptual error with increased flanker depth. However, when the target and fixation

were on the mid screen (1.26 m) and target-flanker spacing is 0.625˚, ES2 does exhibit an

increase in perceptual error when the flanker ring is behind the target and fixation (as do ES1 and

ES3), thus providing an example of between and within subject variation. The variation in

perceptual error between subjects cannot solely be explained by differences in the perceived

position of the target relative to the flanker ring (Figure 10-figure supplements 1 and 2). For

example, when the target and fixation were on the near screen and target-flanker spacing was

0.625˚, the perceived target positions reported by ES2 and ES4 were remarkably similar (Figure

10-figure supplement 2) despite exhibiting the previously mentioned opposite effects of depth on

perceptual error (figure 10).

Figure 10: **Perceptual error results from individual experienced subjects (ES1-4) for Experiments 1**

and 2 in which flanker depth was varied while the target was always at fixation depth. Error bars

show 95% bootstrapped confidence intervals. Note that the 5° target-flanker spacing condition was not

included in this set of experiments. Since the same subjects participated in both experiments, the results

from Experiments 1 and 2 have been combined into a single figure to make it easier to compare the

different conditions. The visual aid below the graph shows the depth of the flanker ring (based on the

same color key as the plot) relative to the depth of the fixation point and target (shown in black) for each

stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring

are to scale. Note that the fixation point is only shown for the 0.625° target-flanker spacing due to limited

space within the figure.

<u>Flanker ring always at fixation depth with variable target depth</u>

Like the results from the repeat of Experiment 1 and 2 with the experienced subjects, Figure 11

shows there was also a relatively large amount of between and within subject variation in

Experiments 3 and 4 when repeated with the experienced subjects. For instance, if we look at the

analogous stimulus condition to the one used in the previous example, i.e., flanker ring and

fixation on the near screen (0.4 m) with a target-flanker spacing of 0.625˚, ES1, ES3 and ES4 all

exhibit an increase in perceptual error with increasing target depth, which is consistent with the

overall effect shown in figure 6B. In contrast, ES2 again shows the opposite effect for the same

stimulus condition, a reduction in perceptual error with increasing target depth. Again, the

variation in perceptual error between subjects cannot solely be explained by differences in the

perceived position of the target relative to the flanker ring (Figure 11-figure supplements 1 and

2). For instance, the perceived target positions reported by ES2 and ES3 were very similar

(Figure 11-figure supplement 2) when the flanker ring and fixation were on the near screen and

target-flanker spacing was 0.625˚, yet the two observers exhibited the opposite effect of

increasing target depth on perceptual error (Figure 11).

Figure 11: **Perceptual error results from individual experienced subjects (ES1-4) for Experiments 3 and 4 in which target depth was varied while the flanker ring was always at fixation depth.** Error bars show 95% bootstrapped confidence intervals. Note that the 5˚ target-flanker spacing condition was not included in this set of experiments. Since the same subjects participated in both experiments, the results from Experiments 3 and 4 have been combined into a single figure to make it easier to compare the different conditions. The visual aid below each graph shows the depth of the target (based on the same color key as the plot) relative to the depth of the fixation point and flanker ring (shown in black) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker spacing due to limited space within the figure.

Target and flanker ring at fixation depth

378 Figure 12 shows there was also some variation between the experienced subjects in Experiment 5

379 with ES2 and ES3 both exhibiting the highest perceptual error when the target, flanker ring and

380 fixation were furthest away (though this is less pronounced at greater eccentricities for ES3).

381 ES1 and ES4 on the other hand both show a less consistent effect of depth.

Figure 12: **Perceptual error results from individual experienced subjects (ES1-4) for Experiment 5**

in which the target and flanker were always presented at the same depth as the fixation point. Error

bars show 95% bootstrapped confidence intervals. Note that the 5˚ target-flanker spacing condition was

not included in this set of experiments. The visual aid below the graph shows the depth of the target,

flanker, and fixation point (based on the same color key as the plot) for each stimulus condition. The

relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that

the fixation point is only shown for the 0.625˚ target-flanker spacing.

Discussion

In all experiments, perceptual error generally decreased as the target-flanker spacing increased, which is in line with previous crowding research (Bouma, 1970; Pelli, 2008). Overall, perceptual error was higher when the target or flanker ring were presented at a different depth from the other and from fixation. The greatest differences in perceptual error among the different depth conditions occurred when the target or flanker ring were presented behind fixation depth. In contrast, when the target or flanker ring were presented in front of fixation depth, the differences in perceptual error among depth conditions were notably smaller. The results also show that when the target, flanker ring and fixation were all at the same depth plane, crowding tended to increase with the absolute distance of the stimuli from the observer. However, in all the main experiments with naïve observers there was a large amount of variation within the data. Follow up experiments with a small sample of highly trained, experienced observers were largely consistent with the main effects observed in the main study. Moreover, the experienced observers also exhibited a high degree of between subject variation, which supports the suggestion that the variability in the main data from the large sample of naïve observers is representative of the natural variation between different individuals. Therefore, although significant effects were found at the population level, the effects of large differences in target-flanker depth can be highly variable at the individual level. This demonstrates the importance of using a large sample size to investigate effects that are not necessarily detectable at the individual level. The discussion of the findings will focus on the overall results from the main study on naïve participants, with the findings from the small sample of experienced observers used to provide additional context and insight.

The main finding that a depth difference between the flankers and the target led to an

increase, rather than a decrease, in perceptual error is different from the findings of several

previous studies that investigated the effect of depth on crowding using a stereoscopic

presentation (Astle et al., 2014; Felisberti et al., 2005; Kooi et al., 1994; Sayim et al., 2008). For

example, Astle et al (2014) found that when the target was presented at fixation depth

(comparable to Experiments 1 and 2 in our study), increasing the disparity between the target and

flankers resulted in less crowding, and ultimately a complete release from crowding. The authors

also reported that crowding was greater when flankers were presented in crossed disparity (in

front of the target and fixation), compared to uncrossed disparity (behind the target and fixation)

(Astle et al., 2014). Similarly, when the flankers were presented at fixation depth (comparable to

Experiments 3 and 4 in our study), Sayim et al (2008) found that foveal crowding was lower

when the target was presented in front of or behind the flanker and fixation.

Several explanations for crowding have been proposed in which crowding occurs because

target and flanking features are inappropriately grouped (Banks et al., 1979; Herzog and

Manassi, 2015; Pachai et al., 2016). Consistent with this explanation, crowding can be alleviated

when the flankers are grouped together and segmented from the target (Herzog and Manassi,

2015). One piece of evidence in support of this account is the reduction of crowding that can

occur when flankers are presented at a different depth from the target (Astle et al., 2014;

Felisberti et al., 2005; Kooi et al., 1994; Sayim et al., 2008). Our results challenge this view of

crowding because it would predict that increasing the depth difference between target and

flankers would decrease the probability of them being inappropriately grouped and increase the

probability of the flankers being segmented from the target, thereby decreasing perceptual error,

the opposite of our finding. Other models of crowding, such as texture processing and feature

integration (Balas et al., 2009; Freeman and Simoncelli, 2011; Parkes et al., 2001; van den Berg

et al., 2012, 2010; Wallis and Bex, 2012), sampling errors (Ester et al., 2015, 2014; Harrison and

Bex, 2017, 2016, 2015), attentional resolution limits (He et al., 1996) or saccade properties

(Nandy and Tjan, 2012) are agnostic concerning the influence of depth differences among targets

and flankers. Therefore, each of these models would require modification to explain the increase

in crowding that we observe with large real depth differences between a target and its flankers.

In contrast with studies that controlled depth with small stereoscopic disparities, our

results do agree with the findings of Eberhardt and Huckauf (2020) who also used a real-depth

display and tested larger differences in depth between the target and flankers than previous

studies had examined, though depth differences were still much smaller than those tested in our

current study. When the target was at fixation depth (comparable to Experiments 1 and 2 in our

study), Eberhardt and Huckauf (2020) found that crowding increased with increasing target-

flanker depth difference. The authors also found that crowding was greater when the flankers

were behind the target and fixation compared to in front of it (Eberhardt and Huckauf, 2020).

Our findings from Experiments 1 and 2 are consistent with these results. However, when the

flankers were at fixation depth (comparable to Experiments 3 and 4 in our study), Eberhardt and

Huckauf (2020) found that crowding was higher when the target was in front of the flankers and

fixation compared to when it was behind them. In contrast, in Experiments 3 and 4 we found that

perceptual error was greatest when the target was presented behind the flanker ring and fixation

rather than in front of it. This difference is likely due to a combination of high defocus blur when

the target was behind fixation combined with the finding from Experiment 5 which found that in

our study a larger absolute distance from the observer resulted in higher perceptual error.

Why the inconsistencies across studies? It is important to consider that most of the

previous studies used very small depth differences, the largest being 0.1 diopters in Eberhardt

and Huckauf (2019, 2020). In comparison, the depth difference in our study ranged from 0.54

diopters (between the mid and far screens) to 2.25 diopters (between the near and far screens).

Our results, together with those from previous studies, suggest that small differences in depth

between the target and flankers reduce crowding (Astle et al., 2014; Felisberti et al., 2005; Kooi

et al., 1994; Sayim et al., 2008), while larger differences in depth can increase it (Eberhardt and

Huckauf, 2020; and results from our study). Although this pattern is quite surprising, there is

precedent for non-linear effects of depth on perceptual processing outside the crowding

literature. In a multiple object tracking task, when the targets and distractors were distributed

equally on two depth planes, tracking accuracy was better than the accuracy during trials in

which targets and distractors appeared on a single depth plane when the depth difference

between planes was small (~6 cm) (Ur Rehman et al., 2015; Viswanathan and Mingolla, 2002),

but worse when the depth difference between planes was large (50 cm) (Ur Rehman et al., 2015).

How could depth affect perception in this nonlinear way? The geometry of the eyes and

the 3D environment create interocular disparity differences between the images in each eye.

When corresponding features in the images in each eye can be identified, the images can be

fused to generate a binocular perceptual experience of depth (for review see Nityananda and

Read, 2017). There is a horizontal and vertical spatial limit to the interocular disparity difference

over which the visual system attempts to identify corresponding retinal points in order to fuse

different 2D interocular images as a single 3D object (Westheimer, 1979). This 2D spatial

correspondence limit creates a 3D volume along the horopter known as Panum's Fusional Area

(Panum, 1858). Outside Panum's Fusional Area, there is a loss of stereoscopic depth perception

and either double vision (diplopia) may be experienced, or single vision may occur if the image

from one eye is partly suppressed (Georgeson and Wallis, 2014; Ono et al., 1977; Spiegel et al.,

2016; Stidwill and Fletcher, 2010). The spatial extent of Panum's Fusional Area depends on the

spatial frequency of the stimuli (Maiello et al., 2020; Pulliam, 1981; Reynaud and Hess, 2017;

Wilcox and Allison, 2009; Yang and Blake, 1991) and increases with eccentricity (Hampton and

Kertesz, 1983; Harrold and Grove, 2019; Qin et al., 2006).

It is possible that differences between fusion and depth perception within Panum's

Fusional Area versus diplopia, and/or local suppression for single vision, outside of it can

explain the patterns of results observed between studies. Small disparity differences between

target and flankers create differences in perceived depth that can alleviate the effects of

crowding, potentially through ungrouping of target and flanking elements (Herzog et al., 2015;

Herzog and Manassi, 2015; Manassi et al., 2012) or by independent processing of target and

flankers in different depth-selective channels (Norcia et al., 1985; Pulliam, 1981; Reynaud and

Hess, 2017; Tyler et al., 1994; Wilcox and Allison, 2009; Yang and Blake, 1991) by any of the

computational models of crowding based on texture processing and feature integration (Balas et

al., 2009; Freeman and Simoncelli, 2011; Parkes et al., 2001; van den Berg et al., 2012, 2010;

Wallis and Bex, 2012), sampling errors (Ester et al., 2015, 2014; Harrison and Bex, 2017, 2016,

2015) attentional resolution limits (He et al., 1996) or saccade properties (Nandy and Tjan,

2012). Large interocular disparity differences on the other hand can induce diplopia that could

increase the perceived number of flankers (Eberhardt and Huckauf, 2019). In theory, this

perceived increase in the number of flanking elements could result in higher perceptual error

because increasing the number of flankers is known to increase crowding under certain

conditions (Chanceaux et al., 2014; Manassi et al., 2012; Pluháček et al., 2021). It is however

worth noting that the opposite can occur, i.e. reduced crowding with increasing number of

flankers, in cases where the additional flankers are grouped together (Manassi et al., 2013, 2012).

Surprisingly, when asked, only a few of the naïve observers reported experiencing diplopia, and

those that did reported only experiencing it in a minority of presentations for which they selected

the 'unsure or no ring' option (see Methods) when reporting the perceived location of the target

(Figure 8-figure supplement 1E and 2E, and Figure 9-figure supplement 1E and 2E). However,

among the four experienced observers, three verbally reported experiencing diplopia during

some of the conditions for which they either selected the 'unsure or no ring' option (Figure 10-

figure supplement 2E and Figure 9-figure supplement 2E), or in cases when the diplopia was

barely visible, they chose the option that was the closest match to the clearest stimulus image.

The fact that the experienced observers were more likely to report experiencing diplopia,

possibly because they were better at recognizing it, raises the possibility that diplopia may have

been more common among the naïve observers than was indicated by asking them. However, it

is worth noting that one of the four experienced observers reported that they never experienced

diplopia in any of the experiments, which is more consistent with the majority of naïve

observers. It is possible that the short presentation time of the stimulus was too fast for most

observers to notice diplopia consciously (i.e., diplopia did not enter awareness), or that diplopia

was suppressed entirely by local suppression to maintain single vision. Attention effects such as

inattentional blindness (Bredemeier and Simons, 2012; Simons and Chabris, 1999) could also

play a role in explaining why so few observers reported diplopia in the main experiments. There

is conflicting evidence as to whether conscious awareness of a flanker is necessary for it to

induce crowding (compare Ho and Cheung, 2011 vs Wallis and Bex, 2011), and thus whether

'unconscious diplopia' could have increased crowding in this study due to the reason highlighted

above is a matter for debate and beyond the scope of our findings. In addition, it is worth noting

that due to the lack of binocular fusion of stimulus areas outside Panum's Fusional Area, we

can't rule out the possible involvement of binocular rivalry which has been shown to interfere

with object recognition not just in isolation but also when combined with crowding (Kim et al.,

2013).

Another explanation for higher perceptual error with increasing target-flanker depth

differences could be perceived target misalignment. As explained in the Methods, at the start of

Experiments 1-4 each observer performed a subjective spatial calibration to ensure that the

angular positions of the stimuli and fixation point relative to primary gaze were consistent across

all three depth planes. Although the main experiments were performed under binocular viewing

conditions, this spatial calibration had to be performed monocularly using the observer's

dominant eye to avoid confusion caused by diplopia or rivalry (see Methods). This means that

the screens were imperfectly aligned for the non-dominant eye. Therefore, as in the real world, at

certain flanker sizes and separations there may be occlusions between target and flanker features

that could affect target report accuracy, which is not the same mechanism as crowding. In our

study, such misalignment could result from local suppression of the calibrated eye to maintain

single vision, and/or head movements following the spatial calibration. To examine how target

misalignment affected the overall results, we recalculated perceptual error for the main

experiments with naïve observers using only trials in which the observer reported seeing the

target inside the flanker ring. We then reanalyzed these data the same way as for perceptual error

calculated from the full data set (see Methods). Apart from Experiment 2, the effect of flanker

and/or target depth remained statistically significant (see Figure 5-figure supplement 1 and

Figure 6-figure supplement 1 for plots and analysis results). Therefore, across the majority of

experiments, misalignment or occlusion of the target by the flanker ring is not enough to explain

the overall results. This supports the suggestion that the increase in perceptual error when the

target and flanker are at different depths is being driven by an increase in crowding.

This current study is the first to investigate how crowding is affected by large, real

differences in depth between a target and flanking features. This is important because objects in

the real world are often flanked by visual features that vary greatly in their distance from an

observer, and many will fall outside the limits of Panum's Fusional Area (Figure 1). A limitation

of our approach is that the use of relatively simple stimuli in a tightly controlled experiment

limits the generalizability of our findings to the real world where clutter is abundant and more

complex. Future research should therefore aim to utilize stimuli that better capture the

complexity and heterogeneity of natural scenes. Future work should also investigate if the

reported effects of depth on crowding in our study hold at different target eccentricities,

particularly given that the range of Panum's Fusional Area changes with eccentricity (Hampton

and Kertesz, 1983; Harrold and Grove, 2019; Qin et al., 2006). Nevertheless, our findings still

show that crowding from clutter outside Panum's Fusional Area has the potential to significantly

impact object recognition and visual perception in the peripheral field. Our findings therefore do

not support the view that depth differences between targets and flankers necessarily reduce

crowding, instead the effect of depth on crowding appears to be more complex than originally

thought. Furthermore, data collected from experienced participants shows that the overall trends

at the population level may not always be apparent at the individual level. Nonetheless, our

results, alongside those of previous studies, suggest that small differences in depth between

targets and flankers do not have the same effect on crowding as larger differences in depth in

which either the target or flankers are presented outside the limits of binocular fusion. This

finding confirms the potential importance and relevance of crowding in three-dimensional

scenes.

Methods

Subject recruitment

For the main study, a total of 110 naïve subjects were recruited from the Northeastern University

undergraduate population in exchange for course credit as compensation for their time. Of these,

seven were excluded immediately following screening because they did not meet the inclusion

criteria of normal or corrected-to-normal vision. An additional five subjects were excluded

because of issues with the experimental code (N=3), calibration issues (N=1) or because they did

not follow the task instructions (N=1). Thus, a total of 98 subjects (32 males, 66 females; mean

age = 18.86 years) contributed usable data across five separate experiments (22 in Experiment 1,

19 in Experiment 2, 21 in Experiments 3 and 4, and 15 in Experiment 5).

In addition to the naïve subjects who each participated in one of the five experiments as

part of the main study, we also separately tested four "experienced subjects" (2 males, 2 females;

mean age = 29 years) who participated in all five experiments in a random order. All of the

experienced subjects (none of whom were the authors) were volunteers from within the

university vision research community and were therefore extremely familiar and well-practiced

in participating in psychophysics experiments such as those in our current study. The purpose of

repeating the study with a small group of highly trained, experienced subjects who participated

in all five experiments was to enable us to better explore the effects of depth at the individual

level.

All subjects read and signed an informed consent form approved by the Northeastern

University Ethics Board before the experiment began. The experimental procedure was approved

by the institutional review board at Northeastern University, and the experiments were

performed in accordance with the tenets of the Declaration of Helsinki.

Multi-depth plane display setup

The study utilized a multi-depth plane display consisting of three screens at 0.4 m (near screen),

1.26 m (mid screen) and 4 m (far screen) from the observer (Figure 2). For the far screen we

used an ultra-short throw projector (Bomaker Polaris 4K Laser Projector) projected onto a 240

cm x 250 cm white wall at the end of the room at a throw distance (measured horizontally from

the projector to the wall) of approx. 50 cm. The mid and near screens were made from 101.5 cm

x 76 cm and 46 cm x 36.5 cm clear 5 mm acrylic respectively held in position by a frame

constructed from extruded aluminum rails. The transparent acrylic was covered on one side with

ClearBright transparent display film (Lux Labs Inc., Boston, MA, USA). The nanoparticle film

enabled us to display images from two projectors (InFocus IN3116) onto the near and mid

screens while preserving their transparency (Hsu et al., 2014). This design enables observers to

simultaneously view adjacent stimuli presented on all three depth planes at the same time. The

throw distances of the near and mid screen projectors (measured from the projector to the bottom

of the screen) were approx. 72 cm and approx. 150 cm respectively. Both projectors were

positioned and angled so that projected light that passed through the transparent screens did not

fall upon the observer or any areas visible by the observer, such as an adjacent screen. Key

stoning was used to digitally correct distortions in the image caused by the projection angle of

the near and mid screen projectors. The screens were temporally synchronized and spatially

aligned to ensure display timing and position were consistent across the three depth planes. All

experiments were conducted in Python using PsychoPy (v2021.2.3 for experiments with the

naïve observers and v2022.2.5 for the experiments with the four experienced observers) (Peirce

et al., 2019).

Spatial calibration

Each observer performed a subjective spatial calibration at the start of Experiments 1-4 to ensure

that the angular positions of the stimuli and fixation point relative to primary gaze were the same

(or as close as realistically possible) across all three depth planes. The center of the far screen

(which was at primary gaze when the observers head was positioned in the chin and head rest)

was used as the fixed reference against which all three screens were calibrated. With their head

in the chin and head rest, observers were instructed to sequentially position a series of targets so

that they were in the same position as a fixed reference target that was displayed on the reference

screen. The calibration was performed under monocular conditions using the observer's

dominant eye as determined by the Miles test (Miles, 1929). The calibration was performed

monocularly because during a preliminary experiment subjects reported finding the task too

difficult to complete under binocular conditions (e.g., because of diplopia, an issue that is

considered further in the Discussion section). For Experiment 5, the spatial calibration was

performed by the experimenter prior to the experiment using a webcam positioned at the

observer's cyclopean point. The main experiment was conducted under binocular viewing

conditions and observers were instructed to maintain the same head position within the chin and

head rest throughout the experiment.

Stimuli and experimental procedure

The target and flankers were inspired by those used in previous studies (Harrison and Bex, 2017,

2015). The target was a white Landolt-C displayed against a black background at 10˚ eccentricity

to the right or left of the fixation point at primary gaze. Following the structure of a Landolt-C

optotype, the target was 1˚ in diameter with a 0.25˚ line width and a 0.25˚ opening. The opening

orientation of the target was random each trial, drawn from a uniform 360˚ distribution. The

target was surrounded by a white flanker ring formed of four circular arcs (Figure 3). In each

depth combination described below, the outer edge of the target was separated from the inner

edge of the flanker ring by 5˚, 2.5˚, 1.25˚ or 0.625˚. These distances of separation represented

1/2, 1/4, 1/8 and 1/16 of the target eccentricity respectively. Note that for the experiments

conducted with the four experienced subjects, the 5˚ target-flanker eccentricity condition was not

included. The line width and size of the four gaps in the flanker ring were kept constant (0.25˚).

For each target depth, a control condition with no flanker ring was also included. The fixation

point consisted of a 1˚ white circle with a black cross (Figure 3). This design was used because it

has been reported to elicit more stable fixation compared to a plain circle or cross hair alone

(Thaler et al., 2013). The background on all screens was always set to the lowest RGB (i.e.

R=G=B=0) value which was transparent on the near and mid screens, and black on the far

screen. Since the mid screen was dimmest (R=G=B=255, white=13.56 cd/m^2), the luminance of

stimuli displayed on the near and far screens were calibrated to match the luminance of the mid

screen using a Photo Research PR-650 spectroradiometer (Photo Research, Chatsworth, CA,

USA). The room lights were switched off for the duration of the experiment.

A summary of the stimulus conditions tested in each experiment is provided in Table 1.

In Experiment 1 the target and fixation point were always on the mid screen while the flanker

ring was displayed in front, behind or at fixation depth. In Experiment 2 the target and fixation

point were presented on the near or far screen while the depth of the flanker ring was varied.

Experiments 3 and 4 were the same as Experiments 1 and 2 respectively except that the flanker

ring was presented at fixation depth while the target depth was varied. For Experiment 5, the

target, flanker ring and fixation were presented together on the same screen at three different

depths. This was to investigate whether there was an effect of absolute viewing distance on

crowding. Each condition was repeated 10 times in Experiments 1, 2, 3 and 5 and eight times in

Experiment 4, with equal number of repeats on either side of the fixation point. The order of

testing was randomized.

Table 1: Summary of the stimulus conditions in each experiment. The same color scheme is used to represent the near (0.4 m), mid (1.26 m) and far (4 m) screens throughout the manuscript. Note that for target-flanker spacing, 'infinity' = no flanker ring. For the experiments conducted with the four experienced subjects, the 5° target-flanker eccentricity condition was not included.

Experiment	Fixation depth (m)	Target depth (m)	Flanker depth (m)	Target-flanker spacing (degrees)	No. of stimulus conditions	Total no. presentations	Sample size
Exp 1	Mid (1.26)	Mid (1.26)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°	13	130 (10 reps per stimulus)	N = 22
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Mid (1.26)	Mid (1.26)	Far (4)	0.625°, 1.25°, 2.5°, 5°			
Exp 2	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	26	260 (10 reps per stimulus)	N = 19
	Near (0.4)	Near (0.4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°			
	Near (0.4)	Near (0.4)	Far (4)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 3	Mid (1.26)	Near (0.4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity	15	150 (10 reps per stimulus)	N = 21
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Mid (1.26)	Far (4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 4	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	30	240 (8 reps per stimulus)	N = 21
	Near (0.4)	Mid (1.26)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Near (0.4)	Far (4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Near (0.4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Mid (1.26)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 5	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	15	150 (10 reps per stimulus)	N = 15
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			

An example trial is illustrated in Figure 4. Briefly, at the start of each trial, observers were instructed to focus on the central fixation point. The fixation point appeared on the fixation screen 1 sec before stimulus onset in Experiments 1, 3 and 5, and 1.5 sec before stimulus onset in Experiments 2 and 4. The fixation point was displayed for slightly longer before stimulus onset in Experiments 2 and 4 to ensure that observers had adequate time to refocus on the fixation point when it switched between the near and far screens. The target and flanker ring (if present)

were displayed for 0.25 sec. The fixation point remained on the screen for 1 sec before being

replaced by a randomly oriented Landolt-C that was the same size as the target Landolt-C. The

observer used the mouse to rotate this Landolt-C until it matched the perceived orientation of the

target and then clicked to report the orientation.

In all experiments except Experiment 5, observers were also asked to report the perceived

location of the target relative to the position of the flanker ring. There were 5 options to choose

from, "target in center of ring", "target inside ring but not in the center", "ring obstructs target",

"target outside ring", and "unsure or no ring". Observers were instructed to select the option that

most closely matched how they perceived the stimulus. If an observer missed the stimulus or

they were unsure of the targets' location relative to the flanker ring, e.g., because of confusion

caused by diplopia, they were instructed to select the "unsure or no ring" option. Collecting data

on the perceived target-flanker location enabled us to examine how differences in target-flanker

depth influenced the perception of the stimulus and investigate the potential effect that factors

such as target misalignment and possible diplopia may have had on perceptual error (see

Discussion). Following this selection, the next trial started immediately.

Before starting an experiment, all participants underwent training during which they

completed as many practice trials as they needed to familiarize themselves with the task. The

main experiment only began once the researcher running the study was confident that the

participant understood the task and was able to perform it correctly. These practice trials did not

contribute to the collected data.

Data processing and statistical analysis

<u>Perceptual error</u>

To quantify crowding for each stimulus condition we first calculated the report error for each

presentation, i.e., the angular difference between the reported gap orientation of the target and

the actual gap orientation. For each subject, this generated 10 report errors per stimulus condition

in Experiments 1, 2, 3 and 5, and eight per stimulus condition in Experiment 4. We then used

these report errors to calculate the circular standard deviation for each stimulus condition using

the Circular Statistics Toolbox (Berens, 2009) in MATLAB R2022a (MathWorks Inc., Natick,

MA, USA). We refer to this measure as perceptual error, which is reported in degrees.

For the main study with naïve observers, the effects of target/flanker depth and target-

flanker spacing on perceptual error were analyzed using mixed effects multiple linear regression,

fitted using the lme4 package (Bates et al., 2015) in R (R Core Team, 2022). Perceptual error

was used as the response (dependent) variable after being natural log transformed to correct for

positive skew. For all five experiments, we included target-flanker spacing as a fixed effect. We

also included flanker depth for Experiments 1 and 2, target depth for Experiments 3 and 4 or

target-flanker-fixation depth for Experiment 5. Target-fixation depth or flanker-fixation depth

were also included as a third fixed effect for Experiments 2 and 4 respectively. All possible

pairwise interactions of fixed effects were fitted in the full model. Subject identification (ID) was

included as a random effect in all experiments to control for repeated measures. We first used

model simplification, whereby models were compared with one another using a likelihood ratio

test (LRT), to remove non-significant interactions sequentially. This enabled us to test the effect

of individual fixed effects that were not involved in an interaction. We then used the minimum

model (i.e., model containing all fixed effects of interest plus any significant two-way

interactions) to report the significance of each main effect or significant interaction(s) by using a

LRT to compare the minimum model with the same model but with the effect/interaction of

interest removed. Individual main effects that were part of a significant interaction were not

tested in this way because their effect was dependent on another variable. Note that data for the

control condition (i.e., no flanker ring) were excluded from statistical analysis. This was to

ensure that a) the fixed effects remained independent and b) the effect of having no flanker ring

did not mask the effect of flanker and target depth.

Perceived target position relative to flanker ring

We used the observers' perceived target position relative to the flanker ring to calculate the

proportion of trials in which observers reported seeing the target inside the flanker ring (i.e., they

chose either "target in center of ring" or "target inside ring but not in the center"). For statistical

analysis, these data were treated as a binary response variable whereby trials in which observers

reported seeing the target inside the flanker ring were assigned a value of 1 while all other trials

(i.e., those for which the observer chose "ring obstructs target", "target outside ring" or "unsure

or no ring") were assigned a value of 0.

For the main study with naïve observers, these data were then analyzed using mixed

effects binary logistic regression, fitted using the lme4 package (Bates et al., 2015) in R (R Core

Team, 2022). We used the DHARMa package (Hartig, 2022) to run residual diagnostics that are

often overlooked in logistic regression. Perceived target position was used as the binary response

variable (i.e., 1 = target inside ring and 0 = target not inside ring). The fixed effect, and pairwise

interaction for each experiment were the same as those described above for the analysis of

perceptual error. Subject ID was included as a random effect in all experiments to control for

repeated measures. Model simplification (to remove non-significant interactions) and

significance testing was the same as described above for the analysis of perceptual error. Data for

the control condition (i.e., no flanker ring) were excluded from statistical analysis for the same

reasons as those outlined above. The separate results showing the proportion of trials for which

observers chose each of the five choices for perceived target position relative to the flanker ring

can be found in the figure supplements (see Results).

Acknowledgements

We thank members of the Bex lab and Eskew lab at Northeastern University for valuable

feedback during pilot experiments. We also thank MiYoung Kwon and Jan Skerswetat for

helpful discussions and the reviewers for their insightful feedback. This work was supported by

NIH grant R01EY032162 (P.J.B.). The Northeastern University IRB approved procedures (IRB

#14-09-16), and all observers consented to participation. The two images in Figure 1 were

generated in Unity (Unity Technologies, San Francisco, CA) using 3D models downloaded from

Sketchfab (Sketchfab Inc., Paris, France) and licensed under Creative Commons Attribution (CC

BY 4.0 https://creativecommons.org/licenses/by/4.0/). The 3D models were "Construction Sign"

by Kyle Burton (https://skfb.ly/6WS8O), "Nathan Animated 003 - Walking 3D Man" by

Renderpeople (https://skfb.ly/6SO7F) and "American Road Intersection" by jimbogies

(https://skfb.ly/o7z8V). The downloaded 3D models were modified in Blender (Blender

Foundation, Amsterdam, Netherlands). Special thanks go to Kerri Walter for her help creating

the images for Figure 1.

Data Availability

798 All data and code required to replicate the analyses in this paper will be made available on an

799 online repository at the stage of publication. It will also be made available to reviewers during

800 peer review.

References

Astle AT, McGovern DP, McGraw P V. 2014. Characterizing the role of disparity information in

alleviating visual crowding. *J Vis* **14**:1–14. doi:10.1167/14.6.8

Balas B, Nakano L, Rosenholtz R. 2009. A summary-statistic representation in peripheral vision

explains visual crowding. *J Vis* **9**:1–18. doi:10.1167/9.12.1

Banks WP, Larson DW, Prinzmetal W. 1979. Asymmetry of visual interference. *Percept*

Psychophys **25**:447–456. doi:10.3758/BF03213822

Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4.

J Stat Softw **67**:1–48. doi:10.18637/jss.v067.i01

Berens P. 2009. CircStat: a MATLAB toolbox for circular statistics. *J Stat Softw* **31**:1–29.

Bouma H. 1970. Interaction effects in parafoveal letter recognition. *Nature* **226**:177–178.

Bredemeier K, Simons DJ. 2012. Working memory and inattentional blindness. *Psychon Bull*

Rev **19**:239–244. doi:10.3758/s13423-011-0204-8

Chanceaux M, Mathôt S, Grainger J. 2014. Effects of number, complexity, and familiarity of

flankers on crowded letter identification. *J Vis* **14**:1–12. doi:10.1167/14.6.7

Coates DR, Ludowici CJH, Chung STL. 2021. The generality of the critical spacing for crowded

optotypes: from Bouma to the 21st century. *J Vis* **21**:1–22. doi:10.1167/jov.21.11.18

Eberhardt LV, Huckauf A. 2019. Crowding in depth for binocular and monocular observation.

Attention, Perception, Psychophys **81**:1951–1961.

Eberhardt LV, Huckauf A. 2017. Examining crowding using a real three-dimensional

experimental setup. *Proc Latv Acad Sci* **71**:313–319. doi:10.1515/prolas-2017-0055

Eberhardt LV, Pittino F, Huckauf A. 2021. Close - but not distant - conditioned flanker emotion

affects crowding. *J Vis* **21**:1–14. doi:10.1167/jov.21.8.22

Eberhardt L V, Huckauf A. 2020. Crowding effects across depth are fixation-centered for

defocused flankers and observer-centered for defocused targets. *Brian Sci* **10**:596.

doi:10.3390/brainsci10090596

Ester EF, Klee D, Awh E. 2014. Visual crowding cannot be wholly explained by feature pooling.

J Exp Psychol Hum Percept Perform **40**:1022–1033. doi:10.1037/a0035377

Ester EF, Zilber E, Serences JT. 2015. Substitution and pooling in visual crowding induced by

similar and dissimilar distractors. *J Vis* **15**:1–12. doi:10.1167/15.1.4

Felisberti FM, Solomon JA, Morgan MJ. 2005. The role of target salience in crowding.

Perception **34**:823–833. doi:10.1068/p5206

Freeman J, Simoncelli EP. 2011. Metamers of the ventral stream. *Nat Neurosci* **14**:1195–1204.

doi:10.1038/nn.2889

Georgeson MA, Wallis SA. 2014. Binocular fusion, suppression and diplopia for blurred edges.

Ophthalmic Physiol Opt **34**:163–185. doi:10.1111/opo.12108

Hampton DR, Kertesz AE. 1983. The extent of Panum's area and the human cortical

magnification factor. *Perception* **12**:161–165. doi:10.1068/p120161

Harrison WJ, Bex PJ. 2017. Visual crowding is a combination of an increase of positional

uncertainty, source confusion, and featural averaging. *Sci Rep* **7**:45551.

doi:10.1038/srep45551

Harrison WJ, Bex PJ. 2016. Ungrouping overcrowded explanations of visual crowding: reply to

Pachai et al. *Curr Biol* **26**:R353–R354. doi:10.1016/j.cub.2016.03.024

Harrison WJ, Bex PJ. 2015. A unifying model of orientation crowding in peripheral vision. *Curr*

Biol **25**:3213–3219. doi:10.1016/j.cub.2015.10.052

Harrold AL, Grove PM. 2019. Size increase in Panum's fusional range is driven by eccentricity

and not separation. *Perception* **48**:820–834. doi:10.1177/0301006619862434

Hartig F. 2022. DHARMa: Residual diagnostics for hierarchical (multi-level/mixed) regression

models. R package version 0.4.6. http://florianhartig.github.io/DHARMa/

He S, Cavanagh P, Lntriligator J. 1996. Attentional resolution and the locus of visual awareness.

Nature **383**:334–337. doi:10.1038/258748a0

Herzog MH, Manassi M. 2015. Uncorking the bottleneck of crowding: a fresh look at object

recognition. *Curr Opin Behav Sci* **1**:86–93. doi:10.1016/j.cobeha.2014.10.006

Herzog MH, Sayim B, Chicherov V, Manassi M. 2015. Crowding, grouping, and object

recognition: a matter of appearance. *J Vis* **15**:1–18. doi:10.1167/15.6.5

Ho C, Cheung SH. 2011. Crowding by invisible flankers. *PLoS One* **6**.

doi:10.1371/journal.pone.0028814

Hoffman DM, Girshick AR, Akeley K, Banks MS. 2008. Vergence-accommodation conflicts

hinder visual performance and cause visual fatigue. *J Vis* **8**:1–30. doi:10.1167/8.3.33

Hsu CW, Zhen B, Qiu W, Shapira O, DeLacy BG, Joannopoulos JD, Soljačić M. 2014.

Transparent displays enabled by resonant nanoparticle scattering. *Nat Commun* **5**:1–6.

doi:10.1038/ncomms4152

Kalpadakis-Smith A V., Goffaux V, Greenwood JA. 2018. Crowding for faces is determined by

visual (not holistic) similarity: Evidence from judgements of eye position. *Sci Rep* **8**:12556.

doi:10.1038/s41598-018-30900-0

Keshvari S, Rosenholtz R. 2016. Pooling of continuous features provides a unifying account of

crowding. *J Vis* **16**:1–15. doi:10.1167/16.3.39

Kim S, Shin E, Chong SC. 2013. When crowding meets binocular rivalry: challenges for object

perception. *Vision Res* **76**:134–143. doi:10.1016/j.visres.2012.10.020

Kooi FL, Toet A, Tripathy SP, Levi DM. 1994. The effect of similarity and duration on spatial

interaction in peripheral vision. *Spat Vis* **8**:255–279. doi:10.1163/156856894X00350

Legge GE. 2007. Psychophysics of reading in normal and low vision. Mahwah, NJ: Lawrence

Erlbaum Associates Publishers.

Levi DM. 2008. Crowding - an essential bottleneck for object recognition: a mini-review. *Vision*

Res **48**:635–654. doi:10.1016/j.visres.2007.12.009

Louie EG, Bressler DW, Whitney D. 2007. Holistic crowding: selective interference between

configural representations of faces in crowded scenes. *J Vis* **7**:1–11. doi:10.1167/7.2.24

Maiello G, Chessa M, Bex PJ, Solari F. 2020. Near-optimal combination of disparity across a

log-polar scaled visual field. *PLoS Comput Biol* **16**:1–28. doi:10.1371/journal.pcbi.1007699

Manassi M, Sayim B, Herzog MH. 2013. When crowding of crowding leads to uncrowding. *J*

Vis **13**:1–10. doi:10.1167/13.13.10

Manassi M, Sayim B, Herzog MH. 2012. Grouping, pooling, and when bigger is better in visual

crowding. *J Vis* **12**, **13**:1–14. doi:10.1167/12.10.13

Manassi M, Whitney D. 2018. Multi-level crowding and the paradox of object recognition in

clutter. *Curr Biol* **28**:R127–R133. doi:10.1016/j.cub.2017.12.051

Miles WR. 1929. Ocular dominance demonstrated by unconscious sighting. *J Exp Psychol*

12:113–126. doi:10.1037/h0075694

Nandy AS, Tjan BS. 2012. Saccade-confounded image statistics explain visual crowding. *Nat*

Neurosci **15**:463–469. doi:10.1038/nn.3021

Nityananda V, Read JCAA. 2017. Stereopsis in animals: evolution, function and mechanisms. *J*

Exp Biol **220**:2502–2512. doi:10.1242/jeb.143883

Norcia AM, Suiter EE, Tyler CW. 1985. Electrophysiological evidence for the existence of

coarse and fine disparity mechanisms in human. *Vision Res* **25**:1603–1611.

doi:10.1016/0042-6989(85)90130-0

Ogata NG, Boer ER, Daga FB, Jammal AA, Stringham JM, Medeiros FA. 2019. Visual

crowding in glaucoma. *Invest Ophthalmol Vis Sci* **60**:538–543.

doi:https://doi.org/10.1167/iovs.18-25150

Ono H, Angus R, Gregor P. 1977. Binocular single vision achieved by fusion and suppression.

Percept Psychophys **21**:513–521. doi:10.3758/BF03198731

Pachai M V., Doerig AC, Herzog MH. 2016. How best to unify crowding? *Curr Biol* **26**:R352–

R353. doi:10.1016/j.cub.2016.03.003

Panum PL. 1858. Physiologische untersuchungen über das sehen mit zwei augen. Kiel:

Schwerssche Buchhandlung.

Parkes L, Lund J, Angelucci A, Solomon JA, Morgan M. 2001. Compulsory averaging of

crowded orientation signals in human vision. *Nat Neurosci* **4**:739–744. doi:10.1038/89532

Peirce J, Gray JR, Simpson S, MacAskill M, Höchenberger R, Sogo H, Kastman E, Lindeløv JK.

2019. PsychoPy2: Experiments in behavior made easy. *Behav Res Methods* **51**:195–203.

909  doi:10.3758/s13428-018-01193-y

910 Pelli DG. 2008. Crowding: a cortical constraint on object recognition. *Curr Opin Neurobiol*

911  **18**:445–451. doi:10.1016/j.conb.2008.09.008

912 Pluháček F, Musilová L, Bedell HE, Siderov J. 2021. Number of flankers influences foveal

913  crowding and contour interaction differently. *Vision Res* **179**:9–18.

914  doi:10.1016/j.visres.2020.11.002

915 Pulliam K. 1981. Spatial frequency analysis of three-dimensional vision. *Proc SPIE Vis Simul*

916  *Image Realis II* **303**:71–77. doi:https://doi.org/10.1117/12.932660

917 Qin D, Takamatsu M, Nakashima Y. 2006. Disparity limit for binocular fusion in fovea. *Opt Rev*

918  **13**:34–38. doi:10.1007/s10043-006-0034-5

919 R Core Team. 2022. R: a language and environment for statistical computing.

920 Reynaud A, Hess RF. 2017. Characterization of spatial frequency channels underlying disparity

921  sensitivity by factor analysis of population data. *Front Comput Neurosci* **11**:63.

922  doi:10.3389/fncom.2017.00063

923 Sayim B, Westheimer G, Herzog MH. 2008. Contrast polarity, chromaticity, and stereoscopic

924  depth modulate contextual interactions in vernier acuity. *J Vis* **8**:1–9. doi:10.1167/8.8.12

925 Shamsi F, Liu R, Kwon MY. 2022. Binocularly asymmetric crowding in glaucoma and a lack of

926  binocular summation in crowding. *Invest Ophthalmol Vis Sci* **63**:36.

927  doi:https://doi.org/10.1167/iovs.63.1.36

928 Simons DJ, Chabris CF. 1999. Gorillas in our midst: sustained inattentional blindness for

929  dynamic events. *Perception* **28**:1059–1074. doi:10.1068/p281059

930 Spiegel DP, Baldwin AS, Hess RF. 2016. The relationship between fusion, suppression, and

931  diplopia in normal and amblyopic vision. *Investig Ophthalmol Vis Sci* **57**:5810–5817.

doi:10.1167/iovs.16-20438

Stidwill D, Fletcher R. 2010. Diplopia and confusion, suppression and rivalryNormal Binocular

Vision: Theory, Investigation and Practical Aspects. Chicester: John Wiley & Sons,

Incorporated. pp. 57–71. doi:10.1002/9781118788684.ch5

Strasburger H, Rentschler I, Jüttner M. 2011. Peripheral vision and pattern recognition: a review.

J Vis **11**:1–82. doi:10.1167/11.5.13.

Thaler L, Schütz AC, Goodale MA, Gegenfurtner KR. 2013. What is the best fixation target?

The effect of target shape on stability of fixational eye movements. *Vision Res* **76**:31–42.

doi:10.1016/j.visres.2012.10.012

Tyler CW, Barghout L, Kontsevich LL. 1994. Computational reconstruction of the mechanisms

of human stereopsis. *Proc SPIE Comput Vis Based Neurobiol* **2054**:52–68.

Ur Rehman A, Kihara K, Matsumoto A, Ohtsuka S. 2015. Attentive tracking of moving objects

in real 3D space. *Vision Res* **109**:1–10. doi:10.1016/j.visres.2015.02.004

van den Berg R, Johnson A, Anton AM, Schepers AL, Cornelissen FW. 2012. Comparing

crowding in human and ideal observers. *J Vis* **12**:13, 1–15.

doi:10.1167/12.6.13.Introduction

van den Berg R, Roerdink JBTM, Cornelissen FW. 2010. A neurophysiologically plausible

population code model for feature integration explains visual crowding. *PLoS Comput Biol*

6:e1000646. doi:10.1371/journal.pcbi.1000646

Viswanathan L, Mingolla E. 2002. Dynamics of attention in depth: Evidence from multi-element

tracking. *Perception* **31**:1415–1437. doi:10.1068/p3432

Wallis TSA, Bex PJ. 2012. Image correlates of crowding in natural scenes. *J Vis* **12**:1–19.

doi:10.1167/12.7.6

Wallis TSA, Bex PJ. 2011. Visual crowding is correlated with awareness. *Curr Biol* **21**:254–258.

doi:10.1016/j.cub.2011.01.011

Watt SJ, Akeley K, Ernst MO, Banks MS. 2005. Focus cues affect perceived depth. *J Vis* **5**:834–

862. doi:10.1167/5.10.7

Westheimer G. 1979. Cooperative neural processes involved in stereoscopic acuity. *Exp Brain*

Res **36**:585–597. doi:10.1007/BF00238525

Whitney D, Levi DM. 2011. Visual Crowding: a fundamental limit on conscious perception and

object recognition. *Trends Cogn Sci* **15**:160–168. doi:10.1016/j.tics.2011.02.005.Visual

Wilcox LM, Allison RS. 2009. Coarse-fine dichotomies in human stereopsis. *Vision Res*

49:2653–2665. doi:10.1016/j.visres.2009.06.004

Xia Y, Manassi M, Nakayama K, Zipser K, Whitney D. 2020. Visual crowding in driving. *J Vis*

20:1–7. doi:10.1167/jov.20.6.1

Yang Y, Blake R. 1991. Spatial frequency tuning of human stereopsis. *Vision Res* **31**:1176–1189.

doi:10.1016/0042-6989(91)90043-5

Supplementary figures

Figure 5-figure supplement 1: Perceptual error for Experiments 1 (A) and 2 (B) calculated only from trials in which the observer reported seeing the target inside the flanker ring. In both experiments, flanker depth was varied while the target was always at fixation depth. Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below each graph shows the depth of the flanker ring (based on the same color key as the box plot) relative to the depth of the fixation point and target (shown in black) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625° target-flanker spacing due to limited space within the figure.

Experiment 1: Target and fixation at 1.26 m with flanker ring presented at each depth. The effect of flanker depth relative to target-fixation depth ($\chi^2_{(2)} = 15.69$, p <0.001) and target-flanker spacing ($\chi^2_{(1)} = 20.03$, p <0.001) both remained significant. There was no significant interaction between flanker depth and target-flanker spacing ($\chi^2_{(2)} = 1.92$, p=0.384).

Experiment 2: Target and fixation at 0.4 m or 4 m with flanker ring presented at each depth. When perceptual error was calculated using only trials in which the observer reported seeing the target inside the flanker ring, flanker depth had no significant effect on perceptual error ($\chi^2_{(2)} =$

5.7, p = 0.058). In contrast, when perceptual error was calculated from the full data set there was a significant interaction between target-fixation depth and flanker depth (see Results section). The interaction between target-flanker spacing and target-fixation depth remained significant ($\chi^2_{(1)}$ = 7.32, p=0.007). There was no significant interaction between target-fixation depth and flanker depth ($\chi^2_{(2)}$ = 3.44, p=0.179) nor between flanker depth and target-flanker spacing ($\chi^2_{(2)}$ = 4.85, p=0.088).

Figure 6-figure supplement 1: Perceptual error for Experiments 3 (A) and 4 (B) calculated only from trials in which the observer reported seeing the target inside the flanker ring. In both experiments, target depth was varied while the flanker was always at fixation depth. Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below each graph shows the depth of the target (based on the same color key as the box plot) relative to the depth of the fixation point and flanker ring (shown in black) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625° target-flanker spacing due to limited space within the figure.

1010 <u>Experiment 3: Flanker ring and fixation at 1.26 m with target presented at each depth.</u> The effect

1011 of target depth relative to flanker-fixation depth ($\chi^2_{(2)}$ = 36.98, p <0.001) and target-flanker

1012 spacing ($\chi^2_{(1)}$ = 50.63, p <0.001) both remained significant. There was no significant interaction

1013 between target depth and target-flanker spacing ($\chi^2_{(2)}$ = 3.59, p=0.166).

1014

1015 <u>Experiment 4: Flanker ring and fixation at 0.4 m or 4 m with target presented at each depth.</u> The

1016 interaction between flanker-fixation depth and target depth ($\chi^2_{(2)}$ = 8.7, p = 0.013), and the effect

1017 of target-flanker spacing ($\chi^2_{(1)}$ = 7.58, p = 0.006) both remained significant. There was no

1018 significant interaction between flanker-fixation depth and target-flanker spacing ($\chi^2_{(1)}$ = 0.64,

1019 p=0.425) nor between target depth and target-flanker spacing ($\chi^2_{(2)}$ = 0.98, p=0.612).

1020

1021

Figure 8-figure supplement 1: **Proportion of trials in Experiment 1 in which observers**

reported seeing the target in the center of the flanker ring (A), within the flanker ring but

not in the center (B), partly or fully obstructed by the flanker ring (C), outside the flanker

ring (D), or they were unsure or there was no flanker ring (E). Flanker depth was varied

while the target was always at fixation depth. Box plots show medians plus the interquartile

range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from

the upper and lower quartiles. Points show the number of trials for each individual observer.

There were 10 repeats per stimulus condition.

1030

Figure 8-figure supplement 2: **Proportion of trials in Experiment 2 in which observers reported seeing the target in the center of the flanker ring (A), within the flanker ring but not in the center (B), partly or fully obstructed by the flanker ring (C), outside the flanker ring (D), or they were unsure or there was no flanker ring (E).** Flanker depth was varied while the target was always at fixation depth. Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the number of trials for each individual observer. There were 10 repeats per stimulus condition.

1039

Figure 9-figure supplement 1: **Proportion of trials in Experiment 3 in which observers**

reported seeing the target in the center of the flanker ring (A), within the flanker ring but

not in the center (B), partly or fully obstructed by the flanker ring (C), outside the flanker

ring (D), or they were unsure or there was no flanker ring (E). Target depth was varied while

the flanker was always at fixation depth. Box plots show medians plus the interquartile range

(IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the

upper and lower quartiles. Points show the number of trials for each individual observer. There

were 10 repeats per stimulus condition.

1048

Figure 9-figure supplement 2: **Proportion of trials in Experiment 4 in which observers reported seeing the target in the center of the flanker ring (A), within the flanker ring but not in the center (B), partly or fully obstructed by the flanker ring (C), outside the flanker ring (D), or they were unsure or there was no flanker ring (E).** Target depth was varied while the flanker was always at fixation depth**.** Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the number of trials for each individual observer. There were eight repeats per stimulus condition.

Figure 10-figure supplement 1: Proportion of trials in which each individual experienced subject (ES1-4) reported seeing the target inside the flanker ring for Experiments 1 and 2.
In both experiments, flanker depth was varied while the target was always at fixation depth.
Error bars show Wilson score intervals. The 5˚ target-flanker spacing condition was not included in this set of experiments. Since the same subjects participated in both experiments, the results from Experiments 1 and 2 have been combined into a single figure to make it easier to compare the different conditions. The visual aid below the graph shows the depth of the flanker ring (based on the same color key as the plot) relative to the depth of the fixation point and target

1066   (shown in black) for each stimulus condition. The relative sizes of, and distances between, the

1067   fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for

1068   the 0.625˚ target-flanker spacing due to limited space within the figure.

1069

Figure 10-figure supplement 2: Proportion of trials in Experiments 1 and 2 in which each individual experienced subject (ES1-4) reported seeing the target in the center of the flanker ring (A), within the flanker ring but not in the center (B), partly or fully obstructed by the flanker ring (C), outside the flanker ring (D), or they were unsure or there was no flanker ring (E). In both experiments' flanker depth was varied while the target was always at fixation depth. The 5˚ target-flanker spacing condition was not included in this set of experiments. There were 10 repeats per stimulus condition in both experiments. Since the same subjects participated in both experiments, the results from Experiments 1 and 2 have been combined into single figures to make it easier to compare the different conditions.

Figure 11-figure supplement 1: Proportion of trials in which each individual experienced subject (ES1-4) reported seeing the target inside the flanker ring for Experiments 3 and 4.
In both experiments' target depth was varied while the flanker ring was always at fixation depth.
Error bars show Wilson score intervals. The 5˚ target-flanker spacing condition was not included in this set of experiments. Since the same subjects participated in both experiments, the results from Experiments 3 and 4 have been combined into a single figure to make it easier to compare the different conditions. The visual aid below each graph shows the depth of the target (based on the same color key as the plot) relative to the depth of the fixation point and flanker ring (shown

1088   in black) for each stimulus condition. The relative sizes of, and distances between, the fixation

1089   point, target, and flanker ring are to scale. Note that the fixation point is only shown for the

1090   0.625˚ target-flanker spacing due to limited space within the figure.

1091

Figure 11-figure supplement 2: Proportion of trials in Experiments 3 and 4 in which each individual experienced subject (ES1-4) reported seeing the target in the center of the flanker ring (A), within the flanker ring but not in the center (B), partly or fully obstructed by the flanker ring (C), outside the flanker ring (D), or they were unsure or there was no flanker ring (E). In both experiments, target depth was varied while the flanker ring was always at fixation depth. The 5˚ target-flanker spacing condition was not included in this set of experiments. There were 10 repeats per stimulus condition in Experiment 3 and eight repeats in Experiments 4. Since the same subjects participated in both experiments, the results from Experiments 3 and 4 have been combined into single figures to make it easier to compare the different conditions.