Large differences in target-flanker depth increase crowding-evidence from a multi-depth plane display

Samuel P. Smithers[1*], Yulong Shao[1], James Altham[1], and Peter J. Bex[1*]

[1] Department of Psychology, Northeastern University, Boston, MA, USA

[*] Corresponding authors: SPS (s.smithers@northeastern.edu) and PJB (p.bex@northeastern.edu)

Abstract

Crowding occurs when the presence of nearby features causes highly visible objects to become unrecognizable. Although crowding has implications for many everyday tasks and the tremendous amounts of research reflect its importance, surprisingly little is known about how depth affects crowding. Most available studies show that stereoscopic disparity reduces crowding, indicating that crowding may be relatively unimportant in three-dimensional environments. However, most previous studies tested only small stereoscopic differences in depth in which disparity, defocus blur, and accommodation are inconsistent with the real world. Using a novel multi-depth plane display, this study investigated how large (0.54 - 2.25 diopters), real differences in target-flanker depth, representative of those experienced between many objects in the real world, affect crowding. Contrary to previous work showing small differences in target-flanker depth reduce crowding, the larger depth differences tested in our study instead increased crowding. While this is likely due to increased clutter in areas outside the limits of binocular fusion, analysis of the stimuli's perceived appearance indicates the increase in crowding occurs even in the absence of reported diplopia. These findings suggest that crowding has a more significant impact on our perception of natural three-dimensional environments than previously estimated with 2D displays.

Introduction

Visual crowding is a phenomenon in which a peripherally viewed object, that is easily

identifiable in isolation, becomes harder to recognize or identify when it is surrounded by other

objects, despite still being clearly visible (Bouma, 1970; Herzog et al., 2015; Levi, 2008; Pelli,

2008; Whitney and Levi, 2011). The importance of crowding is reflected by the tremendous

amount of research on the topic and it has been extensively reviewed over the years (e.g., Coates

et al., 2021; Herzog et al., 2015; Manassi and Whitney, 2018; Pelli, 2008; Strasburger et al.,

2011; Whitney and Levi, 2011). This high level of interest is not surprising given that the natural

environment is usually cluttered and mostly viewed in the peripheral visual field. Consequently,

it is assumed that crowding has a large impact on our daily lives with implications for many

everyday visual tasks including driving (Xia et al., 2020), reading (Legge, 2007) and face

recognition (Kalpadakis-Smith et al., 2018; Louie et al., 2007). Crowding also has important

clinical implications for people with central vision impairment (e.g., macular degeneration),

amblyopia and dyslexia (Whitney and Levi, 2011). Through this body of research, we have been

able to build a good understanding of the factors that mediate and moderate crowding and

develop unifying theories (Balas et al., 2009; Harrison and Bex, 2015; Keshvari and Rosenholtz,

2016). However, despite the large body of literature, one area that remains surprisingly

understudied is how depth affects crowding. This is important because we live in a three-

dimensional environment and so to understand the significance of crowding in the real world it is

essential to understand the role of depth information.

Kooi et al. (1994) were the first to investigate how depth influences crowding. The

authors found that compared to when a peripheral target and flankers were at the same depth,

there was a release from crowding when the target and flankers were stereoscopically presented

in crossed (front) and uncrossed (behind) disparities (Kooi et al., 1994). Sayim et al. (2008)

found that when flankers were at fixation depth, there was less foveal crowding when the target

was stereoscopically presented in front or behind the flankers. Felisberti et al. (2005) also found

an effect of depth on crowding in the parafovea, but to a more limited extent. They reported a

release from crowding in two out of three subjects when the target appeared behind the flankers,

but only in one of the three subjects when the target was in front of the flankers (Felisberti et al.,

2005). Astle et al. (2014) extended on previous research by testing the effect of depth on

crowding over a range of stereoscopically produced target-flanker disparities (±800 arcsec). In

line with previous studies, Astle et al. (2014) found that when the target was at fixation depth,

increasing target-flanker disparity systematically reduced crowding. They also found that the

release from crowding was asymmetric whereby crowding was greater when the flankers were in

front of the target compared to behind it (Astle et al., 2014).

Taken together these studies appear to paint the picture that a difference in depth between

a target and flankers reduces crowding. This challenges the assumption that crowding is a

significant problem in three-dimensional natural environments. However, all the above studies

tested only small depth differences between target and flankers that do not reflect the often-large

differences in depth between objects in the real world. Furthermore, the generalizability of these

studies to the real world is also limited by the fact depth cues were generated stereoscopically.

One potential problem with stereo displays is that disparity information often does not match

with accommodation or defocus blur which can detrimentally effect depth perception (Hoffman

et al., 2008; Watt et al., 2005).

To our knowledge, four studies have investigated the effect of depth on crowding using a

real depth display. All four studies used two orthogonally positioned screens viewed through a

half-transparent mirror allowing stimuli to be displayed on two different depth planes

simultaneously (Eberhardt et al., 2021; Eberhardt and Huckauf, 2019, 2017, 2020). Eberhardt

and Huckauf (2017) kept a target and flankers at the same depth and presented them 0.06

diopters in front, behind or at the same depth as fixation. The authors found that this small

difference in depth between the stimuli and fixation planes had no effect on the amount of

crowding compared to when everything was presented at the fixation plane (Eberhardt and

Huckauf, 2017). Eberhardt and Huckauf (2019) repeated and expanded on this work with the

addition of a ± 0.1 diopter depth difference between fixation and the stimuli in addition to the ±

0.06 diopter condition. The results showed that crowding was greater when the stimuli were

furthest from fixation compared to when they were closer to fixation, regardless of whether the

stimuli were in front or behind the fixation plane (Eberhardt and Huckauf, 2019). More recently,

Eberhardt and Huckauf (2020) used the same approach to investigate the effect of depth on

crowding when the target and flankers were separated in depth. When flanker depth was varied

while the target was at fixation depth, crowding increased with increasing target-flanker depth

difference (Eberhardt and Huckauf, 2020). This result is the opposite to the findings of other

studies in which small differences in depth between the target and flankers were introduced with

stereoscopic disparity (Astle et al., 2014; Felisberti et al., 2005; Kooi et al., 1994; Sayim et al.,

2008). Furthermore, the authors also found that when flanker depth was varied while the target

was at fixation depth, crowding was greater when the flankers were behind the target and

fixation compared to in front of it (Eberhardt and Huckauf, 2020). When target depth was varied

while the flankers were at fixation, there was more crowding when the target was in front of the

flankers and fixation compared to when it was behind them (Eberhardt and Huckauf, 2020).

In light of the above discrepancies, and the prevalence of depth information in real world

scenes, there is a pressing need for a better understanding of how visual crowding is affected by

depth differences representative of those experienced between objects in the real world.

Therefore, using a novel multi-depth plane display consisting of screens at 0.4 m, 1.26 m, and 4

m viewing distances, we investigate how crowding is affected by large, real differences in target-

flanker depth.

Methods

Subject recruitment

A total of 110 naïve subjects were recruited from the Northeastern University undergraduate

population in exchange for course credit as compensation for their time. Of these, seven were

excluded immediately following screening because they did not meet the inclusion criteria of

normal or corrected-to-normal vision. An additional five subjects were excluded because of

issues with the experimental code (N=3), calibration issues (N=1) or because they did not follow

the task instructions (N=1). Thus, a total of 98 subjects (32 males, 66 females; mean age = 18.86

years) contributed usable data across five separate experiments (22 in Experiment 1, 19 in

Experiment 2, 21 in Experiments 3 and 4, and 15 in Experiment 5). All subjects read and signed

an informed consent form approved by the Northeastern University Ethics Board before the

experiment began. The experimental procedure was approved by the institutional review board at

Northeastern University, and the experiments were performed in accordance with the tenets of

the Declaration of Helsinki.

Setup

We developed a multi-depth plane display consisting of three screens at 0.4 m (near screen), 1.26

m (mid screen) and 4 m (far screen) from the observer (Figure 1). These distances were chosen

because 4 m is close to optical infinity while 0.4 m is the clinical standard for near distance, with

1.26 m being in the middle of these two extremes in log units. Given in diopters (i.e., 1/focal

length in meters), the near, mid, and far screens were at 2.5 diopters, 0.79 diopters and 0.25

diopters receptively. For the far screen we used an ultra-short throw projector (Bomaker Polaris

4K Laser Projector) projected onto a 240 cm x 250 cm white wall at the end of the room. The

mid and near screens were made from 101.5 cm x 76 cm and 46 cm x 36.5 cm clear 5 mm

acrylic respectively held in position by a frame constructed from extruded aluminum rails. The

transparent acrylic was covered on one side with ClearBright transparent display film (Lux Labs

Inc., Boston, MA, USA). This nanoparticle film enabled us to display images from two

projectors (InFocus IN3116) onto the near and mid screens while preserving their transparency

(Hsu et al., 2014), thus allowing observers to simultaneously view adjacent stimuli presented on

all three depths planes at the same time. The screens were temporally synchronized and spatially

aligned to ensure display timing and position were consistent across the three depth planes.

Figure 1: Illustration of the multi-depth plane display setup used for this study.

Spatial calibration

Each observer performed a subjective spatial calibration at the start of Experiments 1-4 to ensure

that the angular positions of the stimuli and fixation point relative to primary gaze were the same

(or as close as realistically possible) across all three depth planes. The center of the far screen

(which was at primary gaze when the observers head was positioned in the chin and head rest)

was used as the fixed reference against which all three screens were calibrated. With their head

in the chin and head rest observers were instructed to sequentially position a series of targets so

that they were in the same position as a fixed reference target that was displayed on the reference

screen. The calibration was performed under monocular conditions using the observer's

dominant eye (determined by the Miles test). The calibration was performed monocularly

because during a preliminary experiment subjects reported finding the task too difficult to

complete under binocular conditions (e.g., because of diplopia, an issue that is considered further

in the Discussion section). For Experiment 5, the spatial calibration was performed by the

experimenter prior to the experiment using a webcam positioned at the observer's cyclopean

point. Observers were asked to maintain the same head position within the chin and head rest

throughout the experiment.

Stimuli and experimental conditions

The target and flankers were inspired by those used in previous studies (Harrison and Bex, 2017,

2015). The target was a white Landolt-C displayed against a black background at 10˚ eccentricity

to the right or left of the fixation point at primary gaze. Following the structure of a Landolt-C

optotype, the target was 1˚ in dimeter with a 0.25˚ line width and a 0.25˚ opening. The opening

orientation of the target was random each trial, drawn from a uniform 360˚ distribution. The

target was surrounded by a white flanker ring formed of four circular arcs (Figure 2). In each

depth combination described below, the outer edge of the target was separated from the inner

edge of the flanker ring by 5˚, 2.5˚, 1.25˚ or 0.625˚. These distances of separation represented

1/2, 1/4, 1/8 and 1/16 of the target eccentricity respectively. The line width and size of the four

gaps in the flanker ring were kept constant (0.25˚). For each target depth, a control condition

with no flanker ring was also included. The fixation point consisted of a 1˚ white circle with a

black cross (Figure 2). This design was used because it has been reported to elicit more stable

fixation compared to a plain circle or cross hair alone (Thaler et al., 2013). The background on

all screens was always set to the lowest RGB (i.e. R=G=B=0) value which was transparent on

the near and mid screens, and black on the far screen. Since the mid screen was dimmest

(R=G=B=255, white=13.56 cd/m^2), the luminance of stimuli displayed on the near and far

screens were calibrated to match the luminance of the mid screen using a Photo Research PR-

650 spectroradiometer (Photo Research, Chatsworth, CA, USA). The room lights were switched

off for the duration of the experiment.

Figure 2: Schematic illustration of experimental stimuli and fixation marker. In all conditions, except the

no-flanker condition, the randomly orientated target (a white Landolt-C) was surrounded by a white

flanker ring at one of four possible target-flanker distances. The stimuli was viewed against a black

background.

A summary of the stimulus conditions tested in each experiment is provided in Table 1. Briefly,

in Experiments 1 and 2 the target was always presented at fixation depth while the depth of the

flanker was varied. In Experiment 1 the target and fixation point were always on the mid screen

while each flanker was displayed in front, behind or at fixation depth. In Experiment 2 the target

and fixation point were presented on the near or far screen while the flanker was presented at one

of the three depths across trials for each fixation distance. Experiments 3 and 4 were the same as

Experiments 1 and 2 except that the flanker ring was presented at fixation depth while the target

depth was varied. In Experiment 3, the flanker and fixation point were always on the mid screen

while the target was displayed on one of the three depth planes across trials. In Experiment 4, the

flankers and fixation point were presented on the near or far screen while the target was

presented at each of the three depths across trials for each fixation distance. For Experiment 5,

the target, flanker ring and fixation were presented together on the same screen at three different

depths. This was to investigate whether there was an effect of absolute viewing distance on

crowding.

Table 1: Summary of the stimulus conditions in each experiment. The same color scheme is used to
represent the near (0.4 m), mid (1.26 m) and far (4 m) screens throughout the manuscript. Note that for
target-flanker spacing, 'infinity' = no flanker ring.

Experiment	Fixation depth (m)	Target depth (m)	Flanker depth (m)	Target-flanker spacing (degrees)	No. of stimulus conditions	Total no. presentations	Sample size
Exp 1	Mid (1.26)	Mid (1.26)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°	13	130 (10 reps per stimulus)	N = 22
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Mid (1.26)	Mid (1.26)	Far (4)	0.625°, 1.25°, 2.5°, 5°			
Exp 2	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	26	260 (10 reps per stimulus)	N = 19
	Near (0.4)	Near (0.4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°			
	Near (0.4)	Near (0.4)	Far (4)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 3	Mid (1.26)	Near (0.4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity	15	150 (10 reps per stimulus)	N = 21
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Mid (1.26)	Far (4)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 4	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	30	240 (8 reps per stimulus)	N = 21
	Near (0.4)	Mid (1.26)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Near (0.4)	Far (4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Near (0.4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Mid (1.26)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			
Exp 5	Near (0.4)	Near (0.4)	Near (0.4)	0.625°, 1.25°, 2.5°, 5°, infinity	15	150 (10 reps per stimulus)	N = 15
	Mid (1.26)	Mid (1.26)	Mid (1.26)	0.625°, 1.25°, 2.5°, 5°, infinity			
	Far (4)	Far (4)	Far (4)	0.625°, 1.25°, 2.5°, 5°, infinity			

Each condition was repeated 10 times in Experiments 1, 2, 3 and 5 and eight times in

Experiment 4, with equal number of repeats on either side of the fixation point. The order of

testing was randomized.

An example trial is illustrated in Figure 3. Briefly, at the start of each trial, observers

were instructed to focus on the central fixation point. The fixation point appeared on the fixation

screen 1 sec before stimulus onset in Experiments 1, 3 and 5, and 1.5 sec before stimulus onset in

Experiments 2 and 4. The target and flanker ring (if present) were displayed for 0.25 sec. The

fixation point remained on the screen for 1 sec before being replaced by a randomly oriented

Landolt-C that was the same size as the target Landolt-C. The observer used the mouse to rotate

this Landolt-C until it matched the perceived orientation of the target and then clicked to report

the orientation. In all experiments except Experiment 5, the observer then reported the perceived

location of the target relative to the position of the flanker ring. There were 5 options to choose

from, "target in center of ring", "target inside ring but not in the center", "ring obstructs target",

"target outside ring", and "unsure or no ring". Observers were instructed to select the option that

most closely matched how they perceived the stimulus. If an observer missed the stimulus or

they were unsure of the targets' location relative to the flanker ring, e.g., because of diplopia,

they were instructed to select the "unsure or no ring" option. Following this selection, the next

trial started immediately. Before starting an experiment, all participants underwent training

during which they were encouraged to complete as many practice trials as they needed to

familiarize themselves with the task. These practice trials did not contribute to the collected data.

Figure 3: A step-by-step example of a single trial. Note that step five was not

included in Experiment 5.

Data processing and statistical analysis

<u>Perceptual error</u>

To quantify crowding for each stimulus condition we first calculated the report error for each

presentation, i.e., the angular difference between the reported gap orientation of the target and

the actual gap orientation. For each subject, this generated 10 report errors per stimulus condition

in Experiments 1, 2, 3 and 5, and eight per stimulus condition in Experiment 4. We then used

these report errors to calculate the circular standard deviation for each stimulus condition using

the Circular Statistics Toolbox (Berens, 2009) in MATLAB R2022a (MathWorks Inc., Natick,

MA, USA). We refer to this measure as perceptual error, which is reported in degrees.

The effect of target/flanker depth and target-flanker spacing on perceptual error was

analyzed using mixed effects multiple linear regression, fitted using the lme4 package (Bates et

al., 2015) in R (R Core Team, 2022). Perceptual error was used as the response (dependent)

variable after being natural log transformed to correct for positive skew. For all five experiments,

we included target-flanker spacing as a fixed effect. We also included flanker depth for

Experiments 1 and 2, target depth for Experiments 3 and 4 or target-flanker-fixation depth for

Experiment 5. Target-fixation depth or flanker-fixation depth were also included as a third fixed

effect for Experiment 2 and 4 respectively. All possible pairwise interactions of fixed effects

were fitted in the full model. Subject identification (ID) was included as a random effect in all

experiments to control for repeated measures. We first used model simplification, whereby

models were compared with one another using a likelihood ratio test (LRT), to remove non-

significant interactions sequentially. This enabled us to test the effect of individual fixed effects

that were not involved in an interaction. We then used the minimum model (i.e., model

containing all fixed effects of interest plus any significant two-way interactions) to report the

significance of each main effect or important interaction(s) by using a LRT to compare the

minimum model with the same model but with the effect/interaction of interest removed. Note

that data for the control condition (i.e., no flanker ring) were excluded from statistical analysis.

This was to ensure that a) the fixed effects remained independent and b) the effect of having no

flanker ring did not mask any of the effects we were most interested in (see Supplementary

Material for a more detailed explanation).

<u>Perceived target position relative to flanker ring</u>

For statistical analysis, the results for the perceived target position relative to the flanker ring

were converted to a binary response. A value of 1 meant that the observer perceived the target as

being inside the flanker ring (i.e., they chose either "target in center of ring" or "target inside ring

but not in the center") while a value of 0 meant the subject did not perceive the target as being

inside the flanker ring (i.e., they chose "ring obstructs target", "target outside ring" or "unsure or

no ring").

These data were then analyzed using mixed effects binary logistic regression, fitted using

the lme4 package (Bates et al., 2015) in R (R Core Team, 2022). We used the DHARMa package

(Hartig, 2022) to run residual diagnostics that are often overlooked in logistic regression.

Perceived target position was used as the binary response variable (i.e., 1 = target inside ring and

0 = target not inside ring). The fixed effect, and pairwise interaction for each experiment were

the same as those described above for the analysis of perceptual error. Subject ID was included

as a random effect in all experiments to control for repeated measures. Model simplification (to

remove non-significant interactions) and significance testing was the same as described above

for the analysis of perceptual error. Data for the control condition (i.e., no flanker ring) were

excluded from statistical analysis for the same reasons as those outlined above.

The separate results showing the number of times observers chose each of the five

choices for perceived target position relative to the flanker ring can be found in the

Supplementary Material (Figures S1-S4).

Results

Perceptual error

<u>Target always at fixation depth with variable flanker depth</u>

Experiment 1: Target and fixation at 1.26 m with flanker ring presented at each depth (Figure 4A). Perceptual error was higher when the flanker ring was displayed in front of, or behind, the target and fixation plane (mixed effects multiple linear regression models compared via LRT: $\chi^2_{(2)} = 20.67$, p <0.001). As expected, perceptual error decreased as target-flanker spacing increased ($\chi^2_{(1)} = 31.71$, p <0.001).

Experiment 2: Target and fixation at 0.4 m or 4 m with flanker ring presented at each depth (Figure 4B). Perceptual error tended to increase with increasing flanker depth when the flanker ring was behind the target and fixation plane, however, the effect of flanker depth was much weaker when the flanker ring was in front of the target and fixation (interaction between target-fixation depth and flanker depth: $\chi^2_{(2)} = 17.18$, p<0.001). Perceptual error decreased with increasing target-flanker spacing, in a manner corresponding to target-fixation depth (interaction between target-flanker spacing and target-fixation depth: $\chi^2_{(1)} = 9.2$, p=0.002).

Figure 4: Perceptual error results for Experiments 1 (A) and 2 (B) in which flanker depth was varied while the target was always at fixation depth. Box plots show medians plus the interquartile range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below each graph shows the depth of the flanker ring (based on the same color key as the box plot) relative to the depth of the fixation point and target (shown in black) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker spacing due to limited space within the figure.

Flanker ring always at fixation depth with variable target depth

Experiment 3: Flanker ring and fixation at 1.26 m with target presented at each depth (Figure 5A). Overall, perceptual error was higher when the target was displayed in front of, or behind, the flanker ring and fixation ($\chi^2_{(2)} = 36.93$, p <0.001), with the greatest perceptual error occurring when the target was behind the flanker ring and fixation. Perceptual error decreased as target-flanker spacing increased ($\chi^2_{(1)} = 45.99$, p <0.001).

Experiment 4: Flanker ring and fixation at 0.4 m or 4 m with target presented at each depth (Figure 5B). Perceptual error was greatest when the target was displayed behind the flanker ring and fixation, while the effect of depth was less prominent when the target was displayed in front of the flanker ring and fixation (interaction between flanker-fixation depth and target depth: $\chi^2_{(2)} = 61.87$, p<0.001). Perceptual error decreased as target-flanker spacing increased ($\chi^2_{(1)} = 29.92$, p<0.001).

Figure 5: **Perceptual error results for Experiments 3 (A) and 4 (B) in which target depth was varied**
321

while the flanker ring was always at fixation depth. Box plots show medians plus the interquartile
322

range (IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper
323

and lower quartiles. Points show the perceptual error for each individual observer. The visual aid below
324

each graph shows the depth of the target (based on the same color key as the box plot) relative to the
325

depth of the fixation point and flanker ring (shown in black) for each stimulus condition. The relative
326

sizes of, and distances between, the fixation point, target, and flanker ring are to scale. Note that the
327

fixation point is only shown for the 0.625˚ target-flanker spacing due to limited space within the figure.
328

Target and flanker ring at fixation depth
330

Experiment 5: Target and flanker ring always at fixation depth (Figure 6). Perceptual error
331

decreased as target-flanker spacing increased ($\chi^2_{(1)}$ = 64.8, p<0.001). There was also a significant
332

effect of depth ($\chi^2_{(2)}$ = 8.75, p=0.013) resulting from a higher perceptual error for stimuli
333

displayed at 4 m.
334

Figure 6: **Perceptual error results for Experiment 5 in which the target and flanker were always**

presented at the same depth as the fixation point. Box plots show medians plus the interquartile range

(IQR), whiskers are the lowest and highest values that are within 1.5 times the IQR from the upper and

lower quartiles. Points show the perceptual error for each individual observer. The visual aid below the

graph shows the depth of the target, flanker, and fixation point (based on the same color key as the box

plot) for each stimulus condition. The relative sizes of, and distances between, the fixation point, target,

and flanker ring are to scale. Note that the fixation point is only shown for the 0.625˚ target-flanker

spacing.

Perceived target position relative to the flanker ring

Target always at fixation depth with variable flanker depth

Experiment 1: Target and fixation at 1.26 m with flanker ring presented at each depth

(Figure 7A and supplementary Figure S1). In Experiment 1, most observers consistently

reported seeing the target inside the flanker ring. However, when the flanker ring was away from

target-fixation depth there was a reduction in the proportion of observers who consistently

reported seeing the target within the flanker ring when the target-flanker spacing was small

(mixed effects binary logistic regression models compared via LRT; interaction between target-

flanker spacing and flanker depth: $\chi^2_{(2)} = 36.12$, p <0.001). This effect was greatest when the

flanker ring was displayed behind the target and fixation.

Note that in all four experiments, the response to the control (i.e., no flanker condition)

indicates the probability of an observer making an inaccurate report (e.g., they selected the

wrong option by mistake or missed the stimulus and reported a guess rather than selecting the

"Unsure" option).

Experiment 2: Target and fixation at 0.4 m or 4 m with flanker ring presented at each

depth (Figure 7B and supplementary Figure S2). As with the previous experiments, when the

flanker ring was away from target-fixation depth, there was a reduction in the overall proportion

of trials in which observers reported seeing the target within the flanker ring (interaction between

target-fixation depth and flanker depth: $\chi^2_{(2)} = 374.26$, p < 0.001). The greatest reduction

occurred when target-flanker spacing was smaller ($\chi^2_{(1)} = 178.65$, p < 0.001).

Figure 7: **Proportion of trials in which observers reported seeing the target inside the flanker ring**

for Experiments 1 (A) and 2 (B). In both experiments flanker depth was varied while the target was

always at fixation depth**.** Box plots show medians plus the interquartile range (IQR), whiskers are the

lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points

show the proportion of trials for each individual observer. The visual aid below each graph shows the

depth of the flanker ring (based on the same color key as the box plot) relative to the depth of the fixation

point and target (shown in black) for each stimulus condition. The relative sizes of, and distances

between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown

for the 0.625˚ target-flanker spacing due to limited space within the figure.

<u>Flanker ring always at fixation depth with variable target depth</u>

Experiment 3: Flanker ring and fixation at 1.26 m with target presented at each depth

(Figure 8A and supplementary Figure S3). When the target was away from flanker-fixation

depth, observers were less likely to report seeing the target within the flanker ring. This effect

was greatest when target-flanker spacing was smaller and when the target was displayed in front

of the flanker ring and fixation (interaction between target-flanker spacing and target depth: $\chi^2_{(2)}$

$= 100.38$, p <0.001).

Experiment 4: Flanker ring and fixation at 0.4 m or 4 m with target presented at each

depth (Figure 8B and supplementary Figure S4). As with the previous experiment, the greater

the difference between target depth and flanker-fixation depth, the less likely observers were to

report seeing the target within the flanker ring (interaction between flanker-fixation depth and

target depth: $\chi^2_{(2)} = 779.49$, p < 0.001), particularly when target-flanker spacing was small

(interaction between target-flanker spacing and target depth: $\chi^2_{(2)} = 23.27$, p < 0.001). However,

target-flanker spacing had a weaker effect on the proportion of trials in which observers reported

seeing the target within the flanker ring when the flanker ring and fixation were near compared

to far (interaction between target-flanker spacing and flanker-fixation depth: $\chi^2_{(1)} = 56.23$, p $<$

0.001).

Figure 8: **Proportion of trials in which observers reported seeing the target inside the flanker ring**

for Experiments 3 (A) and 4 (B). In both experiments target depth was varied while the flanker ring was

always at fixation depth**.** Box plots show medians plus the interquartile range (IQR), whiskers are the

lowest and highest values that are within 1.5 times the IQR from the upper and lower quartiles. Points

show the proportion of trials for each individual observer. The visual aid below each graph shows the

depth of the target (based on the same color key as the box plot) relative to the depth of the fixation point

and flanker ring (shown in black) for each stimulus condition. The relative sizes of, and distances

between, the fixation point, target, and flanker ring are to scale. Note that the fixation point is only shown

for the 0.625˚ target-flanker spacing due to limited space within the figure.

Discussion

In all experiments, perceptual error decreased as the target-flanker spacing increased, which is in

line with previous crowding research (Bouma, 1970; Pelli, 2008). Overall, perceptual error was

higher when the target or flanker ring were presented at a different depth from the other and from

fixation. The greatest differences in perceptual error among the different depth conditions

occurred when the target or flanker ring were presented behind fixation depth. In contrast, when

the target or flanker ring were presented in front of fixation depth, the differences in perceptual

error among depth conditions were notably smaller. The results also show that when the target,

flanker ring and fixation were all at the same depth plane, crowding tended to increase with the

absolute distance of the stimuli from the observer.

The main finding that a depth difference between the flankers and the target led to an

increase, rather than a decrease, in perceptual error is different from the findings of several

previous studies that investigated the effect of depth on crowding using a stereoscopic

presentation (Astle et al., 2014; Felisberti et al., 2005; Kooi et al., 1994; Sayim et al., 2008). For

example, Astle et al (2014) found that when the target was presented at fixation depth

(comparable to Experiments 1 and 2 in our study), increasing the disparity between the target and

flankers resulted in less crowding, and ultimately a complete release from crowding. The authors

also reported that crowding was greater when flankers were presented in crossed disparity (in

front of the target and fixation), compared to uncrossed disparity (behind the target and fixation)

(Astle et al., 2014). Similarly, when the flankers were presented at fixation depth (comparable to

Experiments 3 and 4 in our study), Sayim et al (2008) found that foveal crowding was lower

when the target was presented in front of or behind the flanker and fixation.

Several explanations for crowding have been proposed in which crowding occurs because

target and flanking features are inappropriately grouped (Banks et al., 1979; Pachai et al., 2016).

Consistent with this explanation, crowding can be alleviated when the flankers are grouped

together and segmented from the target (Herzog and Manassi, 2015). One piece of evidence in

support of this account is the reduction of crowding that can occur when flankers are presented at

a different depth from the target (Astle et al., 2014; Felisberti et al., 2005; Kooi et al., 1994;

Sayim et al., 2008). Our results challenge this view of crowding because it would predict that

increasing the depth difference between a target and flankers would decrease the probability of

the flankers being grouped and increase the probability of their being segmented from the target,

thereby decreasing crowding, the opposite of our finding. Other models of crowding, such as

texture processing (Balas et al., 2009; Freeman and Simoncelli, 2011; Parkes et al., 2001; Wallis

and Bex, 2012), sampling errors (Ester et al., 2015, 2014; Harrison and Bex, 2017, 2016, 2015),

attentional resolution limits (He et al., 1996) or saccade properties (Nandy and Tjan, 2012) are

agnostic concerning the influence of depth differences among targets and flankers. Therefore,

each of these models would require modification to explain the increase in crowding that we

observe with large real depth differences between a target and its flankers.

In contrast with studies that controlled depth with small stereoscopic disparities, our

results do agree with the findings of Eberhardt and Huckauf (2020) who also used a real-depth

display and tested larger differences in depth between the target and flankers than previous

studies had examined (though depth difference were still much smaller than those tested in our

current study). When the target was at fixation depth (comparable to Experiments 1 and 2 in our

study), Eberhardt and Huckauf (2020) found that crowding increased with increasing target-

flanker depth difference. The authors also found that crowding was greater when the flankers

were behind the target and fixation compared to in front of it (Eberhardt and Huckauf, 2020).

Our findings from Experiments 1 and 2 are consistent with these results. However, when the

flankers were at fixation depth (comparable to Experiments 3 and 4 in our study), Eberhardt and

Huckauf (2020) found that crowding was higher when the target was in front of the flankers and

fixation compared to when it was behind them. In contrast, in Experiments 3 and 4 we found that

perceptual error was greatest when the target was presented behind the flanker ring and fixation

rather than in front of it. This difference is likely due to a combination of high defocus blur when

the target was behind fixation combined with the finding from Experiment 5 which found that in

our study the absolute distance from the observer resulted in higher perceptual error.

Why the inconsistencies across studies? It is important to consider that most of the

previous studies used very small depth differences, the largest being 0.1 diopters in Eberhardt

and Huckauf (2020) and Eberhardt and Huckauf (2019). In comparison, the depth difference in

our study ranged from 0.54 diopters (between the mid and far screens) to 2.25 diopters (between

the near and far screens). Our results, together with those from previous studies, suggest that

small differences in depth between the target and flankers reduce crowding (Astle et al., 2014;

Felisberti et al., 2005; Kooi et al., 1994; Sayim et al., 2008), while larger differences in depth

increase it (Eberhardt and Huckauf, 2020; and results from our study). Although this pattern is

quite surprising, there is precedent for non-linear effects of depth on perceptual processing

outside the crowding literature. In a multiple object tracking task, when the targets and

distractors were distributed equally on two depth planes, tracking accuracy was better than the

accuracy during trials in which targets and distractors appeared on a single depth plane when the

depth difference between planes was small (~6 cm) (Ur Rehman et al., 2015; Viswanathan and

Mingolla, 2002), but worse when the depth difference between planes was large (50 cm) (Ur

Rehman et al., 2015).

How could depth affect perception in this nonlinear way? The geometry of the eyes and

the 3D environment create interocular disparity differences between the images in each eye.

When corresponding features in the images in each eye can be identified, the images can be

fused to generate a binocular perceptual experience of depth (for review see Nityananda and

Read, 2017)). There is a horizontal and vertical spatial limit to the interocular disparity

difference over which the visual system attempts to identify corresponding retinal points in order

to fuse different 2D interocular images as a single 3D object (Westheimer, 1979). This 2D spatial

correspondence limit creates a 3D volume along the horopter known as Panum's Fusional Area

(Panum, 1858). Outside Panum's Fusional Area, there is a loss of stereoscopic depth perception

and either double vision (diplopia) may be experienced, or single vision may occur if the image

from one eye is partly suppressed (Georgeson and Wallis, 2014; Ono et al., 1977; Spiegel et al.,

2016; Stidwill and Fletcher, 2010). The spatial scale of Panum's Fusional Area depends on the

spatial frequency of the stimuli (Maiello et al., 2020; Pulliam, 1981; Reynaud and Hess, 2017;

Wilcox and Allison, 2009; Yang and Blake, 1991) and increases with eccentricity (Hampton and

Kertesz, 1983; Qin et al., 2006).

It is likely that differences between fusion and depth perception within Panum's Fusional

Area versus diplopia, and/or local suppression for single vision, outside of it can explain the

patterns of results observed between studies. Small disparity differences between target and

flankers create differences in perceived depth that can alleviate the effects of crowding,

potentially through grouping (Herzog and Manassi, 2015) or by independent processing of target

and flankers in different depth-selective channels (Norcia et al., 1985; Pulliam, 1981; Reynaud

and Hess, 2017; Tyler et al., 1994; Wilcox and Allison, 2009; Yang and Blake, 1991) by any of

the computational models of crowding based on texture processing (Balas et al., 2009; Freeman

and Simoncelli, 2011; Parkes et al., 2001; Wallis and Bex, 2012), sampling errors (Ester et al.,

2015, 2014; Harrison and Bex, 2017, 2016, 2015) attentional resolution limits (He et al., 1996) or

saccade properties (Nandy and Tjan, 2012).

Large interocular disparity differences on the other hand likely induce diplopia that could

increase perceptual error by increasing the number of features (Eberhardt and Huckauf, 2019).

Surprisingly, when asked, only a few observers reported experiencing diplopia, and those that

did reported only experiencing it in a minority of presentations for which they selected the

'unsure or no ring' option (as instructed) when reporting the perceived location of the target

(supplementary Figures S1E, S2E, S3E and S4E). However, as mentioned in the Methods

sections, observers did report experiencing diplopia when trying to perform the spatial

calibration task under binocular conditions during a pilot experiment. It is possible that the short

presentation time of the stimulus was too fast for observers to notice diplopia consciously, which

could explain why few observers reported diplopia in the main experiments.

Another explanation for higher perceptual error with increasing target-flanker depth

differences could be perceived target misalignment. Recall that to avoid diplopia during the

spatial calibration, the screens were calibrated monocularly with the dominant eye. This means

that the screens were imperfectly aligned for the non-dominant eye. Therefore, at certain flanker

sizes and separations there may be occlusions between target and flanker features that could

affect target report accuracy, which is not the same mechanism as crowding. Such misalignment

could result from diplopia, local suppression of the calibrated eye to maintain single vision,

and/or head movements following the spatial calibration. To examine how target misalignment

(i.e., target obstructed ring, target was outside ring, or they were unsure, e.g., due to diplopia)

affected the overall results, we recalculated perceptual error using only trials in which the

observer reported seeing the target inside the flanker ring. We then reanalyzed these data the

same way as for perceptual error calculated from the full data set (supplementary Figures S5 and

S6). Apart from Experiment 2, the effect of flanker and/or target depth remained statistically

significant (see supplementary materials for analysis results). Therefore, across the majority of

experiments, misalignment or occlusion of the target by the flanker ring is not enough to explain

the overall results. This supports the suggestion that the main driver of the increase in perceptual

error with increasing difference in depth is likely to be an increase in clutter due to a lack of

binocular fusion of stimulus areas outside Panum's Fusional Area. Even if the observer was not

consciously aware of diplopia due to the short presentation time. In addition, it is worth noting

that due to the lack of binocular fusion of stimulus areas outside Panum's Fusional Area, we

can't rule out the possible involvement of binocular rivalry which has been shown to interfere

with object recognition not just in isolation but also when combined with crowding (Kim et al.,

2013).

This current study is the first to investigate how crowding is affected by large, real

differences in depth between a target and flanking features. This is important because objects in

the real world are often flanked by visual features that vary greatly in their distance from an

observer, and many will fall outside the limits of Panum's Fusional Area. Our results show that

crowding from clutter outside Panum's Fusional Area can still have a significant impact on

object recognition and visual perception in the peripheral field. Our findings therefore do not

support the widely held view that depth differences between targets and flankers always reduce

crowding, instead the effect of depth on crowding appears to be more complex than originally

assumed. Our results, alongside those of previous studies, suggest that small differences in depth

between targets and flankers do not have the same effect on crowding as larger differences in

depth in which either the target or flankers are presented outside the limits of Panum's Fusional

Area. We suggest that large differences in depth between target and flankers increase perceptual

error due to an increase in visual features because of a lack of binocular fusion of stimulus areas

outside Panum's Fusional Area, even if diplopia is not consciously experienced. This finding

confirms the importance of crowding in natural three-dimensional scenes.

Acknowledgements

We thank members of the Bex lab and Eskew lab at Northeastern University for valuable

feedback during pilot experiments. We also thank MiYoung Kwon and Jan Skerswetat for

helpful discussions. This work was supported by NIH grant R01EY032162 (P.J.B.). The

Northeastern University IRB approved procedures (IRB #14-09-16), and all observers consented

to participation.

Author Contributions

568   S.P.S. and P.J.B. conceived the study and the idea of the multi-depth plane display and designed

569   the experiments. S.P.S. designed and built the multi-depth plane display, wrote all code, and

570   performed analyses. S.P.S., Y.S., and J.A. conducted the experiments. S.P.S. and P.J.B. wrote the

571   manuscript with input from all authors.

Data Availability

All data and code required to replicate the analyses in this paper will be made available on an online repository at the stage of publication. It will also be made available to reviewers during peer review.

References

Astle AT, McGovern DP, McGraw P V. 2014. Characterizing the role of disparity information in alleviating visual crowding. *J Vis* **14**:1–14. doi:10.1167/14.6.8

Balas B, Nakano L, Rosenholtz R. 2009. A summary-statistic representation in peripheral vision explains visual crowding. *J Vis* **9**:1–18. doi:10.1167/9.12.1

Banks WP, Larson DW, Prinzmetal W. 1979. Asymmetry of visual interference. *Percept Psychophys* **25**:447–456. doi:10.3758/BF03213822

Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *J Stat Softw* **67**:1–48. doi:10.18637/jss.v067.i01

Berens P. 2009. CircStat: a MATLAB toolbox for circular statistics. *J Stat Softw* **31**:1–29.

Bouma H. 1970. Interaction effects in parafoveal letter recognition. *Nature* **226**:177–178.

Coates DR, Ludowici CJH, Chung STL. 2021. The generality of the critical spacing for crowded optotypes: from Bouma to the 21st century. *J Vis* **21**:1–22. doi:10.1167/jov.21.11.18

Eberhardt LV, Huckauf A. 2019. Crowding in depth for binocular and monocular observation. *Attention, Perception, Psychophys* **81**:1951–1961.

Eberhardt LV, Huckauf A. 2017. Examining crowding using a real three-dimensional experimental setup. *Proc Latv Acad Sci* **71**:313–319. doi:10.1515/prolas-2017-0055

Eberhardt LV, Pittino F, Huckauf A. 2021. Close - but not distant - conditioned flanker emotion

affects crowding. *J Vis* **21**:1–14. doi:10.1167/jov.21.8.22

Eberhardt L V, Huckauf A. 2020. Crowding effects across depth are fixation-centered for

defocused flankers and observer-centered for defocused targets. *Brian Sci* **10**:596.

doi:10.3390/brainsci10090596

Ester EF, Klee D, Awh E. 2014. Visual crowding cannot be wholly explained by feature pooling.

J Exp Psychol Hum Percept Perform **40**:1022–1033. doi:10.1037/a0035377

Ester EF, Zilber E, Serences JT. 2015. Substitution and pooling in visual crowding induced by

similar and dissimilar distractors. *J Vis* **15**:1–12. doi:10.1167/15.1.4

Felisberti FM, Solomon JA, Morgan MJ. 2005. The role of target salience in crowding.

Perception **34**:823–833. doi:10.1068/p5206

Freeman J, Simoncelli EP. 2011. Metamers of the ventral stream. *Nat Neurosci* **14**:1195–1204.

doi:10.1038/nn.2889

Georgeson MA, Wallis SA. 2014. Binocular fusion, suppression and diplopia for blurred edges.

Ophthalmic Physiol Opt **34**:163–185. doi:10.1111/opo.12108

Hampton DR, Kertesz AE. 1983. The extent of Panum's area and the human cortical

magnification factor. *Perception* **12**:161–165. doi:10.1068/p120161

Harrison WJ, Bex PJ. 2017. Visual crowding is a combination of an increase of positional

uncertainty, source confusion, and featural averaging. *Sci Rep* **7**:45551.

doi:10.1038/srep45551

Harrison WJ, Bex PJ. 2016. Ungrouping overcrowded explanations of visual crowding: reply to

Pachai et al. *Curr Biol* **26**:R353–R354. doi:10.1016/j.cub.2016.03.024

Harrison WJ, Bex PJ. 2015. A unifying model of orientation crowding in peripheral vision. *Curr*

Biol **25**:3213–3219. doi:10.1016/j.cub.2015.10.052

Hartig F. 2022. DHARMa: Residual diagnostics for hierarchical (multi-level/mixed) regression

models. R package version 0.4.6. http://florianhartig.github.io/DHARMa/

He S, Cavanagh P, Lntriligator J. 1996. Attentional resolution and the locus of visual awareness.

Nature **383**:334–337. doi:10.1038/258748a0

Herzog MH, Manassi M. 2015. Uncorking the bottleneck of crowding: a fresh look at object

recognition. *Curr Opin Behav Sci* **1**:86–93. doi:10.1016/j.cobeha.2014.10.006

Herzog MH, Sayim B, Chicherov V, Manassi M. 2015. Crowding, grouping, and object

recognition: a matter of appearance. *J Vis* **15**:1–18. doi:10.1167/15.6.5

Hoffman DM, Girshick AR, Akeley K, Banks MS. 2008. Vergence-accommodation conflicts

hinder visual performance and cause visual fatigue. *J Vis* **8**:1–30. doi:10.1167/8.3.33

Hsu CW, Zhen B, Qiu W, Shapira O, DeLacy BG, Joannopoulos JD, Soljačić M. 2014.

Transparent displays enabled by resonant nanoparticle scattering. *Nat Commun* **5**:1–6.

doi:10.1038/ncomms4152

Kalpadakis-Smith A V., Goffaux V, Greenwood JA. 2018. Crowding for faces is determined by

visual (not holistic) similarity: Evidence from judgements of eye position. *Sci Rep* **8**:12556.

doi:10.1038/s41598-018-30900-0

Keshvari S, Rosenholtz R. 2016. Pooling of continuous features provides a unifying account of

crowding. *J Vis* **16**:1–15. doi:10.1167/16.3.39

Kim S, Shin E, Chong SC. 2013. When crowding meets binocular rivalry: challenges for object

perception. *Vision Res* **76**:134–143. doi:10.1016/j.visres.2012.10.020

Kooi FL, Toet A, Tripathy SP, Levi DM. 1994. The effect of similarity and duration on spatial

interaction in peripheral vision. *Spat Vis* **8**:255–279. doi:10.1163/156856894X00350

Legge GE. 2007. Psychophysics of reading in normal and low vision. Mahwah, NJ: Lawrence

Erlbaum Associates Publishers.

Levi DM. 2008. Crowding - an essential bottleneck for object recognition: a mini-review. *Vision*

Res **48**:635–654. doi:10.1016/j.visres.2007.12.009

Louie EG, Bressler DW, Whitney D. 2007. Holistic crowding: selective interference between

configural representations of faces in crowded scenes. *J Vis* **7**:1–11. doi:10.1167/7.2.24

Maiello G, Chessa M, Bex PJ, Solari F. 2020. Near-optimal combination of disparity across a

log-polar scaled visual field. *PLoS Comput Biol* **16**:1–28. doi:10.1371/journal.pcbi.1007699

Manassi M, Whitney D. 2018. Multi-level crowding and the paradox of object recognition in

clutter. *Curr Biol* **28**:R127–R133. doi:10.1016/j.cub.2017.12.051

Nandy AS, Tjan BS. 2012. Saccade-confounded image statistics explain visual crowding. *Nat*

Neurosci **15**:463–469. doi:10.1038/nn.3021

Nityananda V, Read JCAA. 2017. Stereopsis in animals: evolution, function and mechanisms. *J*

Exp Biol **220**:2502–2512. doi:10.1242/jeb.143883

Norcia AM, Suiter EE, Tyler CW. 1985. Electrophysiological evidence for the existence of

coarse and fine disparity mechanisms in human. *Vision Res* **25**:1603–1611.

doi:10.1016/0042-6989(85)90130-0

Ono H, Angus R, Gregor P. 1977. Binocular single vision achieved by fusion and suppression.

Percept Psychophys **21**:513–521. doi:10.3758/BF03198731

Pachai M V., Doerig AC, Herzog MH. 2016. How best to unify crowding? *Curr Biol* **26**:R352–

R353. doi:10.1016/j.cub.2016.03.003

Panum PL. 1858. Physiologische untersuchungen über das sehen mit zwei augen. Kiel:

Schwerssche Buchhandlung.

Parkes L, Lund J, Angelucci A, Solomon JA, Morgan M. 2001. Compulsory averaging of

crowded orientation signals in human vision. *Nat Neurosci* **4**:739–744. doi:10.1038/89532

Pelli DG. 2008. Crowding: a cortical constraint on object recognition. *Curr Opin Neurobiol* **18**:445–451. doi:10.1016/j.conb.2008.09.008

Pulliam K. 1981. Spatial frequency analysis of three-dimensional vision. *Proc SPIE Vis Simul Image Realis II* **303**:71–77. doi:https://doi.org/10.1117/12.932660

Qin D, Takamatsu M, Nakashima Y. 2006. Disparity limit for binocular fusion in fovea. *Opt Rev* **13**:34–38. doi:10.1007/s10043-006-0034-5

R Core Team. 2022. R: a language and environment for statistical computing.

Reynaud A, Hess RF. 2017. Characterization of spatial frequency channels underlying disparity sensitivity by factor analysis of population data. *Front Comput Neurosci* **11**:63. doi:10.3389/fncom.2017.00063

Sayim B, Westheimer G, Herzog MH. 2008. Contrast polarity, chromaticity, and stereoscopic depth modulate contextual interactions in vernier acuity. *J Vis* **8**:1–9. doi:10.1167/8.8.12

Spiegel DP, Baldwin AS, Hess RF. 2016. The relationship between fusion, suppression, and diplopia in normal and amblyopic vision. *Investig Ophthalmol Vis Sci* **57**:5810–5817. doi:10.1167/iovs.16-20438

Stidwill D, Fletcher R. 2010. Diplopia and confusion, suppression and rivalryNormal Binocular Vision: Theory, Investigation and Practical Aspects. Chicester: John Wiley & Sons, Incorporated. pp. 57–71. doi:10.1002/9781118788684.ch5

Strasburger H, Rentschler I, Jüttner M. 2011. Peripheral vision and pattern recognition: a review. *J Vis* **11**:1–82. doi:10.1167/11.5.13.

Thaler L, Schütz AC, Goodale MA, Gegenfurtner KR. 2013. What is the best fixation target? The effect of target shape on stability of fixational eye movements. *Vision Res* **76**:31–42.

doi:10.1016/j.visres.2012.10.012

Tyler CW, Barghout L, Kontsevich LL. 1994. Computational reconstruction of the mechanisms

of human stereopsis. *Proc SPIE Comput Vis Based Neurobiol* **2054**:52–68.

Ur Rehman A, Kihara K, Matsumoto A, Ohtsuka S. 2015. Attentive tracking of moving objects

in real 3D space. *Vision Res* **109**:1–10. doi:10.1016/j.visres.2015.02.004

Viswanathan L, Mingolla E. 2002. Dynamics of attention in depth: Evidence from multi-element

tracking. *Perception* **31**:1415–1437. doi:10.1068/p3432

Wallis TSA, Bex PJ. 2012. Image correlates of crowding in natural scenes. *J Vis* **12**:1–19.

doi:10.1167/12.7.6

Watt SJ, Akeley K, Ernst MO, Banks MS. 2005. Focus cues affect perceived depth. *J Vis* **5**:834–

862. doi:10.1167/5.10.7

Westheimer G. 1979. Cooperative neural processes involved in stereoscopic acuity. *Exp Brain*

Res **36**:585–597. doi:10.1007/BF00238525

Whitney D, Levi DM. 2011. Visual Crowding: a fundamental limit on conscious perception and

object recognition. *Trends Cogn Sci* **15**:160–168. doi:10.1016/j.tics.2011.02.005.Visual

Wilcox LM, Allison RS. 2009. Coarse-fine dichotomies in human stereopsis. *Vision Res*

49:2653–2665. doi:10.1016/j.visres.2009.06.004

Xia Y, Manassi M, Nakayama K, Zipser K, Whitney D. 2020. Visual crowding in driving. *J Vis*

20:1–7. doi:10.1167/jov.20.6.1

Yang Y, Blake R. 1991. Spatial frequency tuning of human stereopsis. *Vision Res* **31**:1176–1189.

doi:10.1016/0042-6989(91)90043-5