Genetics and resource availability shape divergence in life history

and behavior between locally-adapted populations of Atlantic mollies

(*Poecilia mexicana*, Poeciliidae)

John L. Coffin[1,2], Bethany L. Williams[3,4] & Michael Tobler[3,4,5]

[1]Division of Biology, Kansas State University, 104 Ackert Hall, Manhattan, KS 66506

[2]Current address: Washburn Rural High School, Topeka, KS 66619, USA

[3]Department of Biology, University of Missouri—St. Louis, St. Louis, MO 63121, USA

[4]Whitney R. Harris World Ecology Center, University of Missouri—St. Louis, St. Louis, MO, USA

[5]WildCare Institute, St. Louis Zoo, St. Louis, MO 63110, USA

Correspondence: tobler@umsl.edu

Abstract

Phenotypic variation is common along environmental gradients, but it is often unknown to what extent it results from genetic differentiation between populations or phenotypic plasticity. We studied populations of a livebearing fish that have colonized streams rich in toxic hydrogen sulfide (H_2S). In nature, there is strong phenotypic differentiation between adjacent sulfidic and nonsulfidic populations. In this study, we varied food availability to pregnant mothers from different populations to induce maternal effects, a form of plasticity, and repeatedly measured life-history and behavioral traits throughout the offspring's ontogeny. Genetic differentiation affected most of the traits we measured, as sulfidic offspring tended to be born larger, mature later, have lower burst swimming performance, be more exploratory, and feed less accurately. In contrast, maternal effects impacted few traits and at a smaller magnitude, even though offspring from poorly provisioned mothers tended to be born larger and be more exploratory. Population differences and maternal effects (when both were present) acted synergistically, and there was no evidence for population differences in plasticity. Overall, our study suggests that phenotypic divergence between these populations in nature is primarily caused by genetic differentiation, and that plasticity mediated by maternal effects accentuates—but does not cause—differences between populations.

Keywords: adaptation; extreme environments; hydrogen sulfide; maternal effects; resource availability; Poeciliidae

Introduction

Phenotypic variation is at the heart of evolutionary analyses because it links the cause (natural selection) to the consequence (genotypic change) of adaptive evolution (Lande & Arnold, 1983). We have known how inheritance causes resemblance between parents and their offspring for well over a century, reflecting a genetic component to phenotypic variation (Stenseth et al. 2022). However, trait variation can also be influenced by phenotypic plasticity, where a single genotype can give rise to alternate phenotypes in response to internal or environmental cues (West-Eberhard, 1989; Pigliucci, 2001). In addition, the environment experienced by parents can affect phenotypes of their offspring (*i.e.*, parental effects; Uller 2008; Badyaev and Uller 2009), representing a case of plasticity that spans generational boundaries. Phenotypic variation in nature can therefore arise from genetic differences among individuals, plasticity induced by individual exposure to different environmental conditions, plasticity induced by parental effects, and their interactions (Scheiner 1993; Dingemanse and Araya-Ajoy 2015). For many natural systems, we know little about the origins of phenotypic variation, even though it critically shapes our inference of adaptation in natural populations.

Plasticity induced by parental effects is particularly strong from mothers due to their higher reproductive investment and, in viviparous species, the physically intimate relationship with their developing young (Lindholm, Hunt, & Brooks, 2006; Wolf & Wade, 2009). Such maternal effects are widespread in nature (Mousseau and Fox 1998) and can impact trait expression and evolution (Rossiter, 1996; Wilson *et al.*, 2005; Beckerman *et al.*, 2006). Maternal effects can be adaptive if the expression of offspring traits is biased to match the environment experienced by the mother (Marshall and Uller 2007), or if mothers in good condition are able to endow phenotypes that provide a competitive advantage to their offspring in any environment (Grafen, 1988; Monaghan, 2008; Van Allen *et al.*, 2021). However, maternal effects can also be maladaptive and produce mismatches between offspring phenotype and environment, as documented in some organisms

responding to anthropogenic climate change (Schuler and Orrock 2012; Leonard and Lancaster

2020). Maladaptive maternal effects can also be related to stress, where physiological stress

responses in mothers have unintended negative side effects on offspring (MacLeod, While, & Uller,

2021). Regardless of whether maternal effects are adaptive, they are important biological phenomena

that warrant careful attention and explicit accounting in evolutionary analyses due to the non-genetic

effects on phenotypic expression.

Phenotypic variation in nature is common along environmental gradients, but it is often

unclear whether it is caused by genetic differentiation among populations or plastic effects that arise

from population-specific environmental exposure histories experienced by mothers or directly by

their offspring. For example, freshwater springs rich in toxic hydrogen sulfide (H_2S) in the Grijalva

River basin of southern Mexico are extreme environments that are connected to adjacent nontoxic

streams, and stark phenotypic gradients can be observed in fish occupying these habitats in as little

as a few meters. Sulfide springs are complex ecosystems with several correlated sources of selection

(Tobler *et al.*, 2016b). H_2S is toxic because it disrupts aerobic ATP production (Reiffenstein, Hulbert,

& Roth, 1992; Cooper & Brown, 2008), but habitats rich in H_2S also differ from nonsulfidic habitats

in other physical and chemical water parameters (dissolved oxygen concentrations, pH, and salinity),

and the presence of competitors and predators (Riesch *et al.*, 2010a; Tobler *et al.*, 2011; Greenway *et*

al., 2014). Resource availability also differs greatly between habitat types; fish in nonsulfidic

environments eat primarily algae and detritus, while fish in sulfidic environments have shifted to

eating primarily sulfide bacteria and invertebrates (Tobler *et al.*, 2015). Populations of Atlantic

mollies (*Poecilia mexicana*), a species of livebearing fish of the family Poeciliidae, have independently

colonized and adapted to sulfidic streams across multiple river drainages, and previous studies have

documented that colonization of sulfide springs has been associated with convergent changes in

morphology, locomotion, and respiration (Tobler & Hastings, 2011; Camarillo, Arias Rodriguez, &

Tobler, 2020), behavior (Plath *et al.*, 2007; Lukas *et al.*, 2021; Doran *et al.*, 2022), physiology (Tobler *et*

al., 2011; Barts *et al.*, 2018; Greenway *et al.*, 2020), and life history traits (Riesch *et al.*, 2011a; Riesch,

Plath, & Schlupp, 2011b; Riesch *et al.*, 2014). Phenotypic divergence between sulfidic and nonsulfidic

mollies likely has a significant genetic component, because it coincides with strong genetic

differentiation between populations, even though there are no physical barriers separating

populations in the different habitat types (Palacios *et al.*, 2013; Plath *et al.*, 2013; Riesch *et al.*, 2016b).

However, trait variation between populations likely also has an environmental component; while

population differentiation persists in captive populations reared in common garden conditions in the

lab (Tobler *et al.*, 2016a; Greenway *et al.*, 2020), there is also evidence for plasticity caused by short-

term exposure to different environmental conditions (Bierbach *et al.*, 2011; Passow *et al.*, 2017a). In

addition, the impact of maternal effects on offspring trait expression remains to be investigated in

these livebearing fish. Hence, we tested how genetic differentiation, maternal effects, and their

interactions shape phenotypic expression in *P. mexicana* populations from sulfidic and nonsulfidic

habitats. To induce maternal effects, we manipulated resource availability to pregnant mothers,

because natural populations vary substantially in nutritional state. Fish in sulfidic habitats are

consistently under food stress, exhibiting significantly reduced body condition (both when inferred

through length-weight regression and body fat content analysis; Plath *et al.*, 2005; Tobler *et al.*, 2006;

Tobler, 2008). Food stress arises as a consequence of constraints associated with resource

acquisition; because H_2S coincides with and exacerbates hypoxia, fish from sulfidic habitats have to

trade off performing aquatic surface respiration, a compensatory behavior to access better-

oxygenated surface waters, with benthic foraging (Tobler *et al.*, 2009). Accordingly, populations in

sulfide springs have adaptations to low resource availability, including reductions in routine

metabolic rates and energetically expensive tissues like the brain (Schulz-Mirbach *et al.*, 2016;

Passow, Arias-Rodriguez, & Tobler, 2017b). In other species, including some poeciliids, resource

availability experienced by mothers has been shown to impact offspring trait expression (Reznick,

Callahan, & Llauredo, 1996a; Altmann & Alberts, 2005; Boots & Roberts, 2012), and different

population histories in terms of exposure to food stress in *P. mexicana* may have caused changes in

resource-induced maternal effects.

To quantify population differentiation and maternal effects in populations of *P. mexicana*, we

followed families of offspring from birth to the onset of maturation and repeatedly quantified a host

of complex phenotypic traits. Focal traits included brood size, size at birth, and age at maturity as

well as ontogenetic trajectories in growth rates, burst swimming, exploratory behavior, and feeding

accuracy. We chose these traits because they likely affect fitness, and we have prior knowledge for

many of them from natural populations, providing us with a framework to make *a priori* predictions.

Specifically, our experiments sought to address four specific questions: (1) Is there evidence for

phenotypic trait differences between populations from sulfidic and nonsulfidic habitats that persist

in fish reared in a common-garden environment for multiple generations? Divergence in phenotypic

traits between populations regardless of maternal food treatments would indicate that trait

differentiation is due to genetic variation between populations. (2) Is there evidence for maternal

effects in response to resource availability? Significant differences in offspring traits between

maternal food treatments, irrespective of population of origin, would suggest resource-induced

maternal effects. (3) How do functional traits vary throughout ontogeny, and how do population

differences and maternal effects interact with ontogeny? Age is a major determinant in the

expression of many traits (Hegyi *et al.*, 2006; Zhang *et al.*, 2015), but how population differences and

maternal effects impact trait expression through ontogeny is less clear. In other poeciliids, maternal

effects tend to be present at birth and decline with age (Lindholm *et al.*, 2006). In contrast,

population differences between sulfidic and nonsulfidic *P. mexicana* are stark in adults, suggesting

differences may emerge early in life and even increase throughout ontogeny (Riesch *et al.*, 2011a).

Accordingly, we predicted that age would impact most of the traits measured, that population

differentiation would increase with age, and that maternal effects would diminish with age. (4) How

do population differences interact with maternal effects? A difference in how each population

responds to variation in maternal resource availability would indicate genotype-by-environment

interactions. Fish from sulfidic habitats generally face constraints in resource levels, while those

from nonsulfidic habitats have access to more abundant resources (Tobler *et al.*, 2006, 2009; Tobler,

2008). Hence, we predicted that trait variation induced by the low-food treatment would occur in

the same direction as trait variation produced by differences between the nonsulfidic and sulfidic

populations (i.e., maternal effects would be aligned with population differences). In this case,

maternal effects would accentuate divergence between populations that resembles patterns of

variation found in the wild. Alternatively, low-resource traits may be canalized in the sulfidic

population because sulfidic individuals are constantly food stressed in nature. In this case, maternal

effects may be weaker in the sulfidic population.

Methods

Experimental overview

For our experiments, we used two lab-reared populations of *Poecilia mexicana* originating from wild-

caught relatives in the Tacotalpa River drainage. One population originated from a sulfide spring

complex called El Azufre I (according to Plath *et al.*, 2013; hereafter referred to as the sulfidic

population or ecotype), and the other population was from a nonsulfidic stream connected to the

mainstem of the Tacotalpa River, called Arroyo Bonita (according to Plath *et al.*, 2013; hereafter

referred to as the nonsulfidic population or ecotype). Both populations were reared in 680-liter stock

tanks filled with filtered tap water. Tanks were fed *ad libitum* twice daily with commercial dry fish

food (Purina), and ~50% of the water was exchanged weekly. Mothers used in this experiment were

raised under common-garden conditions for at least three generations (*i.e.*, they were at least great-

grandchildren from individuals originally collected in the wild).

From each tank, 30 females were caught with a dipnet and isolated in a 20-liter tank with an

aerating filter and a bundle of plastic mesh as shelter for newborn fry. Females were fed twice daily,

once with aquatic gel diet for omnivorous fish (Mazuri) and once with freshly hatched *Artemia*

nauplii (Brine Shrimp Direct). Females were randomly assigned to either a "high-food" diet (0.32 ml

per feeding) or a "low-food" diet (0.08 ml per feeding), with equal numbers of females from each

population being assigned to each group (Figure 1). Each tank was checked daily for newborn fry.

By the end of the experiment, 18 sulfidic females gave birth (12 from the high food treatment and 6

from the low food treatment) and 22 nonsulfidic females produced a brood (12 from the high food

treatment and 10 from the low food treatment). The amount of time each female was in the food

treatment before giving birth varied (mean 28.2 days, range 0-99 days). As some females gave birth

during the acclimation period or immediately after beginning the food treatments, we included

treatment length as a potential covariate in all analytical models. Additionally, for a subset of models

we tested whether using females that were in the treatment for at least seven days would affect the

resulting top models (Table S1). Because the models were largely unaffected, we chose to include all

broods to increase statistical power to detect population differences throughout ontogeny.

Whenever a brood was born, we recorded the brood size, date of birth for each family, the

number of days that the female was in the food treatment, and the mother's standard length

(distance from the anterior tip of the snout to the posterior end of the caudal peduncle, mm).

Mothers were then removed from the tanks, and we randomly selected 15 newborn fry (if available)

from each family to remain in the experimental tanks to minimize density-dependent effects.

From that point on, we followed the developing families through ontogeny. All fry—

irrespective of the mother's food treatment—received the same amount of food; they were fed *ad*

libitum twice daily with a mixture of decapsulated brine shrimp eggs (Brine Shrimp Direct) and dry

food. We assessed offspring phenotypes at approximately weekly intervals by measuring life history

traits (size at birth, weekly growth rate, and age at maturity) and behavioral traits (burst swimming,

exploratory behavior, and feeding accuracy; see Figure 1). We specifically tested whether there were

differences in these phenotypes between maternal food treatments (*i.e.*, maternal effects), between

populations (*i.e.*, sulfidic *vs.* nonsulfidic, indicating effects of evolved population differences), or

both. All analyses were conducted in R 4.0.5 (R Core Team, 2023). Code and data to reproduce all

analyses can be found on GitHub (https://github.com/michitobler/common-garden).

Size at birth and growth rate

To measure size at birth, we photographed each family from above on the day of their birth with a

Nikon D90 digital camera fitted with an AF-S Micro NIKKOR 105mm f/2.8 lens. A ruler was

included in the background of each image. Images were imported to ImageJ v. 1.53 (Schneider,

Rasband, & Eliceiri, 2012) and calibrated by setting the scale. We measured the standard length

(mm) of each offspring in the family. These measurements were averaged across all individuals to

obtain a single mean size at birth for each family. This measurement was completed weekly, and the

measurement for each family was subtracted from that family's measurement from the week prior to

obtain an average growth rate [mm/week]. To account for allometric differences in growth rate, we

converted the average weekly growth rates to a proportional growth rate by dividing the average

weekly growth rate by the mean body size the week prior.

Age at maturity

We estimated the minimum age at maturity for each family using morphological characteristics of

male sexual maturity. While the sexes of juvenile livebearers are difficult to distinguish, at the onset

of sexual maturity, the male anal fin is modified into an intromittent organ (gonopodium), while it

remains unmodified in females (Rosen & Gordon, 1953; Chambers, 1987). We therefore measured

minimum age at maturity (days) as the time it took to for the first male in a group to develop its

complete gonopodium as judged by the presence of a fleshy palp on anal fin ray 3.

Burst swimming

Most fish avoid predation with a highly conserved, reflexive escape response that causes the head to

move away from the stimulus, bending the body into a 'C' shape (Eaton, Bombardieri, & Meyer,

1977). Then, a strong stroke of the caudal fin starts the movement away from the stimulus

(Domenici & Blake, 1997). This process (bending into a 'C' and propelling away from the stimulus)

is known as a C-start response and is frequently used as a metric of escape performance in fish

(Walker, 1997; Ghalambor, Reznick, & Walker, 2004; Langerhans *et al.*, 2004; Camarillo *et al.*, 2020).

To quantify this burst swimming behavior, we adopted the methods and metrics used by prior

studies (Langerhans *et al.*, 2004; Ingley *et al.*, 2016; Camarillo *et al.*, 2020). We placed a haphazardly

chosen individual from each family in a glass petri dish (9 cm in diameter, containing 2 cm of water)

with opaque sides suspended above an angled mirror, providing a ventral view of each fish. After 5

min of acclimation, we struck the surface of the water within a body length of the fish with a probe

and recorded the movement of the fish from below with a Sony NEX-FS700R camcorder at 60

frames per second (fps) and 1080p resolution. We converted the resulting .mts files into .mp4 files

(to enhance compatibility with downstream applications) with FFmpeg (Tomar, 2006).

We used DLTdv8 (Hedrick, 2008) to digitize the 2-D location of the isthmus (*i.e.*, the area

on the ventral surface of the head where the opercula converge) of the fish in each frame. Digitized

points were then used to calculate the maximum velocity (v_{max}, mm/s), maximum acceleration (a_{max},

mm/s^2), and net distance traveled (d_{net}, mm of displacement within 1/12th of a second after the C-

start). To calculate v_{max}, we calculated the straight-line distance between each pair of successive

digitized points, divided this distance by the inverse of the frame rate (60 fps), and found the

maximum value between any two points. a_{max} was calculated by subtracting the velocity value at each

point from the velocity value at the point immediately preceding it and finding the maximum value.

d_{net} was calculated by recording the 2-D position immediately after the fish ends the C-start with a

single stroke of the caudal fin, and then recording the 2-D position $1/12^{th}$ of a second later (5

frames later) and calculating the straight-line distance between the two points. To reduce the

dimensionality of this dataset, we conducted a principal component analysis (PCA) using the

prcomp() function with a correlation matrix. There was one principal component with an eigenvalue

greater than 1 (explaining 85.6 % of the total variance), which was retained as a compound metric of

burst swimming performance. Positive scores along this principal component axis were associated

with higher velocity, acceleration, and distance traveled (Table S2A). All mathematical operations

were conducted using packages contained in the base distribution of R.

Exploratory behavior

We used an open field test to quantify the exploratory tendencies of fry. We filled a Styrofoam cup

(9 cm in diameter) with 3 cm of water and covered the arena with a sheet of glass. We haphazardly

selected one individual from each family and placed it in the arena undisturbed for a 5-min

acclimation period, after which we recorded 5 min of video from above with a GoPro Hero 4

(1080p resolution, linear field of view, 30 fps).

We used idTracker (v. 2.1, bundled with 64-bit Matlab Compiler Runtime 8.3; Pérez-

Escudero *et al.*, 2014) to track the 2-D location of the fish automatically for the entire 5-min

recording. We set the number of individuals to 1 and manually determined the intensity threshold

(0.5–0.8) and minimum size (40–250 pixels) for each video. We also imported a still frame from each

video into ImageJ to measure the centroid coordinates and arena radius.

Using the 2-D coordinates in each frame, we calculated several metrics of motion that we

used as proxies for exploratory behavior. We calculated distance traveled between each pair of

successive points, the velocity, the acceleration, and the total cumulative distance traveled (d_{total},

mm), as described above. We also calculated average velocity (v_{avg}, mm/s), maximum velocity (v_{max},

mm/s), and maximum acceleration (a_{max}, mm/s^2) by finding the means and maxima of all velocity

and acceleration values. Finally, we calculated the proportional average distance from the center of

the arena (d_{center}, dimensionless) by calculating the distance from the fish's location to the centroid of

the arena and dividing this value by the arena radius. Videos were excluded if the fish was

completely still in all frames. To reduce the dimensionality of and observe the correlation structure

within this dataset, we ran a PCA on v_{avg}, v_{max}, a_{max}, d_{total}, and d_{center} with a correlation matrix. We

retained scores along the first PC axis (explaining 56.9 % of the total variance) as a composite

exploratory behavior score. As shown in Table S2B, higher PC1 scores were associated with more

exploratory behavior (positive loadings for all variables).

Feeding accuracy

To measure feeding accuracy, we withheld food from individuals of each tank for 24 h prior to the

experiment and placed one haphazardly selected fry in a viewing tank for a 5-min acclimation

period. The viewing tank was constructed by connecting two 10 cm × 10 cm sheets of glass to either

side of 1 cm diameter glass rods with silicone. The rear wall of the viewing tank was covered with a

black sheet of plastic to enhance contrast between the background and the fish in the tank. The

feeding solution consisted of 1 g of freshly hatched, live *Artemia* nauplii diluted into 100 ml of

filtered tap water. After 5-min of acclimation, we added 0.08 ml of the feeding solution to the

viewing tank and recorded 5 min of video with the camcorder. We analyzed the video frame-by-

frame using BORIS v. 7.13 (Friard & Gamba, 2016) and recorded the number of successful strikes

(a feeding strike that ends in consumption of the food item) and unsuccessful strikes (a feeding

strike that either misses the food item or results in regurgitation of the food item). We divided the

number of successful strikes by the total number of strikes to get an overall proportional estimate of

feeding accuracy, which was arcsine-square-root transformed prior to analysis.

Statistical analyses of individual traits

There were many potential sources of variation in our experiment. Other than the effects of interest

for our study (population, maternal food treatment, and their interaction), the observed variation in

traits could also have arisen from differences in fry age, maternal body size (standard length), the

duration of her food treatment, and brood size. Consequently, we used a model selection approach

to find the models that were best supported by our data for each experiment separately. For each

phenotype, we created a global model that contained all possible effects. For phenotypes that were

only measured once for each family (size at birth, brood size, and age at maturity), we used a general

linear model using the lm() function from the STATS package v. 3.6.2, and for phenotypes that were

measured repeatedly through development (growth rate, burst swimming, exploratory behavior, and

feeding accuracy), we used a linear mixed model implemented with the lmer() function from the

LME4 package v. 1.1-26 (Bates *et al.*, 2015) that included 'family' as a random effect. We then used

the dredge() function from the MUMIN package v. 1.47.1 (Bartón 2009) to create a model selection

table based on the effects contained in the global model, with different models ranked and weighted

based on AIC_c (Johnson & Omland, 2004). Full model selection tables are available for each

phenotype in Table S3. To avoid overfitting, we limited the models to a maximum of four terms. We

chose the top-supported model for each trait and analyzed it with either type-III analysis of variance

(ANOVA) or Wald's chi-square tests (see Table 1). Quantification and visualization of effects was

accomplished by calculating and plotting estimated marginal means for the effects of 'Population'

and/or 'Food Treatment', depending on the best-supported model, using the Effect() function from

the EFFECTS package v. 4.2-2 (Fox & Weisberg, 2018a,b). To aid in drawing inferences from our

models, we generated 95% confidence intervals (CI) for model coefficients in our top models using

the confint() function form the base R distribution. Effect sizes were calculated as partial eta-

squared (η_p^2), which represents the proportion of variance explained by a particular variable after

accounting for the variance explained by all other variables. We calculated η_p^2 with the etasq()

function (Fox et al. 2021) for general linear models or the eta_squared() function (Ben-Shachar,

Lüdecke, & Makowski, 2020) for linear mixed models.

Multivariate analysis

Since selection ultimately acts on complex, multivariate phenotypes (Lande & Arnold, 1983), we

sought to understand how the traits measured for our analyses vary and covary to jointly shape

multivariate phenotypes. To do so, we averaged each phenotype across all ages for each tank. We

used this approach rather than including age as a covariate and analyzing raw phenotypic scores

because of timing and logistical constraints that made it impossible to conduct each experiment on

offspring that were exactly the same age. We analyzed the averaged phenotypes with a PCA

(correlation matrix) and used scores along the first two principal components—which had

eigenvalues greater than 1—as dependent variables. We analyzed principal component scores along

each axis separately because the axes, by definition, were orthogonal. For each axis, we created a

global model containing all possible effects and selected the best-supported model based on AIC$_c$, as

explained above.

Results

We measured seven functional traits in offspring from a sulfidic and a nonsulfidic population of *P.*

mexicana throughout ontogeny. Model selection tables for the analysis of each trait at birth and

separately across all ages can be found in Table S3, and the best-supported models are summarized

in Table 1. For brevity, we will present results in the context of our hypotheses outlined in the

Introduction, focusing on how all traits vary through ontogeny in terms of population

differentiation, maternal effects, and their interaction, rather than presenting the results for each trait

separately.

Is there evidence for population differentiation?

We found evidence for significant population differences in five of the seven traits measured: size at

birth, age at maturity, burst swimming, exploratory behavior, and feeding accuracy (but not brood

size or growth rate). Age at maturity, burst swimming, and feeding accuracy were significantly

different between populations, but not between food treatments, whereas there were significant

effects of 'Population' and 'Food Treatment' (but no interactions) on size at birth and exploratory

behavior (see Table 1 and below). Sulfidic individuals were born 9.2 % larger [$F_{1,35}$ = 20.032; $P <$

0.001; estimated marginal mean for sulfidic fish (EMM$_S$) = 10.03 mm, estimated marginal mean for

nonsulfidic fish (EMM$_{NS}$) = 9.18 mm; η_p^2 = 0.36; Figure 2A], matured an average of 10.5 days later

($F_{1,26}$ = 4.420; P = 0.045; EMM$_S$ = 54.2 days, EMM$_{NS}$ = 43.7 days; η_p^2 = 0.15), had significantly

lower burst swimming performance (χ_1^2 = 4.308; P = 0.038; CI: –0.94, –0.03; η_p^2 = 0.03), were more

exploratory (χ_1^2 = 7.513; P = 0.006; CI: 0.29, 1.70; η_p^2 = 0.28), and were 7.2% less successful during

feeding (χ_1^2 = 4.645; P = 0.031; EMM$_S$ = 0.72, EMM$_{NS}$ = 0.80; η_p^2 = 0.04; Figure 3).

Is there evidence for maternal effects in response to resource availability?

To test whether maternal effects induced by resource availability during pregnancy impact functional

traits in offspring, we compared each phenotype between offspring born to mothers who

experienced a high-food environment and mothers who experienced a low-food environment.

Mothers in low-food treatments produced offspring that were 3.4% larger at birth [$F_{1,35}$ = 4.354; P =

0.044; estimated marginal mean for fish in the low-food treatment (EMM_{low}) = 9.8 mm, estimated

marginal mean for fish in the high-food treatment (EMM_{high}) = 9.4 mm; η_p^2 = 0.11; Figure 2A] and

were more exploratory (χ_1^2 = 4.245; P = 0.039; CI: –0.04, –1.42; η_p^2 = 0.18; Figure 3E). For all

remaining traits (brood size, age at maturity, growth rate, burst swimming, and feeding accuracy),

'Food Treatment' was not included in the best-supported model (see Table 1), suggesting that

maternal effects did not affect the expression of those traits.

How do population differences and maternal effects interact with ontogeny?

To determine how traits changed through offspring development, we compared each phenotype

across age groups. 'Age' was a significant predictor of growth rate and burst swimming, indicating

that these traits changed throughout ontogeny, while the other phenotypes were not affected by age.

Across populations and food treatments, fry grew at a slower relative rate (χ_1^2 = 147.91; P < 0.001,

CI: –0.010, –0.007; η_p^2 = 0.50) and performed better in burst swimming trials (χ_1^2 = 30.178; P <

0.001; CI: 0.024, 0.049; η_p^2 = 0.16) as they got older (Table 1). Burst swimming scores were also

lower in sulfidic individuals (see above), but the interaction term 'Population × Age' was not

significant, demonstrating that both populations exhibited similar changes in burst swimming

throughout ontogeny.

To ensure that signals of population differentiation and maternal effects occurring at birth

were not obscured by measurements later in life, we also subset our dataset for each phenotype to

obtain only the earliest data point for each family. As mentioned above, fry from the sulfidic

population and the low-food maternal treatment had significantly larger size at birth (Figure 2; Table

1). However, there were no population differences or maternal effects on brood size, which was

higher in larger mothers (Mother SL; $F_{1,37}$ = 23.342; P < 0.001; CI: 0.73, 1.79; η_p^2 = 0.38) and

mothers who spent longer in the food treatment (Treatment Length; $F_{1,37}$ = 8.926; P = 0.005; CI:

0.07, 0.40; η_p^2 = 0.19). Growth rate at birth was significantly lower in sulfidic fry ($F_{1,37}$ = 5.025; P =

0.032; CI: –0.12, –0.006; η_p^2 = 0.12; Table 1; Figure 3A), but there was no evidence for maternal

effects on growth rate at birth ('Food Treatment' was not included in best-supported model). Note

that the population difference in growth rate at birth disappeared as fry developed (Figure 3B). All

other traits that were measured throughout ontogeny (burst swimming, exploratory behavior, and

feeding accuracy) did not exhibit significant population differences or maternal effects at birth.

Interestingly, the maternal effect that we detected on exploratory behavior across all ages (see above)

was not evident at birth ($F_{1,31}$ = 2.727; P = 0.108; CI: –0.29, 2.81). These results collectively

demonstrate that—contrary to our hypothesis regarding ontogenetic variation of functional traits—

population differentiation did not increase with age, and maternal effects were not always observable

at birth, nor did they decline with age.

How do population differences interact with maternal effects?

We hypothesized that variation from maternal effects would be aligned with population differences,

but that there would be a significant interaction between population differences and maternal effects

(*i.e.*, different magnitudes of maternal effects in each population due to different evolutionary

histories associated with resource stress). Contrary to our predictions, no 'Population × Food

Treatment' interactions were included in the best-supported model for any of the traits we measured

(Table 1), suggesting a general lack of support for significant interactions between population

differences and maternal effects. Therefore, maternal effects—if present—were similar in direction

and magnitude between populations.

To further address our hypothesis, we asked whether population differences and maternal

effects were, in fact, aligned, and whether they explained a similar proportion of phenotypic variance

when they acted in unison. We compared the signs (positive *vs.* negative coefficient estimates) and

effect sizes (η_p^2) of population differences and maternal effects for the two traits with evidence of

both effects simultaneously impacting trait expression—size at birth and exploratory behavior. For

both traits, the difference between the nonsulfidic and sulfidic population occurred in the same

direction as the trait shifts between the high- and low-food treatments, indicating that the effects

were aligned (Figures 2A and 3E). Effect size estimates for 'Population' and 'Food Treatment' for

both traits indicated that 'Population' had a larger effect on trait expression than 'Food Treatment'

($\eta_\mathrm{p}^2 = 0.36$ *vs.* 0.11 for size at birth and 0.28 *vs.* 0.18 for exploratory behavior).

Multivariate analysis

In addition to the univariate analyses of trait variation, we were also interested in understanding how

the traits covaried with one another, and whether and how multivariate phenotypes were impacted

by population differences and maternal effects. We averaged each phenotype across ages for each

family, conducted a PCA, and then analyzed principal component scores along the first two

principal component axes. The first principal component accounted for 28.5% of variance in

multivariate phenotypes, and scores along the first principal component were positively correlated

with brood size, growth rate, and feeding accuracy, and they were negatively correlated with size at

birth, age at maturity, burst swimming, and exploratory behavior (see Table S2C). The second

principal component explained 21.1% of variance and was positively correlated with burst

swimming and size at birth and negatively correlated with brood size, growth rate, feeding accuracy,

exploratory behavior, and age at maturity (Table S2). After conducting model selection, the best-

supported model for PC1 contained the main effects of 'Population', 'Food Treatment', and

'Treatment Length', and the best-supported model for PC2 contained the main effects of

'Population', 'Mother SL', and 'Treatment Length'.

Sulfidic offspring had significantly lower scores along PC1 ($F_{1,24}$ = 16.266; P < 0.001; CI: –

2.40, –0.78; η_p^2 = 0.40), as did offspring from low-food mothers ($F_{1,24}$ = 4.746; P = 0.039; CI: –1.59,

–0.04), indicating population differences and maternal effects on PC1 scores. As for the univariate

analyses, maternal effects were aligned with population differences in PC1 scores (Figure 4),

'Population' had a larger effect than 'Food Treatment' (η_p^2 for 'Population' = 0.40; η_p^2 for 'Food

Treatment' = 0.17), and there was no evidence for a significant 'Population $\times$ Food Treatment'

interaction (the term was not included in best-supported model). We also found that sulfidic

individuals exhibited marginally lower PC2 scores ($F_{1,24}$ = 3.321; P = 0.081; CI: –1.45, 0.09; η_p^2 =

0.12) but detected no evidence for maternal effects (the 'Food Treatment' term was not included in

the best-supported model) on PC2.

Discussion

Phenotypic variation can be shaped by multiple genetic and nongenetic factors, but the interplay of

genes and environmental effects is rarely disentangled in natural systems, even though it

fundamentally impacts our inference of adaptation. We examined how genetic variation (*i.e.*,

population differences), phenotypic plasticity mediated by maternal effects, and their interactions

shape trait expression in two populations of *Poecilia mexicana* that are exposed to strong divergent

selection in nature. We found significant trait differences between populations even though fish

were housed under common-garden conditions in the laboratory for at least three generations. In

contrast, exposure of mothers to different food treatments impacted relatively few traits in their

offspring and, if they occurred, had weaker effects than the population differences. In addition,

maternal effects tended to be aligned with population differences and act in the same way across

populations. Principal component analysis supported these overall conclusions in multivariate trait

space. We also found no evidence for interactions between populations and food treatments,

suggesting that while populations have diverged in phenotypic traits, they have retained similar

maternal influences on those same traits. Overall, we found that the stark phenotypic differences

between populations of *P. mexicana* that are evident in nature are largely a consequence of genetic

divergence, likely representing local adaptation to the distinct ecological conditions of their habitats.

Maternal effects in response to resource availability, while present in some traits, appear to

accentuate population differences, but not cause them.

Trait variation across populations and maternal food treatments

There is a rich history integrating field-based studies that quantify trait variation of poeciliid fishes in

nature with laboratory-based studies that isolate causative environmental factors (Endler, 1980,

1995; Reznick & Bryga, 1987; Reznick, Bryga, & Endler, 1990; Langerhans, Gifford, & Joseph, 2007;

Tobler *et al.*, 2008; Ghalambor *et al.*, 2015; Ingley & Johnson, 2016). While most studies focus on

single or a few related traits and take a snapshot at a single ontogenetic stage, our study

demonstrated that multiple complex trait differences quantified in common-garden conditions—

across food treatments and across ontogeny—closely mirror trait differences between populations in

nature.

First, our study corroborates a genetic basis for population divergence in reproductive life

history traits. We documented significant population divergence in size at birth, with sulfidic mollies

giving birth to larger offspring (see Figures 2A and 4). This finding is consistent with life history

studies of sulfide spring populations in *P. mexicana* and other poeciliid species (Riesch *et al.*, 2010b;

Riesch, Plath, & Schlupp, 2010c; Riesch *et al.*, 2014), suggesting that divergence in offspring size

between sulfidic and nonsulfidic populations is a repeatable evolutionary consequence of sulfide

spring colonization that is driven by natural selection for larger offspring size. Among-population

variation in size at birth has also been documented in other poeciliids that are exposed to other

sources of divergent selection, including variation in predation (Weeks & Gaggiotti, 1993; Reznick,

Rodd, & Cardenas, 1996b; Schrader & Travis, 2012; Belk, Ingley, & Johnson, 2020). Our

multivariate results (Figure 4) also indicated that, like other poeciliids, the nonsulfidic population of

P. mexicana closely resembles an opportunistic life history strategy (Winemiller & Rose, 1992), which

places a premium on earlier maturity and higher fecundity, at the expense of lower juvenile

survivorship. The population differences we found in the sulfidic population may represent a shift in

the direction of the equilibrium strategy, which is energetically costly to parents but should

ultimately benefit offspring competitive ability. In natural populations, maternal effects induced by

variation in resource availability likely accentuate genetic differences in offspring size across sulfidic

and nonsulfidic habitats. This pattern was also evident across a plethora of life history traits in

guppies; in all ten life history traits with evidence for significant genetic divergence and maternal

effects, these effects always occurred in the same direction (Felmy *et al.*, 2022).

Second, we found patterns of population differentiation in age at maturity that match with

previous work in livebearing fishes. Prior studies have demonstrated that guppies in streams with

high predation pressure on adults mature earlier than conspecifics from low-predation populations

(Reznick & Endler, 1982). Several studies have also documented genetic differences in age at

maturity between populations of *P. mexicana*. For example, cave mollies mature significantly later

than surface populations, and there was also significant plasticity in age at maturity due to resource

availability (Riesch *et al.*, 2016a). Another study of surface mollies that compared size at maturity in

sulfidic and nonsulfidic populations found that sulfidic individuals reached maturity at a significantly

smaller size than nonsulfidic individuals (Riesch *et al.*, 2011b), likely because energy limitation

imposed by constraints associated with compensatory behaviors in sulfidic habitats (Tobler *et al.*,

2009) selects for reduced growth rates and smaller size at maturity (Passow *et al.*, 2017b). Since

poeciliid males exhibit determinate growth that ends at maturity (Stearns & Sage, 1980), size at

maturity should closely mirror age at maturity, assuming growth rates are the same as we found in

our study. However, we found that sulfidic mollies matured at a significantly later age than

nonsulfidic individuals, despite identical growth rates across ontogeny (Figure 3B). This discrepancy

between our findings and those of previous studies suggests that plasticity in age at maturity may be

strong and that variation in experimental design and rearing conditions matters.

Third, we found significant behavioral differences between populations from sulfidic and

nonsulfidic habitats, including exploratory behavior, feeding accuracy, and burst swimming. Previous

work has shown that nonsulfidic mollies—and ones that were better fed—were bolder in their

natural habitat, but behavioral differences also disappeared in the laboratory (Riesch *et al.*, 2009).

While we did not measure boldness *per se*, exploratory behavior as measured in our experiment (*i.e.*,

activity levels in a novel arena) is often characterized as part of a behavioral syndrome that correlates

with boldness (Conrad *et al.*, 2011). Unlike previous work, our study found that individuals from the

sulfidic population and the low-food treatment were more exploratory (Table 1 and Figure 3E).

These results support that maternal resource environment affects exploratory behavior, but also

imply heritable differences between populations. Similar heritable population differences were also

found for feeding accuracy and burst swimming but without any evidence for maternal effects. At

least for burst swimming, the pattern of population differentiation in our experiment again mirrors

findings from adult fish in natural habitats (Camarillo *et al.*, 2020).

Trait variation and adaptive function

Sulfide springs and adjacent nonsulfidic habitats do not only differ in the presence and absence of

H₂S, but they also vary in numerous abiotic and biotic factors that are often unaddressed in studies

of adaptation. Sulfidic habitats have lower dissolved oxygen concentrations, higher temperature,

higher specific conductivity, and lower pH (Tobler *et al.*, 2011), which in turn affect the biotic

communities (Greenway *et al.*, 2014) and selection associated with resource exploitation,

competition, predation, and parasites (Riesch *et al.*, 2010a; Tobler *et al.*, 2014, 2015). Adaptation in

sulfide springs is therefore not solely in response to selection from H₂S but is instead in response to

a multifarious selective regime that has caused multivariate phenotypic differentiation between

populations (Tobler *et al.*, 2018). The complexity of selective regimes and evolutionary responses

make disentangling cause-and-effect relationships difficult, especially because theoretical predictions

for the effects of different, covarying sources of selection are not mutually exclusive. For example,

the evolution of large offspring size at birth could be explained by (1) selection from H₂S, which

should favor larger offspring with a lower surface-to-volume ratio to reduce the influx of toxic H₂S

(Riesch et al. 2014); (2) selection from resource constraints, which favors larger offspring with

higher energy stores (Reznick *et al.*, 1996a); or (3) relaxation of selection from predation, which also

favors larger offspring (Reznick & Endler, 1982; Johnson & Belk, 2001; Jennions *et al.*, 2006).

Similarly, resource constraints and low predation also favor more exploratory individuals that are

better able to locate and exploit resources under those conditions (Teska, Smith, & Novak, 1990;

Kaun *et al.*, 2007; Huang, Sieving, & Mary, 2012). Assessing the adaptive value of trait differences

between sulfidic and nonsulfidic population is consequently nontrivial and remains a work in

progress.

Variation in some traits investigated in our study may also be the consequence of genetic,

developmental, or functional tradeoffs with other traits. Such tradeoffs are common in organisms

inhabiting contrasting environments because divergent selection acting to optimize one trait can

inadvertently influence other traits due to constraints (Ghalambor *et al.*, 2004; Kawecki & Ebert,

2004; Garland, Downs, & Ives, 2022). For example, reductions in burst swimming performance—as

documented in our study for fish from the sulfidic population—may arise as a consequence of

selection for increased steady swimming efficiency, as different body shapes are associated with

optimization of steady *vs.* unsteady swimming (Langerhans, 2007, 2009; Tokić & Yue, 2012). Indeed,

a tradeoff between burst speed and sustained swimming performance has been documented in adult

fish from the same populations we studied here. The tradeoff is likely balanced by the need for

energy-efficient swimming in sulfidic habitats with resource constraints (but low predation) and

selection for efficient predator avoidance in nonsulfidic habitats with high abundances of natural

enemies (but abundant resources) (Camarillo *et al.*, 2020). Our study found a similar reduction in

burst swimming performance in the sulfidic population, even among individuals that had not yet

reached maturity, suggesting population differences in burst swimming arise early in ontogeny. Our

results also matched *a priori* predictions regarding tradeoffs between respiration and feeding.

Habitats rich in H_2S also experience rampant hypoxia, which has selected for the evolution of

craniofacial traits (larger heads and jaws and longer gill filaments) that increase ventilation efficiency

(Camarillo *et al.*, 2020). In addition to changes in morphology, sulfidic individuals also exhibit

decreased foraging efficiency compared to nonsulfidic individuals (Tobler *et al.*, 2009), which was

supported by our findings related to feeding accuracy (Figure 3F). The decreases in feeding

efficiency noted in this and other studies may therefore be a consequence of the craniofacial

modifications that accompany colonization of sulfidic habitats.

Plasticity accentuates genetic trait differentiation in natural populations

Phenotypes in nature are the sum of genetic and environmental effects, but, surprisingly, we found

no evidence for canalization or the evolution of plasticity by genotype × environment interactions.

Our work demonstrated that maternal effects were aligned with population differences, accentuating

trait divergence between populations, and that the trait shifts induced by maternal effects were of a

similar magnitude in both populations. For both traits in which we observed significant population

differentiation and maternal effects (size at birth and exploratory behavior), the lowest phenotypic

scores were found in nonsulfidic fry from high-food mothers, and the highest scores were found in

sulfidic fry from low-food mothers (Figures 2A and 3E). Because sulfidic mollies exhibit reduced

foraging efficiency and body condition as a consequence of hypoxia (Tobler, 2008; Tobler *et al.*,

2009), sulfidic habitats are naturally analogous to our low-food treatment, and nonsulfidic habitats

are similar to the high-food treatment. If maternal effects enhance population differences in natural

populations similarly as in our experiments, then it could explain why stronger trait divergence is

typically observed in nature than in common garden reared fish (Tobler *et al.*, 2008; Passow *et al.*,

2015). Likewise, it is important to note that our experiments only captured a small aspect of

plasticity—the portion controlled by the mother. Future work needs to address genetic effects,

maternal effects, and plasticity in response to variation in environmental factors directly experienced

by the offspring to better understand the forces driving trait variation in nature. We also caution that

the shifts found in our study are only from comparisons of a single population pair. Our

understanding of the relative roles of genetic and plastic effects on phenotypic variation would

benefit greatly from studies that expand to other sulfidic and nonsulfidic population pairs in *P.*

mexicana and other species.

Conflict of Interest Statement

The authors declare no conflict of interest.

Acknowledgements

We would like to thank the members of the community in Tapijulapa, Tabasco for access to field

sites and for their hospitality. We also would like to thank the Mexican government for issuing

permits for past field collections, as well as our collaborators at the Universidad Juárez Autónoma de

Tabasco and the Centro de Investigación E Innovación Para la Enseñanza y El Aprendizaje for

support of research expeditions to obtain the specimens that gave rise to the populations used in

this study. The assistance provided by J. Onnen and Q. La Fon was essential for processing our

diverse datasets. Funding for this research was provided by grants from the National Science

Foundation (IOS-1931657), the U.S. Army Research Office (W911NF-15-1-0175), and the Des Lee

Collaborative Vision in Zoological Studies to MT, and by a Graduate Assistance in Areas of

National Need (GAANN) fellowship from the U.S. Department of Education to JLC.

Data Availability Statement

All datasets and R scripts used to analyze the data, with associated documentation, have been

archived on GitHub (https://github.com/michitobler/common-garden).

References

Altmann J & **Alberts SC**. **2005**. Growth rates in a wild primate population: ecological influences
and maternal effects. *Behavioral ecology and sociobiology* **57**: 490–501.

Badyaev AV & **Uller T**. **2009**. Parental effects in ecology and evolution: mechanisms, processes
and implications. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **364**:
1169–1177.

Bartón K. **2009**. MuMIn: Multi-Model Inference. *Comprehensive R Archive Network (CRAN)*.

Barts N, **Greenway R**, **Passow CN**, **Arias-Rodriguez L**, **Kelley JL** & **Tobler M**. **2018**. Molecular
evolution and expression of oxygen transport genes in livebearing fishes (Poeciliidae) from hydrogen
sulfide rich springs. *Genome* **61**: 273–286.

Bates D, **Mächler M**, **Bolker B** & **Walker S**. **2015**. Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software, Articles* **67**: 1–48.

Beckerman AP, **Benton TG**, **Lapsley CT** & **Koesters N**. **2006**. How effective are maternal
effects at having effects? *Proceedings. Biological sciences* **273**: 485–493.

Belk MC, Ingley SJ & Johnson JB. 2020. Life History Divergence in Livebearing Fishes in
Response to Predation: Is There a Microevolution to Macroevolution Barrier? *Diversity* **12**: 179.

Ben-Shachar M, Lüdecke D & Makowski D. 2020. Effectsize: Estimation of effect size indices
and standardized parameters. *Journal of open source software* **5**: 2815.

Bierbach D, Schulte M, Herrmann N, Tobler M, Stadler S, Jung CT, Kunkel B, Riesch R,
Klaus S, Ziege M, Indy JR, Arias-Rodriguez L & Plath M. 2011. Predator-induced changes of
female mating preferences: innate and experiential effects. *BMC evolutionary biology* **11**: 190.

Boots M & Roberts KE. 2012. Maternal effects in disease resistance: poor maternal environment
increases offspring resistance to an insect virus. *Proceedings. Biological sciences* **279**: 4009–4014.

Camarillo H, Arias Rodriguez L & Tobler M. 2020. Functional consequences of phenotypic
variation between locally adapted populations: swimming performance and ventilation in
extremophile fish. *Journal of evolutionary biology* **33**: 512–523.

Chambers J. 1987. The cyprinodontiform gonopodium, with an atlas of the gonopodia of the fishes
of the genus Limia. *Journal of fish biology* **30**: 389–418.

Conrad JL, Weinersmith KL, Brodin T, Saltz JB & Sih A. 2011. Behavioural syndromes in fishes:
a review with implications for ecology and fisheries management. *Journal of fish biology* **78**: 395–435.

Cooper CE & Brown GC. 2008. The inhibition of mitochondrial cytochrome oxidase by the gases
carbon monoxide, nitric oxide, hydrogen cyanide and hydrogen sulfide: chemical mechanism and
physiological significance. *Journal of Bioenergy and Biomembranes* **40**: 533–539.

Dingemanse NJ & Araya-Ajoy YG. 2015. Interacting personalities: behavioural ecology meets
quantitative genetics. *Trends in ecology & evolution* **30**: 88–97.

Domenici P & Blake RW. 1997. The kinematics and performance of fish fast-start swimming. *The*
Journal of experimental biology **200**: 1165–1178.

Doran C, Bierbach D, Lukas J, Klamser P, Landgraf T, Klenz H, Habedank M, Arias-
Rodriguez L, Krause S, Romanczuk P & Krause J. 2022. Fish waves as emergent collective
antipredator behavior. *Current biology: CB* **32**: 708-714.e4.

Eaton RC, Bombardieri RA & Meyer DL. 1977. The Mauthner-initiated startle response in teleost
fish. *The journal of experimental biology* **66**: 65–81.

Endler JA. 1980. NATURAL SELECTION ON COLOR PATTERNS IN *POECILIA*
RETICULATA. *Evolution; international journal of organic evolution* **34**: 76–91.

Endler JA. 1995. Multiple-trait coevolution and environmental gradients in guppies. *Trends in ecology*
& evolution **10**: 22–29.

Felmy A, Reznick DN, Travis J, Potter T & Coulson T. 2022. Life histories as mosaics: plastic
and genetic components differ among traits that underpin life-history strategies. *Evolution;*
international journal of organic evolution.

Fox J, Friendly M & Monette G. 2021. heplots: Visualizing Hypothesis Tests in Multivariate Linear Models.

Fox J & Weisberg S. 2018a. Visualizing fit and lack of fit in complex regression models with predictor effect plots and partial residuals. *Journal of statistical software* **87**.

Fox J & Weisberg S. 2018b. *An R Companion to Applied Regression*. SAGE Publications.

Friard O & Gamba M. 2016. BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations. *Methods in ecology and evolution / British Ecological Society* **7**: 1325–1330.

Garland T Jr, Downs CJ & Ives AR. 2022. Trade-offs (and constraints) in organismal biology. *Physiological and biochemical zoology: PBZ* **95**: 82–112.

Ghalambor CK, Hoke KL, Ruell EW, Fischer EK, Reznick DN & Hughes KA. 2015. Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature* **525**: 372–375.

Ghalambor CK, Reznick DN & Walker JA. 2004. Constraints on adaptive evolution: the functional trade-off between reproduction and fast-start swimming performance in the Trinidadian guppy (*Poecilia reticulata*). *The American naturalist* **164**: 38–50.

Grafen A. 1988. On the uses of data on lifetime reproductive success. *Reproductive success*: 454–485.

Greenway R, Arias-Rodriguez L, Diaz P & Tobler M. 2014. Patterns of macroinvertebrate and fish diversity in freshwater sulphide springs. *Diversity* **6**: 597–632.

Greenway R, Barts N, Henpita C, Brown AP, Arias Rodriguez L, Rodríguez Peña CM, Arndt S, Lau GY, Murphy MP, Wu L, Lin D, Kelley JL, Shaw JH & Tobler M. 2020. Convergent evolution of conserved mitochondrial pathways underlies repeated adaptation to extreme environments. *Proceedings of the National Academy of Sciences of the United States of America* **117**: 16424–16430.

Hedrick TL. 2008. Software techniques for two- and three-dimensional kinematic measurements of biological and biomimetic systems. *Bioinspiration & biomimetics* **3**: 034001.

Hegyi G, Török J, Tóth L, Garamszegi LZ & Rosivall B. 2006. Rapid temporal change in the expression and age-related information content of a sexually selected trait. *Journal of evolutionary biology* **19**: 228–238.

Huang P, Sieving KE & Mary CMS. 2012. Heterospecific information about predation risk influences exploratory behavior. *Behavioral ecology: official journal of the International Society for Behavioral Ecology* **23**: 463–472.

Ingley SJ, Camarillo H, Willis H & Johnson JB. 2016. Repeated evolution of local adaptation in swimming performance: population-level trade-offs between burst and endurance swimming in Brachyrhaphis freshwater fish. *Biological journal of the Linnean Society. Linnean Society of London* **119**: 1011–1026.

Ingley SJ & **Johnson JB**. **2016**. Divergent natural selection promotes immigrant inviability at early
and late stages of evolutionary divergence. *Evolution; international journal of organic evolution* **70**: 600–616.

Jennions MD, **Wong BBM**, **Cowling A** & **Donnelly C**. **2006**. Life-history phenotypes in a live-
bearing fish *Brachyrhaphis episcopi* living under different predator regimes: seasonal effects?
Environmental biology of fishes **76**: 211–219.

Johnson JB & **Belk MC**. **2001**. Predation environment predicts divergent life-history phenotypes
among populations of the livebearing fish *Brachyrhaphis rhabdophora*. *Oecologia* **126**: 142–149.

Johnson JB & **Omland KS**. **2004**. Model selection in ecology and evolution. *Trends in ecology &*
evolution **19**: 101–108.

Kaun KR, **Riedl CAL**, **Chakaborty-Chatterjee M**, **Belay AT**, **Douglas SJ**, **Gibbs AG** &
Sokolowski MB. **2007**. Natural variation in food acquisition mediated via a Drosophila cGMP-
dependent protein kinase. *The journal of experimental biology* **210**: 3547–3558.

Kawecki TJ & **Ebert D**. **2004**. Conceptual issues in local adaptation. *Ecology letters* **7**: 1225–1241.

Lande R & **Arnold SJ**. **1983**. THE MEASUREMENT OF SELECTION ON CORRELATED
CHARACTERS. *Evolution; international journal of organic evolution* **37**: 1210–1226.

Langerhans RB, **Layman CA**, **Shokrollahi AM** & **DeWitt TJ**. **2004**. Predator-driven phenotypic
diversification in *Gambusia affinis*. *Evolution; international journal of organic evolution* **58**: 2305–2318.

Langerhans RB. **2007**. Evolutionary consequences of predation: avoidance, escape, reproduction,
and diversification. In: Elewa AMT, ed. *Predation in Organisms*. Berlin: Springer.

Langerhans RB. **2009**. Trade-off between steady and unsteady swimming underlies predator-driven
divergence in *Gambusia affinis*. *Journal of evolutionary biology* **22**: 1057–1075.

Langerhans RB, **Gifford ME** & **Joseph EO**. **2007**. Ecological speciation in Gambusia fishes.
Evolution; international journal of organic evolution **61**: 2056–2074.

Leonard AM & **Lancaster LT**. **2020**. Maladaptive plasticity facilitates evolution of thermal
tolerance during an experimental range shift. *BMC evolutionary biology* **20**: 47.

Lindholm AK, **Hunt J** & **Brooks R**. **2006**. Where do all the maternal effects go? Variation in
offspring body size through ontogeny in the live-bearing fish *Poecilia parae*. *Biology letters* **2**: 586–589.

Lukas J, **Auer F**, **Goldhammer T**, **Krause J**, **Romanczuk P**, **Klamser P**, **Arias-Rodriguez L** &
Bierbach D. **2021**. Diurnal changes in hypoxia shape predator-prey interaction in a bird-fish system.
Frontiers in Ecology and Evolution **9**: 151.

MacLeod KJ, **While GM** & **Uller T**. **2021**. Viviparous mothers impose stronger glucocorticoid-
mediated maternal stress effects on their offspring than oviparous mothers. *Ecology and evolution* **11**:
17238–17259.

Marshall DJ & **Uller T**. **2007**. When is a maternal effect adaptive? *Oikos (Copenhagen, Denmark)* **116**:
1957–1963.

Monaghan P. **2008**. Early growth conditions, phenotypic development and environmental change.
Philosophical transactions of the Royal Society of London. Series B, Biological sciences **363**: 1635–1645.

Mousseau T & **Fox C** (Eds.). **1998**. *Maternal effects as adaptations*. New York: Oxford University
Press.

Palacios M, Arias-Rodriguez L, Plath M, Eifert C, Lerp H, Lamboj A, Voelker G & **Tobler**
M. **2013**. The rediscovery of a long described species reveals additional complexity in speciation
patterns of poeciliid fishes in sulfide springs. *PloS one* **8**: e71069.

Passow C, Greenway R, Arias-Rodriguez L, Jeyasingh PD & **Tobler M**. **2015**. Reduction of
energetic demands through modification of body size and metabolic rates in locally adapted
extremophile fishes. *Physiological and biochemical zoology: PBZ* **88**: 371–383.

Passow CN, Henpita C, Shaw JH, Quackenbush CR, Warren WC, Schartl M, Arias-
Rodriguez L, Kelley JL & **Tobler M**. **2017a**. The roles of plasticity and evolutionary change in
shaping gene expression variation in natural populations of extremophile fish. *Molecular ecology* **26**:
6384–6399.

Passow CN, Arias-Rodriguez L & **Tobler M**. **2017b**. Convergent evolution of reduced energy
demands in extremophile fish. *PloS one* **12**: e0186935.

Pérez-Escudero A, Vicente-Page J, Hinz RC, Arganda S & **de Polavieja GG**. **2014**. idTracker:
tracking individuals in a group by automatic identification of unmarked animals. *Nature methods* **11**:
743–748.

Pigliucci M. **2001**. *Phenotypic plasticity: beyond nature and nurture*. Baltimore, MD: The Johns Hopkins
University Press.

Plath M, Heubel KU, De León FJG & **Schlupp I**. **2005**. Cave molly females (*Poecilia mexicana*,
Poeciliidae, Teleostei) like well-fed males. *Behavioral ecology and sociobiology* **58**: 144–151.

Plath M, Tobler M, Riesch R, Garcia de Leon FJ, Giere O & **Schlupp I**. **2007**. Survival in an
extreme habitat: the role of behaviour and energy limitation. *Die Naturwissenschaften* **94**: 991–996.

Plath M, Pfenninger M, Lerp H, Riesch R, Eschenbrenner C, Slattery PA, Bierbach D,
Herrmann N, Schulte M, Arias-Rodriguez L, Rimber Indy J, Passow C & **Tobler M**. **2013**.
Genetic differentiation and selection against migrants in evolutionarily replicated extreme
environments. *Evolution; international journal of organic evolution* **67**: 2647–2661.

R Core Team. **2023**. *R: A language and environment for statistical computing*. Vienna, Austria: R
Foundation for Statistical Computing.

Reiffenstein RJ, Hulbert WC & **Roth SH**. **1992**. Toxicology of hydrogen sulfide. *Annual Reviews of*
Pharmacology and Toxicology **1992**: 109–134.

Reznick DN & **Bryga H**. **1987**. Life-history evolution in guppies (*Poecilia reticulata*): 1. Phenotypic
and genetic changes in an introduction experiment. *Evolution; international journal of organic evolution* **41**:
1370–1385.

Reznick DN, **Bryga HA** & **Endler JA**. **1990**. Experimentally induced life-history evolution in a
natural population. *Nature* **346**: 357–359.

Reznick D, **Callahan H** & **Llauredo R**. **1996a**. Maternal effects on offspring quality in poeciliid
fishes. *American zoologist* **36**: 147–156.

Reznick D & **Endler JA**. **1982**. The impact of predation on life history evolution in trinidadian
guppies (*Poecilia reticulata*). *Evolution; international journal of organic evolution* **36**: 160–177.

Reznick DN, **Rodd FH** & **Cardenas M**. **1996b**. Life-History Evolution in Guppies (*Poecilia*
reticulata: Poeciliidae). IV. Parallelism in Life-History Phenotypes. *The American naturalist* **147**: 319–
338.

Riesch R, **Duwe V**, **Herrmann N**, **Padur L**, **Ramm A**, **Scharnweber K**, **Schulte M**, **Schulz-**
Mirbach T, **Ziege M** & **Plath M**. **2009**. Variation along the shy–bold continuum in extremophile
fishes (*Poecilia mexicana*, *Poecilia sulphuraria*). *Behavioral ecology and sociobiology* **63**: 1515–1526.

Riesch R, **Oranth A**, **Dzienko J**, **Karau N**, **Schiessl A**, **Stadler S**, **Wigh A**, **Zimmer C**, **Arias-**
Rodriguez L, **Schlupp I** & **Plath M**. **2010a**. Extreme habitats are not refuges: poeciliids suffer
from increased aerial predation risk in sulphidic southern Mexican habitats. *Biological journal of the*
Linnean Society. Linnean Society of London **101**: 417–426.

Riesch R, **Plath M**, **Garcia de Leon FJ** & **Schlupp I**. **2010b**. Convergent life-history shifts: toxic
environments result in big babies in two clades of poeciliids. *Die Naturwissenschaften* **97**: 133–141.

Riesch R, **Schlupp I**, **Langerhans RB** & **Plath M**. **2011a**. Shared and unique patterns of embryo
development in extremophile poeciliids. *PloS one* **6**: e27377.

Riesch R, **Plath M**, **Schlupp I**, **Tobler M** & **Langerhans RB**. **2014**. Colonization of toxic springs
drives predictable life-history shift in livebearing fishes (Poeciliidae). *Ecology letters* **17**: 65–71.

Riesch R, **Reznick DN**, **Plath M** & **Schlupp I**. **2016a**. Sex-specific local life-history adaptation in
surface- and cave-dwelling Atlantic mollies (*Poecilia mexicana*). *Scientific reports* **6**: 22968.

Riesch R, **Tobler M**, **Lerp H**, **Jourdan J**, **Doumas T**, **Nosil P**, **Langerhans RB** & **Plath M**.
2016b. Extremophile Poeciliidae: multivariate insights into the complexity of speciation along
replicated ecological gradients. *BMC evolutionary biology* **16**: 136.

Riesch R, **Plath M** & **Schlupp I**. **2010c**. Toxic hydrogen sulfide and dark caves: life-history
adaptations in a livebearing fish (*Poecilia mexicana*, Poeciliidae). *Ecology* **91**: 1494–1505.

Riesch R, **Plath M** & **Schlupp I**. **2011b**. Toxic hydrogen sulphide and dark caves: pronounced
male life-history divergence among locally adapted *Poecilia mexicana* (Poeciliidae). *Journal of evolutionary*
biology **24**: 596–606.

Rosen DE & **Gordon MS**. **1953**. Functional anatomy and evolution of male genitalia in Poeciliid fishes. *Zoologica* **38**: 1–47.

Rossiter M. **1996**. Incidence and consequences of inherited environmental effects. *Annual review of ecology and systematics* **27**: 451–476.

Scheiner SM. **1993**. Genetics and Evolution of Phenotypic Plasticity. *Annual review of ecology and systematics* **24**: 35–68.

Schneider CA, **Rasband WS** & **Eliceiri KW**. **2012**. NIH Image to ImageJ: 25 years of image analysis. *Nature methods* **9**: 671–675.

Schrader M & **Travis J**. **2012**. Variation in offspring size with birth order in placental fish: a role for asymmetric sibling competition? *Evolution; international journal of organic evolution* **66**: 272–279.

Schuler MS & **Orrock JL**. **2012**. The maladaptive significance of maternal effects for plants in anthropogenically modified environments. *Evolutionary ecology* **26**: 475–481.

Schulz-Mirbach T, **Eifert C**, **Riesch R**, **Farnworth MS**, **Zimmer C**, **Bierbach D**, **Klaus S**, **Tobler M**, **Streit B**, **Indy JR**, **Arias-Rodriguez L** & **Plath M**. **2016**. Toxic hydrogen sulphide shapes brain anatomy: a comparative study of sulphide-adapted ecotypes in the *Poecilia mexicana* complex. *Journal of zoology* **300**: 163–176.

Stearns SC & **Sage RD**. **1980**. Maladaptation in a Marginal Population of the Mosquito Fish, *Gambusia affinis*. *Evolution; international journal of organic evolution* **34**: 65–75.

Stenseth NC, **Andersson L** & **Hoekstra HE**. **2022**. Gregor Johann Mendel and the development of modern evolutionary biology. *Proceedings of the National Academy of Sciences of the United States of America* **119**: e2201327119.

Teska WR, **Smith MH** & **Novak JM**. **1990**. Food quality, heterozygosity, and fitness correlates in *Peromyscus polionotus*. *Evolution; international journal of organic evolution* **44**: 1318–1325.

Tobler M, **Schlupp I**, **Heubel KU**, **Riesch R**, **de León FJG**, **Giere O** & **Plath M**. **2006**. Life on the edge: hydrogen sulfide and the fish communities of a Mexican cave and surrounding waters. *Extremophiles: life under extreme conditions* **10**: 577–585.

Tobler M, **Dewitt TJ**, **Schlupp I**, **García de León FJ**, **Herrmann R**, **Feulner PGD**, **Tiedemann R** & **Plath M**. **2008**. Toxic hydrogen sulfide and dark caves: phenotypic and genetic divergence across two abiotic environmental gradients in *Poecilia mexicana*. *Evolution; international journal of organic evolution* **62**: 2643–2659.

Tobler M. **2008**. Divergence in trophic ecology characterizes colonization of extreme habitats. *Biological journal of the Linnean Society. Linnean Society of London* **95**: 517–528.

Tobler M, **Riesch R**, **Tobler CM** & **Plath M**. **2009**. Compensatory behaviour in response to sulfide-induced hypoxia affects time budgets, feeding efficiency, and predation risk. *Evolutionary ecology research* **11**: 935–948.

Tobler M, Palacios M, Chapman LJ, Mitrofanov I, Bierbach D, Plath M, Arias-Rodriguez L,
de León FJG & Mateos M. 2011. Evolution in extreme environments: replicated phenotypic
differentiation in livebearing fish inhabiting sulfidic springs. *Evolution; international journal of organic*
evolution **65**: 2213–2228.

Tobler M, Plath M, Riesch R, Schlupp I, Grasse A, Munimanda GK, Setzer C, Penn DJ &
Moodley Y. 2014. Selection from parasites favours immunogenetic diversity but not divergence
among locally adapted host populations. *Journal of evolutionary biology* **27**: 960–974.

Tobler M, Scharnweber K, Greenway R, Passow CN, Arias-Rodriguez L & García-De-León
FJ. 2015. Convergent changes in the trophic ecology of extremophile fish along replicated
environmental gradients. *Freshwater biology* **60**: 768–780.

Tobler M, Alba DM, Arias-Rodríguez L & Jeyasingh PD. 2016a. Using replicated evolution in
extremophile fish to understand diversification in elemental composition and nutrient excretion.
Freshwater biology **61**: 158–171.

Tobler M, Passow CN, Greenway R, Kelley JL & Shaw JH. 2016b. The evolutionary ecology of
animals inhabiting hydrogen sulfide-rich environments. *Annual review of ecology, evolution, and systematics*
47: 239–262.

Tobler M, Kelley JL, Plath M & Riesch R. 2018. Extreme environments and the origins of
biodiversity: adaptation and speciation in sulphide spring fishes. *Molecular ecology* **27**: 843–859.

Tobler M & Hastings L. 2011. Convergent patterns of body shape differentiation in four different
clades of poeciliid fishes inhabiting sulfide springs. *Evolutionary biology* **38**: 412–421.

Tokić G & Yue DKP. 2012. Optimal shape and motion of undulatory swimming organisms.
Proceedings. Biological sciences **279**: 3065–3074.

Tomar S. 2006. Converting Video Formats with FFmpeg. *Linux Journal* **146**: 10.

Uller T. 2008. Developmental plasticity and the evolution of parental effects. *Trends in ecology &*
evolution **23**: 432–438.

Van Allen B, Jones N, Gilbert B, Carscadden K & Germain R. 2021. Maternal effects and the
outcome of interspecific competition. *Ecology and evolution* **11**: 7544–7556.

Walker JA. 1997. Ecological morphology of lacustrine threespine stickleback *Gasterosteus aculeatus* L.
(Gasterosteidae) body shape. *Biological journal of the Linnean Society. Linnean Society of London* **61**: 3–50.

Weeks SC & Gaggiotti OE. 1993. Patterns of offspring size at birth in clonal and sexual strains of
Poeciliopsis (poeciliidae). *Copeia* **1993**: 1003.

West-Eberhard MJ. 1989. Phenotypic plasticity and the origins of diversity. *Annual review of ecology*
and systematics **20**: 249–278.

Wilson AJ, **Coltman DW**, **Pemberton JM**, **Overall ADJ**, **Byrne KA** & **Kruuk LEB**. **2005**.
Maternal genetic effects set the potential for evolution in a free-living vertebrate population. *Journal*
of evolutionary biology **18**: 405–414.

Winemiller KO & **Rose KA**. **1992**. Patterns of life-history diversification in North American fishes:
Implications for population regulation. *Canadian journal of fisheries and aquatic sciences. Journal canadien des*
sciences halieutiques et aquatiques **49**: 2196–2218.

Wolf JB & **Wade MJ**. **2009**. What are maternal effects (and what are they not)? *Philosophical*
transactions of the Royal Society of London. Series B, Biological sciences **364**: 1107–1115.

Zhang H, **Vedder O**, **Becker PH** & **Bouwhuis S**. **2015**. Age-dependent trait variation: the relative
contribution of within-individual change, selective appearance and disappearance in a long-lived
seabird. *The journal of animal ecology* **84**: 797–807.

Table 1. Model terms in the best-supported model for each dependent variable. Empty cells represent terms that were not present in the best-supported model for that dependent variable. Type-III ANOVA or Wald's Chi-square tests were used to determine whether model terms were significantly different from zero. F-statistics or χ^2 statistics are presented for each included model term, with the degrees of freedom in a subscript. P-values are presented underneath each test statistic.

Dependent Variable	Brood Size	Treatment Length	Mother SL	Age	Population	Food Treatment
Size at Birth	$F_{1,35} = 20.876$ $P < 0.001$		$F_{1,35} = 16.593$ $P < 0.001$		$F_{1,35} = 20.032$ $P < 0.001$	$F_{1,35} = 4.354$ $P = 0.044$
Brood Size		$F_{1,37} = 8.926$ $P = 0.005$	$F_{1,37} = 23.342$ $P < 0.001$			
Age at Maturity					$F_{1,26} = 4.420$ $P = 0.045$	
Growth Rate at Birth					$F_{1,37} = 5.025$ $P = 0.031$	
Overall Growth Rate				$\chi^2_1 = 147.91$ $P < 0.001$		
Burst Swimming at Birth			$F_{1,39} = 5.025$ $P = 0.121$			
Overall Burst Swimming				$\chi^2_1 = 30.178$ $P < 0.001$	$\chi^2_1 = 4.308$ $P = 0.038$	
Exploratory Behavior at Birth						$F_{1,31} = 2.727$ $P = 0.108$
Overall Exploratory Behavior					$\chi^2_1 = 7.513$ $P = 0.006$	$\chi^2_1 = 4.245$ $P = 0.039$
Feeding Accuracy at Birth		$F_{1,38} = 0.775$ $P = 0.384$				
Overall Feeding Accuracy					$\chi^2_1 = 4.645$ $P = 0.031$	
Multivariate Trait Variation PC1		$F_{1,24} = 15.817$ $P < 0.001$			$F_{1,24} = 14.984$ $P < 0.001$	$F_{1,24} = 4.746$ $P = 0.039$
Multivariate Trait Variation PC2		$F_{1,24} = 7.722$ $P = 0.010$	$F_{1,24} = 5.292$ $P = 0.030$		$F_{1,24} = 3.321$ $P = 0.081$	

Figure 1. Overview of our experimental design. We subjected pregnant sulfidic and nonsulfidic

mothers to either a high- or a low-food treatment and measured seven traits in their offspring

throughout their development. These traits include four life history traits (birth size, brood size,

growth rate, and age at maturity) and three behavioral traits (burst swimming, exploratory behavior,

and feeding accuracy).

Figure 2. Plots of average standard length (A), brood size (B), and their relationship (C) for each

family. Data is summarized in boxplots with raw data points overlaid. In both populations, the high-

food treatment is shown in blue, and the low-food treatment is shown in orange. When the best-

supported model for a phenotype contained the terms 'Population' and 'Food Treatment', the

estimated marginal means for those effects were visualized as large points.

Figure 3. Data for each individual phenotype, including (A) growth rate at birth, (B) overall growth

rate, (C) age at maturity, (D) burst swimming, (E) exploratory behavior, and (F) feeding accuracy.

Data is summarized in boxplots with raw data points overlaid. In both populations, the high-food

treatment is shown in blue, and the low-food treatment is shown in orange. When the best-

supported model for a phenotype contained the terms 'Population' and/or 'Food Treatment', the

estimated marginal means for those effects were visualized as large points.

Figure 4. Plot of principal component scores representing linear combinations of all phenotypes in

multivariate space. Scores were plotted along the first two principal components. Nonsulfidic

families are shown in shades of blue, while sulfidic families are shown in shades of yellow. The high-

food treatment within each population is shown in the darker shade (*i.e.*, dark blue or dark yellow).

Radiating from the origin are arrows that represent the correlation between the principal component

scores and each input variable (shown as text at the end of each arrow). Loadings were calculated by

multiplying the eigenvector for each input variable by the square-root of the eigenvalue for that

principal component axis. At the top of the figure are boxplots that show the distribution of

principal component scores along the first principal component axis, which exhibited significant

effects of 'Population' and 'Food Treatment'.