A preclinical model of THC edibles that produces high-dose cannabimimetic responses.

Anthony English[1,2,3], Fleur Uittenbogaard[1,2,3], Alexa Torrens[4], Dennis Sarroza[1], Anna Slaven[1,2], Danielle Piomelli[4], Michael R. Bruchas[1,2,3,4], Nephi Stella*[,1,2,3,6], and Benjamin B. Land*[,1,2,3].

[1]Departments of Pharmacology, [2]UW Center of Excellence in Neurobiology of Addiction, Pain, and Emotion (NAPE), [3]Center for Cannabis Research, University of Washington, Seattle WA 98195.
[4]Department of Anatomy & Neurobiology, University of California Irvine, Irvine CA.
[5]Department of Anesthesiology and [6]Psychiatry & Behavioral Sciences, University of Washington, Seattle WA 98195.

*Co-Corresponding Authors: Nephi Stella nstella@uw.edu and Benjamin Land bbl2@uw.edu

Abstract

No preclinical experimental approach enables the study of voluntary oral consumption of high-concentration Δ^9-tetrahydrocannabinol (**THC**) and its intoxicating effects, mainly owing to the aversive response of rodents to THC that limits intake. Here we developed a palatable THC formulation and an optimized access paradigm in mice to drive voluntary consumption. THC was formulated in chocolate gelatin (**THC-E-gel**). Adult male and female mice were allowed *ad libitum* access for 2 h. Cannabimimetic responses (hypolocomotion, analgesia, and hypothermia) were measured following access. Levels of THC and its metabolites were measured in blood and brain tissue. Acoustic startle responses were measured to investigate THC-induced psychotomimetic behavior.

Adult mice consumed high-dose THC-E-gel (≈30 mg/kg over 2 h) and exhibited robust CB_1 receptor-dependent behavioral responses. High-dose THC-E-gel consumption led to parallel accumulation of THC and its psychoactive metabolite, 11-OH-THC, in brain, a profile that contrasts with the known rapid decline in brain 11-OH-THC levels following intraperitoneal THC injections. High-dose THC-E-gel consumption increased the acoustic startle response preferentially in males, and this psychotomimetic response was remarkably different from the response triggered by intraperitoneal contingent administration of THC. Comparing cannabimimetic responses elicited by intraperitoneal versus oral administration enabled a "predicted dose" model of THC triggered behavioral changes. Significantly, voluntary consumption of high-dose THC-E-gel triggered equivalent cannabimimetic responses in male and female mice but increased acoustic startle response preferentially in males. Thus, THC-E-gel offers a robust preclinical experimental approach to study cannabimimetic responses triggered by voluntary consumption in mice, including sex-dependent psychotomimetic responses.

Introduction

The use of *Cannabis* products containing high concentrations of THC is rapidly increasing despite our limited understanding of its potential impact on physical and mental health [1-3]. These products are typically inhaled as combusted plant matter, vaporized extracts, or consumed in edible formulations. THC acts as a partial agonist at CB_1 receptors (**CB_1R**) to trigger a myriad of responses such as physiological responses (e.g., increase heart rate), altered mood and time perception, inhibition of motor control, and impaired learning and memory [4-10]. Subsequently, a relationship between *Cannabis* and psychotic/affective symptoms and an observable increase in *Cannabis*-associated vehicle crashes has become apparent without an understanding of the neural effects of high dose THC [6,11-14]. Many of these effects translate to preclinical models where in rodents, THC reduces spontaneous locomotion and locomotor control, induces hypersensitivity to tactile and auditory stimuli, ataxia, and sedation; all of which have been shown to be mediated through action at CB_1R [15-18]. Importantly, some cannabimimetic responses are sex-dependent, as exemplified by the finding that THC (5 mg/kg, *i.p.*) triggers a more pronounced reduction in spontaneous locomotion and anxiogenic response in females than in males [19,20]. In addition to these cannabimimetic responses, preclinical investigations have pursued psychosis-related behaviors through the acoustic startle response, finding that involuntary administration of THC impairs psychomotor/sensorimotor gaiting [21-24], emphasizing the translational value in understanding THC's bioactivity in humans.

Understanding the effects of increased THC use in humans through preclinical models of voluntary THC administration has proven difficult to establish due to the aversive behaviors to high doses in rodent models [25,26]. In recent years, progress has been made in promoting

voluntary oral consumption of THC in rodents, but results have been limited to mild, acute CB_1R-dependent cannabimimetic responses [27-29]. This lack of experimental tools to translate high-dose THC intake in humans to preclinical models emphasizes the urgent need to develop and fully characterize a novel experimental approach. To bridge this translational gap, we initially developed an approach where mice are given *ad libitum* access to consume a sugar-water gelatin (**CTR-gel**) containing fixed amounts of THC [28,30]. Matching previous rodent studies, we found that mice consumed more vehicle gelatin than THC gelatin, indicating that they detected and avoided THC [31]. To overcome this limitation, in the current study we developed and characterized a palatable oral gelatin formulation that increases voluntary consumption by formulating THC in a chocolate-flavored nutritional shake, Ensure™ (**E-gel**). Previous work has shown that mice have a preference for chocolate flavor, making it an ideal THC formulant to increase palatability [32,33]. We leveraged this approach to determine whether oral consumption of high THC doses induce commonly studied cannabimimetic responses in mice (hypolocomotion, analgesia, and hypothermia) and then we examined the effects of THC E-gel consumption on acoustic startle response, a preclinical measure of reflexive response rate and psychomotor arousal [22,34]. The model developed here leverages acute voluntary consumption of a sweetened gelatin to investigate psychomotor and reflexive behaviors, pharmacokinetics, and triad responses following consumption of high-dose THC in mice.

Materials and Methods

Animal Studies

Animal studies followed the guidelines established by AAALAC and were approved by IACUC of the University of Washington. Male and female C57BL/6 mice ranging from 8-14 weeks of age were used. Animals were housed with sibling littermates and were provided with standard chow and water, *ad libitum*. Animal procedures were approved by the Institutional Animal Care and Use Committee of the University of Washington and conform to the guidelines on the care and use of animals by the National Institutes of Health.

Pharmacological Agents

Animals received THC (0.1, 0.3, 1, 3, 5, 10, and 30 mg/kg) and SR141716 (**SR1**, 1 mg/kg) *i.p.* or were exposed to THC suspended in gelatin. THC and SR141716 (SR1) were provided by the National Institute of Drugs abuse Drug Supply Program (Bethesda, MD). THC in ethanol (50 mg/ml) was either added to gelatin mixtures (CTR or Ensure®) or prepared for *i.p.* injection. For *i.p.* injection, both THC (0.1, 0.3, 1, 3, 5, 10, and 30 mg/kg) and SR1 (1mg/kg) were dissolved in 95% ethanol and then vortexed thoroughly with equal volume Cremophor and finally dissolved in sterile saline to reach a final 1:1:18 solution consisting of ethanol:cremophor:saline.

Gelatin formulation

Control gelatin (CTR-gel): Deionized water (100 mL) was warmed to 40°C and stirred at a constant rate. 2.5 g of Polycal™ sugar and 3.85 g of Knox™ Gelatin were added and the mixture was maintained at a temperature below 43°C. The mixture was removed from heat, and THC (50 mg/ml in ethanol) was added to reach a concentration of 0.3, 1, 2, or 4 mg/15 ml. An equal volume of ethanol was added to vehicle gelatin (<1% total volume). Gelatin was poured into plastic cups ranging 2-10 ml and set into a 4°C fridge to solidify overnight.

Ensure gelatin (E-gel): Chocolate-flavored Ensure™ (100 ml) was warmed to 40°C and stirred at a constant rate. 3.85 g of Knox™ Gelatin were added and the mixture was maintained at a temperature below 43°C. The mixture was removed from heat and THC (50 mg/15 ml ethanol) was added to reach a concentration of 1, 2, 5, or 10 mg/15 ml. At the 10 mg/15 ml concentration, ethanol was evaporated off to 50% volume before being added to the mixture to reduce total alcohol concentration below 1%. An equal volume of ethanol was added to vehicle gelatin (<1% total volume). Gelatin was poured into plastic cups ranging 2-10 ml and set into a 4°C fridge to solidify overnight. Mice were always exposed to more gelatin than they could consume, smaller volumes were used to conserve THC.

Acute gelatin access: Animals were first habituated to gelatin by receiving an excess of gelatin in their home cage the day before the first timed access. On the first day of access, mice were placed into a home cage-like environment equipped with a vehicle gelatin cup that was stabilized to the cage. Behavior was recorded during the consumption window via an overhead camera. On the second day of access, animals were placed into the same gelatin access cage with either a vehicle or THC gelatin cup. Animals experienced either a triad of behaviors (open field, tail flick, and body temperature) measured immediately preceding and following the consumption window or an acoustic startle trial immediately following consumption. On the third and final day of access animals were placed into the same cage with a vehicle gelatin cup. For all gelatin access days, gelatin cups and animals were weighed before and after the consumption window. Access to gelatin during the consumption window was limited to either 1 or 2 h after which the animals were removed and returned to their home cage.

Triad of Cannabimimetic behaviors

Hypolocomotion, hypothermia, and analgesia were measured 1 h post-*i.p.* injection or immediately following gelatin exposure. Pre-tests were collected immediately prior to injection or gelatin exposure.

Open Field: A 50 cm x 50 cm chamber (25 LUX) was equipped with an overhead camera to record movement. Animals were placed in the chamber for 15 minutes and then returned to their home cage. Total distance traveled (cm) was measured using Noldus Ethovision behavioral tracking software. Locomotion behavior was measured immediately before and after gelatin access to calculate a gelatin dependent difference score (post-pre). Pre-test measurements for CTR-gel were not collected and pre-test values were instead normalized to vehicle post-tests to produce a difference score as Post-VEH.

Tail Flick Analgesia: A hot water bath was set to 52.5°C. Mice were securely held upright in the air with their tail hanging downward. A timer was started as 75% of their tail was submerged into the water. Time was measured once a painful response was presented, marked as a latency to flick their tail out of the hot water. Tail flick responses were measured immediately before and after gelatin access to calculate a gelatin-dependent difference score (post-pre).

Measuring body temperature: Animals were placed on a stable surface with their tails lifted. A rectal thermometer probe (RET-3 Kent Scientific) was inserted into the anus for 10-20 s until the temperature recording stabilized. This test was always performed prior to the Tail Flick test to reduce any potential temperature contamination effects. Body temperatures were measured immediately before and after gelatin access to calculate a gelatin dependent difference score (post-pre).

Histology

Blood and brain tissue collection and quantification

Animals underwent the same gelatin access paradigm for day 1 and 2 described in acute gelatin access. After 2 h of gelatin access, blood was collected by cardiac puncture with a 23-gauge needle and placed on ice. Immediately following, brain tissue was collected and flash frozen in liquid nitrogen. Blood samples were spun in a 4°C centrifuge at 1450 x g for 15 min. Plasma was transferred to another tube and stored alongside brain samples in -80°C until being shipped on dry ice to the Piomelli Lab at UCI for sample analysis. Samples were collected immediately following 1 and 2 h gelatin access and 30 min and 24 h after 2 h gelatin access. THC and its first-pass metabolites 11-OH-THC and 11-COOH-THC were quantified in plasma and brain tissue using a selective isotope-dilution liquid chromatography/tandem mass spectrometry assay (26).

Acoustic startle

Acoustic Startle behaviors were measured after 1 or 2 h THC-E-gel exposure (10 mg/15 ml) and THC-*i.p.* (0.1, 1, 5, 10 mg/kg) injection. Sound-buffered startle chambers (SR-Lab, San Diego Instruments) were used to measure acoustic startle responses, equipped with a holding tube and an accelerometer. Background sound was maintained at 65 dB from a high-frequency speaker producing white noise. Startle tests were conducted 1 h post-THC-*i.p.* injection or immediately following THC-E-gel exposure. Animals were set in the holding tube for 5 min to habituate prior to a series of seven trials presenting escalating sound levels of null, 80, 90, 100, 105, 110, and 120 dB. Tones were presented for 40 ms with an inter-trial interval of 30 s. Animals were only ever exposed to the acoustic startle paradigm once, immediately after gelatin access, to avoid auditory habituation.

Data analysis

All data were analyzed using GraphPad Prism 10-11. All behavioral locomotor tracking was analyzed using Noldus Ethovision software. For all statistical analyses (unpaired t test, one- and two-way ANOVA, and post hoc analyses), alpha level was set to 0.05.

Results

E-gel promotes heightened voluntary oral consumption of THC and induces cannabimimetic behaviors by adult mice.

To incentivize voluntary oral consumption of high-concentration THC, we utilized an E-gel formulation and optimized an exposure paradigm based on previous studies [28,35]. Here, individual mice were exposed to a control (CTR-gel) during a 2 h consumption period (Habituation, Day 1); and the following day exposed to THC formulated in either CTR-gel or E-gel (X mg/15 ml) for 2 h (**Figure 1a**). Gelatin mass was measured before and after access to calculate grams consumed, and gelatin concentrations are expressed as X mg of THC (X = mg of THC/15 ml gelatin) (**Figure 1b**). As expected, higher concentrations of THC-CTR-gel reduced gelatin consumption, an effect significant at 1 mg THC (**Figure 1c**, One-way ANOVA $F_{4,108} = 9.126$, p < 0.001, Sidak's). Remarkably, 7/17 (41%) of mice exposed to 4 mg CTR-gel did not consume any gelatin while all mice consistently consumed 5 mg and 10 mg THC-E-gel (**Figure 1d**). Thus, mice consumed 1.9 ± 0.05 g of VEH-E-gel and 1.0 ± 0.07 g of E-gel containing THC (10 mg/15 ml) (**Figure 1d**, One-way ANOVA $F_{5,75} = 14.10$, p < 0.001, Sidak's). Note that mice consumed similar amounts of VEH-E-gel and VEH-CTR-gel (1.96 ± 0.15 g and 1.92 ± 0.17 g, respectively), indicating that chocolate flavor *per se* does not increase consumption. When calculating the amount of THC consumed in mg/kg, we found that mice consumed more THC when formulated in E-gel (**Figures 1e**). For example, mice consumed

10.5 ± 0.7 mg/kg/2 h when exposed to 2 mg E-gel compared to only 4.6 ± 0.5 mg/kg/2 h when exposed to the same amount of THC formulated in CTR-gel. Using this experimental approach, maximal consumption reached 29.2 ± 1.8 mg of THC per kg/2 h when exposed to E-gel containing THC (10 mg/15 ml), compared to the few mice that consumed only 8.4 ± 1.2 mg of THC per kg over 2 h when exposed to CTR-gel containing THC (4 mg) (**Figures 1e**, Two-way ANOVA $F_{7,94}$ = 73.14, p < 0.001, Sidak's). As previously shown, we found no statistically significant sex-dependent effects in consumption between male and female mice across all treatments (Two-way ANOVA $F_{1,60}$= 3.64, p<0.06) but did see an individual significance between males and females at 1 mg THC-E-gel (**Supplementary Figure S1**, Two-way ANOVA, p<0.05) [28,30]. These data show that mice consistently consume significant quantities of E-gel despite high THC concentrations. Based on these results, we next focused our study on quantifying the pharmacological effects of THC formulated in E-gel.

Considering that THC-CTR-gel triggers mild cannabimimetic responses due to limited consumption [28,30], we determined whether consumption of THC-E-gel could induce cannabimimetic responses measured immediately following the 2 h access period (**Figure 1f**). Thus, we selected three well-described behavioral effects of THC in mice: hypolocomotion, analgesia, and hypothermia (**Figure 1h-j**) [17]. THC was formulated and administered either by *i.p.* injection (grey), CTR-gel (green), or E-gel (purple) and behavioral responses to gelatin consumption were plotted from the average dose consumed, calculated in **Figure 1e**. **Figure 1h** shows that *i.p.* administration of THC reduced locomotion starting at 3 mg/kg, as previously reported [17], and that this response was significant after access to 4 mg THC-CTR-gel and 2 mg THC-E-gel (**Supplementary Figure 2b, e**, One-way ANOVA $F_{5,68}$ = 14.54, p<0.001, Sidak's for CTR-gel and $F_{4,56}$ = 3.24, p=0.02 for E-gel, Sidak's). **Figure 1i** shows the greatest

THC-induced analgesia was reached at 30 mg/kg *i.p.,* after access to 2 mg. *i.p.* injection of THC reduced core body temperature starting at 3 mg/kg, and that this response reached significance at 4 mg THC-CTR-gel and at 5 mg THC-E-gel (**Figure 1i** and **Supplementary Figure S2d, g**, One-way ANOVA $F_{4,104}$ = 9.43, p<0.001, Sidak's for CTR-gel and $F_{7,77}$ = 8.66, p<0.001 for E-gel, Sidak's). **Figure 1j** also shows that reduced core body temperature induced by THC reached a significant effect of -5.84°C at 30 mg/kg *i.p.*, -1.5°C after 4 mg THC-CTR-gel and -1.8°C after 10 mg THC-E-gel (**Supplementary Figure S2d, g**, One-way ANOVA $F_{5,61}$ = 11.16, p<0.001, Sidak's for CTR-gel and $F_{4,114}$ = 6.36, p<0.001 for E-gel, Sidak's). Analgesia and hypothermia did not plateau, matching prior studies which have also shown that 30 mg/kg THC-*i.p.* does not produce a maximal response for these cannabimimetic behaviors [36,37]. Thus, **Figures 1h-j** show that: 1) THC reduces locomotion when administered using these three experimental paradigms, and to a greater extent at high-dose *i.p.* THC and high-concentration THC-E-gel; 2) THC induced analgesia only when administered *i.p.* and using THC-E-gel, though *i.p.* administration is more potent; and 3) THC reduces core body temperature only when administered *i.p.* and using THC-E-gel, though *i.p.* administration is more potent.

Next, we analyzed the cannabimimetic responses of individual mice following THC (10 mg/15 ml) E-gel access and how the CB_1R inverse agonist, SR1, administered 1 h into the consumption window influences these responses (**Figure 1k**). SR1 was administered at 1 h to reach peak circulating concentrations during the behavioral testing (1-2 h post-injection) and to reduce any anorectic effects that would inhibit consumption of THC-E-gel [38,39]. Cannabimimetic responses increased as a function (linear regression) of increasing amount of THC consumed, demonstrating a significant relationship between the amount of THC

consumed and the three cannabimimetic readouts: hypolocomtion (p=0.003), Analgesia (p<0.001), and hypothermia (p<0.001) (**Figure 1l-n**). Confirming the involvement of CB_1R, SR1 blocked the three THC-induced cannabimimetic responses: hypolocomtion (p=0.09), Analgesia (p=0.45), and hypothermia (p=0.44) (**Figure 1l-n**). As expected, SR1-treated mice did not consume maximal THC-E-gel compared to some animals exposed to 10 mg THC-E-gel, likely due in part to the injection introduced 1 h into the consumption window and the anorectic effects of [38,39]. We additionally compared the linear regression of all E-gel tested of animals that consumed enough THC-E-gel within the range of SR1-treated animals and found that THC-E-gel alone still produced a significant correlation to all behaviors. These results indicate that consumption of THC-E-gel evokes robust CB_1R-dependent cannabimimetic behavioral responses in adult mice that are comparable to *i.p.*-THC administration when measuring hypolocomotion, and less potent when compared to *i.p.*-THC administration when measuring analgesia and reduction in core body temperature.

THC-E-Gel reduces locomotion during the exposure period.

We found that mice consumed ~2 g of vehicle E-gel (VEH-E-gel) compared to ~1 g of high-dose THC-E-gel (10 mg/15 ml), indicating a 2-fold reduction in consumption (**Figure 2a**). To investigate the time course of this effect, we weighed gelatin every 10 min during the 2 h access period in a 3-day paradigm: access to VEH-E-gel on Day 1, access to either VEH-E-gel or THC-E-gel Day 2, and access to VEH-E-gel on Day 3 (**Figure 2a**). **Figure 2b** shows that consumption of VEH-E-gel on Day 1 started within 20 min of availability and was constant during the 2 h period. On Day 2, mice consumed comparable amounts of VEH-E-gel and THC-E-gel during the initial 40 min of access (16.3 and 13.0 mg/min, respectively) (**Figure 2c** and **Supplementary Figure S4a**). However, consumption of THC-E-gel plateaued after 40 min to a

rate of 4.2 mg/min (67.7% reduction), while consumption of VEH-E-gel was sustained at 9.9 mg/min (39.3% reduction), producing a significant effect by THC to modify gelatin consumption (**Figure 2c**, Two-way ANOVA, repeated measures $F_{1,14}$ = 7.604, p = 0.015, Sidak's and **Supplementary Figure S4b**, Two-way ANOVA, $F_{1,14}$ = 6.05, p = 0.03, Sidak's). By sharp contrast, mice that had consumed THC-E-gel the day prior consumed VEH-E-gel on Day 3 at a significantly slower rate (6.2 mg/min) during the access period, suggesting an aversive memory to THC-E-gel (**Figure 2d**, Two-way ANOVA, repeated measures $F_{1,14}$ = 4.865, p = 0.045, Sidak's and **Supplementary Figure S4b**). Consequently, mice exposed to THC-E-gel on Day 2 significantly decreased their total VEH-E-gel consumption on Days 2 and 3 (**Figure 2e**, Two-way ANOVA, $F_{1,83}$ = 37.51, p < 0.001, Sidak's). These data suggest that, on Day 2, mice consumed high enough quantities of THC to induce a typically *i.p.*-high-dose (5-10 mg/kg) cannabimimetic response resulting in an avoidance to gelatin on Day 3.

Reduced spontaneous locomotion is a hallmark response to THC in mice. To address whether THC-E-gel consumption impacts spontaneous locomotion, we video-recorded the travelling distance of mice during the gelatin access period (total distance in cm over 2 h) (**Figure 2f**). **Figure 2g** shows that locomotion during the consumption period initially reached (1,200 cm/ 5 min) and then steadily decreased over the 2 h session on Day 1, as expected in mice that are habituating to the environment. On Day 2, spontaneous locomotor between mice that consumed VEH-E-gel and THC-E-gel diverged after 40 min, showing a significant decrease in total locomotion in mice that consumed THC-E-gel (**Figure 2g**, Two-way ANOVA, repeated measures $F_{1,14}$ = 11.18, p = 0.005, Sidak's). Thus, this reduction in locomotion parallels a corresponding reduction in consumption in **Figure 2e** that is significantly different on day 2 (**Figure 2h**, Two-way ANOVA, $F_{1,42}$ = 0.3413, p = 0.562, Sidak's). Importantly,

spontaneous locomotion of mice exposed to VEH-E-gel and THC-E-gel was similar on Day 3. Together, these results show that consumption of THC-E-gel induced hypolocomotion on Day 2 after 40 min of access. Additionally, the decreased consumption of VEH-E-gel on Day 3 is likely due to an aversive memory to THC and not to hypolocomotion. Thus, E-gel incentivizes voluntary THC consumption to induce robust hypolocomotion, a hallmark cannabimimetic response, within 40 min of access.

Consumption of THC-E-gel results in concomitant increases in the levels of THC and its metabolites in brain tissue.

The PK profile of THC (5 mg/kg, *i.p.*) results in peak circulating concentrations of THC (1000pmol/g), its bioactive metabolite 11-OH-THC (300pmol/g), and its inactive metabolite 11-COOH-THC (100pmol/g) in the brain after 2 h that reach approximately [40]. To determine the PK profile of high-concentration THC-E-gel consumption (10 mg) and considering the hypolocomotion behavior occurring during the consumption window, we collected plasma and brain tissue samples after 1 h of consumption, at the end of the 2 h access period, as well as 30 min (2.5 h) and 24 h (26 h) following the 2 h access period (**Figure 3a**). **Figure 3b** shows THC levels in the brain reached 500-600 pmol/g tissue between 1h and post 2.5 h time-point and was below 50 pmol/g tissue after 24 h. Remarkably, 11-OH-THC and 11-COOH-THC levels in brain increased concomitantly to THC levels, reaching 400-500 pmol/g tissue and 200-350 pmol/g tissue, respectively, between 1h and the post 2.5 h time-point, and both were also below 50 pmol/g tissue after 24 h. Thus, levels of both CB_1R agonists THC and 11-OH-THC peaked after 1 h of high-concentration THC-E-gel consumption, matching the hypolocomotion response measured at 40 min during the 2 h consumption period (**Figure 1**).

Furthermore, THC and 11-OH-THC levels in brain tissue were drastically lower after 24 h but still detectable, as previously reported [40-42]. THC levels in plasma reached approximately 400 pmol/g tissue at the 1 h time-point and decreased thereafter (**Figure 3c**). Statistical comparisons between the 1 h and 2 h exposure periods were limited due to different treatment paradigms, prompting the normalization of all PK values to the 1 h exposure period samples (**Figure 3d-e**). Brain samples were all increased at 2 h relative to 1 h exposure but significant differences to 1 h exposure was only found at the 26 h collection time point (**Figure 3d**, One-way ANOVA, $F_{3,139} = 14.03$, $p < 0.001$, Sidak's). Alternatively, plasma samples were significantly decreased at 2 h for THC and COOH-THC while all three compounds were significantly decreased at the 26 h collection time point (**Figure 3d**, One-way ANOVA, $F_{3,141} = 35.23$, $p < 0.001$, Sidak's). Correlation of PK findings with cannabimimetic triad results did not reveal any significant relationships. Note that 11-OH-THC and 11-COOH-THC levels peaked after 2 h of consumption which contrasts with the early onset hypolocomotive response measured in **Figure 1** after 40 min of gelatin access. Thus, PK analysis of high-concentration THC-E-gel consumption demonstrates parallel accumulation of THC and 11-OH-THC in the brain, a unique profile that differs compared to previously established PK profile resulting from THC-*i.p.* injection [40,43].

Modeling *i.p.* THC and THC-E-gel triad cannabimimetic responses predicts THC-E-gel-dependent behaviors.

To further establish the pharmacological relationship between *i.p.* THC injections and THC-E-gel consumption after 1 h and 2 h consumption along with the low variability in the cannabimimetic responses triggered by both routes of administrations, we calculated

"predicted THC doses" by comparing their cannabimimetic responses across experiments (**Figure 4a**). Thus, we extrapolated the relative *i.p.* dose for each cannabimimetic response triggered by consumption by plotting the cannabimimetic response following consumption onto the dose-response curve of THC-*i.p.* as reference (**Figure 4b-d**). **Figure 4b-d** also shows that 1 h access to high-concentration THC-E-gel triggered greater cannabimimetic responses compared to 2 h access. Consequently, this resulted in a higher "predicted *i.p.* dose" shown by dotted lines tracked to the *i.p.* dose-response curves. Of note, 1 h access to high-concentration THC-E-gel triggered stronger hypolocomotion and reduction in core body temperature corresponding to 10.3 and 11.6 mg/kg THC *i.p.*, respectively, and analgesia corresponding to 4.5 mg/kg THC *i.p.* (**Figure 4b-d**). By contrast, 2 h access to high-concentration THC-E-gel triggered a comparable response in the three cannabimimetic behaviors corresponding to 3-4 mg/kg THC *i.p.* (**Figure 4b-d**). **Figure 4e** illustrates the predictive value of these calculations, and the larger variability for the 1 h access predicted dose of 8.81 ± 2.19 mg/kg *i.p.* and 3.69 ± 0.25 mg/kg *i.p.* for 2 h access, a 2.4-fold higher predicted dose after 1 h access. The variability between the cannabimimetic response for the 1 h access results suggest that a difference in the PK profile of THC at 1 h compared to 2 h access (see **Figure 3**). Together, these results indicate that consumption of high-concentration THC-E-gel triggers strong cannabimimetic responses, comparable to *i.p.* injections of THC between 4-12 mg/kg, although this is not necessarily adaptable to all behavioral readouts.

THC injection and high-concentration THC-E-gel consumption reveals sex-dependent acoustic startle responses.

Acoustic startle responses in mice are a well-established preclinical approach to evaluate an unconditional reflex characterized by the rapid contraction of muscles to a sudden and intense startling stimulus. It is an especially useful measure in preclinical research as it is consistent across species and involves simple neural circuitry in sensorimotor gating [44]. It is known that *i.p.* injection of THC (6 and 10 mg/kg) reduces acoustic startle [22,34,45]. However, whether startle response is affected in a sex-dependent manner or is altered by lower dose THC is unknown. Thus, we next sought to extend studies on the effect of *i.p.* injection of THC on acute acoustic startle in male and female mice and compare these results to high-concentration THC-E-gel consumption. We measured the acute startle responses as the peak velocity (Vmax as measured by an accelerometer) following audible tones either 1 h after *i.p.* administration of THC (from 0.1 to 10 mg/kg) or immediately after access to high-concentration THC-E-gel, and immediately after treatment measured the startle response to increasing tone power (80, 90, 100, 105, 110, and 120 dB) (**Figure 5a**). THC administration via *i.p.* induced a significantly increased startle response in males at 1 and 5 mg/kg and a significantly decreased startle response in females at 10 mg/kg (**Figure 5b-c**, males: v. VEH Two-way ANOVA $F_{4,122}$ = 13.89, p < 0.001, Sidak's; females: v. VEH Two-way ANOVA $F_{4,116}$ = 6.761, p < 0.001, Sidak's). The greatest variability in responses came from the 120 dB tone exposures. To this end, we plotted male and female 120 dB tone startle responses to all four doses of *i.p.* THC tested, which produced a biphasic response to increasing doses of *i.p.* THC in males but not females (**Figure 5d**). This resulted in a statistically significant difference between males and females (**Figure 5d**, Two-way ANOVA $F_{3,23}$ = 26.66, p <0.001, Sidak's). THC-E-gel consumption by males and females triggered a remarkably different startle response, here characterized by only males exhibiting an increase in response when allowing access for 2 h to THC-E-gel,

leading to a 2.2-fold increase in the startle response to 120 dB (**Figure 5e-f**, males: v. VEH Two-way ANOVA $F_{2,70} = 26.85$, p < 0.001, Sidak's). These results indicate that THC administered *i.p.* induces an inverted U-shape impairment of acoustic startle responses that is more pronounced in males than in females; and that only males exhibit an increased acoustic startle response when exposed for 2 h to 10 mg THC-E-gel. To emphasize the consumption-dependent effects we correlated the 120 dB startle responses with the respective gelatin dose consumed (10 mg THC-E-gel) which produced a near significant correlation (**Figure 5g**, p = 0.051). Using the male and female-specific dose-responses in **Figure 5d**, and the predicted *i.p.* dose described in **Figure 4e**, an acoustic startle response (Vmax) to a 120 dB tone following 10 mg THC-E-gel consumption was predicted as: in cm/min, 389.9 (1 h) and 708.8 (2 h) for females, and 900.7 (1 h) and 1854.3 (2 h) for males. **Figure 5h-i** compares the acoustic startle response to a 120 dB tone of male and female mice and the *i.p.* predicted response after 1 h and 2 h access to 10 mg THC-E-gel. Predicted startle responses in males exposed to 10 mg THC-E-gel for both 1 and 2 h access was within the standard error of the measured startle response to a 120 dB tone, whereas only the 1 h predicted startle response in females was within the standard error of the measured response (**Figure 5h-i**). This prediction of THC-impaired psychomotor reflexive behavior after a temporally modified access paradigm further illustrates the robust nature of our new THC-E-gel consumption model as an effective behavioral paradigm for investigating voluntary 10 mg THC consumption.

Discussion

Here we report a novel behavioral model that enables the examination of the behavioral impact of voluntary oral consumption of high-concentration THC-E-gel by adult mice. Access to E-gel for 2 h over a two-day period incentivizes robust consumption, and at the highest dose tested

here (10 mg), mice of both sexes consumed ~30 mg/kg THC in 2 h on the second day. Acute consumption of THC triggers commonly established cannabimimetic responses, the potencies of which were right shifted compared to the responses measured with *i.p.* injections. Furthermore, we discovered that acute consumption of 10 mg THC-E-gel increases the acoustic startle response in males to a greater extent than it does in females; and that *i.p.* injection of THC triggers a dose-dependent, inverse U-shaped, impairment of acoustic startle response that was also more pronounced in males than females. Our study provides important translational results at two levels: acute consumption of THC by rodents and its impact on acoustic startle response as a measure for psychomotor reflexive behavior.

Mice of both sexes consumed similar amounts of VEH-CTR-gel and VEH-E-gel, and none consumed more than 20% of their daily caloric intake indicating comparable consumption behaviors. However, consumption of high-concentration THC-CTR-gel (4 mg) was inconsistent, and 41% of the mice completely avoided consumption (as assessed by an unbroken gelatin surface at the end of the 2 h access period) (**Figure 1c**). By contrast, consumption of 10 mg THC-E-gel (10 mg, i.e., 2.5X more concentrated) was consistent with a total consumption rate of 0.95 g/2 h (**Figure 1d**). This difference in consumption between THC-CTR-gel (4 mg) and THC-E-gel (10 mg) is likely due to the chocolate flavoring in Ensure™ that masks the strong odor and bitter taste of high-concentration THC and its aversive properties. Significantly, mice that consumed the high-concentration THC-E-gel on Day 2 consumed remarkably less VEH-E-gel on Day 3 (**Supplementary Figure S2c**). This is potentially due to the development of aversive conditioned associations to high-concentration THC associated with the E-gel. Thus, the novel experimental model reported here also enables the study of aversive memory to high-concentration THC during voluntary oral consumption.

I.p. injection of THC induces hypolocomotion, analgesia, and hypothermia in mice with different median effective doses (ED_{50}, 1.3, 3.9 and 14.4 mg/kg, respectively) (**Figure 2b-d**). By comparison, 1 h access to 10 mg THC-E-gel produced cannabimimetic responses that paralleled the ED_{50} of *i.p.* injections and are equivalent to an *i.p.* dose of ~9 mg/kg. Also, 1 h access to 10 mg THC-E-gel evoked a more pronounced cannabimimetic response compared to 2 h access, agreeing with prior studies which have shown that oral gavage increases brain peak concentration of THC 1-2 h after administration [46]. Oral consumption also increases 11-OH-THC levels in the brain with comparable kinetics and concentration as THC, and the levels of both cannabinoids decrease in parallel (**Figure 3b-c**). Considering that ~600 pmol/g of THC and 11-OH-THC is roughly equivalent to 3 nM of both compounds in the brain that persists over several hours, and both activate CB_1R with comparable potencies, our results suggest that the accumulation of both THC and 11-OH-THC in the brain might contribute to cannabimimetic responses [47]. Interestingly, we found that oral consumption of THC-E-gel produced a higher brain concentration of the primary metabolite 11-OH-THC in the brain compared to previously published concentrations after *i.p.* administration [40]. This suggests oral administration may modify the accumulation of 11-OH-THC or its metabolism in the brain. Finally, considering that voluntary oral consumption of 10 mg THC results in nanomolar concentrations of THC and 11-OH-THC for several hours, the time-dependent reduction in cannabimimetic response that follows their maximal response may be due either to CB_1R desensitization/tolerance or to redistribution of the drug within brain parenchyma.

An *i.p.* injection of THC 6 and 10 mg/kg in male mice reduces acoustic startle behaviors [34,45]. We show here that THC-*i.p.* induces a dose-dependent biphasic behavioral response that is more pronounced in males than females, demonstrating sex-dependent sensorimotor

behaviors, and confirming that THC impacts neurocognitive function in a sex-dependent manner (**Supplementary Figure S3**) [48-50]. THC-E-gel (10 mg) consumption also increased the response to acoustic startle preferentially in males compared to females. Whether the dose of THC formulated in E-gel can be increased to levels that remain palatable to mice and might trigger the pronounced reduced acoustic startle measured with 10 mg THC injection *i.p.* remains an open question. Analysis of the behavioral responses following *i.p.* injection and consumption of THC-E-gel enabled us to propose a model that correlates the doses of THC capable of producing comparable behavioral responses. The flexibility of the THC-E-gel experimental approach may extend its utility as a substitute for traditional *i.p.* injections, bridging the translational gap between preclinical investigations and human use. For example, the THC-E-gel experimental model can be easily modified and implemented to measure, in a less invasive manner, additional mouse behaviors including self-administration and preference/aversion, paradigms that require multiple treatment regimens.

In conclusion, our study outlines a new experimental model that achieves robust voluntary oral consumption of THC in adult mice by formulating THC in a chocolate-flavored sweetened E-gel. Given the recent rise in use of high-dose *Cannabis* products such as high concentration edibles [51], the model allows for the relevant and important translational investigation of sensitive behaviors such as investigations of psychomotor reflexes in mice that voluntarily consume the drug.

Author Contributions: A.E., D.P., M.B., N.S., and B.L. conceived and designed the study. A.E., F.U., A.T., D.S., and A.S. acquired the data. A.E., F.U., A.T., D.S., and A.S. performed the analysis. All authors have edited and approved the final manuscript.

Funding: NIDA F31 DA055448-01 to A.E. with N.S. and M.B. Research support from the UW

Addictions, Drug & Alcohol Institute (ADAI) Small Grants Program and the Washington State

Dedicated Cannabis Fund to A.E., NCCIH R01 AT011524-01A1 to B.L., NIDA R21 DA051558-

02 to N.S. and B.L., NIDA R37 DA033396-10 to M.R.B., Center of Excellence in Neurobiology

of Addiction, Pain, and Emotion NIDA P30 DA048736 to M.R.B. (Charles Chavkin PI), Center

of Excellence ICAL (Impact of Cannabinoids across the Lifespan) P50DA044118-01 to D.P.

Competing Interests: The authors have nothing to disclose.

REFERENCES:

Hasin DS, Shmulewitz D, Sarvet AL. Time trends in US cannabis use and cannabis use disorders overall and by sociodemographic subgroups: a narrative review and new findings. The American journal of drug and alcohol abuse. 2019;45(6):623-43.

Compton WM, Han B, Jones CM, Blanco C, Hughes A. Marijuana use and use disorders in adults in the USA, 2002–14: analysis of annual cross-sectional surveys. The Lancet Psychiatry. 2016;3(10):954-64.

Carlini BH, Schauer GL. Cannabis-only use in the USA: prevalence, demographics, use patterns, and health indicators. Journal of cannabis research. 2022;4(1):1-8.

Martin-Santos R, a Crippa J, Batalla A, Bhattacharyya S, Atakan Z, Borgwardt S, et al. Acute effects of a single, oral dose of d9-tetrahydrocannabinol (THC) and cannabidiol (CBD) administration in healthy volunteers. Current pharmaceutical design. 2012;18(32):4966-79.

Zuurman L, Roy C, Schoemaker R, Hazekamp A, Den Hartigh J, Bender J, et al. Effect of intrapulmonary tetrahydrocannabinol administration in humans. Journal of psychopharmacology. 2008;22(7):707-16.

Morgan CJ, Freeman TP, Hindocha C, Schafer G, Gardner C, Curran HV. Individual and combined effects of acute delta-9-tetrahydrocannabinol and cannabidiol on psychotomimetic symptoms and memory function. Translational psychiatry. 2018;8(1):1-10.

Hollister LE, Gillespie H. Delta-8-and delta-9-tetrahydrocannabinol; Comparison in man by oral and intravenous administration. Clinical Pharmacology & Therapeutics. 1973;14(3):353-57.

Hollister LE. Tetrahydrocannabinol isomers and homologues: contrasted effects of smoking. Nature. 1970;227(5261):968-69.

Weinstein A, Brickner O, Lerman H, Greemland M, Bloch M, Lester H, et al. Brain imaging study of the acute effects of Δ 9-tetrahydrocannabinol (THC) on attention and motor coordination in regular users of marijuana. Psychopharmacology. 2008;196:119-31.

Weinstein A, Brickner O, Lerman H, Greemland M, Bloch M, Lester H, et al. A study investigating the acute dose—response effects of 13 mg and 17 mg Δ 9-tetrahydrocannabinol on cognitive—motor skills, subjective and autonomic measures in regular users of marijuana. Journal of psychopharmacology. 2008;22(4):441-51.

Hindley G, Beck K, Borgan F, Ginestet CE, McCutcheon R, Kleinloog D, et al. Psychiatric symptoms caused by cannabis constituents: a systematic review and meta-analysis. The Lancet Psychiatry. 2020;7(4):344-53.

Barrett FS, Schlienz NJ, Lembeck N, Waqas M, Vandrey R. "Hallucinations" following acute cannabis dosing: a case report and comparison to other hallucinogenic drugs. Cannabis and Cannabinoid Research. 2018;3(1):85-93.

Lira MC, Heeren TC, Buczek M, Blanchette JG, Smart R, Pacula RL, et al. Trends in cannabis involvement and risk of alcohol involvement in motor vehicle crash fatalities in the United States, 2000⬜2018. American journal of public health. 2021;111(11):1976-85.

Moore TH, Zammit S, Lingford-Hughes A, Barnes TR, Jones PB, Burke M, et al. Cannabis use and risk of psychotic or affective mental health outcomes: a systematic review. The Lancet. 2007;370(9584):319-28.

Holtzman D, Lovell RA, Jaffe JH, Freedman DX. 1-Δ9-tetrahydrocannabinol: neurochemical and behavioral effects in the mouse. Science. 1969;163(3874):1464-67.

Beardsley PM, Scimeca JA, Martin BR. Studies on the agonistic activity of! D9–21-tetrahydrocannabinol in mice, dogs and rhesus monkeys and its interactions with! D—9-tetrahydrocannabinol. The Journal of Pharmacology and Experimental Therapeutics. 1987.

Metna-Laurent M, Mondésir M, Grel A, Vallée M, Piazza PV. Cannabinoid-induced tetrad in mice. Current Protocols in Neuroscience. 2017;80(1):9.59. 1-9.59. 10.

Siemens AJ, Doyle OL. Cross-tolerance between Δ9-tetrahydrocannabinol and ethanol: the role of drug disposition. Pharmacology Biochemistry and Behavior. 1979;10(1):49-55.

Ruiz C, Torrens A, Lallai V, Castillo E, Manca L, Martinez M, et al. Pharmacokinetic and pharmacodynamic properties of aerosolized ("vaped") THC in adolescent male and female rats. Psychopharmacology. 2021;238(12):3595-605.

Kasten CR, Zhang Y, Boehm SL. Acute cannabinoids produce robust anxiety-like and locomotor effects in mice, but long-term consequences are age-and sex-dependent. Frontiers in behavioral neuroscience. 2019;13:32.

Ibarra-Lecue I, Mollinedo-Gajate I, Meana JJ, Callado LF, Diez-Alarcia R, Urigüen L. Chronic cannabis promotes pro-hallucinogenic signaling of 5-HT2A receptors through Akt/mTOR pathway. Neuropsychopharmacology. 2018;43(10):2028-35.

Long LE, Chesworth R, Huang X-F, McGregor IS, Arnold JC, Karl T. A behavioural comparison of acute and chronic Δ9-tetrahydrocannabinol and cannabidiol in C57BL/6JArc mice. International Journal of Neuropsychopharmacology. 2010;13(7):861-76.

Boucher AA, Hunt GE, Micheau J, Huang X, McGregor IS, Karl T, et al. The schizophrenia susceptibility gene neuregulin 1 modulates tolerance to the effects of cannabinoids. International Journal of Neuropsychopharmacology. 2011;14(5):631-43.

Hoffman HS, Ison JR. Reflex modification in the domain of startle: I. Some empirical findings and their implications for how the nervous system processes sensory input. Psychological review. 1980;87(2):175.

Burgdorf CE, Jing D, Yang R, Huang C, Hill MN, Mackie K, et al. Endocannabinoid genetic variation enhances vulnerability to THC reward in adolescent female mice. Science Advances. 2020;6(7):eaay1502.

Lepore M, Vorel SR, Lowinson J, Gardner EL. Conditioned place preference induced by Δ9-tetrahydrocannabinol: comparison with cocaine, morphine, and food reward. Life sciences. 1995;56(23-24):2073-80.

Kruse LC, Cao JK, Viray K, Stella N, Clark JC. Voluntary oral consumption of Δ9-tetrahydrocannabinol by adolescent rats impairs reward-predictive cue behaviors in adulthood. Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology. 2019.

Abraham AD, Leung EJ, Wong BA, Rivera ZM, Kruse LC, Clark JJ, et al. Orally consumed cannabinoids provide long-lasting relief of allodynia in a mouse model of chronic neuropathic pain. Neuropsychopharmacology. 2020;45(7):1105-14.

Smoker MP, Mackie K, Lapish CC, Boehm II SL. Self-administration of edible Δ9-tetrahydrocannabinol and associated behavioral effects in mice. Drug and alcohol dependence. 2019;199:106-15.

Kruse LC, Cao JK, Viray K, Stella N, Clark JJ. Voluntary oral consumption of Δ 9-tetrahydrocannabinol by adolescent rats impairs reward-predictive cue behaviors in adulthood. Neuropsychopharmacology. 2019;44(8):1406-14.

Barrus DG, Lefever TW, Wiley JL. Evaluation of reinforcing and aversive effects of voluntary Δ9-tetrahydrocannabinol ingestion in rats. Neuropharmacology. 2018;137:133-40.

Ventura R, Morrone C, Puglisi-Allegra S. Prefrontal/accumbal catecholamine system determines motivational salience attribution to both reward-and aversion-related stimuli. Proceedings of the National Academy of Sciences. 2007;104(12):5181-86.

Barbano MF, Castañé A, Martín-García E, Maldonado R. Delta-9-tetrahydrocannabinol enhances food reinforcement in a mouse operant conflict test. Psychopharmacology. 2009;205(3):475-87.

Nagai H, Egashira N, Sano K, Ogata A, Mizuki A, Mishima K, et al. Antipsychotics improve Δ9-tetrahydrocannabinol-induced impairment of the prepulse inhibition of the startle reflex in mice. Pharmacology Biochemistry and Behavior. 2006;84(2):330-36.

Schindler AG, Tsutsui KT, Clark JJ. Chronic alcohol intake during adolescence, but not adulthood, promotes persistent deficits in risk-based decision making. Alcoholism: clinical and experimental research. 2014;38(6):1622-29.

Falenski KW, Thorpe AJ, Schlosburg JE, Cravatt BF, Abdullah RA, Smith TH, et al. FAAH−/− mice display differential tolerance, dependence, and cannabinoid receptor adaptation after Δ9-tetrahydrocannabinol and anandamide administration. Neuropsychopharmacology. 2010;35(8):1775-87.

Varvel SA, Bridgen DT, Tao Q, Thomas B, Martin B, Lichtman A. Δ9-Tetrahydrocannbinol accounts for the antinociceptive, hypothermic, and cataleptic effects of marijuana in mice. Journal of Pharmacology and Experimental Therapeutics. 2005;314(1):329-37.

Ettaro R, Laudermilk L, Clark SD, Maitra R. Behavioral assessment of rimonabant under acute and chronic conditions. Behavioural brain research. 2020;390:112697.

Wright F, Rodgers R. Low dose naloxone attenuates the pruritic but not anorectic response to rimonabant in male rats. Psychopharmacology. 2013;226:415-31.

Torrens A, Vozella V, Huff H, McNeil B, Ahmed F, Ghidini A, et al. Comparative pharmacokinetics of Δ9-tetrahydrocannabinol in adolescent and adult male mice. Journal of Pharmacology and Experimental Therapeutics. 2020;374(1):151-60.

Kreuz DS, Axelrod J. Delta-9-tetrahydrocannabinol: localization in body fat. Science. 1973;179(4071):391-93.

Johansson E, Norén K, Sjövall J, Halldin MM. Determination of Δ1-tetrahydrocannabinol in human fat biopsies from marihuana users by gas chromatography–mass spectrometry. Biomedical Chromatography. 1989;3(1):35-38.

Vozella V, Zibardi C, Ahmed F, Piomelli D. Fast and sensitive quantification of Δ9-tetrahydrocannabinol and its main oxidative metabolites by liquid chromatography/tandem mass spectrometry. Cannabis and Cannabinoid Research. 2019;4(2):110-23.

Pantoni MM, Herrera GM, Van Alstyne KR, Anagnostaras SG. Quantifying the acoustic startle response in mice using standard digital video. Frontiers in Behavioral Neuroscience. 2020;14:83.

Tournier BB, Ginovart N. Repeated but not acute treatment withΔ 9-tetrahydrocannabinol disrupts prepulse inhibition of the acoustic startle: Reversal by the dopamine D2/3 receptor antagonist haloperidol. European Neuropsychopharmacology. 2014;24(8):1415-23.

Deiana S, Watanabe A, Yamasaki Y, Amada N, Arthur M, Fleming S, et al. Plasma and brain pharmacokinetic profile of cannabidiol (CBD), cannabidivarine (CBDV), Δ9-tetrahydrocannabivarin (THCV) and cannabigerol (CBG) in rats and mice following oral and intraperitoneal administration and CBD action on obsessive–compulsive behaviour. Psychopharmacology. 2012;219(3):859-73.

Dinis-Oliveira RJ. Metabolomics of Δ9-tetrahydrocannabinol: implications in toxicity. Drug metabolism reviews. 2016;48(1):80-87.

Cha YM, Jones KH, Kuhn CM, Wilson WA, Swartzwelder HS. Sex differences in the effects of Δ9-tetrahydrocannabinol on spatial learning in adolescent and adult rats. Behavioural pharmacology. 2007;18(5-6):563-69.

Harte LC, Dow-Edwards D. Sexually dimorphic alterations in locomotion and reversal learning after adolescent tetrahydrocannabinol exposure in the rat. Neurotoxicology and teratology. 2010;32(5):515-24.

Gur RC, Richard J, Calkins ME, Chiavacci R, Hansen JA, Bilker WB, et al. Age group and sex differences in performance on a computerized neurocognitive battery in children age 8– 21. Neuropsychology. 2012;26(2):251.

Freeman TP, Craft S, Wilson J, Stylianou S, ElSohly M, Di Forti M, et al. Changes in delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) concentrations in cannabis over time: systematic review and meta-analysis. Addiction. 2021;116(5):1000-10.

Figure 1: <u>E-gel promotes heightened voluntary oral consumption of THC and induces cannabimimetic behaviors by adult mice.</u>

a) Mice were given free access to vehicle (VEH), or THC formulated in either CTR-gel or E-gel for 2 h on Day 1 and Day 2. **b)** Consumption was determined by weighing gelatin at the end of each session. **c)** Consumption of CTR-gel on Day 2 is decreased after addition of THC. **d)** Consumption of E-gel on Day 2 is maintained after addition of THC. **e)** Dose of THC consumed, in mg/kg, when formulated in either CTR-gel or E-gel on Day 2. Results are mean ± S.E.M. Consumption compared ANOVA and Sidak's, *$p<0.05$, **$p<0.01$, and ***$p<0.001$, N=8-40. **f)** Diagram of behavioral paradigm before and after *i.p.* or gelatin administration. **h-j)** Dose-dependent behavioral responses for hypolocomotion (**h**), analgesia (**i**), and hypothermia (**j**) after THC exposure. Administration by *i.p.* (grey) is plotted on x-axis by single bolus injection while CTR-gel (green) and E-gel (purple) are plotted based on average THC consumed after 2 h exposure window shown in **e**. **k)** Diagram of THC-E-gel exposure, behavioral measurements, and SR1 injection (by *i.p.*) at 1 h into exposure window. **l-n)** Individual behavioral responses for hypolocomotion (**l**), analgesia (**m**), and hypothermia (**n**) for each animal. Individual points are plotted based on individual THC consumption with a linear regression to show correlation between consumed THC and behavioral output (p-values: **l**=0.003, **m**<0.001, **n**<0.001). SR1 treated mice are plotted (red) based on consumed THC after exposure to 10 mg/15 ml THC-E-gel with a linear regression to show no correlation across three behaviors (p-values: **l**=0.09, **m**=0.44, **n**=0.45).

Figure 2: <u>THC-E-gel consumption triggers CB₁R-dependent behaviors</u>

a) Over a 3-day exposure paradigm, mice received 3 days of E-gel with either VEH or THC (10mg/15ml) E-gel on day 2. **b-d)** Cumulative gelatin consumption recorded every 10 minutes throughout the 2 h exposure window over the 3-day paradigm. VEH (black) and THC (purple) groups received access to VEH on day 1 (**b**), VEH or THC on day 2 (**c**), and VEH on day 3 (**d**). **e)** Total gelatin consumption after 2 h of access to gelatin was plotted comparing VEH and THC treatment groups. **f)** Animal consummatory and locomotor behavior was tracked during gelatin exposure window. **g)** Distance traveled recorded every 5 min over the 3-day paradigm, similar as to **b-d. h)** Total distance traveled (cm) after 2 h of gelatin access was plotted

comparing VEH and THC groups. Main effect over 2 h exposure period (**b-d, g**) measured using Two-way ANOVA with repeated measures and Sidak's, main effect on total response (**e, h**)measured by One-way ANOVA and Sidak's (*p<0.05, **p<0.01, ***p<0.001) N=8-16.

Figure 3: Consumption of THC-E-gel results in concomitant increases in the levels of THC and its metabolites in brain tissue

a) Diagram outlining gelatin exposure paradigm where blood and brain samples were collected immediately following 1 h and at 2, 2.5, and 26 hours from the beginning of 2 h access to 10mg/15ml THC-E-gel. **b)** Brain concentration of THC, 11-OH-THC, and COOH-THC after E-gel exposure, 1 h access is separated due to a reduced total access time to THC-E-gel compared to the other time points. **c)** Plasma concentrations for the three compounds plotted similarly to b. **d-e)** PK concentrations in brain **(d)** and plasma **(e)** normalized to the 1 h access period. Statistical comparison to 1 h Two-way ANOVA, Sidak's, *p<0.05, **p<0.01, and ***p<0.001, N=8-15.

Figure 4: Modeling *i.p.* THC and THC-E-gel triad cannabimimetic responses predicts THC-E-gel-dependent behaviors.

a) Diagram of 1 h and 2 h THC-E-gel exposure and *i.p.* administration with behavioral tests. **b-d**) Cannabimimetic responses after THC administration by *i.p.* and subsequent dose-response curve in grey. Responses after 1 h or 2 h exposure to 10 mg THC-E-gel are plotted with dotted lines tracking to relative THC-*i.p.* dose response. **e**) Predicted *i.p.* dose after 1 h and 2 h THC-E-gel exposure window from all three triad behaviors.

Figure 5: Sex-dependent acoustic startle responses after *i.p.* injection of THC and high-concentration THC-E-gel consumption

a) Diagram of THC-E-gel exposure or *i.p.* administration followed by acoustic startle response behavioral testing. **b-c)** Male and female acoustic startle responses after *i.p.* administration of THC in response to escalating tones (80, 90, 100, 105, 110, and 120dB) following *i.p.* administration of THC in males (**b**) and females (**c**). **d)** Male and female acoustic startle dose-

responses to a 120dB tone after *i.p.* THC administration. Results are mean ± S.E.M. One-way ANOVA, Sidak's comparing VEH and *i.p.* THC dose between males and females **p<0.01, and ***p<0.001, N=6-11. **e-f)** Male and female acoustic startle responses after 1 h or 2 h THC E-gel exposure in response to escalating tones (80, 90, 100, 105, 110, and 120dB. **g-h)** Startle response to a 120dB tone for males (**g**) and females (**h**) after 1 h or 2 h access to THC E-gel. Predicted doses calculated from a second order polynomial of *i.p.* dose responses are plotted to show the consistency in predicted dose response after E-gel exposure.

a

CTR-gel

Day 1: VEH-**CTR-gel**
2 h

Day 2: THC-**CTR-gel**
2 h

E-gel

Day 1: VEH-**E-gel**
2 h

Day 2: THC-**E-gel**
2 h

b

E-gel → grams (g) consumed

1, 2, 5, or 10 mg THC/15 ml E-gel → X mg THC

c CTR-gel consumed (g)

[THC] (mg/15ml)
VEH 0.3 1 2 4
** ** ***

d E-gel consumed (g)

[THC] (mg/15ml)
VEH 1 2 5 10
*** ***

e THC consumed (mg/kg)

CTR-gel   E-gel

[THC] (mg/15ml)
0.3 1 2 4   1 2 5 10
*   * *** ***

f

0.3, 1, 2, and 4 mg
OR
1, 2, 5, and 10 mg

Pre-Tests | 2 h | Post-Tests
OR
Pre-Tests | 1 h | Post-Tests

b Open Field   c Tail Flick   d Body Temp.

h Distance traveled (cm, Post-Pre or -VEH)

i.p.
CTR-gel
E-gel

VEH   0.1   1   10
THC

i Tail flick withdrawal (s, Post-Pre)

VEH   0.1   1   10
THC

j Body temperature (Celsius, Post-Pre)

VEH   0.1   1   10
THC

k

1, 2, 5, and 10 mg

Pre-Tests | 1 h | 1 h | Post-Tests
SR1 onboard

SR141617 (SR1)

THC-E-gel   SR1 + THC-E-gel

l Distance traveled (cm, Post-Pre)

p = 0.003
p = 0.09

THC (mg/kg)

m Tail flick withdrawal (s, Post-Pre)

p < 0.001
p = 0.45

THC (mg/kg)

n Body temperature (Celsius, Post-Pre)

p < 0.001
p = 0.44

THC (mg/kg)

a

	Day 1	Day 2	Day 3
	VEH 2 h	VEH 2 h	VEH 2 h
	VEH 2 h	THC 2 h	VEH 2 h

f

VEH E-gel or
10mg/15ml THC E-gel

b-d **Consumption**

b Day 1: VEH

c Day 2: VEH or 10mg THC

d Day 3: VEH

Total gelatin consumed (g)

Time (min)

g **Locomotion**

Day 1: VEH

Day 2: VEH or 10mg THC

Day 3: VEH

Distance traveled (cm/5 min)

Time (min)

e

Total gelatin consumed (g)

VEH / VEH or THC / VEH

Exposure days

1   2   3

***   ***

VEH   THC

h

Distance traveled (cm)

VEH / VEH or THC / VEH

Exposure days

1   2   3

*

VEH   THC

a
Pre-Tests | 1 h | Post-Tests

OR

Pre-Tests | 2 h | Post-Tests

OR

Pre-Tests | 1 h | Post-Tests

b

Distance traveled (% control)

THC (mg/kg)

○ i.p.
● E-gel 2 h
● E-gel 1 h

4.27 **11.6**

c

Tail flick withdrawal (s, Post-Pre)

THC (mg/kg)

3.42 **4.50**

d

Body temperature (DegC, Post-Pre)

THC (mg/kg)

3.91 **10.34**

e

Predicted Dose THC (mg/kg) by *i.p.*

Exposure window

■ E-gel 2 h
■ E-gel 1 h