Caught in the Act: targeted mutagenesis of the herpesvirus glycoprotein B central helix

captures fusion transition states

Momei Zhou[1*], Ben Vollmer [4], Emily Machala[4], Muyuan Chen[6], Kay Grünewald [4], Ann M.

Arvin[1,2,3], Wah Chiu [2,5,6], and Stefan L. Oliver[1]

*Corresponding author: mzhou6@stanford.edu

[1] Department of Pediatrics, Stanford University School of Medicine, Stanford, California, USA.

[2] Department of Microbiology & Immunology, Stanford University School of Medicine,

Stanford, California, USA.

[3] Vir Biotechnology Inc, USA.

[4] Centre for Structural Systems Biology, University of Hamburg, Leibniz Institute of Virology

(LIV), Hamburg, Germany.

[5] Bioengineering, Stanford University School of Medicine, Stanford, CA 94305, USA.

[6] Division of Cryo-EM and Bioimaging SSRL, SLAC National Accelerator Laboratory, Menlo

Park, CA 94025, USA.

Key words

Herpesvirus, glycoprotein B, gB central helix structure, prefusion gB, gB biosynthesis, gB fusion

function, viral replication

Running title

Central helix determines VZV gB structure, fusion function and biosynthesis

Abstract

Herpesviruses remain a burden for animal and human health, including the medically

important varicella-zoster virus (VZV). Membrane fusion mediated by conserved core

glycoproteins, the fusogen gB and the heterodimer gH-gL, enables herpesvirus cell entry. The

ectodomain of gB orthologs has five domains and is proposed to transition from a prefusion to

postfusion conformation but the functional relevance of the domains for this transition remains

poorly defined. Structure-function studies of the VZV gB DIII central helix were performed

targeting residues ^{526}EHV^{528}. Critically, a H527P mutation captured gB in a prefusion

conformation as determined by cryo-EM, a loss of membrane fusion in a virus free assay, and

failure of recombinant VZV to spread in cell monolayers. Importantly, two predominant cryo-

EM structures of gB[H527P] were identified by 3D classification and focused refinement,

suggesting they represented gB conformations in transition. These studies reveal gB DIII as a

critical element for herpesvirus gB fusion function.

Introduction

The *Herpesviridae* encompass pathogens that cause a wide range of medically and economically important diseases of humans and animals. All herpesviruses are enveloped and use glycoproteins to initiate and drive membrane fusion between the virion and the host cell during entry. The three glycoproteins, gB, gH, and gL, required for this process are conserved across the *Herpesviridae* [1,2]. Structure-function studies have revealed that the gB trimer is the primary fusogen and the gH-gL heterodimer has roles in cell tropism and gB regulation [3-10]. Incorporation of gB into the envelope of herpesvirus particles is complex. gB is transported to the plasma membrane of the infected cell, endocytosed, and trafficked to the trans-Golgi network for insertion into the herpesvirus envelope during morphogenesis [11]. To achieve fusion with cellular membranes, gB on the virion envelope is thought to undergo a conformational transition from a meta-stable prefusion structure to an energetically more stable postfusion state [12].

VZV, a member of the *Alphaherpesvirinae* and human host restricted pathogen, causes varicella (chickenpox), establishes latency in sensory ganglion neurons, and can reactivate to manifest as zoster (shingles) [13]. In addition to membrane fusion required for virion entry, gB dependent cell-cell fusion is a hallmark of VZV pathogenesis. Characteristic polykaryocytes form within infected tissues *in vivo* and are modeled *in vitro* by syncytia formation between MeWo cells [3]. VZV gB, together with gH-gL, are necessary and sufficient to induce cell-cell fusion *in vitro* [6,14-17]. Unlike its HSV gB orthologues, VZV gB is modified by furin cleavage post-translationally, which is needed for optimal skin lesion formation *in vivo* [14]. Critically, VZV induced fusion between differentiated cells has been linked to adverse health effects, including fusion of ganglion neurons and satellite cells associated with postherpetic neuralgia (PHN), and fusion of endothelial cells in the brain vasculature linked to strokes [18,19].

Despite substantial amino acid sequence differences, the gB orthologues from alpha-, beta-

and gamma-herpesvirus subfamilies adopt remarkably similar structures in their postfusion

ectodomain conformations, consisting of homotrimers with five domains, DI-DV [2,20-27]. This

architecture led to the classification of gB orthologues as type III fusogens based on similarities

to postfusion forms of vesicular stomatitis virus glycoprotein G (VSV-G) and the baculovirus

gp64 [28,29]. In the gB trimer, DI contains the fusion loops, DII has a pleckstrin homology domain,

DIII forms a central helical bundle, DIV forms the crown, and DV connects the crown to the

transmembrane domain. Our recent single particle cryogenic electron microscopy (cryo-EM)

studies of full-length VZV gB have demonstrated the importance of gB DIV in fusion initiation

[2,20]. In contrast, the rapid conversion of gB to its postfusion form upon deletion of the

transmembrane domain or removal from a lipid bilayer has hindered deriving high-resolution

structures of prefusion gB [21,27,30,31]. Extracellular vesicles (EVs) were used as a surrogate for

structural analyses of virion envelope-associated glycoproteins, overcoming effects on the native

topology of the protein after detergent solubilization and reconstitution into an artificial

membrane [32]. Cryogenic electron tomography (cryo-ET) and single particle cryo-EM have

revealed prefusion structures for herpes simplex virus 1 (HSV-1) and human cytomegalovirus

(HCMV) gB anchored in native membranes [31,33-36]. Expression on EVs of a HSV-1 mutant with

a proline substitution at histidine 516 in the DIII central helix yielded a condensed form of gB

that allowed assignment of DI-DV within a 9Å resolution cryo-ET structure [31]. In this structure,

DI and DII from each of three protomers created a tripod that encapsulated DIV, which was

inverted compared to the postfusion structure, with DIII protruding from the top of the prefusion

gB trimer while DV was centrally located and connected the ectodomain via transmembrane

helices to the cytoplasmic domain. A 3.6Å resolution single particle cryo-EM structure of

HCMV gB extracted from purified virus particles with an architecture similar to HSV-1

gB[H516P] was achieved using a small molecule fusion inhibitor (WAY-174865) and chemical

cross-linking [36]. Although these structural studies provide information about the stationary

phases of herpesvirus gB in pre- and postfusion conformations, the transition between these two

forms is not understood.

Here, we demonstrate the critical role of the gB DIII central helix using structure-function

studies of gB from the model pathogen VZV. Individual proline substitutions of ^{526}EHV^{528}

within VZV gB DIII inhibited gB/gH-gL mediated cell-cell fusion. Cryo-ET revealed a

condensed form of gB[H527P] expressed on EVs. Importantly, molecular classification of the

gB trimers on EVs revealed two condensed forms, suggesting a stalled process of conformational

change instigated by substitution of the conserved histidine at this position. In addition, the

prefusion state of VZV gB[H527P] was supported by binding of anti-gB neutralizing antibody,

mAb 93k, in cells transiently expressing this mutant. Critically, proline substitutions of

^{526}EHV^{528} in the VZV pOka genome were fully compatible with intracellular assembly of virus

particles and their release at cell surfaces, as demonstrated by a novel flow cytometry-based

transmission electron microscopy (FC-TEM) assay. However, only the pOka-gB[E526P] mutant

yielded infectious VZV although gB-dependent cell-cell spread was markedly impaired by this

mutation. Inactivation of pOka-gB[V528P] was explained by disrupted ER egress, preventing gB

maturation and trafficking to the cell surface, associated with a drastically distorted central helix

structure. Detection of complete viral particles despite gB ER retention suggested that virus

assembly can proceed without gB incorporation into the virion envelope. Thus, the structural

integrity of the gB DIII central helix is necessary not only for the molecular transition of gB for

fusogenic function but also for essential steps in gB biosynthesis required to produce infectious

herpesviruses.

Results

The gB[H527P] mutation constrains VZV gB in a prefusion conformation. VZV gB

residues, [526]EHV[528], were within the upper region of the DIII central helix based on our

postfusion structure, and in the predicted prefusion structure modeled on HSV-1 prefusion gB

[2,20,31] (**Fig. 1A-1C**). Conservation of the histidine at position 527 across the *alphaherpesvirinae*

(**Fig. S1**), suggested that a proline substitution might fix VZV gB in a prefusion form as

observed with HSV-1 gB[H516P] [31]. The valine at position 528 is conserved in the

varicelloviruses, the simplexviruses, and several more distant alphaherpesviruses whereas the

glutamic acid at 526 is unique to VZV and one non-human primate varicellovirus (**Fig. S1**).

Although gB was detectable in all cell lysates from BHK-21 cells transiently transfected with

vectors expressing WT gB or proline substitutions at [526]EHV[528], only WT gB and gB[H527P]

were incorporated into EVs (**Fig. S2**). A preliminary cryo-ET dataset revealed that WT gB and

gB[H527P] formed different conformations in the membranes of purified EVs from transfected

BHK-21 cells (**Fig. 1D**). WT gB was 160Å in height, and resembled the 2.8 Å postfusion

structure of gB purified from VZV infected cells [2]. DI-IV were discernable, but DV was not,

consistent with its location in postfusion gB. In contrast, the gB[H527P] trimers formed squat

structures of 100Å in height and 100Å in width that resembled those of HSV-1 gB[H516P],

indicating that the H527P substitution prevented the transition of gB to a postfusion

conformation structure (**Fig. 1D**).

To achieve a higher resolution structure of gB[H527P] on EVs, a more extensive cryo-ET

dataset was generated and coupled with subtomogram averaging (STA) (**Fig. S3; Table. S1**).

Clusters of gB[H527P] were evident with extensive edge contacts between the gB molecules on

the surface of EVs (**Fig. 2A**). STA of gB[H527P] on EVs produced an initial cryo-EM map of

10Å that also revealed extensive intermolecular contacts (**Fig. 2B**). The overall architecture of

gB[H527P] suggested a squat form similar to HSV-1 gB[H516P] where DI and DII formed a

tripod around DIV, which was inverted toward the membrane compared to the postfusion form

of gB, and DIII underwent a slight bend and twist that exposed this domain at the top of the

trimeric molecule (**Fig. 1B and 2B**). However, the decline of the "gold-standard" Fourier Shell

Correlation (FSC) curve at low resolution (20Å), in addition to the local resolution estimation

suggested that large scale conformational heterogeneity might be present in the system (**Fig. 2C**).

Furthermore, the upper central region of the gB trimer, where DIII was predicted to emerge, had

a lower local resolution compared to the rest of the protein, also indicative of heterogeneity in

the reconstruction. Two predominant classes with distinct architectures were revealed by

subtomogram classification and focused refinement of the central region of the gB trimer (**Fig.**

2D). The separation of the data into two classes improved the consistency of the cryo-EM map

resolution (12-15Å), which is also evident by the improved shape of the FSC curves, suggesting

the presence of two gB architectures on the surface of gB[H527P] induced EVs. The VZV gB

prefusion model predicted from the HSV-1 gB[H516P] structure fit only partially into the STA

maps for each class (**Fig. 2E and 2F; Movie. S1**). Although the tripod shape of the squat form of

gB was retained, a considerable lateral movement of the gB fusion loops in DI (**Fig. 2E and 2F**

box i; Movie. S1) and rotation in the region of gB DII and III (**Fig. 2E and 2F box ii; Movie.**

S1) was noted by comparison of the two classes. Thus, the substitution of proline at gB H527

prevents the transition of VZV gB to a postfusion conformation but allows movement of the gB

trimer between a closed (class I) and open (class II) squat form. These data provide evidence for

the first transitional steps from a pre- to postfusion conformation.

VZV neutralizing mAb 93k supports the prefusion conformation of gB[H527P].

Because production of EVs from gB[E526P] was limited and gB[V528P] yielded none, effects of

proline substitutions on the DIII central helix structure were modeled using AlphaFold2 (**Table.**

S2). Both E526P and H527P induced a modest deformation whereas V528P was predicted to

bend the helix by almost 90 degrees compared to WT gB DIII (**Fig. 3A**). Alanine substitutions of

^{526}EHV^{528} modeled to assess the contribution of E, H, and V side chains to DIII structure did not

significantly affect the central helix architecture.

The functional consequences of changes in DIII architecture due to ^{526}EHV^{528} proline

mutations was assessed using the human VZV neutralizing mAb 93k, which targets an epitope

mapped to gB DIV (the crown), and prevents VZV infection via fusion inhibition, indicating the

presence of its epitope on prefusion gB [2,20]. When the accessibility of this functional epitope was

quantified by flow cytometry to determine the frequency of CHO-DSP1 cells expressing the gB

^{526}EHV^{528} mutants and median fluorescence intensity (MFI) with levels of gB expression

normalized to WT gB (**Fig. 3B**; **Fig. S4**), the frequency of gB-expressing cells was comparable

to WT gB for all proline and alanine mutants with the exception of gB[V528P] (**Fig. 3C**). The

frequency of CHO-DSP1 cells expressing the ^{526}EHV^{528} proline mutants was consistent with the

relative band densities in blots of transfected BHK-21 cell lysates (**Fig. S2**), reflecting the steady

state abundance of those three mutant proteins. The gB[E526P] mutant had reduced MFI

compared to WT gB (**Fig. 3C**), suggesting that this mutation might generate an allosteric

conformational change in the epitope even though the DIII central helix structure was predicted

to be affected only modestly (**Fig. 3A**). MFI for gB[V528P] was markedly reduced, attributable

to disruption of the mAb 93k epitope due to gB misfolding caused by distortion of the helix (**Fig.**

3A), the significantly decreased abundance of this mutant protein (**Fig. 3C and Fig. S2**), or

likely both. Notably, 93k mAb binding to gB[H527P] was comparable to WT gB (**Fig. 3C**),

indicating preservation of the conformational 93k epitope and confirming that the gB[H527P]

cryo-EM structures represent forms of gB locked in prefusion conformations.

Targeted mutagenesis of gB ^{526}EHV^{528} disrupts gB/gH-gL dependent membrane fusion.

Given the structural changes in gB imposed by gB[H527P] and predicted for gB[E526P] and

gB[V528P] (**Fig. 2 and 3A**), together with the differences in 93k binding, the effect of these

mutations on membrane fusion was assessed using a stable reporter fusion assay (SRFA). This

assay measures gB/gH-gL mediated fusion as a surrogate for membrane fusion during virus entry

and cell-cell fusion in the absence of other VZV proteins [6,14-17]. The fusion activity for all three

proline substitutions was barely detectable compared to WT gB (**Fig. 3D; Table. 1**). Failure to

induce gB/gH-gL fusion was consistent with the inability of gB[H527P] to transition to a

postfusion conformation. Alanine mutants, gB[E526A], gB[H527A] and gB[V528A], retained

fusion activity of approximately 47%, 80% and 70% respectively compared to WT gB (**Fig. 3D;**

Table. 1). These data indicate that the DIII central helix architecture is necessary whereas the

amino acid side chain interactions of ^{526}EHV^{528} support but are not critical for gB/gH-gL

mediated fusion. This was consistent with minimal deformation of DIII by ^{526}EHV^{528} alanine

substitutions predicted by AlphaFold2 (**Fig. 3A**) and supported by the preservation of mAb 93k

epitope (**Fig. 3C**).

VZV is inactivated by gB[H527P] and gB[V528P] but not gB[E526P]. To determine the

role of gB DIII in VZV replication and cell-cell spread, proline or alanine substitutions at

^{526}EHV^{528} were transferred to pOka-TKGFP BACs. As expected, infectious VZV was recovered

from MeWo cells transfected with BACs containing the fusion competent alanine substitutions,

gB[E526A], gB[H527A] and gB[V528A], and plaque sizes were comparable to the BAC with

WT gB (**Fig. 3E and 3F; Table. 1**). However, for the proline substitutions, only pOka-TKGFP-

gB[E526P] yielded infectious VZV (**Fig. 3E; Table. 1**). Although plaque sizes of pOka-TKGFP-

gB[E526P] were significantly smaller than WT pOka-TKGFP plaques (**Fig. 3F**), expression of

this gB mutant allowed the production of virus particles that could enter neighboring cells even

though gB fusion function was severely impaired (**Fig. 3D; Table. 1**). Infectious VZV was not

recovered from the pOka-TKGFP-gB[H527P] or pOka-TKGFP-gB[V528P] BACs despite five

separate transfections and at least six passages to detect residual function or emergence of

compensatory mutations.

Like other herpesviruses, VZV genes are expressed at immediate early (IE), early (E), and

late (L) times after infection. The essential IE protein, IE62, thymidine kinase, an E protein, and

the L proteins ORF23, a minor capsid protein, and ORF31 (gB), were all produced in MeWo

cells transfected with the pOka BAC and the mutant BACs with proline substitutions of

^{526}EHV^{528} (**Fig. 4A and Fig. S5A**).

To investigate whether VZV particle assembly was disrupted by the inactivating mutations,

gB[H527P] and gB[V528P], VZV particle morphogenesis was assessed using a flow cytometry-

based transmission electron microscopy (FC-TEM) assay; VZV infected cells were enriched by

cell sorting based on the GFP fluorescence signal of TKGFP (**Fig. 4B**). Nucleocapsids were

observed in pOka-TKGFP-gB[E526P] and pOka-TKGFP-gB[H527P] BAC transfected cells at

48 hours with morphologies like those present in pOka-TKGFP BAC transfected cells, including

viral DNA containing C-capsids, and scaffold containing B-capsids or empty A-capsids (**Fig.**

4C), while none were found in pOka-TKGFP-gB[V528P] BAC transfected cells (**Fig. 4C**).

Herpesviruses produce complete particles with viral DNA containing capsids and light particles

that are enveloped and contain tegument proteins but no capsids. Both light and complete

particles were observed on the surfaces of pOka-TKGFP-gB[E526P], pOka-TKGFP-gB[V528P]

and pOka-TKGFP BAC transfected cells (**Fig. 4C**), demonstrating that proline substitutions at

E526 and V528 in gB DIII did not inhibit secondary envelopment of capsids or particle egress

across the plasma membrane. The complete virus particle on the surface of the pOka-TKGFP-

gB[V528P] BAC transfected cell confirmed that nucleocapsids could be produced despite the

mutation. Notably, a genome-containing particle was identified in the cytoplasm of pOka-

TKGFP-gB[H527P] BAC transfected cells (**Fig. S5B**), together with the light particles observed

(**Fig. 4C**), indicating that gB[H527P] did not interfere with virus particle assembly. Thus,

enveloped VZV particles could be assembled even when gB mutations prevented VZV

replication.

Disrupted biosynthesis renders gB[V528P] fusion defective. To further dissect the role

of the gB DIII central helix, the intracellular localization of gB ^{526}EHV^{528} mutants was evaluated

by confocal microscopy after transient expression in MeWo cells. Mutants with proline or

alanine substitutions were compared to WT gB, which traffics from the endoplasmic reticulum

(ER) to the trans-Golgi network (anti-TGN46) and has a punctate distribution within early

endosomes (anti-EEA1), as we reported previously [4,7] (**Fig. 5A**). All gB mutants exhibited

expected localization compared to WT gB, except for gB[V528P] which failed to move to either

the TGN or early endosomes (**Fig. 5A**), indicating severe disruption of gB intracellular

trafficking.

Next, the effects of the ^{526}EHV^{528} mutations on gB synthesis and steps in its post-

translational maturation were compared to WT gB using glycosylase endoglycosidase H (Endo

H), which removes high-mannose N-linked glycans added to proteins in the ER but not the

complex glycan modifications acquired in the TGN; peptide-N-glycosidase F (PNGase F)

removes all N-linked glycans. Without treatment, full-length gB was ~124 kDa, and the furin

cleaved C-terminal fragments were 66 kDa and 60 kDa (**Fig. 5B**). Endo H reduced full-length gB

to approximately 95 kDa, the same size as gB treated by PNGase F, indicating ER localization of

the full-length immature gB. The C-terminal cleavage fragments were Endo H resistant, with

molecular weights of 62 kDa and 50 kDa, whereas the fragment treated by PNGase F was less

than 50 kDa, indicating that proteolytic cleavage during maturation occurred after ER egress

(**Fig. 5B**). In particular, the detection of a single C-terminal gB fragment after PNGase F

treatment implies that the 66 kDa and 60 kDa fragments for untreated gB are differentially N-

linked glycoforms and not distinct cleavage products.

The alanine substitutions of ^{526}EHV^{528} and proline substitutions at E526 and H527 did not

alter effects of Endo H or PNGase F treatment compared to WT gB, indicating that protein

synthesis and post-translational modification were not significantly disrupted (**Fig. 5B**). In

contrast, only immature, ER associated gB was produced by the gB[V528P] mutant and gB was

substantially reduced in quantity compared to WT gB (**Fig. 5B**). Absence of the C-terminal

cleaved glycoforms showed gB[V528P] was not transported out of the ER for further processing

and maturation. These data were consistent with its absence on EVs (**Fig. S2**) and significantly

reduced abundance on cell surfaces (**Fig. S4**).

Disruption of the canonical biosynthesis of gB, required for gB incorporation into the

envelope of VZV particles, explained why gB[V528P] did not induce cell fusion (**Fig. 3D;**

Table. 1) and pOka-TKGFP-gB[V528P] failed to replicate (**Fig. 3E; Table. 1**). Since

gB[V528A] did not differ from WT gB, the consequences are not due to an amino acid side

chain change but can be attributed to distortion of the gB DIII central helix structure by

gB[V528P] as predicted by AlphaFold2 (**Fig. 3A**). The amino acids equivalent to V528 in VZV

gB were relatively conserved among all the human herpesviruses (**Fig. S6A**) and AlphaFold2

predicted a similar distortion in the gB DIII central helix by proline substitutions at this position

(**Fig. S6B; Table. S3**) except for HHV8 gB, which required an additional G471Y mutation to

allow the bend caused by I472P (**Fig. S6B; Table. S3)** and all betaherpesviruses have a tyrosine

residue equivalent to G471 (**Fig. S6B**). The predicted deformation by proline substitution to

valine or isoleucine is a conserved feature, supporting the important role of the equivalent V528

position in maintaining the helical structure of herpesvirus gB DIII.

Discussion

Like other viral fusogens, a conformational change of herpesvirus gB from a pre- to

postfusion structure is thought to be the driver for fusion of the virion envelope with the host cell

membrane for entry [1]. This is based on end-point structures for the energetically stable

postfusion gB and for gB stabilized by either mutagenesis of DIII or using a fusion inhibitor and

chemical cross-linking [2,21-27,31,36]. Identifying transitional states of the trimer is essential for

understanding the molecular dynamics of gB as a fusogen. The present study visualizes, for the

first time, two such presumed transitional states using cryo-ET STA and focused classification of

the VZV gB mutant, H527P. The compact and more expanded conformations of VZV

gB[H527P] and the poor rigid body fitting of a prefusion homology model based on HSV-1

gB[H516P] into the two cryo-EM STA maps provide insights into the first steps in this dynamic

process. A critical observation about the related HSV-1 gB[H516P] mutant was that the

ectodomain still transitions to a postfusion conformation as shown by X-ray crystallography,

supporting the concept that prefusion gB stability requires additionally the transmembrane and

C-terminal domains along with the lipid membrane [21,27,30,31]. In contrast to WT HSV-1 gB, which

shows a partial transition, WT VZV gB on EVs fully transitions to the postfusion conformation

suggesting that VZV gB is a highly metastable fusogen. In addition to the membrane, our

previous studies have demonstrated the likely requirement of the gH-gL heterodimer to keep

VZV gB locked in place [4-7], which is also supported by recent findings with pseudorabies virus

(PrV) and HSV-1 gB [8-10]. The overall requirement for the structural integrity of the gB DIII

central helix was demonstrated by consequences of the proline mutations at E526 and V528 for

gB-mediated fusion, gB maturation, viral replication and cell-cell spread.

The two VZV gB[H527P] classes imply that the previous structures for HSV-1

gB[H516P] or HCMV might not represent the complete array of prefusion gB conformations.

The architecture of the HSV-1 gB[H517P] structure [31] is similar to the more open conformation

of the class II structure for VZV gB[H527P] in the present study. In addition, the soluble,

truncated HSV-1 gB[H517P] ectodomain forms a postfusion structure, inferring that the full-

length proline mutant only becomes trapped during its transition in the context of the membrane

[31]. Furthermore, HCMV was captured using a thiourea fusion inhibitor, WAY-174865, and

chemical cross-linking. WAY-174865 binds to the gB fusion loops, membrane proximal region

(MPR), and transmembrane domain (TM) through interactions with hydrophobic and aromatic

residues, including the critical fusion loop residues Y153 and Y155 [36]. Importantly, the WAY-

174865 binding pockets in the HCMV gB trimer are located internally in between the α8 and α9

glycine-rich amphipathic helices, indicating that the gB trimer must be in an open configuration

for the compound to gain access to the MPR, TM, and fusion loops as seen for the 3.6Å

resolution cryo-EM structure [36]. While the activity data for WAY-174865 is not available,

previous studies on two closely related thiourea compounds, CF102 and WY1768, demonstrated

that these HCMV fusion inhibitors do not stably interact with cell-free virus particles and are

only effective once virus particles bind to the cell, preventing a complete conformational change

of gB triggered by receptor interaction [37,38]. CF102 inhibition escape mutations have also been

identified in the WAY-174865 binding pocket, including α9 amphipathic helix [36], providing

structural evidence for the shared mechanism of action of WAY-174865 and related thiourea

compounds. These studies indicate that thiourea fusion inhibitors target exposed regions in gB

transitional states rather than a more closed prefusion conformation similar to VZV gB[H527P]

class I in the present study. Together, this supports the notion that additional gB prefusion

conformations can populate herpesvirus envelopes.

The absence of gB[E526P] on EVs and the predominance of immature gB in transfected

cells (**Fig. S2**) was consistent with a moderate decrease in cell surface trafficking (**Fig. S4B**),

which might contribute to disrupted gB/gH-gL mediated cell-cell fusion activity. In addition,

although the effect of E526P on the central helix structure was minimal as predicted by

AlphaFold2, the mutation might induce an allosteric change in DIV since binding of neutralizing

antibody mAb 93k to its conformational DIV epitope was reduced (**Fig. 3B and 3C**). The barrier

to gB/gH-gL dependent fusion is higher under cell-cell fusion assay conditions compared to

VZV infected cells in culture, as observed with gB[E526P] inhibition of cell-cell fusion but

recovery of pOka-TKGFP-gB[E526P] virus and plaque formation, albeit limited. These findings

corroborated the importance of assessing the contributions of gB/gH-gL to fusion function by

testing mutations in the viral genome. Furthermore, although pOka-TKGFP-gB[E526P]

produced infectious virus, the marked reduction in plaque sizes was in the range associated with

failure of VZV to replicate in human skin xenografts in the SCID mouse model of VZV

pathogenesis, supporting the conclusion that even slight deviations in the gB central helix core

will be detrimental for herpesviruses as pathogens in their native hosts [4,6,14].

In line with the severe DIII central helix distortion predicted by AlphaFold2, the proline

substitution at V528 had the greatest affect as gB[V528P] did not exit the ER, severely limiting

intracellular trafficking necessary for gB maturation required to execute its fusion function.

However, while lethal for replication, defective biogenesis of gB[V528P] was compatible with

VZV particle assembly, showing that morphogenesis can occur independently of gB maturation

and trafficking, as reported for other herpesviruses [39,40]. The fate of gB[V528P] was likely

proteolysis as proteins misfolded in the ER eventually translocate to the proteasome for

degradation [41,14]. V528 is fully conserved in all alphaherpesvirus orthologues of gB except for

the distantly related TeHV3 (Testudinid alphaherpesvirus 3) which has an asparagine (**Fig. S1**).

The betaherpesviruses and gammaherpesviruses have a conserved change to isoleucine at the

residue equivalent to V528 (**Fig. S6**). A similar severe distortion in DIII was predicted for the

equivalent gB proline substitution in all human herpesvirus, implying a critical role in gB

biogenesis likely conserved for the *Herpesviridae*.

The induction of neutralizing antibodies that inhibit virus entry is a cornerstone of

vaccine design to protect against disease caused by enveloped viruses, first achieved with live

attenuated vaccines. Since neutralizing antibodies that target epitopes accessible on the prefusion

conformation of viral fusogens might be more potent, prefusion immunogens have become a

strategy for subunit vaccine design. Proline substitutions targeting the helical structure of the

protein (i.e., the heptad repeat in the coiled-coil region) have been used to stabilize several class I

fusion proteins for evaluation as candidate vaccines, including the HIV-1 envelope, the

respiratory syncytial virus fusion protein, and the spike proteins of the coronaviruses, Middle

East respiratory syndrome (MERS)-CoV, severe acute respiratory syndrome coronavirus (SARS-

CoV), and SARS-CoV-2 [42-46]. Our work indicates that proline stabilized VZV gB has the

potential to elicit neutralizing antibodies predominantly to prefusion gB and impair the transition

to the postfusion conformation effectively rendering VZV incapable of membrane fusion and cell

entry. Due to the highly conserved nature of gB, such an approach in vaccine design could be

applicable to all herpesviruses to reduce the burden of disease in animal or human populations.

Acknowledgements

We thank the National Institutes of Health (NIH) for their valuable support through

grants R01-AI102546 and R21-MH125285. We thank the Stanford Cell Sciences Imaging

Facility for supporting the confocal microscopy and the transmission electron microscopy (TEM)

work, the project described was supported, in part, by Award Number 1S10OD010580-01A1

from the National Center for Research Resources (NCRR) (confocal microscopy work), and by

NIH S10 Award Number 1S10OD028536-01, titled "OneView 4kX4k sCMOS camera for

transmission electron microscopy applications" from the Office of Research Infrastructure

Programs (ORIP) (TEM work). Its contents are solely the responsibility of the authors and do not

necessarily represent the official views of the NCRR or the NIH. Cell sorting was performed in

the Stanford Shared FACS Facility on an instrument obtained using NIH S10 Shared Instrument

Grant S10RR025518-01. Molecular graphics and analyses performed with UCSF ChimeraX,

developed by the Resource for Biocomputing, Visualization, and Informatics at the University of

California, San Francisco, with support from NIH R01-GM129325 and the Office of Cyber

Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.

Competing interests

The authors have declared that no competing interests exist.

Author contributions

M.Z. constructed mutant expressing plasmids, BAC recombinant viruses, carried out

sequence alignments, flow cytometry, stable reporter fusion assays, plaque assays,

immunoprecipitation, western blotting, confocal microscopy, and FC-TEM. B.V. generated EVs,

prepared Cryo-ET grids, carried out Cryo-ET data acquisition. M.C. performed subtomogram

averaging analysis on the Cryo-ET data. S.L.O. conducted the AlphaFold2 simulation and

performed model building, together with M.C. A.M.A., K.G., S.L.O., B.V., and M.Z.,

conceptualized the project. All the authors discussed the data and contributed intellectually to the

manuscript.

 Figure legend

Figure 1. The point mutation H527P constrains VZV gB in a prefusion conformation. A –

A diagram of the linear structure for VZV gB highlighting the location of the DIII central helix.

The signal sequence is depicted by the crossed white box. Colored regions (residues 115–736)

correspond to domains in the cryo-EM structure; DI (cyan), DII (green), DIII (yellow), DIV

(orange), DV (red) and linker regions (hot pink). Amino acids [526]EHV[528] (bold) were targeted for

mutagenesis. B and C – Trimer and monomer models for VZV gB prefusion (B) by SWISS

Model (previously reported in Oliver et al. 2021) and postfusion (C) from the 2.8A cryo-EM

structure [2]. The inset shows the [526]EHV[528] site in the DIII central helix. D – Single plane views of

representative tomographic reconstructions from cryo-ET of EVs produced by BHK-21 cells

transfected with WT gB or gB[H527P] expressing vectors. Upper and lower panels are different

planes from the same tomogram to show side and top views of WT gB or gB[H527P] expressed

on EVs. Boxes with dashed lines are magnified areas shown in the insets. Scale bars shown are

for each panel and inset.

Figure 2. Cryo-ET structures of VZV gB[H527P] resident on the membranes of EVs. A

and B – A representative slice view of a tomogram (A) and an EMAN2 cryo-EM map generated

by subtomogram averaging (STA) (B) from EVs purified from BHK-21 cells transiently

transfected with gB[H527P]. The white box in the upper panel of A highlights the area shown in

the lower panel. Arrow heads point to gB[H527P] on the membrane of EVs. C – A 10Å

resolution cryo-EM map derived from a local refinement of a tight mask and the associated

Fourier Shell Correlation (FSC) curve. The coloring represents resolution from 9.8Å (blue) to

12.4Å (red). D – STA maps and FSC curves for two classes (I – blue; II – red) of gB[H527P]

identified on the surface of EVs using EMAN2. E and F – STA maps of Class I (E) and Class II

(F) gB[H527P] molecules with the right-hand panels showing fitting of the homology model of

the VZV gB (previously published in Oliver et al. 2021) based on HSV-1 gB[H516P] [31] into the

cryo-EM maps. Snapshots from Movie S1 are shown, the VZV gB domains are colored as for

Fig. 1; DI (cyan), DII (green), DIII (yellow), DIV (orange), DV (red) and linker regions (hot

pink). Boxed areas of the STA maps (i and ii) highlight regions of structural differences between

the two maps. Arrows (blue to red gradient) represent the proposed movement between the class

I and class II maps.

Figure 3. Targeted mutagenesis of gB ^{526}EHV^{528} disrupts gB/gH-gL dependent membrane

fusion and viral replication. A- The predicted structures for the WT and mutant VZV gB DIII

central helices by AlphaFold2 [47,48] and the mutant structures aligned to WT gB using

Matchmaker [49] are shown; WT gB: yellow, alanine mutants: green, proline mutants: red. B-Flow

cytometry analysis of CHO-DSP1 cells transfected with vectors expressing WT gB or gB proline

and alanine mutants, permeabilized, and immunostained with Cy5-conjugated mAb 93k; non-

antigen control: grey, WT gB: blue, mutant gB: orange. C- Frequency of gB-expressing cells

detected by mAb 93k and median florescence intensity (MFI) of mAb 93k binding normalized to

WT gB (mean ± SEM). Statistical differences were evaluated by one-way ANOVA (**, $p <$

0.01; ***, $p < 0.001$; ****, $p < 0.0001$). D- Cell-cell fusion measured by the stable reporter

fusion assay (SRFA) using CHO-DSP1 cells transfected with vectors expressing WT gB or gB

mutants together with gH-gL expressing vectors, mixed with MeWo-DSP2 reporter cells. Fusion

efficiency was normalized to that mediated by gB[WT]/gH-gL (%). Data from at least three

independent experiments is shown in Box (25-75 percentile) and Whisker (10-90 percentile)

plots; dots are outliers, horizontal bar is the median and the + indicates the mean. Statistical

differences were analyzed by two-way ANOVA (*, $p < 0.1$; **, $p < 0.01$; ***, $p < 0.001$). E-

Plaque sizes in MeWo cells infected with pOka (WT), pOka-TKGFP-gB[E526P], pOka-

TKGFP-gB[E526A], pOka-TKGFP-gB[H527A], or pOka-TKGFP-gB[V528A] at 4 dpi; lethal

mutant viruses pOka-TKGFP-gB[H527P] and pOka-TKGFP-gB[V528P] are indicated by N/A

(not available). Comparison of thirty plaques per virus (mm^2) from three independent

experiments shown by Box and Whisker plots. Statistical differences between plaque sizes of

WT and mutant viruses were evaluated by one-way ANOVA (****, $p < 0.0001$). F-

Immunohistochemistry staining of VZV plaques in MeWo cells, scale bar = 1 mm.

Figure 4. Disrupted gB fusion function did not interfere with VZV immediate early, early

or late protein synthesis or with virus particle assembly. A- MeWo cells transfected with

pOka-TKGFP (WT), pOka-TKGFP-gB[E526P], pOka-TKGFP-gB[H527P], or pOka-TKGFP-

gB[V528P] BACs were imaged by confocal microscopy at 48 hrs post transfection. Immediate-

early protein (IE62, red) and late protein (capsid-ORF23, violet); early protein, thymidine kinase

(TK) expressed as a GFP fusion protein (green); nuclei stained with Hoechst 33342 (blue). Areas

highlighted by a small white box are magnified in the larger white boxes, white arrows indicate

cells at the early stage of infection identified by punctate expression of IE62 in nuclei with little

to no TK-GFP expression. Scale bar = 20 µm. B- Diagram of the flow cytometry-based

transmission electron microscopy (FC-TEM) assay. MeWo cells were transfected with VZV

BACs expressing TK-GFP, harvested at 48 hrs posttransfection, and sorted for GFP-positive

cells; cells were fixed, embedded in 10% gelatin, infiltrated with resin, negative staining and

mounted on the grid. C- TEM images of MeWo cells transfected with pOka-TKGFP (WT),

pOka-TKGFP-gB[E526P], pOka-TKGFP-gB[H527P], or pOka-TKGFP-gB[V528P] BACs.

Panels at the right are magnifications of areas indicated by boxes (N: nuclei; C: cytoplasm).

Boxes outlined in red show representative viral capsids, with solid red arrowheads indicating

viral DNA containing C-capsid and open red arrowheads indicating empty A-, or scaffold

containing B-capsids. Boxes outlined in blue show representative viral particles, with solid blue

arrowheads indicating complete virus particles and open blue arrowheads indicating light

particles. In the magnified images, scale bar = 0.1 µm.

Fig 5. Canonical biosynthesis of gB was disrupted by V528P. A- MeWo cells transiently

expressing WT gB and gB mutants were imaged by confocal microscopy at 24 hrs

posttransfection. Cells were stained with mouse anti-gB mAb SG2 (red), trans-Golgi network

marker, TGN46 (green), or early endosome antigen, EEA1 (green), and nuclei were stained with

Hoechst 33342 (blue). Representative images are from experiments performed at least in

duplicate. Areas in white boxes are shown at higher magnification in panels to the right; white

arrows indicate gB colocalization with TGN46 or EEA1 respectively. Scale bar = 15 µm. B-

Intracellular trafficking of gB was assessed by glycosylation status. MeWo cells transfected with

vectors expressing WT gB, or gB mutants were lysed at 24 hrs posttransfection and

immunoprecipitated with anti-gB mAb SG2, left untreated (-), or treated with Endo H (H) or

PNGase F (F). Proteins were separated by SDS-PAGE gel (7.5%) under reducing conditions, and

gB was detected with rabbit anti-gB 746-867. The arrows and vertical solid lines indicate the

glycosylated forms of full-length and cleaved gB respectively, the vertical dashed line indicates

the Endo H-resistant glycoforms of gB, and arrowheads indicate forms of gB deglycosylated by

both Endo H and PNGase F. Mass markers (kDa) are shown on the left.

Table 1. Properties of the mutant protein and recombinant viruses bearing the proline and alanine substitutions to ^{526}EHV^{528} in VZV gB DIII central helix.

Material and Methods

Cells. CHO-DSP1 cells [17], derived from Chinese Hamster Ovary (CHO) K1 cells (CCL-61, ATCC) that express Dual Split Protein (DSP)$_{1-7}$ of the chimeric reporter protein composed of split GFP and split *Renilla* luciferase, were propagated using F-12K nutrient mixture medium (Corning) supplemented with 10% fetal bovine serum (FBS, Gibco), penicillin (100 U/ml, Gibco), and streptomycin (100 µg/ml, Gibco). Human melanoma MeWo cells (HTB-65, ATCC) and MeWo-DSP2 derived from MeWo cells that express DSP$_{8-11}$ [17] were propagated in minimal essential medium (MEM, Corning) supplemented with 10% FBS, nonessential amino acids (1X, Corning), penicillin (100 U/ml, Gibco), streptomycin (100 µg/ml, Gibco) and amphotericin B (0.5 µg/ml, Corning). CHO-DSP1 and MeWo-DSP2 cells were kept under puromycin selection (8 µg/ml for CHO-DSP1 and 5 µg/ml for MeWo-DSP2; Gibco). BHK-21 cells were grown in GMEM supplemented with 20 mM Hepes pH 7.4, 2% (v/v) TPB and 2% (v/v) FBS.

Sequence alignment. The amino acid sequence of alphaherpesvirus gB orthologues and human herpesvirus gB orthologues were obtained from the NCBI (https://www.ncbi.nlm.nih.gov/protein). The amino acids that aligned with VZV gB DIII central helix were defined using Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo).

Construction of gB expressing vectors with mutations. gB mutant constructs were generated from the pCAGGS-gB[WT] vector, which was a gift from Tadahiro Suenaga and Hisashi Arase (Osaka University, Osaka, Japan) [16]. Primers pairs XmaI-Forward/gB-E526P-Reverse, XmaI-Forward/gB-H527P-Reverse, XmaI-Forward/gB-V528P-Reverse, XmaI-

Forward/gB-E526A-Reverse, XmaI-Forward/gB-H527AP-Reverse, XmaI-Forward/gB-V528A-

Reverse, gB-EHV-Forward/AgeI-Reverse containing the desired mutation were used to amplify

two PCR products using AccuPrime Taq (Invitrogen). The amplicons were digested with either

XmaI or AgeI restriction enzymes (New England Biolabs), and ligation was performed with the

digested sticky end and the 5' phosphate blunt end. Ligated products were electroporated into

TOP10F' Electrocomp E. Coli (Invitrogen) and plated under ampicillin selection. Clones were

confirmed by AgeI/XmaI digestion and sequencing using primers gB-pCAGGS-Seq-F and gB-

pCAGGS-Seq-R. All the primers used were list in **Table S4**.

Generation of pOka-DX Bacterial Artificial Chromosomes (BACs) with mutations

in gB (ORF31). The pOka-TK-GFP virus was engineered from the parental Oka strain of VZV

(pOka) derived from a self-excisable bacterial artificial chromosome (BAC) [50] to express EGFP

conjugated to the thymidine kinase (TK) [6]. A pPOKA-ΔORF31-TKGFP BAC lacking ORF31

was constructed previously [4]. Briefly, the shuttle vector, gB-Kan containing ORF31[4] was

digested by BgIII and NdeI enzymes (New England Biolabs). pCAGGS-gB plasmids containing

the desired mutation were also digested with BgIII and NdeI enzymes, and the fragment of

interest was inserted into the shuttle vector gB-Kan by ligation, and were electroporated into

TOP10F' Electrocomp E. Coli (Invitrogen) and plated under kanamycin selection. Clones were

confirmed by AgeI/XmaI digestion and sequencing using gB-Kan-Seq-F and gB-Kan-Seq-R.

Then the gB-Kan vector carrying the mutations were used to generate BACs using GS1783 cells

containing the pPOKA-ΔORF31-TKGFP BAC by Red recombination [11,50,51]. Fragment insertion

was confirmed by PCR using primers [31]F56625-56645 and [31]R59697-59717. All BACs

were purified using large-construct purification kit (Qiagen), and digested by HindIII to verify

that spurious recombination had not occurred. To generate BAC-derived viruses, MeWo cells

were transfected in at least triplicate with two BAC clones for each mutation. Viruses were

typically recovered at 5-10 days posttransfection and passed on MeWo cells to propagate and

titrated on MeWo cells. All the primers used were list in **Table S4**.

EVs Production. EVs were prepared as described [32]. In brief, BHK-21 cells, at around

70% confluency, were transiently transfected with pCAGGS vectors expressing proline

substitutions to [526]EHV[528], or His-tagged WT gB or gB[H527P]. Cells were grown for an

additional 48 hours with a media exchange to serum free GMEM after 24 hours. Vesicles were

harvested from the supernatant by centrifugation through a 20% sucrose cushion and

resuspended in 20 mM Hepes pH 8, 130 mM NaCl.

Cryo-ET data collection using the Titan Krios. For grid preparation, 3.5 µL vesicles

were mixed with 5 nm gold fiducials on Quantifoil R2/1 grids and plunge frozen in a

propane/ethane mixture using a manual plunge freezer. Microscopy of WT gB, gB[H527P], and

gB[H527P]-His EVs was performed at 300 keV on a Titan Krios electron microscope (Thermo

Fisher scientific) equipped with a Gatan K3 (5 μm/pixel) camera and Bioquantum post-column

energy filter controlled using SerialEM [52] for micrograph and automated batch tomography data

collection. Movie data (11,283 total stacks) were captured in counted mode with a dose rate of

~3.22-3.44 $e^-/Å^2$/s per frame and 200-300 millisecond exposure time per frame and 0.8-1.8

second total exposure time at a nominal magnification of 81,000x and a pixel size of 1.095

Å/pixel on the specimen. The defocus was set to -2.0µm.

Reconstruction of the cryo-ET maps for gB[H527P]. The motion correction, damage

compensation, and tilt stacks for all movie-mode data were generated using WARP [53]. All data

processing described was performed using the tomography workflow in EMAN 2.99 [54,55]. Forty

tilt series were automatically reconstructed, and 2,074 particles were manually selected from the

tomograms. The CTF were estimated from the tilt images before unbinned, phase flipped

subtomograms were extracted. The initial orientation of each particle was assigned using a vector

pointing from the center of the vesicle that the particle was located on, to the coordinates of that

particle. The initial model was directly generated from the particles at 30Å resolution and used as

a reference "gold-standard" subtomogram to perform subtilt refinement on all particles. A soft

cylinder mask that includes the protein and membrane density was used during the refinement,

and C3 symmetry was applied to the averaged structure. After each iteration of the refinement,

local resolution was measured based on the two half-maps, and the averaged structures were

filtered according to the local resolution. Then, to resolve the structural heterogeneity at the core

of the protein revealed by the local resolution, focused classification was performed based on the

orientation determined from the single model refinement, using a mask that covers the central

part of the protein. Finally, separate subtomogram and subtilt refinements were performed for the

particles of each class from the refinement, resulting in the final averaged structures. All images

and movies were generated using UCSF ChimeraX 1.14 [49].

AlphaFold2 predictions of herpesvirus gB DIII central helix. The gB DIII central

helix for eight human herpesvirus gB orthologs was defined by amino acid alignment of the

complete gB proteins using Clustal X and comparison to the native VZV gB structure (PDB

6VN1) DIII central helix. AlphFold2 [47,48] was compiled following the installation instructions

(https://github.com/deepmind/alphafold) on a Colfax ProEdge DX935 Workstation with an Intel

Xeon W-2195 18C/36T 2.3Ghz CPU, 4 PNY GeForce RTX2080Ti 11GB GPUs, 256MB RAM

and Centos 8 operating system.

Stable-reporter fusion assays. CHO-DSP1 were transfected with vectors pCAGGS-

gB[WT], pCAGGS-gB[E526P], pCAGGS-gB[H527P], pCAGGS-gB[V528P], pCAGGS-

gB[E526A], pCAGGS-gB[H527A], or pCAGGS-gB[V528A]; pCAGGS-gB[WT]-His-tagged, or

pCAGGS-gB[H527P]-His-tagged, with or without vectors expressing gH-gL, pME18S-gH[TL]

and pcDNA3.1-gL using Lipofectamine 2000. At 6 hrs post-transfection, the transfected cells

were harvested and mixed with MeWo-DSP2 cells for additional 48 hrs. The activity of *Renilla*

luciferase was read immediately after adding substrate h-Coelenterazine (Nanolight Technology)

on a Synergy H1 Hybrid Multi-Mode Reader (BioTek). Experiments were performed in triplicate

at least. The fusion activity for His-tagged WT gB and gB[H527P] was shown in **Fig. S7**.

581   **Quantitation of cell surface and total expression of gB.** The quantity of cell surface

VZV glycoproteins were measured as described previously [4]. Briefly, CHO-DSP1 cells were

transfected with vectors pCAGGS-gB[WT], or pCAGGS-gB[E526P], pCAGGS-gB[H527P],

pCAGGS-gB[V528P], pCAGGS-gB[E526A], pCAGGS-gB[H527A], and pCAGGS-

gB[V528A], or vectors expressing gH-gL, pME18S-gH[TL] and pcDNA3.1-gL. At 24 hrs later,

cells were dislodged using an enzyme-free cell dissociation buffer (Gibco), fixed with 1%

paraformaldehyde (PFA, Boston Bioproducts), and resuspended in FACS staining buffer [PBS

with 0.2% IgG-free bovine serum albumin (Jackson ImmunoResearch), and 0.1% NaN_3 (Sigma

Aldrich)]. Cell surface gB were detected with primary antibodies mouse anti-VZV gB mAb

(93k; human)[2] conjugated with Cy5, at 1:100 dilution. Total amounts of viral glycoprotein

production were performed by the same staining protocol, except cells were permeabilized using

Cytofix/Cytoperm kit (BD Biosciences) before adding the antibodies and during the following

procedure. Stained cells were analyzed using a DXP multi-color FACScan analyzer (Cytek

Biosciences), and data were processed with FlowJo (TreeStar) to determine the quantity of total

and surface levels of gB respectively. The ratio of cell surface to total quantity was also

calculated. Experiments were performed in triplicate.

Immunoprecipitation and Western blotting. Whole cell lysate from MeWo cells

transiently expressing WT gB or gB with proline and alanine substitutions, or gH-gL, a non-

antigen negative control was extracted at 24 hrs posttransfection using extraction buffer [0.1 M

NaCl, 5 mM KCl (Fisher Scientific), 1 mM $CaCl_2$ (Fisher Scientific), 0.5 mM $MgCl_2$ (Fisher

Scientific), 1% IGEPAL CA-630 (Sigma Aldrich), 1% Deoxycholate (Sigma Aldrich) in 0.1 M

Tris buffer, pH 7.2], with cOmplete EDTA-free protease inhibitor, for 30 min on ice. The sample

was centrifuged at 3,000 RCF for 10 min at 4°C to get rid of cell debris. The supernatant was

immunoprecipitated (IP) with mouse anti-VZV gB mAb (SG2-2E6, Meridian Life Sciences) that

was crosslinked to protein A beads (Pierce) at 20 µg antibody/30 µl beads slurry. Samples were

then either left untreated or treated with Endo H (New England Biolabs) or PNGase F (New

England Biolabs) according to the manufacture's protocol. Proteins were denatured under

reducing condition and separated by 7.5% SDS PAGE gel (BioRad), transferred to Immobilon-P

PVDF membranes (Millipore) using Trans-blot SD Semi-dry Transfer cell (BioRad) in transfer

buffer [48 mM Tris Base (Fisher Scientific), 39 mM Glycine (Fisher Scientific) and 0.01% SDS

(Sigma Aldrich), with 20% methanol (Sigma Aldrich)]. The transferred proteins were probed for

gB using rabbit polyclonal antiserum, 746-868, that recognizes the epitope

^{833}PEGMDPFAEKPNAT^{846} in the cytoplasmic domain of gB (GenScript Corp) [14], followed by

anti-rabbit IgG horseradish peroxidase-linked secondary antibodies (GE Healthcare Life

Sciences, UK Ltd., 1:10,000) and Pierce ECL Plus Western Blotting Substrate (ThermoFisher).

Chemiluminescence was detected using BioMax MR Film (Carestream). For EVs analysis, after

supernatants were removed for vesicle preparations, cells were washed with cold PBS and

detached using cell scrapers. Cells were pelleted by centrifugation (5 minutes, 4500 RCF, 4°C),

transferred into 1.5 ml microfuge tubes and washed in cold PBS again before resuspension in

RIPA buffer, 100 µL per T175 flask (50 mM Tris pH 8, 1% NP-40, 0.1% SDS, 150 mM NaCl,

0.5% Sodium Deoxycholate, 5 mM EDTA, 1 mM PMSF). Samples were shaken at 4°C for 30

min before centrifugation at 500 RCF for 10 minutes. Supernatants were mixed in SDS sample

buffer and run in parallel with vesicle samples in SDS PAGE, followed by Coomassie staining or

western blotting with the rabbit anti-VZV gB antibody 746-868[14] followed by anti-rabbit-HRP

(Sigma-Aldrich Chemie GmbH). For loading control, western blots were re-probed using a

mouse anti-GAPDH antibody (Sigma-Aldrich Chemie GmbH) followed by anti-mouse-HRP

(Sigma-Aldrich Chemie GmbH) The Coomassie staining and Western blotting for His-tagged

WT gB and gB[H527P] was shown in **Fig. S7**.

Confocal microscopy. MeWo cells transfected with gB expressing vectors were fixed 24

hrs posttransfection with 4% formaldehyde for 10 min. MAb SG2 (Meridian Life Sciences) was

used to stain for VZV gB, and primary antibodies to cellular proteins were used as markers for

early endosomes (EEA-1; rabbit polyclonal antibody; Santa Cruz Biotechnology Inc), and the

trans-Golgi (TGN46; sheep polyclonal antibody; BioRad). Primary antibodies were detected

with secondary antibodies, anti-mouse Alexa Fluor 555 (Invitrogen), anti-rabbit Alexa 488, and

anti-sheep Alexa Fluor 647 (Invitrogen). Nuclei were stained with Hoechst 33342 (Invitrogen).

MeWo cells transfected with pOka, pOka-TKGFP-gB[E526P], pOka-TKGFP-gB[H527P], or

pOka-TKGFP-gB[V528P] BACs were fixed at 48 hrs or 72 hrs post transfection with 4%

formaldehyde for 10 min, and cells were probed using primary antibodies to VZV proteins IE62

(mouse; Chemicon international), capsid-ORF23 (polyclonal rabbit) [56] and gB (93k; human)[2] and

nuclei (Hoechst 33342). Primary antibodies were detected with secondary antibodies, anti-mouse

Alexa Fluor 555 (Invitrogen), and anti-rabbit or anti-human Alexa Fluor 647 (Invitrogen). Nuclei

were stained with Hoechst 33342 (Invitrogen). Images were captured with a Leica SP8,

Whitelight Laser Confocal Microscope. Channel merging and image processing was performed

with ImageJ and Photoshop.

Plaque size assay. Monolayers of MeWo cells (1×10^5 cell/cm^2) were inoculated with

50 plaque forming unit (PFU) of cell-associated pOka-TKGFP viruses or pOka-TKGFP-

gB[E526P] for 2 hrs, followed by change of medium. Cells were fixed with 4% PFA at 4 days

post-infection and processed with Immunohistochemistry (IHC) using mouse monoclonal anti-

VZV antibody (Mixed, GeneTex #GTX38720, 1:2000 diluted in PBS), followed by incubation

with biotinylated anti-mouse IgG (Vector Laboratories #BA-9200), and alkaline phosphatase

streptavidin (Jackson ImmunoResearch #016-050-084). The plaques were visualized by staining

with a mixture of FastRed salt (Sigma Aldrich) and Naphthol AS-MX phosphate (Sigma

Aldrich). Images of plaques were taken with Axio microscope (Zeiss) using 2.5X lens. The

plaques were outlined and the area of plaque was calculated using Image J. Experiments were

performed at least in triplicate.

FC-TEM of BAC transfected cells. Monolayers of MeWo cells (1.6×10^5 cell/cm^2)

were seeded onto P100 dish and were transfected with 29 µg pOka-TKGFP, pOka-TKGFP-

gB[E526P], pOka-TKGFP-gB[H527P], or pOka-TKGFP-gB[V528P] BACs by Lipofectamine

2000. Cells were harvested at 48hrs posttransfection, and resuspended in FACS staining buffer

[2.5% FBS, 2 mM EDTA in PBS] at 5×10^6 cell/ml. The GFP positive cells were sorted for

enrichment by negative gating at 0.5% of the mock sample that was MeWo cells transfected

without BAC. The sorted cells were pelleted and re-suspended in 10% Gelatin in 0.1M Sodium

Cacodylate buffer pH 7.4 at 37°C and allowed to equilibrate 5 min. Cells were pelleted again,

excess gelatin removed, then chilled in cold blocks and covered with cold 1% Osmium tetroxide

(EMS Cat# 19100) for 2 hrs rotating in a cold room, washed 3 times with cold ultrafiltered

water, then en bloc stained overnight in 1% Uranyl Acetate at 4°C while rotating. Samples were

then dehydrated in a series of ethanol washes for 20 minutes each at 4°C beginning at 30%, 50%,

70%, 95% where the samples were then allowed to rise to RT, changed to 100% ethanol, then

Propylene Oxide (PO) for 15 min. Samples were infiltrated with EMbed-812 resin (EMS

Cat#14120) mixed 1:2, 1:1, and 2:1 with PO, and then were placed into neat EMbed-812 for 2 to

4 hrs followed by placement into molds at 65° C oven overnight. Sections were taken

approximately 80nm, picked up on formvar/Carbon coated 100 mesh Cu grids, stained for 30 sec

in 3.5% Uranyl Acetate in 50% Acetone followed by staining in 0.2% Lead Citrate for 3 min.

Observed in the JEOL JEM-1400 120kV and photos were taken using a Gatan Orius 4k X 4k

digital camera. The yield efficiency of FC-TEM was summarized in **Table S5.**

Supplemental information

Fig S1. Sequence alignment of the amino acids equivalent to VZV gB DIII central helix

from the five genera of the subfamily alphaherpesvirinae. Virus host species and the

glycoprotein gB NCBI accession are listed and the residues corresponding to [526]EHV[528] are

colored in blue. Conservation among residues is presented based on Clustal Omega: "." indicates

weak similarity; ":" indicates strong similarity; "*" indicates full conservation. Histidine

corresponding to H527 in VZV is highlighted in pink. VZV: Varicella-zoster virus; CeHV-9:

Cercopithecine alphaherpesvirus 9; PRV: pseudorabies virus; EHV-1: Equine alphaherpesvirus

1; EHV-3: Equine alphaherpesvirus 3; EHV-4: Equine alphaherpesvirus 4; EHV-8: Equine

alphaherpesvirus 8; EHV-9: Equine alphaherpesvirus 9; FeHV-1: Felid alphaherpesvirus 1;

CaHV-1: Canid alphaherpesvirus 1; BoHV-1: Bovine alphaherpesvirus 1; BoHV-5: Bovine

alphaherpesvirus 5; BuHV-1: Bubaline alphaherpesvirus 1; CpHV-1: Caprine alphaherpesvirus

1; CvHV-1: Cervid alphaherpesvirus 1; CvHV-2: Cervid alphaherpesvirus 2; CvHV-3: Cervid

alphaherpesvirus 3; MoHV-1: Monodontid alphaherpesvirus 1; PhHV-1: Phocid

alphaherpesvirus 1; HSV-1: Herpes simplex virus 1; HSV-2: Herpes simplex virus 2; CeHV-2:

Cercopithecine alphaherpesvirus 2; AtHV-1: Ateline alphaherpesvirus 1; McHV-1: Macacine

alphaherpesvirus 1; PnHV-3: Panine alphaherpesvirus 3; SaHV-1: Saimiriine alphaherpesvirus 1;

PaHV-2: Papiine alphaherpesvirus 2; BoHV-2: Bovine alphaherpesvirus 2; LeHV-4: Leporid

alphaherpesvirus 4; MaHV-1: Macropodid alphaherpesvirus 1; MaHV-2: Macropodid

alphaherpesvirus 2; PtHV-1: Pteropodid alphaherpesvirus 1; AnHV-1: Anatid alphaherpesvirus

1; CoHV-1: Columbid alphaherpesvirus 1; MeHV-1: Meleagrid alphaherpesvirus 1; GaHV-2:

Gallid alphaherpesvirus 2; GaHV-3: Gallid alphaherpesvirus 3; SpAHV-1: Spheniscid

alphaherpevirus 1; GaHV-1: Gallid alphaherpesvirus 1; PsHV-1: Psittacid alphaherpesvirus 1;

ChHV-5: Chelonid alphaherpesvirus 5; TeHV-3: Testudinid alphaherpesvirus 3.

Fig S2. Expression of gB proline substitutions to ^{526}EHV^{528} residues in DIII on EVs. A-

Coomassie blue stain of EVs purified from BHK-21 cells transfected with vectors expressing

VZV WT gB, gB ^{526}EHV^{528} proline mutants, HSV-1 gB or empty vector. Green arrow heads

highlight the uncleaved form of HSV-1 gB and the cleaved form of VZV gB. B- Western blot of

VZV gB from purified EVs detected with rabbit anti-gB 746-867 and whole cell lysates from

transiently transfected BHK-21 cells; GAPDH loading control. The combined band density of

the immature full-length and the cleaved fragment of gB in cell lysates was normalized to that of

GAPDH and shown as a percentage of WT gB.

Figure S3. Classification scheme of gB[H527P] particles in EVs membranes identified in cryo-ET micrographs.

Fig S4. The analysis of gB expression by flow cytometry. A- Surface or total expression of gB was analyzed by flow cytometry of CHO-DSP1 cells transfected with vectors expressing WT gB or gB mutants, nonpermeabilized for surface gB detection or permeabilized for total gB detection, and immunostained with Cy5-conjugated anti-gB mAb 93k. The population of WT gB-expressing cells (blue) and mutant gB-expressing cells (orange) was gated against cells transfected with gH-gL (grey). The frequency of gB-expressing cells and median of fluorescence intensity (MFI) were compared to that in the WT gB and presented as percentage of WT in the table (%WT; mean $\pm$ SD). B- The ratio of surface to total gB expression from A normalized to WT gB (% WT) (mean $\pm$ SEM) was calculated. Statistical differences from at least three independent experiments were evaluated by two-way ANOVA (*, $p < 0.1$; ****, $p < 0.0001$).

Fig S5. Proline substitutions to [526]EHV[528] residues in DIII does not interfere with gB expression and virus assembly in BAC transfected cells. A- Confocal microscopy images of MeWo cells transfected with pOka-TKGFP (WT), pOka-TKGFP-gB[E526P] (E526P), pOka-TKGFP-gB[H527P] (H527P), or pOka-TKGFP-gB[V528P] (V528P) BACs at 72 hrs post transfection. All the BACs expressed thymidine kinase (TK) fused with GFP fluorescence protein at the C-terminus. Cells were stained for immediate-early protein (IE62; red), gB (violet), and nuclei (Hoechst 33342; blue). Scale bar = 20 µm. B- Transmission Electron Microscopy (TEM) images of MeWo cells transfected with pOka-TKGFP-gB[H527P] BAC at 48 hrs post transfection. The transfected cells were harvested at 48 hrs post transfection and enriched for

GFP-expressing cells by sorting prior to negative stain TEM. Panels at the right are

magnifications of areas indicated by boxes in the larger images (N: nuclei; C: cytoplasm). Boxes

outlined in red show representative viral capsids, with solid red arrowheads indicating viral DNA

containing C-capsid and open red arrowheads indicating empty A-, or scaffold containing B-

capsids. Boxes outlined in blue show representative viral particles, with solid blue arrowhead

indicating complete virus particle in the process of assembly in the cytoplasm and open blue

arrowheads indicating light particles on the cell surface or in the process of assembly in the

cytoplasm. In the magnified images, scale bar = 0.1 µm.

Figure S6. Predicted structural deformations induced by proline substitutions to residues

corresponding to VZV ^{526}EHV^{528} residues in gB DIII central helix of the eight human

herpesviruses. A- Sequence alignment of the amino acids equivalent to VZV gB DIII central

helix from human herpesviruses. gB NCBI accession are listed. Residues corresponding to VZV

^{526}EHV^{528} are colored in blue. Conservation among residues is presented based on Clustal

Omega: "." indicates weak similarity; ":" indicates strong similarity; "*" indicates full

conservation. B- All structures for the wild type (WT) and mutant the gB DIII central helices

were predicted using AlphaFold2 [47,48] and the mutant structures aligned to WT gB using

Matchmaker [49]. WT gB is colored yellow and each of the proline substitutions is colored as for

VZV gB E526 (orange), H527 (orange red), V528 (maroon). For HHV8, a fourth proline

mutant, gB[G471Y/I472P] (purple) was modeled to assess the effect of tyrosine at the -1 position

for I472, which had an RMSD of 7.109. The amino acids at the beginning and end of the gB DIII

central helix are indicated. HHV1: HSV-1; HHV2: HSV-2; HHV3: VZV; HHV4: EBV; HHV5:

HCMV; HHV8: KSHV.

Figure S7. Characterization of EVs expressing His-tagged gB[H527P] and the fusion

function of His-tagged gB[H527P]. A- Coomassie Blue staining of an SDS-PAGE gel with

increasing quantities of EVs produced from BHK-21cells transfected with vectors expressing the

His-tagged gB[H527P] (H527P-His). N-terminal and C-terminal cleavage products of VZV

gB[H527P] are indicated by blue and red arrowheads respectively. B- Western blot of EVs and

cell lysates for gB (anti-His tag) and GAPDH (anti-GAPDH); C-terminal cleavage product of

VZV gB[H527P] is indicated by red arrowhead. C- Box and whisker plots for cell-cell fusion

measured by the stable reporter fusion assay (SRFA) using CHO-DSP1 cells transfected with

vectors expressing untagged (WT) or His-tagged (WT-His) WT gB or the corresponding H527P

mutants with gH-gL, and mixed with MeWo-DSP2 cells for 48hrs. Fusion efficiency was

normalized to that mediated by gB[WT]/gH-gL (%). Box represents 25-75 percentile, whiskers

extend to 10-90 percentile, dots represent outliers, the median is the horizontal band and the

mean is the +. Data from three independent experiments is shown. The statistical difference

compared to WT gB was analyzed by two-way ANOVA (**, $p < 0.01$).

Table S1. Cryo-ET data collection parameters for WT gB and gB[H527P] EVs.

Table S2. Comparison of the structure deviation for proline and alanine mutants in the

VZV gB DIII central helix predicted with AlphaFold2.

Table S3. Comparison of the structure deviation for proline mutants in the gB DIII central

helix for all human herpesviruses predicted with AlphaFold2.

Table S4. Primer list

Table S5. Yield efficiency of the FC-TEM experiment.

Movie S1. Cryo-EM maps of the two conformations of gB[H527P]. This movie shows the

fitting of the homology modelled prefusion VZV gB into 9.0Å cryo-EM structure of HSV-1

gB[H517P] [31], followed by the class I and class II cryo-EM maps of VZV gB[H527P]. Boxed

areas of the VZV gB[H527P] STA maps (i and ii) highlight regions of structural differences

between the two maps. Arrows (blue to red gradient) represent the proposed movement between

the class I and class II maps.

References

1   Connolly, S. A., Jardetzky, T. S. & Longnecker, R. The structural basis of herpesvirus

entry. *Nat Rev Microbiol* **19**, 110-121, doi:10.1038/s41579-020-00448-w (2021).

2   Oliver, S. L. *et al.* A glycoprotein B-neutralizing antibody structure at 2.8 A uncovers a

critical domain for herpesvirus fusion initiation. *Nat Commun* **11**, 4141,

doi:10.1038/s41467-020-17911-0 (2020).

3   Arvin AM and Abendroth A, Varicella-zoster Virus, Chapter 13, in Howley Peter M and

David M Knipe, Editors, *Fields virology. Volume 2,* 7th Edition, Lippincott Williams &

Wilkins, 2021.

4   Oliver, S. L. *et al.* An immunoreceptor tyrosine-based inhibition motif in varicella-zoster

virus glycoprotein B regulates cell fusion and skin pathogenesis. *Proc Natl Acad Sci U S*

A **110**, 1911-1916, doi:10.1073/pnas.1216985110 (2013).

5   Heineman, T. C. & Hall, S. L. Role of the varicella-zoster virus gB cytoplasmic domain

in gB transport and viral egress. *J Virol* **76**, 591-599, doi:10.1128/jvi.76.2.591-599.2002

(2002).

6   Yang, E., Arvin, A. M. & Oliver, S. L. The cytoplasmic domain of varicella-zoster virus

glycoprotein H regulates syncytia formation and skin pathogenesis. *PLoS Pathog* **10**,

e1004173, doi:10.1371/journal.ppat.1004173 (2014).

7   Yang, E., Arvin, A. M. & Oliver, S. L. The Glycoprotein B Cytoplasmic Domain Lysine

Cluster Is Critical for Varicella-Zoster Virus Cell-Cell Fusion Regulation and Infection. *J*

Virol **91**, doi:10.1128/JVI.01707-16 (2017).

8   Vallbracht, M. *et al.* In Vitro Viral Evolution Identifies a Critical Residue in the

Alphaherpesvirus Fusion Glycoprotein B Ectodomain That Controls gH/gL-Independent

Entry. *mBio* **12**, doi:10.1128/mBio.00557-21 (2021).

9   Pataki, Z., Rebolledo Viveros, A. & Heldwein, E. E. Herpes Simplex Virus 1 Entry

Glycoproteins Form Complexes before and during Membrane Fusion. *mBio*, e0203922,

doi:10.1128/mbio.02039-22 (2022).

10   Pataki, Z., Sanders, E. K. & Heldwein, E. E. A surface pocket in the cytoplasmic domain

of the herpes simplex virus fusogen gB controls membrane fusion. *PLoS Pathog* **18**,

e1010435, doi:10.1371/journal.ppat.1010435 (2022).

11   Oliver, S. L., Yang, E. & Arvin, A. M. Varicella-Zoster Virus Glycoproteins: Entry,

Replication, and Pathogenesis. *Curr Clin Microbiol Rep* **3**, 204-215, doi:10.1007/s40588-

016-0044-4 (2016).

12   Harrison, S. C. Viral membrane fusion. *Virology* **479**, 498-507,

doi:10.1016/j.virol.2015.03.043 (2015).

13   Zerboni, L., Sen, N., Oliver, S. L. & Arvin, A. M. Molecular mechanisms of varicella

zoster virus pathogenesis. *Nat Rev Microbiol* **12**, 197-210, doi:10.1038/nrmicro3215

(2014).

14   Oliver, S. L. *et al.* Mutagenesis of varicella-zoster virus glycoprotein B: putative fusion

loop residues are essential for viral replication, and the furin cleavage motif contributes to

pathogenesis in skin tissue in vivo. *J Virol* **83**, 7495-7506, doi:10.1128/JVI.00400-09

(2009).

15   Vleck, S. E. *et al.* Structure-function analysis of varicella-zoster virus glycoprotein H

identifies domain-specific roles for fusion and skin tropism. *Proc Natl Acad Sci U S A*

108, 18412-18417, doi:10.1073/pnas.1111333108 (2011).

16   Suenaga, T. *et al.* Myelin-associated glycoprotein mediates membrane fusion and entry of

neurotropic herpesviruses. *Proc Natl Acad Sci U S A* **107**, 866-871,

doi:10.1073/pnas.0913351107 (2010).

17   Yang, E., Arvin, A. M. & Oliver, S. L. Role for the alphaV Integrin Subunit in Varicella-

Zoster Virus-Mediated Fusion and Infection. *J Virol* **90**, 7567-7578,

doi:10.1128/JVI.00792-16 (2016).

18   Reichelt, M., Zerboni, L. & Arvin, A. M. Mechanisms of varicella-zoster virus

neuropathogenesis in human dorsal root ganglia. *J Virol* **82**, 3971-3983,

doi:10.1128/JVI.02592-07 (2008).

19   Nagel, M. A. & Bubak, A. N. Varicella Zoster Virus Vasculopathy. *J Infect Dis* **218**,

S107-S112, doi:10.1093/infdis/jiy425 (2018).

20   Oliver, S. L. *et al.* The N-terminus of varicella-zoster virus glycoprotein B has a

functional role in fusion. *PLoS Pathog* **17**, e1008961, doi:10.1371/journal.ppat.1008961

(2021).

21   Heldwein, E. E. *et al.* Crystal structure of glycoprotein B from herpes simplex virus 1.

Science **313**, 217-220, doi:10.1126/science.1126548 (2006).

22   Backovic, M., Longnecker, R. & Jardetzky, T. S. Structure of a trimeric variant of the

Epstein-Barr virus glycoprotein B. *Proc Natl Acad Sci U S A* **106**, 2880-2885,

doi:10.1073/pnas.0810530106 (2009).

23   Chandramouli, S. *et al.* Structure of HCMV glycoprotein B in the postfusion

conformation bound to a neutralizing human antibody. *Nat Commun* **6**, 8176,

doi:10.1038/ncomms9176 (2015).

24   Burke, H. G. & Heldwein, E. E. Crystal Structure of the Human Cytomegalovirus

Glycoprotein B. *PLoS Pathog* **11**, e1005227, doi:10.1371/journal.ppat.1005227 (2015).

25   Vallbracht, M. *et al.* Structure-Function Dissection of Pseudorabies Virus Glycoprotein B

Fusion Loops. *J Virol* **92**, doi:10.1128/JVI.01203-17 (2018).

26   Li, X. *et al.* Two classes of protective antibodies against Pseudorabies virus variant

glycoprotein B: Implications for vaccine design. *PLoS Pathog* **13**, e1006777,

doi:10.1371/journal.ppat.1006777 (2017).

27   Cooper, R. S., Georgieva, E. R., Borbat, P. P., Freed, J. H. & Heldwein, E. E. Structural

basis for membrane anchoring and fusion regulation of the herpes simplex virus fusogen

gB. *Nat Struct Mol Biol* **25**, 416-424, doi:10.1038/s41594-018-0060-6 (2018).

28   Roche, S., Bressanelli, S., Rey, F. A. & Gaudin, Y. Crystal structure of the low-pH form

of the vesicular stomatitis virus glycoprotein G. *Science* **313**, 187-191,

doi:10.1126/science.1127683 (2006).

29   Kadlec, J., Loureiro, S., Abrescia, N. G., Stuart, D. I. & Jones, I. M. The postfusion

structure of baculovirus gp64 supports a unified view of viral fusion machines. *Nat Struct*

Mol Biol **15**, 1024-1030, doi:10.1038/nsmb.1484 (2008).

30   Vitu, E., Sharma, S., Stampfer, S. D. & Heldwein, E. E. Extensive mutagenesis of the

HSV-1 gB ectodomain reveals remarkable stability of its postfusion form. *J Mol Biol*

425, 2056-2071, doi:10.1016/j.jmb.2013.03.001 (2013).

31 Vollmer, B. *et al.* The prefusion structure of herpes simplex virus glycoprotein B. *Sci Adv*

6, doi:10.1126/sciadv.abc1726 (2020).

32 Zeev-Ben-Mordehai, T., Vasishtan, D., Siebert, C. A., Whittle, C. & Grunewald, K.

Extracellular vesicles: a platform for the structure determination of membrane proteins by

Cryo-EM. *Structure* **22**, 1687-1692, doi:10.1016/j.str.2014.09.005 (2014).

33 Zeev-Ben-Mordehai, T. *et al.* Two distinct trimeric conformations of natively membrane-

anchored full-length herpes simplex virus 1 glycoprotein B. *Proc Natl Acad Sci U S A*

113, 4176-4181, doi:10.1073/pnas.1523234113 (2016).

34 Fontana, J. *et al.* The Fusion Loops of the Initial Prefusion Conformation of Herpes

Simplex Virus 1 Fusion Protein Point Toward the Membrane. *mBio* **8**,

doi:10.1128/mBio.01268-17 (2017).

35 Si, Z. *et al.* Different functional states of fusion protein gB revealed on human

cytomegalovirus by cryo electron tomography with Volta phase plate. *PLoS Pathog* **14**,

e1007452, doi:10.1371/journal.ppat.1007452 (2018).

36 Liu, Y. *et al.* Prefusion structure of human cytomegalovirus glycoprotein B and structural

basis for membrane fusion. *Sci Adv* **7**, doi:10.1126/sciadv.abf3178 (2021).

37 Jones, T. R. *et al.* Specific inhibition of human cytomegalovirus glycoprotein B-mediated

fusion by a novel thiourea small molecule. *J Virol* **78**, 1289-1300,

doi:10.1128/jvi.78.3.1289-1300.2004 (2004).

38 Patrone, M., Secchi, M., Bonaparte, E., Milanesi, G. & Gallina, A. Cytomegalovirus

UL131-128 products promote gB conformational transition and gB-gH interaction during

entry into endothelial cells. *J Virol* **81**, 11479-11488, doi:10.1128/JVI.00788-07 (2007).

39   Farnsworth, A. *et al.* Herpes simplex virus glycoproteins gB and gH function in fusion

910   between the virion envelope and the outer nuclear membrane. *Proc Natl Acad Sci U S A*

911   **104**, 10187-10192, doi:10.1073/pnas.0703790104 (2007).

912   40   Klupp, B., Altenschmidt, J., Granzow, H., Fuchs, W. & Mettenleiter, T. C. Glycoproteins

913   required for entry are not necessary for egress of pseudorabies virus. *J Virol* **82**, 6299-

914   6309, doi:10.1128/JVI.00386-08 (2008).

915   41   Braakman, I. & Hebert, D. N. Protein folding in the endoplasmic reticulum. *Cold Spring*

916   *Harb Perspect Biol* **5**, a013201, doi:10.1101/cshperspect.a013201 (2013).

917   42   Sanders, R. W. *et al.* Stabilization of the soluble, cleaved, trimeric form of the envelope

918   glycoprotein complex of human immunodeficiency virus type 1. *J Virol* **76**, 8875-8889,

919   doi:10.1128/jvi.76.17.8875-8889.2002 (2002).

920   43   Krarup, A. *et al.* A highly stable prefusion RSV F vaccine derived from structural

921   analysis of the fusion mechanism. *Nat Commun* **6**, 8143, doi:10.1038/ncomms9143

922   (2015).

923   44   Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion

924   conformation. *Science* **367**, 1260-1263, doi:10.1126/science.abb2507 (2020).

925   45   Pallesen, J. *et al.* Immunogenicity and structures of a rationally designed prefusion

926   MERS-CoV spike antigen. *Proc Natl Acad Sci U S A* **114**, E7348-E7357,

927   doi:10.1073/pnas.1707304114 (2017).

928   46   Kirchdoerfer, R. N. *et al.* Stabilized coronavirus spikes are resistant to conformational

929   changes induced by receptor recognition or proteolysis. *Sci Rep* **8**, 15701,

930   doi:10.1038/s41598-018-34171-7 (2018).

47   Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding the

structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids*

Res **50**, D439-D444, doi:10.1093/nar/gkab1061 (2022).

48   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583-589, doi:10.1038/s41586-021-03819-2 (2021).

49   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators,

and developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).

50   Tischer, B. K. *et al.* A self-excisable infectious bacterial artificial chromosome clone of

varicella-zoster virus allows analysis of the essential tegument protein encoded by ORF9.

J Virol **81**, 13200-13208, doi:10.1128/JVI.01148-07 (2007).

51   Tischer, B. K., von Einem, J., Kaufer, B. & Osterrieder, N. Two-step red-mediated

recombination for versatile high-efficiency markerless DNA manipulation in Escherichia

coli. *Biotechniques* **40**, 191-197, doi:10.2144/000112096 (2006).

52   Mastronarde, D. N. Automated electron microscope tomography using robust prediction

of specimen movements. *J Struct Biol* **152**, 36-51, doi:10.1016/j.jsb.2005.07.007 (2005).

53   Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data preprocessing with

Warp. *Nat Methods* **16**, 1146-1152, doi:10.1038/s41592-019-0580-y (2019).

54   Tang, G. *et al.* EMAN2: an extensible image processing suite for electron microscopy. *J*

Struct Biol **157**, 38-46, doi:10.1016/j.jsb.2006.05.009 (2007).

55   Chen, M. *et al.* A complete data processing workflow for cryo-ET and subtomogram

averaging. *Nat Methods* **16**, 1161-1168, doi:10.1038/s41592-019-0591-8 (2019).

56   Chaudhuri, V., Sommer, M., Rajamani, J., Zerboni, L. & Arvin, A. M. Functions of

953   Varicella-zoster virus ORF23 capsid protein in viral replication and the pathogenesis of

954   skin infection. *J Virol* **82**, 10231-10246, doi:10.1128/JVI.01890-07 (2008).

Table 1. Properties of the mutant protein and recombinant viruses bearing the proline and alanine substitutions to ^{526}EHV^{528} in VZV gB DIII central helix.

	Protein				Recombinant virus		
	Fusion activity *	**Intracellular localization** **	**Protease cleavage** **	**EVs production**	**Viral gene expression**	**TEM morphology**	**Growth phenotype**
E526P	3.8%	ER, trans-Golgi, early endosome	Yes	No	IE, E, L	Capsids; virus particles	Severely impaired; Small plaque
H527P	4.1%	ER, trans-Golgi, early endosome	Yes	Yes	IE, E, L	Capsids; virus particles	Lethal
V528P	3.8%	ER only	No	No	IE, E, L	virus particles	Lethal
E526A	46.8%	ER, trans-Golgi, early endosome	Yes	ND	ND	ND	WT
H527A	80.2%	ER, trans-Golgi, early endosome	Yes	ND	ND	ND	WT
V528A	70.0%	ER, trans-Golgi, early endosome	Yes	ND	ND	ND	WT

*Fusion activity was meaused by SRFA and normalized to WT gB (%WT).

**WT gB localizes to ER, trans-Golgi network and early endosomes and undergoes cleavage.

IE: immediately Early gene; E: early gene; L: late gene.

ND: experiment was not performed.

Fig 1

Fig 2

Fig 4

A

gB/TGN46   gB/EEA1

WT

E526P

H527P

V528P

E526A

H527A

V528A

B

(-)   H   F

Fig 5

50