Caught in the Act: targeted mutagenesis of the herpesvirus fusogen central helix

captures transition states

Momei Zhou [1*], Benjamin Vollmer [2,3,4], Emily Machala [2,3,4], Muyuan Chen [5], Kay

Grünewald [2,3,4], Ann M. Arvin [1,6,7], Wah Chiu [5,6,8], and Stefan L. Oliver [1]

*Corresponding author: mzhou6@stanford.edu

[1] Department of Pediatrics, Stanford University School of Medicine, Stanford, California, USA.

[2] Centre for Structural Systems Biology (CSSB), Hamburg, Germany.

[3] Department of Chemistry, University of Hamburg, Hamburg, Germany.

[4] Leibniz Institute of Virology (LIV), Hamburg, Germany.

[5] Division of Cryo-EM and Bioimaging SSRL, SLAC National Accelerator Laboratory, Menlo

Park, CA 94025, USA.

[6] Department of Microbiology & Immunology, Stanford University School of Medicine,

Stanford, California, USA.

[7] Vir Biotechnology Inc, USA.

[8] Bioengineering, Stanford University School of Medicine, Stanford, CA 94305, USA.

Abstract

Herpesviruses remain a burden for animal and human health, including the medically important varicella-zoster virus (VZV). Membrane fusion mediated by conserved core glycoproteins, the fusogen gB and the heterodimer gH-gL, enables herpesvirus cell entry. The ectodomain of gB orthologs has five domains and is proposed to transition from a prefusion to postfusion conformation but the functional relevance of the domains for this transition remains poorly defined. Structure-function studies of the VZV gB DIII central helix were performed targeting residues [526]EHV[528]. Critically, a H527P mutation captured gB in a prefusion conformation as determined by cryo-EM, a loss of membrane fusion in a virus free assay, and failure of recombinant VZV to spread in cell monolayers. Importantly, two predominant cryo-EM structures of gB[H527P] were identified by 3D classification and focused refinement, suggesting they represented gB conformations in transition. These studies reveal gB DIII as a critical element for herpesvirus gB fusion function.

Introduction

The *Herpesviridae* encompass pathogens that cause a wide range of medically and economically important diseases of humans and animals [1]. All herpesviruses are enveloped and use glycoproteins to initiate and drive membrane fusion between the virion and the host cell during entry [2]. The three glycoproteins, gB, gH, and gL, required for this process are conserved across the *Herpesviridae* [2,3]. Structure-function studies have revealed that the gB trimer is the primary fusogen and the gH-gL heterodimer has roles in cell tropism and gB regulation [1,4-10]. Incorporation of gB into the envelope of herpesvirus particles is complex. gB is transported to the plasma membrane of the infected cell, endocytosed, and trafficked to the trans-Golgi network for insertion into the herpesvirus envelope during morphogenesis [11]. To achieve fusion with cellular membranes, gB on the virion envelope is thought to undergo a conformational transition from a meta-stable prefusion structure to an energetically more stable postfusion state [2,12].

VZV, a member of the *Alphaherpesvirinae* and human host restricted pathogen, causes varicella (chickenpox), establishes latency in sensory ganglion neurons, and can reactivate to manifest as zoster (shingles) [13]. In addition to membrane fusion required for virion entry, gB dependent cell-cell fusion is a hallmark of VZV pathogenesis [1]. Characteristic polykaryocytes form within infected tissues *in vivo* and are modeled *in vitro* by syncytia formation between MeWo cells [1]. VZV gB, together with gH-gL, are necessary and sufficient to induce cell-cell fusion *in vitro* [6,14-17]. Unlike its HSV gB orthologues, VZV gB is modified by furin cleavage post-translationally, which is needed for optimal skin lesion formation *in vivo* [14]. Critically, VZV induced fusion between differentiated cells has been linked to adverse health effects, including fusion of ganglion neurons and satellite cells associated with postherpetic neuralgia (PHN), and fusion of endothelial cells in the brain vasculature linked to strokes [18,19].

Despite substantial amino acid sequence differences, the gB orthologues from alpha-, beta-

and gammaherpesvirus subfamilies adopt remarkably similar structures in their postfusion

ectodomain conformations, consisting of homotrimers with five domains, DI-DV [3,20-27]. This

architecture led to the classification of gB orthologues as type III fusogens based on similarities

to postfusion forms of vesicular stomatitis virus glycoprotein G (VSV-G) and the baculovirus

gp64 [28,29]. In the gB trimer, DI contains the fusion loops, DII has a pleckstrin homology domain,

DIII forms a central helical bundle, DIV forms the crown, and DV connects the crown to the

transmembrane domain. Our recent single particle cryogenic electron microscopy (cryo-EM)

studies of full-length VZV gB have demonstrated the importance of gB DIV in fusion initiation

[3,20]. In contrast, the rapid conversion of gB to its postfusion form upon deletion of the

transmembrane domain or removal from a lipid bilayer has hindered deriving high-resolution

structures of prefusion gB [21,27,30,31]. Extracellular vesicles (EVs) were used as a surrogate for

structural analyses of virion envelope-associated glycoproteins, overcoming effects on the native

topology of the protein after detergent solubilization and reconstitution into an artificial

membrane [32]. Cryogenic electron tomography (cryo-ET) and single particle cryo-EM have

revealed prefusion structures for herpes simplex virus 1 (HSV-1) and human cytomegalovirus

(HCMV) gB anchored in native membranes [31,33-36]. Expression on EVs of a HSV-1 mutant with a

proline substitution at histidine 516 in the DIII central helix yielded a condensed form of gB that

allowed assignment of DI-DV within a 9Å resolution cryo-ET structure [31]. In this structure, DI

and DII from each of three protomers created a tripod that encapsulated DIV, which was inverted

compared to the postfusion structure, with DIII protruding from the top of the prefusion gB

trimer while DV was centrally located and connected the ectodomain via transmembrane helices

to the cytoplasmic domain. A 3.6Å resolution single particle cryo-EM structure of HCMV gB

extracted from purified virus particles with an architecture similar to HSV-1 gB[H516P] was

achieved using a small molecule fusion inhibitor and chemical cross-linking [36]. Although these

structural studies provide information about the stationary phases of herpesvirus gB in pre- and

postfusion conformations, the transition between these two forms is not understood.

Here, we demonstrate the critical role of the gB DIII central helix using structure-function

studies of gB from the model pathogen VZV. Individual proline substitutions of ^{526}EHV^{528}

within VZV gB DIII inhibited gB/gH-gL mediated cell-cell fusion. Cryo-ET revealed that

gB[H527P] was present in the condensed form on the membranes of EVs. Importantly,

molecular classification of the gB trimers further revealed two classes of conformations,

suggesting a stalled process of conformational change instigated by substitution of the conserved

histidine at this position. In addition, the prefusion state of VZV gB[H527P] was supported by

binding of an anti-gB neutralizing antibody, mAb 93k, in cells transiently expressing this mutant.

Critically, proline substitutions of ^{526}EHV^{528} in the VZV pOka genome were fully compatible

with intracellular assembly of virus particles and their release at cell surfaces, as demonstrated

by a flow cytometry-based transmission electron microscopy (FC-TEM) assay. However, only

the pOka-gB[E526P] mutant yielded infectious VZV although gB-dependent cell-cell spread was

markedly impaired by this mutation. As expected, the fusion-arrested gB[H527P] mutation was

lethal to the virus. Inactivation of pOka-gB[V528P] was explained by disrupted ER egress,

preventing gB maturation and trafficking to the cell surface, associated with a drastically

distorted central helix structure predicted by AlphaFold2. Detection of complete viral particles

despite gB ER retention suggested that virus assembly can proceed without gB incorporation into

the virion envelope. Thus, the structural integrity of the gB DIII central helix is necessary not

only for the molecular transition of gB for fusogenic function but also for essential steps in gB

biosynthesis required to produce infectious herpesviruses.

Results

The gB[H527P] mutation constrains VZV gB in a prefusion conformation. VZV gB

residues, [526]EHV[528], were within the upper region of the DIII central helix of the HSV-1

gB[H516P]-based prefusion VZV gB homology model and the lower region of DIII in our gB

postfusion structure [3,20,31] (**Fig. 1a-1c**). Conservation of the histidine at position 527 across the

alphaherpesvirinae (**Fig. S1**), suggested that a proline substitution might fix VZV gB in a

prefusion form as observed with HSV-1 gB[H516P] [31]. The valine at position 528 is conserved in

the varicelloviruses, the simplexviruses, and several evolutionary divergent alphaherpesviruses,

whereas the glutamic acid at 526 is unique to VZV and one non-human primate varicellovirus

(**Fig. S1**). Although gB was detectable in cells transiently transfected with vectors expressing

WT gB or proline substitutions at [526]EHV[528], only WT gB and gB[H527P] were incorporated

into EVs (**Fig. S2a-S2d**). A preliminary cryo-ET dataset revealed that WT gB and gB[H527P]

formed different conformations in the membranes of purified EVs from transfected BHK-21

cells (**Fig. 1d**). WT gB was 160Å in height, and resembled the 2.8 Å postfusion structure of gB

purified from VZV infected cells [3]. DI-IV were discernable, but DV was not, consistent with its

location in postfusion gB. In contrast, the gB[H527P] trimers formed condensed structures of

100Å in height and 100Å in width that resembled those of HSV-1 gB[H516P], indicating that the

H527P substitution prevented the transition of gB to a postfusion conformation structure (**Fig.**

1d). Notably, WT VZV gB was predominantly in a postfusion conformation, while gB[H527P]

was exclusively in a prefusion conformation (**Fig. 1d**).

To achieve a higher resolution structure of gB[H527P] on EVs, a more extensive cryo-ET

dataset was generated and coupled with subtomogram averaging (STA) (**Fig. S2f; Table. S1**).

Clusters of gB[H527P] were evident with edge contacts between the gB molecules on the surface

of EVs (**Fig. 2a**). STA of gB[H527P] on EVs produced an initial cryo-EM map of 10Å that also

revealed intermolecular contacts (**Fig. 2b**). The overall architecture of gB[H527P] suggested a

condensed form similar to HSV-1 gB[H516P] where DI and DII formed a tripod around DIV,

which was inverted toward the membrane compared to the postfusion form of gB, and DIII

underwent a slight bend and twist that exposed this domain at the top of the trimeric molecule

(**Fig. 2b**). However, the decline of the "gold-standard" Fourier Shell Correlation (FSC) curve at

low resolution (20Å), in addition to the local resolution estimation suggested that large scale

conformational heterogeneity might be present in the system (**Fig. 2c and Fig. S2f**).

Furthermore, the upper central region of the gB trimer, where DIII was predicted to emerge, had

a lower local resolution compared to the rest of the protein (**Fig. 2c**), also indicative of

heterogeneity in the reconstruction. Initial classification of the gB[H527P] particles on the

membranes of EVs identified three classes that were randomly distributed in the EV membranes

(**Fig. S2f-S2h**). However, class III was discarded due to low resolution in the center region of the

gB trimer. The remaining two classes (class I and class II) with distinct architectures were

revealed by further subtomogram classification and focused refinement of the central region of

the gB trimer (**Fig. 2d and Fig. S2f**). The separation of the data into two classes improved the

consistency of the cryo-EM map resolution (9.8-12.4Å), which is also evident by the improved

shape of the FSC curves (**Fig. 2d**), suggesting the presence of two predominant gB architectures

on the surface of gB[H527P] induced EVs. Molecular dynamics flexible fitting (MDFF) of the

VZV gB prefusion model predicted from the HSV-1 gB[H516P] structure was used to fit the

domains DI-V into the STA maps for each class (**Fig. 2e and 2f; Movie. S1**). The locations of

the ^{526}EHV^{528} residues in the gB DIII central helix in class I and class II of gB[H527P] was

modeled and a slight bend of the central helix was seen similar to that in HSV-1 gB[H516P]

(**Fig. S3a**). Although the tripod shape of the squat form of gB was retained, there was

considerable lateral movement of the gB DI fusion loop region (**Fig. 2e and 2f box i; Movie. S1**)

and rotation of gB DIII between the two classes (**Fig. 2e and 2f box ii; Movie. S1**) as predicted

by the MDFF of the two classes. The modeled trajectories for DI and DIII between the two

classes paralleled the low resolution observed for the discarded class III, suggesting the presence

of further intermediate structures transitioning between the two predominant classes of VZV

gB[H527P]. Thus, the substitution of proline at gB H527 prevents the transition of VZV gB to a

postfusion conformation but allows movement of the gB trimer between a closed (class I) and

open (class II) condensed form. These data provide evidence for the potential transitional steps

from a pre- to postfusion conformation.

The prefusion gB[H527P] maintains the epitope of a neutralizing mAb. Because

production of EVs from gB[E526P] was limited and gB[V528P] yielded none, effects of proline

substitutions on the DIII central helix structure were modeled using AlphaFold2 (**Table. S2**).

Both E526P and H527P induced a modest deformation whereas V528P was predicted to bend the

helix by almost 90 degrees compared to WT gB DIII (**Fig. 3a**). Alanine substitutions of

^{526}EHV^{528} modeled to assess the contribution of E, H, and V side chains to DIII structure did not

significantly affect the central helix architecture (**Fig. 3a**).

The functional consequences of changes in DIII architecture due to ^{526}EHV^{528} proline

mutations was assessed using the human VZV neutralizing mAb 93k. Our previous 2.8 Å cryo-

EM structure of postfusion VZV gB in complex with mAb 93k revealed a conformational

epitope mapped to gB DIV (the crown) [3]. Neutralization of VZV infection by mAb 93k via

fusion inhibition indicates accessibility of this epitope on prefusion gB, which was supported by

our VZV gB prefusion homology model [3,20]. When the accessibility of this functional epitope

was quantified by flow cytometry to determine the frequency of transiently transfected cells

expressing the gB ^{526}EHV^{528} mutants and median fluorescence intensity (MFI) with levels of gB

expression normalized to WT gB (**Fig. 3b and Fig. S4**), the frequency of gB-expressing cells

was comparable to WT gB for all proline and alanine mutants with the exception of gB[V528P]

(**Fig. 3c**). The frequency of cells expressing the ^{526}EHV^{528} proline mutants was consistent with

the relative total protein abundance detected in western blots (**Fig. S2b**), reflecting the steady

state level of those three mutant proteins. The gB[E526P] mutant had reduced MFI compared to

WT gB (**Fig. 3c**), suggesting that this mutation might generate an allosteric conformational

change in the epitope even though the DIII central helix structure was predicted to be affected

only modestly (**Fig. 3a**). MFI for gB[V528P] was markedly reduced, attributable to disruption of

the mAb 93k epitope due to gB misfolding caused by drastic distortion of the helix (**Fig. 3a**), the

significantly decreased abundance of this mutant protein (**Fig. 3c and Fig. S2b**), or likely both.

Notably, 93k mAb binding to gB[H527P] was comparable to WT gB (**Fig. 3c**), indicating

preservation of the conformational 93k epitope and confirming that the gB[H527P] cryo-EM

structures represent forms of gB locked in prefusion conformations. Consistently, the 93k

epitope was accessible when Fab binding was modeled for the class I and class II gB[H527P]

based on 93k footprint on the gB DIV postfusion structure (**Fig. S3b**).

Targeted mutagenesis of gB ^{526}EHV^{528} disrupts gB fusion function. Given the structural

changes in gB imposed by gB[H527P] and predicted for gB[E526P] and gB[V528P] (**Fig. 2 and**

3a), together with the differences in 93k binding, the effect of these mutations on membrane

fusion was assessed using a stable reporter fusion assay (SRFA). This assay measures gB/gH-gL

mediated fusion as a surrogate for membrane fusion during virus entry and cell-cell fusion in the

absence of other VZV proteins [6,14-17]. The fusion activity for all three proline substitutions was

barely detectable compared to WT gB (**Fig. 3d; Table. 1**). Failure to induce gB/gH-gL fusion

was consistent with the inability of gB[H527P] to transition to a postfusion conformation.

Alanine mutants, gB[E526A], gB[H527A] and gB[V528A], retained fusion activity of

approximately 48%, 90% and 68% respectively compared to WT gB (**Fig. 3d; Table. 1**). These

data indicate that the DIII central helix architecture is necessary whereas the amino acid side

chain interactions of ^{526}EHV^{528} support but are not critical for gB/gH-gL mediated fusion. This

was consistent with minimal deformation of DIII by ^{526}EHV^{528} alanine substitutions predicted by

AlphaFold2 (**Fig. 3a**) and supported by the preservation of mAb 93k epitope (**Fig. 3b and 3c**).

VZV is inactivated by gB[H527P] and gB[V528P] but not gB[E526P]. To determine the

role of gB DIII in VZV replication and cell-cell spread, proline or alanine substitutions at

^{526}EHV^{528} were transferred to pOka-TKGFP Bacterial Artificial Chromosomes (BACs). As

expected, infectious VZV was recovered from MeWo cells transfected with BACs containing the

fusion competent alanine substitutions, gB[E526A], gB[H527A] and gB[V528A], and plaque

sizes were comparable to the BAC with WT gB (**Fig. 3e and 3f; Table. 1**). However, for the

proline substitutions, only pOka-TKGFP-gB[E526P] yielded infectious VZV (**Fig. 3e; Table. 1**).

Although plaque sizes of pOka-TKGFP-gB[E526P] were significantly smaller than WT pOka-

TKGFP plaques (**Fig. 3f**), expression of this gB mutant allowed the production of virus particles

that could enter neighboring cells even though gB fusion function was severely impaired (**Fig.**

3d; Table. 1). Infectious VZV was not recovered from the pOka-TKGFP-gB[H527P] or pOka-

TKGFP-gB[V528P] BACs despite five separate transfections and at least six passages to detect

residual function or emergence of compensatory mutations.

Like other herpesviruses, VZV genes are expressed at immediate early (IE), early (E), and

late (L) times after infection. The essential IE protein, IE62, thymidine kinase, an E protein, and

the L proteins ORF23, a minor capsid protein, and ORF31 (gB), were all produced in MeWo

cells transfected with the pOka BAC and the mutant BACs with proline substitutions of

^{526}EHV^{528} (**Fig. 4a and Fig. S5a**). To investigate whether VZV particle assembly was disrupted

by the inactivating mutations, gB[H527P] and gB[V528P], VZV particle morphogenesis was

assessed using a flow cytometry-based transmission electron microscopy (FC-TEM) assay; VZV

infected cells were enriched by cell sorting based on the GFP fluorescence signal of TKGFP

(**Fig. 4b**). Nucleocapsids were observed in pOka-TKGFP-gB[E526P] and pOka-TKGFP-

gB[H527P] BAC transfected cells at 48 hours with morphologies like those present in pOka-

TKGFP BAC transfected cells, including viral DNA containing C-capsids, and scaffold

containing B-capsids or empty A-capsids (**Fig. 4c**), while none were found in pOka-TKGFP-

gB[V528P] BAC transfected cells (**Fig. 4c**). Herpesviruses produce complete particles with viral

DNA containing capsids and light particles that are enveloped and contain tegument proteins but

no capsids. Both light and complete particles were observed on the surfaces of pOka-TKGFP-

gB[E526P], pOka-TKGFP-gB[V528P] and pOka-TKGFP BAC transfected cells (**Fig. 4c**),

demonstrating that proline substitutions at E526 and V528 in gB DIII did not inhibit secondary

envelopment of capsids or particle egress across the plasma membrane. The complete virus

particle on the surface of the pOka-TKGFP-gB[V528P] BAC transfected cell confirmed that

nucleocapsids could be produced despite the mutation. Notably, a genome-containing particle

was identified in the cytoplasm of pOka-TKGFP-gB[H527P] BAC transfected cells (**Fig. S5b**),

together with the light particles observed (**Fig. 4c**), indicating that gB[H527P] did not interfere

with virus particle assembly. Thus, enveloped VZV particles could be assembled even when gB

mutations prevented VZV replication.

Disrupted biosynthesis renders gB[V528P] fusion defective. To further dissect the role of

the gB DIII central helix, the intracellular localization of gB ^{526}EHV^{528} mutants was evaluated by

confocal microscopy after transient expression. Mutants with alanine substitutions were

compared to WT gB, which traffics from the endoplasmic reticulum (ER) to the trans-Golgi

network (anti-TGN46) and has a punctate distribution within early endosomes (anti-EEA1), as

we reported previously [4,7] (**Fig. 5a**). The proline substitutions at E526 and H527 exhibited

expected localization compared to WT gB, whereas gB[V528P] failed to move to either the TGN

or early endosomes (**Fig. 5a**), indicating severe disruption of gB intracellular trafficking.

Next, the effects of the ^{526}EHV^{528} mutations on gB synthesis and steps in its post-

translational maturation were compared to WT gB using glycosylase endoglycosidase H (Endo

H), which removes high-mannose N-linked glycans added to proteins in the ER but not the

complex glycan modifications acquired in the TGN; peptide-N-glycosidase F (PNGase F)

removes all N-linked glycans. Without treatment, full-length gB was ~124 kDa, and the furin

cleaved C-terminal fragments were 66 kDa and 60 kDa (**Fig. 5b**). Endo H reduced full-length gB

to approximately 95 kDa, the same size as gB treated by PNGase F, indicating ER localization of

the full-length immature gB. The C-terminal cleavage fragments were Endo H resistant, with

molecular weights of 62 kDa and 50 kDa, whereas the fragment treated by PNGase F was less

than 50 kDa, indicating that proteolytic cleavage during maturation occurred after ER egress

(**Fig. 5b**). In particular, the detection of a single C-terminal gB fragment after PNGase F

treatment implies that the 66 kDa and 60 kDa fragments for untreated gB are differentially N-

linked glycoforms and not distinct cleavage products.

The alanine substitutions of ^{526}EHV^{528} and proline substitutions at E526 and H527 did not

alter effects of Endo H or PNGase F treatment compared to WT gB, indicating that protein

synthesis and post-translational modification were not significantly disrupted (**Fig. 5b**). In

contrast, only immature, ER associated gB was produced by the gB[V528P] mutant and gB was

substantially reduced in quantity compared to WT gB (**Fig. 5b and Fig. S2b**). Absence of the C-

terminal cleaved glycoforms showed gB[V528P] was not transported out of the ER for further

processing and maturation. These data were consistent with its absence on EVs (**Fig. S2a-S2b**)

and significantly reduced abundance on cell surfaces (**Fig. S4b-S4c**).

Disruption of the canonical biosynthesis of gB, required for gB incorporation into the

envelope of VZV particles, explained why gB[V528P] did not induce cell fusion (**Fig. 3d;**

Table. 1) and pOka-TKGFP-gB[V528P] failed to replicate (**Fig. 3e; Table. 1**). Since

gB[V528A] did not differ from WT gB, the consequences are not due to an amino acid side

chain change but can be attributed to distortion of the gB DIII central helix structure by

gB[V528P] as predicted by AlphaFold2 (**Fig. 3a**). The amino acids equivalent to V528 in VZV

gB were relatively conserved among all the human herpesviruses (**Fig. S6a**) and AlphaFold2

predicted a similar distortion in the gB DIII central helix by proline substitutions at this position

(**Fig. S6b; Table. S3**) except for HHV8 gB, which required an additional G471Y mutation to

allow the bend caused by I472P (**Fig. S6b; Table. S3)** and all betaherpesviruses have a tyrosine

residue equivalent to G471 (**Fig. S6a**). The predicted deformation by proline substitution to

valine or isoleucine is a conserved feature, supporting the important role of the equivalent V528

position in maintaining the helical structure of herpesvirus gB DIII.

Discussion

Like other viral fusogens, a conformational change of herpesvirus gB from a pre- to postfusion structure is thought to drive fusion of the virion envelope with the host cell membrane for entry [2]. This is based on end-point structures for the energetically stable postfusion gB and for gB stabilized by either mutagenesis of DIII or using a fusion inhibitor and chemical cross-linking [3,21-27,31,36]. Identifying transitional states of the trimer is essential for understanding the molecular dynamics of gB as a fusogen for rational drug design and vaccine development. The present study visualizes two such presumed transitional states using cryo-ET STA and focused classification of the VZV gB mutant, H527P. The closed (class I) and open (class II) conformations of VZV gB[H527P] and the MDFF fitting of a prefusion homology model based on HSV-1 gB[H516P] into the two cryo-EM STA maps provide insights into the first steps in this dynamic process.

A critical observation about the related HSV-1 gB[H516P] mutant was that the soluble ectodomain transitions to a postfusion conformation as shown by X-ray crystallography, demonstrating that this mutation does not fully lock gB [31]. This supports the concept that prefusion gB stability requires the transmembrane domain and C-terminal domain (CTD) along with the lipid membrane [21,27,30,31]. Consistent with this notion, the CTD of VZV gB contains elements that regulate membrane fusion [4,7,37]. In addition, herpesvirus gB requires the gH-gL heterodimer for its fusion function and regulation [2,38,39]. Whether gH-gL relays a gB activation signal from cellular receptors or other viral proteins, to trigger gB-mediated membrane fusion, and whether gH-gL acts as a "gatekeeper" to spatially and temporally regulate gB from premature fusion remain active areas of investigation. Our previous studies have demonstrated the likely involvement of the gH-gL heterodimer to regulate VZV gB fusion; truncation of the

VZV gH CTD yields a hyperfusognic phenotype detrimental for skin replication [6]. Similar

findings were reported for HSV-1 where truncation of the gH CTD also increased fusion [10,40].

The regulation of gB by gH-gL is further supported by the physical interaction of the trimer with

the heterodimer [35,41]. Critically, for VZV, the present study demonstrated that predominant WT

gB on EVs adopts the postfusion conformation in the absence of gH-gL, providing further

evidence that gH-gL is required to maintain VZV gB in a prefusion conformation. Although 70%

of WT HSV-1 gB was reported to remain in a prefusion conformation on EVs in the absence of

gH-gL, the gB[H516P] mutation was still required to stably lock the trimer in a prefusion

conformation on EVs [31]. Together, these data support our observations that without the

constraints of gH-gL, the VZV gB[H527P] is caught between to kinetically trapped

conformations, presumably early transitions states.

The two VZV gB[H527P] classes imply that the previously reported prefusion structures for

HSV-1 gB[H516P] or HCMV might not represent the complete array of prefusion gB

conformations. The architecture of the HSV-1 gB[H516P] structure is similar to the more open

conformation of the class II structure for VZV gB[H527P] in the present study [31]. In addition, the

soluble, truncated HSV-1 gB[H516P] ectodomain forms a postfusion structure, inferring that the

full-length proline mutant only becomes trapped during its transition in the context of the

membrane [31]. Furthermore, HCMV gB was captured using a thiourea fusion inhibitor, WAY-

174865, and chemical cross-linking. WAY-174865 binds to the gB fusion loops, membrane

proximal region (MPR), and transmembrane domain (TM) through interactions with

hydrophobic and aromatic residues, including the critical fusion loop residues Y153 and Y155 [36].

Importantly, the WAY-174865 binding pockets in the HCMV gB trimer are located internally in

between the α8 and α9 glycine-rich amphipathic helices, indicating that the gB trimer must be in

an open configuration for the compound to gain access to the MPR, TM, and fusion loops as

seen for the 3.6Å resolution cryo-EM structure [36]. While the activity data for WAY-174865 is

not available, previous studies on two closely related thiourea compounds, CF102 and WY1768,

demonstrated that these HCMV fusion inhibitors do not stably interact with cell-free virus

particles and are only effective once virus particles bind to the cell, preventing a complete

conformational change of gB triggered by receptor interaction [42,43]. CF102 inhibition escape

mutations have also been identified in the WAY-174865 binding pocket, including α9

amphipathic helix [36], providing structural evidence for the shared mechanism of action of WAY-

174865 and related thiourea compounds. These studies indicate that thiourea fusion inhibitors

target exposed regions in gB transitional states rather than a more closed prefusion conformation

similar to VZV gB[H527P] class I in the present study. This concept was proposed for HCMV

gB where a "snapshot" of a "breathing" molecule was locked into an intermediate conformation

by the fusion inhibitor and by the cross-linking agent [44]. Together, this supports the notion that

additional gB prefusion conformations can populate herpesvirus envelopes.

The absence of gB[E526P] on EVs and the predominance of immature gB in transfected

cells (**Fig. S2a-S2b**) was consistent with a moderate decrease in cell surface trafficking (**Fig.

S4b-S4c**), which might contribute to disrupted gB/gH-gL mediated cell-cell fusion activity. In

addition, although the effect of E526P on the central helix structure was minimal as predicted by

AlphaFold2, the mutation might induce an allosteric change in DIV since binding of neutralizing

antibody mAb 93k to its conformational DIV epitope was reduced (**Fig. 3c**). The barrier to

gB/gH-gL dependent fusion is higher under cell-cell fusion assay conditions compared to VZV

infected cells in culture, as observed with gB[E526P] inhibition of cell-cell fusion but recovery

of pOka-TKGFP-gB[E526P] virus and plaque formation, albeit limited. These findings

corroborated the importance of assessing the contributions of gB/gH-gL to fusion function by

testing mutations in the viral genome. Furthermore, although pOka-TKGFP-gB[E526P]

produced infectious virus, the marked reduction in plaque sizes was in the range associated with

failure of VZV to replicate in human skin xenografts in the SCID mouse model of VZV

pathogenesis, supporting the conclusion that even slight deviations in the gB central helix will be

detrimental for herpesviruses as pathogens in their native hosts [4,6,14].

In line with the severe DIII central helix distortion predicted by AlphaFold2, the proline

substitution at V528 had the greatest affect as gB[V528P] did not exit the ER, severely limiting

intracellular trafficking necessary for gB maturation required to execute its fusion function.

However, while lethal for replication, defective biogenesis of gB[V528P] was compatible with

VZV particle assembly, showing that morphogenesis can occur independently of gB maturation

and trafficking, as reported for other herpesviruses [45,46]. The fate of gB[V528P] was likely

proteolysis as proteins misfolded in the ER eventually translocate to the proteasome for

degradation [47]. V528 is fully conserved in all alphaherpesvirus orthologues of gB except for the

distantly related TeHV3 (Testudinid alphaherpesvirus 3) which has an asparagine (**Fig. S1**). The

betaherpesviruses and gammaherpesviruses have a conserved change to isoleucine at the residue

equivalent to V528 (**Fig. S6a**). A similar severe distortion in DIII was predicted for the

equivalent gB proline substitution in all human herpesvirus, implying a critical role in gB

biogenesis likely conserved for the *Herpesviridae*.

The induction of neutralizing antibodies that inhibit virus entry is a cornerstone of vaccine

design to protect against disease caused by enveloped viruses, first achieved with live attenuated

vaccines. Since neutralizing antibodies that target epitopes accessible on the prefusion

conformation of viral fusogens might be more potent, prefusion immunogens have become a

strategy for subunit vaccine design. Proline substitutions targeting the helical structure of the

protein (i.e., the heptad repeat in the coiled-coil region) have been used to stabilize several class I

fusion proteins for evaluation as candidate vaccines, including the HIV-1 envelope, the

respiratory syncytial virus fusion protein, and the spike proteins of the coronaviruses, Middle

East respiratory syndrome (MERS)-CoV, severe acute respiratory syndrome coronavirus (SARS-

CoV), and SARS-CoV-2 [48-52]. Our work indicates that proline stabilized VZV gB has the

potential to elicit neutralizing antibodies predominantly to prefusion gB and impair the transition

to the postfusion conformation effectively rendering VZV incapable of membrane fusion and cell

entry. Due to the highly conserved nature of gB, such an approach in vaccine design could be

applicable to all herpesviruses to reduce the burden of disease in animal or human populations.

Methods

Cells. CHO-DSP1 cells [17], derived from Chinese Hamster Ovary (CHO) K1 cells (CCL-61,

ATCC) that express Dual Split Protein (DSP)$_{1-7}$ of the chimeric reporter protein composed of

split GFP and split *Renilla* luciferase, were propagated using F-12K nutrient mixture medium

(Corning) supplemented with 10% fetal bovine serum (FBS, Gibco), penicillin (100 U/ml,

Gibco), and streptomycin (100 µg/ml, Gibco). Human melanoma MeWo cells (HTB-65, ATCC)

and MeWo-DSP2 derived from MeWo cells that express DSP$_{8-11}$ [17] were propagated in minimal

essential medium (MEM, Corning) supplemented with 10% FBS, nonessential amino acids (1X,

Corning), penicillin (100 U/ml, Gibco), streptomycin (100 µg/ml, Gibco) and amphotericin B

(0.5 µg/ml, Corning). CHO-DSP1 and MeWo-DSP2 cells were kept under puromycin selection

(8 µg/ml for CHO-DSP1 and 5 µg/ml for MeWo-DSP2; Gibco). BHK-21 cells were grown in

GMEM supplemented with 20 mM Hepes pH 7.4, 2% (v/v) TPB and 2% (v/v) FBS.

Sequence alignment. The amino acid sequence of alphaherpesvirus gB orthologues and

human herpesvirus gB orthologues were obtained from the NCBI

(https://www.ncbi.nlm.nih.gov/protein). The amino acids that aligned with VZV gB DIII central

helix were defined using Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo).

Construction of gB expressing vectors with mutations. gB mutant constructs were

generated from the pCAGGS-gB[WT] vector, which was a gift from Tadahiro Suenaga and

Hisashi Arase (Osaka University, Osaka, Japan) [16]. Primers pairs XmaI-Forward (5'-

ATGTGTAAAGGAGGAAGCCCGGGC-3')/gB-E526P-Reverse (5'pho-

AACATGCGGTTGAATGTGGTCATATG-3'), XmaI-Forward/gB-H527P-Reverse (5'pho-

AACTGGCTCTTGAATGTGGTCATATG-3'), XmaI-Forward/gB-V528P-Reverse (5'pho-

AGGATGCTCTTGAATGTGGTCATATG-3'), XmaI-Forward/gB-E526A-Reverse (5'pho-

AACATGGGCTTGAATGTGGTCATATG-3'), XmaI-Forward/gB-H527A-Reverse (5'pho-

AACCGCCTCTTGAATGTGGTCATATG-3'), XmaI-Forward/gB-V528A-Reverse (5'pho-

TGCATGCTCTTGAATGTGGTCATATG-3'), gB-EHV-Forward (5'pho-

AATGAAATGTTGGCACGTATCTCC-3') /AgeI-Reverse (5'-

AAGTTTAAGCACGTACCGGTACGC-3') containing the desired mutation were used to

amplify two PCR products using AccuPrime Taq (Invitrogen). The amplicons were digested with

either XmaI or AgeI restriction enzymes (New England Biolabs), and ligation was performed

with the digested sticky end and the 5' phosphate blunt end. Ligated products were

electroporated into TOP10F' Electrocomp E. Coli (Invitrogen) and plated under ampicillin

selection. Clones were confirmed by AgeI/XmaI digestion and sequencing using primers gB-

pCAGGS-Seq-F (5'-TTCGCGATGAGTATGCACAC-3') and gB-pCAGGS-Seq-R (5'-

TTGGGTTTCTCGGCAAAGGGATCC-3').

Generation of pOka-DX BACs with mutations in gB (ORF31). The pOka-TK-GFP virus

was engineered from the parental Oka strain of VZV (pOka) derived from a self-excisable

bacterial artificial chromosome (BAC) [53] to express EGFP conjugated to the thymidine kinase

(TK) [6]. A pPOKA-ΔORF31-TKGFP BAC lacking ORF31 was constructed previously [4]. Briefly,

the shuttle vector, gB-Kan containing ORF31was digested by BgIII and NdeI enzymes (New

England Biolabs). pCAGGS-gB plasmids containing the desired mutation were also digested

with BgIII and NdeI enzymes, and the fragment of interest was inserted into the shuttle vector

gB-Kan by ligation, and were electroporated into TOP10F' Electrocomp E. Coli (Invitrogen) and

plated under kanamycin selection. Clones were confirmed by AgeI/XmaI digestion and

sequencing using gB-Kan-Seq-F (5'-TAATCATTCCCACATCATGGACTGC-3') and gB-Kan-

Seq-R (5'-CCCCTTGCATTACTGTTTATGTAAG-3'). Then the gB-Kan vector carrying the

mutations were used to generate BACs using GS1783 cells containing the pPOKA-ΔORF31-

TKGFP BAC by Red recombination [11,53,54]. Fragment insertion was confirmed by PCR using

primers [31]F56625-56645 (5'-AGGTATAGGCAGTTCCCACGG-3') and [31]R59697-59717

(5'- TTTCATTGAGACTTGAAGCGC-3'). All BACs were purified using large-construct

purification kit (Qiagen), and digested by HindIII to verify that spurious recombination had not

occurred. To generate BAC-derived viruses, MeWo cells were transfected in at least triplicate

with two BAC clones for each mutation. Viruses were typically recovered at 5-10 days

posttransfection and passed on MeWo cells to propagate and titrated on MeWo cells.

EVs Production. EVs were prepared as described [33]. In brief, BHK-21 cells, at around 70%

confluency, were transiently transfected with pCAGGS vectors expressing proline substitutions

to $^{526}EHV^{528}$, or C-terminal His-tagged WT gB or gB[H527P]. Cells were grown for an

additional 48 hours with a media exchange to serum free GMEM after 24 hours. Vesicles were

harvested from the supernatant by centrifugation through a 20% sucrose cushion and

resuspended in 20 mM Hepes pH 8, 130 mM NaCl.

Cryo-ET data collection using the Titan Krios. For grid preparation, 3.5 µL vesicles

were mixed with 1.5 µL 5 nm gold fiducials on Quantifoil R2/1 grids and plunge frozen in a

propane/ethane mixture after blotting using a manual plunge freezer. Microscopy of WT gB,

gB[H527P], and gB[H527P]-His EVs was performed at 300 keV on a Titan Krios electron

microscope (Thermo Fisher scientific) equipped with a Gatan K3 (5 µm/pixel) camera and

Bioquantum post-column energy filter controlled using SerialEM [55] for micrograph and

automated batch tomography data collection. Movie data (11,283 total stacks) were captured in

counted mode with a dose rate of $\sim$10.75 – 11.9 e$^-$/Å^2/s and 50 millisecond exposure time per

frame and 0.2-0.3 second total exposure time per tilt at a nominal magnification of 81,000x and a

pixel size of 1.095 Å/pixel on the specimen. The defocus was set to -2.0µm.

Reconstruction of the cryo-ET maps for gB[H527P]. The motion correction, damage

compensation, and tilt stacks for all movie-mode data were generated using WARP [56]. All data

processing described was performed using the tomography workflow in EMAN 2.99 [57,58]. Forty

tilt series were automatically reconstructed, and 2,074 particles were manually selected from the

tomograms. The CTF were estimated from the tilt images before unbinned, phase flipped

subtomograms were extracted. The initial orientation of each particle was assigned using a vector

pointing from the center of the vesicle that the particle was located on, to the coordinates of that

particle. The initial model was directly generated from the particles at 30Å resolution and used as

a reference "gold-standard" subtomogram to perform subtilt refinement on all particles. A soft

cylinder mask that includes the protein and membrane density was used during the refinement,

and C3 symmetry was applied to the averaged structure. After each iteration of the refinement,

local resolution was measured based on the two half-maps, and the averaged structures were

filtered according to the local resolution. Then, to resolve the structural heterogeneity at the core

of the protein revealed by the local resolution, focused classification was performed based on the

orientation determined from the single model refinement, using a mask that covers the central

part of the protein. Finally, separate subtomogram and subtilt refinements were performed for the

particles of each class from the refinement, resulting in the final averaged structures. The HSV-

1-based VZV gB prefusion homology model [20] was fitted into the class I and class II cryo-ET

maps by domain restricted Molecular Dynamics Flexible Fitting (MDFF) using Visual Molecular

Dynamics (VMD 1.9.4a29; http://www.ks.uiuc.edu/Research/vmd/) [59] to prepare the input and

configuration files and the Nanoscale Molecular Dynamics (NAMD 2.14;

http://www.ks.uiuc.edu/Research/namd/) [60] software to perform simulations with the

CHARMM36 force field [61]. All images and movies were generated using UCSF ChimeraX 1.14

[62].

AlphaFold2 predictions of herpesvirus gB DIII central helix. The gB DIII central helix

for eight human herpesvirus gB orthologs was defined by amino acid alignment of the complete

gB proteins using Clustal X and comparison to the native VZV gB structure (PDB 6VN1) DIII

central helix. AlphFold2 [63,64] was compiled following the installation instructions

(https://github.com/deepmind/alphafold) on a Colfax ProEdge DX935 Workstation with an Intel

Xeon W-2195 18C/36T 2.3Ghz CPU, 4 PNY GeForce RTX2080Ti 11GB GPUs, 256MB RAM

and Centos 8 operating system.

Stable-reporter fusion assays. CHO-DSP1 were transfected with vectors pCAGGS-

gB[WT], pCAGGS-gB[E526P], pCAGGS-gB[H527P], pCAGGS-gB[V528P], pCAGGS-

gB[E526A], pCAGGS-gB[H527A], or pCAGGS-gB[V528A]; pCAGGS-gB[WT]-His-tagged, or

pCAGGS-gB[H527P]-His-tagged, with or without vectors expressing gH-gL, pME18S-gH[TL]

and pcDNA3.1-gL using Lipofectamine 2000. At 6 hrs post-transfection, the transfected cells

were harvested and mixed with MeWo-DSP2 cells for additional 48 hrs. The activity of *Renilla*

luciferase was read immediately after adding substrate h-Coelenterazine (Nanolight Technology)

on a Synergy H1 Hybrid Multi-Mode Reader (BioTek). Experiments were performed in triplicate

at least. The fusion activity for His-tagged WT gB and gB[H527P] was shown in **Fig. S2e**.

Quantitation of cell surface and total expression of gB. The quantity of cell surface VZV

glycoproteins were measured as described previously [4]. Briefly, CHO-DSP1 cells were

transfected with vectors pCAGGS-gB[WT], or pCAGGS-gB[E526P], pCAGGS-gB[H527P],

pCAGGS-gB[V528P], pCAGGS-gB[E526A], pCAGGS-gB[H527A], and pCAGGS-

gB[V528A], or vectors expressing gH-gL, pME18S-gH[TL] and pcDNA3.1-gL. At 24 hrs later,

cells were dislodged using an enzyme-free cell dissociation buffer (Gibco), fixed with 1%

paraformaldehyde (PFA, Boston Bioproducts), and resuspended in FACS staining buffer [PBS

with 0.2% IgG-free bovine serum albumin (Jackson ImmunoResearch), and 0.1% NaN_3 (Sigma

Aldrich)]. Cell surface gB were detected with primary antibodies mouse anti-VZV gB mAb

(93k; human) [3] conjugated with Cy5, at 1:100 dilution. Total amounts of viral glycoprotein

production were performed by the same staining protocol, except cells were permeabilized using

Cytofix/Cytoperm kit (BD Biosciences) before adding the antibodies and during the following

procedure. Stained cells were analyzed using a DXP multi-color FACScan analyzer (Cytek

Biosciences), and data were processed with FlowJo (TreeStar) to determine the quantity of total

and surface levels of gB respectively. The ratio of cell surface to total quantity was also

calculated. Experiments were performed in triplicate for proline substitution mutants and in

duplicate for alanine substitution mutants.

Immunoprecipitation and Western blotting. Whole cell lysate from MeWo cells

transiently expressing WT gB or gB with proline and alanine substitutions, or gH-gL, a non-

antigen negative control was extracted at 24 hrs posttransfection using extraction buffer [0.1 M

NaCl, 5 mM KCl (Fisher Scientific), 1 mM $CaCl_2$ (Fisher Scientific), 0.5 mM $MgCl_2$ (Fisher

Scientific), 1% IGEPAL CA-630 (Sigma Aldrich), 1% Deoxycholate (Sigma Aldrich) in 0.1 M

Tris buffer, pH 7.2], with cOmplete EDTA-free protease inhibitor, for 30 min on ice. The sample

was centrifuged at 3,000 RCF for 10 min at 4°C to get rid of cell debris. The supernatant was

immunoprecipitated (IP) with mouse anti-VZV gB mAb (SG2-2E6, Meridian Life Sciences) that

was crosslinked to protein A beads (Pierce) at 20 µg antibody/30 µl beads slurry. Samples were

then either left untreated or treated with Endo H (New England Biolabs) or PNGase F (New

England Biolabs) according to the manufacture's protocol. Proteins were denatured under

reducing condition and separated by 7.5% SDS PAGE gel (BioRad), transferred to Immobilon-P

PVDF membranes (Millipore) using Trans-blot SD Semi-dry Transfer cell (BioRad) in transfer

buffer [48 mM Tris Base (Fisher Scientific), 39 mM Glycine (Fisher Scientific) and 0.01% SDS

(Sigma Aldrich), with 20% methanol (Sigma Aldrich)]. The transferred proteins were probed for

gB using rabbit polyclonal antiserum, 746-868, that recognizes the epitope

[833]PEGMDPFAEKPNAT[846] in the cytoplasmic domain of gB (GenScript Corp) [14], followed by

anti-rabbit IgG horseradish peroxidase-linked secondary antibodies (GE Healthcare Life

Sciences, UK Ltd., 1:10,000) and Pierce ECL Plus Western Blotting Substrate (ThermoFisher).

Chemiluminescence was detected using BioMax MR Film (Carestream). For EVs analysis, after

supernatants were removed for vesicle preparations, cells were washed with cold PBS and

detached using cell scrapers. Cells were pelleted by centrifugation (5 minutes, 4500 RCF, 4°C),

transferred into 1.5 ml microfuge tubes and washed in cold PBS again before resuspension in

RIPA buffer, 100 µL per T175 flask (50 mM Tris pH 8, 1% NP-40, 0.1% SDS, 150 mM NaCl,

0.5% Sodium Deoxycholate, 5 mM EDTA, 1 mM PMSF). Samples were shaken at 4°C for 30

min before centrifugation at 500 RCF for 10 minutes. Supernatants were mixed in SDS sample

buffer and run in parallel with vesicle samples in SDS PAGE, followed by Coomassie staining or

western blotting with the rabbit anti-VZV gB antibody 746-868 followed by anti-rabbit-HRP

(Sigma-Aldrich Chemie GmbH). For loading control, western blots were re-probed using a

mouse anti-GAPDH antibody (Sigma-Aldrich Chemie GmbH) followed by anti-mouse-HRP

(Sigma-Aldrich Chemie GmbH). The Coomassie staining and Western blotting for His-tagged

WT gB and gB[H527P] using rabbit anti-His6 antibody (Abcam) was shown in **Fig. S2c and**

S2d. Band density was analyzed using ImageJ.

Confocal microscopy. MeWo cells transfected with gB expressing vectors were fixed 24

hrs posttransfection with 4% formaldehyde for 10 min. mAb SG2 (Meridian Life Sciences) was

used to stain for VZV gB (1:200), and primary antibodies to cellular proteins were used as

markers for early endosomes (EEA-1; rabbit polyclonal antibody; Santa Cruz Biotechnology Inc;

1:200), and the trans-Golgi (TGN46; sheep polyclonal antibody; BioRad; 1:250). Primary

antibodies were detected with secondary antibodies, anti-mouse Alexa Fluor 555 (Invitrogen),

anti-rabbit Alexa 488, and anti-sheep Alexa Fluor 647 (Invitrogen). Nuclei were stained with

Hoechst 33342 (Invitrogen). MeWo cells transfected with pOka, pOka-TKGFP-gB[E526P],

pOka-TKGFP-gB[H527P], or pOka-TKGFP-gB[V528P] BACs were fixed at 48 hrs or 72 hrs

post transfection with 4% formaldehyde for 10 min, and cells were probed using primary

antibodies to VZV proteins IE62 (mouse mAb; EMD Millipore MAB8616; 1:200), capsid-

ORF23 (polyclonal rabbit; 1:200) [65] and gB (93k; human mAb; 1:200) [3] and nuclei (Hoechst

33342). Primary antibodies were detected with secondary antibodies, anti-mouse Alexa Fluor

555 (Invitrogen), and anti-rabbit or anti-human Alexa Fluor 647 (Invitrogen). Nuclei were

stained with Hoechst 33342 (Invitrogen). Images were captured with a Leica SP8, Whitelight

Laser Confocal Microscope. Channel merging and image processing was performed with ImageJ

and Photoshop.

Plaque size assay. Monolayers of MeWo cells (1×10^5 cell/cm^2) were inoculated with 50

plaque forming unit (PFU) of cell-associated pOka-TKGFP viruses or pOka-TKGFP-gB[E526P]

for 2 hrs, followed by change of medium. Cells were fixed with 4% PFA at 4 days post-infection

and processed with Immunohistochemistry (IHC) using mouse monoclonal anti-VZV antibody

(Meridian Life Sciences C05108MA, 1:2000 diluted in PBS), followed by incubation with

biotinylated anti-mouse IgG (Vector Laboratories #BA-9200), and alkaline phosphatase

streptavidin (Jackson ImmunoResearch #016-050-084). The plaques were visualized by staining

with a mixture of FastRed salt (Sigma Aldrich) and Naphthol AS-MX phosphate (Sigma

Aldrich). Images of plaques were taken with Axio microscope (Zeiss) using 2.5X lens. The

plaques were outlined, 15 plaques were measured for each experiment, and the area of plaque

were measured using ImageJ. Experiments were performed in triplicate.

FC-TEM of BAC transfected cells. Monolayers of MeWo cells (1.6×10^5 cell/cm^2) were

seeded onto P100 dish and were transfected with 29 µg pOka-TKGFP, pOka-TKGFP-

gB[E526P], pOka-TKGFP-gB[H527P], or pOka-TKGFP-gB[V528P] BACs by Lipofectamine

2000. Cells were harvested at 48hrs posttransfection, and resuspended in FACS staining buffer

[2.5% FBS, 2 mM EDTA in PBS] at 5×10^6 cell/ml. The GFP positive cells were sorted for

enrichment by negative gating at 0.5% of the mock sample that was MeWo cells transfected

without BAC. The sorted cells were pelleted and re-suspended in 10% Gelatin in 0.1M Sodium

Cacodylate buffer pH 7.4 at 37^{o}C and allowed to equilibrate 5 min. Cells were pelleted again,

excess gelatin removed, then chilled in cold blocks and covered with cold 1% Osmium tetroxide

(EMS Cat# 19100) for 2 hrs rotating in a cold room, washed 3 times with cold ultrafiltered

water, then en bloc stained overnight in 1% Uranyl Acetate at 4^{o}C while rotating. Samples were

then dehydrated in a series of ethanol washes for 20 minutes each at 4^{o}C beginning at 30%, 50%,

70%, 95% where the samples were then allowed to rise to RT, changed to 100% ethanol, then

Propylene Oxide (PO) for 15 min. Samples were infiltrated with EMbed-812 resin (EMS

Cat#14120) mixed 1:2, 1:1, and 2:1 with PO, and then were placed into neat EMbed-812 for 2 to

4 hrs followed by placement into molds at 65^{o} C oven overnight. Sections were taken

approximately 80nm, picked up on formvar/Carbon coated 100 mesh Cu grids, stained for 30 sec

in 3.5% Uranyl Acetate in 50% Acetone followed by staining in 0.2% Lead Citrate for 3 min.

Observed in the JEOL JEM-1400 120kV and photos were taken using a Gatan Orius 4k X 4k

digital camera. The yield efficiency of FC-TEM was summarized in **Table S4.**

Data availability

592   Data generated and/or analyzed during the current study are available in the paper or are

593 appended as supplementary data. The source data underlying Figs. 3c, 3d, 3e, 5b, S2b, S2d, S2e,

594 S4c have been provided as a Source Data file. Source data are provided with this paper. All

595 primary data will be provided by the corresponding author upon reasonable request.

597 **References**

598 1  Arvin, A. M., & Abendroth, A. Varicella-zoster Virus, Ch. 13, in Howley, P. M., Knipe,

599   D. M. Editors. *Fields virology. Volume 2,* 7[th] edition (Lippincott Williams & Wilkins,

600   Philadephia, 2021).

2   Connolly, S. A., Jardetzky, T. S. & Longnecker, R. The structural basis of herpesvirus

entry. *Nat Rev Microbiol* **19**, 110-121, doi:10.1038/s41579-020-00448-w (2021).

3   Oliver, S. L. *et al.* A glycoprotein B-neutralizing antibody structure at 2.8 A uncovers a

critical domain for herpesvirus fusion initiation. *Nat Commun* **11**, 4141,

doi:10.1038/s41467-020-17911-0 (2020).

4   Oliver, S. L. *et al.* An immunoreceptor tyrosine-based inhibition motif in varicella-zoster

virus glycoprotein B regulates cell fusion and skin pathogenesis. *Proc Natl Acad Sci U S*

A **110**, 1911-1916, doi:10.1073/pnas.1216985110 (2013).

5   Heineman, T. C. & Hall, S. L. Role of the varicella-zoster virus gB cytoplasmic domain

in gB transport and viral egress. *J Virol* **76**, 591-599, doi:10.1128/jvi.76.2.591-599.2002

(2002).

6   Yang, E., Arvin, A. M. & Oliver, S. L. The cytoplasmic domain of varicella-zoster virus

glycoprotein H regulates syncytia formation and skin pathogenesis. *PLoS Pathog* **10**,

e1004173, doi:10.1371/journal.ppat.1004173 (2014).

7   Yang, E., Arvin, A. M. & Oliver, S. L. The Glycoprotein B Cytoplasmic Domain Lysine

Cluster Is Critical for Varicella-Zoster Virus Cell-Cell Fusion Regulation and Infection. *J*

Virol **91**, doi:10.1128/JVI.01707-16 (2017).

8   Vallbracht, M. *et al.* In Vitro Viral Evolution Identifies a Critical Residue in the

Alphaherpesvirus Fusion Glycoprotein B Ectodomain That Controls gH/gL-Independent

Entry. *mBio* **12**, doi:10.1128/mBio.00557-21 (2021).

9   Pataki, Z., Rebolledo Viveros, A. & Heldwein, E. E. Herpes Simplex Virus 1 Entry

Glycoproteins Form Complexes before and during Membrane Fusion. *mBio*, e0203922,

doi:10.1128/mbio.02039-22 (2022).

10   Pataki, Z., Sanders, E. K. & Heldwein, E. E. A surface pocket in the cytoplasmic domain

of the herpes simplex virus fusogen gB controls membrane fusion. *PLoS Pathog* **18**,

e1010435, doi:10.1371/journal.ppat.1010435 (2022).

11   Oliver, S. L., Yang, E. & Arvin, A. M. Varicella-Zoster Virus Glycoproteins: Entry,

Replication, and Pathogenesis. *Curr Clin Microbiol Rep* **3**, 204-215, doi:10.1007/s40588-

016-0044-4 (2016).

12   Harrison, S. C. Viral membrane fusion. *Virology* **479**, 498-507,

doi:10.1016/j.virol.2015.03.043 (2015).

13   Zerboni, L., Sen, N., Oliver, S. L. & Arvin, A. M. Molecular mechanisms of varicella

zoster virus pathogenesis. *Nat Rev Microbiol* **12**, 197-210, doi:10.1038/nrmicro3215

(2014).

14   Oliver, S. L. *et al.* Mutagenesis of varicella-zoster virus glycoprotein B: putative fusion

loop residues are essential for viral replication, and the furin cleavage motif contributes to

pathogenesis in skin tissue in vivo. *J Virol* **83**, 7495-7506, doi:10.1128/JVI.00400-09

(2009).

15   Vleck, S. E. *et al.* Structure-function analysis of varicella-zoster virus glycoprotein H

identifies domain-specific roles for fusion and skin tropism. *Proc Natl Acad Sci U S A*

108, 18412-18417, doi:10.1073/pnas.1111333108 (2011).

16   Suenaga, T. *et al.* Myelin-associated glycoprotein mediates membrane fusion and entry of

neurotropic herpesviruses. *Proc Natl Acad Sci U S A* **107**, 866-871,

doi:10.1073/pnas.0913351107 (2010).

17   Yang, E., Arvin, A. M. & Oliver, S. L. Role for the alphaV Integrin Subunit in Varicella-

Zoster Virus-Mediated Fusion and Infection. *J Virol* **90**, 7567-7578,

doi:10.1128/JVI.00792-16 (2016).

18   Reichelt, M., Zerboni, L. & Arvin, A. M. Mechanisms of varicella-zoster virus

neuropathogenesis in human dorsal root ganglia. *J Virol* **82**, 3971-3983,

doi:10.1128/JVI.02592-07 (2008).

19   Nagel, M. A. & Bubak, A. N. Varicella Zoster Virus Vasculopathy. *J Infect Dis* **218**,

S107-S112, doi:10.1093/infdis/jiy425 (2018).

20   Oliver, S. L. *et al.* The N-terminus of varicella-zoster virus glycoprotein B has a

functional role in fusion. *PLoS Pathog* **17**, e1008961, doi:10.1371/journal.ppat.1008961

(2021).

21   Heldwein, E. E. *et al.* Crystal structure of glycoprotein B from herpes simplex virus 1.

Science **313**, 217-220, doi:10.1126/science.1126548 (2006).

22   Backovic, M., Longnecker, R. & Jardetzky, T. S. Structure of a trimeric variant of the

Epstein-Barr virus glycoprotein B. *Proc Natl Acad Sci U S A* **106**, 2880-2885,

doi:10.1073/pnas.0810530106 (2009).

23   Chandramouli, S. *et al.* Structure of HCMV glycoprotein B in the postfusion

conformation bound to a neutralizing human antibody. *Nat Commun* **6**, 8176,

doi:10.1038/ncomms9176 (2015).

24   Burke, H. G. & Heldwein, E. E. Crystal Structure of the Human Cytomegalovirus

Glycoprotein B. *PLoS Pathog* **11**, e1005227, doi:10.1371/journal.ppat.1005227 (2015).

25   Vallbracht, M. *et al.* Structure-Function Dissection of Pseudorabies Virus Glycoprotein B

Fusion Loops. *J Virol* **92**, doi:10.1128/JVI.01203-17 (2018).

26   Li, X. *et al.* Two classes of protective antibodies against Pseudorabies virus variant

glycoprotein B: Implications for vaccine design. *PLoS Pathog* **13**, e1006777,

doi:10.1371/journal.ppat.1006777 (2017).

27   Cooper, R. S., Georgieva, E. R., Borbat, P. P., Freed, J. H. & Heldwein, E. E. Structural

basis for membrane anchoring and fusion regulation of the herpes simplex virus fusogen

gB. *Nat Struct Mol Biol* **25**, 416-424, doi:10.1038/s41594-018-0060-6 (2018).

28   Roche, S., Bressanelli, S., Rey, F. A. & Gaudin, Y. Crystal structure of the low-pH form

of the vesicular stomatitis virus glycoprotein G. *Science* **313**, 187-191,

doi:10.1126/science.1127683 (2006).

29   Kadlec, J., Loureiro, S., Abrescia, N. G., Stuart, D. I. & Jones, I. M. The postfusion

structure of baculovirus gp64 supports a unified view of viral fusion machines. *Nat Struct

Mol Biol* **15**, 1024-1030, doi:10.1038/nsmb.1484 (2008).

30   Vitu, E., Sharma, S., Stampfer, S. D. & Heldwein, E. E. Extensive mutagenesis of the

HSV-1 gB ectodomain reveals remarkable stability of its postfusion form. *J Mol Biol*

425, 2056-2071, doi:10.1016/j.jmb.2013.03.001 (2013).

31   Vollmer, B. *et al.* The prefusion structure of herpes simplex virus glycoprotein B. *Sci Adv*

6, doi:10.1126/sciadv.abc1726 (2020).

32   Zeev-Ben-Mordehai, T., Vasishtan, D., Siebert, C. A., Whittle, C. & Grunewald, K.

Extracellular vesicles: a platform for the structure determination of membrane proteins by

Cryo-EM. *Structure* **22**, 1687-1692, doi:10.1016/j.str.2014.09.005 (2014).

33   Zeev-Ben-Mordehai, T. *et al.* Two distinct trimeric conformations of natively membrane-

anchored full-length herpes simplex virus 1 glycoprotein B. *Proc Natl Acad Sci U S A*

113, 4176-4181, doi:10.1073/pnas.1523234113 (2016).

34   Fontana, J. *et al.* The Fusion Loops of the Initial Prefusion Conformation of Herpes

692   Simplex Virus 1 Fusion Protein Point Toward the Membrane. *mBio* **8**,

693   doi:10.1128/mBio.01268-17 (2017).

694   35   Si, Z. *et al.* Different functional states of fusion protein gB revealed on human

695   cytomegalovirus by cryo electron tomography with Volta phase plate. *PLoS Pathog* **14**,

696   e1007452, doi:10.1371/journal.ppat.1007452 (2018).

697   36   Liu, Y. *et al.* Prefusion structure of human cytomegalovirus glycoprotein B and structural

698   basis for membrane fusion. *Sci Adv* **7**, doi:10.1126/sciadv.abf3178 (2021).

699   37   Oliver, S. L., Zhou, M. & Arvin, A. M. Varicella-zoster virus: molecular controls of cell

700   fusion-dependent pathogenesis. *Biochem Soc Trans* **48**, 2415-2435,

701   doi:10.1042/BST20190511 (2020).

702   38   Gonzalez-Del Pino, G. L. & Heldwein, E. E. Well Put Together-A Guide to

703   Accessorizing with the Herpesvirus gH/gL Complexes. *Viruses* **14**,

704   doi:10.3390/v14020296 (2022).

705   39   Stampfer, S. D. & Heldwein, E. E. Stuck in the middle: structural insights into the role of

706   the gH/gL heterodimer in herpesvirus entry. *Curr Opin Virol* **3**, 13-19,

707   doi:10.1016/j.coviro.2012.10.005 (2013).

708   40   Rogalin, H. B. & Heldwein, E. E. Interplay between the Herpes Simplex Virus 1 gB

709   Cytodomain and the gH Cytotail during Cell-Cell Fusion. *J Virol* **89**, 12262-12272,

710   doi:10.1128/JVI.02391-15 (2015).

711   41   Vanarsdall, A. L., Howard, P. W., Wisner, T. W. & Johnson, D. C. Human

712   Cytomegalovirus gH/gL Forms a Stable Complex with the Fusion Protein gB in Virions.

713   *PLoS Pathog* **12**, e1005564, doi:10.1371/journal.ppat.1005564 (2016).

42 Jones, T. R. *et al.* Specific inhibition of human cytomegalovirus glycoprotein B-mediated

fusion by a novel thiourea small molecule. *J Virol* **78**, 1289-1300,

doi:10.1128/jvi.78.3.1289-1300.2004 (2004).

43 Patrone, M., Secchi, M., Bonaparte, E., Milanesi, G. & Gallina, A. Cytomegalovirus

UL131-128 products promote gB conformational transition and gB-gH interaction during

entry into endothelial cells. *J Virol* **81**, 11479-11488, doi:10.1128/JVI.00788-07 (2007).

44 Dormitzer, P. R. N., NY, US), Che, Ye (Niantic, CT, US), Chi, Xiaoyuan Sherry

(Tenafly, NJ, US), Han, Seungil (Mystic, CT, US), Heim, Kyle Paul (Boulder, CO, US),

Jones, Thomas Richard (Bluffton, SC, US), Liu, Yuhang (South Glastonbury, CT, US),

Qiu, Xiayang (Mystic, CT, US), Yang, Xinzhen (Woodcliff, NJ, US), Yao, Xiaojie (East

Lyme, CT, US), Griffor, Matthew Curtis (North Stonington, CT, US), Nicki, Jennifer

Anne (Gales Ferry, CT, US). HUMAN CYTOMEGALOVIRUS GB POLYPEPTIDE.

United States patent (2020).

45 Farnsworth, A. *et al.* Herpes simplex virus glycoproteins gB and gH function in fusion

between the virion envelope and the outer nuclear membrane. *Proc Natl Acad Sci U S A*

104, 10187-10192, doi:10.1073/pnas.0703790104 (2007).

46 Klupp, B., Altenschmidt, J., Granzow, H., Fuchs, W. & Mettenleiter, T. C. Glycoproteins

required for entry are not necessary for egress of pseudorabies virus. *J Virol* **82**, 6299-

6309, doi:10.1128/JVI.00386-08 (2008).

47 Braakman, I. & Hebert, D. N. Protein folding in the endoplasmic reticulum. *Cold Spring*

Harb Perspect Biol **5**, a013201, doi:10.1101/cshperspect.a013201 (2013).

48   Sanders, R. W. *et al.* Stabilization of the soluble, cleaved, trimeric form of the envelope

736   glycoprotein complex of human immunodeficiency virus type 1. *J Virol* **76**, 8875-8889,

737   doi:10.1128/jvi.76.17.8875-8889.2002 (2002).

738   49   Krarup, A. *et al.* A highly stable prefusion RSV F vaccine derived from structural

739   analysis of the fusion mechanism. *Nat Commun* **6**, 8143, doi:10.1038/ncomms9143

740   (2015).

741   50   Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion

742   conformation. *Science* **367**, 1260-1263, doi:10.1126/science.abb2507 (2020).

743   51   Pallesen, J. *et al.* Immunogenicity and structures of a rationally designed prefusion

744   MERS-CoV spike antigen. *Proc Natl Acad Sci U S A* **114**, E7348-E7357,

745   doi:10.1073/pnas.1707304114 (2017).

746   52   Kirchdoerfer, R. N. *et al.* Stabilized coronavirus spikes are resistant to conformational

747   changes induced by receptor recognition or proteolysis. *Sci Rep* **8**, 15701,

748   doi:10.1038/s41598-018-34171-7 (2018).

749   53   Tischer, B. K. *et al.* A self-excisable infectious bacterial artificial chromosome clone of

750   varicella-zoster virus allows analysis of the essential tegument protein encoded by ORF9.

751   *J Virol* **81**, 13200-13208, doi:10.1128/JVI.01148-07 (2007).

752   54   Tischer, B. K., von Einem, J., Kaufer, B. & Osterrieder, N. Two-step red-mediated

753   recombination for versatile high-efficiency markerless DNA manipulation in Escherichia

754   coli. *Biotechniques* **40**, 191-197, doi:10.2144/000112096 (2006).

755   55   Mastronarde, D. N. Automated electron microscope tomography using robust prediction

756   of specimen movements. *J Struct Biol* **152**, 36-51, doi:10.1016/j.jsb.2005.07.007 (2005).

56   Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data preprocessing with

Warp. *Nat Methods* **16**, 1146-1152, doi:10.1038/s41592-019-0580-y (2019).

57   Tang, G. *et al.* EMAN2: an extensible image processing suite for electron microscopy. *J*

Struct Biol **157**, 38-46, doi:10.1016/j.jsb.2006.05.009 (2007).

58   Chen, M. *et al.* A complete data processing workflow for cryo-ET and subtomogram

averaging. *Nat Methods* **16**, 1161-1168, doi:10.1038/s41592-019-0591-8 (2019).

59   Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol*

Graph **14**, 33-38, 27-38, doi:10.1016/0263-7855(96)00018-5 (1996).

60   Phillips, J. C. *et al.* Scalable molecular dynamics on CPU and GPU architectures with

NAMD. *J Chem Phys* **153**, 044130, doi:10.1063/5.0014475 (2020).

61   Huang, J. & MacKerell, A. D., Jr. CHARMM36 all-atom additive protein force field:

validation based on comparison to NMR data. *J Comput Chem* **34**, 2135-2145,

doi:10.1002/jcc.23354 (2013).

62   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators,

and developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).

63   Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding the

structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids*

Res **50**, D439-D444, doi:10.1093/nar/gkab1061 (2022).

64   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583-589, doi:10.1038/s41586-021-03819-2 (2021).

65   Chaudhuri, V., Sommer, M., Rajamani, J., Zerboni, L. & Arvin, A. M. Functions of

Varicella-zoster virus ORF23 capsid protein in viral replication and the pathogenesis of

skin infection. *J Virol* **82**, 10231-10246, doi:10.1128/JVI.01890-07 (2008).

Acknowledgements

We thank the National Institutes of Health (NIH) for their valuable support through grants

R01-AI102546 and R21-MH125285. We thank the Stanford Cell Sciences Imaging Facility for

supporting the confocal microscopy and the transmission electron microscopy (TEM) work, the

project described was supported, in part, by Award Number 1S10OD010580-01A1 from the

National Center for Research Resources (NCRR) (confocal microscopy work), and by NIH S10

Award Number 1S10OD028536-01, titled "OneView 4kX4k sCMOS camera for transmission

electron microscopy applications" from the Office of Research Infrastructure Programs (ORIP)

(TEM work). Its contents are solely the responsibility of the authors and do not necessarily

represent the official views of the NCRR or the NIH. Cell sorting was performed in the Stanford

Shared FACS Facility on an instrument obtained using NIH S10 Shared Instrument Grant

S10RR025518-01. Molecular graphics and analyses performed with UCSF ChimeraX,

developed by the Resource for Biocomputing, Visualization, and Informatics at the University of

California, San Francisco, with support from NIH R01-GM129325 and the Office of Cyber

Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.

NAMD was developed by the Theoretical and Computational Biophysics Group in the Beckman

Institute for Advanced Science and Technology at the University of Illinois at Urbana-

Champaign.

We gratefully acknowledge funding from the DFG [a Cluster of Excellence RESIST (EXC

2155) project and the BMBF (05K18BHA) to K.G. and a Marie Curie Individual Fellowship

(HSV1ENTRYPROTEINMAP) to E.M.. Part of this work was performed at the CryoEM

Facility at CSSB, supported by the UHH and DFG grant numbers (INST 152/772-1|152/774-

1|152/775-1|152/776-1).

Author contributions

M.Z. constructed mutant expressing plasmids, BAC recombinant viruses, carried out

sequence alignments, flow cytometry, stable reporter fusion assays, plaque assays,

immunoprecipitation, western blotting, confocal microscopy, and FC-TEM. B.V. generated EVs,

prepared Cryo-ET grids, carried out Cryo-ET data acquisition. M.C. performed subtomogram

averaging analysis on the Cryo-ET data. S.L.O. conducted the AlphaFold2 simulation and

performed model building, together with M.C. A.M.A., K.G., S.L.O., B.V., and M.Z.,

conceptualized the project. All the authors discussed the data and contributed intellectually to the

manuscript.

Competing interests

K.G. and B.V. are inventors on the U.S. patent application 17/484,583 and international

patent application PCT/EP2022/076579, "Prefusion-Stabilized Herpesvirus Glycoprotein B".

The authors declare that they have no other competing interests.

Figure legends

Figure 1. The point mutation H527P constrains VZV gB in a prefusion conformation. a A

diagram of the linear structure for VZV gB highlighting the location of the DIII central helix.

The signal sequence is depicted by the crossed white box. Colored regions (residues 115–736)

correspond to domains in the cryo-EM structure; DI (cyan), DII (green), DIII (yellow), DIV

(orange), DV (red) and linker regions (hot pink). Amino acids [526]EHV[528] (bold) were targeted for

mutagenesis. **b and c** Trimer and monomer models for VZV gB prefusion (**b**) by SWISS Model

[20] and postfusion (**c**) from the 2.8 Å cryo-EM structure [3]. The inset shows the [526]EHV[528] site in

the DIII central helix. **d** Single plane views of representative tomographic reconstructions from

cryo-ET of EVs produced by BHK-21 cells transfected with gB[H527P] or WT gB expression

vectors. Upper and lower panels are different planes from the same tomogram to show side and

top views of gB[H527P] or WT gB expressed on EVs. Boxes with dashed lines are magnified

areas shown in the insets. Scale bars shown are for each panel and inset.

Figure 2. Cryo-ET structures of VZV gB[H527P] resident on the membranes of EVs. a

and b A representative slice view of a tomogram (**a**) and an EMAN2 cryo-EM map generated by

subtomogram averaging (STA) (**b**) from EVs purified from BHK-21 cells transiently transfected

with gB[H527P]. The white box in the upper panel of (**a**) highlights the area shown in the lower

panel. Arrow heads point to gB[H527P] on the membrane of EVs. **c** A 10Å resolution cryo-EM

map derived from a local refinement of a tight mask and the associated

Fourier Shell Correlation (FSC) curve. The coloring represents resolution from 9.8Å (blue) to

12.4Å (red). **d** STA maps and FSC curves for two classes (I – blue; II – red) of gB[H527P]

identified on the surface of EVs using EMAN2. **e** STA maps of Class I and Class II gB[H527P]

molecules (inset) with each panel showing MDFF of the homology model of the VZV gB [20]

based on HSV-1 gB[H516P] [31] into the cryo-EM maps. Snapshots from Movie. S1 are shown,

the VZV gB domains are colored as for Fig. 1; DI (cyan), DII (green), DIII (yellow), DIV

(orange), DV (red) and linker regions (hot pink). **f** Molecular models from the MDFF of the

VZV gB homology model with boxed areas (i and ii) highlighting regions of structural

differences between the two models. Arrows (blue to red gradient) represent the proposed

movement between the class I and class II maps.

Figure 3. Targeted mutagenesis of gB ^{526}EHV^{528} disrupts gB/gH-gL dependent membrane

fusion and viral replication. a The predicted structures for the WT and mutant VZV gB DIII

central helices by AlphaFold2 [63,64] and the mutant structures aligned to WT gB using

Matchmaker [62] are shown; WT gB: yellow, alanine mutants: green, proline mutants: red. **b** Flow

cytometry analysis of CHO-DSP1 cells transfected with vectors expressing WT gB or gB proline

and alanine mutants, permeabilized, and immunostained with Cy5-conjugated mAb 93k;

nonspecific antigen control: grey, WT gB: blue, mutant gB: orange. **c** Frequency of gB-

expressing cells detected by mAb 93k and median florescence intensity (MFI) of mAb 93k

binding normalized to WT gB (mean ± SEM). Statistical differences were evaluated by one-way

ANOVA (**, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$). **d** Cell-cell fusion measured by the

stable reporter fusion assay (SRFA) using CHO-DSP1 cells transfected with vectors expressing

WT gB or gB mutants together with gH-gL expression vectors, mixed with MeWo-DSP2

reporter cells. Fusion efficiency was normalized to that mediated by gB[WT]/gH-gL (%). Data

from at least three independent experiments is shown in Box (25-75 percentile) and Whisker (10-

90 percentile) plots; horizontal bar is the median and the + indicates the mean. Statistical

differences were analyzed by one-way ANOVA (***, $p < 0.001$; ****, $p < 0.0001$). **e** Plaque

sizes in MeWo cells infected with pOka (WT), pOka-TKGFP-gB[E526P], pOka-TKGFP-

gB[E526A], pOka-TKGFP-gB[H527A], or pOka-TKGFP-gB[V528A] at 4 dpi; lethal mutant

viruses pOka-TKGFP-gB[H527P] and pOka-TKGFP-gB[V528P] are indicated by N/A (not

available). Comparison of 45 plaques per virus (mm^2) from three independent experiments is

shown by Box (25-75 percentile) and Whisker (10-90 percentile) plots; dots are outliers,

horizontal bar is the median and the + indicates the mean. Statistical differences between plaque

sizes of WT and mutant viruses were evaluated by one-way ANOVA (****, $p < 0.0001$). **f**

Immunohistochemistry staining of VZV plaques in MeWo cells, scale bar = 1 mm.

Figure 4. Disrupted gB fusion function did not interfere with VZV immediate early, early

or late protein synthesis or with virus particle assembly. a MeWo cells transfected with

pOka-TKGFP (WT), pOka-TKGFP-gB[E526P], pOka-TKGFP-gB[H527P], or pOka-TKGFP-

gB[V528P] BACs were imaged by confocal microscopy at 48 hrs post transfection. Immediate-

early protein (IE62, red), early protein, thymidine kinase (TK) expressed as a GFP fusion protein

(green) and late protein (capsid-ORF23, violet); nuclei stained with Hoechst 33342 (blue). Areas

highlighted by a small white box are magnified in the larger white boxes, white arrows indicate

cells at the early stage of infection identified by punctate expression of IE62 in nuclei with little

to no TK-GFP expression. Scale bar = 20 µm. **b** Diagram of the flow cytometry-based

transmission electron microscopy (FC-TEM) assay. MeWo cells were transfected with VZV

BACs expressing TK-GFP, harvested at 48 hrs posttransfection, and sorted for GFP-positive

cells; cells were fixed, embedded in 10% gelatin, infiltrated with resin, heavy metal staining and

mounted on the grid. **c** TEM images of MeWo cells transfected with pOka-TKGFP (WT), pOka-

TKGFP-gB[E526P], pOka-TKGFP-gB[H527P], or pOka-TKGFP-gB[V528P] BACs. Panels at

the right are magnifications of areas indicated by boxes (N: nuclei; C: cytoplasm). Boxes

outlined in red show representative viral capsids, with solid red arrowheads indicating viral DNA

containing C-capsid and open red arrowheads indicating empty A-, or scaffold containing B-

capsids. Boxes outlined in blue show representative viral particles, with solid blue arrowheads

indicating complete virus particles and open blue arrowheads indicating light particles. In the

magnified images, scale bar = 0.1 µm.

Fig 5. Canonical biosynthesis of gB was disrupted by V528P. a MeWo cells transiently

expressing WT gB and gB mutants were imaged by confocal microscopy at 24 hrs

posttransfection. Cells were stained with mouse anti-gB mAb SG2 (red), trans-Golgi network

marker, TGN46 (green), or early endosome antigen, EEA1 (green), and nuclei were stained with

Hoechst 33342 (blue). Representative images are from experiments performed at least in

duplicate. Areas in white boxes are shown at higher magnification in panels to the right; white

arrows indicate gB colocalization with TGN46 or EEA1 respectively. Scale bar = 15 µm. **b**

Intracellular trafficking of gB was assessed by glycosylation status. MeWo cells transfected with

vectors expressing WT gB, or gB mutants were lysed at 24 hrs posttransfection and

immunoprecipitated with anti-gB mAb SG2, left untreated (-), or treated with Endo H (H) or

PNGase F (F). Proteins were separated by SDS-PAGE gel (7.5%) under reducing conditions, and

gB was detected with rabbit anti-gB 746-867. The arrows and red vertical lines indicate the

glycosylated forms of full-length and cleaved gB respectively, the blue vertical lines indicate the

Endo H-resistant glycoforms of cleaved gB, and arrowheads indicate forms of gB deglycosylated

by both Endo H and PNGase F. Mass markers (kDa) are shown on the left.

917 **Table 1. Properties of the mutant protein and recombinant viruses bearing the proline and**

918 **alanine substitutions to ^{526}EHV^{528} in VZV gB DIII central helix.**

	Protein				Recombinant virus		
	Fusion activity *	Intracellular localization **	Protease cleavage **	EVs production	Viral gene expression	TEM morphology	Growth phenotype
E526P	3.8%	ER, trans-Golgi, early endosome	Yes	No	IE, E, L	Capsids; virus particles	Severely impaired; Small plaque
H527P	4.1%	ER, trans-Golgi, early endosome	Yes	Yes	IE, E, L	Capsids; virus particles	Lethal
V528P	3.8%	ER	No	No	IE, E, L	virus particles	Lethal
E526A	48.0%	ER, trans-Golgi, early endosome	Yes	ND	ND	ND	WT
H527A	89.6%	ER, trans-Golgi, early endosome	Yes	ND	ND	ND	WT
V528A	67.9%	ER, trans-Golgi, early endosome	Yes	ND	ND	ND	WT

919 *Fusion activity was meaused by SRFA and normalized to gB[WT]/gH-gL (%)

920 **WT gB localizes to ER, trans-Golgi network and early endosomes and undergoes cleavage.

921 EVs: extracellular vesicles

922 IE: immediately Early gene; E: early gene; L: late gene.

923 ND: experiment was not performed.

a

| | | | | | | | | | | |
|1|100|200|300|400|500|600|700|800|900|

DI DII DIII DIV DV

N' 115 DI DII DIII DIV DV C'
1 115 736 931

⁵¹¹SVEFAMLQFTYDHIQ**EHV**NEMLARISSSWCQLQNRERALWSGLFPI⁵⁵⁶

b

90°

V528 H527 E526

c

90°

H527 V528 E526

d

gB[H527P]   WT gB

50nm   50nm