Progressive neural engagement within the IFG-pMTG circuit as gesture and speech entropy and MI advances

Wanying Zhao[1*]; Zhouyi Li[1,4]; Xiang Li[1,2]; Yi Du[1,2,3*]

[1] State Key Laboratory of Cognitive Science and Mental Health, Institute of Psychology, Chinese Academy of Sciences, Beijing, China 100101

[2] Department of Psychology, University of Chinese Academy of Sciences, Beijing, China 100049

[3] Chinese Institute for Brain Research, Beijing, China 102206

[4] School of Psychology, Central China Normal University, Wuhan, China 430079

* Corresponding author:

Dr. Yi Du, Email: duyi@psych.ac.cn

16 Lincui Road, Chaoyang district, Beijing, China 100101

Dr. Wanying Zhao, Email: zhaowy@psych.ac.cn

Lincui Road, Chaoyang District, Beijing, China 100101

Abstract

Semantic representation emerges from distributed multisensory modalities, yet a

comprehensive understanding of the functional changing pattern within convergence zones or

hubs integrating multisensory semantic information remains elusive. In this study, employing

information-theoretic metrics, we quantified gesture and speech information, alongside their

interaction, utilizing entropy and mutual information (MI). Neural activities were assessed via

interruption effects induced by High-Definition transcranial direct current stimulation (HD-

tDCS). Additionally, chronometric double-pulse transcranial magnetic stimulation (TMS) and

high-temporal event-related potentials were utilized to decipher dynamic neural changes

resulting from various information contributors. Results showed gradual inhibition of both

inferior frontal gyrus (IFG) and posterior middle temporal gyrus (pMTG) as degree of gesture-

speech integration, indexed by MI, increased. Moreover, a time-sensitive and staged

progression of neural engagement was observed, evidenced by distinct correlations between

neural activity patterns and entropy measures of speech and gesture, as well as MI, across

early sensory and lexico-semantic processing stages. These findings illuminate the gradual

nature of neural activity during multisensory gesture-speech semantic processing, shaped by

dynamic gesture constraints and speech encoding, thereby offering insights into the neural

mechanisms underlying multisensory language processing.

Keywords: gesture-speech integration; pMTG-IFG circuit; information theory; multisensory;

semantic; dual-stage modal

Introduction

Semantic representation, distinguished by its cohesive conceptual nature, emerges from distributed modality-specific regions. Consensus acknowledges the presence of 'convergence zones' within the temporal and inferior parietal areas [1], or the 'semantic hub' located in the anterior temporal lobe[2], pivotal for integrating, converging, or distilling multimodal inputs. Contemporary theories frame the semantic processing as a dynamic sequence of neural states[3], shaped by systems that are finely tuned to the statistical regularities inherent in sensory inputs[4]. These regularities enable the brain to evaluate, weight, and integrate multisensory information, optimizing the reliability of individual sensory signals[5]. However, sensory inputs available to the brain are often incomplete and uncertain, necessitating adaptive neural adjustments to resolve these ambiguities[6]. In this context, neuronal activity is thought to be linked to the probability density of sensory information, with higher levels of uncertainty resulting in the engagement of a broader population of neurons, thereby reflecting the brain's adaptive capacity to handle diverse possible interpretations[7,8]. Although the role of 'convergence zones' and 'semantic hubs' in integrating multimodal inputs is well established, the precise functional patterns of neural activity in response to the distribution of unified multisensory information—along with the influence of unisensory signals—remain poorly understood.

To this end, we developed an analytic approach to directly probe the cortical engagement during multisensory gesture-speech semantic integration. Even though gestures convey information in a global-synthetic way, while speech conveys information in a linear segmented way, there exists a bidirectional semantic influence between the two modalities[9,10]. Gesture is regarded as 'part of language'[11] or functional equivalents of lexical units that alternate and integrate with speech into a 'single unification space' to convey a coherent meaning[12-14]. Empirical studies have investigated the semantic integration between gesture and speech by manipulating their semantic relationship[15-18] and revealed a mutual interaction between them[19-21] as reflected by the N400 latency and amplitude[14] as well as common neural underpinnings in the left inferior frontal gyrus (IFG) and posterior middle temporal gyrus (pMTG)[15,22,23].

Building on these insights, the present study quantified the amount of information from

both sources and their interaction adopting the information-theoretic complexity metrics of

entropy and *mutual information* (MI). Unisensory Entropy measures the disorder or

randomness of information and serves as an index of the uncertainty in modality-specific

representations of gesture or speech activated by an event[24]. MI assesses share information

between modalities[25], indicating multisensory convergence and acting as an index of gesture-

speech integration.

To investigate the neural mechanisms underlying gesture-speech integration, we

conducted three experiments to assess how neural activity correlates with distributed

multisensory integration, quantified using information-theoretic measures of MI. Additionally,

we examined the contributions of unisensory signals in this process, quantified through

unisensory entropy. **Experiment 1** employed high-definition transcranial direct current

stimulation (HD-tDCS) to administer Anodal, Cathodal and Sham stimulation to either the IFG

or the pMTG. HD-tDCS induces membrane depolarization with anodal stimulation and

membrane hyperpolarization with cathodal stimulation[26], thereby increasing or decreasing

cortical excitability in the targeted brain area, respectively. This experiment aimed to

determine whether the overall facilitation (Anodal-tDCS minus Sham-tDCS) and/or inhibitory

(Cathodal-tDCS minus Sham-tDCS) of these integration hubs is modulated by the degree of

gesture-speech integration, as measure by MI.

Given the differential involvement of the IFG and pMTG in gesture-speech integration,

shaped by top-down gesture predictions and bottom-up speech processing [23], **Experiment 2**

was designed to further assess whether the activity of these regions was associated with

relevant informational matrices. Specifically, we applied inhibitory chronometric double-pulse

transcranial magnetic stimulation (TMS) to specific temporal windows associated with

integration processes in these regions[23], assessing whether the inhibitory effects of TMS were

correlated with unisensory entropy or the multisensory convergence index (MI).

Experiment 3 complemented these investigations by focusing on the temporal dynamics

of neural responses during semantic processing, leveraging high-temporal event-related

potentials (ERPs). This experiment investigated how distinct information contributors

modulated specific ERP components associated with semantic processing. These

components included the early sensory effects as P1 and N1–P2[27,28], the N400 semantic

conflict effect[14,28,29], and the late positive component (LPC) reconstruction effect[30,31]. By

integrating these ERP findings with results from Experiments 1 and 2, Experiment 3 aimed to

provide a more comprehensive understanding of how gesture-speech integration is

modulated by neural dynamics.

Material and methods

Participants

Ninety-eight young Chinese participants signed written informed consent forms and took part

in the present study (Experiment 1: 29 females, 23 males, age = 20 ± 3.40 years; Experiment

2: 11 females, 13 males, age = 23 ± 4.88 years; Experiment 3: 12 females, 10 males, age =

21 ± 3.53 years). All of the participants were right-handed (Experiment 1: laterality quotient

(LQ)[32] = 88.71 ± 13.14; Experiment 2: LQ = 89.02 ± 13.25; Experiment 3: LQ = 88.49 ±

12.65), had normal or corrected-to-normal vision and were paid ¥100 per hour for their

participation. All experiments were approved by the Ethics Committee of the Institute of

Psychology, Chinese Academy of Sciences.

Stimuli

Twenty gestures (**Appendix Table 1**) with 20 semantically congruent speech signals taken

from previous study[23] were used. The stimuli set were recorded from two native Chinese

speakers (1 male, 1 female). To validate the stimuli, 30 participants were recruited to replicate

the multisensory index of semantic congruency effect, hypothesizing that reaction times for

semantically incongruent gesture-speech pairs would be significantly longer than those for

congruent pairs. The results confirmed this hypothesis, with a significantly ($t(29)$ = 7.16, $p <$

0.001) larger reaction time when participants were asked to judge the gender of the speaker if

gesture contained incongruent semantic information with speech (a 'cut' gesture paired with

speech word '喷 pen1 (spray)': mean = 554.51 ms, SE = 11.65) relative to when they were

semantically congruent (a 'cut' gesture paired with '剪 jian3 (cut)' word: mean = 533.90 ms,

SE = 12.02)[23].

Additionally, two separate pre-tests with 30 subjects in each (pre-test 1: 16 females, 14

males, age = 24 ± 4.37 years; pre-test 2: 15 females, 15 males, age = 22 ± 3.26 years) were

conducted to determine the comprehensive values of gesture and speech. Participants were

presented with segments of increasing duration, beginning at 40 ms, and were prompted to

provide a single verb to describe either the isolated gesture they observed (pre-test 1) or the

isolated speech they heard (pre-test 2). For each gesture or speech, the action verb

consistently provided by participants across four to six consecutive repetitions—with the

number of repetitions varied to mitigate learning effects—was considered the comprehensive

response for the gesture or speech. The initial instance duration was marked as the

discrimination point (DP) for gesture (mean = 183.78 ± 84.82ms) or the identification point (IP)

for speech (mean = 176.40 ± 66.21ms) (**Figure 1A top**).

To quantify information content, comprehensive responses for each item were converted

into Shannon's entropy (H) as a measure of information richness (**Figure 1A bottom**). With

no significant gender differences observed in both gesture ($t(20) = 0.21$, $p = 0.84$) and speech

($t(20) = 0.52$, $p = 0.61$), responses were aggregated across genders, resulting in 60 answers

per item (**Appendix Table 2**). Here, p(xi) and p(yi) represent the distribution of 60 answers for

a given gesture (**Appendix Table 2B**) and speech (**Appendix Table 2A**), respectively. High

entropy indicates diverse answers, reflecting broad representation, while low entropy

suggests focused lexical recognition for a specific item (**Figure 2B**). MI was used to measure

the overlap between gesture and speech information, calculated by subtracting the entropy of

the combined gesture-speech dataset (Entropy(gesture + speech)) from the sum of their

individual entropies (Entropy(gesture) + Entropy(speech)) (see **Appendix Table 2C**). For

specific gesture-speech combinations, equivalence between the combined entropy and the

sum of individual entropies (gesture or speech) indicates absence of overlap in response sets.

Conversely, significant overlap, denoted by a considerable number of shared responses

between gesture and speech datasets, leads to a noticeable discrepancy between combined

entropy and the sum of gesture and speech entropies. Elevated MI values thus signify

substantial overlap, indicative of a robust mutual interaction between gesture and speech.

Additionally, the number of responses provided for each gesture and speech, as well as

the total number of combined responses, were also recorded. The quantitative data for each

stimulus, including gesture entropy, speech entropy, joint entropy, MI, and the respective

counts, are presented in **Appendix Table 3.**

To determine whether entropy or MI values corresponds to distinct neural changes, the

current study first aggregated neural responses (including inhibition effects of tDCS and TMS

or ERP amplitudes) that shared identical entropy or MI values, prior to conducting

correlational analyses.

Experimental procedure

Given that gestures induce a semantic priming effect on concurrent speech[33], this study

utilized a semantic priming paradigm in which speech onset was aligned with the DP of each

gesture[23,33], the point at which the gesture transitions into a lexical form[34]. The gesture itself

began at the stroke phase, a critical moment when the gesture conveys its primary semantic

content[34].

An irrelevant factor of gender congruency (e.g., a man making a gesture combined with a

female voice) was created[22,23,35]. This involved aligning the gender of the voice with the

corresponding gender of the gesture in either a congruent (e.g., male voice paired with a male

gesture) or incongruent (e.g., male voice paired with a female gesture) manner. This

approach served as a direct control mechanism, facilitating the investigation of the automatic

and implicit semantic interplay between gesture and speech[35]. In light of previous findings

indicating a distinct TMS-disruption effect on the semantic congruency of gesture-speech

interactions[23], both semantically congruent and incongruent pairs were included in

Experiment 1 and Experiment 2. Experiment 3, conversely, exclusively utilized semantically

congruent pairs to elucidate ERP metrics indicative of nuanced semantic progression.

Gesture–speech pairs were presented randomly using Presentation software

(www.neurobs.com). Participants were asked to look at the screen but respond with both

hands as quickly and accurately as possible merely to the gender of the voice they heard.

The RT and the button being pressed were recorded. The experiment started with a fixation

cross presented on the center of the screen, which lasted for 0.5-1.5 sec.

Experiment 1: HD-tDCS protocol and data analysis

Participants were divided into two groups, with each group undergoing HD-tDCS stimulation

at different target sites (IFG or pMTG). Each participant completed three experimental

sessions, spaced one week apart, during which 480 gesture-speech pairs were presented

across various conditions. In each session, participants received one of three types of HD-

tDCS stimulation: Anodal, Cathodal, or Sham. The order of stimulation site and type was

counterbalanced using a Latin square design to control for potential order effects.

HD-tDCS protocol employed a constant current stimulator (The Starstim 8 system)

delivering stimulation at an intensity of 2000mA. A 4 * 1 ring-based electrode montage was

utilized, comprising a central electrode (stimulation) positioned directly over the target cortical

area and four return electrodes encircling it to provide focused stimulation. Building on a

meta-analysis of prior fMRI studies examining gesture-speech integration[22], we targeted

Montreal Neurological Institute (MNI) coordinates for the left IFG at (-62, 16, 22) and the

pMTG at (-50, -56, 10). In the stimulation protocol for HD-tDCS, the IFG was targeted using

electrode F7 as the optimal cortical projection site[36], with four return electrodes placed at AF7,

FC5, F9, and FT9. For the pMTG, TP7 was selected as the cortical projection site[36], with

return electrodes positioned at C5, P5, T9, and P9. The stimulation parameters included a 20-

minute duration with a 5-second fade-in and fade-out for both Anodal and Cathodal conditions.

The Sham condition involved a 5-second fade-in followed by only 30 seconds of stimulation,

then 19'20 minutes of no stimulation, and finally a 5-second fade-out (**Figure 1B**). Stimulation

was controlled using NIC software, with participants blinded to the stimulation conditions.

All incorrect responses (702 out of the total number of 24960, 2.81% of trials) were

excluded. To eliminate the influence of outliers, a 2SD trimmed mean for every participant in

each session was also calculated. To examine the relationship between the degree of

information and neural responses, we conducted Pearson correlation analyses using a

sample of 20 sets. Neural responses were quantified based on the effects of HD-tDCS (active

tDCS minus sham tDCS) on the semantic congruency effect, defined as the difference in

reaction times between semantic incongruent and congruent conditions (Rt(incongruent) -

Rt(congruent)). This effect served as an index of multisensory integration[35] within the left IFG

and pMTG. The variation in information was assessed using three information-theoretic

metrics. To account for potential confounds related to multiple candidate representations, we

conducted partial correlation analyses between the tDCS effects and gesture entropy, speech

entropy, and MI, controlling for the number of responses provided for each gesture and

speech, as well as the total number of combined responses. Given that HD-tDCS induces

overall disruption at the targeted brain regions, we hypothesized that the neural activity within

the left IFG and pMTG would be progressively affected by varying levels of multisensory

convergence, as indexed by MI. Moreover, we hypothesized that the modulation of neural

activity by MI would differ between the left IFG and pMTG, as reflected in the differential

modulation of response numbers in the partial correlations, highlighting their distinct roles in

semantic processing[37]. False discovery rate (FDR) correction was applied for multiple

comparisons.

Experiment 2: TMS protocol and data analysis

Experiment 2 involved 800 gesture-speech pairs, presented across 15 blocks over three days,

with one week between sessions. Stimulation was administered at three different sites (IFG,

pMTG, or Vertex). Within the time windows (TWs) spanning the gesture-speech integration

period, five TWs that exhibited selective disruption of integration were selected: TW1 (-120 to

-80 ms relative to the speech identification point), TW2 (-80 to -40 ms), TW3 (-40 to 0 ms),

TW6 (80 to 120 ms), and TW7 (120 to 160 ms)[23] (**Figure 1C**). The order of stimulation site

and TW was counterbalanced using a Latin square design.

At an intensity of 50% of the maximum stimulator output, double-pulse TMS was

delivered via a 70 mm figure-eight coil using a Magstim Rapid² stimulator (Magstim, UK).

High-resolution (1 × 1 × 0.6 mm) T1-weighted MRI scans were obtained using a Siemens 3T

Trio/Tim Scanner for image-guided TMS navigation. Frameless stereotaxic procedures

(BrainSight 2; Rogue Research) allowed real-time stimulation monitoring. To ensure precision,

individual anatomical images were manually registered by identifying the anterior and

posterior commissures. Subject-specific target regions were defined using trajectory markers

in the MNI coordinate system. Vertex was used as control.

All incorrect responses (922 out of the total number of 19200, 4.8% of trials) were

excluded. We focused our analysis on Pearson correlations of the TMS interruption effects

(active TMS minus vertex TMS) of the semantic congruency effect with the gesture entropy,

speech entropy or MI. To control for potential confounds, partial correlations were also

performed between the TMS effects and gesture entropy, speech entropy, and MI, controlling

for the number of responses for each gesture and speech, as well as the total number of

combined responses. By doing this, we can determine how the time-sensitive contribution of

the left IFG and pMTG to gesture–speech integration was affected by gesture and speech

information distribution. FDR correction was applied for multiple comparisons.

Experiment 3: Electroencephalogram (EEG) recording and data analysis

Experiment 3, comprising a total of 1760 gesture-speech pairs, was completed in a single-day

session. EEG were recorded from 48 Ag/AgCl electrodes mounted in a cap according to the

10-20 system[38], amplified with a PORTI-32/MREFA amplifier (TMS International B.V.,

Enschede, NL) and digitized online at 500 Hz (bandpass, 0.01-70 Hz). EEGLAB, a MATLAB

toolbox, was used to analyze the EEG data[39]. Vertical and horizontal eye movements were

measured with 4 electrodes placed above the left eyebrow, below the left orbital ridge and at

bilateral external canthus. All electrodes were referenced online to the left mastoid. Electrode

impedance was maintained below 5 KΩ. The average of the left and right mastoids was used

for re-referencing. A high-pass filter with a cutoff of 0.05 Hz and a low-pass filter with a cutoff

of 30 Hz were applied. Semi-automated artifact removal, including independent component

analysis (ICA) for identifying components of eye blinks and muscle activity, was performed

(**Figure 1D**). Participants with rejected trials exceeding 30% of their total were excluded from

further analysis.

All incorrect responses were excluded (147 out of 1760, 8.35% of trials). To eliminate the

influence of outliers, a 2 SD trimmed mean was calculated for every participant in each

condition. Data were epoched from the onset of speech and lasted for 1000 ms. To ensure a

clean baseline with no stimulus presented, a 200 ms pre-stimulus baseline correction was

applied before gesture onset.

To consolidate the data, we conducted both a traditional region-of-interest (ROI) analysis,

with ROIs defined based on a well-established work[40], and a cluster-based permutation

approach, which utilizes data-driven permutations to enhance robustness and address

multiple comparisons.

For the traditional ROI analysis, grand-average ERPs at electrode Cz were compared

between the higher (≥50%) and lower (<50%) halves for gesture entropy (**Figure 5A1**),

speech entropy (**Figure 5B1**), and MI (**Figure 5C1**). Consequently, four ERP components

were determined: the P1 effect observed within the time window of 0-100 ms[27,28], the N1-P2

effect observed between 150-250ms[27,28], the N400 within the interval of 250-450ms[14,28,29],

and the LPC spanning from 550-1000ms[30,31]. Additionally, seven regions-of-interest (ROIs)

were defined in order to locate the modulation effect on each ERP component: left anterior

(LA): F1, F3, F5, FC1, FC3, and FC5; left central (LC): C1, C3, C5, CP1, CP3, and CP5; left

posterior (LP): P1, P3, P5, PO3, PO5, and O1; right anterior (RA): F2, F4, F6, FC2, FC4, and

FC6; right central (RC): C2, C4, C6, CP2, CP4, and CP6; right posterior (RP): P2, P4, P6,

PO4, PO6, and O2; and midline electrodes (ML): Fz, FCz, Cz, Pz, Oz, and CPz[40].

Subsequently, cluster-based permutation tests[41] in Fieldtrip was further used to determine

the significant clusters of adjacent time points and electrodes of ERP amplitude between the

higher and lower halves of gesture entropy, speech entropy and MI, respectively. The

electrode-level type I error threshold was set to 0.025. Cluster-level statistic was estimated

through 5000 Monte Carlo simulations, where the cluster-level statistic is the sum of T-values

for each stimulus within a cluster. The cluster-level type I error threshold was set to 0.05.

Clusters with a p-value less than the critical alpha-level are considered to be conditionally

different.

Paired t-tests were conducted to compare the lower and upper halves of each information

model for the averaged amplitude within each ROI or cluster across the four ERP time

windows, separately. Pearson correlations were computed between each model value and

the averaged ERP amplitudes in each ROI or cluster. Additionally, partial correlations were

conducted, accounting for the number of responses for each respective metric. FDR

correction was applied for multiple comparisons.

Results

Experiment 1: Modulation of left pMTG and IFG engagement by gradual changes in

gesture-speech semantic information

In the IFG, one-way ANOVA examining the effects of three tDCS conditions (Anodal, Cathodal,

or Sham) on semantic congruency (RT (semantic incongruent) − RT (semantic congruent))

demonstrated a significant main effect of stimulation condition ($F_{(2, 75)}$ = 3.673, p = 0.030,

ηp2 = 0.089). Post hoc paired t-tests indicated a significantly reduced semantic congruency

effect between the Cathodal condition and the Sham condition ($t_{(26)}$ = -3.296, p = 0.003, 95%

CI = [-11.488, 4.896]) (**Figure 3A left**). Subsequent Pearson correlation analysis revealed that

the reduced semantic congruency effect was progressively associated with the MI, evidenced

by a significant correlation between the Cathodal-tDCS effect (Cathodal-tDCS minus Sham-

tDCS) and MI (r = -0.595, p = 0.007, 95% CI = [-0.995, -0.195]) (**Figure 3B**). Additionally, a

similar correlation was observed between the Cathodal-tDCS effect and the total response

number (r = -0.543, p = 0.016, 95% CI = [-0.961, -0.125]).

However, partial correlation analysis, controlling for the total response number, revealed

that the initially significant correlation between the Cathodal-tDCS effect and MI was no

longer significant (r = -0.303, p = 0.222, 95% CI = [-0.770, 0.164]). This suggests that the

observed relationship between Cathodal-tDCS and MI may be confounded by semantic

control difficulty, as reflected by the total number of responses. Specifically, the reduced

activity in the IFG under Cathodal-tDCS may be driven by variations in the difficulty of

semantic control rather than a direct modulation of MI.

In the pMTG, a one-way ANOVA assessing the effects of three tDCS conditions on

semantic congruency also revealed a significant main effect of stimulation condition ($F_{(2, 75)}$

= 3.250, p = 0.044, ηp2 = 0.080). Subsequent paired t-tests identified a significantly reduced

semantic congruency effect between the Cathodal condition and the Sham condition ($t_{(25)}$ = -

2.740, p = 0.011, 95% CI = [-11.915, 6.435]) (**Figure 3A right**). Moreover, a significant

correlation was observed between the Cathodal-tDCS effect and MI (r = -0.457, p = 0.049, 95%

CI = [-0.900, -0.014]) (**Figure 3B**).

Importantly, the reduced activity in the pMTG under Cathodal-tDCS was not influenced

by the total response number, as indicated by the non-significant correlation (r = -0.253, p =

0.295, 95% CI = [-0.735, 0.229]). This finding was further corroborated by the unchanged

significance in the partial correlation between Cathodal-tDCS and MI, when controlling for the

total response number (r = -0.472, p = 0.048, 95% CI = [-0.903, -0.041]).

RTs of congruent and incongruent trials of IFG and pMTG in each of the stimulation

conditions were shown in **Appendix Table 4A.**

Experiment 2: Time-sensitive modulation of left pMTG and IFG engagements by

gradual changes in gesture-speech semantic information

A 2 (TMS effect: active - Vertex) × 5 (TW) ANOVA on semantic congruency revealed a

significant interaction between TMS effect and TW ($F_{(3.589, 82.538)}$ = 3.273, p = 0.019, ηp2

= 0.125). Further t-tests identified a significant TMS effect over the pMTG in TW1 ($t_{(23)}$ = -

3.068, p = 0.005, 95% CI = [-6.838, 0.702]), TW2 ($t_{(23)}$ = -2.923, p = 0.008, 95% CI = [-6.490,

0.644]), and TW7 ($t_{(23)}$ = -2.005, p = 0.047, 95% CI = [-5.628, 1.618]). In contrast, a

significant TMS effect over the IFG was found in TW3 ($t_{(23)}$ = -2.335, p = 0.029, 95% CI = [-

5.928, 1.258]), and TW6 ($t_{(23)}$ = -4.839, p < 0.001, 95% CI = [-7.617, -2.061]) (**Figure 4A**).

Raw RTs of congruent and incongruent trials were shown in **Appendix Table 4B.**

Additionally, a significant negative correlation was found between the TMS effect (a larger

negative TMS effect signifies a greater disruption of the integration process) and speech

entropy when the pMTG was inhibited in TW2 (r = -0.792, p = 0.004, 95% CI = [-1.252, -

0.331]). Meanwhile, when the IFG activity was interrupted in TW6, a significant negative

correlation was found between the TMS effect and gesture entropy (r = -0.539, p = 0.014, 95%

CI = [-0.956, -0.122]), speech entropy (r = -0.664, p = 0.026, 95% CI = [-1.255, -0.073]), and

MI (r = -0.677, p = 0.001, 95% CI = [-1.054, -0.300]) (**Figure 4B**).

Notably, inhibition of pMTG activity in TW2 was not influenced by the number of speech

responses (r = -0.539, p = 0.087, 95% CI = [-1.145, 0.067]). However, the number of speech

responses did affect the modulation of speech entropy on the pMTG inhibition effect in TW2.

This was evidenced by the non-significant partial correlation between pMTG inhibition and

speech entropy when controlling for speech response number (r = -0.218, p = 0.545, 95% CI

= [-0.563, 0.127]).

In contrast, the interrupted IFG activity in TW6 appeared to be consistently influenced by

the confound of semantic control difficulty. This was reflected in the significant correlation with

both gesture response number (r = -0.480, p = 0.032, 95% CI = [-904, -0.056]), speech

response number (r = -0.729, p = 0.011, 95% CI = [-1.221, -0.237]), and total response

number (r = -0.591, p = 0.008, 95% CI = [-0.993, -0.189]). Additionally, partial correlation

analyses revealed non-significant relationship between interrupted IFG activity in TW6 and

gesture entropy (r = -0.369, p = 0.120, 95% CI = [-0.810, -0.072]), speech entropy (r = -0.455,

p = 0.187, 95% CI = [-1.072, 0.162]), and MI (r = -0.410, p = 0.091, 95% CI = [-0.856, -0.036])

when controlling for response numbers.

Experiment 3: Temporal modulation of P1, N1-P2, N400 and LPC components by

gradual changes in gesture-speech semantic information

Topographical maps illustrating amplitude differences between the lower and higher halves of

speech entropy demonstrate a central-posterior P1 amplitude (0-100 ms, **Figure 5B**). Aligning

with prior findings[27], the paired t-tests demonstrated a significantly larger P1 amplitude within

the ML ROI (t(22) = 2.510, p = 0.020, 95% confidence interval (CI) = [1.66, 3.36]) when

contrasting stimuli with higher 50% speech entropy against those with lower 50% speech

entropy (**Figure 5D1 left**). Subsequent correlation analyses unveiled a significant increase in

the P1 amplitude with the rise in speech entropy within the ML ROI (r = 0.609, p = 0.047, 95%

CI = [0.039, 1.179], **Figure 5D1 right**). Furthermore, a cluster of neighboring time-electrode

samples exhibited a significant contrast between the lower 50% and higher 50% of speech

entropy, revealing a P1 effect spanning 16 to 78 ms at specific electrodes (FC2, FCz, C1, C2,

Cz, and CPz, **Figure 5D2 middle**) (t(22) = 2.754, p = 0.004, 95% confidence interval (CI) =

[1.65, 3.86], **Figure 5D2 left**), with a significant correlation with speech entropy (r = 0.636, p =

0.035, 95% CI = [0.081, 1.191], **Figure 5D2 right**).

Additionally, topographical maps comparing the lower 50% and higher 50% gesture

entropy revealed a frontal N1-P2 amplitude (150-250 ms, **Figure 5A**). In accordance with

previous findings on bilateral frontal N1-P2 amplitude[27], paired t-tests displayed a significantly

larger amplitude for stimuli with lower 50% gesture entropy than with higher 50% entropy in

both ROIs of LA (t(22) = 2.820, p = 0.011, 95% CI = [2.21, 3.43]) and RA (t(22) = 2.223, p =

0.038, 95% CI = [1.56, 2.89]) (**Figure 5E1 left**). Moreover, a negative correlation was found

between N1-P2 amplitude and gesture entropy in both ROIs of LA (r = -0.465, p = 0.039, 95%

CI = [-0.87, -0.06]) and RA (r = -0.465, p = 0.039, 95% CI = [-0.88, -0.05]) (**Figure 5E1 right**).

Additionally, through a cluster-permutation test, the N1-P2 effect was identified between 184

to 202 ms at electrodes FC4, FC6, C2, C4, C6, and CP4 (**Figure 5E2 middle**) (t(22) = 2.638,

p = 0.015, 95% CI = [1.79, 3.48], (**Figure 5E2 left**)), exhibiting a significant correlation with

gesture entropy (r = -0.485, p = 0.030, 95% CI = [-0.91, -0.06], **Figure 5E2 right**).

Furthermore, in line with prior research[42], a left-frontal N400 amplitude (250-450 ms) was

discerned from topographical maps of gesture entropy (**Figure 5A**). Specifically, stimuli with

lower 50% values of gesture entropy elicited a larger N400 amplitude in the LA ROI compared

to those with higher 50% values (t(22) = 2.455, p = 0.023, 95% CI = [1.95, 2.96], **Figure 5F1**

left). Concurrently, a negative correlation was noted between the N400 amplitude and gesture

entropy (r = -0.480, p = 0.032, 95% CI = [-0.94, -0.03], **Figure 5F1 right**) within the LA ROI.

The identified clusters showing the N400 effect for gesture entropy (282 – 318 ms at

electrodes FC1, FCz, C1, and Cz, **Figure 5F2 middle**) (t(22) = 2.828, p = 0.010, 95% CI =

[2.02, 3.64], **Figure 5F2 left**) also exhibited significant correlation between the N400

amplitude and gesture entropy (r = -0.445, p = 0.049, 95% CI = [-0.88, -0.01], **Figure 5F2**

right).

Similarly, a left-frontal N400 amplitude (250-450 ms)[42] was discerned from topographical

maps for MI (**Figure 5C)**. A larger N400 amplitude in the LA ROI was observed for stimuli with

lower 50% values of MI compared to those with higher 50% values (t(22) = 3.00, p = 0.007,

95% CI = [2.54, 3.46], **Figure 5G1 left**). This was accompanied by a significant negative

correlation between N400 amplitude and MI (r = -0.504, p = 0.028, 95% CI = [-0.97, -0.04],

Figure 5G1 right) within the LA ROI. The N400 effect for MI, observed in the 294–306 ms

window at electrodes F1, F3, Fz, FC1, FC3, FCz, and C1 (**Figure 5G2 middle**) (t(22) = 2.461,

p = 0.023, 95% CI = [1.62, 3.30], **Figure 5G2 left**), also showed a significant negative

correlation with MI (r = -0.569, p = 0.011, 95% CI = [-0.98, -0.16], **Figure 5G2 right**).

Finally, consistent with previous findings[30], an anterior LPC effect (550-1000 ms) was

observed in topographical maps comparing stimuli with lower and higher 50% speech entropy

(**Figure 5B**). The reduced LPC amplitude was evident in the paired t-tests conducted in ROIs

of LA (t(22) = 2.614, p = 0.016, 95% CI = [1.88, 3.35]); LC (t(22) = 2.592, p = 0.017, 95% CI =

[1.83, 3.35]); RA (t(22) = 2.520, p = 0.020, 95% CI = [1.84, 3.24]); and ML (t(22) = 2.267, p =

0.034, 95% CI = [1.44, 3.10]) (**Figure 5H1 left**). Simultaneously, a marked negative

correlation with speech entropy was evidenced in ROIs of LA (r = -0.836, p =   0.001, 95% CI

= [-1.26, -0.42]); LC (r = -0.762, p = 0.006, 95% CI = [-1.23, -0.30]); RA (r = -0.774, p = 0.005,

95% CI = [-1.23, -0.32]) and ML (r = -0.730, p = 0.011, 95% CI = [-1.22, -0.24]) (**Figure 5H1**

right). Additionally, a cluster with the LPC effect (644 - 688 ms at electrodes Cz, CPz, P1, and

Pz, **Figure 5H2 middle**) (t(22) = 2.754, p = 0.012, 95% CI = [1.50, 4.01], **Figure 5H2 left**)

displayed a significant correlation with speech entropy (r = -0.699, p = 0.017, 95% CI = [-1.24,

-0.16], **Figure 5H2 right**).

To clarify potential confounds of semantic control difficulty, partial correlation analyses

were conducted to examine the relationship between the elicited ERP components and the

relevant information matrices, controlling for response numbers. Results consistently

indicated modulation by response numbers in the relationship of ERP components with the

information matrix, as evidenced by the non-significant partial correlations between the P1

amplitude (P1 component over ML: r = -0.574, p = 0.082, 95% CI = [-1.141, -0.007]) and the

P1 cluster (r = -0.503, p = 0.138, 95% CI = [-1.102, 0.096]) with speech entropy; the N1-P2

amplitude (N1-P2 component over LA: r = -0.080, p = 0.746, 95% CI = [-0.554, 0.394]) and

N1-P2 cluster (r = -0.179, p = 0.464, 95% CI = [-0.647, 0.289]) with gesture entropy; the N400

amplitude (N400 component over LA: r = 0.264, p = 0.247, 95% CI = [-0.195,0.723]) and

N400 cluster (r = 0.394, p = 0.095, 95% CI = [-0.043, 0.831]) with gesture entropy; the N400

amplitude (N400 component over LA: r = -0.134, p = 0.595, 95% CI = [-0.620, 0.352]) and

N400 cluster (r = -0.034, p = 0.894, 95% CI = [-0.524,0.456]) with MI; and the LPC amplitude

(LPC component over LA: r = -0.428, p = 0.217, 95% CI = [-1.054, 0.198]) and LPC cluster (r

= -0.202, p = 0.575, 95% CI = [-0.881, 0.477]) with speech entropy.

Discussion

Through mathematical quantification of gesture and speech information using entropy and

mutual information (MI), we examined the functional pattern and dynamic neural structure

underlying multisensory semantic integration. Our results, for the first time, revealed that the

inhibition effect of cathodal-tDCS on the pMTG and IFG correlated with the degree of gesture-

speech multisensory convergence, as indexed by MI (**Experiment 1**). Moreover, the gradual

neural engagement was found to be time-sensitive and staged, as evidenced by the

selectively interrupted time windows (**Experiment 2**) and the distinct correlated ERP

components (**Experiment 3**), which were modulated by different information contributors,

including unisensory entropy or multisensory MI. These findings significantly expand our

understanding of the cortical foundations of statistically regularized multisensory semantic

information.

It is widely acknowledged that a single, amodal system mediates the interactions among

perceptual representations of different modalities[5,43,44]. Moreover, observations have

suggested that semantic dementia patients experience increasing overregularization of their

conceptual knowledge due to the progressive deterioration of this amodal system[45].

Consistent with this, the present study provides robust evidence, through the application of

HD-tDCS and TMS, that the integration hubs for gesture and speech—the pMTG and IFG—

operate in an incremental manner. This is supported by the progressive inhibition effect

observed in these brain areas as the entropy and mutual information of gesture and speech

advances.

Moreover, by dividing the potential integration period into eight time windows (TW)

relative to the speech identification point (IP) and administering inhibitory double-pulse TMS

across each TW, the current study attributed the gradual TMS-selective regional inhibition to

distinct information sources. In TW2 of gesture-speech integration, which precedes the

speech identification point[23] and represents a pre-lexical stage, the suppression effect

observed in the pMTG was correlated with speech entropy. Conversely, during TW6, which

follows the speech identification point[23] and represents a post-lexical stage, the IFG

interruption effect was influenced by both gesture entropy, speech entropy, and their MI. A

dual-stage pMTG-IFG-pMTG neurocircuit loop during gesture-speech integration has been

proposed previous[23]. As an extension, the present study unveils a staged accumulation of

engagement within the neurocircuit linking the transmodal regions of pMTG and IFG, arising

from distinct contributors of information.

Furthermore, we disentangled the sub-processes of integration with high-temporal ERPs,

when representations of gesture and speech were variously presented. Early P1-N1 and P2

sensory effects linked to perception and attentional processes[28,46] was comprehended as a

reflection of the early audiovisual gesture-speech integration in the sensory-perceptual

processing chain[47]. Note that a semantic priming paradigm was adopted here to create a top-

down prediction of gesture over speech. The observed positive correlation of the P1 effect

with speech entropy and the negative correlation of the N1-P2 effect with gesture entropy

suggest that the early interaction of gesture-speech information was modulated by both top-

down gesture prediction and bottom-up speech processing. Additionally, the lexico-semantic

effect of the N400 and the LPC were differentially mediated by top-down gesture prediction,

bottom-up speech encoding and their interaction: the N400 was negatively correlated with

both the gesture entropy and MI, but the LPC was negatively correlated only with the speech

entropy.

The varying contributions of unisensory gesture-speech information and the convergence

of multisensory inputs, as reflected in the correlation between distinct ERP components and

TMS time windows (TMS TWs), are consistent with recent models suggesting that

multisensory processing involves parallel detection of modality-specific information and

hierarchical integration across multiple neural levels[4,48]. These processes are further

characterized by coordination across multiple temporal scales[49]. Building on this, the present

study offers additional evidence that the multi-level nature of gesture-speech processing is

statistically structured, as measured by information matrix of unisensory entropy and

multisensory convergence index of MI, the input of either source would activate a distributed

representation, resulting in progressively functioning neural responses.

Given that control processes are intrinsically integrated with semantic processing[50], a

distributed semantic representation enables dynamic modulation of access to and

manipulation of meaningful information, thereby facilitating flexible control over the diverse

possibilities inherent in a concept. Accordingly, an increased number of candidate responses

amplifies the control demands necessary to resolve competing semantic representations. This

effect was observed in the present study, where the association of the information matrix with

the tDCS effect in IFG, the inhibition of pMTG activity in TW2, disruption of IFG activity in

TW6, and modulation of four distinct ERP components collectively demonstrated that

response quantity modulates neural activity. These results underscore the intricate interplay

between the difficulty of semantic representation and the control pressures that shape the

resulting neural responses.

The IFG and pMTG, central components of the semantic control network, have been

extensively implicated in previous research [50-52]. While the role of the IFG in managing both

unisensory information and multisensory convergence remains consistent, as evidenced by

the confounding difficulty results across Experiments 1 and 2, the current study highlights a

more context-dependent function for the pMTG. Specifically, although the pMTG is well-

established in the processing of distributed speech information, the multisensory convergence,

indexed by MI, did not evoke the same control-related modulation in pMTG activity. These

findings suggest that, while the pMTG is critical to semantic processing, its engagement in

control processes is likely modulated by the specific nature of the sensory inputs involved.

Considering the close alignment of the ERP components with the TWs of TMS effect, it is

reasonable to speculate the ERP components with the cortical involvements (**Figure 6**).

Consequently, referencing the recurrent neurocircuit connecting the left IFG and pMTG for

semantic unification[53], we extended the previously proposed two-stage gesture-speech

integration circuit[23] into sequential steps. First, bottom-up speech processing mapping

acoustic signal to its lexical representation was performed to the pMTG. The larger speech

entropy was, the greater effort was made during the matching of the acoustic input with its

stored lexical representation, thus leading to a larger involvement of the pMTG at pre-lexical

stage (TW2) and a larger P1 effect (**Figure 6①**). Second, the gesture representation was

activated in the pMTG and further exerted a top-down modulation over the phonological

processing of speech [54]. The higher the certainty of gesture is, a larger modulation of gesture

would be made upon speech, as indexed by a smaller gesture entropy with an enhanced N1-

P2 amplitude (**Figure 6②**). Third, information was relayed from the pMTG to the IFG for

sustained activation, during which a semantic constraint from gesture has been made on the

semantic retrieval of speech. Greater TMS inhibitory effect over the IFG at post-lexical stage

(TW6) accompanying with a reduced N400 amplitude were found with the increase of gesture

entropy, when the representation of gesture was wildly distributed and the constrain over the

following speech was weak **(Figure 6③)**. Fourth, the activated speech representation was

compared with that of the gesture in the IFG. At this stage, the larger, overlapped neural

populations activated by gesture and speech as indexed by a larger MI is, a greater TMS

disruption effect of the IFG and a reduced N400 amplitude indexing easier integration and

less semantic conflict were observed (**Figure 6④**). Last, the activated speech representation

would disambiguate and reanalyze the semantic information and further unify into a coherent

comprehension in the pMTG[12,37]. As speech entropy increases, indicating greater uncertainty

in the information provided by speech, more cognitive effort is directed towards selecting the

targeted semantic representation. This leads to enhanced involvement of the IFG and a

corresponding reduction in LPC amplitude (**Figure 6⑤**).

Note that the sequential cortical involvement and ERP components discussed above are

derived from a deliberate alignment of speech onset with gesture DP, creating an artificial

priming effect with gesture semantically preceding speech. Caution is advised when

generalizing these findings to the spontaneous gesture-speech relationships, although

gestures naturally precede speech[34]. Furthermore, MI quantifies overlap in gesture-speech

integration, primarily when gestures convey redundant meaning. Consequently, the

conclusions drawn in this study are constrained to contexts in which gestures serve to

reinforce the meaning of the speech. Future research should aim to explore the neural

responses in cases where gestures convey supplementary, rather than redundant, semantic

information.

Limitations exist. ERP components and cortical engagements were linked through

intermediary variables of entropy and MI. Dissociations were observed between ERP

components and cortical engagement. Importantly, there is no direct evidence of the brain

structures underpinning the corresponding ERPs, necessitating clarification in future studies.

Additionally, not all influenced TWs exhibited significant associations with entropy and MI.

While HD-tDCS and TMS may impact functionally and anatomically connected brain

regions[55,56], whether the absence of influence in certain TWs can be attributed to

compensation by other connected brain areas, such as angular gyrus[57] or anterior temporal

lobe[58], warrants further investigation. Therefore, caution is needed when interpreting the

causal relationship between inhibition effects of brain stimulation and information-theoretic

metrics (entropy and MI). Finally, the current study incorporated a restricted set of entropy

and MI measures. The generalizability of the findings should be assessed in future studies

using a more extensive range of matrices.

In summary, utilizing information-theoretic complexity metrics such as entropy and mutual

information (MI), our study demonstrates that multisensory semantic processing, involving

gesture and speech, gives rise to dynamically evolving representations through the interplay

between gesture-primed prediction and speech presentation. This process correlates with the

progressive engagement of the pMTG-IFG-pMTG circuit and various ERP components.

These findings significantly advancing our understanding of the neural mechanisms

underlying multisensory semantic integration.

Acknowledgments

This research was supported by grants from the STI 2030—Major Projects 2021ZD0201500,

the National Natural Science Foundation of China (31822024, 31800964), the Scientific

Foundation of Institute of Psychology, Chinese Academy of Sciences (E2CX3625CX), and the

Strategic Priority Research Program of Chinese Academy of Sciences (XDB32010300).

Author contributions

Conceptualization, W.Y.Z. and Y.D.; Investigation, W.Y.Z. and Z.Y.L.; Formal Analysis, W.Y.Z.

and Z.Y.L.; Methodology, W.Y.Z. and Z.Y.L.; Validation, Z.Y.L. and X.L.; Visualization, W.Y.Z.

and Z.Y.L. and X.L.; Funding Acquisition, W.Y.Z. and Y.D.; Supervision, Y.D.; Project

administration, Y.D.; Writing – Original Draft, W.Y.Z.; Writing – Review & Editing, W.Y.Z., Z.Y.L.,

X.L., and Y.D.

Competing interests

The authors declare no competing interests.

References

References

1.   Damasio, H., Grabowski, T.J., Tranel, D., Hichwa, R.D., and Damasio, A.R. (1996). A

neural basis for lexical retrieval. Nature *380*, 499-505. DOI 10.1038/380499a0.

2.   Patterson, K., Nestor, P.J., and Rogers, T.T. (2007). Where do you know what you

know? The representation of semantic knowledge in the human brain. Nature

Reviews Neuroscience *8*, 976-987. 10.1038/nrn2277.

3.   Brennan, J.R., Stabler, E.P., Van Wagenen, S.E., Luh, W.M., and Hale, J.T. (2016).

Abstract linguistic structure correlates with temporal activity during naturalistic

comprehension. Brain and Language *157*, 81-94. 10.1016/j.bandl.2016.04.008.

4.   Benetti, S., Ferrari, A., and Pavani, F. (2023). Multimodal processing in face-to-face

interactions: A bridging link between psycholinguistics and sensory neuroscience.

Front Hum Neurosci *17*, 1108354. 10.3389/fnhum.2023.1108354.

5.   Noppeney, U. (2021). Perceptual Inference, Learning, and Attention in a Multisensory

World. Annual Review of Neuroscience, Vol 44, 2021 *44*, 449-473. 10.1146/annurev-

neuro-100120-085519.

6.   Ma, W.J., and Jazayeri, M. (2014). Neural coding of uncertainty and probability. Annu

Rev Neurosci *37*, 205-220. 10.1146/annurev-neuro-071013-014017.

7.   Fischer, B.J., and Pena, J.L. (2011). Owl's behavior and neural representation

predicted by Bayesian inference. Nat Neurosci *14*, 1061-1066. 10.1038/nn.2872.

8.   Ganguli, D., and Simoncelli, E.P. (2014). Efficient sensory encoding and Bayesian

inference with heterogeneous neural populations. Neural Comput *26*, 2103-2134.

10.1162/NECO_a_00638.

9.   Hostetter, A., and Mainela-Arnold, E. (2015). Gestures occur with spatial and Motoric

knowledge: It's more than just coincidence. Perspectives on Language Learning and

Education *22*, 42-49. doi:10.1044/lle22.2.42.

10.   McNeill, D. (2005). Gesture and though (University of Chicago Press).

10.7208/chicago/9780226514642.001.0001.

11.   Kendon, A. (1997). Gesture. Annu Rev Anthropol *26*, 109-128.

10.1146/annurev.anthro.26.1.109.

12.   Hagoort, P. (2005). On broca, brain, and binding: a new framework. Trends in

Cognitive Sciences *9*, 416-423. 10.1016/j.tics.2005.07.004.

13.   Hagoort, P., Hald, L., Bastiaansen, M., and Petersson, K.M. (2004). Integration of

word meaning and world knowledge in language comprehension. Science *304*, 438-

441. 10.1126/science.1095455.

14.   Ozyurek, A., Willems, R.M., Kita, S., and Hagoort, P. (2007). On-line integration of

semantic information from speech and gesture: Insights from event-related brain

potentials. J Cognitive Neurosci *19*, 605-616. 10.1162/jocn.2007.19.4.605.

15.   Willems, R.M., Ozyurek, A., and Hagoort, P. (2009). Differential roles for left inferior

frontal and superior temporal cortex in multimodal integration of action and language.

Neuroimage *47*, 1992-2004. 10.1016/j.neuroimage.2009.05.066.

16.   Drijvers, L., Jensen, O., and Spaak, E. (2021). Rapid invisible frequency tagging

reveals nonlinear integration of auditory and visual information. Human Brain

Mapping *42*, 1138-1152. 10.1002/hbm.25282.

17.   Drijvers, L., and Ozyurek, A. (2018). Native language status of the listener modulates

637   the neural integration of speech and iconic gestures in clear and adverse listening

638   conditions. Brain and Language *177*, 7-17. 10.1016/j.bandl.2018.01.003.

639   18.   Drijvers, L., van der Plas, M., Ozyurek, A., and Jensen, O. (2019). Native and non-

640   native listeners show similar yet distinct oscillatory dynamics when using gestures to

641   access speech in noise. Neuroimage *194*, 55-67. 10.1016/j.neuroimage.2019.03.032.

642   19.   Holle, H., and Gunter, T.C. (2007). The role of iconic gestures in speech

643   disambiguation: ERP evidence. J Cognitive Neurosci *19*, 1175-1192.

644   10.1162/jocn.2007.19.7.1175.

645   20.   Kita, S., and Ozyurek, A. (2003). What does cross-linguistic variation in semantic

646   coordination of speech and gesture reveal?: Evidence for an interface representation

647   of spatial thinking and speaking. J Mem Lang *48*, 16-32. 10.1016/S0749-

648   596x(02)00505-3.

649   21.   Bernardis, P., and Gentilucci, M. (2006). Speech and gesture share the same

650   communication system. Neuropsychologia *44*, 178-190.

651   10.1016/j.neuropsychologia.2005.05.007.

652   22.   Zhao, W.Y., Riggs, K., Schindler, I., and Holle, H. (2018). Transcranial magnetic

653   stimulation over left inferior frontal and posterior temporal cortex disrupts gesture-

654   speech integration. Journal of Neuroscience *38*, 1891-1900. 10.1523/Jneurosci.1748-

655   17.2017.

656   23.   Zhao, W., Li, Y., and Du, Y. (2021). TMS reveals dynamic interaction between inferior

657   frontal gyrus and posterior middle temporal gyrus in gesture-speech semantic

658   integration. The Journal of Neuroscience, 10356-10364. 10.1523/jneurosci.1355-

21.2021.

24.   Shannon, C.E. (1948). A mathematical theory of communication. Bell Syst Tech J *27*,

379-423. 10.1002/j.1538-7305.1948.tb01338.x.

25.   Tremblay, P., Deschamps, I., Baroni, M., and Hasson, U. (2016). Neural sensitivity to

syllable frequency and mutual information in speech perception and production.

Neuroimage *136*, 106-121. 10.1016/j.neuroimage.2016.05.018.

26.   Bikson, M., Inoue, M., Akiyama, H., Deans, J.K., Fox, J.E., Miyakawa, H., and

Jefferys, J.G.R. (2004). Effects of uniform extracellular DC electric fields on

excitability in rat hippocampal slices. J Physiol-London *557*, 175-190.

10.1113/jphysiol.2003.055772.

27.   Federmeier, K.D., Mai, H., and Kutas, M. (2005). Both sides get the point:

hemispheric sensitivities to sentential constraint. Memory & Cognition *33*, 871-886.

10.3758/bf03193082.

28.   Kelly, S.D., Kravitz, C., and Hopkins, M. (2004). Neural correlates of bimodal speech

and gesture comprehension. Brain and Language *89*, 253-260. 10.1016/s0093-

934x(03)00335-3.

29.   Wu, Y.C., and Coulson, S. (2005). Meaningful gestures: Electrophysiological indices

of iconic gesture comprehension. Psychophysiology *42*, 654-667. 10.1111/j.1469-

8986.2005.00356.x.

30.   Fritz, I., Kita, S., Littlemore, J., and Krott, A. (2021). Multimodal language processing:

How preceding discourse constrains gesture interpretation and affects gesture

integration when gestures do not synchronise with semantic affiliates. J Mem Lang

117, 104191. 10.1016/j.jml.2020.104191.

31.   Gunter, T.C., and Weinbrenner, J.E.D. (2017). When to take a gesture seriously: On

how we use and prioritize communicative cues. J Cognitive Neurosci *29*, 1355-1367.

10.1162/jocn_a_01125.

32.   Oldfield, R.C. (1971). The assessment and analysis of handedness: the Edinburgh

inventory. Neuropsychologia *9*, 97-113. 10.1016/0028-3932(71)90067-4.

33.   Zhao, W. (2023). TMS reveals a two-stage priming circuit of gesture-speech

integration. Front Psychol *14*, 1156087. 10.3389/fpsyg.2023.1156087.

34.   McNeill, D. (1992). Hand and mind : what gestures reveal about thought (University of

Chicago Press). 10.2307/1576015.

35.   Kelly, S.D., Creigh, P., and Bartolotti, J. (2010). Integrating speech and iconic

gestures in a Stroop-like task: Evidence for automatic processing. Journal of

Cognitive Neuroscience

22, 683-694. 10.1162/jocn.2009.21254.

36.   Koessler, L., Maillard, L., Benhadid, A., Vignal, J.P., Felblinger, J., Vespignani, H.,

and Braun, M. (2009). Automated cortical projection of EEG sensors: Anatomical

correlation via the international 10-10 system. Neuroimage *46*, 64-72.

10.1016/j.neuroimage.2009.02.006.

37.   Tesink, C.M.J.Y., Petersson, K.M., van Berkum, J.J.A., van den Brink, D., Buitelaar,

J.K., and Hagoort, P. (2009). Unification of speaker and meaning in language

comprehension: An fMRI study. J Cognitive Neurosci *21*, 2085-2099.

10.1162/jocn.2008.21161.

38.   Nuwer, M.R., Comi, G., Emerson, R., Fuglsang-Frederiksen, A., Guerit, J.M., Hinrichs,

704   H., Ikeda, A., Luccas, F.J., and Rappelsberger, P. (1999). IFCN standards for digital

705   recording of clinical EEG. The International Federation of Clinical Neurophysiology.

706   Electroencephalogr Clin Neurophysiol Suppl *52*, 11-14. 10.1016/S0013-

707   4694(97)00106-5.

708   39.   Delorme, A., and Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of

709   single-trial EEG dynamics including independent component analysis. J Neurosci

710   Methods *134*, 9-21. 10.1016/j.jneumeth.2003.10.009.

711   40.   Habets, B., Kita, S., Shao, Z.S., Ozyurek, A., and Hagoort, P. (2011). The Role of

712   Synchrony and Ambiguity in Speech-Gesture Integration during Comprehension. J

713   Cognitive Neurosci *23*, 1845-1854. 10.1162/jocn.2010.21462.

714   41.   Oostenveld, R., Fries, P., Maris, E., and Schoffelen, J.-M. (2011). FieldTrip: Open

715   Source Software for Advanced Analysis of MEG, EEG, and Invasive

716   Electrophysiological Data. Computational Intelligence and Neuroscience *2011*,

717   156869. 10.1155/2011/156869.

718   42.   Kutas, M., and Federmeier, K.D. (2011). Thirty Years and Counting: Finding Meaning

719   in the N400 Component of the Event-Related Brain Potential (ERP). Annual Review

720   of Psychology, Vol 62 *62*, 621-647. 10.1146/annurev.psych.093008.131123.

721   43.   Rogers, T.T., Ralph, M.A.L., Garrard, P., Bozeat, S., McClelland, J.L., Hodges, J.R.,

722   and Patterson, K. (2004). Structure and deterioration of semantic memory: A

723   neuropsychological and computational investigation. Psychological Review *111*, 205-

724   235. 10.1037/0033-295x.111.1.205.

44.   Ralph, M.A.L., Jefferies, E., Patterson, K., and Rogers, T.T. (2017). The neural and

computational bases of semantic cognition. Nature Reviews Neuroscience *18*, 42-55.

10.1038/nrn.2016.150.

45.   Rogers, T.T., Hodges, J.R., Ralph, M.A.L., and Patterson, K. (2003). Object

recognition under semantic impairment: The effects of conceptual regularities on

perceptual decisions. Lang Cognitive Proc *18*, 625-662.

10.1080/01690960344000053.

46.   Fadiga, L., Craighero, L., and Olivier, E. (2005). Human motor cortex excitability

during the perception of others' action. Current Opinion in Neurobiology *15*, 213-218.

10.1016/j.conb.2005.03.013.

47.   Giard, M.H., and Peronnet, F. (1999). Auditory-visual integration during multimodal

object recognition in humans: A behavioral and electrophysiological study. J

Cognitive Neurosci *11*, 473-490. 10.1162/089892999563544.

48.   Meijer, G.T., Mertens, P.E.C., Pennartz, C.M.A., Olcese, U., and Lansink, C.S. (2019).

The circuit architecture of cortical multisensory processing: Distinct functions jointly

operating within a common anatomical network. Prog Neurobiol *174*, 1-15.

10.1016/j.pneurobio.2019.01.004.

49.   Senkowski, D., and Engel, A.K. (2024). Multi-timescale neural dynamics for

multisensory integration. Nat Rev Neurosci *25*, 625-642. 10.1038/s41583-024-00845-

7.

50.   Jackson, R.L. (2021). The neural correlates of semantic control revisited. Neuroimage

224, 117444. 10.1016/j.neuroimage.2020.117444.

51.   Jefferies, E. (2013). The neural basis of semantic cognition: converging evidence

from neuropsychology, neuroimaging and TMS. Cortex *49*, 611-625.

10.1016/j.cortex.2012.10.008.

52.   Noonan, K.A., Jefferies, E., Visser, M., and Lambon Ralph, M.A. (2013). Going

beyond inferior prefrontal involvement in semantic control: evidence for the additional

contribution of dorsal angular gyrus and posterior middle temporal cortex. J Cogn

Neurosci *25*, 1824-1850. 10.1162/jocn_a_00442.

53.   Hagoort, P. (2013). MUC (Memory, Unification, Control) and beyond. Frontiers in

Psychology *4*, 416. 10.3389/fpsyg.2013.00416.

54.   Bizley, J.K., Maddox, R.K., and Lee, A.K.C. (2016). Defining auditory-visual objects:

Behavioral tests and physiological mechanisms. Trends in Neurosciences *39*, 74-85.

10.1016/j.tins.2015.12.007.

55.   Hartwigsen, G., Bzdok, D., Klein, M., Wawrzyniak, M., Stockert, A., Wrede, K.,

Classen, J., and Saur, D. (2017). Rapid short-term reorganization in the language

network. Elife *6*. 10.7554/eLife.25964.

56.   Jackson, R.L., Hoffman, P., Pobric, G., and Ralph, M.A.L. (2016). The semantic

network at work and rest: Differential connectivity of anterior temporal lobe

subregions. Journal of Neuroscience *36*, 1490-1501. 10.1523/JNEUROSCI.2999-

15.2016.

57.   Humphreys, G. F., Lambon Ralph, M. A., & Simons, J. S. (2021). A Unifying Account of
Angular Gyrus Contributions to Episodic and Semantic Cognition. Trends in
neurosciences, 44(6), 452–463. https://doi.org/10.1016/j.tins.2021.01.006

58.   Bonner, M. F., & Price, A. R. (2013). Where is the anterior temporal lobe and what

does it do?. The Journal of neuroscience : the official journal of the Society for

Neuroscience, 33(10), 4213–4215. https://doi.org/10.1523/JNEUROSCI.0041-13.2013

Figure 1. Experimental design, and stimulus characteristics.

(**A**) **Experimental stimuli.** Twenty gestures were paired with 20 relevant speech stimuli. Two

separate Pre-tests were executed to define the minimal length of each gesture and speech

required for semantic identification, namely, the discrimination point (DP) of gesture and the

identification point (IP) of speech. Overall, a mean of 183.78 ms (SD = 84.82) was found for

the DP of gestures and the IP of speech was 176.40 ms (SD = 66.21). The onset of speech

was set at the gesture DP. Responses for each item were assessed utilizing information-

theoretic complexity metrics to quantify the information content of both gesture and speech

during integration, employing entropy and MI.

(**B**) **Procedure of Experiment 1.** HD-tDCS, including Anodal, Cathodal, or Sham conditions,

was administered to the IFG or pMTG) using a 4 * 1 ring-based electrode montage. Electrode

F7 targeted the IFG, with return electrodes placed on AF7, FC5, F9, and FT9. For pMTG

stimulation, TP7 was targeted, with return electrodes positioned on C5, P5, T9, and P9.

Sessions lasted 20 minutes, with a 5-second fade-in and fade-out, while the Sham condition

involved only 30 seconds of stimulation.

(**C**) **Procedure of Experiment 2.** Eight time windows (TWs, duration = 40 ms) were

segmented in relative to the speech IP. Among the eight TWs, five (TW1, TW2, TW3, TW6,

and TW7) were chosen based on the significant results in our prior study[23]. Double-pulse

TMS was delivered over each of the TW of either the pMTG or the IFG.

(**D**) **Procedure of Experiment 3.** Semantically congruent gesture-speech pairs were

presented randomly with Electroencephalogram (EEG) recorded simultaneously. Epochs

were time locked to the onset of speech and lasted for 1000 ms. A 200 ms pre-stimulus

baseline correction was applied before the onset of gesture stoke. Various elicited

components were hypothesized.

Figure 2. Quantification formulas (A) and distributions of each stimulus in Shannon's entropy (B).

Two separate pre-tests (N = 30) were conducted to assign a single verb for describing each of the isolated 20 gestures and 20 speech items. Responses provided for each item were transformed into Shannon's entropy using a relative quantification formula. Gesture (**B left**) and speech (**B right**) entropy quantify the randomness of gestural or speech information, representing the uncertainty of probabilistic representation activated when a specific stimulus occurs. Joint entropy (**B middle**) captures the widespread nature of the two sources of information combined. Mutual information (MI) was calculated as the difference between joint entropy with gesture entropy and speech entropy combined (**A**), thereby capturing the overlap of gesture and speech and representing semantic integration.

Figure 3. tDCS effect over semantic congruency.

(A) tDCS effect was defined as active-tDCS minus sham-tDCS. The semantic congruency

effect was calculated as the reaction time (RT) difference between semantically incongruent

and semantically congruent pairs (Rt(incongruent) - Rt(congruent)).

(B) Correlations of the tDCS effect over the semantic congruency effect with three information

models (gesture entropy, speech entropy and MI) are displayed with best-fitting regression

lines. Significant correlations are marked in red**.** * $p < 0.05$, ** $p < 0.01$ after FDR correction.

Figure 4. TMS impacts on semantic congruency effect across various time windows (TW).

 (A) Five time windows (TWs) showing selective disruption of gesture-speech integration were chosen: TW1 (-120 to -80 ms relative to speech identification point), TW2 (-80 to -40 ms), TW3 (-40 to 0 ms), TW6 (80 to 120 ms), and TW7 (120 to 160 ms). TMS effect was defined as active-TMS minus vertex-TMS. The semantic congruency effect was calculated as the reaction time (RT) difference between semantically incongruent and semantically congruent pairs.

(B) Correlations of the TMS effect over the semantic congruency effect with three information models (gesture entropy, speech entropy and MI) are displayed with best-fitting regression lines. Significant correlations are marked in red**.** * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ after FDR correction.

838
839
Figure 5. ERP results of gesture entropy (A), speech entropy (B) or MI (C).
Four ERP components were identified from grand-average ERPs at the Cz electrode,
contrasting trials with the lower 50% (red lines) and the higher 50% (blue lines) of gesture
entropy, speech entropy or MI. Clusters of adjacent time points and electrodes were
subsequently identified within each component using a cluster-based permutation test.
Topographical maps depict amplitude differences between the lower and higher halves of
each information model, with significant ROIs (**D1-H1 middle**) or electrode clusters (**D2-H2**
middle) highlighted in black. Solid rectangles delineating the ROIs that exhibited the maximal
correlation and paired t-values (**D1-H1 middle**). T-test comparisons with normal distribution
lines (**left**) and correlations with best-fitting regression lines (**right**) are calculated and
illustrated between the average ERP amplitude within the rectangular ROI (**D1-H1**) or the

elicited clusters (**D2-H2**) and the three information models individually. $* p < 0.05$, $** p < 0.01$

after FDR correction.

Figure 6. Progressive processing stages of gesture–speech information within the pMTG-IFG loop.

Correlations between the TMS disruption effect of pMTG and IFG with three information models are represented by the orange line and the green lines, respectively. Black lines denote the strongest correlations of ROI averaged ERP components with three information models. * $p < 0.05$, ** $p < 0.01$ after FDR correction.