Epilepsy-linked kinase CDKL5 phosphorylates voltage-gated calcium channel Cav2.3, altering inactivation kinetics and neuronal excitability

Marisol Sampedro-Castañeda[1,5]*, Lucas L Baltussen[1,4,5], Andre T Lopes[1], Yichen Qiu[2], Liina Sirvio[1], Simeon R. Mihaylov[1], Suzanne Claxton[1], Jill C Richardson[3], Gabriele Lignani[2], Sila K Ultanir[1]*

1. Kinases and Brain Development Lab, The Francis Crick Institute, 1 Midland Road, London NW1 1AT, United Kingdom

2. Department of Clinical and Experimental Epilepsy, UCL Queen Square Institute of Neurology, 808 Queen Square House, London WC1N 3BG, United Kingdom

3. UK Neuroscience Group, MSD R&D Innovation Centre, 120 Moorgate, London, EC2M 6UR, United Kingdom

*Correspondence: sila.ultanir@crick.ac.uk

marisol.sampedro-castaneda@crick.ac.uk

4. Current address: Laboratory of Synapse Biology (VIB-KU Leuven), Department of Neurosciences, ON5 Herestraat 49, 3000 Leuven, Belgium

5. These authors contributed equally

SUMMARY

Developmental and epileptic encephalopathies (DEEs) are a group of rare childhood disorders characterized by severe epilepsy and related cognitive deficits. Numerous DEE genes have been discovered thanks to advances in genomic diagnosis, yet putative molecular links between these disorders are not known. CDKL5 deficiency disorder (CDD, DEE2) is one of the most common forms of genetic epilepsy; it is caused by loss-of-function mutations in the brain-enriched kinase CDKL5. To elucidate CDKL5 function, we looked for CDKL5 substrates using a SILAC based phosphoproteomic screen. We identified the voltage-gated Ca^{2+} channel Cav2.3 (encoded by *CACNA1E*) as a novel physiological target of CDKL5 in mice and humans. Recombinant channel electrophysiology and interdisciplinary characterization of Cav2.3 phosphomutant mice revealed that the loss of Cav2.3 phosphorylation leads to channel gain-of-function via slower channel inactivation and enhanced acetylcholine-induced stimulation, resulting in increased neuronal excitability. These changes in Cav2.3 closely resemble those described for gain-of-function point-mutations in *CACNA1E* that cause DEE69, a disorder sharing clinical features with CDD. Our results show that these two single-gene disorders are mechanistically related. We suggest that CDD is partly a channelopathy with Cav2.3 gain-of-function, thus Cav2.3 inhibition could be therapeutic in these DEEs.

KEYWORDS

CDKL5, CACNA1E, Cav2.3, epilepsy, developmental and epileptic encephalopathy, DEE2, DEE69

HIGHLIGHTS

- CDKL5 phosphorylates Cav2.3 at Ser15 and reduces channel inactivation time
- Loss of CDKL5 causes Cav2.3 gain-of-function, mimicking pathological *CACNA1E* mutations
- Hippocampal neurons from Cav2.3 S15A mutant mice have an enhanced excitability response to cholinergic stimulation
- A common molecular mechanism connects CDKL5 and *CACNA1E* epileptic encephalopathies

INTRODUCTION

Developmental and epileptic encephalopathies (DEE) are characterized by severe early-onset epileptic activity accompanied by global developmental and cognitive impairments. The genetic aetiology for 100 DEEs have been identified, but targeted therapies remain scarce. Elucidating shared pathways in infantile-onset epilepsies will reveal molecular links between disease genes and greatly advance common targeted treatments.

Cyclin dependent kinase like-5 (CDKL5) is a brain-enriched serine-threonine kinase. *De novo* loss-of-function mutations in the X-linked *CDKL5* gene, including missense, nonsense and insertions/deletions, cause CDKL5 deficiency disorder (CDD) (OMIM 300672, 300203) [1-4]. Pathogenic missense mutations, almost exclusively located in the kinase domain, indicate that kinase activity is critical for CDD pathology [5-8]. CDD is characterized by infantile-onset, intractable seizures, profound neurodevelopmental impairment in motor and sensory function, impaired language acquisition and autonomic disturbances [2, 4]. The estimated incidence for CDD is 1/42,000 live births each year [9-11] with approximately 80% female patients, making it the second most common type of genetic childhood epilepsy [12, 13].

CDKL5 is highly enriched throughout the forebrain with expression starting at late embryonic stages in rodents and humans [14-18]. Low levels of expression were reported in other tissues [15]. Phosphorylation targets include microtubule binding proteins EB2 and MAP1S [5, 19] and transcriptional regulators ELOA [20] and Sox9 [21], showing its diverse cellular roles. CDKL5 KO mouse neurons have deficits in synaptic transmission, dendritic spine development, neurite outgrowth, cilia elongation and microtubule dynamics [22-27]. However, molecular mechanisms linking CDKL5 loss to neuronal hyperexcitability remain unknown. The identification of key physiological targets of CDKL5 directly involved in the regulation of cellular excitability may help elucidate epileptogenic mechanisms and lead to targeted therapies.

Voltage-gated calcium channels (VGCC) have a key role in physiology, driving neuronal excitability and allowing influx of second messenger Ca^{2+} ions in response to membrane depolarization [28-30]. VGCC subtype Cav2.3 is formed by the ion-conducting α1E subunit (encoded by *CACNA1E*) in complex with accessory subunits from the β and α2δ families. Cav2.3 is highly expressed in the CNS, where it is localized at both pre- and post-synaptic sites in neurons, as shown functionally and by immuno-EM [31-33]. *CACNA1E* knockout mice demonstrated that Cav2.3 mediates neuronal R-type Ca^{+2} currents [33-35], so-called due to their "resistance" to organic VGCC channel inhibitors [36, 37]. Cav2.3 is dynamically regulated by phosphorylation at multiple sites [38, 39]. In general, VGCC phosphorylation is an important mechanism by which neuromodulators and their G-protein coupled receptors (GPCRs) exert their effect on cellular excitability [40].

De novo heterozygous missense mutations in *CACNA1E* cause an infantile epileptic encephalopathy (OMIM 618285) [41, 42]. Cav2.3 variants studied *in vitro* have altered gating and/or inactivation kinetics, leading to gain-of-function (GoF) of Cav2.3 [41, 42]. Patients with *CACNA1E* mutations have overlapping clinical phenotypes with CDD, such as intractable seizures, profound intellectual disability and hypotonia. Single-gene DEEs such as this one are estimated to impact approximately 1/2100 live births [9], but their real prevalence is unknown as more genes are being identified rapidly due to recent advances in genetic diagnosis [43, 44]. Despite, overlapping clinical features, potential molecular links between each of these rare disorders remain to be elucidated.

In this work, we apply for the first time a global phosphoproteomics approach in CDKL5 knockout mice and identify and validate Cav2.3 as a novel substrate of CDKL5 in human and mouse neurons. Through our analysis of Cav2.3 function and characterization of a novel Cav2.3 phosphomutant mouse line, in which CDKL5 phosphorylation site is mutated, we reveal that lack of phosphorylation leads to Cav2.3 gain-of-function and hyperexcitability. Our results indicate that developmental and epileptic encephalopathies caused by *CDKL5* and *CACNA1E* are related at the molecular level. We propose Cav2.3 as a converging therapeutic target for these disorders.

RESULTS

Cav2.3 is a CDKL5 phosphorylation substrate in mice and humans

We used stable isotope labelling of amino acids in cell culture (SILAC) [45] to differentially tag newly synthesized proteins in primary cortical cultures from wild type (WT) and CDKL5 knockout (KO) mice [46] and compare changes in protein phosphorylation levels (Fig. 1a). As expected in CDKL5 KO neurons, we observe a significant reduction in CDKL5 pS407 and in EB2 pS222, a known CDKL5 phosphorylation target [19]. Phosphorylation of EB2 S221 was also decreased, indicating co-regulation of these neighbouring sites. More importantly, a potential new target protein, the ion channel Cav2.3, showed strongly reduced phosphorylation at S15 (pS15), in CDKL5 KO neurons. Strikingly, the Cav2.3 S15 site matches the RPXS/T* consensus motif [5, 19], suggesting direct CDKL5 phosphorylation (Fig. 1a, b). S15, located in the N-terminus of the α1E channel pore subunit (Fig. 1b), is conserved in humans (S14) but absent in the related Cav2.1 (*CACNA1A*) and Cav2.2 (*CACNA1B*) channels (Fig. 1b).

To test if CDKL5 can phosphorylate Cav2.3 S15, we generated a phosphospecific antibody targeting this site. Overexpression of Cav2.3 (& ancillary subunits) together with the CDKL5

kinase domain in HEK293 cells shows strong Cav2.3 phosphorylation *in vitro* (Cav2.3 pS14, Fig. 1c, f). This phosphorylation is absent when overexpressing phosphomutant Cav2.3 or kinase-dead CDKL5, demonstrating the specificity of the antibody and the dependence of phosphorylation at this site on CDKL5 kinase activity. Importantly, full-length CDKL5 also directly phosphorylated Cav2.3 *in vitro* (Fig. S1a, b). Cav2.3 phosphorylation is consistently reduced in the cortex of CDKL5 KO mice at postnatal day P20 (Cav2.3 pS15), validating the substrate *in vivo* (Fig. 1d, g). Finally, Cav2.3 pS14 was decreased in human iPSC-derived neurons from CDD patients (Fig. 1e, h), demonstrating altered phospho-regulation of this target in CDD. Together, these results show that Cav2.3 is a *bona fide* CDKL5 phosphorylation target in mice and humans.

Loss of Cav2.3 Ser14 phosphorylation slows baseline channel kinetics and amplifies muscarinic regulation

Cav2.3 is causally linked to epilepsy by two main lines of evidence. First, channel deletion renders mice resistant to tonic-clonic seizures [47-49]. On the other hand, Cav2.3 GoF leads to severe epileptic encephalopathy in humans [41, 42]. The remarkable overlap in clinical manifestations between CACNA1E and CDD patients, including seizures and neurodevelopmental delays, raises the possibility that Cav2.3 regulation by CDKL5 might affect channel function and in turn neuronal excitability.

To test the effect of pS14 on Cav2.3 channel properties we used a HEK293 cell line stably expressing the human auxiliary subunits β3 and α2δ1 and transiently transfected human α1E and full-length CDKL5 plasmids (Fig. S1a, b). Whole-cell patch-clamp recordings of Cav2.3 Ba^{2+} currents revealed slower decay kinetics (inactivation τ) in S14A mutant channels (Fig. 2a-c). Notably, in absence of CDKL5, WT Cav2.3 decay kinetics were indistinguishable from S14A Cav2.3 mutant (Fig. 2d). We also found a statistically significant increase in whole-cell current density at maximum activation voltages in S14A when compared to WT Cav2.3 in presence of CDKL5 (Fig. 2e). To account for differences in biophysical properties that may be introduced by α1E association with a different β subunit [50, 51], we conducted parallel experiments in a HEK293 cell line stably expressing human α2δ1 and β1b, another Cavβ subunit associated with α1E in the brain [52]. We used Ca^{2+} as charge carrier to examine channel properties in a more physiological context. Decay kinetics of WT Cav2.3 currents with β1b were substantially slower than currents with β3 (Fig. S2 a, b *vs.* Fig 1a, b), as previously reported [50]. In absence of CDKL5 phosphorylation (S14A α1E mutant or kinase-dead CDKL5 conditions), Cav2.3 current inactivation was further slowed (Fig. S2a-c), recapitulating our results with β3 (Fig. 2a-c). Current density at maximal activation voltages did not reach significance in the β1-expressing cells, where currents were generally larger (Fig. S2d).

Finally, we observed no changes in half maximal activation and inactivation potential or time
course of recovery from fast inactivation in β3- or β1-expressing cells (Fig. 2f & S2e,f).

Altogether, our results indicate that CDKL5-mediated Ser14 phosphorylation of α1E speeds
up the inactivation of human Cav2.3 and suppresses current amplitude, independent of Cavβ
subtype or charge carrier. Consequently, absence of CDKL5 leads to Cav2.3 GoF, with larger
and more prolonged currents. A subset of *CACNA1E* GoF mutations described in human DEE
patients also cause slower inactivation kinetics and/or increased current density [41], mirroring
the phenotypes observed in the unphosphorylated channel. Thus, the functional deficits in
Cav2.3 observed in absence of CDKL5 phosphorylation match those caused by pathological
point mutations in Cav2.3 and could partly explain the pathophysiology of CDD.

Cav2.3 channels are downstream targets of muscarinic acetylcholine receptors, which modify
channel function predominantly via PKC-mediated phosphorylation, leading to current
enhancement in HEK293 cells and neurons [53-55]. Given the observed GoF in S14A Cav2.3,
we hypothesized that muscarinic receptor enhancement may also be affected by CDKL5
phosphorylation. To test this, we co-expressed Cav2.3, CDKL5 and muscarinic acetylcholine
receptor type 3 or 1 (M3 or M1), which are prominent brain subtypes known to regulate
neuronal Cav2.3 post-synaptically. As expected, application of the non-hydrolysable agonist
carbachol substantially increased current amplitude regardless of Ser14 phosphorylation in
both β3 and β1 stable cell lines (Fig. 2g-l). The current increase with carbachol was 1.5- to 2-
fold greater in cells without phospho-Ser14 at -10 mV (Fig. 2g, h, j, l), reflecting Cav2.3 gain-
of-function in absence of phosphorylation. The half-maximal activation potential ($V_{1/2}$) was left-
shifted by 5 mV for WT channels in both cell lines, a described effect of muscarinic receptor
activation [54, 56]. Interestingly, without Ser14 phosphorylation this shift was significantly greater,
implying an effect of phospho-Ser14 on the modulation of Cav2.3 gating (Fig. 2m-p). This
represents an additional gain-of-function due to loss of CDKL5, a phenotype comparable to
pathogenic *CACNA1E* mutations that alter $V_{1/2}$ [41].

Upon GPCR activation, Cav2.3 channels can be suppressed by $G_{\beta\gamma}$ subunits. This effect is
has been observed in Gαq-coupled receptors including M1/M3 in HEK293 cells [54] and CA1
neurons [53]. The Cav2.3 N-terminus and Cavβ are involved in $G_{\beta\gamma}$ regulation [57, 58]. For this
reason, we investigated the possibility that the enhanced carbachol effects in absence of N-
terminal pS14 could be due to reduced $G_{\beta\gamma}$ suppression. To distinguish between PKC-
mediated enhancement and $G_{\beta\gamma}$ suppression, we boosted PKC activity prior to carbachol
application using phorbol 12-myristate 13-acetate (PMA), reliably increasing Cav2.3 currents
[54, 59]. With further addition of carbachol we found no evidence of muscarinic inhibition of human
WT Cav2.3 channels (Fig. S2g,h). Moreover, co-expression of Cav2.3 and Gi-coupled
dopamine D2 receptors to further probe $G_{\beta\gamma}$ inhibition in our system, revealed significant
reduction in peak currents upon D2 activation with quinpirole [57]. However, there was no

difference in inhibitory modulation with or without Ser14 phosphorylation (Fig. S2i,j). The
above manipulations suggest that $G_{\beta\gamma}$ is not involved in pathological Cav2.3 GoF.

Our results describe two Cav2.3 gain-of-function (GoF) mechanisms in the absence of CDKL5
phosphorylation: slower decay kinetics and a left shift in voltage-dependence of activation
($V_{1/2}$) upon muscarinic activation, both reflected in increased Ca^{2+} influx through Cav2.3
channels. These GoF features are reminiscent of those described for pathological Cav2.3
variants.

Altered R-type current inactivation and increased excitability in neurons from Cav2.3
S15A phosphomutant mice

To investigate the role of Cav2.3 S15 phosphorylation in neurons, we generated Cav2.3 S15A
phosphomutant mice using CRISPR-Cas9 genome editing. We found that the total S15
phosphorylation levels detected by our phosphoantibody were reduced to <20% of control
levels in homozygous S15A mice (HOM S15A) while total Cav2.3 levels were not changed
(Fig. 3a-c). Cav2.3 is highly expressed in the somato-dendritic region of CA1 hippocampal
neurons [32] where it underlies most of the R-type current and regulates intrinsic excitability [33,
53]. We performed whole-cell somatic recordings in acute slices from young mice and recorded
pharmacologically isolated Cav2.3 Ca^{2+} currents in these cells. When compared to wild type
littermates (WT), HOM S15A mice showed significantly slower inactivation τ at maximal
conductance voltages, recapitulating HEK293 cell results (Fig. 3 d,e). Current amplitudes at
the soma did not differ (Fig. 3f).

We next tested the effect of Cav2.3 S15A on neuronal excitability and cholinergic
neuromodulation in response to depolarising current injections in adult CA1 pyramidal cells.
Resting membrane potential (V_{rest}), input resistance (R_{in}) and firing properties were identical
between WT and S15A mice under baseline conditions (Fig. 3g-i). Following bath application
of carbachol, we observed a 4mV depolarisation in V_{rest} and a 25% increase in R_{in} (Fig 3g),
changes that were comparable between WT and S15A neurons (Fig. S3a). These alterations
are likely due to conductances other than Cav2.3, in line with previous reports [56, 60, 61].

In neurons, cholinergic activation can elicit sustained depolarizing plateau potentials (DPPs):
all-or-none, large amplitude depolarizations that outlast stimulus-evoked firing [56, 62, 63]. A
muscarinic medium duration afterdepolarization (ADP) is also reported in CA1 pyramidal cells
[64]. Up-regulation of Cav2.3-mediated R-type currents is required for DPPs and ADPs in
hippocampus [65] and cortex [63]. We inspected these Cav2.3-dependent hyper-excitability
measures in S15A mice. Carbachol application significantly increased action potential firing
frequency in WT and S15A neurons (Fig. 3 h-k). At current injections greater than 250 pA,

however, phosphomutant neurons responded with fewer spikes and depolarization block towards the end of the stimulus step, indicating greater underlying depolarization. The percentage of cells exhibiting at least one DPP or all-or-none afterdepolarization (ADP) >8 mV upon carbachol treatment was significantly higher in S15A mice, showing an enhanced depolarization response (Fig. 3j, k). Importantly, the occurrence of DPP/large ADPs was shifted towards lower current injections in S15A cells (Fig. 3l). Greater perisomatic depolarization in S15A neurons was also reflected in a larger ADP at 100ms post stimulus and more spike attenuation (Fig. S3b, c). Together with the enhanced carbachol-mediated left-shift in $V_{1/2}$ observed in phosphomutant channels in HEK293 cells (Fig. 2 j,l), these increased neuronal depolarizations in Cav2.3 phosphomutant mice suggest facilitation of gating of endogenous Cav2.3 without S15 phosphorylation upon cholinergic agonist application.

We tested if other conductances known to be regulated by carbachol are altered in Cav2.3 phosphomutants. The medium- and slow-afterhyperpolarizations (AHPs) are critical for spike frequency regulation and their underlying K^+ and mixed-cationic currents are targets of the muscarinic system [60, 66, 67]. Baseline AHPs elicited by a short spike burst were not different between WT and S15A mice and were equally suppressed by carbachol application (Fig. S3d). We also analysed baseline small-conductance Ca^{2+}-activated K^+ (SK) currents and spontaneous excitatory synaptic currents (EPSCs), known to be functionally coupled to [68] or modulated by [69] Cav2.3 in CA1 neurons. We hypothesized that these may be affected given the prolonged Cav2.3 currents in S15A mice, but our measurements show no differences between genotypes (Fig. S3e-h). Therefore, we suggest that the increased carbachol-induced excitability of CA1 neurons in Cav2.3 S15A knock-in mice is specifically due to Cav2.3 GoF.

Cav2.3 S15A phosphomutant mice have behavioural and EEG deficits

Multiple CDKL5 KO mouse models have been generated [14, 17, 46] by deleting exons 4, 2 or 6, all of which lead to loss of CDKL5 protein. Several behavioural deficits were reported in these mice [46, 70, 71]. We investigated behavioural phenotypes in Cav2.3 S15A phosphomutant mice and tested if they recapitulate reported CDKL5 KO impairments. Initially, we observed no overt behavioural alterations or spontaneous seizures, as reported for CDKL5 KO mice [14, 17, 72], but see [73, 74]. However, when monitored in a home-cage environment for prolonged periods, S15A showed reduced locomotion and voluntary wheel use (Fig. 4a, b), prompting closer behavioural examination. We observed minor changes in locomotion and zone preference in the open field test, but only towards the end of a 30 min trial (Fig. 4c, S4a), potentially indicating fatigue rather than anxiety. Unlike CDKL5 KOs, S15A mice did not present hindlimb clasping (mean scores: WT 0.041, HOM S15A 0, p=0.334, unpaired t test) and performed equally in the rotarod test (Fig. 4d) when compared to WT littermates. Cognitive abilities in the

Y- and Barnes maze were also unchanged in phosphomutants (Fig. S4b-e). On the other hand, S15A mice had mild impairments in sociability (Fig. 4e). A consistently reported phenotype of CDKL5 KO mice is altered contextual fear conditioning responses [14, 71]. Strikingly, memory formation, retention and extinction in this task were also defective in Cav2.3 S15A (Fig. 4f), indicating a clear overlap with CDKL5 KOs.

Finally, given the high epilepsy incidence in human CDKL5 and CACNA1E DEE patients, seizure susceptibility was analysed in a separate cohort using wireless electrocorticogram (ECoG) transmitters [75] and repeated low-dose kainic-acid (KA) injections [76] (Fig. 4g-i). We found no changes in baseline ECoG (Fig 4g) or overall KA-induced seizure susceptibility in S15A mice (Fig. 4h), as reported for CDKL5 KO mouse models [17, 35]. Interestingly, however, female phosphomutant mice had a reduced threshold to stage 5 behavioural seizures with an increase in ECoG activity during KA (Fig. 4h,i).

In summary, our *in vivo* analyses of S15A mutant mice demonstrate similarities with CDD mouse models and patients, including social, motor and cognitive impairments, and to some extent increased seizure susceptibility. These observations suggest that loss of pS15 is a key contributor to the phenotypic features of CDD.

DISCUSSION

There are currently no disease-targeting therapies for CDD and the causes of aberrant excitability in absence of CDKL5 are unknown. Here we identify the alpha subunit of Cav2.3 channel complex as a CDKL5 phosphorylation substrate in mouse and human neurons. *CACNA1E* mutations cause severe early onset epilepsy in humans and there is remarkable overlap between CDD and patients with *CACNA1E* mutations. These include intractable seizures, global developmental delay, intellectual disability, hypotonia, hyperkinetic movements and sleep disturbances. A consistent feature of Cav2.3 variants associated with epilepsy is GoF of the channel, by left-shifting the activation voltage, altering maximal current and/or by slowing inactivation [41]. Our data shows that deletion of CDKL5 results in loss of Cav2.3 S15 phosphorylation, leading to GoF due to slowed inactivation and amplified gating modulation, similar to the effects observed in Cav2.3 variants. Our results establish Cav2.3 overactivity as a common feature of CACNA1E (DEE69) and CDKL5 (DEE2) epileptic encephalopathies, potentially explaining overlapping clinical features of these diseases.

Cytoplasmic N-terminal S15 phosphorylation of Cav2.3 has been previously detected in global phosphoproteomics screens from brain (phosphosite.org), but the responsible kinase and the role of this phosphorylation were unknown. N-terminal deletions in Cav2.3 [57] or the related Cav2.2 [77, 78], result in channels with normal gating and conduction properties. Inactivation

kinetics were not measured in these studies, yet interestingly Cav2.2 N-terminal deletion mutants seemed to have slower decaying currents, indicating that this region participates in the regulation of inactivation [77]. Analogously, loss of N-terminal Ser14/15 phosphorylation slows Cav2.3 inactivation in HEK293 cells and neurons, underscoring a previously overlooked regulatory role of the Cav2.3 N-terminus. This domain may interact with Cavβ, as previously suggested for Cav2.2 [78].

We also find that cholinergic stimulation amplifies Cav2.3 GoF in absence of CDKL5 phosphorylation, by prominently shifting activation $V_{1/2}$ and increasing currents at maximal activation voltages, thus revealing a second molecular mechanism by which loss of pS14/15 enhances Cav2.3 function. It is known that muscarinic enhancement of Cav2.3 depends on Gαq signalling and PKC phosphorylation at multiple sites, including the intracellular loop I-II [59], the most prominent site of α1-β interaction [51]. We show that loss of pS14 interferes with PKC enhancement upon carbachol application, but the mechanisms of this interplay are yet to be established.

Our recordings from hippocampal neurons in Cav2.3 S15A mice demonstrate that S15 regulates Cav2.3 function in neurons. We show that pS15 is critical for R-type current inactivation and regulation of firing by cholinergic stimulation. Isolating Cav2.3 currents is challenging due to space clamp limitations and the need for inhibitors of other voltage-gated channels. These drugs may have promiscuous effects on Cav2.3 [79] or incompletely block other Ca^{2+} currents (Tai et al., 2006). We observed enhanced neuronal excitability and increased pro-epileptiform cholinergic ADPs/DPPs [56, 61] in Cav2.3 phosphomutant mice, an effect that was not linked to carbachol suppression of AHPs or other intrinsic excitability properties in these cells. Instead, we suggest that the GoF of unphosphorylated Cav2.3 underlies the hyper-excitability seen in response to muscarinic stimulation. Interestingly, CDKL5 KO mice have altered cholinergic tone [80] and the muscarinic antagonist Solifenacin was shown to ameliorate network defects in CDD patient-derived neurons [81], supporting an enhanced cholinergic modulation in this human model.

We find striking behavioural similarities between our Cav2.3 phosphomutant mice and CDKL5 KOs. Specifically, S15A mice showed reduced encoding and retention of fear memory, a robust phenotype identified by several groups in CDKL5 KO models [14, 71, 72]. In contrast, Cav2.3 KO mice display enhanced fear [35]. Home-cage behaviour and social interaction phenotypes are also common between Cav2.3 phosphomutant and CDKL5 KO mice [46, 71]. The deficits observed in voluntary wheel in Cav2.3 phosphomutants could be indicative of mild hypotonia [82, 83]. Importantly, these phosphomutant mouse phenotypes mirror human clinical manifestations of CDD, arguing that slowed inactivation and enhanced cholinergic modulation of unphosphorylated Cav2.3 may contribute to these symptoms.

Cav2.3 KO mice exhibit reduced susceptibility to chemical absence seizures [84-86] and

hippocampal KA-induced seizures [49]. In agreement with Cav2.3's role in epilepsy, we find

increased seizure susceptibility in females in the low-dose KA injection paradigm in our Cav2.3

S15A mice. The difference between males and females, observed in both ECoG and

behavioural seizure analysis, may arise from increased basal function of Cav2.3 in the female

brain downstream of hormonal signals. Gender should be taken into consideration in future

mouse CDKL5 KO studies, where KA-induced seizure susceptibility changes have not been

observed [17, 46] and susceptibility to other chemoconvulsants were reported [72] but not widely

reproduced [17]. Disturbance-associated seizure-like events have been observed in aged

heterozygous females in two different CDD mouse models (PMID: 33197557), an occurrence

that may be linked to developmental changes in Cav2.3 regulation.

In rodents, it is possible to reverse adult phenotypes with reintroduction of CDKL5 genetically

[70] or using AAV viruses [87], indicating an open therapeutic window for treatment. Here, we

report a novel functional regulation of Cav2.3 by CDKL5 phosphorylation, which impinges of

neuronal excitability. Thus Cav2.3 GoF in absence of CDKL5 may be an important molecular

mechanism in CDD pathology. In addition to CACNA1E encephalopathy, increased Ca^{2+} influx

through Cav2.3 may contribute to epileptogenesis in Fragile X syndrome [88] and juvenile

myoclonic epilepsy [89]. Cav2.3 also increases susceptibility to drug-induced Parkinson's

disease [90, 91]. We propose that Cav2.3 inhibitors/ modulators could be beneficial for CDD

patients and potentially for a broader population of neurological patients in the future.

Acknowledgements

We thank Ultanir lab members for valuable discussion and thoughtful comments. We thank
Helen Flynn for valuable assistance with mass spectrometry. We thank Crick GEMS team for
creation of CACNA1E S15A mice. This work was supported by the Francis Crick Institute
which receives its core funding from Cancer Research UK (CC2037), the UK Medical
Research Council (CC2037), and the Wellcome Trust (CC2037); Loulou Foundation Project
Grant (11015); Crick-MSD Framework collaboration grant (11202). For the purpose of Open
Access, the author has applied a CC BY public copyright licence to any Author Accepted
Manuscript version arising from this submission.

Author contributions

MSC, LLB, GL and SKU designed the experiments. MSC performed and analysed
electrophysiology and biochemistry experiments. LLB performed mass spectrometry. YQ and
GL conducted/ analysed EEG and kainic acid-induced seizure experiments. ATL and MSC
performed behaviour experiments/ analysis. LS, SM, SC provided technical assistance in
molecular biology and electrophysiology. MSC and SKU wrote the initial version of the
manuscript. JR contributed to the original draft and review/editing along with all authors.

Declaration of interests

401 The authors declare no competing interest.

Inclusion and diversity

403 We ensured the sex balance in our mouse cohorts. Authors in this paper have balanced
404 gender representation. This manuscript includes authors with diverse race, ethnicity, and
405 sexual orientation.

References

1.   Kalscheuer, V.M*., et al.* Disruption of the serine/threonine kinase 9 gene causes severe X-linked infantile spasms and mental retardation. *American journal of human genetics* **72**, 1401-1411 (2003).
2.   Olson, H.E*., et al.* Cyclin-Dependent Kinase-Like 5 Deficiency Disorder: Clinical Review. *Pediatr Neurol* **97**, 18-25 (2019).
3.   Van Bergen, N.J*., et al.* CDKL5 deficiency disorder: molecular insights and mechanisms of pathogenicity to fast-track therapeutic development. *Biochemical Society transactions* **50**, 1207-1224 (2022).
4.   Leonard, H*., et al.* CDKL5 deficiency disorder: clinical features, diagnosis, and management. *Lancet Neurol* **21**, 563-576 (2022).
5.   Munoz, I.M*., et al.* Phosphoproteomic screening identifies physiological substrates of the CDKL5 kinase. *The EMBO journal* **37** (2018).
6.   Hector, R.D*., et al.* CDKL5 variants: Improving our understanding of a rare neurologic disorder. *Neurol Genet* **3**, e200 (2017).
7.   Krishnaraj, R., Ho, G. & Christodoulou, J. RettBASE: Rett syndrome database update. *Human mutation* **38**, 922-931 (2017).
8.   Bahi-Buisson, N*., et al.* Recurrent mutations in the CDKL5 gene: genotype-phenotype relationships. *American journal of medical genetics. Part A* **158A**, 1612-1619 (2012).
9.   Symonds, J.D*., et al.* Incidence and phenotypes of childhood-onset genetic epilepsies: a prospective population-based national cohort. *Brain : a journal of neurology* **142**, 2303-2318 (2019).
10.   Lindy, A.S*., et al.* Diagnostic outcomes for genetic testing of 70 genes in 8565 patients with epilepsy and neurodevelopmental disorders. *Epilepsia* **59**, 1062-1071 (2018).
11.   Kothur, K*., et al.* Diagnostic yield of targeted massively parallel sequencing in children with epileptic encephalopathy. *Seizure* **59**, 132-140 (2018).
12.   Bahi-Buisson, N. & Bienvenu, T. CDKL5-Related Disorders: From Clinical Description to Molecular Genetics. *Mol Syndromol* **2**, 137-152 (2012).
13.   Siri, B*., et al.* CDKL5 deficiency disorder in males: Five new variants and review of the literature. *Eur J Paediatr Neurol* **33**, 9-20 (2021).
14.   Wang, I.T*., et al.* Loss of CDKL5 disrupts kinome profile and event-related potentials leading to autistic-like phenotypes in mice. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 21516-21521 (2012).
15.   Hector, R.D*., et al.* Characterisation of CDKL5 Transcript Isoforms in Human and Mouse. *PloS one* **11**, e0157758 (2016).
16.   Hector, R.D*., et al.* Characterisation of Cdkl5 transcript isoforms in rat. *Gene* **603**, 21-26 (2017).
17.   Okuda, K*., et al.* CDKL5 controls postsynaptic localization of GluN2B-containing NMDA receptors in the hippocampus and regulates seizure susceptibility. *Neurobiology of disease* **106**, 158-170 (2017).
18.   Rusconi, L*., et al.* CDKL5 expression is modulated during neuronal development and its subcellular distribution is tightly regulated by the C-terminal tail. *The Journal of biological chemistry* **283**, 30101-30111 (2008).
19.   Baltussen, L.L*., et al.* Chemical genetic identification of CDKL5 substrates reveals its role in neuronal microtubule dynamics. *The EMBO journal* **37** (2018).
20.   Khanam, T*., et al.* CDKL5 kinase controls transcription-coupled responses to DNA damage. *The EMBO journal* **40**, e108271 (2021).
21.   Kim, J.Y*., et al.* A kinome-wide screen identifies a CDKL5-SOX9 regulatory axis in epithelial cell death and kidney injury. *Nat Commun* **11**, 1924 (2020).
22.   Zhou, A., Han, S. & Zhou, Z.J. Molecular and genetic insights into an infantile epileptic encephalopathy - CDKL5 disorder. *Front Biol (Beijing)* **12**, 1-6 (2017).

23.   Katayama, S., Sueyoshi, N., Inazu, T. & Kameshita, I. Cyclin-Dependent Kinase-Like 5 (CDKL5):
Possible Cellular Signalling Targets and Involvement in CDKL5 Deficiency Disorder. *Neural plasticity*
2020, 6970190 (2020).
24.   Gurgone, A.*, et al.* mGluR5 PAMs rescue cortical and behavioural defects in a mouse model of
CDKL5 deficiency disorder. *Neuropsychopharmacology* (2022).
25.   La Montanara, P.*, et al.* Cyclin-dependent-like kinase 5 is required for pain signaling in human
sensory neurons and mouse models. *Sci Transl Med* **12** (2020).
26.   Di Nardo, A.*, et al.* Phenotypic characterization of Cdkl5-knockdown neurons establishes
elongated cilia as a functional assay for CDKL5 Deficiency Disorder. *Neuroscience research* **176**, 73-78
(2022).
27.   Kontaxi, C., Davenport, E.C., Kind, P.C. & Cousin, M.A. Epilepsy-related CDKL5 deficiency slows
synaptic vesicle endocytosis in central nerve terminals. *bioRxiv*, 2022.2003.2015.484308 (2022).
28.   Halpain, S. & Dehmelt, L. The MAP1 family of microtubule-associated proteins. *Genome
biology* **7**, 224 (2006).
29.   Huang, J. & Zamponi, G.W. Regulation of voltage gated calcium channels by GPCRs and post-
translational modification. *Curr Opin Pharmacol* **32**, 1-8 (2017).
30.   Zamponi, G.W. Targeting voltage-gated calcium channels in neurological and psychiatric
diseases. *Nature reviews. Drug discovery* **15**, 19-34 (2016).
31.   Dietrich, D.*, et al.* Functional specialization of presynaptic Cav2.3 Ca2+ channels. *Neuron* **39**,
483-496 (2003).
32.   Parajuli, L.K.*, et al.* Quantitative regional and ultrastructural localization of the Ca(v)2.3
subunit of R-type calcium channel in mouse brain. *J Neurosci* **32**, 13555-13567 (2012).
33.   Sochivko, D.*, et al.* The Ca(V)2.3 Ca(2+) channel subunit contributes to R-type Ca(2+) currents
in murine hippocampal and neocortical neurones. *The Journal of physiology* **542**, 699-710 (2002).
34.   Zhang, J.F.*, et al.* Distinctive pharmacology and kinetics of cloned neuronal Ca2+ channels and
their possible counterparts in mammalian CNS neurons. *Neuropharmacology* **32**, 1075-1088 (1993).
35.   Lee, S.C.*, et al.* Molecular basis of R-type calcium channels in central amygdala neurons of the
mouse. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 3276-
3281 (2002).
36.   Schneider, T.*, et al.* Molecular analysis and functional expression of the human type E neuronal
Ca2+ channel alpha 1 subunit. *Recept Channels* **2**, 255-270 (1994).
37.   Soong, T.W.*, et al.* Structure and functional expression of a member of the low voltage-
activated calcium channel family. *Science* **260**, 1133-1136 (1993).
38.   Neumaier, F., Alpdogan, S., Hescheler, J. & Schneider, T. Protein phosphorylation maintains
the normal function of cloned human Cav2.3 channels. *J Gen Physiol* **150**, 491-510 (2018).
39.   Schneider, T., Alpdogan, S., Hescheler, J. & Neumaier, F. In vitro and in vivo phosphorylation
of the Cav2.3 voltage-gated R-type calcium channel. *Channels (Austin)* **12**, 326-334 (2018).
40.   Proft, J. & Weiss, N. G protein regulation of neuronal calcium channels: back to the future.
Mol Pharmacol **87**, 890-906 (2015).
41.   Helbig, K.L.*, et al.* De Novo Pathogenic Variants in CACNA1E Cause Developmental and
Epileptic Encephalopathy with Contractures, Macrocephaly, and Dyskinesias. *American journal of
human genetics* **103**, 666-678 (2018).
42.   Heyne, H.O.*, et al.* De novo variants in neurodevelopmental disorders with epilepsy. *Nature
genetics* **50**, 1048-1053 (2018).
43.   Happ, H.C. & Carvill, G.L. A 2020 View on the Genetics of Developmental and Epileptic
Encephalopathies. *Epilepsy Curr* **20**, 90-96 (2020).
44.   Steward, C.A.*, et al.* Re-annotation of 191 developmental and epileptic encephalopathy-
associated genes unmasks de novo variants in SCN1A. *NPJ Genom Med* **4**, 31 (2019).
45.   Mann, M. Functional and quantitative proteomics using SILAC. *Nature reviews. Molecular cell
biology* **7**, 952-958 (2006).
46.   Amendola, E.*, et al.* Mapping pathological phenotypes in a mouse model of CDKL5 disorder.
PloS one **9**, e91613 (2014).

47.   Weiergraber, M.*, et al.* Altered seizure susceptibility in mice lacking the Ca(v)2.3 E-type Ca2+
channel. *Epilepsia* **47**, 839-850 (2006).
48.   Weiergraber, M., Kamp, M.A., Radhakrishnan, K., Hescheler, J. & Schneider, T. The Ca(v)2.3
voltage-gated calcium channel in epileptogenesis--shedding new light on an enigmatic channel.
Neurosci Biobehav Rev **30**, 1122-1144 (2006).
49.   Weiergraber, M., Henry, M., Radhakrishnan, K., Hescheler, J. & Schneider, T. Hippocampal
seizure resistance and reduced neuronal excitotoxicity in mice lacking the Cav2.3 E/R-type voltage-
gated calcium channel. *Journal of neurophysiology* **97**, 3660-3669 (2007).
50.   Jones, L.P., Wei, S.K. & Yue, D.T. Mechanism of auxiliary subunit modulation of neuronal
alpha1E calcium channels. *J Gen Physiol* **112**, 125-143 (1998).
51.   Buraei, Z. & Yang, J. The ss subunit of voltage-gated Ca2+ channels. *Physiol Rev* **90**, 1461-1506
(2010).
52.   Muller, C.S.*, et al.* Quantitative proteomics of the Cav2 channel nano-environments in the
mammalian brain. *Proceedings of the National Academy of Sciences of the United States of America*
107, 14950-14957 (2010).
53.   Tai, C., Kuzmiski, J.B. & MacVicar, B.A. Muscarinic enhancement of R-type calcium currents in
hippocampal CA1 pyramidal neurons. *J Neurosci* **26**, 6249-6258 (2006).
54.   Bannister, R.A., Melliti, K. & Adams, B.A. Differential modulation of CaV2.3 Ca2+ channels by
Galphaq/11-coupled muscarinic receptors. *Mol Pharmacol* **65**, 381-388 (2004).
55.   Meza, U., Bannister, R., Melliti, K. & Adams, B. Biphasic, opposing modulation of cloned
neuronal alpha1E Ca channels by distinct signaling pathways coupled to M2 muscarinic acetylcholine
receptors. *J Neurosci* **19**, 6806-6817 (1999).
56.   Fraser, D.D. & MacVicar, B.A. Cholinergic-dependent plateau potential in hippocampal CA1
pyramidal neurons. *J Neurosci* **16**, 4113-4128 (1996).
57.   Page, K.M., Canti, C., Stephens, G.J., Berrow, N.S. & Dolphin, A.C. Identification of the amino
terminus of neuronal Ca2+ channel alpha1 subunits alpha1B and alpha1E as an essential determinant
of G-protein modulation. *J Neurosci* **18**, 4815-4824 (1998).
58.   Canti, C., Page, K.M., Stephens, G.J. & Dolphin, A.C. Identification of residues in the N terminus
of alpha1B critical for inhibition of the voltage-dependent calcium channel by Gbeta gamma. *J
Neurosci* **19**, 6855-6864 (1999).
59.   Fang, H.*, et al.* Role of alpha1 2.3 subunit I-II linker sites in the enhancement of Ca(v) 2.3
current by phorbol 12-myristate 13-acetate and acetyl-beta-methylcholine. *The Journal of biological
chemistry* **280**, 23559-23565 (2005).
60.   Buchanan, K.A., Petrovic, M.M., Chamberlain, S.E., Marrion, N.V. & Mellor, J.R. Facilitation of
long-term potentiation by muscarinic M(1) receptors is mediated by inhibition of SK channels. *Neuron*
68, 948-963 (2010).
61.   Park, J.Y. & Spruston, N. Synergistic actions of metabotropic acetylcholine and glutamate
receptors on the excitability of hippocampal CA1 pyramidal neurons. *J Neurosci* **32**, 6081-6091 (2012).
62.   Kawasaki, H., Palmieri, C. & Avoli, M. Muscarinic receptor activation induces depolarizing
plateau potentials in bursting neurons of the rat subiculum. *Journal of neurophysiology* **82**, 2590-2601
(1999).
63.   Williams, S.R. & Fletcher, L.N. A Dendritic Substrate for the Cholinergic Control of Neocortical
Output Neurons. *Neuron* **101**, 486-499 e484 (2019).
64.   Park, J.Y.*, et al.* A post-burst after depolarization is mediated by group i metabotropic
glutamate receptor-dependent upregulation of Ca(v)2.3 R-type calcium channels in CA1 pyramidal
neurons. *PLoS biology* **8**, e1000534 (2010).
65.   Kuzmiski, J.B., Barr, W., Zamponi, G.W. & MacVicar, B.A. Topiramate inhibits the initiation of
plateau potentials in CA1 neurons by depressing R-type calcium channels. *Epilepsia* **46**, 481-489
(2005).
66.   Brown, D.A. & Passmore, G.M. Neural KCNQ (Kv7) channels. *Br J Pharmacol* **156**, 1185-1195
(2009).
67.   Stocker, M., Hirzel, K., D'Hoedt, D. & Pedarzani, P. Matching molecules to function: neuronal
Ca2+-activated K+ channels and afterhyperpolarizations. *Toxicon* **43**, 933-949 (2004).

68.   Bloodgood, B.L. & Sabatini, B.L. Nonlinear regulation of unitary synaptic signals by CaV(2.3)
voltage-sensitive calcium channels located in dendritic spines. *Neuron* **53**, 249-260 (2007).
69.   Murphy, J.G.*, et al.* R-type voltage-gated Ca(2+) channels mediate A-type K(+) current
regulation of synaptic input in hippocampal dendrites. *Cell Rep* **38**, 110264 (2022).
70.   Terzic, B.*, et al.* Temporal manipulation of Cdkl5 reveals essential postdevelopmental
functions and reversible CDKL5 deficiency disorder-related deficits. *J Clin Invest* **131** (2021).
71.   Okuda, K.*, et al.* Comprehensive behavioral analysis of the Cdkl5 knockout mice revealed
significant enhancement in anxiety- and fear-related behaviors and impairment in both acquisition
and long-term retention of spatial reference memory. *PloS one* **13**, e0196587 (2018).
72.   Yennawar, M., White, R.S. & Jensen, F.E. AMPA Receptor Dysregulation and Therapeutic
Interventions in a Mouse Model of CDKL5 Deficiency Disorder. *J Neurosci* **39**, 4814-4828 (2019).
73.   Terzic, B.*, et al.* X-linked cellular mosaicism underlies age-dependent occurrence of seizure-
like events in mouse models of CDKL5 deficiency disorder. *Neurobiology of disease* **148**, 105176
(2021).
74.   Wang, H.T.*, et al.* CDKL5 deficiency in forebrain glutamatergic neurons results in recurrent
spontaneous seizures. *Epilepsia* **62**, 517-528 (2021).
75.   Colasante, G.*, et al.* In vivo CRISPRa decreases seizures and rescues cognitive deficits in a
rodent model of epilepsy. *Brain : a journal of neurology* **143**, 891-905 (2020).
76.   Tse, K., Puttachary, S., Beamer, E., Sills, G.J. & Thippeswamy, T. Advantages of repeated low
dose against single high dose of kainate in C57BL/6J mouse model of status epilepticus: behavioral
and electroencephalographic studies. *PloS one* **9**, e96622 (2014).
77.   Page, K.M.*, et al.* N terminus is key to the dominant negative suppression of Ca(V)2 calcium
channels: implications for episodic ataxia type 2. *The Journal of biological chemistry* **285**, 835-844
(2010).
78.   Stephens, G.J., Page, K.M., Bogdanov, Y. & Dolphin, A.C. The alpha1B Ca2+ channel amino
terminus contributes determinants for beta subunit-mediated voltage-dependent inactivation
properties. *The Journal of physiology* **525 Pt 2**, 377-390 (2000).
79.   Williams, M.E.*, et al.* Structure and functional characterization of neuronal alpha 1E calcium
channel subtypes. *The Journal of biological chemistry* **269**, 22347-22357 (1994).
80.   Artoni, P.*, et al.* Deep learning of spontaneous arousal fluctuations detects early cholinergic
defects across neurodevelopmental mouse models and patients. *Proceedings of the National*
Academy of Sciences of the United States of America **117**, 23298-23303 (2020).
81.   Negraes, P.D.*, et al.* Altered network and rescue of human neurons derived from individuals
with early-onset genetic epilepsy. *Molecular psychiatry* **26**, 7047-7068 (2021).
82.   Kamaludin, A.A.*, et al.* Muscle dysfunction caused by loss of Magel2 in a mouse model of
Prader-Willi and Schaaf-Yang syndromes. *Human molecular genetics* **25**, 3798-3809 (2016).
83.   Ruiz, A.*, et al.* Improvement of muscle strength in a mouse model for congenital myopathy
treated with HDAC and DNA methyltransferase inhibitors. *eLife* **11** (2022).
84.   Weiergraber, M.*, et al.* Altered thalamocortical rhythmicity in Ca(v)2.3-deficient mice.
Molecular and cellular neurosciences **39**, 605-618 (2008).
85.   Zaman, T.*, et al.* Cav2.3 channels are critical for oscillatory burst discharges in the reticular
thalamus and absence epilepsy. *Neuron* **70**, 95-108 (2011).
86.   Wormuth, C.*, et al.* Review: Cav2.3 R-type Voltage-Gated Ca(2+) Channels - Functional
Implications in Convulsive and Non-convulsive Seizure Activity. *Open Neurol J* **10**, 99-126 (2016).
87.   Gao, Y.*, et al.* Gene replacement ameliorates deficits in mouse and human models of cyclin-
dependent kinase-like 5 disorder. *Brain : a journal of neurology* **143**, 811-832 (2020).
88.   Gray, E.E.*, et al.* Disruption of GpI mGluR-Dependent Cav2.3 Translation in a Mouse Model of
Fragile X Syndrome. *J Neurosci* **39**, 7453-7464 (2019).
89.   Suzuki, T.*, et al.* Mutations in EFHC1 cause juvenile myoclonic epilepsy. *Nature genetics* **36**,
842-849 (2004).
90.   Benkert, J.*, et al.* Cav2.3 channels contribute to dopaminergic neuron loss in a model of
Parkinson's disease. *Nat Commun* **10**, 5094 (2019).

91.   Siller, A*., et al.* beta2-subunit alternative splicing stabilizes Cav2.3 Ca(2+) channel activity during continuous midbrain dopamine neuron-like activity. *eLife* **11** (2022).

92.   Shakeri, B., Bourdin, B., Demers-Giroux, P.O., Sauve, R. & Parent, L. A quartet of leucine residues in the guanylate kinase domain of CaVbeta determines the plasma membrane density of the CaV2.3 channel. *The Journal of biological chemistry* **287**, 32835-32847 (2012).

93.   Iannello, F. Non-intrusive high throughput automated data collection from the home cage. *Heliyon* **5**, e01454 (2019).

Figure 1

Figure 1. Identification and validation of Cav2.3 as substrate of CDKL5 kinase. A)
Volcano plot of differential phosphoprotein levels between WT and CDKL5 KO primary
neurons, obtained using SILAC based quantitative proteomics (5 embryos/genotype). Each
point represents one peptide; those with CDKL5 consensus motif RPXS/T are highlighted in
magenta. The x axis shows log2 transformed fold change in phosphopeptide levels between
genotypes and the y axis shows significance as -log10 transformed p value. Significant was
established as indicated by the dotted lines (p<0.01, fold change 1.5, one sample t test). B)
Top: Schematic depiction of Cav2.3 formed of α1E channel pore subunit and associated
proteins Cavβ and α2δ. CDKL5 phosphorylation site at α1E S15/14 is highlighted in green.
Bottom: Alignment of proximal N-terminus of mouse and human Cav2.3 and related human
Cav2.1 and 2.2 channel proteins. *S is the site of CDKL5 phosphorylation. C) Western blot of

HEK293 cells stably expressing human $\beta3/\alpha2\delta1$ subunits and transiently co-transfected with

human HA-α1E (Cav2.3: WT or S14A mutant) and HA-CDKL5 kinase domain (kd CDKL5: WT

or K42A mutant). S14 phosphorylation (pS14 Cav2.3) was detected with a custom made

phosphoantibody. D) Western blot of P20 cortices from WT and CDKL5 KO mice. E) Western

blot of iPSC-derived forebrain neurons from three CDD patients and related controls (parents)

at 6 weeks of differentiation. Gender and *CDKL5* mutation are specified [81]. F) Quantification

for immunoblot in C (**p<0.01 Kruskal-Wallis ANOVA & Dunn's test, n=11, 4 blots, 2

transfections). G) Quantification for immunoblot in D (**p<0.01 unpaired t test, n=9, 3 blots, 6

mice/genotype). H) Quantification for WB in E (**p<0.01 paired t test, n=6, 2 blots, 2 technical

replicates). For antibodies used see Materials and methods.

Figure 2

671 **Figure 2. Functional characterization of phospho-Ser14 Cav2.3 in HEK 293 cells.** A)

672 Representative depolarization-evoked current responses in HEK293 cells stably expressing

673 human β3/α2δ1 subunits and co-transfected with human Cav2.3 (WT α1E or S14A α1E) and

FLAG-CDKL5 full length (WT CDKL5). Construct combinations are presented in different colours. Traces show steps from -10 to +30mV from a holding potential of -80mV and Ba^{2+} as charge carrier B) Open channel inactivation tau (τ_{inact}) for Cav2.3 with (WT/WT, n=13-15) and without (S14A/WT, n=11-13) CDKL5 phosphorylation (*p<0.05, TW ANOVA, Fisher's LSD). Data were acquired using 100ms voltage steps from -80mV in +10mV increments every 10s. C) Individual data points for τ_{inact} at +20mV (*p<0.05, unpaired t test). D) WT or phophomutant S14A Cav2.3 inactivation in the same cell line in absence of CDKL5 (n=7-9; p>0.05 TW ANOVA). Ca^{2+} was charge carrier. E) Current voltage relationship for cells in B. Inset: comparison of current density at 0mV (inset, *p<0.05, unpaired t test). F) Normalised Cav2.3 current conductance and voltage dependence of inactivation for the same transfection conditions. Activation $V_{1/2}$, n: WT/WT -6±1 mV, 15; S14A/WT -6±1 mV,12; Inactivation $V_{1/2}$, n: WT/WT -61±3 mV,10; S14A/WT -62±2 mV,11 (p>0.05, unpaired t test). For inactivation protocol see Methods. Solid lines are Boltzman fits to the average data points G) Normalised Ba^{2+} currents in the $\beta3/\alpha2\delta1$-cell line or H) Ca^{2+} currents in the $\beta1/\alpha2\delta1$-cell line, co-transfected with $\alpha1E$ (WT or S14A), WT CDKL5 and muscarinic type 3 receptor (M3), in presence and absence of carbachol (CCh) 30 and 10 μM, respectively. HP= -100mV. I) IV curves and J) CCh-induced change in current amplitude for experiment in G (pS14 n=8, no pS14 n=7, TW ANOVA, Fisher's LSD). K) IV curves and L) CCh-induced change in current amplitude for experiment in H (pS14 n=7, no pS14 n=16 , TW ANOVA, Fisher's LSD). M) Conductance plots for experiment in G and N) corresponding change in activation ($V_{1/2}$: -5±0.6 *vs*. -10±1.9 mV, Kolmogorov-Smirnov). O) Conductance plots for experiment in H and P) corresponding change in activation ($V_{1/2}$: -4±0.5 *vs* -9±1 mV, Welch's t test). From I-P, 'pS14' refers to WT $\alpha1E$/WT CDKL5; 'no pS14' refers to S14A $\alpha1E$ /WT CDKL5 and $\alpha1E$ WT /K42R* CDKL5 conditions pulled together.

Figure 3

Figure 3. **Physiological properties and cholinergic neuromodulation of hippocampal CA1 neurons from WT and phosphomutant mice.** A) WB validation of CRISPR-generated Cav2.3 Ser15Ala mutant mice using a pS15 phosphoantibody in whole-brain lysates from 12-week-old WT, Heterozygous S15A and Homozygous S15A mice. B) Quantification of total

Cav2.3 (p>0.05 one way ANOVA) and C) relative phospho-S15 Cav2.3 (*p<0.05, **0.01 one-

way ANOVA, Fisher's LSD) expression (n=5, 2 blots, 3 mice/genotype). D) Normalised R-type

Ca^{2+} current in neurons from young mice of each genotype during a step from -60 to 0 mV in

presence of Na^+, K^+, Ca^{2+} and synaptic channel inhibitors. E) R-type current inactivation tau

(τ_{inact}) at maximal activation voltages (*p<0.05 TW RM ANOVA, Fisher's LSD) and F) average

IV curve for all P10-11cells recorded (p>0.05), (n=13 neurons, 4-5 mice/genotype). Data for -

10mV step is shown in the inset (p=0.06, unpaired t test). G) Membrane resting potential (V_{rest},

left) and input resistance (R_{input}, right) of adult neurons (WT n=13, HOM S15A n=17; 7

mice/genotype) in control conditions and after 10μM carbachol (CCh) application (**p<0.01,

paired t test). H) Input/output curves for the same neurons (WT n=9, HOM n=13, *p<0.05 TW

RM ANOVA) and percentage of cells with CCh-induced depolarization block (inset, WT n=13,

HOM n=21, *p<0.05, binomial test). I) Representative control action potential trains evoked in

WT (top) and HOM S15A (bottom) CA1 cells by 1s-long current injections. Membrane was

held at -65mV. J) Same cells as I upon application of CCh and representative depolarising

plateau potential (DPP), observed at some current injections in both genotypes; insets: fraction

of cells with at least one DPP in each genotype group (WT n=13, HOM n=21, p>0.05 binomial

test). K) Same cells as J at higher stimulation illustrating sustained depolarization and AP

block (top) or attenuation (bottom) towards the end of the stimulus and long-lasting large

amplitude afterdepolarizations (ADP). Like DPPs, these ADPs were present at some current

injections in a fraction of cells in both groups of mice (inset, WT n=13, HOM n=21, **p<0.01,

binomial test). L) Percentage of cells displaying sustained DPPs or ADPs upon stimulus

termination at each current injection step (non-linear fit comparison).

Figure 4

Figure 4. *In vivo* phenotypic characterization of Cav2.3 S15A mice. A) Average home cage night-time locomotion of WT and HOM S15A mice (N=9) per day and overall locomotion index (inset) in a three-week period (**$p<0.01$, Welch's t test). B) left: wheel rotations per hour during voluntary exercise activity for the same mouse cohort as A, over a two-week period; right: overall dark cycle average rotations in each group for three consecutive weeks ($p<0.05$ TW ANOVA, Fisher's LSD). C) Total distance travelled by WT (N=8) and HOMS15A (n=7) mice in an open field arena ($p<0.05$ at 30 min TW ANOVA, Fisher's LSD). D) Accelerating rotarod performance for the same cohort as C ($p>0.05$, TW ANOVA). Animal weights were equal (WT=25±4 g, HOM=25±5 g, $p>0.05$, unpaired t test). E) Social, non-social and total exploration times during a 10min three-chamber sociability test ($p<0.05$, unpaired t test). In habituation trials, chamber occupancy was equal between groups. F) Freezing behaviour during learning day 1 (left), associative memory test day 2+3 (middle) and memory extinction test days 4-7 (right), in a fear-conditioning experiment with tactile and olfactory cues (*$p<0.05$ or **0.01, TW RM ANOVA, Fisher's LSD or unpaired t test). G) left: two-week baseline ECoG spectral analysis and total power (inset) in WT (N=8) and HOMS15A (N=10) 9-12 week-old mice; corresponding coastline index (middle) and kurtosis (right) ($p>0.05$, TW ANOVA and unpaired t tests). No sex differences were observed. H) left: behavioural scoring of kainic acid (KA)-induced seizures (WT N=7, HOMS15A N=10, $p>0.05$, TW ANOVA) using repetitive 5mg/kg i.p. injections (inset); right: latency to stage 5 tonic clonic seizures per gender (males, $p>0.05$; females *$p<0.05$, unpaired t test). I) Corresponding cumulative ECoG coastline during KA seizures in females (WT N=3, HOM S15A N=4, *$p<0.05$, TW ANOVA).

 Supplementary Figures

Figure S1

FigureS1. Full-length CDKL5 phosphorylates Cav2.3. A) Western blot of HEK293 cells
stably expressing human $\beta3/\alpha2\delta1$ subunits and transiently co-transfected with human WT HA-
$\alpha1E$ (Cav2.3) and FLAG-CDKL5 full length (FL CDKL5: WT, K42A, K42R, K42R*+DFG-AFG
kinase dead mutants). (-) transfection condition excludes CDKL5. The K42A mutation in FL-
CDKL5 retains some kinase activity on $\alpha1E$, while K42R and K42R* mutants are fully inactive
and were thus selected for electrophysiology experiments. B) Quantification of relative
phospho-$\alpha1E$ (pS14 Cav2.3) levels for the experiment in A (**p<0.01, Brown-Forsythe &
Welch ANOVA, Dunnet's test; n=4-6, 3 blots, 2 transfections).

Figure S2

Figure S2. Functional characterization of phospho-Ser14 Cav2.3 in HEK 293 cells expressing β1 accessory subunit. A) Representative depolarization-evoked current responses in HEK293 cells stably expressing human $\beta1/\alpha2\delta1$ subunits and co-transfected with human Cav2.3 (WT α1E or S14A α1E mutant) and FLAG-CDKL5 full length (WT CDKL5 or K42R*+DFG-AFG kinase dead CDKL5 mutant). Construct combinations are presented in different colours. Traces show steps from -10 to +20mV from a holding potential of -100mV and Ca^{2+} as charge carrier. B) Open channel inactivation tau (τ_{inact}) for Cav2.3 with (WT/WT, n=28-32) and without (S14A/WT, n=34-37; WT/K45R, n=25-26) CDKL5 phosphorylation. WT/WT vs. S14A/WT: *p<0.05, **p<0.01 as indicated; WT/WT vs. WT/K42R*, p<0.01 at +40mV, p<0.05 at -10,10,20,30mV, TW ANOVA, Fisher's LSD. Some of the cells included in this plot co-expressed M3 receptors; this was *per se* not found to affect inactivation kinetics.

C) Individual data points for τ_{inact} at +10mV (p<0.05, unpaired t tests). D) Current-voltage

relationship for experiments in A (WT/WT, n=15; S14A/WT n=15-17; WT/K42R* n=12-15) and

current density at +10mV (inset, p>0.05, one way ANOVA). Data was acquired with 100ms

voltage steps from -100mV in +10mV increments every 10s. E) Normalised Cav2.3

conductance and voltage dependence of inactivation for the same three transfection

conditions. Activation $V_{1/2}$, n: WT/WT -3±1 mV, 14; S14A/WT -3±1 mV,17; WT/K42R* -2±1

mV, 14; Inactivation $V_{1/2}$, n: WT/WT -66±2 mV,14; S14A/WT -63±1 mV,14; WT/K42R* -65±2

mV,11 (p>0.05, one way ANOVA). For inactivation protocol see Methods. Solid lines are

Boltzman fits to the average data points. F) Inactivation recovery time using a double pulse

protocol with variable inter-pulse recovery time (inset). Recovery tau, n: WT/WT 265±21 ms,

4; S14A/WT 263±21 ms, 5; WT/K42R* 290±15 ms, 4, p>0.05 Kruskall-Wallis, Dunn's test). G)

Example time course of Cav2.3 current recording at 0mV in the $\beta1/\alpha2\delta1$-stable cell line co-

transfected with α1E WT, CDKL5 and muscarinic receptor type 3 (M3). PKC activation with

PMA did not occlude CCh enhancing effect. H) Relative change in current amplitude for same

experiment as G) with either M1 or M3 receptors. No muscarinic inhibition was observed. I)

Representative Cav2.3 current traces at +10mV in $\beta1/\alpha2\delta1$ stable cell line co-transfected with

α1E (WT or S14A), CDKL5 (WT) and dopamine receptor type 2 (D2). Quinpirole 100nM (QP)

application resulted in reliable and reversible inhibition of WT and S14A currents. J) Quantified

quinpirole inhibition of Cav2.3 with (n=7) or without pS14 (n=12-13) (p>0.05 TW ANOVA,

Fisher's LSD). Data for α1E S14A phosphomutant and K42R* CDKL5 conditions (no

phosphorylation) were pooled together.

A

B

C

D

E

F

G

H

Figure S3. Additional CA1 neuron responses to carbachol and other intrinsic properties

in adult WT and HOM S15A mice. A) Relative changes in membrane potential (V_{rest}) and

resistance (R_{input}) upon 10 μM CCh application in WT (13) and HOM S15A (17) neurons (7

mice/genotype, p>0.05, unpaired t tests). B) CCh-evoked medium duration afterdepolarization

(ADP) measured 100ms post 1s-long stimulus (WT n=13, HOM n=21; p=0.07 TW ANOVA;

*p<0.05 Fisher's LSD). C) Spike attenuation in control (dotted lines) and upon CCh-induced

depolarization (markers) shown as ratio between last and first spike amplitude in 1s train (WT

n=13, HOM n=19, TW ANOVA; *p<0.05 Fisher's LSD). D) Medium and slow duration

afterhyperpolarizations (AHPs) quantified at the peak and 500ms post stimulus, respectively

(inset). AHPs evoked at -65mV by a 100Hz burst of somatic current injections were equal in

WT (13) and HOM S15A (18-19) mice (mAHP, -1.5 *vs* -1.2mV; sAHP, -0.2 *vs* -0.3mV, p>0.05,

unpaired t test). CCh suppression of burst-evoked AHPs (**p<0.01 paired t tests) was

unchanged between genotypes (mAHP ~Δ1.5mV, sAHP ~ Δ 1mV, p>0.05 unpaired t test). E) Example traces of isolated small-conductance Ca^{2+}-activated SK currents and full inhibition by d-tubocurarine (dTC, grey). F) dTC-sensitive current amplitude (left) and charge (right) were not different between groups (WT n=8, HOM n=10 cells, 4 mice/genotype, p>0.05 unpaired t tests). G) Example traces of spontaneous excitatory postsynaptic currents in WT and HOMS15A mice H) Distribution of measured sEPSC properties for each genotype (WT: 1422-34 events, n=8 cells, HOM: 1455-65 events, n=7 cells; mean amp=14.0±0.8 *vs* 14.7±0.5 pA; instantaneous frequency=4.0±0.4 *vs* 4.3±0.3 Hz; N=3 mice/genotype; p>0.05, unpaired t tests).

Figure S4

Figure S4. Behavioural characterization of Cav2.3 S15A phosphomutant mice. A) left: total time in the central zone (CZ) of the open field arena in WT (N=8) and HomS15A (N=7) mice (30 min test) (p>0.05, TW RM ANOVA; p=0.08 at 30 min, Fisher's LSD); right: mean displacement velocity (p>0.05, TW RM ANOVA). B) Percentage of successful spontaneous alternations (left) and total arm entries (right) in the Y-maze test show no deficits in arm recognition and exploration in phosphomutants (p>0.05, unpaired t test). C) Acquisition phase of the Barnes maze test with improved performance for both groups in the first 4 training days: distance travelled (left), latency (middle) and number of errors committed (right) during the 1st

845   visit to the escape box were not significantly different ($p > 0.05$, TW RM ANOVA). D) memory

846   assessment on probe day 5 compared using the same parameters as in C ($p > 0.05$, unpaired

847   t tests). E) Quantification of total time spent around the expected location of the goal box (left)

848   and total distance travelled (right) ($p > 0.05$ unpaired t tests).

Materials and Methods

Mouse handling and mutant generation

Animals were housed in a controlled environment with 12-hour light/dark cycle. They were fed *ad libitum* and used in accordance with the principles of the "3Rs" and the Animals (Scientific Procedures) Act 1986 of the United Kingdom. Protocols and procedures were approved by institutional evaluations. With the exception of some *in vivo* experiments, mice were housed in groups. Mouse strains were backcrossed into C57 Bl/6 (Jackson) and none of the experimental mice were immunocompromised. Male and female mice aged embryonic day (E)16.5 to postnatal 30 weeks were used as specified.

CDKL5 KO animals were a kind gift from Cornelius Gross [46]. Mutant Cav2.3 Ser15Ala animals were created in-house using the CRISPR/Cas9 system. The guide contained the sequence 5'AGGCUCAGGCGAUGGAGACU3'. This replaced the original DNA sequence of 5' AGGCGCAGGCGATGGAGACT 3' in CACNA1E. Genetically altered C57Bl6 embryos were obtained using electroporation as part of Francis Crick Institute's Genetic Modification Services. Genotype was determined by in-house PCR or Sanger sequencing (Source Bioscience).

SILAC phosphoproteomics

Sample preparation

Mouse cortical neurons were cultured from individual male E16.5 embryos from a heterozygous Cdkl5 female. Cdkl5 +/Y (WT) or -/Y (KO) genotype of embryos was determined afterwards. Neurons were plated at a density of 8x10^6 cells per 10 cm dish coated with 60 µg/ml poly-D-lysine (Sigma) and 2.5 µg/ml laminin (Sigma). Cells were grown in neurobasal media free of L-Arginine and L-Lysine (Invitrogen) supplied with either Lysine 8 (U-13C6, U-15N2) and Arginine 10 (U-13C6, U-15N4) or Lysine 4 (4, 4, 5, 5-D4) and Arginine 6 (U-13C6). L-Proline (200mg/ml) was supplemented to prevent Arginine to Proline conversion. At DIV4, 1 µM Ara-C was added to inhibit glia growth. For every embryo, one dish was labelled with K8R10 (heavy) and one dish was labelled with K4R6 (light).

At DIV12, SILAC labelled primary neurons from 3 Cdkl5 WT and 3 Cdkl5 KO animals were lysed in lysis buffer containing 20 mM Tris pH7.5, 100 mM NaCl, 10 mM MgCl$_2$, 0.25% IGEPAL NP40, 0.5 mM DTT, 1x Protease Inhibitor Cocktail, 1 µM Okadaic Acid. Each sample was checked for SILAC label incorporation and Arginine to Proline conversion. Lysates were mixed to obtain 3x heavy WT/light KO and 3x heavy KO/light WT samples containing 1 mg total

protein. Proteins were digested overnight with Sequencing Grade Modified Trypsin/Lys-C (Promega) at 37^0C, desalted using Sep-Pak Classic C18 Cartridges (Waters) and vacuum dried completely. Samples were enriched for phosphopeptides using Titansphere titanium dioxide beads (GL Science) in batch mode with the following buffers: Loading buffer: 80% ACN, 5% TFA, 1M Glycolic Acid, Wash buffer 1: 80% ACN 1% TFA, Wash buffer 2: 10% ACN, 0.2% TFA, Elution buffer 1: 1% Ammonium Hydroxide, Elution buffer 2: 5% Ammonium Hydroxide. Phosphopeptides were cleaned up using C18 Stage Tips with a centrifuge protocol and vacuum dried completely. Samples were stored at -80^0C until required for analysis by mass spectrometry.

Mass spectrometry

Each sample was resuspended in 35 µl 1% TFA, sonicated for 15 minutes and injected 3 times (10 µl per injection). Peptide mixtures were separated on a 50 cm, 75 µm I.D. Pepmap column over a 3 hr gradient and eluted directly into the mass spectrometer (Orbitrap Velos). Xcalibur software was used to control data acquisition. The instrument ran in data dependent acquisition mode with the top 10 most abundant peptides selected for MS/MS by CID, MSA or HCD fragmentation techniques (one fragmentation technique per replicate).

Data processing was performed using the MaxQuant bioinformatics suite (v 1.3.0.5) and protein database searching was done by the Andromeda search engine using a Uniprot database of *Mus musculus* proteins amended with common contaminants. Default search settings were used including an FDR of 1% on the phosphosite, peptide and protein level and matching between runs for peptide identification. K4R6 and K8R10 were set as labels with a maximum of 3 labeled AAs. Phosphorylation at serine, threonine or tyrosine residues, oxidation (Met) and protein N-terminal acetylation were set as variable modifications. Carbamidomethylation of Cys-residues was set as a fixed modification.

Data was further processed in Perseus v.1.6.15.0. Phospho (STY) sites from the six experimental samples in the MaxQuant search were loaded and site tables expanded. Potential contaminants and reversed hits were removed. Normalized H/L ratios were log2 transformed and filtered for rows containing at least three valid values. Potential CDKL5 substrates were annotated using the R-P-X-pS/pT motif [19]. A one-sample t-test was performed comparing the normalized ratios to the null hypothesis and visualized as a volcano plot.

Mutagenesis

Point mutations were introduced using QuickChange site-directed mutagenesis (Agilent). Complimentary primers contained the desired mutations flanked by at least 18 bp. To ensure mutagenesis efficiency for each construct, two separate PCR reactions with each primer were

set up: 0.2 μM primer, 100 ng template DNA, 0.2 mM dNTP mix, 1x Pfu buffer and 2.5 U Pfu

Ultra HF. The PCR programme was run 4x, after which complementary samples were

combined and the PCR run repeated 20x. PCR products were treated with DpnI for 1hr at

37^0C and transformed into XL-10 Gold competent cells. At least 4 colonies were used for each

DNA miniprep (QIAprep, Qiagen) and mutations were confirmed by Sanger sequencing

(Source Bioscience). The constructs used were as follow: 1) HA-tagged human α1E subunit

(gene bank L27745.2, gift from L. Parent, Université de Montréal), in a commercial vector

under control of CMV promoter [92]. 2) The N-terminal FLAG-tagged full-length human $CDKL5_{107}$

and N-terminal HA-tagged human $CDKL5_{1-352}$ kinase domain construct kinase as previously

described [19].

Cell line maintenance and transfection

Tetracycline-inducible HEK293 cells expressing human Cavβ3 and α2δ1 subunits were

obtained from SB Drug Discovery (Glasgow, UK). HEK293 cells stably expressing human Cav

β1b and α2δ1 subunits were kindly provided by Andrew Powell (GSK Ltd., Middlesex, UK).

Naïve HEK293T cells were supplied by Francis Crick Institute's Cell Services.

All cells were cultured in high glucose DMEM with 10% Fetal Bovine Serum (tetracycline free,

Clontech or Gibco) and penicillin/streptomycin 50units/ml-50 mg/ml. They were passaged

using TrypLE Express and maintained under selection antibiotics as appropriate: HEKβ3/

α2δ1, Zeocin 300mg/ml (Invivogen) & Blasticidin 5mg/ml (Invivogen); HEKβ1b/α2δ1,

Puromycin 1mg/ml and Hygromycin-B 200mg/ml. Cell culture reagents were obtained from

Gibco/Invitrogen unless specified.

Selection antibiotics were removed before transfection and doxycycline 1μg/μl added in the

case of β3-expressing cells. Transient transfection was carried out using X-tremeGENE™ 9

reagent (Roche) following manufacturer's instructions. Total DNA was 1.5-1.6μg. For Western

blot experiments, human α1E-HA subunit and CDKL5 (HA-kinase domain or FLAG-full length)

were co-transfected in a 1:1 ratio, substituting kinase for empty pcdna3 vector in some cases.

For electrophysiological recordings α1E-HA, FLAG-CDKL5 full length and GFP (pcdna3) were

co-transfected at a ratio of 10:4:1 (HEK β3 cell line) or 8:4:1 (HEKβ1b cell line). To study

muscarinic regulation, human α1E-HA, FLAG-CDKL5, muscarinic receptor type 3/1

(CHRM3/1, Addgene) and GFP were co-expressed at a ratio of 8:4:3:1. For dopamine

modulation experiments, human α1E-HA, FLAG-CDKL5 and dopamine D2 receptor (gfp-

DRD2, Addgene) were co-transfected at a ratio of 2:0.75:1.

Western Blotting

Three different types of sample were used for Western blot experiments: 1) HEK293 cells 48 hr post-transfection, 2) Mouse brain tissue collected after cervical dislocation, followed by dissection and snap-freezing in liquid nitrogen, 3) Human iPSC derived neurons obtained as frozen pellets from Cleber Trujillo (UCSD, USA). Protocols for iPSC generation, patient *CDKL5* mutations and related control details are described in [81]. All lysates were prepared in 1x/2x sample buffer (Invitrogen) with 0.1-0.2 M dithiothreitol (Sigma). This was followed by sonication, centrifugation at 13000g and denaturation at 70^0C, 10 min. Protein was loaded onto NuPAGE 8% Bis-Tris polyacrylamide gels (Invitrogen) for electrophoresis and subsequently transferred to polyvinylidene difluoride membranes (Millipore) for 15-22 hr at 20V in 10% MetOH, Tris Glycine buffer. Membranes were blocked for 30 min in 5-10% skimmed milk and incubated with primary antibodies overnight at 4^0C or 1hr at room temperature (RT). Secondary incubation in horseradish peroxidase-conjugated (HRP) antibodies was for 1hr at RT. Chemiluminescence signals were detected using Amersham ECL (Cytiva) and Amersham Imager (GE Healthcare). Analysis was perfomed using Fiji 2.1. Phospho-S14/15 channel levels were measured as the ratio between pS14/15 Cav2.3 signal and total Cav2.3 signal. Total channel levels are expressed as the ratio over loading control GAPDH. Each replicate in a blot constitutes a data point normalized to the average internal control signal in each blot. At least two independent immunoblot experiments with technical replicates were used for quantification, as specified in figure legends.

Primary antibodies used in each sample were: mouse anti-HA 1:2000 (Biolegend, 901513; HEK293), rabbit anti Cav2.3 N-terminus 1:250 (Covalab, custom 1, HEK293 & brain), rabbit anti pS15 Cav2.3 1:500 (Covalab, custom 2, HEK293, brain & human neurons), mouse anti Cav α1E 1:1000 (Synaptic Systems, 152411, human neurons), rabbit anti CDKL5 1:2000 (Atlas, HPA002847, HEK293), mouse anti GAPDH 1:50000 (Abcam, ab8245, brain & human neurons). Custom made Cav2.3 Ser15 phosphospecific antibodies were raised by immunizing rabbits with peptide PRPG(pS)GDGDSDQSRNC. Phosphorylated Cav2.3 antibody was obtained after double purification. First, they were selected against non-phosphorylated peptide PRPGSGDGDSDQSRNC. This antibody is used as total Cav2.3 antibody (custom 1). Next the eluted fraction was purified again by binding to PRPG(pS)GDGDSDQSRNC -linked beads, obtaining pSer15 Cav2.3 (custom 2). Secondary antibodies used were: donkey anti rabbit HRP 1:10000 (Jackson, 711-035-152), donkey anti mouse HRP 1:10 000 (Jackson, 715-035-151).

In vitro **electrophysiology**

HEK293 cells

Cells were split 24hr after transfection and replated on glass coverslips and allowed to recover

overnight. Whole-cell patch clamp recordings in isolated cells were performed 48-72 hr post-

transfection at room temperature. Transfected cells were identified by GFP fluorescence. Data

were sampled at 20kHz and filtered at 1-2 kHz. Cells were continuously perfused (1ml/min)

with extracellular solution containing (in mM): NaCl 120, TEACl 20, HEPES 10, glucose 10,

KCl 5, MgCl$_2$ 1, CaCl$_2$ or BaCl$_2$ 5, pH 7.4 with NaOH. Pipettes (2.5-3.5 ΩM) were pulled from

borosilicate glass (1.2x0.69mm, Harvard Bioscience) and filled with the following intracellular

solutions adapted from [38] to minimize current run-down (in mM): CsMeS 140, EGTA 5, MgCl$_2$

0.5, MgATP 5, HEPES 10, for GPCR experiments; CsMethanesulfonate 125, TEACl 5, EGTA

5, MgATP 5, Na$_3$GTP 0.3, Na-phosphocreatine 5, Na-pyruvate 2.5, HEPES 10, pH 7.25, 285-

295 mOsm, all other recordings. Upon seal break, currents were evoked every 5/10s with a

step from -80/-100mV (holding potential, HP) to 0/+10mV. After an initial run-up period, steady

state current voltage (IV) relationships were obtained with 100ms depolarising test steps to

+60/+80 in 10mV increments every 10s. For voltage dependence of inactivation the protocol

consisted of a 1s pulse from -140 to +20mV in 10 mV increments, followed by a 50 ms test

pulse to 0 mV, every 10s. Peak currents (I$_{peak}$) in response to test voltages (V$_{test}$) were used

to plot IVs and measure the voltage dependence of activation and inactivation. Reversal

voltage (E$_{rev}$) was estimated by extrapolating a linear fit to the IV curve from +20-+40 mV.

Conductance (g) was calculated using the equation (g=(I$_{peak}$)/(V$_{test}$-E$_{rev}$)). Voltage of half-

maximal activation or inactivation (V$_{1/2}$) were measured by fitting with a Boltzmann equation

(Y= A+(B-A)/(1+exp((V-V$_{1/2}$)/K)) where Y is the conductance or the current, and A and B are

the minimum and maximum amplitudes of the fit. Time course of open state inactivation (τ$_{inact}$)

was estimated by fitting a single exponential function to current decay in response to test

pulses. The average of 3 fits at each voltage was used. For all transfection conditions, cells

with outward currents >40% of the peak current at +10mV or with τ$_{inact}$ that did not decay

uniformly with depolarization were excluded from analysis. Similarly, cells were not used for

current density or τ$_{inact}$ comparisons if run-down was >30% of the maximal current recorded in

the experiment as this is known to affect inactivation time [38]. To examine recovery from

inactivation, an inactivating pulse to +10mV of 0.8-1s (pulse 1), was followed by a variable

recovery period (Δ 150-200ms) and a test pulse to +10mV (pulse 2), at 0.1-0.3 Hz. Time

course of recovery was calculated using the ratio of I$_{peak}$ in response to pulse 2/pulse 1 plotted

against duration of recovery period and fitted with single exponential equation. Carbachol,

quinpirole and PMA were bath at ~3 ml/min whilst activating Cav2.3 currents with brief

monitoring (25ms) test steps to 0 or +10mV, every 2, 5 or 10s, depending on speed of run-

down. Upon steady state, IVs were obtained. Cells included in the analysis showed no

significant changes in series resistance and a consistent shift in activation V$_{1/2}$, with current

amplitude matching between IV and brief monitoring steps. Leak and capacitance subtraction

was applied to all recordings using -P/5 protocol and series resistance compensated from 75-
95% to keep voltage error <5mV.

1050

Ex vivo **electrophysiology**

Hippocampal slices

Mice older than P10 were anaesthetized by intraperitoneal injection of ketamine (80mg/kg)
and xylazine (10mg/kg). To avoid bias, the genotype was unknown at the time of experiment.
Animals were decapitated and the brain transferred to ice cold artificial cerebrospinal fluid
(standard aCSF) containing (in mM): NaCl 125, NaH_2PO_4 1.25, $NaHCO_3$ 26, KCl 2.5, glucose
25, $MgCl_2$ 1, $CaCl_2$ 1, saturated with carbogen. Cerebellum and olfactory bulb were discarded.
Coronal or transverse hippocampal slices (300µm) were prepared with a Leica VTS 1200S
vibratome and transferred to an immersion storage chamber containing bubbled aCSF at 35^0C
for 30 min. Slices were allowed to recover for a further 30 min at room temperature (RT) before
electrophysiology experiments. Somatic whole-cell patch clamp recordings were obtained
from CA1 pyramidal neurons using borosilicate electrodes (3-4.5 MΩ). Series resistance (R_s)
and input resistance (R_{in}) were monitored throughout recordings using a 100ms, -5mV step
from -60mV to evoke passive membrane responses. R_s was calculated from the amplitude of
the capacitive transient and R_{in} from the steady state currents. These were sampled at 20kHz
and filtered at 10kHz. The maximum change in Rs permitted was 25%.

R-type Ca^{2+} currents - Recordings were obtained from young neurons (P10-11) no longer than
6h post slicing using the blind technique. Currents were recorded at RT from P10-11 coronal
slices during bath application of a modified aCSF solution (in mM): NaCl 115, TEACl 10,
$NaHCO_3$ 26, KCl 2.5, NaH_2PO_4 1.25, glucose 25, $MgCl_2$ 1, $CaCl_2$ 2; and in presence of the
following inhibitors (in µM): TTX 0.5, gabazine 1, APV 25, NBQX 5, 4AP 5, Nifedipine 10
(Tocris), w-conotoxin GVIA 2, w-agatoxin IVA 0.2, w-conotoxin MVIIC 2 (Alomone) and BSA
1mg/ml. Drug stock solutions were prepared in water and stored at -20^0C. External solution
was constantly bubbled and re-circulated. Slices were exposed to ion channel inhibitors for at
least 20 min before voltage clamp acquisition. Patch pipettes were filled with (in mM):
CsMethanesulfonate 130, TEACl 10, Na-phosphocreatine 5, $MgATP_2$ 4, Na_3GTP 0.3, HEPES
10, EGTA 2 and biocytin 0.2% pH 7.3 with CsOH. After 10 min stabilization in the whole-cell
configuration, IV curves were obtained with voltage steps from -80 to +40, HP=-70mV. A -P/5
leak and capacitance subtraction protocol was used and R_s was compensated 65-85%, to
minimize voltage error. Capacitance (C_m) was calculated using equation: C_m [pF] = τ_m
[ms]/R_s[MΩ], where τ_m is the decay time constant of capacitative transients. Time course of
open state inactivation was estimated by iterative single exponential fit to current decay. A
positive correlation was found between τ_{inact} and C_m for C_m<80pF, thus this was established as

the cut off cell size criteria for inclusion in τ_{inact} comparisons. Cells with large cells outward

currents at depolarising voltages (>50% of the peak current at +10mV) were also excluded.

Small-conductance Ca^{2+}-activated (SK) currents and excitatory postsynaptic currents (EPSC)-

Recordings were obtained from adult neurons (5-6 weeks) in transverse slices. SK currents

were measured as tail currents following a 100ms step from -50 mV to +10mV (0.3 Hz) to

engage VGCCs. Standard aCSF (CaCl$_2$ 2mM, RT) was supplemented with (in µM): TTX 0.5,

TEA 1000 and XE991 5, to isolate SK currents by inhibiting temporally overlapping voltage-

gated Na$^+$ and K$^+$ conductances. The intracellular solution was (in mM): KGluconate 135, KCl

10, Na-phosphocreatine 10, MgATP$_2$ 2, Na$_3$GTP 0.3, HEPES 10, pH 7.3, supplemented with

8-cloro-phenylthio cAMP 50µM, to inhibit the slow afterhyperpolarizing current. SK current

identity was confirmed at the end of all experiments by reversible d-tubocurarine (dTC, 50µM)

inhibition. dTC-subtracted currents were used for quantification. Traces were sampled at 5

kHz and filtered at 1kHz. SK current amplitude was measured at the peak of the after-current

and charge as the integral 500ms post stimulus. EPSCs were isolated at -70 mV in standard

aCSF (1) (CaCl$_2$ 2.5mM, 30-32^{0}C) with gabazine 1 µM and Kgluconate-based internal solution.

Traces were sampled at 10kHz and filtered at 2kHz. Upon 5 min stabilization in whole-cell, a

2 min gap free trace was analysed using template-based event detection and threshold match

>3.

Current clamp recordings - Recordings were obtained from adult neurons (5-6 weeks) in

coronal slices in standard aCSF (2mM Ca^{2+}, 30-32^{0}C) and Kgluconate-based internal solution.

Membrane potential (V$_{rest}$) was measured throughout the recording with 0 current injection and

maintained around -65mV during stimulation. Input-output relationships were obtained with 1s

current injections from -100 to 450 pA in 50 pA increments. Traces were sampled at 10kHz

and filtered at 5kHz. Action potentials were identified with threshold-based event detection

(0mV) and amplitude measured from V$_{rest}$. Spike attenuation during the train was calculated

as ratio between last and first spike amplitudes. Depolarizing plateau potentials (DPPs) were

defined as sustained depolarizations >20mV upon stimulus termination. Medium duration

afterdepolarizations (ADPs) were measured 100ms post-1s step depolarization. Firing

patterns were compared using binomial tests or comparisons of non-linear least squares

regression fits to the data distribution. The effect of carbachol on the medium and slow

afterhyperpolarizations (mAHP and sAHP) was used as positive control for muscarinic

receptor activation. AHPs were evoked at -65mV by a burst of five 2nA, 2ms somatic current

injections at 100Hz every 20s and measured at the peak and 500ms post stimulus

respectively.

All patch clamp data was obtained with Multiclamp 700B, Digidata 1440A and pClamp^{TM} 10

software (Molecular Devices). Data were analysed using Clampfit 10.7/11.2 and OriginPro

9.8. Liquid junction potential was not corrected. Statistical analysis was performed in

GraphPad Prism 9. Student's T tests and one- or two-way ANOVA (TW ANOVA) with Geisser-Greenhouse correction followed by Fisher's LSD test were used for most statistical comparisons. Repeated measures (RM) ANOVA was used as appropriate if there were no missing values. Data are presented as mean ± S.E.M. Box and whiskers (min to max) or violin plots are used to illustrate data spread and frequency distribution respectively; percentile 25/75, median and mean are indicated by lines/markers.

In vivo electrophysiology

Chronic electrocorticogram recordings

Surgical procedures were performed in adult male and female mice (8-10 weeks old) using a stereotaxic frame (Kopf) under isoflurane anaesthesia and temperature control. Each animal was subcutaneously implanted with an electrocorticogram (ECoG) transmitter (A3022B-CC-B45-B, Open Source Instruments, Inc.). The recording electrode was placed above the visual cortex (AP -2, ML 1.5) and the ground electrode in the contralateral prefrontal hemisphere (AP 1.8, ML 1.5). Implanted WT and Hom S15A animals (total 18) were housed individually and no drug treatment was given. The ECoG was recorded wirelessly (sampling frequency 512 Hz, band-pass filter 1-160 Hz) for two weeks using software from Open Source Instruments, Inc. Simultaneous video recordings were obtained (6x/hour) using IP cameras from Microseven (https://www.microseven.com/index.html). The coastline, kurtosis and power spectrum analysis (1-160Hz) were performed using Python. Researchers were blinded to animal genotype during data acquisition and analysis.

Kainic acid susceptibility

At the end of the chronic recordings, low dose 5mg/kg kainic acid (Tocris Bioscience) dissolved in sterile saline was administrated intraperitoneally every 30 minutes to assess brain susceptibility to increasing dosage of chemo convulsant. The clock started at the first injection from 0 minute. Seizure severity was initially assessed while the experiment was ongoing at 10-minute intervals, using a modified Racine scale: 1. immobility; 2, hunched position with facial jerks, 3. rearing and forelimb clonus, 4. persistent rearing and forelimb clonus, falling, 5. generalized tonic clonic convulsions, or wild jumping [76]. These assessments were confirmed by re-analyzing the video recordings. Latency measures the length of time from the start of the experiment to the first indication of generalized seizure (Stage 5 on Racine scale). Both kainic acid inductions and analysis were performed by a researcher blinded to the genotypes. A total of 17 implanted mice were used for susceptibility experiments, one male mouse was removed from the study due to surgical complications with the transmitter. In a subset of these where transmitter performance was not compromised, ECoG recordings during seizure

1156 inductions were recovered and used for quantification. Traces were analysed using semi-
1157 automated seizure detection as described previously [75]. The cumulative coastline was
1158 calculated from when the first epileptiform activity was detected until the mice reached
1159 generalized seizure (stage 5).

1160

Behavioural analysis

Home Cage monitoring

Two separate cohorts of WT and homozygous S15A Cav2.3 mice of both sexes and aged
between 12-16 weeks were used. Animals were housed individually and monitored using the
Digital Ventilated Cage system (DVC®,Techniplast, Italy). The DVC consists of standard size,
barcoded IVC units equipped with a 3x4 arrangement of cage-floor electrodes. Capacitance
sensing-based detection of animal position allows for real-time tracking of locomotion activity
24/7. Following a 2-week-acclimatization period, baseline activity was recorded and analysed.
Rotating wheels (Ø 4.4 in) were then introduced, and animals allowed to familiarize
themselves for another week before data collection. All activity metrics were calculated using
the DVC Analytics 3.4.0 platform [93]. For locomotion index, the average signal of all 12 cage
electrodes was used. Data were grouped in 60 min bins for each day.

Behaviour tests

Behaviour experiments were conducted in 8-week-old WT and homozygous S15A Cav2.3
littermates of both sexes, housed under an inverted light cycle. Mice were handled daily for
one week prior to experiments. All tests were video recorded for off-line analysis. Tracking
data were analysed using Ethovision XT 15 (Noldus Information Technology, Netherlands)
equipped with three-point (nose, body centre, and tail) detection settings. One cohort of mice
was used for all 7 tests, carried out in the order described below. Mice were allowed to
habituate to the testing room for at least an hour before the behaviour test.

Hind-Limb clasping – Mice were suspended by the middle of the tail and lifted 15 cm above
the ground; the extent of hind-limb clasping was recorded for 15 seconds. The room lighting
was kept at 90 Lux. Hind-limb clasping was scored from 0 to 3: score 0 = both hind limbs were
splayed outward away from the abdomen; score 1 = one hind limb retracted inward toward the
abdomen for at least 50% of the observation period; score 2 = both hind limbs partially
retracted inward toward the abdomen for at least 50% of the observation period; score 3 =
both hind limbs completely retracted inward toward the abdomen for at least 50% of the
observation period. Hind-limb extension reflex severity scores were calculated by averaging
three trials separated by 1 min.

Open field test - The apparatus consisted of an illuminated (70 lux) white floor PVC foam arena (50 x 50 x 40 cm). A central zone (16 x 16 cm) was pre-defined. Test mice were placed in the centre of the open field and allowed to explore the arena for 30 minutes. At the end of each trial, the mouse was returned to the home cage and the arena was cleaned. Total distance travelled in 30 minutes and percentage of time spent in the central zone were used as measures of locomotor activity and anxiety levels, respectively.

Accelerating Rotarod test - Day 1: Mice walked on a rotating rod (Ugo Basile model 47650, Italy) at constant speed (4 rpm) for three minutes for two acclimatization trials. Day 2: Each mouse was placed on the rotating rod for three test trials, during which the rotation speed gradually increased from 4 to 40 rpm within four minutes. The inter trial interval was 1 hr. Performance was evaluated by measuring the latency to fall.

Three-chamber social interaction test – The apparatus consisted of a white rectangular PVC foam arena (62x42x23 cm) divided into one central and two adjacent compartments of equal size compartments connected by square openings. Empty cylindrical wire cages (Ø 8 cm, height 18 cm) were placed in the lateral chambers. Each mouse was placed in the central chamber and allowed to explore the arena for 10 min. The brightness of the room was kept at 10 lux. Chamber occupancy during habituation was equal between groups. After this, the mouse was returned to the waiting cage for 3 minutes. To assess social preference, an unfamiliar WT mouse of the same age, sex and strain (stranger 1) was gently introduced inside one of the wire cages to serve as a social stimulus. An unfamiliar and inanimate object was added to the other wire cage as the non-social stimulus. The test mouse was placed in the apparatus containing the social and non-social stimuli for 10 minutes. After that, the test mouse was removed and kept in the waiting cage for 3 minutes. All compartments were cleaned with 30% ethanol solution before the next test phase. To assess preference for social novelty, the non-social object from the previous test phase was replaced by an unfamiliar mouse (stranger 2) and the previous stranger 1 (now familiar mouse) was placed into the wire cage of the opposite compartment. The test mouse was placed in the central chamber and allowed to explore the arena for the next 10 minutes, after which it was returned to the home cage. The location of the novel and familiar mice with respect to the side compartments was counterbalanced across trials. The exploration of the target mouse/object was scored when the mouse's nose was detected within 2 cm from the cylindrical wire cage.

Y-maze spontaneous alternation test – Testing occurs in a Y-shaped maze with three arms (1,2,3) made of opaque plastic (34 x 9 x 14 cm) oriented in a 120^0 angle. Each mouse was randomly placed into one arm and allowed to explore the maze for 10 minutes. Room lighting was kept at 10 lux. An arm entry was recorded manually when the mouse moved beyond the central triangle of the maze and entered an arm with all four paws. Alternation behaviour was defined as consecutive entries into each of the three arms in overlapping triplet sets (e.g.: 1,

2, 3 or 2, 1, 3 or 3, 2, 1). The percentage of alternations was calculated as the number of actual alternations divided by the maximum possible number of arm entries.

Barnes-maze test – Twenty-four hours before training, the animal was habituated to the apparatus and escape box for a minute. During the training phase (4 days), mice were trained to locate the target hole (with an escape chamber underneath) from among 20 holes evenly spaced around the perimeter of an elevated circular open field (Ø 96 cm). Each animal was initially placed in the centre of the arena covered by an opaque cylinder, which was removed 10-20s after the start of the trial. Room lighting was kept at 635 lux. The mouse is then free to explore the platform for 3 minutes using four visual cues to aid navigation. If the mouse does not enter the escape chamber within 3 minutes the experimenter guides the mouse gently to the escape box (20x11x7cm) and leaves the mouse inside for 1 min. This step is repeated two more times a day (3 trials in total per day) with at least 20 minutes in between where the mouse is placed back in its home cage. To assess learning, a probe test in carried out 24h after the last training, where the escape box has been removed from the target hole. The mouse is placed on the platform and free to explore it for 3 minutes. Distance to first: distance walked from the centre of the arena to the target hole (centre-point detection); latency to first: time to reach the target hole from the centre of the arena (centre-point detection); errors to first are defined as checking any hole before reaching the escape box (nose-point detection).

Contextual fear conditioning – The fear conditioning chamber (27x27x35 cm, Ugo Basile, Italy) was equipped with a shock grid floor and a digital Near Infrared Red Video Fear Conditioning system. Mice from each genotype and sex were examined in four successive phases comprising: conditioned acquisition (day 1), memory of the conditioned background context A (day 2), exploration in new background context B (day 3) and memory extinction in the conditioned context A (day 4-7).

Context A was characterized by a cubic shape, illuminated at 100 Lux and the presence of a vanilla odour. The floor consisted of metal rods, mediating the foot shock. Context B was also cubic shaped, not illuminated and without additional odour cues. On day 1: Mice were introduced in the conditioning chambers scented with vanilla odour for 8 min and received 3 unconditioned stimuli (US: 1s, 0.25 mA foot shock, 2 min inter-trials interval). To examine the conditioned response to the context, mice were reintroduced in the training context 24 h later (day 2) and monitored for 5 minutes in the absence of the US. On day 3, the mice were introduced to a new context (context B) for 5 min to examine whether freezing behaviour is associated to a specific context (context A). During days 4-7 mice were placed back in context A without US to assess extinction of the association of the specific context A and the US (which is described as a form of new learning).