Effects of cardiac function alterations on the risk of postoperative thrombotic complications in patients receiving endovascular aortic repair

Xiaoning Sun[†,a,b], Siting Li[†,a,b], Yuan He[†,c], Yuxi Liu[†,c], Tianxiang Ma[c], Rong Zeng[a], Zhili Liu[a], Yu Chen[c], Yuehong Zheng[*,a,b], Xiao Liu[*,c]

a Department of Vascular Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100730, China

b Department, State Key Laboratory of Complex Severe and Rare Diseases, Peking Union Medical College Hospital, Chinese Academy of Medical Science and Peking Union Medical College, Beijing, China

c Key Laboratory of Biomechanics and Mechanobiology (Beihang University), Ministry of Education, Beijing Advanced Innovation Center for Biomedical Engineering, School of Biological Science and Medical Engineering, Beihang University, Beijing, 100083, China

*Correspondence:

Yuehong Zheng, zhengyuehong2022@outlook.com

Xiao Liu, liuxiao@buaa.edu.cn

Keywords: Thrombus, Hemodynamics, Computational model, Ventricular systolic function, Abdominal aortic aneurysms

Abstract

Chronic heart disease (CHD) is a common comorbidity of patients receiving endovascular aneurysm repair (EVAR) for abdominal aortic aneurysms (AAA). The ventricular systolic function determines the hemodynamic environments in aorta, and thus regulating the formation of postoperative thrombus. However, the explicit relationship between ventricular systolic function and EVAR complication of thrombotic events is unknown. Here, we proposed a three-dimensional numerical model coupled with the lumped-elements heart model, which is capable of simulating thrombus formation in diverse systolic functions. The computational results show that thrombus tended to occur at the interior side of the aorta arch and iliac branches, which is consistent with the post-operative imaging follow-up. In addition, we found that both an increase in maximum ventricular contractile force and a decrease in minimum ventricular contractile force inhibit thrombus formation, whereas the effect of heart rate on thrombus formation is not significant. In conclusion, changes in ventricular systolic function may alter the risk of thrombotic events after EVAR repair, which could provide insight into the selection of adjuvant therapy strategies for AAA patients with CHD.

Introduction

Abdominal aortic aneurysm (AAA) is one of the most common degenerative aortic diseases and is often complicated with various levels of cardiac dysfunction and cardiovascular comorbidities (1). For advanced and/or rapid-growing aneurysms, prompt surgical or endovascular interventions are indicated, among which endovascular aortic aneurysm repair (EVAR) is widely accepted as the first-line treatment option for anatomically feasible cases (2).

Intraluminal thrombus formation is commonly observed not only in major arteries with atherosclerotic and aneurysmal lesions but also, to a lesser degree, intra-prosthetically after endovascular stenting. Despite its lower rate of occurrence, intra-prosthetic thrombus formation may lead to post-operative distal embolism, branch occlusion, and critical end-organ malperfusion (3-5). The formation of intraluminal thrombus and thrombogenic atherosclerotic plaques are believed to be driven by hemodynamic factors. Pathogenic blood flow that generates oscillatory and non-physiological wall shear stress may trigger the accumulation of lipid and pro-thrombotic factors (6). By modeling the process of mass transport in the coagulation cascade, the patient-specific patterns of formation and distribution of intraluminal/intra-prosthetic thrombus formation can be simulated or predicted with acceptable accuracy (7, 8). Simulation of thrombus-forming reactions can facilitate the risk evaluation of post-operative thrombotic events and aid the tailoring of follow-up strategies.

However, the outcomes of numerical simulation of fluid dynamics are sensitive to the settings of boundary conditions (9, 10), which may vary, in real-world scenarios, significantly among patients with different levels of cardiac output (11). We noted that only limited and contradicted evidence exists on whether peri- and postoperative use of cardiac supportive drugs such as β-blockers could be beneficial for AAA patients after EVAR in light of aneurysmal sac shrinking and clinical outcomes (12-14). The effects of variations in cardiac function on the risk of intraluminal and intra-prosthetic thrombus formation are yet to be further elaborated.

In this pilot study, by coupling a lumped parameters model of cardiac output and aortic/peripheral outflow resistance with a 3D numerical mass transport model of thrombus formation, we focused on investigating the relation of cardiac functions and the predicted risk of thrombus formation in the aorta and/or endograft of 4 patients who underwent EVAR. Relative risks for thrombus formation were identified using machine-learning algorithms.

Methods

Imaging acquiring and 3D model reconstruction

Four patients diagnosed with AAA who received EVAR and regular follow-ups at the
department of vascular surgery of Peking Union Medical College Hospital were included. DICOM
format of CTA images of each patient from preoperative and at least 6 months post-operative was
obtained. Mimics 12.0 (Materialise, Belgium) and CRIMSON (15) were used to reconstruct 3D
models of the aorta from ascending aorta to aortoiliac bifurcations, with preservation of major
branches in the arches and the thoracoabdominal segments (Figure 1a). Models were subsequently
smoothed with WRAP (v2017, Geomagic) and loaded into SOLIDWORKS (v18, Solidworks Co.).
Vessel outlets were trimmed and extended sufficiently to allow for the fully developed flow pattern.
The geometric model was meshed with a tetrahedral mesh using ICEM (v17.0, ANSYS, Inc.). The
global meshing size was chosen at 0.02 mm to accommodate the smallest components. Elements of
the boundary layers were set to be hexahedral with a primary thickness of 0.001 mm and would
gradually develop over 5 layers (Figure 1b). Under steady-state conditions, grid independence is
considered to be achieved when the mean difference between WSS and platelet concentrations in two
continuous simulations is less than 1%. The study followed the Helsinki Declaration and was
approved by the ethical committee of Peking Union Medical College Hospital. Written informed
consent was obtained from all participants.

Mathematical modeling for 0D model

The aorta was separated into three sections and represented by three RLC elements. The outlets
of the aorta are modeled as the three elements Windkessel model(16). The parameters in the RLC
element and Windkessel model were calculated by solving the differential equations based on the
blood flow pressure and velocity. In addition, a function imitating ventricular systole is used in the
lumped parameter model of the heart module. Through animal tests, Suga and Sagawa(17) developed
a ventricular pressure-volume relationship that can be expressed as a time-varying function E(t), as
indicated in Equation (1):

90
$$E(t) = \frac{P_{SV}(t)}{V_{SV}(t) - V_0} \tag{1}$$

91   where E(t) is the time-varying function (mmHg/ml), P_{sv}(t) is the time function of ventricular pressure
92   (mmHg), V_{sv}(t) is a time function of ventricular volume (ml), and V_0 is the ventricular reference
93   volume (ml), which is the theoretical volume relative to ventricular zero pressure. Boston(18)

94 proposed a mathematical fit to determine the function of ventricular systole according to equation
95 (2):

$$E(t) = (E_{max} - E_{min}) \cdot E_n(t_n) + E_{min} \tag{2}$$

97 where E_{max} refers to the ventricular pressure-volume relation at end-systole, E_{min} refers to the
98 ventricular pressure-volume relation at end-diastolic. $E_n(t_n)$ represents the normalized function of
99 ventricular elasticity, which is described as the Hill equation as follows[19]:

$$E_n(t_n) = 1.55 \left[\frac{(\frac{t_n}{0.7})^{1.9}}{1+(\frac{t_n}{0.7})^{1.9}} \right] \left[\frac{1}{1+(\frac{t_n}{1.17})^{1.9}} \right] \tag{3}$$

101 where t_n is t/T_{max}, and the T_{max} can be calculated from the cardiac cycle:

$$T_{max} = 0.2 + 0.15T \tag{4}$$

Mathematical modeling for 3D simulation of thrombus formation

In this article, we simulate the 3D formation of thrombus by solving the convection-diffusion-
reaction equations. The thrombus formation is tracked through the formation of bounded platelets
(BPs), which was induced by localized high concentration of activated platelets (APs), coagulation
enzymes (C), and prolonged flow residence time (RRT). The schematic of the thrombus formation is
illustrated in Figure 1 g. We only show the general form of the equation here, and the complete
reactions and parameters adapted from the published models (8, 20, 21) are explained in detail in
Table S2 and Table S3:

$$\frac{\partial[C_i]}{t} + \nabla \cdot (\boldsymbol{u} \cdot [C_i]) = \nabla \cdot (D_i \cdot \nabla[C_i]) + S_i \#(5)$$

where $[C_i]$ is the concentration of species i; $\boldsymbol{u}$ represents the blood flow velocity vector; D_i refers to
the diffusivity of species i in blood; and S_i is a local reaction source term for species i. The species
considered in this model include rested platelets [RPs], activated platelets [APs], coagulation
enzymes[C], and flow residence time [RT].

The bounded platelets (BPs) attached to the reactive surface have neither convective nor
diffusive motion; therefore, in this case, Equation (6) is simplified and the deposition of platelets are
governed by the following concentration rate equations:

$$\frac{\partial[C_{BP}]}{t} = S_{BP} \#(6)$$

118   where S_{BP} is the local generation rate of bounded platelets (BPs).

119   Considering the resistance of bounded platelets to blood flow, the momentum source term was
120   added to the Navier-Stokes equations, and the viscosity coefficient was modified. Equation (7) shows
121   the modified Navier-Stokes equations. and Eq (8) and (9) show the modified viscosity and source
122   term:

$$\rho\frac{\partial \boldsymbol{u}}{\partial t} + (\boldsymbol{u} \cdot \nabla)\boldsymbol{u} = -\nabla p + \mu\Delta\boldsymbol{u} - S_M \#(7)$$

$$S_M = k_m \frac{[BP]^2}{[BP]^2 + [BP]_t^2}\boldsymbol{u}\#(8)$$

$$\mu = k_\mu \left(1 + \frac{[BP]^2}{[BP]^2 + [BP]_t^2}\right)\#(9)$$

where ρ represents the blood density equals $1050kg/m^3$, $\boldsymbol{u}$ represents the blood flow velocity vector,
p represents blood pressure, μ represents the blood viscosity, and S_M represents the local source
term of flow momentum. Detailed value of coefficient k_m, k_μ can be found in supplement material
(Supplementary Table S3).

Multiscale simulation methods and procedures

Our model coupled the 3D simulation with the lumped-elements model. First, based on the
clinically measured blood flow pressure and velocity(8), we transformed the aortic model from 3D to
lumped elements and coupled the boundary nodes to Windkessel elements(22) (Figure 1c, d, detailed
parameters in Supplementary Table S1). Subsequently, the lumped-elements model representing the
heart was coupled to the entrance node in the lumped-elements aortic model (Figure 1e). We first
adjusted the heart model's parameters so that the averaged cardiac output and aortic inlet blood
pressure levels matched the clinically determined values. To mimic the various heart functions, we
swept the parameters E_{max} (end-systolic elastance), E_{min} (end-diastolic elastance), and T (heart rate)
from 10% to -10% compared to the reference values and got the corresponding aortic blood flow
velocity in different heart functions. Finally, we simulated the formation of thrombus in the 3D aorta
under various cardiac functions, the boundary conditions of which are derived from the blood flow

velocity at the boundary nodes in the lumped-elements model (Figure 1f, g). The 3D simulations of

blood flow and thrombus formation were performed in FLUENT (v17.0, ANSYS, Inc.) with user-

defined functions, whereas the lumped-elements simulations were performed in MATLAB (R2021b,

MathWorks, Inc.) The results were analyzed and visualized using CFD-Post (v17.0, ANSYS, Inc.).

Statistical Analysis

Data were analyzed using Prism software (v9.0, GraphPad Prism, Inc). Statistical significance

for samples was determined using two-tailed unpaired Student's t-test. Data were considered

statistically significant if $P < 0.05$. The Parallel coordinate analysis and the Multiple correspondence

analysis are performed in MATLAB (R2021b, MathWorks, Inc.).

Results

Demographic characteristics and clinical verification of thrombus formation region

The baseline information of 4 patients were shown in table 1. There were 3 male and 1 female

patients, with an average age of 74.3 ± 7.6 years. Pre-operatively, laboratory results indicated normal

cardiac and renal function for the 4 subjects. All 4 patients underwent the operation successfully, and

follow-up CTA suggested patency of stents and visceral branches. Concerning thrombus formation,

CTA results in Figure 2 (A1-D1) illustratrd that thrombus was found at the root of the left subclavian

artery for patient number 3. For patient number 1, 2, and 4, CTA images all showed thrombus within

the iliac stents. Thrombus was not observed at other parts of the arterial or stent model from clinical

images. These regions with thrombus formation were defined as the region of interests (ROI) in the

following analysis.

The hemodynamic parameters and BP levels at ROI were demonstrated in Figure 2 and Figure 3.

The TAWSS, oscillatory shear index (OSI), and relative residence time (RRT) profiles of 4 models

were illustrated in Figure 3. In general, TAWSS was highest at the visceral branch level in all 4

patients, which indicated higher frictional force exerted on these vessel areas by the blood flow and

prevented activated platelets from binding to coagulant to form thrombus in our model. Descending

aorta and iliac branches had higher RRT, which indicated a higher degree of retention for a substance

in the bloodstream at the region and induced the development of thrombus in our model (See

supplementary material). RRT also increased at the oversizing area for the iliac branches. However,

zooming in on the ROI in Figure 2 did not indicate distinct abnormal local CFD parameters. In

addition, no significant difference in TAWSS, OSI, and RRT was observed for the ROI compared to

the average levels of the whole model (right channel in Figure 3). On the other hand, patterns of BP

distribution from the thrombus model simulation (Figure A2-D2) corresponded well with the

thrombus results from ROI.

Impact of cardiac function alterations on thrombosis

The impact of heart function alteration on thrombus formation in the ROI as well as the aortic

inlet velocity were shown in Figure 4. For patient 1, 2, and 4, the iliac branches were shown, while

the arch and descending aorta was illustrated for patient 3.

The aortic velocity profile demonstrated that the magnitude and shape of the blood flow profile

are affected by the heart function parameters of E_{max}, E_{min}, and T. Then we analyzed the formation of

thrombus, which was induced by the altered heart functions. First of all, the increased E_{max} (end-

systolic elastance, representing maximum ventricular contractile force) could maintain or alleviate

thrombus formation levels in the ROI of four patients. However, the effects of decreased E_{max} are

heterogeneous. For patient 1, a decrease in Emax would result in an increase in thrombus

development at the iliac branches. However, for patients 2-4, the lower Emax would result in less

thrombus development at the arch and descending aorta. In addition, the increased E_{min}, which

represents ventricular compliance, would maintain or exacerbate the thrombus formation in the ROI

of four patients. The decreased E_{min} would maintain or alleviate the thrombus formation in these

regions. Furthermore, the influences of heart rate (T) on thrombus formation are less significant.

We evaluated the area-averaged BP concentration and area-averaged TAWSS (traditional

indicator) of the four patients to quantitatively highlight the associations between heart function and

thrombus formation in ROI. We discovered that TAWSS is largely affected by E_{max} level, while E_{min}

and T have little impact by showing the Parallel coordinate charts classified by the heart function

(Emax, Emin, T ranges from +10% to -10% compared with the reference condition). In addition, we

found that increasing E_{max} and decreasing E_{min} could reduce overall thrombus formation levels

(Figure 5a-5b), which is in agreement with the qualitative analysis. Besides, both the increase and

decrease of heart rate (T) would exacerbate the thrombus formation level (Figure 5c). The effects of

Emin and t on TAWSS, however, were not significant, indicating the limitations of using

hemodynamic features to predict thrombus development. Then, we performed the multidimensional

analysis method to quantitatively reveal the correlations among E_{max}, E_{min}, T, and thrombus

formation. We found that the BP concentration was negatively correlated with E_{max}, and was
positively related with E_{min} (Figure 5c). However, there was a lack of correlation between heart rate
(T) and BP concentration. In addition, we found that the ROI (iliac in patient 1,2,4 or arch in patient
3) showed the highest correlations with E_{max} and E_{min}, which indicated that the thrombus formation in
these regions was mostly sensitive to the alteration in heart function.

Collectively, these data indicated that the increase in E_{max} and decrease in E_{min} would both
inhibit thrombus formation, and the effect of T was not significant.

Discussion

Endograft mural thrombus accumulation had reported rates up to 33% and was detected as early
as 1 week after EVAR (3). It might be due to cytokine and prothrombotic factors released from the
intramural thrombus of the aneurysm triggered by the operation. Whether the risk of long-term
thrombotic events could be increased is debatable (23). Mestres et al. reported that endograft mural
thrombotic deposits were related to device occlusion during 24 months of follow-up (24), while
Melson et al. did not find a significant association (25). Previous studies on CFD-based thrombus
simulation mainly focused on the intra-luminal thrombosis in the sac of AAA. Abnormal wall shear
stress, platelet activation, vortical structures, and morphological parameters were all found to play a
potential role (26-29). Regarding intra-prothetic thrombosis, Nauta and colleagues investigated the
alteration of PLAP in one patient receiving 3 virtual interventions including TEVAR (30). Liu et al.
explored the TAWSS, OSI, and RRT levels in 3 patients who underwent multibranched endovascular
repair (31). Most of these studies referred to previous works or idealized values for boundary blood
flow and pressure profiles.

Thrombosis involves complex interlinked interactions between platelets, coagulation cascades,
and the vascular wall (32). In this study, traditional hemodynamics parameters including WSS, OSI,
and RRT did not precisely identified regions of thrombus formation, probably due to the lack of
reflection of this complex reactions. In a former study, we used a numeric thrombus prediction model
to calculate the vascular remodeling and thrombotic events in one patient receiving hybrid repair for
the middle aortic syndrome, which was consistent with follow-up images (8). The continuum-
macroscopic scale model could capture the clotting patterns taking both activated platelets, local
hemodynamic conditons, and residence time into account (7). Herein, this numeric model was
applied to 4 patients receiving EVAR. In line with CTA results, thrombus formation was observed at

iliac branches in 3 patients and at the opening of LSCA in 1 patient. Iliac endografts are commonly

reported places for thrombosis, and the aorto-uni-iliac configuration was confirmed as a risk factor

for intra-prosthetic thrombus (IPT) deposit in a meta-analysis (4). The low-density mural thrombus in

the descending aorta, on the other hand, may originate from the disruption of vulnerable

atherosclerotic plaques (33). Our result suggested that this numeric model was suitable for both intra-

luminal and intra-prosthetic thrombosis prediction.

Previous works investigating the relationship between cardiac function and EVAR complications

were mainly retrospective cohorts or reviews (13, 34). To the best of our knowledge, this is the first

attempt at using a cardiac numeric model for post-EVAR patients. Lumped heart models used simple

parameters such as resistance and capacitance to simulate the relaxation, filling, contraction, and

ejection phases of the heart, and have been widely studied for their interaction with the arterial vessel

systems (35). In this study, we used a 0-D cardiac model adapted from Kim et al. to simulate the

change in heart function especially ventricular pressure/volume depending on the time-varying

elastance curve (36). Emax represented the maximum ventricular contractility, Emin represented

ventricular compliance or end-diastolic pressure, and t stood for the cardiac cycle. Heart failure with

reduced ejection fraction (HFrEF) or with preserved ejection fraction (HFpEF) could be mimicked

under the circumstances. The influence of cardiac parameters on thrombosis was reflected by altered

BP levels.

Concerning the actual thrombus forming area, both elevated or reduced E_{min} and t promoted

thrombus formation, yet changes of Emax induced distinct results. It should be noted that BP levels

in patients 1 and 3 were higher and more concentrated around the actual thrombus-forming area.

Despite thrombus deposition in the local iliac stent, the BP in patients 2 and 4 had relatively low

absolute value and occupied a more extensive spatial range. Morphological factors such as increasing

diameter because of stent oversize may have a stronger influence on thrombosis in these patients, and

the impact of cardiac parameter alteration should be cautiously interpreted. The heterogeneous results

from different patients emphasized the importance of individualized analysis.

Quantitation calculation of BP for general aortic parts revealed that Emax was negatively

associated with thrombus formation while E_{min} was positively related. It has been reported that male

AAA patients had reduced ventricular systolic and diastolic function (37). Our result indicated that

for patients with CHD (both HFrEF and HFpEF) in clinical practice, we should pay more attention to

their increased risk of thrombus formation after EVAR. Additionally, preservation of maximum

ventricular contractile force and stabilization of ventricular compliance or heart rate was also a
crucial consideration when selecting supporting drugs.

This study has some limitations. Patient-specific outflow data of each aortic branch may not
always be available, especially for supra-arch branches in AAA patients, and assumptions were made
for blood distribution in a limited number of branches according to the literature. The simulation
process should be applied in a larger patient cohort to validate our findings. Besides, we ignored the
arterial deformation in the 3D calculation. Future studies may combine fluid-structure interaction
(FSI) models with mass transport models of thrombus to more realistically predict thrombus
formation in deformed vessels(38, 39).

Conclusion

In conclusion, patients with CHD after EVAR may have an increased risk of thrombosis.
Postoperative cardiac support recommends drugs that support maximum ventricular contractile force,
maintain left ventricular pressure, and stabilize heart rate.

Author Contributions

X.S, S.T., Y.H., Y.L., and T.M. led the modelling and calculation of thrombus formation. X.S., T.M.
Y.Z., X.L. conceived, designed, and led the interpretation of the project. X.S, S.T., R.Z., Z.L.
provides and analyses the clinical data. Y.C. constructed the lumped-elements model. All authors
contributed to the writing of the manuscript.

Funding

This work was supported by the National Natural Science Research Foundation of China (grant no.
11827803, 31971244, 31570947, 32071311, U20A20390, 51890894, 82070492, and 82100519), the
Fundamental Research Funds for the General Universities (KG16186101) and the 111 Project
(B13003). The CAMS Innovation fund for Medical Science (CIFMS, Grant No.2021-I2M-C&T-A-
006 and 2021-I2M-1-016), and the National High Level Hospital Clinical Research Funding (Grant
No.2022-PUMCH-B-100 and 2022-PUMCH-A-190). The funders had no role in study design, data
collection, analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

None.

References

1. Lim S, Halandras PM, Park T, Lee Y, Crisostomo P, Hershberger R, et al. Outcomes of endovascular abdominal aortic aneurysm repair in high-risk patients. Journal of vascular surgery. 2015;61(4):862-8.

2. Isselbacher EM, Preventza O, Black JH, 3rd, Augoustides JG, Beck AW, Bolen MA, et al. 2022 ACC/AHA Guideline for the Diagnosis and Management of Aortic Disease: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. Circulation. 2022.

3. Maleux G, Koolen M, Heye S, Heremans B, Nevelsteen A. Mural thrombotic deposits in abdominal aortic endografts are common and do not require additional treatment at short-term and midterm follow-up. Journal of vascular and interventional radiology : JVIR. 2008;19(11):1558-62.

4. Perini P, Bianchini Massoni C, Azzarone M, Ucci A, Rossi G, Gallitto E, et al. Significance and Risk Factors for Intraprosthetic Mural Thrombus in Abdominal Aortic Endografts: A Systematic Review and Meta-analysis. Annals of vascular surgery. 2018;53:234-42.

5. Banno H, Kawai Y, Sato T, Tsuruoka T, Sugimoto M, Kodama A, et al. Low-density vulnerable thrombus/plaque volume on preoperative computed tomography predicts for spinal cord ischemia after endovascular repair for thoracic aortic aneurysm. Journal of vascular surgery. 2021;73(5):1557-65.e1.

6. Cunningham KS, Gotlieb AI. The role of shear stress in the pathogenesis of atherosclerosis. Laboratory investigation; a journal of technical methods and pathology. 2005;85(1):9-23.

7. Menichini C, Xu XY. Mathematical modeling of thrombus formation in idealized models of aortic dissection: initial findings and potential applications. Journal of mathematical biology. 2016;73(5):1205-26.

8. Sun X, Ma T, Liu Z, Wu X, Zhang B, Zhu S, et al. Sequential numerical simulation of vascular remodeling and thrombosis in unconventional hybrid repair of ruptured middle aortic syndrome. Medical engineering & physics. 2021;94:87-95.

9. Pirola S, Cheng Z, Jarral OA, O'Regan DP, Pepper JR, Athanasiou T, et al. On the choice of outlet boundary conditions for patient-specific analysis of aortic flow using computational fluid dynamics. Journal of biomechanics. 2017;60:15-21.

10. Alastruey J, Xiao N, Fok H, Schaeffter T, Figueroa CA. On the impact of modelling assumptions in multi-scale, subject-specific models of aortic haemodynamics. Journal of the Royal Society, Interface. 2016;13(119).

11. Pagoulatou S, Adamopoulos D, Rovas G, Bikia V, Stergiopulos N. The effect of left ventricular contractility on arterial hemodynamics: A model-based investigation. PloS one. 2021;16(8):e0255561.

12. Kim W, Gandhi RT, Peña CS, Herrera RE, Schernthaner MB, Tsoukas A, et al. Effect of β-blocker on aneurysm sac behavior after endovascular abdominal aortic repair. Journal of vascular surgery. 2017;65(2):337-45.

13. Salata K, Syed M, Hussain MA, Eikelboom R, de Mestral C, Verma S, et al. Renin-angiotensin system blockade does not attenuate abdominal aortic aneurysm growth, rupture rate, or perioperative mortality after elective repair. Journal of vascular surgery. 2018;67(2):629-36.e2.

14.   Saratzis A, Saratzis N, Melas N, Kiskinis D. Pharmacotherapy before and after endovascular
repair of abdominal aortic aneurysms. Current vascular pharmacology. 2008;6(4):240-9.

15.   Arthurs CJ, Khlebnikov R, Melville A, Marčan M, Gomez A, Dillon-Murphy D, et al.
CRIMSON: An open-source software framework for cardiovascular integrated modelling and
simulation. PLoS computational biology. 2021;17(5):e1008881.

16.   Westerhof N, Lankhaar J-W, Westerhof BE. The arterial windkessel. Medical & biological
engineering & computing. 2009;47(2):131-41.

17.   Suga H, Sagawa K. Instantaneous pressure-volume relationships and their ratio in the excised,
supported canine left ventricle. Circulation research. 1974;35(1):117-26.

18.   Goldberger AL, Amaral LA, Glass L, Hausdorff JM, Ivanov PC, Mark RG, et al. PhysioBank,
PhysioToolkit, and PhysioNet: components of a new research resource for complex physiologic
signals. circulation. 2000;101(23):e215-e20.

19.   Stergiopulos N, Meister J-J, Westerhof N. Determinants of stroke volume and systolic and
diastolic aortic pressure. American Journal of Physiology-Heart and Circulatory Physiology.
1996;270(6):H2050-H9.

20.   Menichini C, Cheng Z, Gibbs RG, Xu XY. Predicting false lumen thrombosis in patient-
specific models of aortic dissection. Journal of The Royal Society Interface. 2016;13(124):20160759.

21.   Menichini C, Xu XY. Mathematical modeling of thrombus formation in idealized models of
aortic dissection: initial findings and potential applications. Journal of mathematical biology.
2016;73(5):1205-26.

22.   Stergiopulos N, Westerhof BE, Westerhof N. Total arterial inertance as the fourth element of
the windkessel model. The American journal of physiology. 1999;276(1):H81-8.

23.   Kapetanios DM, Karkos CD, Papazoglou KO. Changes in circulating markers of coagulation
and fibrinolysis after EVAR. International angiology : a journal of the International Union of
Angiology. 2018;37(6):444-50.

24.   Mestres G, Maeso J, Fernandez V, Allegue N, Constenla I, Matas M. Incidence and evolution
of mural thrombus in abdominal aortic endografts. Annals of vascular surgery. 2009;23(5):627-33.

25.   Oliveira NF, Bastos Gonçalves FM, Hoeks SE, Ten Raa S, Ultee KH, Rouwet E, et al.
Clinical outcome and morphologic determinants of mural thrombus in abdominal aortic endografts.
Journal of vascular surgery. 2015;61(6):1391-8.

26.   Hansen KB, Arzani A, Shadden SC. Mechanical platelet activation potential in abdominal
aortic aneurysms. Journal of biomechanical engineering. 2015;137(4):041005.

27.   Zambrano BA, Gharahi H, Lim C, Jaberi FA, Choi J, Lee W, et al. Association of
Intraluminal Thrombus, Hemodynamic Forces, and Abdominal Aortic Aneurysm Expansion Using
Longitudinal CT Images. Annals of biomedical engineering. 2016;44(5):1502-14.

28.   Bhagavan D, Di Achille P, Humphrey JD. Strongly Coupled Morphological Features of
Aortic Aneurysms Drive Intraluminal Thrombus. Scientific reports. 2018;8(1):13273.

29.   Zambrano BA, Gharahi H, Lim CY, Lee W, Baek S. Association of vortical structures and
hemodynamic parameters for regional thrombus accumulation in abdominal aortic aneurysms.
International journal for numerical methods in biomedical engineering. 2022;38(2):e3555.

30.   Nauta FJH, Lau KD, Arthurs CJ, Eagle KA, Williams DM, Trimarchi S, et al. Computational
Fluid Dynamics and Aortic Thrombus Formation Following Thoracic Endovascular Aortic Repair.
The Annals of thoracic surgery. 2017;103(6):1914-21.

31.   Liu MY, Jiao Y, Liu J, Zhang S, Li W. Hemodynamic Parameters Predict In-stent
Thrombosis After Multibranched Endovascular Repair of Complex Abdominal Aortic Aneurysms: A
Retrospective Study of Branched Stent-Graft Thrombosis. Frontiers in cardiovascular medicine.
2021;8:654412.

32.   Chong MY, Gu B, Armour CH, Dokos S, Ong ZC, Xu XY, et al. An integrated fluid-structure
interaction and thrombosis model for type B aortic dissection. Biomechanics and modeling in
mechanobiology. 2022;21(1):261-75.

33.   Virmani R, Burke AP, Farb A, Kolodgie FD. Pathology of the vulnerable plaque. Journal of
the American College of Cardiology. 2006;47(8 Suppl):C13-8.

34.   Kurzencwyg D, Filion KB, Pilote L, Nault P, Platt RW, Rahme E, et al. Cardiac medical
therapy among patients undergoing abdominal aortic aneurysm repair. Annals of vascular surgery.
2006;20(5):569-76.

35.   Hunter PJ, Pullan AJ, Smaill BH. Modeling total heart function. Annual review of biomedical
engineering. 2003;5:147-77.

36.   Kim HJ, Vignon-Clementel IE, Figueroa CA, LaDisa JF, Jansen KE, Feinstein JA, et al. On
coupling a lumped parameter heart model and a three-dimensional finite element aorta model. Annals
of biomedical engineering. 2009;37(11):2153-69.

37.   Åström Malm I, De Basso R, Engvall J, Blomstrand P. Males with abdominal aortic
aneurysm have reduced left ventricular systolic and diastolic function. Clinical physiology and
functional imaging. 2022;42(1):1-7.

38.   Ma T, Liu X, Ren Q, Zhang Z, Sun X, Zheng Y, et al. Flow-mediated dilation analysis
coupled with nitric oxide transport to enhance the assessment of endothelial function. Journal of
Applied Physiology. 2021;131(1):1-14.

39.   Chong MY, Gu B, Armour CH, Dokos S, Ong ZC, Xu XY, et al. An integrated fluid–
structure interaction and thrombosis model for type B aortic dissection. Biomechanics and Modeling
in Mechanobiology. 2022;21(1):261-75.

Table 1. Clinical information and total BP levels of patients under different heart conditions.

	Patient 1	Patient 2	Patient 3	Patient 4	Average
Age, years	80	70	64	83	74.3 ± 7.6
Sex	Male	Female	Male	Male	-
Hypertension	Yes	Yes	Yes	Yes	-
Diabetes	No	Yes	No	No	
Smoking	Yes	No	Yes	No	-
Laboratory results					
LVEF%	68	74	67	77	71.5 ± 4.2
NT-proBNP, pg/ml	104	112	81	77	93.5 ± 14.8
D-dimer, mg/L	6.31	5.3	1.67	1.41	3.7 ± 2.2
HGB, g/L	123	121	150	120	128.5 ± 12.5
Creatinine, μmol/L	78	70	93	111	88.0 ± 15.6
Follow-up, months	35.5	47.7	34.8	11.9	32.5 ± 14.9

407   BP, bound platelets; E_{max}, maximum elastase; E_{min}, minimum elastase; LVEF, left ventricular ejection fraction;

408   NT-proBNP, N-terminal pro-brain natriuretic peptide; HGB, hemoglobulin.

Figure 1. Flowchart of patient-specific modeling

A, Geometry of postoperative aorta was modeled from CTA data, starting from the ascending aorta to bilateral common iliac arteries. **B,** Meshing the model and vessel outlets were sufficiently extended to allow fully developed flow boundary. **C, D,** Building the Windkessel model, lumped-parameter aortic model based on the clinical measured blood flow pressure and velocity. **E,** The lumped-elements model representing the heart was coupled to the entrance node in the lumped-elements aortic model. **F,** Coupling the three kinds of lumped-parameter model to give out the aortic blood flow velocity in different heart functions. **G,** The simulation of thrombosis formation involving the activation of resting platelets (RPs), the deposition of coagulant (C) and the aggregation of activated platelets (APs) to form the bounded platelets (BPs). **H,** Thrombus concentration profile of one patient. CTA, computed tomography angiography.

Figure 2. Clinical verification of the thrombus forming model.

Red arrows indicated places of thrombus formation on CTA images.

Figure 3. CFD characteristics of patients' original models.

TAWSS, time average wall shear stress; OSI, oscillatory shear index; RRT, relative residence time; ns, non-significant. The fourth columns indicated RRT levels at ROI.

Figure 4. Impact of heart function alteration on thrombosis at critical positions.

430 Bounded platelets (BP) levels under different heart conditions were shown for each patient. Aortic

431 blood flow values over a cardiac cycle were also shown.

Figure 5. Heart function alteration and change of thrombosis or TAWSS at different aortic parts.

A-C. Parallel coordinate charts of BP concentration and TAWSS classified by heart function (E_{max}, E_{min}, T ranges from +10% to -10% compared with the reference state). BP concentration and TAWSS have been rescaled ranges from 0 to 1. The arrowhead in Figure 5A indicates the significant role of E_{max} on thrombus formation. D. Multiple correspondence analysis (MCA) of E_{max}, E_{min}, T, and BP concentration in different aortic sections.