Precision RNAi using synthetic shRNAmir target sites

Thomas Hoffmann[1,2,5,#], Alexandra Hörmann[3,#], Maja Corcokovic[3,#], Jakub Zmajkovic[1], Matthias Hinterndorfer[1], Jasko Salkanovic[3], Fiona Spreitzer[3], Anna Köferle[3], Katrin Gitschtaler[3], Alexandra Popa[3], Sarah Oberndorfer[3], Florian Andersch[1], Markus Schäfer[1], Michaela Fellner[1], Nicole Budano[3], Jan G. Ruppert[3], Paolo Chetta[3], Melanie Wurm[3], Johannes Zuber[1,4*] & Ralph A. Neumüller[3*]

[1] Institute of Molecular Pathology (IMP), Campus-Vienna-Biocenter 1, 1030 Vienna, Austria

[2] Vienna BioCenter PhD Program, Doctoral School of the University at Vienna and Medical University of Vienna, Vienna BioCenter (VBC), 1030 Vienna, Austria

[3] Boehringer Ingelheim RCV GmbH & Co KG, Doktor-Boehringer-Gasse 5-11, 1120 Vienna, Austria

[4] Medical University of Vienna, Vienna BioCenter, 1030 Vienna, Austria

[5] Present address: Advantage Therapeutics GmbH, Karl-Farkas-Gasse 22, 1030 Vienna, Austria

[#] These authors contributed equally to this work

[*] Corresponding authors: J.Z.: johannes.zuber@imp.ac.at; R.A.N.: ralph.neumueller@boehringer-ingelheim.com

Abstract

Loss-of-function genetic tools are widely applied for validating therapeutic targets, but their utility remains limited by incomplete on- and uncontrolled off-target effects. We describe artificial RNA interference (ARTi) based on synthetic, ultra-potent, off-target-free shRNAs that enable efficient and inducible suppression of any gene upon introduction of a synthetic target sequence into non-coding transcript regions. ARTi establishes a scalable loss-of-function tool with full control over on- and off-target effects.

Introduction

Drug development is guided by genetic loss-of-function (LOF) experiments that validate a therapeutic target, study its general and disease-specific functions, and thereby model and benchmark expected activities of inhibitory molecules. Applied genetic tools include RNAi, CRISPR/Cas9, and site-specific recombination technologies such as the Cre-Lox or Flp-FRT systems[1,2]. While these technologies have undoubtedly revolutionized genetic screening, target identification and validation, each method is associated with drawbacks that limit the usability for certain aspects of target validation. Specifically, RNAi is prone to off-target effects[3–5] and insufficient knock-down levels, while CRISPR/Cas9-based methods are associated with off-target effects and incomplete LOF across cell populations. These limitations are particularly problematic for candidate targets in oncology, which should ideally be validated genetically

in cancer cell lines and tumor models *in vivo*. In both cases, insufficient LOF or off-target effects can lead
to far-reaching misconceptions about target suppression effects. Outgrowth of wild-type clones that
retain gene function upon CRISPR- and recombination-based gene editing can result in transient
phenotypes that complicate data interpretation. Similarly, insufficient knock-down or uncontrolled off-
target effects induced by RNAi can lead to an unjustified de-prioritization or pursuit of candidate targets,
respectively. Prior to initiating the time- and resource-intense process of drug development, more
informative target validation assays would be highly desirable.

Results

To develop such an assay system, we reasoned that instead of using gene-specific LOF triggers for every
new candidate gene, the expression of any gene could be efficiently suppressed by engineering the target
site of a pre-validated, highly potent, synthetic short-hairpin RNA (shRNA) into its exonic sequence (Figure
1A). Besides ensuring efficient target knockdown in a highly standardized manner, such an approach
would also provide control over off-target activities through expressing the shRNA side-by-side in target-
site engineered and wildtype cells. This approach leaves the genome engineering procedure, needed to
integrate the pre-validated artificial RNA interference (ARTi) target site into a gene of interest, as the only
potential source of off-target effects. As suitable RNAi system, we chose optimized micro-RNA embedded
shRNAs (shRNAmirs) in the miR-E backbone[6], which do not interfere with endogenous miRNA processing[7]
and can be expressed from tet-responsive elements and other Pol-II promoters in the 3'-UTR of
fluorescent reporter genes, thus providing a versatile system for inducible RNAi[8]. To identify potent ARTi
sequences with minimal off-target activity, we analyzed the nucleotide composition of shRNAs that reach
exceptionally high performance scores in common shRNA design algorithms[9] and of miRNA seed
sequences with exceptionally low off-target scores according to siSPOTR[10]. By merging nucleotide biases
identified in both analyses, we derived a 22-nt base composition matrix for the design of ARTi shRNAmirs
(TTCGWWWNNAHHWWCATCCGGN; W = A/T, H = A/T/C; N = A/T/G/C) (Figure 1B; Figure 1 - figure
supplement 1A,B). To further reduce possible off-target effects, we eliminated all shRNAs whose extended
seed sequence (guide positions 2-14) had a perfect match in the human or mouse transcriptome and,
finally, selected six top-scoring ARTi predictions for experimental validation.

We tested these ARTi-shRNAmirs using an established knockdown reporter assay[6] for their ability to
suppress expression of a GFP transgene that harbors the respective target sites in its 3'-UTR. In all six
cases, ARTi-shRNAmir matched or outperformed previously validated highly potent shRNAmirs targeting
Renilla luciferase or PTEN[6] (Figure 1 - figure supplement 1C). We compared these results to a panel of 592
gene-specific shRNAs that were tested using the same assay and found that ARTi-shRNAmirs ranked
among top-performing shRNAmirs overall (Figure 1C). Next, we selected the three top-performing ARTi-
shRNAmirs and evaluated possible off-target activities using competitive proliferation assays and
transcriptome profiling. ARTi-shRNAmir expression had no effects on proliferation or survival in four
human and three mouse cell lines (Figure 1D; Figure 1 - figure supplement 1D), with the exception
ARTi.6634 and ARTi.6786, which induced a mild fitness defect in the murine leukemia cell line RN2 and
MV4-11 respectively. In contrast to effects of a MEK inhibitor trametinib, which we included as a positive
control, stable expression of ARTi-shRNAmirs had only marginal effects on the transcriptome in two
commonly used human cell lines (Figure 1E,F; Figure 1 - figure supplement 1E,F) and no effect in
proliferation assays (Figure 1 - figure supplement 1G), indicating that they do not trigger major off-target

effects, even in the absence of their respective target site. Together, these studies established a set of
highly potent, off-target-free ARTi-shRNAmirs, among which we selected ARTi.6570 (ARTi-shRNAmir) for
further investigations.

To establish ARTi as a method for target validation, we performed ARTi-based LOF experiments in cancer
cell lines and xenograft models for three prominent oncology targets: EGFR and KRAS, which act as driving
oncogenes in various cancer types[11,12], and STAG1, which has been identified as a synthetic lethal
interaction with recurrent LOF mutations of STAG2[13–16]. To establish an ARTi-repressible version of
oncogenic EGFR^{del19}, we cloned ARTi-shRNAmir target sequences into an expression construct encoding
EGFR^{del19} fused to dsRed (Figure 2A), which rendered Ba/F3 cells cytokine-independent and sensitive to
EGFR inhibition (Figure 2 – figure supplement 2A). We introduced this construct into human EGFR^{del19}-
dependent PC-9 lung adenocarcinoma cells and subsequently knocked out the endogenous *EGFR* gene.
The EGFR^{del19}::V5::dsRed::ARTi transgene fully rescued the loss of endogenous EGFR^{del19}, while doxycycline
(dox)-induced expression of the ARTi-shRNAmir (ARTi.6570) strongly inhibited proliferation of PC-9 cells
and triggered a near-complete suppression of the EGFR^{del19}::V5::dsRed::ARTi protein (Figure 2B,C; Figure
2 – figure supplement 2B). Consistently, in RNA-sequencing we observed an almost complete drop of
reads mapping to the codon-optimized *EGFR^{del19}::V5::dsRed::ARTi* transgene upon dox-inducible
expression of the ARTi-shRNAmir, as well as a downregulation of *DUSP6* and other canonical targets of
RAF-MEK-ERK signaling, which acts as key effector pathway of EGFR^{del19} (Figure 2 – figure supplement 2C-
E).

To evaluate whether ARTi can recapitulate drug activities *in vivo*, we xenotransplanted
EGFR^{del19}::V5::dsRed::ARTi engineered PC-9 cells harboring the dox-inducible ARTi-shRNAmir and treated
recipient mice upon tumor formation with either dox or the clinically approved EGFR inhibitor
Osimertinib. Dox-induced expression of the ARTi-shRNAmir led to rapid and durable tumor regression that
was indistinguishable from the effects of Osimertibib (Figure 2d) and not observed in parental PC-9 cells
that lack the ARTi target site (Figure 2 – figure supplement 2F). We therefore conclude that ARTi-induced
phenotypes are to be attributed to on-target effects that predict the activity of advanced small-molecule
inhibitors *in vivo*.

In a second study, we used ARTi to investigate KRAS^{G12R}, an oncogenic KRAS variant with no available *in-
vivo* compatible inhibitors. To establish a suitable KRAS^{G12R}-driven model, we engineered a
dsRed::KRAS^{G12R}-ARTi transgene and the dox-inducible ARTi-shRNAmir into KRAS^{G12C}-dependent MIA
PaCa-2 pancreatic adenocarcinoma cells and subsequently knocked out endogenous *KRAS* alleles (Figure
2E; Figure 2 – figure supplement 3A). As expected, switching the driving oncogene from KRAS^{G12C} to
KRAS^{G12R} rendered MIA PaCa-2 cells resistant to the KRAS^{G12C} inhibitor AMG-510[17–19] (Figure 2 – figure
supplement 3B). ARTi-shRNAmir induction led to strong suppression of dsRed::KRAS^{G12R} expression at the
mRNA and protein level (Figure 2F; Figure 2 – figure supplement 3C,D) and marked anti-proliferative
effects (Figure 2G). In xenograft experiments, ARTi-mediated suppression of KRAS^{G12R} led to full tumor
regression in the absence of KRAS^{G12C} (Figure 2H), while tumors harboring both oncogenic KRAS alleles
only displayed a delay in tumor progression, suggesting that both oncogenes contribute to tumor growth
in vivo.

To evaluate STAG1 as synthetic-lethal dependency that is relevant in wide range of STAG2-deficient
cancers (Figure 2 – figure supplement 4A)[13–16], we engineered an isogenic pair of STAG2-wildtype and -
deficient HCT 116 colon carcinoma cells and homozygously inserted ARTi target sites (besides an AID::V5

tag; explained in the materials and methods section) into the 3'-UTR of endogenous *STAG1* (Figure 2I).
Western blotting confirmed the knockout of *STAG2*, the insertion of the targeting cassette into the *STAG1*
locus, and potent suppression of STAG1 following dox-induced ARTi-shRNAmir expression (Figure 2J,K;
Figure 2 – figure supplement 4B). Suppression of STAG1 in STAG2-deficient HCT 116 cells impaired their
proliferation *in vitro* (Figure 2 – figure supplement 4C) and the progression of xenografted tumors *in vivo*
(Figure 2L), while dox-induced expression of the ARTi-shRNAmir had no anti-proliferative effects in the
presence of STAG2.

Discussion

Together, these studies establish ARTi as a versatile and precise LOF method for target validation in
oncology and beyond. Instead of designing gene-specific LOF reagents that remain prone to off-target
effects, ARTi involves a simple, highly standardized experimental procedure that provides full control over
on- and off-target effects and can be applied to any coding and non-coding gene. In cells that are pre-
engineered to express the dox-inducible ARTi-shRNAmir, candidate genes can be converted into ARTi
target genes and subsequently evaluated head-to-head in a highly standardized manner, either through
knocking in ARTi target sites into endogenous loci or through knock-out rescue approaches. Placement of
ARTi target sites in non-coding transcript regions leaves the endogenous target protein unaltered, which
is a key advantage over chemical-genetic LOF methods relying on the introduction of degron tags[20,21].
Targeting endogenously engineered ARTi target sites through tetracycline (Tet)-inducible ARTi-shRNAmir
expression enables inducible, titratable, and likely reversible (not tested in this study) LOF perturbations
of candidate genes without altering their endogenous transcriptional regulation, which is not possible
using conventional Tet-expression systems. In principle, by engineering multiple target sites in the same
cell, ARTi enables combinatorial LOF perturbations, e.g. for modeling synergistic target interactions,
without multiplying the risk for off-target effects. Although ARTi requires the design of gene-specific
sgRNAs and genome engineering steps that can be non-trivial, the method enables researchers to reach
well-interpretable, comparable, and unambiguous experimental results that can guide larger investments
in targets of high medical interest. Beyond providing a scalable method for early target validation, we
foresee that ARTi can be used to establish on-target benchmark phenotypes for guiding the development
and optimization of inhibitory molecules.

Figure Legends

Figure 1: Design and selection ARTi-shRNAmirs. A, Schematic outline of the ARTi approach. **B,** Sequence
logo (https://weblogo.berkeley.edu) displaying nucleotide position biases of 2161 shRNAs with
exceptionally high DSIR scores (>105). Inlay depicts miRNA seed sequence biases. **C,** Reporter assay
comparing gene-specific shRNAs to ARTi-shRNAmirs. **D,** Competitive proliferation assays in human cell
lines after transduction with ARTi-shRNAmirs, neutral (shRen.713) and essential control shRNAs
(shRPL9.324 or shPSMA3.164). **E,** Principal component (PC) analysis of gene expression profiling upon
stable expression of indicated shRNAs or treatment with MEK inhibitor (trametinib) in RKO cells. **F,**

Volcano plots visualizing de-regulated genes upon expression of indicated shRNAs and trametinib
treatment in RKO cells, compared to empty vector control.

Figure 2: Experimental validation of the ARTi approach. A, Schematic of EGFR^{del19}::V5::dsRed::ARTi
engineering in PC-9 cells. Blue color denotes overexpressed ARTi variant. Green denotes endogenous
gene. **B,** Western Blot demonstrating knock-down of EGFR^{del19}::V5::dsRed::ARTi. Western Blot is a
representative example of three independent biological repeat experiments. **C,** Proliferation assay and
crystal violet staining of parental and engineered PC-9 cells in absence or presence of dox. Crystal violet
staining is a representative example of two independent biological repeat experiments. **D,** *In vivo*
experiment comparing dox-induced EGFR^{del19}::V5::dsRed::ARTi knockdown to pharmacological EGFR^{del19}
inhibition. Mean tumor volume and +/- s.e.m. is plotted for all *in vivo* experiments. **E,** Schematic of MIA
PaCa-2 engineering. Blue color denotes overexpressed ARTi variant. Green denotes endogenous gene. **F,**
Western blot for KRAS and Actin in indicated engineered MIA PaCa-2 cells in the presence and absence of
dox. Western Blot is a representative example of two independent biological repeat experiments. **G,**
Proliferation assay and crystal violet staining of parental and engineered MIA PaCa-2 cells in absence or
presence of dox. Crystal violet staining is a representative example of two independent biological repeat
experiments. **H,** Growth curves of tumors implanted with engineered MIA PaCa-2 cells in absence and
presence of dox. In vivo **I,** Schematic of C-terminal endogenous tagging of STAG1. Green color denotes
endogenous genes. **J,** Western Blot demonstrating knock-down of STAG1-ARTi. Western Blot is a
representative example of three independent biological repeat experiments. **K,** Immunohistochemistry
staining of STAG1 in engineered HCT 116 cells in absence and presence of dox. Asterisk marks an area of
murine fibroblasts that serve as an internal positive control. **I,** Growth curves of tumors implanted with
engineered HCT 116 cells in the absence and presence of dox.

Figure 1 – figure supplement 1: Design and selection ARTi-shRNAmirs. A, Sequence logo displaying
nucleotide position biases of seed sequences (guide position 2-8) with the lowest off-target score in
siSPOTR analysis (top 1%). **B,** Sequence logo displaying position biases in sequences from a that harbor a
T in guide position 2. **C,** Knockdown efficiency of ARTi-shRNAmirs. Flow cytometric quantification of GFP
knockdown efficiency in immortalized MEFs two days after transduction with indicated ARTi-shRNAmir or
control shRNAs. Percent knockdown is normalized to shRen.713 control. shRen.660 served as a neutral
control whose target site was not included in the reporter. Red asterisks indicate ARTi-shRNAmirs that
were selected for follow-up studies. **D,** Toxicity of ARTi-shRNAmirs. Competitive proliferation assays of
three murine cell lines after transduction with ARTi-shRNAmirs, shRen.713 or shMyc.1834 control,
showing the relative fraction of shRNA-expressing cells at indicated time points following the initial
measurement (day 4 after shRNA transduction). **E,** Principal component analysis of gene expression
profiling in HT-1080 cells. Respective shRNAmirs and treatments are indicated in the respective colors. X-
axis: principal component 1; Y-axis: principal component 2. **F,** Volcano plots visualizing de-regulated genes
in HT 1080 cells. All shRNAmirs and treatments were tested against the empty vector control. X-axis: -
log$_{10}$(p-value); Y-axis: log$_2$ fold change. **G,** Cell growth assay for ARTi-shRNAmir transduced cells and their
parental controls in the presence and absence of dox.

Figure 2 – figure supplement 1: Validation of ARTi *in vitro* and *in vivo* - EGFR. A, Functional validation of
the EGFR[del19] construct in Ba/F3 cells. Provided are GI_{50} values (nM) for indicated conditions and
compounds (parental cells IL-3 dependent; other cells dependent on transgenes (no IL-3)). Numbers
represent GI_{50} value calculated from three technical repeats. Three biological repeat experiments were
conducted, and one representative experiment is shown in panel A. **B,** Western Blot analysis confirming
expression of EGFR[del19]::V5::dsRed::ARTi constructs, knockout of endogenous EGFR and dox-induced
knockdown or EGFR[del19]::V5::dsRed::ARTi. Western Blot is a representative example of three
independent biological repeat experiments. **C,** Expression levels (y-axis: normalized counts) of the
EGFR[del19]::V5::dsRed::ARTi construct in parental and engineered PC-9 cells after 4 and 8 days of dox
treatment or no treatment. Individual datapoints of three biological replicates overlay the boxplots. **D,**
Expression levels (y-axis: normalized counts) of the MAPK target gene *DUSP6* in parental and engineered
PC-9 cells after 4 and 8 days of dox treatment or no treatment. Individual datapoints of three biological
replicates overlay the boxplots. **E,** Heatmap visualizing the gene expression changes in MAPK pathway
target and ERBB genes in engineered cells in the presence and absence of dox. **F,** *In vivo* experiment
comparing dox-induced expression of the ARTi-shRNAmir to drinking water control in PC-9 parental
cells. Lines connect mean tumor volume data and +/- s.e.m..

Figure 2 – figure supplement 2: Validation of ARTi *in vitro* and *in vivo* - KRAS. A, Schematic of MIA PaCa-
2 genome engineering. **B,** AMG-510 GI_{50} values in MIA PaCa-2 parental and engineered cells. Numbers
represent GI_{50} value calculated from three technical repeats. Three biological repeat experiments were
conducted, and one representative experiment is shown in panel B. **C,** Expression levels (y-axis:
normalized counts) of the dsRed::KRAS[G12R] transgene in parental and engineered MIA PaCa-2 cells after 4
and 8 days of dox treatment or no treatment. Individual datapoints of three biological replicates overlay
the boxplots. **D,** Western blot for KRAS (green) and Actin (magenta) (upper plot) and dsRed (green) and
Actin (magenta) (bottom plot) for indicated engineered MIA PaCa-2 cells in the presence and absence of
dox. Western Blot is a representative example of three independent biological repeat experiments.

Figure 2 – figure supplement 3: Validation of ARTi *in vitro* and *in vivo* – STAG1. A, Schematic of synthetic
lethal interaction between STAG1 and STAG2. Cells survive loss of either paralog but are incapable of
growing upon combined loss of STAG1 and STAG2. **B,** Western blot confirmation of STAG2 knockout and
ARTi-shRNAmir induced knockdown of endogenous STAG1::V5::ARTi. Western Blot is a representative
example of three independent biological repeat experiments. **C,** Proliferation assay of ARTi engineered
HCT 116 cells *in vitro*, visualized by crystal violet staining following a 9-day dox or control treatment.
Staining is a representative example of three independent biological repeat experiments. **D**,
Quantification of nuclear Stag1 level (Figure 2k) using engineered HCT 116 cells in an in vivo
xenotransplantation experiment the control group (-dox) and doxycycline (+dox) treated group.

Figure 2 – source data 1. Original blots for Figure 2B and Figure2 - figure supplement 1B.

Figure 2 – source data 2. Original blots for Figure 2F.

Figure 2 – source data 3. Original blots for Figure 2J.

Figure 2 – figure supplement 2 – source data 1. Original blots for Figure2 - figure supplement 2D.

Figure 2 – figure supplement 3 – source data 1. Original blots for Figure2 - figure supplement 3B.

Material and Methods

Design and cloning of ARTi shRNAs

To design pairs of artificial shRNAs and matching target sites that trigger effective and selective target
suppression with minimal off-target effects, nucleotide composition of shRNAs that reach exceptionally
high performance scores in a well-established siRNA prediction tool (DSIR=Designer of Small Interfering
RNA[9]) and contain no A or T in guide position 20 to eliminate shRNAmirs that produce RISC-loadable small
RNAs from the passenger strand were analyzed. To establish criteria for ARTi design, siRNA predictions
for the human and mouse genome were retrieved from DSIR[9] using default parameters. Top scoring
predictions (DSIR score > 105) harboring G or C in position 20 were analyzed for nucleotide biases, which
were used to define basic design criteria at the 5'-end the 3'-half. Next, possible off-targets were
minimized using siSPOTR[10], an siRNA-based prediction tool that assesses off-target potential of different
siRNA seed sequences (guide positions 2-8) in the human and mouse genome. Seed sequences with the
lowest predicted off-target activity (top 1%) showed biases for C and/or G in guide positions 2-6, but 23%
contained a T at the 5' end of the seed sequence that is required in our design (Fig. 1b). Among these, we
observed particularly strong biases for CG in the following two positions (Fig. 1b), which were found in
73% of all seed sequences harboring a 5' T. Overall, 17% of all top-scoring seed sequences harbored TCG
at their 5' end, making it the second most common triplet (after CGC, which was found in 20%). Based on
these analyses, we fixed the first 4 nucleotides of the guide to TTCG. For the following positions we
reasoned that introducing additional GC biases would destroy 5'-3' asymmetry of small RNA duplexes that
is critically required for efficient RISC loading. To maintain sufficient asymmetry, we therefore decided to
bias the next 3 positions towards A or T, which in most positions is in alignment with nucleotide biases
associated with knockdown efficacy.

For the remaining sequence 3' of the seed region, we adhered to nucleotide features associated with
knockdown efficacy based on our DSIR analysis, which are remarkably prominent and cannot all be
explained through established processing requirements. Altogether, this established a 22-nt matrix for
the design of ARTi shRNAmirs (TTCGWWWWNNAHHWWCATCCGGN; W = A/T, H = A/T/C; N = A/T/G/C). In a
last step, we further reduced possible off-target effects by eliminating all guides whose extended seed
sequence (guide positions 2-14) had a perfect match in the human or mouse transcriptome and, finally,
selected the following six top-scoring ARTi predictions for experimental validation:

ARTi.6588 – target site: TCCGGATGAAGTTTATATCGAA / shRNAmir (97mer):
TGCTGTTGACAGTGAGCGCCCGGATGAAGTTTATATCGAATAGTGAAGCCACAGATGTATTCGATATAAACTTCAT
CCGGATGCCTACTGCCTCGGA

ARTi.6570 – target site: TCCGGATGATATTGTTATCGAA / shRNAmir (97mer):
TGCTGTTGACAGTGAGCGCCCGGATGATATTGTTATCGAATAGTGAAGCCACAGATGTATTCGATAACAATATCAT
CCGGATGCCTACTGCCTCGGA

ARTi.6634 – target site: TCCGGATGATGTTTTAATCGAA / shRNAmir (97mer):
TGCTGTTGACAGTGAGCGCCCGGATGATGTTTTAATCGAATAGTGAAGCCACAGATGTATTCGATTAAAACATCAT
CCGGATGCCTACTGCCTCGGA

ARTi.6786 – target site: TCCGGATGATATTGTATACGAA / shRNAmir (97mer):
TGCTGTTGACAGTGAGCGCCCGGATGATATTGTATACGAATAGTGAAGCCACAGATGTATTCGTATACAATATCAT
CCGGATGCCTACTGCCTCGGA

ARTi.6834 – target site: TCCGGATGATATTGCATACGAA / shRNAmir (97mer):
TGCTGTTGACAGTGAGCGCCCGGATGATATTGCATACGAATAGTGAAGCCACAGATGTATTCGTATGCAATATCAT
CCGGATGCCTACTGCCTCGGA

ARTi.6516 – target site: TCCGGATGAAGTTTAATTCGAA / shRNAmir (97mer):
TGCTGTTGACAGTGAGCGCCCGGATGAAGTTTAATTCGAATAGTGAAGCCACAGATGTATTCGAATTAAACTTCAT
CCGGATGCCTACTGCCTCGGA

The following ARTi target sequences were used for experimental validation studies:

Insertion into the coding sequence before the STOP codon:

ATCCGGATGATATTGTATACGAATCCGGATGATATTGTTATCGAA (with the first "A" being inserted to retain
the reading frame)

Insertion after the STOP codon:

TCCGGATGATATTGTATACGAATCCGGATGATGTTTTAATCGAATCCGGATGATATTGTTATCGAA

shRNAs were ordered as single stranded DNA Ultramer oligonucleotides (Integrated DNA Technologies),
amplified by PCR and cloned into different retroviral or lentiviral miRE/miRF shRNAmir expression vectors
(LT3GFPIR[6]) using EcoR/XhoI restriction digest or Gibson assembly.

299
Cell Culture

Human HCT 116 cells (ATCC: CCL-247) were cultured in RPMI 1640 medium (Thermo Fisher) and PC-9 cells
in McCoy's 5A medium (Thermo Fisher), supplemented with 10% FBS and 1 x GlutaMAX (Thermo Fisher).
RKO (ATCC: CRL-2577) and MOLM-13 cells (DSMZ: ACC 584) were cultured in RPMI 1640, supplemented
with 10% FBS (Sigma-Aldrich), 4mM L-glutamine (Thermo Fisher), 1mM sodium pyruvate (Sigma-Aldrich),
and penicillin/streptomycin (100 U ml^{-1}/100 µg ml^{-1}, Sigma-Aldrich). HT-1080 (ATCC: CCL-121) and LentiX
lentiviral packaging cells (Clontech, cat. no. 632180) were cultivated in DMEM (Thermo Fisher) with 10%
FBS, 4mM L-glutamine, 1mM sodium pyruvate, and penicillin/streptomycin (100 U ml^{-1}/100 µg ml^{-1}).
MV4-11 cells (ATCC: CRL-9591) were cultured in IMDM with 10% FBS, 4mM L-glutamine, 1mM sodium
pyruvate, and penicillin/streptomycin (100 U ml^{-1}/100 µg ml^{-1}).

Murine MLL-AF9^{OE}, Nras^{G12D} AML cells (RN2; Zuber et al. 2011) were cultured in RPMI 1640 medium
supplemented with 10% FBS, 20 mM L-glutamine, 10 mM sodium pyruvate, 10 mM HEPES (pH 7.3),
penicillin/streptomycin (100 U ml^{-1}/100 µg ml^{-1}), and 50 µM β-ME. *Kras^{G12D}*, *Trp53$^{-/-}$*, *MYC^{OE}* PDAC cells
(EPP2), SV40 large T antigen immortalized mouse embryonic fibroblasts (RRT-MEF[22]; and NIH/3T3 (ATCC:
CRL-1658) were cultured in DMEM supplemented with 10% FBS, 20 mM glutamine, 10 mM sodium
pyruvate, and penicillin/streptomycin (100 U ml^{-1}/100 µg ml^{-1}). MIA PaCa-2 (ATCC: CRL-1420) and GP2d
(Ecacc: 95090714) cells were cultured in DMEM supplemented with 10% FBS. Ba/F3 (DSMZ: ACC300) cells
were cultured in RPMI 1640 medium supplemented with 10% FBS, 10ng/ml IL-3 (R&D systems) and Ls513

cells were cultured in RPMI 1640 medium supplemented with 10% FBS. All cell lines were maintained at
37°C with 5% CO_2, routinely tested for mycoplasma contamination and authenticated by short tandem
repeat analysis.

Reporter Assay

SFFV-GFP-P2A-Puro-ARTi-target sensor was cloned into pRSF91 retroviral plasmid[23] using Gibson
assembly. RRT-MEFs were transduced with retroviruses expressing a GFP-reporter harboring the target
sites for validated shRNAs and one ARTi-shRNA in its 3'-UTR. For each reporter cell line single cells were
FACS-sorted into 96-well plates using a FACSAria III cell sorter (BD Bioscience) to obtain single-cell derived
clones. These clones were transduced with retrovirus constructs in pSin-TRE3G-mCherry-miRE-PGK-Neo
(TCmPNe) backbone expressing either the respective doxycycline (dox)-inducible ARTi shRNA or validated
shRNAs and mCherry fluorescence marker. shRNA expression was induced with dox and GFP levels were
quantified via flow cytometry 2 days post induction. Knockdown efficiency was calculated as 1 minus the
ratio of mean GFP signal in mCherry$^+$ (shRNA$^+$) cells over mCherry$^-$ cells and normalized to Renilla
luciferase specific neutral control shRNA (Ren.713).

Competitive proliferation assay

To investigate the effect of ARTi constructs in the absence of the endogenous target gene, competitive
proliferation assays were performed in several human and murine cell lines. Human HT-1080, RKO,
MOLM-13, and MV4-11 cell lines were lentivirally transduced with shRNAmir expression constructs cloned
into pRRL-SFFV-GFP-miRF-PGK-Neo (SGFN) backbone at 20-60% efficiency. Initial infection efficiency was
determined at day 4 post transfection (day 0) by measuring GFP expression as a readout using iQue
Screener Plus flow cytometer (IntelliCyt). Percentage of shRNA$^+$ cells (GFP positive) was monitored by flow
cytometry in regular intervals and results were normalized to day 0.

Human GP2d, Ls513, and MIA PaCa-2 cell lines were lentivirally transduced with shRNAmir constructs
cloned into pRRL-TRE3G-GFP-miRE-PGK-Puro-IRES-rtTA3 backbone (LT3GEPIR, Addgene plasmid
#111177). 500 cells were seeded in duplicates in 96-well plates and treated with 1 µg/ml dox for 9-10 days
and analyzed with Incucyte (Sartorius). Untreated cells served as reference.
Murine NIH-3T3, EPP2 and RN2 cells were retrovirally transduced with shRNAmir constructs cloned into
pMSCV-miR-E-PGK-Neo-IRES-mCherry backbone (LENC; Addgene plasmid #111163), and initial infection
levels were determined by flow cytometry based on mCherry expression 4 days post transduction (day 0).
349

Crystal Violet staining

To visualize ARTi-shRNAmir's effect, crystal violet staining assays were performed. 25'000 HCT 116 cells
and 15'000 PC-9 cells per well were seeded in a 6-well plate containing 2 mL tetracycline-free growth
medium and 1 µg/mL dox. Medium was exchanged every 2-3 days. After 9 days, wells were washed with
ice-cold PBS and subsequently stained with 1 mL of 2.3% crystal violet solution for 10 minutes.
Subsequently, wells were washed with ultrapure water and dried overnight. Images were obtained with
a scanner.

Transcriptional profiling

For the unbiased identification of ARTi shRNA off-targets, RKO and HT-1080 shRNA[+] cells from the
competition proliferative assay experiment were selected with Geneticin/G418 Sulfate (Gibco) for 7 days
and checked for GFP expression. On day 7, one arm of the empty vector control group was treated with
IC$_{50}$ concentration of trametinib (MedChem Express: HY-10999) for 24 hours based on the data in
Genomics of Drug Sensitivity in Cancer (GDSC) database (https://www.cancerrxgene.org)[24]. Subsequently,
cells were trypsinized, washed with ice-cold PBS, pelleted and snap frozen. Total RNA was isolated using
in-house magnetic beads kit and King Fisher Duo Prime Purification System (Thermo Fisher). NGS libraries
were prepared with QuantSeq 3' mRNA-Seq Library Prep Kit (FWD) HT for Illumina (Lexogen) and UMI
Second Strand Synthesis Module for QuantSeq FWD (Lexogen). Samples were sequenced on Illumina
NovaSeq platform with 100bp single-read protocol.

Engineered MIA PaCa-2 and PC-9 cells were cultured in the presence of 1 µg/ml dox to induce expression
of the ARTi shRNA. dox-containing media were replenished twice weekly and on day 4 and day 8 after the
initial treatment. 2x10^6 dox-treated and untreated control cells were harvested, washed with PBS, lysed,
treated with DNAse I (QIAGEN) and total RNA was extracted using the RNeasy Mini Kit (QIAGEN). NGS
libraries were prepared as above. Samples were sequenced on an Illumina NextSeq 2000 platform with a
75bp protocol.

Bioinformatic analyses of 3' mRNA-Seq

For 3' mRNA-Seq reads derived from the human cell lines HT-1080 and RKO, the six nucleotide long 5'
UMIs were attached to each read name with umi-tools (v1.0.0)[25]. Subsequently, the UMIs plus the next
four nucleotides (UMI spacer), as well as 3' adapters (stringency of 3) and bases with low quality (threshold
of 25) were trimmed away using cutadapt (v1.18)[26] and its wrapper tool trimgalore (v0.6.2). Read quality
control was performed with FastQC (v0.11.8). The remaining reads were sample-wise aligned to the
human (GRCh38.p13; GCA_000001405.28) reference genome. Mapping and subsequent filtering of 3' UTR
mapped reads was performed with slamdunk (v0.4.3)[27] in QuantSeq mode (slamdunk map -5 12 -n 100 -
q). 3' UTR regions were assembled based on the description in Muhar et al.[28]. Aligned and filtered reads
were deduplicated with umi-tools (v1.0.0) (2), based on the mapping coordinate and the UMI attached to
the read name, prior to quantifying read abundances within 3' UTR regions using featureCounts (v2.0.1)[29].
Differential expression analysis (DEA) was performed with DESeq2 (v1.30.1)[30] for each ARTi shRNA to
empty vector control. Here, the number of up- and downregulated genes were calculated by filtering the
DEA results for genes with a log2 fold-change ≥ 2 (up) or ≤ 2 (down) and a -log10 p-value ≥ 5. Principal
component analysis was performed on the 1000 most variable expressed genes with the prcomp function
from the stats (v4.2.0) R package.

For transcriptional profiling of the human cell lines MIA PaCa-2 and PC-9, the genome reference file and
annotations were constructed based on the GRCh38 assembly and the Ensembl 86 version, respectively.
The sequences of the shRNA construct (ARTi.6570 (RN_v_76, RN_v_118)) as well as the

EGFR[del19]::V5::dsRed::ARTi (RN_v_108) and the dsRed::Linker::KRAS[G12R]-ARTi (RN_v_287) were also
included. Mapping of the sequencing reads derived from the human cell lines was performed with STAR
(v2.5.2b)[31] aligner allowing for soft clipping of adapter sequences. Quantification of read counts to
transcript annotations was implemented using RSEM (v1.3.0)[32] and featureCounts (v1.5.1)[29].
Normalization of read counts and differential analysis was implemented with the limma[33] and voom[34] R
packages.
402

Genome engineering of EGFR in PC-9 and Ba/F3

Genome engineering of PC-9 cells was done as previously described[35]. In brief: PC-9_RIEN cells were
transduced with an ecotropic pMSCV-EGFRdel19_V5_dsRed_ARTi-PGK-Blasticidin retrovirus cloned at
GenScript and produced in Platinum E cells (Cell Biolabs) in the presence of 8 µg/mL Polybrene (Merck
Millipore). After 24 hours, stable transgenic cell pools were selected using 10 µg/mL Blasticidin (Sigma-
Aldrich). Subsequently, cells were diluted to obtain single cell clones. After 14 days of culture, single cell-
derived colonies were transferred to 6-well plates and analyzed by Western Blot. Identified homozygous
PC-9_ EGFR[del19]-ARTi clones were further engineered by cutting endogenous EGFR with a CRISPR all-in-
one vector pX458_Exon20_gRNA TAGTCCAGGAGGCAGCCGAA (GenScript) using X-tremeGENE 9 DNA
transfection reagent (Roche) according to the protocol supplied by the vendor. 48 hours after
transfection, GFP positive cells were sorted by FACS (SONY cell sorter S800Z) and diluted to obtain single
cell clones. Positive clones, which contained only the exogenous EGFR[del19]-ARTi, but not the endogenous
EGFR, were identified by Western Blot. Next, the selected EGFR clone was transduced with a pantropic
LT3GEPIR_Puro_ARTi-shRNA TTCGATAACAATATCATCCGGA retrovirus cloned at GenScript, China and
produced via the Lenti-X Single Shot system (Clontech). 72 hours later, stable transgenic cell pools were
selected using 0.5µg/mL Puromycin (Sigma-Aldrich). Following the selection, cells were diluted to obtain
single cell clones. After 14 days of culture, single cell-derived colonies were transferred to 6-well plates
and induced via 1µg/ mL dox. Positive clones were characterized by a strong GFP-induction that was
identified by flow cytometry.

422
423   Ba/F3 cells were transduced with an ecotropic pMSCV-EGFRdel19_V5_dsRed_ARTi-PGK-Blasticidin
424   retrovirus cloned at GenScript and produced in Platinum E cells in the presence of 4 µg/mL Polybrene.
425   After 72 hours, stable transgenic cells were selected by using 50 µg/mL Blasticidin, without adding IL-3.
426

Genome engineering of KRAS in MIA PaCa-2

MIA PaCa-2 ARTi-shRNAmir expressing cells were transduced with an ecotropic pMSCV-dsRed::KRAS[G12R]-
ARTi-PGK-Blasticidin retrovirus cloned at GenScript, China and produced in Platinum E cells in the
presence of 8 µg/mL Polybrene. After 24 hours, stable transgenic cell pools were selected using 10 µg/mL
Blasticidin. Subsequently, endogenous KRAS was knocked out by transient transfection of three gRNAs
targeting exon 2 and the region containing the G12C variant (present in MIA PaCa-2 cells) were used in a
co-transfection (gRNA#3: *GAATATAAACTTGTGGTAGT*; gRNA#6: *CTTGTGGTAGTTGGACTTG*; gRNA#7:
GTAGTTGGAGCTTGTGGCGT). Knockout clones were identified by the absence of the endogenous KRAS
protein using Western Blot.

Genome engineering of STAG1/STAG2 in HCT 116

HCT 116 cell line was engineered by cutting STAG1 with gRNAs targeting the region close to the STOP codon. Guide RNAs *TTCTTCAGACTTCAGAACAT* or *CTGAAGAAAATTTACAAATC* were cloned into the pSpCas9(BB)-2A-GFP plasmid (pX458; Addgene plasmid 48138) and used in a co-transfection. Simultaneously, a STAG1_AID_V5_P2A_Blasti_STOP_ARTi repair template with 800bp of left and right homologous arms of the STAG1 genomic locus (in pUC57-Simple backbone) was transfected into HCT 116 cells using Lipofectamine 3000 transfection reagent (Thermo Fisher) according to the manufacturer's instructions. A stable transgenic cell pool was selected 48 hours after transfection using 5µg/ml Blasticidin and diluted to obtain single cell clones. Positive clones were identified by Western Blot.

Identified homozygous HCT 116_STAG1_ARTi clones were further engineered by disrupting STAG2 gene with a CRISPR all-in-one vector Hs0000077505_U6gRNA-Cas9-2A-GFP and Hs0000077502_U6gRNA-Cas9-2A-GFP, respectively. Cells were transfected using X-tremeGENE 9 DNA transfection reagent according to the manufacturer's instructions, sorted 48 hours post transfection for GFP positive cells and diluted to obtain single cell clones. Positive clones were identified by Western Blot.

Next, the selected HCT 116 clone was transduced with a pantropic LT3GEPIR_Puro_ARTi-shRNA *TTCGATAACAATATCATCCGGA* retrovirus cloned at GenScript, China and produced via the Lenti-X Single Shot system (Clontech). 72 hours later, stable transgenic cell pools were selected using 2 µg/mL Puromycin (SIGMA, P9620). Following the selection, cells were diluted to obtain single cell clones. After 14 days of culture, single cell-derived colonies were transferred to 6-well plates and induced via 1µg/ mL dox. Positive clones were characterized by a strong GFP-induction that was identified by flow cytometry.

Western Blot

The following primary antibodies were used for immunoblot analyses: EGFR (Cell Signaling, #4267); STAG1 (GeneTex, GTX129912); STAG2 (Bethyl, A300-159A); b-actin (Sigma, A5441); KRAS (LSbio, LS-C17566); V5 (Sigma, V8012). PC-9, MIA PaCa-2, and HCT 116 cell pellets harboring EGFR, KRAS, and STAG1/STAG constructs respectively were lysed in Triton X-100 lysis buffer, sonicated and stored at -80°C. For protein detection, the pellets were thawed on ice, followed by 15 minutes centrifugation at 13,000 rpm and 4°C. Further, cell lysates were loaded onto a pre-casted SDS–polyacrylamide gel (4-12%) and proteins were transferred onto a nitrocellulose or PVDF-membrane. Membranes were probed with the respective primary antibodies overnight. The next day, secondary antibodies conjugated with fluorescent dye were added and the proteins were detected by the Odyssey detection system.

Compound treatment

To investigate sensitivity to EGFR-targeting compounds, cell viability was determined using the Cell Titer Glo assay (Promega). For this purpose, 10 mM stock solutions in DMSO of afatinib[36], Osimertinib[37,38] [37] and poziotinib[39,40] were used. 5'000 cells per well were seeded in 150µL of the medium in technical triplicates in 96-well plates and incubated at 37°C and 5% CO_2 for 5 hours, followed by the compound addition. Cells were treated with seven different concentrations of inhibitors in a serial eight-fold dilutions starting with the highest concentration of 3 µM. For comparability, DMSO normalization to the highest added volume was performed. Subsequently, cells were cultivated for 96 hours at 37°C and 5% CO_2. 50 µL of Cell Titer Glo reagent was added to each well, incubated for 10 minutes in the dark and luminescence

478 was measured using the multilabel Plate Reader VICTOR X4. The measurement time was set to 0.2
479 seconds. Luminescence values relative to DMSO-treated cells were plotted in GraphPad Prism and fitted
480 using nonlinear regression with a variable slope to calculate IC50 values at 50% inhibition. MIA PaCa-2
481 KRAS G12C inhibitor (AMG-510)[17–19] treatments were performed as described for EGFR with the following
482 modifications: 2'000 cells per well, and 0.5nM-3µM concentration range of AMG-510.

In vivo experiments

The PC-9 EGFR k.o., EGFR[del19]::ARTi study was performed at the AAALAC accredited animal facility of
CrownBio Leicestershire, UK. Female NSG® (NOD.Cg-PrkdcscidII2rgtm1wjl/SzJ) Crl mice were obtained
from Charles River. Age of animals at study initiation was 7-8 weeks and hat an acclimatization period of
≥14 days. Mice were groups housed in IVCs. The study complies with the UK Animals Scientific Procedures
Act 1986 (ASPA) in line with Directive 2010/63/EU of the European Parliament and the Council of 22
September 2010 on the protection of animals used for scientific purposes.

All other in vivo experiments were performed at the AAALAC accredited animal facility of Boehringer
Ingelheim RCV GmbH & CoKG. Female BomTac:NMRI-*Foxn1*[nu] mice were obtained from Taconic Denmark
at 6-8 weeks of age. After the arrival, mice were allowed to adjust to the housing conditions at least for 5
days before the start of the experiment. Mice were housed in pathogen-free and controlled
environmental conditions (open cage housing), and handled according to the institutional, governmental
and European Union guidelines (Austrian Animal Protection Laws, GV-SOLAS and FELASA guidelines).

Studies were approved by the internal ethics committee (called "ethics committee") of Boehringer
Ingelheim RCV GmbH & Co KG in the department of Cancer Pharmacology and Disease Positioning.
Furthermore, all protocols were approved by the Austrian governmental committee (MA 60 Veterinary
office; approval numbers GZ: 903122/2017/21 and GZ: 416181-2020-29).

To establish subcutaneous tumors, mice were injected with 2×10^6 HCT 116 in PBS, 5×10^6 MIA PaCa-2
in 1:2 Matrigel : PBS with 5% FBS or with 1×10^7 PC-9 cells in PBS with 5% FBS.

Tumor diameters were measured with a caliper three times a week. The volume of each tumor [in mm^3]
was calculated according to the formula "tumor volume = length * diameter^2 * $\pi/6$." Mice were
randomized into the treatment groups when tumor size reached between ~130 and 190 mm^3. Group sizes
were calculated for each tumor model based on tumor growth during model establishment experiments.
A power analysis was performed using a sample size calculator
(https://www.stat.ubc.ca/~rollin/stats/ssize/n2.html). For all models used in the studies, 10 mice per
group were used. 2mg/ml doxycycline hyclate (Sigma) and 5mg/ml sucrose were added to the drinking
water of the treatment groups, the control group received water with 5mg/ml sucrose only. Osimertinib
(Tagrisso, AstraZeneca, 40 mg tablet) was dosed per os daily at a dose of 25mg/kg in Natrosol and control
mice were dosed per os daily with Natrosol. To monitor side effects of treatment, mice were inspected
daily for abnormalities and body weight was determined three times per week. Animals were sacrificed
when the tumors reached a size of 1'500 mm^3. Food and water were provided ad libitum. *In vivo*
experiments were not repeated.

Immunohistochemistry

Xenograft samples were fixed in 4% formaldehyde for 24 h and embedded in paraffin. Two-micrometer-
thick sections were cut using a microtome. STAG1 was stained with Polink 2 Plus HRP rat NM detection
system (GBI Labs #D46-6) according to manufacturer's instructions using a recombinant rat monoclonal
STAG1 antibody (Abcam ab241544, Lot:GR3334172-1; 1:200) after cooking 10 minutes at 110°C in
antigen unmasking solution (Vector #H3301). After staining the slides were digitalized (scanner: Leica
Aperio AT2). All slides were reviewed and evaluated for quality by a board-certified MD Pathologist.
Imaging analysis was performed using the digital pathology platform HALO (Indica Labs). A tissue-
classifying algorithm was trained to selectively recognize viable tumor tissue against stroma, necrosis,
and skin. The tissue classification output for each scan was reviewed and manually edited as necessary.
A cell detection and scoring algorithm was trained to measure DAB optical density (OD) in the nuclei of
tumor cells. A positivity threshold for DAB OD was determined by normalization with respect to the DAB
OD as calculated from *bona fide* negative tissue (e.g., murine stroma as background). The object data for
background-normalized nuclear ODs for each tumor cell were exported from three control and three
doxycycline-treated cases (207945 and 33331 pooled objects, respectively). The cumulative distribution
of the background-normalized DAB OD for each tumor cell nucleus was then plotted for control and
doxycycline-treated cases, separately (supplemental figure 4, panel K).

Reagents, code and data availability

DNA sequences of the transgenes used in this study and the code used to generate the figures are
available upon request. The workflow for the RNA-seq bioinformatics analyses is described in the
Materials and Methods section. RNA sequencing data were uploaded to GEO with the accession
numbers: GSE218404 and GSE218617. The data will be made publicly available upon acceptance of this
manuscript. All ARTi cell lines and constructs described in this study are available upon request.

Cell lines

All newly generated ARTi cell lines are available upon request. The following parental cell lines were
used:

Cell line name	species	Supplier name	Catalog number	Clone number
PC-9	human	ECACC	90071810	Lot# 14A030
MIA PaCa-2	human	ATCC	CRL-1420	Lot# 1350798
HT 1080	human	ATCC	CCL-121	Lot# 63835574
HCT 116	human	ATCC	CCL-247	Lot# 6028631
RKO	human	ATCC	CRL-2577	Lot# 59354083
MV4-11	human	ATCC	CRL-9591 (old number)	Lot# 58352230
MOLM-13	human	DSMZ	ACC 554	Lot# 11
NIH-3T3	human	ATCC	CRL-1658	Lot# 63292276
EPP2		KrasG12D, Trp53-/-, MYCOE PDAC cells (EPP2)	NA	NA

RN2	murine	MLL-AF9OE, NrasG12D AML cells (RN2; Zuber et al. 2011)	NA	NA
Gp2d	human	ECACC	95090714	Lot# 11E013
Ls513	human	ATCC	CRL-2134	Lot# 57761342
Ba/F3	murine	DSMZ	ACC300	
293T-Lenti-X	human	Clontech	632180	Lot# 1404558A

Competing interests

Alexandra Hörmann, Maja Corcokovic, Jasko Salkanovic, Fiona Spreitzer, Anna Köferle, Katrin
Gitschtaler, Alexandra Popa, Sarah Oberndorfer, Nicole Budano, Jan G. Ruppert, Paolo Chetta, Melanie
Wurm and Ralph Neumüller: Are or were full time employees Boehringer Ingelheim. Johannes Zuber:
Quantro Therapeutics GmbH (Co-founder, Scientific Advisor, Shareholder); Mirimus Inc. (Scientific
Advisor, Shareholder); AstraZeneca-CRUK Functional Genomics Consortium (Scientific Advisor);
Boehringer Ingelheim (Research Support).

Funding

J.Z. is supported by the Swiss National Science Foundation (Early Postdoc Mobility Fellowship
P2BSP3_188110) and European Union's Horizon 2020 research and innovation program under the Marie
Skłodowska-Curie grant agreement No. 847548 (VIP2) and No. 101032582 (AML-SynergyX). Research at
the IMP is supported by Boehringer Ingelheim and the Austrian Research Promotion Agency
(Headquarter grant FFG-852936). MS is a member of the Boehringer Ingelheim Discovery Research
global post-doc program.

Acknowledgements

564 The authors thank all colleagues in the labs of Johannes Z. and R.A.N. for critical discussions and
565 technical support.

References

1. Mohr, S. E., Smith, J. A., Shamu, C. E., Neumüller, R. A. & Perrimon, N. RNAi screening comes of age: improved techniques and complementary approaches. *Nat Rev Mol Cell Bio* 15, 591–600 (2014).

2. Housden, B. E. *et al.* Loss-of-function genetic tools for animal models: cross-species and cross-platform differences. *Nat Rev Genet* 18, 24–40 (2017).

3. Jackson, A. L. *et al.* Expression profiling reveals off-target gene regulation by RNAi. *Nat Biotechnol* 21, 635–637 (2003).

4. Scacheri, P. C. *et al.* Short interfering RNAs can induce unexpected and divergent changes in the levels of untargeted proteins in mammalian cells. *P Natl Acad Sci Usa* 101, 1892–1897 (2004).

5. Lin, X. *et al.* siRNA-mediated off-target gene silencing triggered by a 7 nt complementation. *Nucleic Acids Res* 33, 4527–4535 (2005).

6. Fellmann, C. *et al.* An Optimized microRNA Backbone for Effective Single-Copy RNAi. *Cell Reports* 5, 1704–1713 (2013).

7. Premsrirut, P. K. *et al.* A Rapid and Scalable System for Studying Gene Function in Mice Using Conditional RNA Interference. *Cell* 145, 145–158 (2011).

8. Zuber, J. *et al.* Toolkit for evaluating genes required for proliferation and survival using tetracycline-regulated RNAi. *Nat Biotechnol* 29, 79–83 (2011).

9. Vert, J.-P., Foveau, N., Lajaunie, C. & Vandenbrouck, Y. An accurate and interpretable model for siRNA efficacy prediction. *Bmc Bioinformatics* 7, 520 (2006).

10. Boudreau, R. L. *et al.* siSPOTR: a tool for designing highly specific and potent siRNAs for human and mouse. *Nucleic Acids Res* 41, e9–e9 (2013).

11. Hynes, N. E. & MacDonald, G. ErbB receptors and signaling pathways in cancer. *Curr Opin Cell Biol* 21, 177–184 (2009).

12. Punekar, S. R., Velcheti, V., Neel, B. G. & Wong, K.-K. The current state of the art and future trends in RAS-targeted cancer therapies. *Nat Rev Clin Oncol* 1–19 (2022) doi:10.1038/s41571-022-00671-9.

13. Bailey, M. L. *et al.* Paralogous synthetic lethality underlies genetic dependencies of the cancer-mutated gene STAG2. *Life Sci Alliance* 4, e202101083 (2021).

14. Lelij, P. van der *et al.* STAG1 vulnerabilities for exploiting cohesin synthetic lethality in STAG2-deficient cancers. *Life Sci Alliance* 3, e202000725 (2020).

15. Benedetti, L., Cereda, M., Monteverde, L., Desai, N. & Ciccarelli, F. D. Synthetic lethal interaction
between the tumour suppressor STAG2 and its paralog STAG1. *Oncotarget* 5, 37619–37632 (2014).

16. Lelij, P. van der *et al.* Synthetic lethality between the cohesin subunits STAG1 and STAG2 in diverse
cancer contexts. *Elife* 6, e26980 (2017).

17. Fakih, M. *et al.* CodeBreak 100: Activity of AMG 510, a novel small molecule inhibitor of KRAS G12C ,
in patients with advanced colorectal cancer. *J Clin Oncol* 38, 4018–4018 (2020).

18. Lanman, B. A. *et al.* Discovery of a Covalent Inhibitor of KRASG12C (AMG 510) for the Treatment of
Solid Tumors. *J Med Chem* 63, 52–65 (2020).

19. Fakih, M. *et al.* Phase 1 study evaluating the safety, tolerability, pharmacokinetics (PK), and efficacy
of AMG 510, a novel small molecule KRAS G12C inhibitor, in advanced solid tumors. *J Clin Oncol* 37,
3003–3003 (2019).

20. Lai, A. C. & Crews, C. M. Induced protein degradation: an emerging drug discovery paradigm. *Nat Rev*
Drug Discov 16, 101–114 (2017).

21. Cromm, P. M. & Crews, C. M. Targeted Protein Degradation: from Chemical Biology to Drug
Discovery. *Cell Chem Biol* 24, 1181–1190 (2017).

22. Fellmann, C. *et al.* Functional Identification of Optimized RNAi Triggers Using a Massively Parallel
Sensor Assay. *Mol Cell* 41, 733–746 (2011).

23. Galla, M. *et al.* Avoiding cytotoxicity of transposases by dose-controlled mRNA delivery. *Nucleic Acids*
Res 39, 7147–7160 (2011).

24. Yang, W. *et al.* Genomics of Drug Sensitivity in Cancer (GDSC): a resource for therapeutic biomarker
discovery in cancer cells. *Nucleic Acids Res* 41, D955–D961 (2013).

25. Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in Unique Molecular
Identifiers to improve quantification accuracy. *Genome Res* 27, 491–499 (2017).

26. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *Embnet J*
17, 10–12 (2011).

27. Neumann, T. *et al.* Quantification of experimentally induced nucleotide conversions in high-
throughput sequencing datasets. *Bmc Bioinformatics* 20, 258 (2019).

28. Muhar, M. *et al.* SLAM-seq defines direct gene-regulatory functions of the BRD4-MYC axis. *Science*
360, 800–805 (2018).

29. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* 30, 923–930 (2014).

30. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol* 15, 550 (2014).

31. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21 (2013).

32. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a
reference genome. *Bmc Bioinformatics* 12, 323 (2011).

33. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* 43, e47–e47 (2015).

34. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights unlock linear model analysis
tools for RNA-seq read counts. *Genome Biol* 15, R29–R29 (2014).

35. Wilding, B. *et al.* Discovery of potent and selective HER2 inhibitors with efficacy against HER2 exon
20 insertion-driven tumors, which preserve wild-type EGFR signaling. *Nat Cancer* 3, 821–836 (2022).

36. Li, D. *et al.* BIBW2992, an irreversible EGFR/HER2 inhibitor highly effective in preclinical lung cancer
models. *Oncogene* 27, 4702–4711 (2008).

37. Liu, S. *et al.* Targeting HER2 Aberrations in Non–Small Cell Lung Cancer with Osimertinib. *Clin Cancer
Res* 24, 2594–2604 (2018).

38. Soria, J.-C. *et al.* Osimertinib in Untreated EGFR -Mutated Advanced Non–Small-Cell Lung Cancer.
New Engl J Med 378, 113–125 (2018).

39. Kim, T.-Y. *et al.* A phase I/II study of poziotinib combined with paclitaxel and trastuzumab in patients
with HER2-positive advanced gastric cancer. *Gastric Cancer* 22, 1206–1214 (2019).

40. Robichaux, J. P. *et al.* Pan-Cancer Landscape and Analysis of ERBB2 Mutations Identifies Poziotinib as
a Clinically Active Inhibitor and Enhancer of T-DM1 Activity. *Cancer Cell* 36, 444-457.e7 (2019).

Design and selection of ARTi-shRNAmirs

A
I. ARTi shRNA transduction
TRE^3G → GFP
(once per model)

II. ARTi target site engineering
TRE^3G → GFP
Target A
(once per target)

III. Loss-of-function experiment (fully controlled)
+Dox
TRE^3G → GFP
Target A

+Dox
TRE^3G → GFP
Control

tumor size vs time p.i. +Dox

B
miRNA seed
bits vs guide position (5' → 3')

C Knockdown reporter assay
ARTi design (6 shRNAs): ARTi.6634, ARTi.6570, ARTi.6786
Gene specific (592 shRNAs)
reporter knockdown (% of Ren.713)

D
HT-1080 RKO MV4-11 MOLM-13
%GFP vs days
Ren.713, empty, ARTi.6570, ARTi.6634, ARTi.6786, RPL9.324, PSMA3.164

E
empty, ARTi.6570, ARTi.6634, ARTi.6786, Trametinib
PC2: 15.3 % variance vs PC1: 72.1 % variance

F
ARTi.6570 — down 0, up 0
ARTi.6634 — down 0, up 0
ARTi.6786 — down 0, up 0
Trametinib — down 272, up 259
-Log₁₀ P vs Log₂ fold change

Hoffmann et al., Figure 1

Experimental validation of the ARTi approach