Deep Domain Adversarial Neural Network for the Deconvolution of Cell Type Mixtures in Tissue Proteome Profiling

Abstract

Deconvolution is a computational method for the determination/resolution of cell type proportions from bulk RNA sequence data. This technique is frequently used for the analysis of divergent cell types in tumor tissue samples. However, deconvolution technology is still in its infancy for the analysis of cell types using proteomic data sets due to challenges with repeatability/reproducibility, variable reference standards and the issues relating to single-cell proteomic reference data sets. Here, we developed a novel deep learning-based deconvolution method (scpDeconv) tailored to single-cell proteomic data sets. scpDeconv uses an autoencoder to leverage the information from bulk proteomic data to improve the quality of single-cell proteomic data, and employs a domain adversarial architecture to bridge the single-cell and bulk data distributions and transfer labels from single-cell data set to bulk data set. Extensive experiments validated the performance of scpDeconv in the deconvolution of proteomic data from various species/sources and produced using different proteomic technologies. This method should find broad applicability to areas including tumor microenvironment interpretation and clinical diagnosis/classification.

Introduction

The quantitative differentiation of cell types in a tumor microenvironment provides critical information for chemo- and immune-therapy, which is of important clinical and prognostic significance[1–3]. Bulk sequencing technology generally refers to lysing and sequencing all cells in a particular tissue, organ or organ system, and measures the average transcription/expression of genes/proteins in a cell mixture and, thus, ignores the cellular heterogeneity[4]. The development of single-cell (transcriptome/proteome) sequencing technology allows the isolation of single cells from tissue(s) for analysis[5,6]. Despite the advantages of this technology, applying it to large-scale cohorts to explore cellular heterogeneity in tumor microenvironments is impractical due to the relatively high cost and incompatibility to formalin-fixed or paraffin-embedded tissue samples[7,8]. Therefore, some computational methods have been developed to infer cell type proportions from transcription/expression profiles of genes from bulk tissue samples[7–10], providing a relatively low-cost and convenient way of studying tumor microenvironments in a clinical context.

The deconvolution of cell types from transcriptomic data from bulk tissue samples is feasible [4,8,11], and

generally assumes that the mixture of cells can be modelled as a system of linear weights equations,

therefore non-negative matrix factorization can be applied to estimate cell type proportions[8]. However, in

reality, multiple single-cell subpopulations within each cell type do not always share exactly the same

transcription/expression profile or marker genes due to the influence of cell cycle, cell state,

microenvironment and background in sequence data, where variability in these features prevents high

quality modelling using a linear distribution assumption. For this reason, alternative deep-learning-based

methods were proposed following these steps: collecting scRNA-seq data and their corresponding cell

type labels; training deep learning model with labelled scRNA-seq data as reference; and then inferring

the corresponding cell type proportions of bulk data with the trained model[9,12,13]. However, these deep

learning models were prone to overfitting on the reference datasets and failed on unseen bulk data from

another source[9]. As a remedy, some methods were proposed to finetune the model on new datasets with

known cell type proportions to improve performance[9]. However, the finetuning strategy hinders these

methods' generalization and is infeasible in real scenarios since clinical bulk data usually lack reliable

cell type proportions beforehand.

Existing deconvolution methods only focused on transcriptomics or genomics, but no attention has been

paid to proteomics. However, there is a need to develop proteome-based deconvolution algorithms due for

the following reasons: (1), proteins are the final executor of cell functions, and proteomics has

irreplaceable advantages in biological and clinical studies ranging from understanding biological

mechanisms to discovering drug targets [14]; (2), the concentration of most proteins cannot be accurately

represented by the transcription/expression level of corresponding genes due to complex post-

transcriptional regulation, RNA/protein degradation and post-translation modification[14,15];(3), there is

substantial demand for deconvolution tools for the proteomic re-analysis of microenvironments within

cancers to explore and understand tumorigenesis[16].

The advent of single-cell proteomics has greatly expanded both the number and types of proteins detected,

which has laid the foundation for the deconvolution of bulk proteomic data[17–19]. Compared with

transcriptomic data, there are a number of challenges for the analysis of proteomic data sets – which

include significant background noise, poor data quality and marked variation depending on analysis run

and/or technology. Other challenges relate to existing deconvolution methods being designed for

transcriptomic data rather than proteomic data, and include: (i) distinct distributions between proteomic

data and transcriptomic data, with most existing deconvolution methods for transcriptomic data being

based on an assumed distribution, such as a Poisson or negative binomial distribution, which are

recognized as unsuitable for proteomic data[4,11]; (ii) huge mismatch between types of proteins detected by

single cell proteomics and bulk proteomics[20,21], while these mismatched proteins are ignored by existing

deconvolution methods; (iii) for single-cell proteomic data and bulk proteomic data, "batch effects"

(variability among distributions) are pronounced, which is rarely considered by existing deconvolution

methods[22].

Here, we tackled these challenges by developing a novel deep learning architecture. Based on the

framework of mix-up, autoencoder and domain-adversarial neural network, this method can circumvent

generating huge simulation datasets, making full use of single-cell proteomic data efficiently, impute the

missing low abundant proteins with autoencoder, and establish a deconvolution module by domain

adversary training, thereby improving the model's performance in a real-world scenario. The

deconvolution approach developed gives new insights and potential to re-analyze clinical tissue proteomic

data, reveal tumor microenvironment from the proteome level, and discover new clinical diagnostic

markers.

Results

Overview of the deconvolution architecture

The overall framework of scpDeconv is illustrated in Figure 1. Firstly, for deconvolution task, we

randomly mix-up single-cell proteomic data to construct pseudo-bulk data with corresponding cell type

proportions as reference data, and use pseudo-bulk or real-bulk proteomic data as target data to be

deconvoluted.

Then an autoencoder model is applied to impute low-abundant proteins in reference data by borrowing

information from target data. To be specific, this autoencoder consists of an encoder, a predictor and a

decoder, uses both reference data and target data as input, and is trained with the reconstruction loss

calculated between the raw input and the reconstructed output and the prediction loss calculated between

the ground truth ratios and the predicted cell type proportions for reference data. In this way, by joint

training on labelled reference data and unlabelled target data, we can close the gap between reference data

and target data, improve the quality of reference data, and impute the low abundant proteins which are

missing in reference data.

Finally, the imputed reference data and raw target data are sent to a shared encoder and the extracted

features are passed through a domain discriminator for generative adversarial training, thereby facilitating

the learning of domain-invariant features by the encoder. A deconvolution head is employed to predict

cell type proportions from the extracted domain-invariant features. In this way, this method can efficiently

improve the domain generalizability to overcome diverse distributions from single-cell proteomic data

and bulk proteomic data.

Quantitative evaluation of deconvolution performance with proteome profiling

To quantitatively evaluate scpDeconv's performance on cell type mixtures as well as on cell cycle state

mixtures, we collected 8 independent single cell proteomic datasets (including 2 murine cell lines datasets

with 3 shared cell types, 5 human cell lines datasets with 4 shared cell types, and 1 human breast atlas

dataset with 6 cell types) from up-to-date technologies, and 2 bulk proteomic datasets designed for cell

cycles (including melanoma cell cycle dataset, and monocyte cell cycle dataset) (Supplementary Table 1).

We constructed pseudo-bulk proteomic data with known cell type composition as ground truth ratios from

these single-cell or pure-bulk proteomic data to rigorously quantify the performance of scpDeconv, and

then compared with four representative and state-of-the-art deconvolution methods developed for

transcriptomic data, involving deep learning based method (Scaden)[23], linear regression-based method

(MuSiC)[4], Bayes-based method (BayesPrism)[24] and distribution hypothesis based method (DestVI)[25]

(Figure 2a, Extended Data Figure 1a). Evaluating by Lin's concordance correlation coefficient (CCC),

Root Mean Square Error (RMSE), and Pearson product-moment correlation coefficient (*r*), scpDeconv

significantly surpassed Scaden, MuSiC, BayesPrism and DestVI on a broad range of proteomic datasets,

due to the rationale design of imputation to overcome the missing low abundant proteins in reference

single-cell proteomic data and the domain adversary training tailored for distribution differentiation

between reference data and target data. In the contrast, compared with human breast atlas data from the

same source, the performance of Scaden dropped in the target data with unseen distributions, such as

murine cell line data from two different sources, indicating its limited generalizability (Figure 2a,

Extended Data Figure 1a). The performance of MuSiC was hindered by its heavy dependency on gene

weights' consistency across subjects, which might be highly inconsistent between reference and target

data. BayesPrism had poor performance due to relying on parameter estimation which required high-

quality data and was unsuitable for current single-cell proteomics. By reason of the rigorous assumption

of distribution on transcriptomics, DestVI obtained the worst performance and was hard to migrate to

proteomics.

The human breast atlas is a large-scale clinical dataset with scRNA-seq data and CyTOF data sharing 33

genes/proteins and 6 cell types. To compare the distribution between transcriptomics and proteomics, we

selected paired scRNA-seq data(A) and CyTOF data(A) from the same donor(A) and CyTOF data(B)

from another donor(B). As shown in Figure 2b and Extended Data Figure 1b, there are significant

differences between the distributions of scRNA-seq data(A) and CyTOF data(A), although they were

sampled from the same donor. While the profiles of CyTOF data(A) and CyTOF data(B) are highly

similar. Furthermore, scpDeconv, Scaden and MuSiC achieved better performance on deconvoluting

pseudo-bulk proteomic data generated by CyTOF data(B) with CyTOF data(A) as reference than that with

scRNA-seq data(A) as reference, while DestVI performed better on deconvoluting proteomics with

transcriptomics as reference due to its distribution hypothesis tailored for transcriptome, and MuSiC

obtained the worst performance on both experiments, indicating the necessity of deconvoluting bulk

proteomic data with single-cell proteomic data as reference (Figure 2c-d and Extended Data Figure 1c).

We also extended scpDeconv's application to *Mus musculus*. The murine cell lines datasets contain two

independent datasets generated from two advanced single-cell proteomic technologies (N2 and nanoPOTs)

with 3 common cell types. These two datasets have less overlap in protein types (Figure 2e) and different

distributions (Extended Data Figure 1d). In this case, we used mixed N2 dataset as reference data and

mixed nanoPOTS dataset as target data to validate scpDeconv's performance on samples from different

sources (Figure 2f). Furthermore, with the changing of reference sample numbers and sizes on murine cell

lines datasets, scpDeconv achieved stable performance (Figure 2g-h). We also validated the effect of the

imputation module, and the performance dropped after removing imputation module on the deconvolution

task of murine cell lines data (Figure 2f and Extended Data Figure 1g).

The imputation step designed for proteomic dataset

Step 2 of scpDeconv is specifically designed for proteomics deconvolution, where an autoencoder

imputes missing proteins of reference data (especially the key proteins of target data). To systematically

analyze the effect of the imputation module, we integrated 3 single cell human cell lines datasets

(SCoPE2_Leduc, pSCoPE_Huffman, and pSCoPE_Leduc datasets) from data-independent acquisition

analysis mode as the reference group, and 2 single cell human cell lines datasets (T-SCP and plexDIA

datasets) from data-dependent acquisition analysis mode as the target group, which suffered from severe

batch effects (Figure 3a-b) and strikingly different protein types (Figure 3c, Extended Data Figure 2a). In

particular, the reference group missed the highly variable proteins (HVPs) specific in the target group,

which are, to some extent, informative for deconvolution task (Figure 3d and Extended Data Figure 2b).

Therefore, scpDeconv introduced imputation module in *Step 2* to tackle with this problem. The ablation

study on human cell line datasets showed that, with the help of the imputation module, adding input

features with those target specific HVPs improved the model's performance, especially in the prediction

of PDAC cell line proportion (Figure 3e-g). In addition, we also explored the effect of the number of

target HVPs on deconvolution task by changing them from 400 to 700, and found that they all achieved

better performance than deconvolution without adding target specific HVPs, and got the best CCC score

and RMSE value when target HVPs' number is 500 (Figure 3e). As shown in Extended data figure 2c-e,

after imputation module, missing features within reference data can be complemented on the basis of

target data.

Inference of cell cycle states

Besides the cell type deconvolution task, inference of cell cycle composition is also critical for studying

biological progress, tumor microenvironment, and cancer therapy[26–28]. Thanks to the ultra-high sensitivity

of mass spectrometry, proteomic technologies can detect the subtle changes of functional proteins

involved in cell cycles[29,30]. In this regard, we quantitatively evaluated scpDeconv's performance on

deconvoluting cell cycle mixtures simulated by bulk proteomic data under pure melanoma/monocyte cell

cycle states. To make model focus on cell cycles not cell types, only marker proteins corresponding to

certain cell cycles were chosen as input features (Figure 4a-b). Moreover, we noticed that marker proteins

of cell cycle states are highly conserved in melanoma cell line and monocyte cell line (Figure 4c), which

means cell cycle deconvolution across cell types/lines is feasible. For this reason, we deconvoluted the

cell cycle mixtures of melanoma with monocyte mixtures as reference and vice versa. As shown in Figure

4d-e, Extended Data Figure 3, and Extended Data Figure 4, scpDeconv achieved state-of-the-art

performance compared to other deconvolution methods for both cell cycle deconvolution experiments.

Application on clinical tissue proteomic profiling

Melanoma is a rare but invasive cancer of the skin, with high rates of growth and metastasis[31]. Previous

clinical proteomic researches of melanoma samples are limited to marker proteins and phenotypes, hard

to look into the tumor microenvironment and the effect of cellular heterogeneity on clinical outcomes[32].

Here, we explored the capacity of scpDeconv to re-analyze the representative clinical proteomic data of

174 metastatic melanoma samples at Grade IV stage and 4884 proteins with high quality[33] (Figure 5a). As

shown in Figure 5a and Extended Data Figure 5a, scpDeconv used the most advanced spatial proteomics

(Deep Visual Proteomics, DVP) on melanoma tissues as reference, which contained 7 cell classes

(vertical growth melanoma, radial growth melanoma, CD146-high melanoma, CD146-low melanoma, in

situ melanoma, melanocytes, and stroma) in the tumor microenvironment[34]. Following the procedure of

scpDeconv, the proportions of 7 cell classes were estimated from the tissue proteomic profiles of 174

melanoma patients (Figure 5b-c). And obviously, the vertical growth melanoma class was dominant in the

late stage metastasis melanoma samples, while in situ melanoma class was the minority, which is

consistent with previous clinical research, that is, the presence of vertical growth phage signifies a

melanoma acquires the capacity of metastasis[35]. We further performed survival analysis with the

predicted cell classes' proportions for 174 melanoma samples (Figure 5d-f, Extended Data Figure 5b-e).

Compared with other cell classes, the proportion of vertical growth melanoma and radial growth

melanoma exhibited significant indications for survival risk of melanoma patients. The patients with

higher vertical growth melanoma proportions tended to gain lower survival probability, while those with

higher radial growth melanoma proportions tended to gain higher survival probability, which is consistent

with previous studies, that is, in contrast to melanoma in radial growth stage, melanoma with vertical

growth pattern usually accompanies with strong aggressivity and poor prognosis[36,37].

In addition, as BRAF mutation is reported to reflect the tumorigenesis of melanoma, we further combined

the indication of predicted cell class proportions with the occurrence of BRAF mutation (Extended Data

Figure 6a). In this cohort, we can observe that BRAF mutation's indicator power is much weaker than our

predicted vertical/radial growth melanoma proportions (Extended Data Figure 6b). By grouping samples

into BRAF[(+)] (mutation) and BRAF[(-)] (non-mutation), the predicted vertical or radial growth melanoma

proportion showed more significant prognostic indication in BRAF[(+)] samples (Extended Data Figure 6c-

f).

scpDeconv's application on clinical proteome cohort indicated that tumor microenvironment could be

reflected by predicted cell type proportions, and demonstrated its potential to be a valuable tool for cancer

prognosis.

Discussion

Single-cell proteomic technology is rapidly developing and taking the center stage in biological and

clinical researches[20], paving the way for the deconvolution of tissue proteome profiling. By fully

considering the inherent characteristics of the recently developed single-cell proteomics technologies, we

developed a novel deconvolution method, scpDeconv, tailored for proteomic data. By feat of autoencoder

network, scpDeconv's imputation module can fill up missing low-abundant proteins in mixed-up single-

cell proteomic data by joint-learning on both mixed-up single-cell proteomic data and target bulk

proteomic data. Through domain-adversary training, scpDeconv can extract the domain-invariant latent

features of imputed reference data and transfer labels to target data. Rigorous and extensive experiments

evaluated scpDeconv's superiority and robustness on cell type deconvolution and inferring cell cycle

states for tissue proteomics. Furthermore, re-analysis of an existing bulk proteomic dataset from clinical

melanoma samples showed the clinical application value of scpDeconv in cancer diagnosis and prognosis.

Compared to the existing deconvolution methods tailored for transcriptomics, scpDeconv enjoys four-fold

advantages: (1), scpDeconv infers cell type proportions based on joint learning of reference data and

target data in a data-driven manner, thereby unrestricted by the prior assumptions of data distribution and

able to be flexibly applied to a broad range of proteomic data from different experiments or technology

background; (2), scpDeconv brings in domain adaptation strategy, which can efficiently tackle the batch

effect and close the distribution gaps between reference data and target data; (3), scpDeconv can impute

missing low-abundant proteins in reference data and improve the quality of reference data with the help of

imputation module; (4), we have validated that the output of scpDeconv is capable to reflect tumor

microenvironment and play roles in disease diagnosis and prognosis. However, there is still limitation of

scpDeconv. Reference proteomic data with known cell type composition needed for model training are

constructed from single-cell proteomic data with labelled cell types. While single-cell proteomics is still

in its infancy and has not been widely applied to a variety of biological samples. The single-cell

proteomic data we exhaustively collected in this study only involves a minority of cell types, thereby

limiting our method's applicability at present.

This first algorithm, tailored for the deconvolution of proteomic data using single-cell proteomic data as a

reference, will have numerous key applications including deconvolution of cell type proportions,

estimation of cell cycle states, and re-analysis of the microenvironment of multiple cancers. We hope

scpDeconv will benefit the proteomics researcher community for more valuable discoveries in biological

and clinical researches from the perspective of proteomics. With the exploration of the publicly available

single-cell proteomic atlas in the future, it will be possible to re-analyse existing tissue proteomic data

within the broader context. Furthermore, with a set of reliable microenvironment references of a vast

range of cancer types, we believe that scpDeconv will be capable of analysing all the components of

certain tumor samples at low cost and benefiting for basic medical research.

Methods

The model

The overall model architecture is shown in Figure 1, containing 4 steps: (1) Reference data preparation; (2)

Missing features imputation and data quality improvement; (3) Domain-adversarial training and

deconvolution; (4) Cell type proportion inference for target samples.

Step 1. Reference data preparation

As a supervised deconvolution method, single-cell proteomic data labelled with cell types are needed to

prepare reference data with known cell type fractions. And the bulk proteomic dataset from the same

cohort is termed the target data to be deconvoluted. The reference data and target data are usually sourced

from different sequencing platforms and experiments, thereby having gaps in distribution. Furthermore,

bulk proteomics can usually detect more proteins compared to single-cell proteomics, in which the low

abundant proteins are hard to detect.

Mix-up strategy. Mix-up is an efficient way of data augmentation by a random combination of inputs[38].

Here we employed it on single-cell proteomic data to make up pseudo-bulk proteomic data before training.

The random combination of single-cell proteomic data increases the data diversity of the reference dataset

for training, thereby improving the model's generalizability. In this step, we generated the pseudo bulk

proteomic data $\widetilde{X_r}$ with their cell type proportions $Y_r^{(1)}$ as reference data, and then sent it to the

autoencoder module for data imputation.

Data alignment. Before model training, we need to align proteins of reference data and target data for

model input. We firstly extract the proteins shared between reference and target data. And then we union

these shared proteins with HVPs specific in target data as input feature set, since the target data are

supposed to contain more proteins than the reference data, and those target specific HVPs may be

informative for deconvolution task. The missing proteins of the input feature set within reference data are

initialized to zeros and will be imputed in *Step 2*.

Step 2. Missing features imputation and data quality improvement.

Due to the immaturity of single-cell proteomics technology, the mixed-up reference data have strong

noises, large-scale missing low abundant proteins, and an obvious distribution gap with the target bulk

proteomic data. To compensate for the above obstacles caused by single-cell proteomics technology, we

employed an autoencoder module to close the gap between the reference data and the target data, impute

missing proteins of the mixed-up reference data, and improve the quality of reference data. As shown in

Figure 1b, the target data and the mixed-up reference data generated in *Step 1* are inputted into Encoder1.

And then the latent embedding generated by Encoder1 is sent to decoder to reconstruct the input. The loss

between the original input and the reconstructed output is MSE loss. The latent embedding is also sent to

a deconvolution head to estimate cell type proportions of reference data, and the L1 loss is calculated

between the predictions and its known proportions. The formulations of the architecture in *Step 2* are:

$$\text{E1}_r = \text{Encoder1}\left(\widetilde{X_r}\right) \#(1)$$

$$\widetilde{X_r}' = \text{Decoder}(\text{E1}_r) \#(2)$$

$$Y_r^{(2)} = \text{Predictor1}(E1_r) \#(3)$$

$$E1_t = \text{Encoder1}(X_t) \#(4)$$

$$X_t^{'} = \text{Decoder}(E1_t) \#(5)$$

$$\text{Loss}_{deconvolution1} = \text{loss}_{L1}\left(Y_r^{(2)}, Y_r^{(1)}\right) \#(6)$$

$$\text{Loss}_{reconstruction} = \text{loss}_{MSE}\left(\widetilde{X_r}, X_r^{'}\right) + \text{loss}_{MSE}\left(X_t, X_t^{'}\right) \#(7)$$

Where $E1_r$, $E1_t$, $Y_r^{(2)}$, $\widetilde{X_r}$, $\widetilde{X_r^{'}}$, X_t and $X_t^{'}$ are the embedding of mixed up reference data, the embedding of

the target data, the output of the predictor1 used for deconvolution, the reference proteomic data, the

output of decoder with the embedding of mixed up reference data, the target proteomic data, and the

output of decoder with the embedding of the target data, respectively. After the convergence of the

training stage, the imputed mixed-up reference data ($\widetilde{X_r^{'}}$) are used as the input of *Step 3*.

Step 3. Domain-adversarial training and deconvolution.

As shown in Figure 1c, the imputed reference data from *Step 2* and the target data from *Step 1* are sent to

the Encoder2 in *Step 3*, and the encoded features are sent to the domain discriminator and the

deconvolution head. The domain discriminator is employed to discriminate whether the feature is encoded

from the reference data or the target data. By joint training of the Encoder2 and the discriminator in this

step, the Encoder2 is prone to extracting domain-invariant features, which is suitable for the

deconvolution task with the reference data generated from different sources. The deconvolution head

consists of 2 fully connected layers and is used for predicting the cell proportions of target data. The

Encoder2 and deconvolution head are joint training with the imputed reference data as input and the

known proportions as supervision signals. The two kinds of joint training (Encoder2 and

discriminator/Encoder2 and deconvolution head) are then executed iteratively until both losses are

converged.

As for domain adaptation, we calculate BCE loss and optimize our model's weights by minimizing the

distributions between the reference and target domains. The formulations of the architecture in *Step3* are:

$$E2_r = Encoder2\left(\widetilde{X_r^{'}}\right) \#(8)$$

$$D_r = Discriminator(E2_r) \#(9)$$

$$E2_t = Encoder2(X_t) \#(10)$$

$$D_t = Discriminator(E2_t) \#(11)$$

$$Y_r^{(3)} = Predictor2(E2_r) \#(12)$$

$$Loss_{discriminator} = loss_{BCE}\left(D_t, D_t^{'}\right) + loss_{BCE}\left(D_r, D_r^{'}\right) \#(13)$$

$$Loss_{deconvolution2} = loss_{L1}\left(Y_r^{(3)}, Y_r^{(1)}\right) \#(14)$$

Where $E2_r$, $E2_t$, $Y_r^{(3)}$, D_r and D_t are the embedding of imputed reference data by Encoder2, the

embedding of the target data by Encoder2, the output of the Predictor2 used for deconvolution, the output

of discriminator with the embedding of imputed reference data, and the output of discriminator with the

embedding of target data, respectively.

Step 4. Cell type proportion inference for target samples.

As shown in Figure 1d, after domain-adversary training in *Step 3*, the trained Encoder2 and deconvolution

head are used to infer cell type proportions of target samples. The formulation of the architecture

employed in model inference is:

$$Y_t = Predictor2(E2_t) \#(15)$$

303   Where $E2_t$ is the output in *Formula* (10) with X_t as input, and Y_t is the final predicted cell type

304   proportions of the target data.

Data

The datasets used in this study are listed in Supplemental Table 1.

Human breast atlas datasets. In the human breast atlas dataset[39], scRNA-seq was performed on 52,681

cells from 16 breast tissues. CyTOF was performed on 38 breast tissues (751,970 cells, 40 protein

markers involved in breast development and tumorigenesis). The scRNA-seq and CyTOF datasets have 6

cell types in common (alveolar, hormone sensing, basal, fibroblast, vascular lymphatic, and immune

cells).

Human cell line datasets. The human cell line datasets include five different single cell proteomic datasets,

generating from T-SCP, plexDIA, pSCoPE_Leduc, SCoPE2_Leduc, and pSCoPE_Huffman,

respectively[30,40–43]. These five datasets involve 4 cell types in total, consisting of 1,755 melanoma, 434

Hela cells, 579 PDAC cells, and 163 U-937.

Murine cell line datasets. The murine cell line datasets come from two single-cell proteomic technologies,

N2 and nanoPOTS, applied in 3 identical murine cell lines (RAW 264.7, a macrophage cell line; C10, a

respiratory epithelial cell line; SVEC, an endothelial cell line)[44,45]. Murine_N2 dataset (Murine cell lines

dataset generated by N2) includes 108 cells (36 RAW cells, 36 C10 cells, and 36 SVEC cells) with 1437

proteins high quality detected. Murine_nanoPOTS dataset (Murine cell lines dataset generated by

nanoPOTS) includes 72 cells (24 RAW cells, 24 C10 cells, and 24 SVEC cells) with 1032 proteins high

quality detected. These two datasets have 762 proteins in common.

Bulk proteomic cell cycle datasets. Pure bulk proteomic datasets of melanoma G1/S/G2 and monocyte

G1/S/G2 were generated in the same experiment of pSCoPE_Leduc dataset[43]. The top 20 highly

expressed marker proteins of each cell cycle state were selected as model's input features.

Deep Visual Proteomics dataset. Deep Visual Proteomics (DVP) dataset links protein abundance of

melanoma samples to complex cellular or subcellular phenotypes while preserving spatial context,

involving average 1910 proteins[34]. This dataset contains 7 cell classes (vertical growth melanoma, radial

growth melanoma, CD146-high melanoma, CD146-low melanoma, in situ melanoma, melanocytes, and

stroma) within the tumor microenvironment of melanoma, defined by spatial location, SOX10 and CD146

staining intensity.

The clinical proteomic dataset of metastatic melanoma. The clinical proteomic dataset of metastatic

melanoma was collected from the previous study in 174 melanoma metastasis patients, with 4883 proteins

high quality detected[46].

Training details

For all cases, the batch size of training process in stage 2 and stage 3 was set to 50, and the learning rate

was set to 0.0001. The other training details, such as pseudo-sample number and pseudo-sample size (cell

number in each pseudo-sample) of reference data or target data, number of HVPs identified in target data,

and number of training epochs used in each case can be found in Supplemental Table 2.

Furthermore, in the case of murine cell lines data, to explore scpDeconv's stability, we set different

reference sample numbers (2000, 4000, and 6000) with fixed sample size, and different reference sample

sizes (5, 10, 15, and 20) with fixed sample number.

Comparison of methods

Scaden. Scaden is a deep learning-based deconvolution algorithm that uses RNA-seq data and paired

scRNA-seq data to infer the cellular composition of tissues[9]. The basic architecture of Scaden is a neural

network with 4 fully connected layers which takes the pseudo bulk RNA-seq data as input and the

predicted cell type fraction as output, using L1 loss during training to minimize the error between the

predicted cell fractions and the real cell fractions. The final Scaden model is ensembled by three best-

performing models with different parameter combinations.

MuSiC. MuSiC uses cell-type specific gene expression from scRNA-seq data to deconvolute bulk RNA-

seq data in tissues[4]. MuSiC first reweighs genes based on cross-subject variance (informativeness), and

then uses a recursively clustering strategy to close related cell types. The proportions of cell types are

estimated by non-negative least squares regression based on a series of assumptions.

BayesPrism. BayesPrism is a Bayesian deconvolution method for cancer tissues depending on patient-

derived scRNA-seq as prior information[24]. The Bayesian algorithm models prior distribution learned from

the paired scRNA-seq data and then infers joint posterior distribution of cell proportions as the

deconvolution results.

DestVI. DestVI is a Bayesian model that learns cell type specific profiles and latent variations using a

conditional deep generative model[25]. The method assumes that transcriptomic data follows a negative

binomial distribution to model the reference scRNA-seq data and spatial RNA-seq data.

Evaluation experiments

Cell type mixture decomposition with proteome profiling.

As for testing the deconvolution performance with reference data and target data from different modalities,

we deconvoluted the simulated bulk proteomic data constructed from CyTOF data with scRNA-seq data

in the same cohort as reference or with CyTOF data in the same cohort as reference in the human breast

atlas dataset, splitting the dataset into reference and target data by donor.

There are apparent batch effects between different proteomic technologies and a huge gap between the

distributions of bulk proteomic data and single-cell proteomic data. Therefore, we evaluated scpDeconv's

deconvolution performance on the simulated bulk proteomic data constructed from murine_nanoPOTS

dataset with the mixed-up murine_N2 dataset as reference to mimic the situation where the bulk/single

cell proteomic data come from totally different proteomic technologies and different experiments.

Furthermore, we employed the integrated human cell line datasets to evaluate scpDeconv's performance

on integrated multi-centre datasets. The Melanoma, PDAC, U-937, and Hela cell lines appeared in

different experiments were selected for testing.

Systematic analysis.

To explore the efficiency of the imputation of missing proteins in *Step 2*, we executed cell type

deconvolution experiments on the integrated human cell line datasets with adding target specific HVPs or

without adding target specific HVPs, and also compared deconvolution performance between model with

imputation stage and model without imputation stage. For the experiments without adding target specific

HVPs, we only used the overlap proteins from both reference and target datasets as input features. For the

experiments with adding target specific HVPs, we changed the number of HVP in target data from 400 to

700.

Application

Cell cycle inference with proteome profiling.

The cell cycles reflect the replication process of cancer cells and can be represented by the subtle changes

of cell-cycle-relevant proteins. We collected the bulk proteomic data of melanoma or monocyte samples

with pure cell subpopulations within G1, S, and G2 stage. Then the melanoma cell cycle mixtures were

constructed from the melanoma cell subpopulations within G1, S, and G2 stages, and monocyte cell cycle

mixtures were constructed from the monocyte subpopulations within G1, S, and G2 stages. We

deconvoluted the melanoma cell cycle mixtures with monocyte pure cell cycle subpopulations as

reference, and deconvoluted the monocyte cell cycle mixtures with melanoma pure cell cycle

subpopulations as reference. In this way, we can estimate scpDeconv's performance on the cell cycle

inference task across cell types/cell lines.

Investigation of microenvironment and clinical outcomes with clinical tissue proteome profiling.

The establishment of scpDeconv tailored for cell type deconvolution on proteomic data opens up an

opportunity to mine the tumor microenvironment of previously released clinical tissues with provided

bulk proteomic data. We deconvoluted the proteomic data from metastasis melanoma cohort and analysed

the role of cell type proportions in cancer prognosis.

Evaluation metrics

401   The linear concordance between ground truth ratios and predicted proportions of cell types or cell states is

402   estimated by Lin's concordance correlation coefficient (CCC), Root Mean Square Error (RMSE), and

403   Person product-moment correlation coefficient (*r*). The formulas are as follows,

$$CCC(y, \hat{y}) = \frac{2r\sigma_y\sigma_{\hat{y}}}{\sigma_y^2 + \sigma_{\hat{y}}^2 + (\mu_y - \mu_{\hat{y}})} \#(16)$$

$$RMSE(y, \hat{y}) = \sqrt{avg(y - \hat{y})^2} \#(17)$$

$$r(y, \hat{y}) = \frac{cov(y, \hat{y})}{\sigma_y\sigma_{\hat{y}}} \#(18)$$

404   Where y are the ground truth cell type (or cell cycle) proportions, $\hat{y}$ are the prediction of cell type (or cell

405   cycle) proportions, σ_y is the standard deviation of y, $\sigma_{\hat{y}}$ is the standard deviation of $\hat{y}$, $cov(y, \hat{y})$ is the

406   covariance of y and $\hat{y}$, and μ_y, $\mu_{\hat{y}}$ respectively are the mean of y and $\hat{y}$.

Statistical analysis

408   The group comparisons are estimated by the two-sided Wilcoxon test. Kaplan-Meier survival analysis is

409   conducted to assess if the predicted cell type proportions are associated to outcomes (overall survival) in

410   the clinical cohort. The P values are calculated using the log-rank test and are considered statistically

411   significant if they are two-tailed less than 0.05.

Biological analysis

413   Highly variable proteins are selected by Scanpy[47]. The Gene Ontology (GO) and pathway enrichment are

414   performed using DAVID platform[48].

Data Availability

416   All data used in this study are publicly available and the usages were fully illustrated in the Method

417   section.

Code Availability

419   The codes are implemented in Python and released on https://github.tencentailab/scpDeconv with detailed

420   instruction. The comparison methods are implemented under their official guidance and the reference

421   codes are listed in the Method section.

Acknowledgements

Author Contributions

F.Y. and J.Y. conceived and designed the project. F.W. developed the algorithm and conducted

experiments under the supervision of F.Y. and J.Y.. F.W. and F.Y. analysed the results. F.Y. and F.W.

write the manuscript. F.W. finish the figure under the guidance of F.Y. and J.Y.. J.Y., G.W., J.Q. revised

the manuscript. L.H. gave suggestion for the domain adaptation and improvement of the manuscript. J.S.

gave suggestion for the project. All authors reviewed and approved the manuscript. F.W. and F.Y.

contributed equally.

Competing Interests

The authors declare no competing interests.

Figure Legends

Figure 1. The architecture of our proposed method.

a, Stage 1: preparation of training data. The mix-up process that combines the proteome profiling of
different cell types by random proportions.

b, Stage 2: domain transfer. The encoder-decoder scheme to close the gap between the reference data
and the target data, and imputation of the mixed up single-cell proteomic data generated in **Stage 1**.
Reconstruction loss is applied for model training.

c, Stage 3: deconvolution model training. The domain-adversary training architecture of encoder. The
domain discriminator is used to discriminate whether the latent feature is encoded from the mixed up
reference data or from the target bulk proteomic data. The deconvolution head is used to predict the
proportions of cell type.

d, Stage 4:inference stage. Target bulk proteomic data are inputted to the trained encoder and the

deconvolution head to infer the proportions of cell types.

Figure 2. Overall performance comparation of cell type (or cell cycle state) mixture

deconvolution.

a, CCC (top) and RMSE values (bottom) of scpDeconv, Scaden, MuSiC, BayesPrism, and DestVI on

human breast atlas datasets, murine cell lines datasets, human cell lines datasets, and cell cycle datasets.

Boxplots show the median (center lines), interquartile range (hinges), and 1.5 times the interquartile range

(whiskers).

b, Correlation plots between donor A's proteome and donor A's transcriptome (left) or donor A's

proteome and donor B's proteome (right) within 6 cell types studied in human breast atlas datasets. AV:

alveolar, HS: hormone sensing, BA: basal, FI: fibroblast, VA: vascular lymphatic, IM: immune cells.

c, Scatter plots of true (y axis) and predicted cell type proportions (x axis) by deconvoluting pseudo-bulk

proteomic data from donor B with single-cell transcriptomic data from donor A as reference (top) or with

single-cell proteomic data from donor B as reference (bottom) in human breast atlas datasets.

d, CCC values of scpDeconv and the comparison methods by deconvoluting pseudo-bulk proteomic data

from donor B with single-cell transcriptomic data from donor A as reference (triangle) or with single-cell

proteomic data from donor B as reference (circle) in human breast atlas datasets.

e, Venn diagram showing the overlap of proteins between two murine cell line datasets (N2 dataset and

nanoPOTS dataset).

f, Scatter plots of true (y axis) and predicted proportions (x axis) of three cell types (C10, SVEC, and

RAW) in murine cell lines datasets using scpDeconv.

g, CCC values of scpDeconv on murine cell lines datasets with different reference sample numbers in

stage 1.

h, CCC values of scpDeconv on murine cell lines datasets with different reference sample sizes in stage 1.

Figure 3. Deconvolution performance on integrated human cell lines datasets with different

HVP numbers.

a, Integrated t-SNE plots colored by data sources (left panel) and by cell types (right panel) of the

reference group (SCoPE2_Leduc, pSCoPE_Huffman, and pSCoPE_Leduc datasets).

b, Integrated t-SNE plots colored by data sources (left panel) and by cell types (right panel) of the target

group (T-SCP and plexDIA datasets).

c, Venn diagram showing the overlap of proteins from five different human cell lines datasets.

d, Venn diagram showing the overlap between shared proteins in all datasets and highly variable proteins

(HVPs) in target group. Heatmap showing the protein expression of specific HVPs in the target group.

e, CCC (left panel) and RMSE values (right panel) of scpDeconv with different target HVP numbers on

integrated human cell line datasets.

f, Scatter plots of true (y axis) and predicted proportions (x axis) of four cell types (Melanoma, PDAC, U-

937, and Hela cell) in human cell lines datasets using scpDeconv without adding target specific HVPs.

g, Scatter plots of true (y axis) and predicted proportions (x axis) of four cell types (Melanoma, PDAC,

U-937, and Hela cell) in human cell lines datasets using scpDeconv with setting target HVP number as

500.

Figure 4. Inference of cell cycle across cell lines.

a, Heatmap of cell cycle specific marker proteins in melanoma and monocytes.

b, GO enrichment of marker proteins of G1, S, and G2 phages.

c, Heatmap showing the correlation of marker proteins' abundance in melanoma and monocytes.

d, Scatter plots of true (y axis) and predicted proportions (x axis) of three cell cycle states (G1, S, and G2)

by deconvoluting monocytes (mon) pseudo samples with melanoma (mel) sample as reference (top) and

vise versa (bottom) using scpDeconv.

e, CCC (top) and RMSE values (bottom) of scpDeconv and the comparison methods by deconvoluting

monocytes pseudo samples with melanoma samples as reference (circle) and vise versa (triangle).

Figure 5. Application on clinical tissue proteomic data of metastasis melanoma.

a, The pipeline of applying scpDeconv on deconvoluting clinical tissue proteome of metastasis melanoma

patients with melanoma DVP dataset as reference.

b, Heatmap of proteomic profile (top) and histogram of predicted cell class proportions (bottom) for each

sample in clinical melanoma dataset.

c, Distribution of predicted cell class proportions of melanoma samples. Box plots show the median

(center lines), interquartile range (hinges), and 1.5 times the interquartile range (whiskers).

d, KM plot of the predicted proportions of vertical growth melanoma for 174 melanoma patients.

e, KM plot of the predicted proportions of radial growth melanoma for 174 melanoma patients.

f, KM plot of the predicted proportions of in situ melanoma for 174 melanoma patients.

Extended Data Figure 1. Overall performance comparation of cell type (or cell cycle state)
mixture deconvolution.

a, Pearson's correlation values of scpDeconv, Scaden, MuSiC, BayesPrism, and DestVI on human breast
atlas datasets, murine cell lines datasets, human cell lines datasets, and cell cycle datasets. Boxplot show
the median (center lines), interquartile range (hinges), and 1.5 times the interquartile range (whiskers).

b, Heatmap of transcriptomic/proteomic profile from donor A and proteomic profile from donor B on 33
shared markers (x axis) and 6 cell types (y axis) studied in human breast atlas datasets. AV: alveolar, HS:
hormone sensing, BA: basal, FI: fibroblast, VA: vascular lymphatic, IM: immune cells.

c, RMSE values of scpDeconv and the comparison methods by deconvoluting pseudo-bulk proteomic
data from donor B with single-cell transcriptomic data from donor A as reference (triangle) or with
single-cell proteomic data from donor B as reference (circle) in human breast atlas datasets.

d, t-SNE plot showing the distribution of N2 dataset and nanoPOTS dataset.

e, RMSE values of scpDeconv on murine cell lines datasets with different reference sample numbers in
stage 1.

f, RMSE values of scpDeconv on murine cell lines datasets with different reference sample sizes in stage
1.

g, Scatter plots of true (y axis) and predicted proportions (x axis) of three cell types (C10, SVEC, and
RAW) in murine cell lines datasets after removing stage 2 of scpDeconv.

Extended Data Figure 2. Systematical analysis of missing feature imputation in human cell
lines datasets.

a, Venn diagram showing the overlap of proteins from different sources in the reference group (left) or
target group (middle), and the overlap between reference groups' shared proteins and target groups'
shared proteins (right) in the human cell lines datasets.

b, Heatmap showing the z-score of target specific HVPs in the target group.

c, Heatmap showing the raw expression of all proteins and missing features in reference pseudo-bulk data

after stage 1 of scpDeconv. Proteins and samples were ordered by hierarchical clustering on raw

expression.

d, Heatmap showing the imputed expression of all proteins and missing features in reference pseudo-bulk

data after stage 2 of scpDeconv. Proteins and samples kept the same order with above.

e, Heatmap showing the raw expression of all proteins in the target data. Proteins kept the same order

with above and samples were ordered by hierarchical clustering.

Extended Data Figure 3. Performance of cell cycle state inference by deconvoluting

monocytes pseudo samples with melanoma samples as reference.

a, Scatter plot of true (y axis) and predicted cell cycle proportions (x axis) of all stages, G1 stage, S stage,

and G2 stage for monocytes pseudo samples (from left to right) by scpDeconv with melanoma samples as

reference.

b, Scatter plot of true (y axis) and predicted cell cycle proportions (x axis) of all stages, G1 stage, S stage,

and G2 stage for monocytes pseudo samples (from left to right) by Scaden with melanoma samples as

reference.

c, Scatter plot of true (y axis) and predicted cell cycle proportions (x axis) of all stages, G1 stage, S stage,

and G2 stage for monocytes pseudo samples (from left to right) by MuSiC with melanoma samples as

reference.

Extended Data Figure 4. Performance of cell cycle state inference by deconvoluting

melanoma pseudo samples with monocytes as reference.

a, Scatter plot of true (y axis) and predicted cell cycle proportions (x axis) of all stages, G1 stage, S stage,

and G2 stage for melanoma pseudo samples (from left to right) by scpDeconv with monocytes samples as

reference.

b, Scatter plot of true (y axis) and predicted cell cycle proportions (x axis) of all stages, G1 stage, S stage,

and G2 stage for melanoma pseudo samples (from left to right) by Scaden with monocytes samples as

reference.

c, Scatter plot of true (y axis) and predicted cell cycle proportions (x axis) of all stages, G1 stage, S stage,

and G2 stage for melanoma pseudo samples (from left to right) by MuSiC with monocytes samples as

reference.

Extended Data Figure 5. Survival analysis of the predicted cell class proportions of clinical melanoma samples.

a, Three cases showing the scpDeconv's pipeline on deconvoluting clinical tissue proteome of metastasis melanoma patients.

b, KM plot of the predicted proportions of CD146-high melanoma for 174 melanoma patients.

c, KM plot of the predicted proportions of CD146-low melanoma for 174 melanoma patients.

d, KM plot of the predicted proportions of melanocytes for 174 melanoma patients.

e, KM plot of the predicted proportions of stroma for 174 melanoma patients.

Extended Data Figure 6. Comprehensive survival analysis by combining BRAF mutation and the predicted cell class proportions of clinical melanoma samples.

a, Distribution of predicted cell class proportions of melanoma samples grouped by BRAF mutation. Boxplots show the median (center lines), interquartile range (hinges), and 1.5 times the interquartile range (whiskers).

b, Hazard ratios of BRAF mutation, vertical growth melanoma and radial growth melanoma in melanoma patients.

c, KM plot of the predicted proportions of vertical growth melanoma for melanoma samples without BRAF mutation.

d, KM plot of the predicted proportions of radial growth melanoma for melanoma samples without BRAF mutation.

e, KM plot of the predicted proportions of vertical growth melanoma for melanoma samples with BRAF mutation.

f, KM plot of the predicted proportions of radial growth melanoma for melanoma samples with BRAF mutation.

References

1.   Galon, J. *et al.* Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science (1979)* **313**, 1960–1964 (2006).

2.   Ino, Y. *et al.* Immune cell infiltration as an indicator of the immune microenvironment of pancreatic cancer. *British Journal of Cancer 2013 108:4* **108**, 914–923 (2013).

3.   Shang, B., Liu, Y., Jiang, S. J. & Liu, Y. Prognostic value of tumor-infiltrating FoxP3+ regulatory T
cells in cancers: a systematic review and meta-analysis. *Scientific Reports 2015 5:1* **5**, 1–9 (2015).

4.   Wang, X., Park, J., Susztak, K., Zhang, N. R. & Li, M. Bulk tissue cell type deconvolution with multi-
subject single-cell expression reference. *Nature Communications 2019 10:1* **10**, 1–9 (2019).

5.   Qian, J. *et al.* A pan-cancer blueprint of the heterogeneous tumor microenvironment revealed by
single-cell profiling. *Cell Research 2020 30:9* **30**, 745–762 (2020).

6.   Azizi, E. *et al.* Single-Cell Map of Diverse Immune Phenotypes in the Breast Tumor
Microenvironment. *Cell* **174**, 1293-1308.e36 (2018).

7.   Frishberg, A. *et al.* Cell composition analysis of bulk genomics using single-cell data. *Nature*
Methods 2019 16:4 **16**, 327–332 (2019).

8.   Newman, A. M. *et al.* Determining cell type abundance and expression from bulk tissues with
digital cytometry. *Nature Biotechnology 2019 37:7* **37**, 773–782 (2019).

9.   Menden, K. *et al.* Deep learning-based cell composition analysis from tissue expression profiles.
Sci Adv **6**, (2020).

10.   Cable, D. M. *et al.* Robust decomposition of cell type mixtures in spatial transcriptomics. *Nature*
Biotechnology 2021 40:4 **40**, 517–526 (2021).

11.   Patrick, E. *et al.* Deconvolving the contributions of cell-type heterogeneity on cortical gene
expression. *PLoS Comput Biol* **16**, e1008120 (2020).

12.   Lv, X. *et al.* Deconvolute Gene Expression Based on Deep Learning in Scrna-Seq. *Proceedings -*
2021 17th International Conference on Computational Intelligence and Security, CIS 2021 70–74
(2021) doi:10.1109/CIS54983.2021.00023.

13.   Lin, Y. *et al.* DAISM-DNNXMBD: Highly accurate cell type proportion estimation with in silico data
augmentation and deep neural networks. *Patterns* **3**, 100440 (2022).

14.   Marguerat, S. *et al.* Quantitative Analysis of Fission Yeast Transcriptomes and Proteomes in
Proliferating and Quiescent Cells. *Cell* **151**, 671–683 (2012).

15.   Gygi, S. P., Rochon, Y., Franza, B. R. & Aebersold, R. Correlation between Protein and mRNA
Abundance in Yeast. *Mol Cell Biol* **19**, 1720–1730 (1999).

16.   Edwards, N. J. *et al.* The CPTAC data portal: A resource for cancer proteomics research. *J*
Proteome Res **14**, 2707–2713 (2015).

17.   Strack, R. Spatial proteomics with subcellular resolution. *Nature Methods 2022 19:7* **19**, 780–780
(2022).

18.   Petelski, A. A. *et al.* Multiplexed single-cell proteomics using SCoPE2. *Nature Protocols 2021*
16:12 **16**, 5398–5425 (2021).

19.   Woo, J. *et al.* High-throughput and high-efficiency sample preparation for single-cell proteomics
using a nested nanowell chip. *Nature Communications 2021 12:1* **12**, 1–11 (2021).

20.   Perkel, J. M. Single-cell proteomics takes centre stage. *Nature* **597**, 580–582 (2021).

21.   Doerr, A. Single-cell proteomics. *Nature Methods 2018 16:1* **16**, 20–20 (2018).

22.   Vanderaa, C. & Gatto, L. Replication of single-cell proteomics data reveals important
computational challenges. *https://doi.org/10.1080/14789450.2021.1988571* **18**, 835–843 (2021).

23.   Menden, K. *et al.* Deep learning–based cell composition analysis from tissue expression profiles.
Sci Adv **6**, (2020).

24.   Chu, T., Wang, Z., Pe'er, D. & Danko, C. G. Cell type and gene expression deconvolution with
BayesPrism enables Bayesian integrative analysis across bulk and single-cell RNA sequencing in
oncology. *Nature Cancer 2022 3:4* **3**, 505–517 (2022).

25.   Lopez, R. *et al.* DestVI identifies continuums of cell types in spatial transcriptomics data. *Nature*
Biotechnology 2022 40:9 **40**, 1360–1369 (2022).

26.   Warrener, R. *et al.* Tumor cell-specific cytotoxicity by targeting cell cycle checkpoints. *The FASEB*
Journal **17**, 1–21 (2003).

27.   Li, J. & Stanger, B. Z. Cell Cycle Regulation Meets Tumor Immunosuppression. *Trends Immunol* **41**,
859–863 (2020).

28.   Schwartz, G. K. & Shah, M. A. Targeting the cell cycle: A new approach to cancer therapy. *Journal*
of Clinical Oncology **23**, 9408–9421 (2005).

29.   Brunner, A.-D. *et al.* Ultra-high sensitivity mass spectrometry quantifies single-cell proteome
changes upon perturbation. *Mol Syst Biol* **18**, e10798 (2022).

30. Leduc, A., Huffman, R. G. & Slavov, N. Droplet sample preparation for single-cell proteomics
applied to the cell cycle. doi:10.1101/2021.04.24.441211.

31. Balch, C. M. *et al.* Final Version of 2009 AJCC Melanoma Staging and Classification. *Journal of*
Clinical Oncology **27**, 6199 (2009).

32. Betancourt, L. H. *et al.* The Human Melanoma Proteome Atlas—Complementing the melanoma
transcriptome. *Clin Transl Med* **11**, e451 (2021).

33. Beck, L. *et al.* Clinical proteomics of metastatic melanoma reveals profiles of organ specificity and
treatment resistance. *Clinical Cancer Research* **27**, 2074–2086 (2021).

34. Mund, A. *et al.* Deep Visual Proteomics defines single-cell identity and heterogeneity. *Nature*
Biotechnology 2022 40:8 **40**, 1231–1240 (2022).

35. Crowson, A. N., Magro, C. M. & Mihm, M. C. Prognosticators of melanoma, the melanoma report,
and the sentinel lymph node. *Modern Pathology 2006 19:2* **19**, S71–S87 (2006).

36. Prognosis and survival for melanoma skin cancer | Canadian Cancer Society.
https://cancer.ca/en/cancer-information/cancer-types/skin-melanoma/prognosis-and-survival.

37. Ciarletta, P., Foret, L. & Amar, M. ben. The radial growth phase of malignant melanoma: multi-
phase modelling, numerical simulations and linear stability analysis. *J R Soc Interface* **8**, 345
(2011).

38. Zhang, H., Cisse, M., Dauphin, Y. N. & Lopez-Paz, D. mixup: Beyond Empirical Risk Minimization.
Preprint at https://github.com/facebookresearch/mixup-cifar10. (2022).

39. Gray, G. K. *et al.* A human breast atlas integrating single-cell proteomics and transcriptomics. *Dev*
Cell **57**, 1400-1420.e7 (2022).

40. Brunner, A. *et al.* Ultra-high sensitivity mass spectrometry quantifies single-cell proteome
changes upon perturbation. *Mol Syst Biol* **18**, (2022).

41. Derks, J. *et al.* Increasing the throughput of sensitive proteomics by plexDIA. *Nature*
Biotechnology 2022 1–10 (2022) doi:10.1038/s41587-022-01389-w.

42.   Huffman, R. G. *et al.* Prioritized single-cell proteomics reveals molecular and functional
666   polarization across primary macrophages. *bioRxiv* 2022.03.16.484655 (2022)
667   doi:10.1101/2022.03.16.484655.

668   43.   Leduc, A., Huffman, R. G. & Slavov, N. Droplet sample preparation for single-cell proteomics
669   applied to the cell cycle. doi:10.1101/2021.04.24.441211.

670   44.   Dou, M. *et al.* High-Throughput Single Cell Proteomics Enabled by Multiplex Isobaric Labeling in a
671   Nanodroplet Sample Preparation Platform. *Anal Chem* **91**, 13119–13127 (2019).

672   45.   Woo, J. *et al.* High-throughput and high-efficiency sample preparation for single-cell proteomics
673   using a nested nanowell chip. *Nature Communications 2021 12:1* **12**, 1–11 (2021).

674   46.   Beck, L. *et al.* Clinical Proteomics of Metastatic Melanoma Reveals Profiles of Organ Specificity
675   and Treatment Resistance. doi:10.1158/1078-0432.CCR-20-3752.

676   47.   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: Large-scale single-cell gene expression data
677   analysis. *Genome Biol* **19**, 1–5 (2018).

678   48.   Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional
679   annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216–W221 (2022).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

ED Figure 1

ED Figure 2

ED Figure 3

ED Figure 4

ED Figure 5