Arabidopsis transcriptome responses to low water potential using high throughput plate assays

Stephen Gonzalez [1*], Joseph Swift [1*†], Adi Yaaran [2], Jiaying Xu [1], Charlotte Miller [1], Natanella Illouz-Eliaz [1], Joseph R. Nery [3], Wolfgang Busch [1], Yotam Zait [2], and Joseph R. Ecker [1,3,4†]

1. Plant Biology Laboratory, The Salk Institute for Biological Studies, 10010 N Torrey Pines Rd, La Jolla, CA 92037, United States.
2. The Robert H. Smith Institute of Plant Sciences and Genetics in Agriculture, The Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot, Israel.
3. Genomic Analysis Laboratory, The Salk Institute for Biological Studies, 10010 N Torrey Pines Rd, La Jolla, CA 92037, United States.
4. Howard Hughes Medical Institute, The Salk Institute for Biological Studies, 10010 N Torrey Pines Rd, La Jolla, CA 92037, United States.

* These authors contributed equally to this work

† corresponding author

Abstract

Soil-free assays that induce water stress are routinely used to investigate drought responses in the plant *Arabidopsis thaliana*. Due to their ease of use, the research community often relies on polyethylene glycol (PEG), mannitol and salt (NaCl) treatments to reduce the water potential of agar media, and thus induce drought conditions in the laboratory. However, while these types of stress can create phenotypes that resemble those of water deficit experienced by soil-grown plants, it remains unclear how these treatments compare at the transcriptional level. Here, we demonstrate that these different methods of lowering water potential elicit both shared and distinct transcriptional responses in *Arabidopsis* shoot and root tissue. When we compared these transcriptional responses to those found in *Arabidopsis* roots subject to vermiculite drying, we discovered many genes induced by vermiculite drying were repressed by low water potential treatments on agar plates (and vice versa). Additionally, we also tested another method for lowering water potential of agar media. By increasing the nutrient content and tensile strength of agar, we show the 'hard agar' (HA) treatment can be leveraged as a high-throughput assay to investigate natural variation in *Arabidopsis* growth responses to low water potential.

Introduction

As climate change advances, improving crop drought tolerance will be key for ensuring food security (*1, 2*). This has led to intense research at the molecular level to find novel loci and alleles that drive plant responses to drought conditions. Such investigations benefit from simple assays that can reproduce drought phenotypes at both the physiological and molecular levels. While some researchers use soil-based assays, these are cumbersome. For example, extracting intact root systems from the soil is difficult, and reproducing the rate at which water evaporates from the soil can be challenging (*3*). In light of this, chemical agents such as polyethylene glycol (PEG), mannitol, or salt (NaCl) are often employed to induce drought stress. When present in aqueous or agar media, they allow precise and dose-dependent control of water potential (*4, 5*). When exposed to these media types, plants exhibit the hallmarks of drought physiology, such as reduced growth rate, reduced stomatal conductance, and increased leaf senescence (*4, 6, 7*). While each of these methods lower water potential and thus induce a drought stress, each method exerts additional and distinct effects. For example NaCl not only induces osmotic stress, but can cause salt toxicity (*6*). Since it is not metabolized by most plants mannitol is considered less toxic (*3*), however evidence suggests it may act as a signaling molecule (*5, 8*). Since both NaCl and mannitol can enter the pores of plant cell walls (*9, 10*), they can induce plasmolysis, a process that does not typically occur under mild water deficit (*9*). Due to its higher molecular weight, PEG treatment avoids this, and instead causes cytorrhysis (*9*), a physiology more common under drought settings (*10, 11*).

The unique impacts PEG, mannitol and NaCl have on plant physiology may also extend to the level of gene expression. Indeed, a broad spectrum of transcriptional changes are documented in response to low water potential, which may be attributed to the specific method employed (*4, 12, 13*). Here, we examine the transcriptional responses to PEG, mannitol and NaCl in *Arabidopsis*, and compare these responses to those elicited when plants are exposed to vermiculite drying. Furthermore, we explore a new approach for reducing water potential.

Comparing differential gene expression responses elicited by PEG, mannitol and NaCl treatment to vermiculite drying

To understand the impact PEG, mannitol and NaCl treatment have on gene expression, we first tested their physiological effects across a range of doses. To this end, we grew *Arabidopsis* seedlings on agar plates supplemented with Linsmaier & Skoog (LS) nutrients for 14 days on three different doses of each stress type. Dose ranges were chosen based on published literature, and ranged from mild to severe stress levels (*3, 4, 11*). As the dose of each stress type increased, the media's water potential significantly decreased in a dose-dependent manner (Pearson, $p < 8.5 \times 10^{-5}$). Across the doses tested, we found that each stress type's impact on water potential was not statistically different from one other (ANCOVA post-hoc, $p > 0.05$). In response to these treatments, we found shoot biomass significantly decreased in a dose-dependent manner (Pearson, $p < 2 \times 10^{-6}$) (**Figure 1A - C**, **Figure 1-figure supplement 1**, **Supplementary File 1**).

Genes that change their expression in response to an environmental signal often do so in a dose-responsive manner (*4, 14*). In light of this, we sought to discover genes whose expression was dose-responsive to the amount of PEG, mannitol, or NaCl applied. By identifying stress-responsive genes across a range of doses, we ensured such genes responded to and were directional with the stress as a whole and not induced or repressed at an individual dose. Taking this approach, we sequenced root and shoot bulk transcriptomes by RNA-seq, and associated each gene's expression with the dose of stress with a linear model. To ensure we captured steady-state differences in gene expression, and avoided trainsent ones, we sequenced root and shoot transcriptome profiles after 14 days of stress exposure (*3, 6, 15*). By these means, we found hundreds of genes that were dose-responsive to each treatment within root and shoot tissue (**Figure 1E - 1H, Figure 1-figure supplement 2, Supplementary File 2**) (adj. $p < 0.05$). We found that a portion of these dose-responsive genes were shared across treatments, suggesting a common response to low water potential (**Figure 1D**, **Figure 1-supplement Figure 3**). Conversely, we also found a portion of dose-responsive genes were unique to each stress type.

Next, we wanted to compare these different methods of lowering water potential to a pot-based water deficit assay. To perform this experiment, we subjected mature *Arabidopsis* plants grown in pots on vermiculite supplemented with LS media to mild water stress by withholding water for 5 days. During this period, field capacity was reduced from 100 % to 41 %. This treatment led to

a reduction in plant biomass ($p = 1.8 \times 10^{-3}$), as well as seed yield ($p = 1.2 \times 10^{-4}$), but did not

induce visible signs of senescence or wilting (**Figure 1-figure supplement 4**, **Supplementary**

File 3). We assayed root and shoot gene expression responses daily during water loss by RNA-

seq. We observed a dose-dependent relationship between a decrease in field capacity and gene

expression responses in both roots and shoots, identifying 1,949 differentially expressed genes

in roots and 1,792 in shoots (adj. $p < 0.01$) (**Figure 1I, Figure 1-figure supplement 2,**

Supplementary File 2). We ensured these genes' expression patterns recovered upon

rewatering (**Figure 1I**). We note that while vermiculite has greater aeration than soil, we found

that the genes differentially expressed in roots in response to vermiculite drying largely agreed

with a previous report detailing transcriptional responses to soil drying (*16*) (**Figure 2-figure**

supplement 1). We also found differentially expressed genes responsive to vermiculite drying

agreed with those responsive to transient treatment with Abscisic Acid (ABA), a stress hormone

whose levels rise in response to water deficit (*4*) (**Figure 2-figure supplement 2**).

To assess how PEG, mannitol and NaCl treatments compared to the vermiculite drying response

described above, we overlapped genes found differentially expressed in each experiment. For

shoot tissue, we found genes that were differentially expressed during vermiculite drying

overlapped significantly with genes that were differentially expressed by either PEG, mannitol and

NaCl treatments (Fisher test, adj. $p < 0.05$). Additionally, there was 88 – 99 % directional

agreement within these overlaps, indicating that genes induced or repressed by vermiculite drying

were similarly induced or repressed by low water potential treatments on agar (**Figure 2A** & **2B**).

Along these lines, across all conditions we saw differential expression of the desiccation-

associated genes *RESPONSE TO DESSICATION 20;29B (RD20;29B) (17, 18)*, the osmo-

protectant gene *DELTA1-PYRROLINE-5-CARBOXYLATE SYNTHASE 1* (*P5CS1*), and ABA

signaling   and   biosynthesis   genes   *HOMEOBOX   7*   (HB7)   (*19*)   and   *NINE-CIS-*

EPOXYCAROTENOID DIOXYGENASE (*NCED3) (20*) (**Figure 2-figure supplement 3**). We note

that while we observed agreement in the direction of gene expression across assays, there were

differences in the amplitude of gene expression (**Figure 2-figure supplement 3**). This may be

due to confounding factors, such as differences in the ranges of water potential tested (**Figure**

1B), or through comparing seedlings grown on plates with mature *Arabidopsis* plants grown on

vermiculite.

In root tissue, we found greater variability in transcriptomic responses to the different methods of

lowering water potential. In particular, we found a number of genes that were upregulated during

vermiculite drying were downregulated by PEG, mannitol and NaCl treatments (and vice versa)
(**Figure 2A**). This trend persisted when we assessed genes found differentially expressed at each
discrete dose of stress (**Figure 2B**). For example, 27 % of PEG dose-responsive genes shared
the same direction of expression seen in vermiculite drying responses. We note that previously
published PEG transcriptome datasets largely agreed with our own (**Figure 2-figure supplement**
1). Such differential regulation in comparison to vermiculite drying is exemplified by the expression
of genes such as *HOMEOBOX 12* (*HB12*) (*19*) (**Figure 2C**), *GRC2-LIKE 1* (*GCL1*) (*21*), and
RESPONSE TO DEHYDRATION 21 (*RD21*) (*22*) (**Figure 2-figure supplement 3**). We found
genes downregulated by PEG are over-represented in the 'monooxygenase activity', and 'oxygen
binding' Gene Ontology (GO) Terms ($p < 1 \times 10^{-15}$, **Supplementary File 4**). Mannitol and NaCl
held a 48 % and 57 % agreement in gene expression direction with vermiculite drying respectively.
Examples of genes that followed this pattern of differential regulation in mannitol and NaCl
treatments were *DROUGHT HYPERSENSITIVE 2* (*DRY2*) (*23*) (**Figure 2D**) and *ROOT HAIR*
SPECIFIC 18 (RHS18) (*24*). NaCl-responsive GO Terms included a specific downregulation of
'phosphorous metabolic processes' ($p = 5.2 \times 10^{-6}$), suggesting that the roots were changing
phosphate levels in response to NaCl, a process known to help maintain ion homeostasis (*25*).
For mannitol, we observed a specific downregulation of 'cell wall organization or biogenesis' and
'microtubule-based processes' ($p < 7.8 \times 10^{-3}$) (**Supplementary File 4**).
187
188
189   **Examining differential gene expression responses to 'hard agar' (HA) treatment**
190
In addition to examining PEG, mannitol and NaCl transcriptional responses, we also tested a new
way of lowering water potential on an agar plate. We hypothesized that we could induce stress
by increasing both the agar and nutrient concentration. We called this media 'hard agar' (HA). By
testing three different doses (1.25×, 1.67× and 2.5× fold increase in both agar and LS
concentration, where 1.0× is 2 % agar and 1 X LS), we found that it limited plant shoot dry weight
and media water potential in a similar way to PEG, mannitol and NaCl treatment (**Figure 1A - C**).
Additionally, we found that HA treatment limited *Arabidopsis* primary root growth rate, shoot water
potential, and photosynthesis efficiency (**Figure 2-figure supplement 4**). At the molecular level,
RNA-seq revealed 1,376 and 1,921 genes that were dose-responsive to the level of HA stress in
roots and shoots respectively (**Figure 1E & Figure1-figure supplement 2**). We found that these
gene expression responses overlapped significantly with those found differentially expressed in

response to vermiculite drying (Fisher test, $p < 1 \times 10^{-32}$, 87 % directional agreement) (**Figure 2A**
& 2B). HA's impact can be seen in the gene expression responses of *HB12* (**Figure 2C**), *GCL1*
and *RD21* (**Figure2-figure supplement 3**).

An increase in nutrient concentration can induce salt-like stress while increasing agar
concentration will increase tensile stress (*26*). We tested each of these variables separately to
understand the role each played in eliciting the gene expression responses found in the HA assay.
To do this, we repeated our HA dose experiment, but now increasing only the concentration of
LS nutrients (1, 1.25, 1.67 and 2.5 X) or the concentration of agar (2, 2.5, 3.3 and 5 %) (**Figure**
2E). We found a significant decrease in shoot area size in response to an increase in nutrient
concentration (Pearson $p = 1.7 \times 10^{-7}$) and agar concentration ($p = 3.9 \times 10^{-13}$), where the latter
more closely phenocopied the effect of HA (**Figure 2E**, **Figure 2-figure supplement 5,**
Supplementary File 5). Since the increase in nutrient concentration alone was responsible for
changing media water potential, the phenotypic response to increased agar concentration was
not in response to a lower water potential (**Figure 2-figure supplement 5**). Next, we examined
the transcriptional responses underlying nutrient and agar responses by sequencing root tissue
across each dose tested. Through linear modeling, we found 1,043 genes and 938 genes that
were dose-responsive to nutrient or agar concentration, respectively. Then, we investigated how
these genes compared to those found in vermiculite drying responses. We found that genes
differentially expressed in response to an increase in agar or nutrient concentration overlapped
12 % and 17 % of vermiculite drying responsive gene expression respectively (permutation test,
$p < 0.05$) (**Figure 2F, Figure 2-figure supplement 5, Supplementary File 6**). However, we found
genes differentially expressed in response to HA treatment led to a higher overlap (26 %),
suggesting that both nutrient and agar concentration contribute to the similarity between HA
treatment and vermiculite drying.

Finally, we tested if our HA assay was sensitive enough to detect phenotypic variability. To
achieve this, we grew 20 different *Arabidopsis* ecotypes on 2× HA agar (4% agar, 2 X LS), where
ecotypes were selected from a previous drought study that assessed fitness in a common garden
experiment (*27*). By comparing the total shoot area after three weeks of growth, we found that our
assay revealed variability in shoot growth responses (**Figure 2G** & **2H**, **Supplementary File 7**).
Furthermore, we found that the greater the impact HA agar had on reducing an accession's
relative shoot size, the better the accession's fitness was, as measured under field conditions (*27*)
(Spearman $p = 0.04$, **Figure 2-figure supplement 6**). This is likely because smaller shoot

systems have a better chance of survival and reproduction (*28*). This suggests that our assay
may be useful for screening for novel drought-associated loci among a wider group of accessions
or mutants (*29, 30*).
239
In summary, our investigation has assessed the shared and unique impacts of agar-based low
water potential treatments on gene expression. We also compared these effects with the
expression patterns elicited during vermiculite drying. We found each plate-based assay
generated similar responses in shoot tissue, but more varied responses in root tissue. We note
that our comparative analysis focuses largely on transcriptomic responses in *Arabidopsis.* We
suggest investigating gene expression responses in other species as future work. Here, we also
introduce another method for lowering water potential. By increasing nutrient and agar
concentration, our HA approach also induced gene expression responses comparable to
vermiculite drying. We describe how to make this media within the **Materials and Methods**.
249
250
251 Supplementary
252
Supplementary File 1 – Plant physiological measurements
Supplementary File 2 – Differentially expressed genes and normalized counts in HA, PEG,
Mannitol, NaCl or vermiculite drying experiments
Supplementary File 3 – Vermiculite drying assay measurements
Supplementary File 4 – GO Term enrichment of differentially expressed genes
Supplementary File 5 – Shoot area of seedlings grown under different agar and nutrient
concentrations
Supplementary File 6 – Differentially expressed genes and normalized counts in response to
changes in nutrient or agar concentration
Supplementary File 7 – Shoot area of different *Arabidopsis* accessions grown on HA media
263
264 **Acknowledgements**
265

266 We thank Renee Garza for critical reading of the manuscript. J.S is an Open Philanthropy
267 awardee of Life Science Research Foundation, as well as recipient of the Pratt Industries
268 American-Australian Association Scholarship. J.R.E is an Investigator of the Howard Hughes
269 Medical Institute.

Data Availability

Raw sequencing data is available at the National Center for Biotechnology Information Sequence

Read Archive (accession number PRJNA904764). Normalized read counts and raw phenotypic

datasets can be found in the Supplementary Material.

Materials and Methods

Hard Agar Assay – *Arabidopsis* seedlings were grown on vertical plates for 8 days under short

day conditions (8 h light, 21 C, 150 umoles light) on agar media (1 × Linsmaier & Skoog (LS)

(Cassion LSP03) media, 1% sucrose, 2% agar, pH 5.7). We note LS media is identical to

Murashige & Skoog (MS) media in inorganic salt content, but lacks Glycine, Nicotinic Acid and

Pyridoxine HCl. After 8 days, plants were transferred to 'hard agar' (HA) plates. The 1.0× plate

consisted of 2 % and 1 X LS media, with no sucrose (pH 5.7) at a final volume of 75 mL.

Subsequent doses of increased nutrient and agar concentration (1.25×, 1.67× and 2.5× fold

increase) were made by preparing the same media but reducing the amount of water present. For

example, the 1.25×, treatment plate contained 60 mL of 2.5% agar and 1.25× LS media. We note

that the volume of HA itself has minimal impact gene expression responses (**Figure 2-figure**

supplement 7). On day 14, 2 hours after subjective dawn, shoot and root samples were flash

frozen (6 plants per replicate). In total, we collected 16 samples for RNA-seq analysis (2 organs,

2-3 biological replicates, 3 treatment levels). We also collected a non-treated control set (2

biological replicates).

To test different *Arabidopsis* accessions on HA agar, plants were sown on either 1×, or 2×

treatments as described above, however supplemented with 0.5% or 1% sucrose respectively to

encourage germination. Seedlings were grown for 3 weeks under short day conditions in before

imaging plates in duplicate (n = 2 - 5 plants per plate) (**Supplementary File 7**). Shoot area was

calculated from images using Plant Growth Tracker (GitHub -

https://github.com/jiayinghsu/plant_growth_tracker).

<u>Vermiculite Drying Assay</u> – *Arabidopsis* seedlings were grown on vertical plates for 17 days under

short day conditions (8 h light, 21 C, 150 umoles light) on agar media (1×LS, 1% sucrose, 2%

agar, pH 5.7), before transfer to vermiculite saturated with $0.75 \times$ LS media. We note at the timing

of transfer lateral root formation had begun. Plants were then grown on vermiculite at 100% field

capacity (FC) for 12 days (8 h light, 21 C, 150 umoles light). On the 13[th] day, the first time point

was sampled (4.5 hours after subjective dawn) where tissue was flash frozen in liquid nitrogen.

After this, excess aqueous solution was drained from each pot, and then each pot was calibrated

to $1 \times$ FC. Plant tissue was harvested each day on subsequent days at the same time of day.

Each day, pots were weighed to measure extent of evaporation. By these means, FC was

measured (**Figure 1-figure supplement 4**). After the 5[th] day sample was taken, water was re-

added to the remaining pots to an excess of $1\times$ FC. ~ 15 plants were sampled per time point. In

total, we harvested 78 tissue samples for RNA-seq (3 biological replicates, 2 organ types, 7 days,

2 treatments). Plants were then left to grow under long day conditions until flowering. Seeds were

harvested, dried, and weighed (n = 50 plants per treatment).

<u>Polyethylene Glycol (PEG) Stress Assay</u> – *Arabidopsis* seedlings were grown on vertical plates

for 8 days under short-day conditions (8 h light, 21 C, 150 umoles light) on agar media (1×LS, 1%

sucrose, 2% agar, pH 5.7), before transfer to polyethylene glycol (PEG) media of varying

concentrations. PEG media plates were prepared by dissolving crystalline 6000 MW PEG into

freshly autoclaved $1\times$ LS media pH 5.7 and pouring 50 mL of PEG media solution onto $1\times$ LS, 2%

agar, media plates (pH 5.7), letting the PEG solution diffuse into the solid media overnight, then

pouring off excess and transferring seedlings to PEG infused media plates as described in (*11*).

Plants were grown under 3 different treatments (12%, 20%, and 28% PEG solution w/v) for 14

days. On day 14, 2 hours after subjective dawn, shoot and root samples were flash frozen (6

plants per replicate). In total, we collected 16 samples for RNA-seq analysis (2 organs, 2-3

biological replicates, 3 treatment levels).

<u>Mannitol and NaCl Osmotic Stress Assays</u> – *Arabidopsis* seedlings were grown on vertical plates

for 8 days under short-day conditions (8 h light, 21 C, 150 umoles light) on agar media ($1\times$ LS,

1% sucrose, 2% agar, pH 5.7), before transfer to either mannitol or salt (NaCl) media of varying

concentrations. Mannitol and NaCl media plates were prepared by adding respective volume of

stock solution to $1\times$ LS, 2% agar, pH 5.7 media before autoclaving. Plants were grown under 3

different treatments of mannitol or NaCl (50 mM, 100 mM and 200 mM for mannitol, 30 mM, 75

mM, and 150 mM for NaCl) for 14 days. On day 14, 2 hours after subjective dawn, shoot and root

samples were flash frozen (6 plants per replicate). In total, for either Mannitol or NaCl treatment

experiments, we collected 18 samples for RNA-seq analysis (2 organs, 3 biological replicates, 3

treatment levels).

ABA Exogenous Treatment Assay – *Arabidopsis* seedlings were grown on vertical plates for 8

days under short-day conditions (8 h light, 21 C, 150 umoles light) on agar media (1× LS, 1%

sucrose, 2% agar, pH 5.7), before transfer to 1× LS, 2% agar, pH 5.7 control media and grown

for 14 days. On day 14, abscisic acid (ABA) solutions of 1 uM, 5 uM and 10 uM were prepared

from 10 mM ABA dissolved in ethanol stock, as well as a mock treatment solution containing 0.1%

ethanol concentration. 30 min after subjective dawn, 15 mL of each solution was dispersed onto

the roots of the seedlings. After 1 min of treatment, the ABA solution was removed from the plates,

and the plates returned to the growth chamber. 2 hours after subjective dawn, shoot and root

samples were flash frozen (6 plants per replicate). In total, we collected 8 samples for RNA-seq

analysis (root tissue only, 2 biological replicates, 4 conditions).

Osmotic Potential Measurements: The water potential of media was determined considering it

equivalent to the osmotic potential (Ψs). Osmotic potential was measured using a vapor pressure

osmometer (Model 5600, ELITech Group; Puteaux, France). Readings were taken from melted

agar media constituted with the particular stress type. Osmolality readings for each sample

obtained were converted to megapascals (MPa) using the equation Ψs = -CRT, where C is the

molar concentration, R is the universal gas constant, T is the temperature in Kelvin. We note that

to measure the water potential of PEG treatment media, we infiltrated the PEG solution into plates

as described above, and then melted the PEG-infiltrated agar for measurement with the

osmometer. We assessed the osmotic potential of shoot tissue two weeks after transplanting the

seedlings to HA agar media, After immersion in liquid nitrogen 3 shoots were placed into 0.5-ml

tubes and centrifuged to extract the tissue sap. The osmotic potential (Ψs) of the extracted sap

was determined using a vapor pressure osmometer.

Chlorophyll fluorescence measurements: Chlorophyll fluorescence was assessed in eight

seedlings of each plate using the Walz PAM IMAGING PAM M-series IMAG-K7 (MAXI)

fluorometer. For every experiment, leaves were pre-conditioned in the dark for 1 h. The maximum

quantum yield of PSII (F_v/F_m) was calculated using the formula:

368                                  $$F_v/F_m = (F_m - F_0)/F_m$$

Where F_v is the variable fluorescence, F_m is the maximal fluorescence following 1 h of dark
adaptation and F_0 is the minimal fluorescence level of a dark-adapted leaf when all photosystem
II (PSII) reaction centers are open.

Root Growth Rate Measurements – *Arabidopsis* seedlings were grown on vertical plates for 8
days under short day conditions (8 h light, 21 C, 150 umoles light) on agar media (1× LS, 1%
sucrose, 2% agar), before transfer to 2.5× HA, 28% PEG, 150 mM mannitol or 150 mM NaCl
treatment plates as described above. Root images were acquired every two days for a total of 8
days using scanners. Primary root length, defined as the length (scaled to
cm) from hypocotyl base to root tip, was quantified using image J. For each treatment we
screened 4 plates, with each plate holding 4 individual plants.

RNA-extraction and Library Preparation – Plant tissue was crushed using the TissueLyser
(Agilent) and RNA extracted using RNeasy Mini Kit (Qiagen). Number of biological replicates per
library ranged between RNA quality was assessed using Tape station High Sensitivity RNA assay
(Agilent). 0.5 - 1 ug of total RNA proceeded to library preparation, where libraries were prepared
using TruSeq stranded mRNA kit (Illumina). Resulting libraries were sequenced on the NovaSeq
6000 (Illumina) with 2x150 bp paired-end read chemistry. Read sequences were aligned to the
Arabidopsis TAIR10 genome using HISAT2 (*31*), and gene counts called using HT-seq (*32*), by
relying on Araport11 annotation (*33*). Normalized counts can be found in **Supplementary File 2**.
For each organ, libraries from all experiments were normalized together before calling differential
expression.

Statistical Analysis – To detect differential expression in our drought assay on vermiculite, we
called differential expression using a linear model using the DESeq2 LRT function to associate a
change in field capacity with change in gene expression. The same statistical approach was used
to associate a change in a gene's expression to changes in dose of HA, PEG, mannitol, and NaCl,
as well as changes in agar concentration, nutrient concentration and volume of agar used.
Resulting model p-values were adjusted to account for false discovery (*p*-value < 0.05). The
complete list of differentially expressed genes for each experiment can be found in
Supplementary File 2 and **Supplementary File 6**. Pairwise differential gene expression was
called using DESeq2 (*34*). Specifically, for plate based assays, we called differential expression
by comparing the control treatment to each treatment dose, using an adjusted *p*-value threshold
of 0.05. Overlap analyses were performed using Fisher exact tests, with an adjusted *p*-value

threshold of 0.05. The background for these intersects was all expressed genes within the

respective organ. Permutation tests and GO Term enrichment analyses was performed in

VirtualPlant (*35*), with all expressed genes within the respective organ used as background.

419
420

Figure 1 - **Benchmarking the impact different stress assays have on *Arabidopsis* gene expression. A:** 22-day-old *Arabidopsis* growth on plates under either 1.67× hard agar (HA), 20 % PEG, 100 mM mannitol, or 75 mM NaCl treatments. **B**: Water potential measurements of treatment media (n = 3 - 4). **C:** Dry weight of 22-day-old *Arabidopsis* seedlings under different doses of each stress treatment (n = 11 - 12). **D:** Number and intersect of differentially expressed genes (DEGs) that are dose-responsive to each stress treatment within root and shoot tissue. **E - I:** Heatmaps displaying the top 50 most significant upregulated or downregulated genes in response to **(E)** HA, **(F)** PEG, **(G)** mannitol, **(H)** NaCl and **(I)** vermiculite drying in the *Arabidopsis* root (n = 2 - 3 biological replicates). Membership of Gene Ontology (GO) Terms for 'response to stress', 'response to chemical stimulus' or 'response to ABA stimulus' are indicated.

Figure 2 – Comparing HA, PEG, mannitol, and NaCl gene expression responses to vermiculite drying. A: Heatmap displaying genes differentially expressed in response to vermiculite drying in shoot or root tissue compared to their dose-responsive expression within each plate-based assay. Level of 'directional agreement' (i.e., differentially expressed in the same direction) found within each assay reported. **B:** Overlap analysis of genes found differentially expressed due to vermiculite drying, compared to those found differentially expressed within each dose of PEG, mannitol, NaCl or HA agar assays in either shoot and root. **C - D:** Expression patterns of *HOMEOBOX12* (*HB12*) and *DROUGHT HYPERSENSITIVE 2* (*DRY2*) across each assay in root tissue. **E:** Shoot area of seedlings grown under increasing doses of HA, agar or nutrient concentrations (n = 19). **F:** Number and percent overlap of genes found differentially expressed in response to increasing doses of HA, agar or nutrient concentrations with those differentially expressed in response to vermiculite drying. **G:** Total shoot area of *Arabidopsis* accessions grown under either 1× or 2× HA agar treatment (n = 5 - 12). **H:** Images of *Arabidopsis Trs-0* or *UKSE06-325* accessions grown on either 1× or 2× HA treatment.

A
| HA 1.25× | PEG 12 % | Mannitol 50 mM | NaCl 30 mM |

B
| no treatment | HA 1.67× | PEG 20 % | Mannitol 100 mM | NaCl 75 mM |

C
| HA 2.5× | PEG 28 % | Mannitol 200 mM | NaCl 150 mM |

Figure 1 – figure supplement 1

Plant growth responses to stress assays. A – C: Images of 22-day-old *Arabidopsis* seedlings grown under different doses of each agar stress assay.

490
491
Figure 1 - figure supplement 2
492
493
494 **Shoot gene expression responses to each stress assay are dose responsive.** Heatmap
495 displaying the top 50 most significant upregulated or downregulated genes in shoots in response
496 to **(A)** HA, **(B)** PEG, **(C)** mannitol, **(D)** NaCl and **(E)** vermiculite drying (n = 2 - 3 biological
497 replicates). Membership of Gene Ontology (GO) Terms for 'response to stress', 'response to
498 chemical stimulus' or 'response to ABA stimulus' indicated.

Figure 1 - figure supplement 3

Overlapping differentially expressed genes (DEGs) responsive to different assay types.
Overlap of dose-responsive differentially expressed genes in shoot **(A)** and root **(B)** in response
to either HA, PEG, mannitol or NaCl (replicated from Figure 1). Number of upregulated or
downregulated dose-responsive genes in response to each treatment type in shoot **(C)** and root
(D). Overlapping gene sets in **(E)** shoot or **(F)** root tissue (permutation test, p < 0.001).

522
523
524 **Figure 1 - figure supplement 4**
525
526 **Treating vermiculite-grown *Arabidopsis* plants to mild drought stress. A:** Field capacity
527 measurements of vermiculite as water evaporated over a 5-day period (n = 6 - 12). **B:** Shoot dry
528 weight of *Arabidopsis* rosettes as they grew either under well-watered conditions or drought
529 conditions over a 5-day period (t-test, n = 12). **C:** Images of plants after 5 days of water stress.
530 **D:** Seed yield resulting from *Arabidopsis* plants after drought recovery (t-test, n = 50).

539
540
541
Figure 2-figure supplement 1

Comparing gene expression response to vermiculite drying and PEG treatment with
previous studies. A: Intersect analysis of root or shoot genes found differentially expressed in
response to vermiculite drying within this study, and genes found differentially expressed in
response to soil drying by Lozano-Elena *et al.* (*16*) or by Wilkins *et al. (36)* (permutation test, p <
0.001). **B - C:** Intersect analysis of genes found differentially expressed in response to PEG
treatment in shoot **(B)** or root **(C)** in this study, with those found differentially expressed in
response to PEG treatment by Wong *et al.* (*37*) and Wang *et al.* (*38*) (permutation test, p < 0.001).
D - E: Heatmap displaying direction of shoot **(D)** or root **(E)** differentially expressed in response
to vermiculite drying (this study) and Wilkins *et al. (36)* or Lozano-Elena *et al.* (*16*) respectively.
Directional agreement with this study's vermiculite drying response indicated. **F - G :** Heatmap
displaying direction of genes differentially expressed in response to PEG treatment across each
study. We note that both Wong *et al.* and Wang *et al.* assess transcriptomic responses of whole
seedlings (both root and shoot), and thus we compare our shoot **(F)** and root **(G)** data separately.
E: Examining the 119 genes that were differentially expressed in response to drought (this study),
PEG treatment (this study) and PEG treatment reported in Wong *et al.*

Figure 2-figure supplement 2

Comparing ABA-induced differential expression to vermiculite drying and HA induced gene expression patterns. A: Overlap analysis of genes found differentially expressed in response to vermiculite drying, compared to those within each dose of either transient ABA treatment, PEG, mannitol, NaCl, or HA agar assays in both root and shoot (Fisher exact test, adj. $p < 0.05$). **B:** Heatmap displaying genes differentially expressed under vermiculite drying in root tissue compared to their dose-responsive expression within each stress assay. Direction of gene expression agreement with vermiculite-drying responsive gene expression (i.e., 'directional agreement') indicated.

Figure 2-figure supplement 3

Gene expression profiles of individual genes. A - E: Expression patterns of individual genes under doses of each assay in shoot tissue: **(A)** RD29B, **(B)** RD20, **(C)** HB7, **(D)** NCED3 and **(E)** P5CS1. **F - H:** Expression patterns of individual genes under doses of each assay in root tissue: **(F)** GCL1, **(G)** RD21 and **(H)** RHS18.

602
603
604 **Figure 2-figure supplement 5**
605
Comparing the separate effects of nutrient concentration and agar concentration on
seedling growth. Image of *Arabidopsis* seedlings grown on either **(A)** 1× HA agar (i.e. 1X LS,
2% agar) or **(B)** 2.5× HA, which increased both nutrient and agar concentrations to 2.5X and 5%,
respectively. **C:** Image of seedlings grown on an increased 2.5X nutrient concentration (without a
change in agar concentration). **D:** Image of seedlings grown on an increased 5% agar
concentration (without a change in nutrient concentration). **E:** Water potential measurements of
media presented in **A – D** (n = 3). **F:** Intersection of differentially expressed genes responsive to
either agar concentration, nutrient concentration, HA treatment or drought stress (permutation
test, $p < 0.001$).

631 **Figure 2-figure supplement 6**

633 **Associating hard agar's (HA) impact on shoot size with plant fitness.** Comparing the impact
634 HA treatment has on shoot size of 20 different *Arabidopsis* accessions to the change in their
635 fitness found under drought conditions in the field, as reported in (27).

652
653   **Figure 2-figure supplement 7**
654
The volume of HA agar has minimal impact on gene expression. Image of *Arabidopsis*
seedlings grown on either **(A)** 75 mL of 1× HA agar, **(B)** 30 mL of 2.5× HA agar, or **(C)** 75 mL of
2.5× HA agar. Number of genes found differentially expressed is reported for comparisons **(A)**
and **(B),** as well as **(B)** and **(C)** (DESeq, adj. *p* < 0.01). Of the 29 genes found differentially
expressed between **(B)** and **(C)**, 13 are found in comparison **(A)** and **(B)**. **(E)** Heatmap of genes
found differentially expressed in either comparison.

References

1. H. C. J. Godfray *et al.*, Food Security: The Challenge of Feeding 9 Billion People. *Science* **327**, 812-818 (2010).
2. D. S. Battisti, R. L. Naylor, Historical warnings of future food insecurity with unprecedented seasonal heat. *Science* **323**, 240-244 (2009).
3. M. Dubois, D. Inze, Plant growth under suboptimal water conditions: early responses and methods to study them. *J Exp Bot* **71**, 1706-1722 (2020).
4. H. Claeys, S. Van Landeghem, M. Dubois, K. Maleux, D. Inze, What Is Stress? Dose-Response Effects in Commonly Used in Vitro Stress Assays. *Plant Physiol* **165**, 519-527 (2014).
5. M. Hohl, P. Schopfer, Water Relations of Growing Maize Coleoptiles : Comparison between Mannitol and Polyethylene Glycol 6000 as External Osmotica for Adjusting Turgor Pressure. *Plant Physiol* **95**, 716-722 (1991).
6. R. Munns, Comparative physiology of salt and water stress. *Plant Cell Environ* **25**, 239-250 (2002).
7. K. C. Jisha, J. T. Puthur, Halopriming of seeds imparts tolerance to NaCl and PEG induced stress in Vigna radiata (L.) Wilczek varieties. *Physiol Mol Biol Plants* **20**, 303-312 (2014).
8. C. Trontin *et al.*, A pair of receptor-like kinases is responsible for natural variation in shoot growth response to mannitol treatment in Arabidopsis thaliana. *Plant J* **78**, 121-133 (2014).
9. P. E. Verslues, M. Agarwal, S. Katiyar-Agarwal, J. Zhu, J. K. Zhu, Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* **45**, 523-539 (2006).
10. T. E. Juenger, P. E. Verslues, Time for a drought experiment: Do you know your plants' water status? *Plant Cell* **35**, 10-23 (2023).
11. C. M. van der Weele, W. G. Spollen, R. E. Sharp, T. I. Baskin, Growth of Arabidopsis thaliana seedlings under water deficit studied by control of water potential in nutrient-agar media. *J Exp Bot* **51**, 1555-1562 (2000).
12. G. Zeller *et al.*, Stress-induced changes in the Arabidopsis thaliana transcriptome analyzed using whole-genome tiling arrays. *Plant J* **58**, 1068-1082 (2009).
13. J. A. Kreps *et al.*, Transcriptome changes for Arabidopsis in response to salt, osmotic, and cold stress. *Plant Physiol* **130**, 2129-2141 (2002).
14. J. Swift, J. M. Alvarez, V. Araus, R. A. Gutierrez, G. M. Coruzzi, Nutrient dose-responsive transcriptome changes driven by Michaelis-Menten kinetics underlie plant growth rates. *Proc Natl Acad Sci U S A* **117**, 12531-12540 (2020).
15. N. Nikonorova *et al.*, Early mannitol-triggered changes in the Arabidopsis leaf (phospho)proteome reveal growth regulators. *J Exp Bot* **69**, 4591-4607 (2018).
16. F. Lozano-Elena, N. Fabregas, V. Coleto-Alcudia, A. I. Cano-Delgado, Analysis of metabolic dynamics during drought stress in Arabidopsis plants. *Sci Data* **9**, 90 (2022).
17. J. Msanne, J. Lin, J. M. Stone, T. Awada, Characterization of abiotic stress-responsive Arabidopsis thaliana RD29A and RD29B genes and evaluation of transgenes. *Planta* **234**, 97-107 (2011).
18. S. Takahashi, T. Katagiri, K. Yamaguchi-Shinozaki, K. Shinozaki, An Arabidopsis gene encoding a Ca2+-binding protein is induced by abscisic acid during dehydration. *Plant Cell Physiol* **41**, 898-903 (2000).
19. A. E. Valdes, E. Overnas, H. Johansson, A. Rada-Iglesias, P. Engstrom, The homeodomain-leucine zipper (HD-Zip) class I transcription factors ATHB7 and ATHB12

modulate abscisic acid signalling by regulating protein phosphatase 2C and abscisic
acid receptor gene activities. *Plant Mol Biol* **80**, 405-418 (2012).

20.   B. C. Tan *et al.*, Molecular characterization of the Arabidopsis 9-cis epoxycarotenoid
dioxygenase gene family. *Plant J* **35**, 44-56 (2003).

21.   Y. Gao *et al.*, Genetic characterization reveals no role for the reported ABA receptor,
GCR2, in ABA control of seed germination and early seedling development in
Arabidopsis. *Plant J* **52**, 1001-1013 (2007).

22.   M. Koizumi, K. Yamaguchi-Shinozaki, H. Tsuji, K. Shinozaki, Structure and expression of
two genes that encode distinct drought-inducible cysteine proteinases in Arabidopsis
thaliana. *Gene* **129**, 175-182 (1993).

23.   D. Pose *et al.*, Identification of the Arabidopsis dry2/sqe1-5 mutant reveals a central role
for sterols in drought tolerance and regulation of reactive oxygen species. *Plant J* **59**, 63-
76 (2009).

24.   O. P. Ponce *et al.*, Transcriptome profiling shows a rapid variety-specific response in two
Andigenum potato varieties under drought stress. *Front Plant Sci* **13**, 1003907 (2022).

25.   K. Miura, A. Sato, M. Ohta, J. Furukawa, Increased tolerance to salt stress in the
phosphate-accumulating Arabidopsis mutants siz1 and pho2. *Planta* **234**, 1191-1199
(2011).

26.   S. Verger, Y. Long, A. Boudaoud, O. Hamant, A tension-adhesion feedback loop in plant
epidermis. *Elife* **7**, (2018).

27.   M. Exposito-Alonso *et al.*, Natural selection on the Arabidopsis thaliana genome in
present and future climates. *Nature* **573**, 126-129 (2019).

28.   A. Skirycz *et al.*, Survival and growth of Arabidopsis plants given limited water are not
equal. *Nat Biotechnol* **29**, 212-214 (2011).

29.   J. M. Alonso *et al.*, Genome-wide insertional mutagenesis of Arabidopsis thaliana.
Science **301**, 653-657 (2003).

30.   Genomes Consortium,1,135 Genomes Reveal the Global Pattern of Polymorphism in
Arabidopsis thaliana. *Cell* **166**, 481-491 (2016).

31.   D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-
915 (2019).

32.   S. Anders, P. T. Pyl, W. Huber, HTSeq-a Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**, 166-169 (2015).

33.   C. Y. Cheng *et al.*, Araport11: a complete reannotation of the Arabidopsis thaliana
reference genome. *Plant J* **89**, 789-804 (2017).

34.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**, (2014).

35.   M. S. Katari *et al.*, VirtualPlant: a software platform to support systems biology research.
Plant Physiol **152**, 500-515 (2010).

36.   O. Wilkins, K. Brautigam, M. M. Campbell, Time of day shapes Arabidopsis drought
transcriptomes. *Plant J* **63**, 715-727 (2010).

37.   M. M. Wong *et al.*, Phosphoproteomics of Arabidopsis Highly ABA-Induced1 identifies
AT-Hook-Like10 phosphorylation required for stress growth regulation. *Proc Natl Acad
Sci U S A* **116**, 2354-2363 (2019).

38.   Y. Wang, Z. Fang, L. Yang, Z. Chan, Transcriptional variation analysis of Arabidopsis
ecotypes in response to drought and salt stresses dissects commonly regulated
networks. *Physiol Plant* **172**, 77-90 (2021).

765