Neural alpha oscillations and pupil size differentially index cognitive demand under competing audio-visual task conditions

Frauke Kraus [1,2], Sarah Tune [1,2], Jonas Obleser [1,2]*, & Björn Herrmann [3,4]*

[1] Department of Psychology, University of Lübeck, Lübeck, Germany

[2] Centre for Brain, Behavior, and Metabolism, University of Lübeck, Lübeck, Germany

[3] Rotman Research Institute, Baycrest Health Sciences, Toronto, ON, Canada

[4] University of Toronto, Toronto, ON, Canada

joint senior authorship

Correspondence concerning this article should be addressed to Björn Herrmann, Rotman Research Institute, Baycrest, 3560 Bathurst St, North York, ON, M6A 2E1, Canada. bherrmann@research.baycrest.org or Frauke Kraus, Maria-Goeppert-Str. 9a, 23562 Lübeck, Germany. f.kraus@uni-luebeck.de

Keywords: Cognitive demand, pupil dilation, alpha oscillations, audio-visual stimulation, multiple object tracking task, gap detection, dual task

Number of pages (29), Number of figures (6)

Word counts: Abstract (213/250), Introduction (717/650), Discussion (1564/1500)

Conflict of interest statement:

The authors declare no competing financial interests.

Acknowledgments:

We thank Larissa Scheller and Hannah Schewe for assisting with the data collection. This work was supported by Deutsche Forschungsgesellschaft (DFG; grant number HE 7857/1-1) awarded to BH.

F.K., J.O. and B.H. designed research, F.K. performed research, F.K., J.O., S.T. and B.H. analyzed data, F.K., J.O., S.T. and B.H. wrote the paper.

31   Abstract

Cognitive demand is thought to modulate two often used, but rarely combined, measures: pupil size and

neural alpha (8–12 Hz) oscillatory power. However, it is unclear whether these two measures capture

cognitive demand in a similar way, and whether a measured 'cognitive demand' response in pupil size

or brain activity is dependent on sensory modality by which the demand is induced. Here we recorded

pupil size and neural alpha power (using electroencephalography), while human participants of both

sexes concurrently performed a visual multiple object-tracking task and an auditory gap-detection task.

Difficulties of the two tasks were manipulated independent of each other. Participants' performance

decreased in accuracy and speed with increasing cognitive demand. Pupil size increased with increasing

demand for both the auditory and the visual task. Alpha power emerging from two cortical sources

showed counteracting neural dynamics: Alpha power in visual cortex decreased with increasing demand

in the visual task, whereas alpha power in parietal cortex increased with increasing demand in the

auditory task. Furthermore, demand-related changes in pupil size and alpha power were uncorrelated,

suggesting that they index different aspects of cognitive demand. Overall, the current study

demonstrates that the dynamics of the neurophysiological indices of cognitive demand and associated

effort are multi-faceted and potentially modality-dependent under complex audio-visual task conditions.

47   Significance Statement

Pupil size and oscillatory alpha power are associated with cognitive demand and effort, but their

sensitivity under complex audio-visual task conditions is unclear and so is the extent to which they share

underlying mechanisms. Using an audio-visual dual-task paradigm, we show that pupil size increases

with increasing cognitive demands for both audition and vision. In contrast, oscillatory alpha power

changes depend on sensory modality: Power in auditory cortex increased with auditory demand,

whereas alpha power in visual cortex decreased with visual demand. Hence, pupil size and alpha power

show different sensitivity to cognitive demands, which is also indicated by the absence of a correlation

between both measures, suggesting different underlying neural mechanisms.

56   Introduction

Many situations in everyday life require the integration of information from different sensory modalities.

However, cognitive resources are limited (Kahneman, 1973; Lavie, 1995), and, depending on the induced

cognitive demand, such situations may be experienced as effortful (Björn Herrmann & Johnsrude, 2020;

Pichora-Fuller et al., 2016). This cognitive demand and related effort have practical relevance, because

people who experience effort may avoid challenging situations and be at risk of social isolation (Mick et

al., 2014; Shukla et al., 2020; Sung et al., 2016). A better understanding of how individuals manage

cognitive resources under complex multi-sensory conditions, and of the underlying psychophysiology

are critical to objectively measuring and identifying effort in people who struggle.

65   Dual-task paradigms are often used in behavioral research to investigate how perceptual or

cognitive demands in one modality affect performance in another modality (Desjardins & Doherty, 2013;

Gagné et al., 2017; Picou & Ricketts, 2014; Wu et al., 2016). A hallmark result of dual-task paradigms

indicating cognitive constraint lies in the deterioration of behavioral performance for concurrent tasks

that would be performed with high accuracy and speed if carried out separately (Desjardins & Doherty,

2013; Gosselin & Gagné, 2014; Picou & Ricketts, 2014; Wu et al., 2016). In addition, manipulating the

difficulty of the individual tasks allows tapping into the dynamic allocation of cognitive resources to

either task as a function of the respective task difficulty combinations. The current study aims to

understand how physiological responses change with varying degrees of task difficulty and thus cognitive

demand in single- and dual-task conditions combining the auditory and visual domain.

75   At least two neurophysiological measures have recently and repeatedly become associated with

changes in cognitive demand: pupil size (Joshi & Gold, 2020; Koelewijn et al., 2012; Ohlenforst et al.,

2018; Wendt et al., 2016) and neural oscillatory activity in the alpha-frequency band (8–12 Hz; Obleser

et al., 2012; Paul et al., 2021; Wisniewski et al., 2017). Pupil size variations are mainly driven via

noradrenergic pathways from Locus Coeruleus (Aston-Jones & Cohen, 2005; Joshi et al., 2016; Joshi &

Gold, 2020) that is modulated by attention (Vazey et al., 2018) or arousal (Breton-Provencher & Sur,

2019; Murphy et al., 2011). Pupil size is long known to vary with the degree to which a person engages

cognitively in a task (Kahneman & Beatty, 1966) and has recently gained substantial interest especially

in the hearing sciences, as a potential indicator of listening effort: pupil size increases with increasing

speech-comprehension difficulty induced by acoustic degradation (Miles et al., 2017; Winn et al., 2016),

acoustic masking (Koelewijn et al., 2012; Zekveld et al., 2010), or linguistic complexity (Kadem et al.,

2020; Wendt et al., 2016). Apart from the sensitivity of pupil size to auditory-related demands, pupil size

further increases with the degree of cognitive demand during visual stimulation (Martin et al., 2020;

Porter et al., 2007; Stolte et al., 2020) and memory performance (Kahneman & Beatty, 1966; Miller et

al., 2019). However, it is currently unknown how sensitive pupil size remains as an objective marker of

cognitive effort under more realistically complex audio-visual conditions of varying demands.

The second neurophysiological measure that may enable segregating different contributions

associated with cognitive demand is neural alpha power (8–12 Hz; Dimitrijevic et al., 2017; 2021;

Petersen et al., 2015; Wöstmann et al., 2017). Alpha power in parietal cortex increases when auditory-

induced cognitive demand increases (Henry et al., 2017; Björn Herrmann et al., 2022; Winneke et al.,

2020), such as with acoustic degradation of speech (Obleser et al., 2012; Wöstmann et al., 2015). In

contrast, increased cognitive demand in a visual task leads to a decrease in alpha power, often in visual

rather than parietal areas (Erickson et al., 2019; Magosso et al., 2019; Roijendijk et al., 2013). A recently

observed associated of the Locus Coeruleus activity with neural oscillatory activity (Dahl et al., 2020,

2022) raises the possibility that both pupil size and neural oscillatory activity are driven by a common

underlying neural process, which might be noradrenergically mediated.

The present study investigates this hypothesis. We investigate how pupil size and source-

localized alpha power in various sensory and executive brain areas covaries with varying levels of

cognitive demand in an audio-visual, dual-task setting. If both measures are proxies for a common

underlying neural process, they should co-vary with levels of cognitive demand.

Material and Methods

Participants

Twenty-four adults (age range: 19-30 years; mean = 23.7 years; SD = 3.09 years; 7 males, 17 females; all

right-handed) were recruited for the current study via the participant database of the Department of

Psychology at the University of Lübeck. They were native speakers of German and reported no history

of neural disorders nor hearing problems.

Each participant took part in two sessions. In the first session, participants separately performed

two single tasks. In the second session, participants performed the same tasks in a dual-task procedure.

Task procedures are described in detail below. The two sessions were conducted on different days,

separated by at least one day (median: 7 days; range: 1-18 days).

Participants gave written informed consent prior to participation and were financially

compensated with €10/hour or received course credits. The study was conducted in accordance with

the Declaration of Helsinki and was approved by the local ethics committee of the University of Lübeck.

Experimental environment

Participants were seated in a comfortable chair in a sound-attenuated booth. Participants placed their

head on a chinrest positioned at about 70 cm distance from a computer monitor (ViewSonic TD2421,

refresh rate 60 Hz). The experimental stimulation was controlled by a desktop computer (Windows 7)

running Psychtoolbox-3 in MATLAB and an external RME Fireface UC sound card. Visual stimulation was

mirrored from the stimulation computer to the computer monitor in the sound booth. Sound was

delivered binaurally via in-ear headphones (EARTONE 3A, 3M). Responses were given via a four-button

response box (The Black Box Toolkit, Sheffield, UK).

Experimental Design

In all task conditions, participants were simultaneously presented with auditory and visual stimulation.

The auditory stimulation consisted of a 7-s white noise sound in which a single gap occurred at one of

70 randomly selected and linearly spaced time points at 4–6 sec post noise onset (see Figure 1A). The

task for participants was to press a button on a response box as soon as they detected the gap.

The auditory gap-detection task comprised two difficulty levels: In the hard condition, gap

duration was titrated for each individual participant to about 75% gap-detection performance in training

blocks prior to the main experimental blocks (4–6 training blocks, each about 2 min). In the easy

condition, the estimated gap duration was doubled. Auditory stimuli were presented at 50 dB sensation

level estimated using a methods of limits procedure (cf. Herrmann et al., 2018). All button responses

occurring between 0.1–1s post gap onset were counted as hits. Response times were calculated as the

time elapsed between gap onset and button press. Response speed was calculated as the inverse of

response times (1/RT). Response speed was averaged across trials, separately for each condition and

participant.

The visual stimulation consisted of a multiple object-tracking (MOT) display (Cavanagh & Alvarez,

2005; Björn Herrmann & Johnsrude, 2018; Pylyshyn & Storm, 1988; Scholl, 2009). The computer monitor

displayed a 14 × 14 cm white edged black rectangle on black background at a distance of 70 cm from

participants' eyes (approximately 13° visual angle). A small, yellow fixation square was presented at the

center of the rectangle. The critical stimuli were 16 dots presented within the borders of a rectangle

(Figure 1A). At the beginning of each trial (prior to sound onset), a stationary display of the 16 dots was

shown for 1 second. One or five of the 16 dots were colored red (target dots) whereas the rest of the

dots were white (distractor dots). We refer to the one-dot condition as the 'easy' visual condition,

whereas we refer to the five-dot condition as the 'hard' visual condition. After 1 second, the dots that

were marked in red turned to white, and all 16 dots started to move for 7 seconds, simultaneously with

the presentation of the white noise auditory stimulus. Participants were instructed to follow the target

dot(s) (previously marked in red) over the 7-second period. After 7 seconds, all dots stopped moving,

and one target dot and two distractor dots were colored green. The three colored dots were each also

overlaid by the numbers 1, 2 and 3. Participants had to decide which of the three dots was a target dot

by pressing the respective button on the response box with no explicit time limit. Response times for

the visual task were calculated as the time elapsed between the onset of the answer screen and the

occurrence of a button press. Again, response speed was calculated and averaged across trials for each

participant and condition.

In the single-task session, participants were presented with a total of 70 trials for each of the

four resulting task and difficulty combinations (Figure 1B, left panel). Trials were split across 8 blocks of

35 trials each. Task and difficulty levels were varied block-by-block resulting in two blocks per task and

difficulty condition. The order of auditory-task blocks and visual-task blocks alternated. Starting block

and difficulty condition were counterbalanced across participants. Difficulty levels of the to-be-ignored

modality varied orthogonally to the difficulty of the target modality across the stimulation blocks. Each

block started with written instructions, indicating the level of difficulty and target modality in the

upcoming block. Prior to the main experimental blocks, participants performed the tasks separately to

familiarize them with the tasks and procedures.

Similarly, in the dual-task session, participants were presented with 70 trials for each of the four

difficulty combinations (Figure 1B, right panel), distributed across eight blocks. Participants performed

two blocks per difficult combination which remained fixed throughout each block of 35 trials. Block order

ensured that all four difficulty combinations were presented once before the presentation of the second

block per difficulty combination. Condition order was counterbalanced across participants. Each block

started with instructions written on the screen, stating the difficulty combination of the trials in the

upcoming block.

Figure 1. Experimental design. A: Trial design. Auditory and visual stimuli were presented concurrently. Auditory gap-detection task: Participant had to detect a gap within 7 s of white noise (the gap could occur within the 4–6-s time window). For the 'hard' condition, the gap duration was individually titrated to 75% correct. For the 'easy' condition, the gap duration was doubled. Multiple object-tracking task: Participants viewed 16 moving dots and were asked to follow the initially cued (red) dots in a moving-dot scene. After 7 s, dots stopped moving and three dots were marked green and labelled 1, 2, and 3. Participants had to decide which of the three dots was among the cued dots. Participants had to follow one (easy) or five (hard) dots. Analyses focused on the 3–4-s time window (pre-gap window; and additionally on the 5–6-s window for pupil size due to its slow response). **B:** Design for the single-task session (left) and the dual-task session (right). In the single-task session, participants performed the auditory and visual tasks separately (but were always presented with the audio-visual stimulation). In the dual-task condition, participants performed both tasks simultaneously. **C:** Schematic of hypotheses. If a physiological measure indexes cognitive demand independent of modality, the effect of increased difficulty should be the same across modalities (left panel). Alternatively, the effect of increased difficulty may differ between sensory modalities (right panel).

Pupil data recording and preprocessing

Eye movements and pupil size of the right eye were continuously recorded using an Eyelink 1000 Plus

eye tracker (SR Research) at a sampling rate of 500 Hz.

Data were preprocessed and analyzed using MATLAB (MathWorks, Inc.). Time points at which

the pupil size was more than three standard deviations above or below the mean pupil size calculated

over the whole block were categorized as blinks and marked as invalid ('missing') data in the time

window spanning 100 ms prior to and 100 ms following an identified blink. Subsequently, missing data

in the pupil size time series were linearly interpolated. To control for the potential influence of eye

movement-related changes, the x- and y-coordinates were regressed out of the pupil data (multiple

linear regression; Fink et al., 2021) and the resultant residual pupil size time course was used for all

further analyses. Data were then low-pass filtered at 4 Hz (Butterworth, 4th order) and segmented into

trials ranging from −2 to 8 s relative to noise onset. Trials were excluded if more than 40% of data points

within a trial had to be interpolated. The full dataset of a participant was excluded from analysis if more

than 50% of trials were excluded in any of the conditions (N=1).

Pupil-size data were downsampled to 50 Hz. For each trial, the mean pupil size was calculated in

the time window -2 to -1.1 s time window prior to noise onset and subtracted from the data at each

time point (baseline correction). Pupil size was averaged across trials, separately for each condition. To

account for the sluggishness of changes in pupil size, we averaged data both within the main time

window of interest also used in EEG analysis (3–4 s) along with a later time window 5–6 s post noise

onset (Montefusco-Siegmund et al., 2022).

EEG recording and preprocessing

We recorded participants' electroencephalogram (EEG) from 64 electrodes (ActiChamp, Brain Products,

München) at a sampling rate of 1,000 Hz, referenced to electrode TP9 (280 Hz online low-pass filter).

EEG data were analyzed with the FieldTrip toolbox (version 2019-09-20, (Oostenveld et al., 2011) for

Matlab, MathWorks, Inc.). Data were re-referenced to the average across electrodes, high-pass filtered

at 0.7 Hz (Hann window, 2391 points), and low-pass filtered at 100 Hz (Hann window, 89 points). Data

were filtered with a 50-Hz elliptic band-stop filter to suppress line noise.

Independent components analysis (ICA) was calculated to remove artifacts due to blinks, lateral

eye movements, and muscle activity. To this end, data were divided into 1-s segments, and segments

with non-stereotypical artefacts were removed on the basis of visual inspection prior to ICA calculation.

Noisy channels were removed prior to ICA. Artifact components were identified through visual

inspection. The filtered, continuous data were projected to ICA space using the unmixing matrix (i.e.,

that was calculated using the 1-s segments for ICA). The components previously identified to contain

artifacts were removed and the mixing matrix was used to project the data back to original 64 EEG-

channels. Noisy channels removed prior to ICA (for six participants, one channel each), were interpolated

following ICA as the average across neighboring channels. Afterwards, data were low-pass filtered at 30

Hz (Hann window, 111 pts) and divided into trials of 12 seconds (-3 s to 9 s time-locked to the

simultaneous onset of sound and dot movement). Finally, data were downsampled to 500 Hz and trials

that exceeded a signal change of more than 200 μV across the entire epoch were excluded from

analyses. Pooled across all participants, 0.8% of trials during the single task 0.7% of trials during the dual

task were excluded.

Analysis of time-frequency power

In order to analyze oscillatory activity, single-trial time-domain EEG signals were convolved with Morlet

wavelets. Complex wavelet coefficients were calculated for frequencies ranging from 1–20 Hz in steps

of 0.5 Hz and time from −2 s to 8 s time-locked to noise onset, separately for each trial, electrode, and

participant. Power was calculated by squaring the magnitude of the complex wavelet coefficients,

separately for each trial, electrode, and time-frequency bin. Time-frequency power representations

were then averaged across trials, separately for each condition. Power was baseline corrected to dB

power change: Trial-averaged data at each time point were divided by the mean power in the baseline

time window (−2 to −1.1 s), and subsequently log10 transformed.

224   Since we were primarily interested in changes in alpha power, we calculated alpha-power time

225   courses by averaging across frequencies ranging from 8 to 12 Hz (Jensen & Mazaheri, 2010; Klimesch et

226   al., 2007; Weisz et al., 2011). To avoid gap-related changes in parietal alpha power, we focused the

227   analysis on the time window right before the gap. We thus averaged power across parietal electrodes

228   (CPz, CP1, CP2, CP3, CP4, Pz, P1, P2, P3, P4, POz, PO3, PO4; Figure 4) and across the 3–4 s time window

229   – that is the pre-gap window (Figure 1A).

Source localization

To localize the underlying sources related to alpha power, Fieldtrip's MRI template was used as source

model (Holmes et al., 1998) along with a 3-layer boundary element model (BEM, Oostenveld et al., 2003)

as volume conductor. This head-model was used to estimate individual leadfields (Nolte, 2003). A cross-

spectral density matrix was calculated based on a fast Fourier transform using all trials per participants

centered on 10 Hz (±2Hz spectral smoothing; multitaper) for the -2–8 s time window. The cross-spectral

density matrix was used to calculate spatial filters for each source location using dynamic imaging of

coherent sources (DICS, Gross et al., 2001).

The spatial-filter coefficients resulting from the DICS calculation were multiplied with the single-

trial wavelet coefficients that were calculated in the sensor-level time-frequency analysis. Similar to the

sensor-level analysis, source localized single-trial time-frequency power was calculated by squaring the

magnitude of the complex wavelet coefficients. Time-frequency power representations were averaged

across trials, separately for each condition. Decibel (dB) power change was calculated relative to the

baseline time window of −2 to −1.1 s.

Finally, we separately averaged individual source-projected power within three pre-defined

regions of interest (ROI; auditory, visual, and parietal region; Figure 5) using functional parcels defined

by the Human Connectome Project parcellation template (Glasser et al., 2017; Keitel & Gross, 2016).

Similar to the analysis in sensor space, power was averaged in the time window of interest (3–4 s time-

locked to noise onset).

Statistical analysis

For the analysis of behavioral performance in the single task, we tested for differences in performance

accuracy and speed between the easy and the hard task difficulty using paired samples t-tests,

separately for each sensory modality. For the analysis of behavioral data in the dual-task, we used a

repeated-measures analysis of variance (rmANOVA) with the factors Auditory Difficulty (easy, hard) and

Visual Difficulty (easy, hard), separately for the auditory and visual performance measures (accuracy and

speed). Note that auditory and visual performance measures were treated separately for the behavioral

analysis, because the scales and chance levels were different between modalities. Moreover, this

allowed us to investigate the effect of visual difficulty on auditory performance and auditory difficulty

on visual performance for the dual-task data.

For the analysis of pupil size in the single task, a rmANOVA with the factors Modality (auditory,

visual) and Difficulty (easy, hard) was calculated. For the analysis of pupil size in the dual task, we

calculated a rmANOVA with the two factors Auditory Difficulty (easy, hard) and Visual Difficulty (easy,

hard). Analyses were conducted separately for the 3–4 s and the 5–6 s time window.

The details of the statistical analysis of alpha power data mirrored those for the analysis of pupil

size. For the analysis of the single task, a rmANOVA with the factors Modality (auditory, visual) and

Difficulty (easy, hard) was calculated. For the analysis of alpha power in the dual task, we calculated a

rmANOVA with the two factors Auditory Difficulty (easy, hard) and Visual Difficulty (easy, hard). For the

analysis of alpha power in source space, the additional within-participants factor ROI (auditory, parietal,

visual) was added to the rmANOVA.

We also investigated the relation between pupil size and alpha power. To this end, we used the

dual-task data for each participant and calculated the difference between the auditory & visual easy and

the auditory & visual hard condition, separately for pupil size and alpha power. The relationship between

the difference (hard minus easy) in pupil size and alpha power was analyzed using Pearson correlation.

Effect sizes for t-tests are reported as Cohen's d (Cohen, 1988). Effect sizes for ANOVAs are

reported as generalized eta square (η_g^2) (Bakeman, 2005). For null results, Bayes Factors are reported.

All statistical analyses were calculated in MATLAB (MathWorks, Inc).

Results

Behavioral performance declines with increasing task demand

For the analysis of behavioral performance (Figure 2) in the single task, we found that accuracy was

lower for the hard compared to the easy difficulty level, as expected (auditory: t_{23} = -8.60, p = 1.21×10^{-8},

d = -1.76; visual: t_{23} = -13.5, p = 2.04×10^{-12}, d = -2.76). Response speed was also slower for the hard

compared to the easy difficulty level (auditory: t_{23} = -8.32, p = 2.19×10^{-8}, d = -1.70; visual: t_{23} = -10.97,

p = 1.24×10^{-10}, d = -2.24).

The analysis of performance in the dual task was carried out separately for the auditory and

visual task. For the auditory-task performance, we observed lower accuracy and speed when the

auditory task was hard compared to easy (main effect of Auditory Difficulty; accuracy: $F_{1,23}$ = 56.48, p =

1.23×10^{-7}, η_g^2 = 0.46; speed: $F_{1,23}$ = 36.51, p = 3.67×10^{-6}, η_g^2 = 0.15), whereas performance was not

significantly affected by the concurrent visual-task difficulty (main effect of Visual Difficulty: for both

accuracy and speed p > 0.3; Auditory Difficulty × Visual Difficulty interaction: for both accuracy and speed

p > 0.1).

For the visual-task performance, we observed lower accuracy and speed when the visual task

was hard compared to easy (main effect of Visual Difficulty; accuracy: $F_{1,23}$ = 113.99, p = 2.19×10^{-10}, η_g^2

= 0.65; speed: $F_{1,23}$ = 387.78, p = 6.66 × 10^{-16}, η_g^2 = 0.43). However, visual-task accuracy was also affected

by the difficulty in the auditory task (main effect of Auditory Difficulty: $F_{1,23}$ = 9.8, p = 0.005, η_g^2 = 0.02):

accuracy in the visual task was lower when the concurrent auditory task was hard compared to easy (no

main effect of Auditory Difficulty for speed: p > 0.5). The data perhaps suggest that participants prioritize

the auditory task over the visual task (i.e., visual performance dropped when the auditory task was hard,

whereas auditory performance was unaffected by visual-task difficulty). The Auditory Difficulty × Visual

Difficulty interaction for visual performance was not significant (for both accuracy and speech p > 0.1).

Figure 2. Effects of task difficulty on performance in the single and the dual task. A: Single task. Results for the auditory gap-
detection task (left) and visual MOT task (right). Accuracy as proportion correct responses (top) and speed as inverse response
time (bottom). For both modalities, performance was lower for the hard compared to the easy condition. P-values are
obtained from paired samples t-tests. **B:** Dual task. Order is the same as in A. For both modalities, performance was lower for
the hard than the easy condition. For accuracy in the visual task, a significant main effect of auditory difficulty was observed
as well. Only significant effects of the rmANOVAs are presented. Error bars reflect the standard error of the mean. Gray lines
show data from individual participants.

Pupil size reflects demand manipulation independent of sensory modality

Single task

Pupil-size time courses for each task and difficulty condition are shown in Figure 3A. Descriptively,

difficulty-induced changes in pupil size followed modality-dependent trajectories: When participants

performed the visual MOT task, pupil size increased relatively early during the trial for difficult compared

to easy trials, whereas in the auditory gap-detection task, pupil size increased later for the hard relative

to the easy condition. This is consistent with the nature of the different tasks. The visual MOT task

requires attention throughout, whereas the auditory gap-detection task requires participants to focus

to a specific point in time.

More formally, in the 3-4 s time window, pupil size was larger for the hard compared to the easy

condition (main effect Difficulty: $F_{1,22}$ = 9.98, p = 0.005, η_g^2 = 0.03), but there was no difference in pupil

size between modalities (p > 0.07) and no interaction (p > 0.3). In the 5-6 s time window, pupil size was

also larger for the difficult relative to the easy condition (main effect of Difficulty: $F_{1,22}$ = 13.85, p = 0.001,

η_g^2 = 0.04) and for the auditory compared to the visual task (main effect of Modality: $F_{1,22}$ = 35.93, p =

4.94 x 10^{-6}, η_g^2 = 0.21). We did not find evidence for any interactive effects of task difficulty and modality

(p > 0.7).

Figure 3. Pupil-size variation in the single and dual tasks. A: Single task. Top: Pupil-size time courses. Gray areas reflect the
time windows of interest for statistical analysis, vertical gray lines indicate the time window during which a gap could occur.
Bottom: Averaged pupil size for time windows of interest. Left: Time window 3-4 s. Pupil size increased with task difficulty
independent of modality. Right: Time window 5-6 s. In addition to the difficulty effect, pupil size was larger for the auditory
compared to the visual task. Inset: 45-degree plot for main effect difficulty. Black dots show individual averaged pupil size for
each difficulty level averaged across modalities. The 45-degree line indicates no difference between conditions. **B:** Dual task.
Order is the same as in panel A. Pupil size was larger when the visual task was hard compared to easy. The Interaction
indicates that the auditory difficulty effect was greater when the visual task was easy. Inset: Black dots show individual data
points for the visual easy condition, showing the driving effect of the interaction. Error bands reflect the within-subject error.
Error bars indicate standard error of the mean. Crosshairs indicate 95% CI.

Dual task

Pupil-size time courses in the dual task are shown in Figure 3B. Time courses in the dual task visually

appear to resemble the combination of the time courses in the auditory and visual single tasks.

In the 3-4 s time window, we observed a larger pupil size for the hard than the easy condition

(main effect of Visual Difficulty: $F_{1,22}$ = 16.59, p = 0.001, η_g^2 = 0.08), whereas there was no effect of

Auditory Difficulty (p > 0.07). The Auditory Difficulty × Visual Difficulty interaction was marginally

significant ($F_{1,22}$ = 4.12, p = 0.055, η_g^2 = 0.008), showing that the increase in pupil size with auditory-task

difficulty was greater when the visual task was easy compared to difficult.

Changes in pupil size in the 5-6 s time window were similar to those in the 3-4 s time window:

pupil size was larger for the hard than the easy conditions, for both the visual and auditory task (main

effect of Visual Difficulty: $F_{1,22}$ = 10.97, p = 0.003, η_g^2 = 0.03; main effect of Auditory Difficulty: $F_{1,22}$ =

7.86, p = 0.01, η_g^2 = 0.01). The Auditory Difficulty × Visual Difficulty interaction was significant ($F_{1,22}$ =

5.10, p = 0.034, η_g^2 = 0.006), indicating again that the increase in pupil size with auditory-task difficulty

was greater when the visual task was easy compared to difficult. Pupil size only differed between the

auditory easy and hard conditions when the visual task was easy (t_{22} = 4.28, p = 3.03 × 10^{-4}, d = -0.89),

but not when the visual task was hard (p > 0.4).

Task difficulty affects alpha power differently for the auditory and visual modality

Single task

As shown in Figure 4A, alpha power was lower when participants performed the visual task compared

to the auditory task (main effect of Modality: $F_{1,23}$ = 22.35, p = 9.19 × 10^{-5}, η_g^2 = 0.28), and when task

difficulty was hard compared to easy (main effect of Difficulty: $F_{1,23}$ = 11.94, p = 0.002, η_g^2 = 0.03).

Critically, the decrease in alpha power with task difficulty was more pronounced for the visual compared

to the auditory task (Modality × Difficulty interaction: $F_{1,23}$ = 11.89, p = 0.002, η_g^2 = 0.03). Alpha power

was only more suppressed for the hard compared to the easy conditions for the visual task (t_{23} = -4.76,

p = 8.53 × 10-5, d = -0.97), whereas there was no task-difficulty effect for the auditory task (p > 0.7).

Dual task

Time courses for alpha power in the dual task are shown in Figure 4B. Alpha power was lower for the

hard compared to the easy conditions in the visual task (main effect of Visual Difficulty: $F_{1,23}$ = 15.81, p

= 0.001, η_g^2 = 0.05) and lower for the easy compared to the hard condition in the auditory task (main

effect of Auditory Difficulty: $F_{1,23}$ = 7.18, p = 0.013, η_g^2 = 0.008). Critically, the Auditory Difficulty × Visual

Difficulty interaction ($F_{1,23}$ = 8.82, p = 0.007, η_g^2 = 0.006) shows that, when the visual task was hard and

leading to an overall suppression of alpha power, this suppression was reduced when the concurrent

auditory task was hard relative to when it was easy (t_{23} = 3.87, p = 7.75 × 10^{-4}, d = 0.79; no auditory-

difficulty effect when the visual task was easy: p > 0.7). Hence, the data demonstrate that, in a competing

audio-visual situation, rising demands in the visual modality decrease alpha power (i.e., greater power

suppression), whereas rising demands in the auditory modality increase alpha power.

Figure 4. Alpha-power results in the single and the dual task. A: Single task. Top row: Alpha power time courses. Gray areas
reflect the time window of interest for statistical analysis. Right: Topographies for the time window of interest. Bottom row:
Averaged data across participants for the time window of interest. The difficulty effect was only present for visual but not for
auditory task. Inset: Difficulty effect of the visual task shown in a 45-degree plot. Black dots show averaged alpha power per
difficulty level for visual task, separately for each participant. The 45-degree line indicates no difference between conditions.
B: Dual task. Order is the same as in A. Increasing the visual demand led to greater alpha-power suppression. This suppression
was reduced when the auditory demands were high. Inset: Black dots show averaged alpha power for each participant in this
hard visual condition, showing the driving effect for the interaction. Error bands reflect the within-subject error. Error bars
indicate standard error of the mean.

Cortical regions differentially reflect modality specific demand

378 We were interested in characterizing the region specificity of the alpha oscillatory dynamics. To this end,

379 we projected alpha-power data to source space to differentiate between alpha oscillatory activity in

380 auditory, parietal, and visual cortices (Figure 5; see Methods for details).

Single Task

Source-localized alpha power was overall more suppressed during the visual compared to the auditory

task (main effect of Modality: $F_{1,23}$ = 13.45, p = 0.001, η_g^2 = 0.15). Critically, the Modality × Difficulty

interaction was significant ($F_{1,23}$ = 6.51, p = 0.018, η_g^2 = 0.009), showing that alpha power was larger for

the hard compared to the easy auditory task, whereas alpha power was smaller (i.e., more suppressed)

for the hard compared to the easy visual task (Figure 5C).

There was also a main effect of ROI ($F_{2,46}$ = 6.06, p = 0.005, η_g^2 = 0.03), showing that the

suppression of alpha power in auditory cortex was lower compared to the suppression in visual cortex

(t_{23} = 2.5, p = 0.02, d = 0.51) and parietal cortex (t_{23} = 2.5, p = 9.74 × 10^{-6}, d = 1.15). No other interaction

was significant.

Figure 5. Source-localized alpha power in three brain regions of interest. A: Single-task data for the auditory, parietal and
visual regions of interest (ROIs). Data are averaged in the pre-gap time window (3-4 s). Areas on the brain surfaces show the
ROIs. The same ROIs were used for the single- and the dual-task analyses. B: Dual-task data. C: Mean difference between the
hard and easy conditions for the auditory and the visual single task (Modality × Difficulty interaction). Alpha power was larger
for the hard compared to the easy condition during the auditory task, whereas the reverse effect was present for the visual
task. D: Condition differences to visualize significant interactions for dual-task data. Left: The difference between the hard
and easy conditions in the auditory task was larger when the visual task was difficult compared to easy. Middle: The auditory
difficulty effect (i.e., hard minus easy) was greatest in the parietal cortex. Right: Alpha-power suppression for the hard
compared to the easy visual task was greatest in visual cortex.

Dual Task

The analysis of source-localized alpha power in the dual task revealed that all three two-way interactions

were significant, whereas the three-way interaction was not significant (p > 0.89). Specifically, the

Auditory Difficulty × Visual Difficulty interaction ($F_{1,23}$ = 6.63, p = 0.017, η_g^2 = 0.007) demonstrates that

the power increase for the hard relative to the easy condition associated with the auditory task was

more pronounced when the concurrent visual task was hard compared to easy (Figure 5D, left). The ROI

× Auditory Difficulty interaction ($F_{2,46}$ = 3.27, p = 0.047, η_g^2 = 0.001) shows that the difficulty effect (i.e.,

hard minus easy) associated with the auditory task was greatest in parietal cortex (Figure 5D, middle),

and significantly larger than the difficulty effect in visual cortex (t_{23} = 2.21, p = 0.04, d = 0.45). Finally,

alpha-power suppression for the hard compared to the easy condition in the visual task was greatest in

visual cortex (ROI × Visual Difficulty interaction $F_{2,46}$ = 4.76, p = 0.013, η_g^2 = 0.005; Figure 5D, right), and

significantly smaller than in parietal cortex (t_{23} = -2.46, p = 0.02, d = -0.50).

Overall, the results of the source-localized alpha power in the dual task suggest that the greater

alpha-power suppression associated with visual-task difficulty originates primarily from visual cortex. In

contrast, the relative diminishment of this alpha-power suppression associated auditory-task difficulty

originates primarily from parietal cortex (Figure 5D).

Task difficulty influences pupil size and alpha power largely independently

In the previous sections, we have reported the effects of task difficulty and sensory modality separately

for pupil size and alpha power. One key question in the field is whether these two neurophysiological

measures of cognitive demand (or effort) are driven by a common underlying mechanism (Ala et al.,

2020; Alhanbali et al., 2019; Miles et al., 2017). If the same mechanism were to underlie changes in pupil

size and alpha power, we would expect these measures to correlate.

In order to investigate the relationship between pupil size and alpha power, we calculated

individual difficulty effects (auditory–visual hard vs. auditory–visual easy conditions in the dual-task

data) and did so separately for pupil size and alpha power. This allowed us to calculate a simple Pearson

correlation between the pupil-size difficulty effect and the alpha-power difficulty effect. Note that we

did so separately for each of the three regions of interest: auditory, parietal, and visual cortex (Figure 6).

There was no significant correlation between pupil size and alpha power in any of the three brain regions

(p > 0.05). Visual cortex exhibits an indecisive r = −.41 correlation (Bayes Factor BF_{10} ~1.5), in the auditory

and parietal cortices, however, Bayes factors (BF_{10} ~ 1/3) provide tentative evidence for the absence of

a true correlation. The overriding picture emerging is one of no unitary, underlying process driving the

effect in these two neurophysiological measures of cognitive demand.

Figure 6. No correlation between pupil size and alpha power for three regions of interest. Data points reflect the difference between the auditory & visual hard and auditory & visual easy condition for dual-task data, for pupil size and alpha power. The gray line reflects the best fitting line (correlation). Left: Auditory region of interest (ROI). Middle: Parietal ROI. Right: Visual ROI.

Discussion

In the current study, we investigated how two key and often used neurophysiological measures, pupil size and neural alpha oscillatory power, change depending on different degrees of cognitive demand under complex audio-visual dual-task conditions. First, our behavioral data suggest that individuals prioritized auditory over visual task performance. Second, we observed that changes in cognitive demand affected pupil size and alpha power differently. Pupil size increased with increasing demands in both the auditory and the visual modality, and the temporal dynamics of the pupil size indicated the specific temporal evolution of the required cognitive task demands in each modality. In contrast, changes in alpha power with increasing cognitive demand, on the other hand, differed as a function of task modality and appeared to originate from two different neural sources: Alpha power in parietal cortex increased with auditory cognitive demand, whereas alpha power in occipital cortex decreased with visual cognitive demand. Lastly, our data suggest that pupil size and alpha power index different facets of cognitive demand under complex audio-visual conditions.

Behavioral performance suggests prioritization of the auditory over the visual task

As expected, based on previous work (Björn Herrmann et al., 2022; Björn Herrmann & Johnsrude, 2018; Ritz et al., 2020; Wutz et al., 2020), behavioral performance decreased with increasing cognitive demand in the auditory and the visual task, as well as under single- and dual-task conditions. In previous uses of

dual tasks, participants were explicitly instructed to prioritize one task over the other (Gagné et al., 2017;

Picou & Ricketts, 2014; Wu et al., 2016). We avoided such prioritization instruction to investigate how

different degrees of cognitive demand in either task affect performance. Our behavioral results suggest

that participants prioritized the auditory task over the visual task, as indicated by the performance

decline in the visual task with increasing difficulty in the auditory task (Figure 1B). The data suggest that

the available cognitive resources were insufficient to perform both tasks well concurrently. This is

broadly in line with the dual-task literature showing performance decline in a secondary task with

increasing demand in a primary task (Desjardins & Doherty, 2013; Gagné et al., 2017; Gosselin & Gagné,

2014; Picou & Ricketts, 2014).

One potential reason for why participants may have prioritized the auditory over the visual task

in the current study may be related to the different timing of when cognitive investment was required

in the auditory and in the visual task. For the auditory task, participants were required to detect a single

event in time with some degree of temporal predictability (Björn Herrmann et al., 2022) as it always

occurred within 4-6 s after sound onset, whereas the visual MOT task required participants to attentively

focus throughout a trial (Cavanagh & Alvarez, 2005; Björn Herrmann & Johnsrude, 2018; Pylyshyn &

Storm, 1988; Scholl, 2009; Wutz et al., 2020). Participants may have lost the dots they were asked to

track during the moments they fully focused on and responded to the auditory gap-detection task.

Pupil size tracks cognitive demand in the auditory and visual tasks

Pupil size increased with the degree of cognitive demand for both the auditory and the visual task (Figure

3). This increase in pupil size with cognitive demand is generally consistent with previous work (Kadem

et al., 2020; Kahneman & Beatty, 1966; Koelewijn et al., 2012; Martin et al., 2020; Ohlenforst et al., 2018;

Porter et al., 2007; Stolte et al., 2020; Wendt et al., 2016; Winn et al., 2018; Zekveld et al., 2010; Zekveld

& Kramer, 2014; Zhao et al., 2019).

The time course of the pupil size mirrored the temporal evolution of the cognitive demand

manipulation in each task. The pupil-size time courses for the auditory task diverged late between task-

difficulty conditions and peaked late during a trial, mirroring the temporal occurrence of the gap. For

the visual task, pupil-size time courses diverged between task-difficulty conditions right from the

beginning of a trial and remained different throughout, potentially reflecting the need to track the

relevant dots from the beginning to the end of a trial (Cavanagh & Alvarez, 2005; Bjorn Herrmann &

Johnsrude, 2018; Björn Herrmann & Johnsrude, 2018; Pylyshyn & Storm, 1988; Scholl, 2009; Wutz et al.,

2020). Our data thus suggest high sensitivity of pupil-size changes to when in time and to what degree

participants invest cognitively during an auditory or visual task.

Under dual-task conditions, we further observed that auditory difficulty effects were only

present when the concurrent visual task was easy but not hard. This may suggest that the respective

highly demanding task is driving the pupil response. Therefore, the pupil-size time courses under dual-

task setting are more similar to single-task time courses of the auditory task when the concurrent visual

task was easy, but more similar to the visual single-task time courses when the visual task was hard.

Cognitive demand modulated alpha power differently under auditory and visual task conditions

In contrast to variations in pupil size, most likely linked to changes in global arousal or vigilance state,

alpha-oscillatory responses should offer more nuanced insights into the cognitive-demand sensitivity of

different neural systems. Indeed, alpha power in parietal cortex increased with auditory cognitive

demand, whereas alpha power in occipital cortex decreased with visual cognitive demand (Figure 5).

These findings shed light on the differential cognition-related changes in alpha power in vision

and audition. An increase in alpha power has been observed when auditory demands increase (Obleser

et al., 2012; Winneke et al., 2020; Wisniewski et al., 2017; Wöstmann et al., 2015), and this effect may

originate from an oscillator in parietal cortex (Björn Herrmann et al., 2022). In contrast, a decrease in

alpha power – that is, alpha-power suppression – has been observed when demands in a visual task

increase (Erickson et al., 2019; Magosso et al., 2019), and this effect may originate from an oscillator in

occipital cortex (Bonnefond & Jensen, 2012).

Although we observed a distinction based on sensory modality, others have suggested that

demand-dependent changes in alpha power may differ with the degree of internal (e.g., audition) or

external processing (e.g., vision) requirements (Palva & Palva, 2011). Regardless of this functional

distinction, our data emphasize the presence of multiple alpha oscillators in the brain (see also Başar et

al., 1997; Bollimunta et al., 2008; Herrmann et al., 2022; Mo et al., 2011; Wisniewski et al., 2021) that

are modulated differently by cognitive demand, depending on task modality.

A dominant line of thinking assumes alpha oscillatory activity to reflect functional inhibition, such

that brain regions in which alpha power increases are inhibited (Foxe & Snyder, 2011; Jensen &

Mazaheri, 2010; Klimesch et al., 2007; Weisz et al., 2011). Our observations that alpha power decreases

with increasing difficulty in the visual task is consistent with this function inhibition view, potentially

reducing inhibition in visual cortices.

In contrast, the increase in alpha power in parietal cortex with increased cognitive demand in

the auditory task may be less consistent with function inhibition, except if we were to assume that

parietal cortex is selectively inhibited when individuals invest cognitively in the auditory task.

Nonetheless, this increase in alpha power under auditory attention is consistent with other previous

work (Obleser et al., 2012; Winneke et al., 2020; Wisniewski et al., 2017; Wöstmann et al., 2015), and

more generally aligns with recent suggestions that the functional inhibition hypothesis may not

generalize across sensory modalities (Ai & Ro, 2014; Björn Herrmann et al., 2016; Linkenkaer-Hansen et

al., 2004).

Pupil size and alpha power may reflect different facets of cognitive investment

Pupil size and alpha power were both modulated by cognitive demand, although in somewhat different

ways. Perhaps not surprisingly, given the different sensitivity to cognitive demand of the two measures,

the degree to which cognitive demand affected pupil size was not significantly correlated with the

degree to which cognitive demand affected alpha power (Figure 6). The absence of a correlation

between pupil size and alpha power is consistent with previous work that also did not find a relationship

(Ala et al., 2020; Alhanbali et al., 2019; Miles et al., 2017). One potential reason for the missing

relationship might be that the two measures largely reflect separable neural pathways associated with

cognitive resource recruitment (Alhanbali et al., 2019; Strand et al., 2018). The present study shows that

the pupil size tracked the cognitive demand induced by the different tasks more closely. Pupil size may

thus be taken as the more intuitive index for cognitive demand, also in terms of sign and magnitude of

the pupil signal: A greater pupil size means higher cognitive demand (Kahneman & Beatty, 1966; Pichora-

Fuller et al., 2016). As discussed above, assigning an equally parsimonious interpretation to alpha-power

changes is not an accurate representation of the extant empirical data.

In order to interpret alpha-power changes, the different alpha oscillators and their spatial mixing

in EEG recordings (perhaps less in magnetoencephalography recordings) need to be considered and

could explain the lack of the relation between pupil size and alpha power. Alpha power may perhaps

index more clearly the cognitive demands in auditory tasks when stimulation is devoid of visual input

(Henry et al., 2017; Björn Herrmann et al., 2022; Paul et al., 2021; Wöstmann et al., 2015), and vice versa

in visual tasks (Bonnefond & Jensen, 2012; Erickson et al., 2019; Magosso et al., 2019; Roijendijk et al., 2013). The current data provide a detailed picture of the multi-faceted, modality-dependent changes in alpha power under complex audio-visual task conditions.

Conclusions

Our results show that pupil size and parietal alpha power are not two interchangeable neurophysiological measures of cognitive investment or effort, but reflect different neurophysiological facets of task-induced cognitive demands. Specifically, we demonstrate that pupil size tracks increase in cognitive demand independently of the sensory modality that induces the demand. However, the changes in magnitude of neural alpha power associated with task demands depend greatly on the sensory modality from which the demands originate. Alpha power in parietal cortex increases with auditory cognitive demand, whereas alpha power in occipital cortex decreases with visual cognitive demand. Finally, our data add on the amounting evidence that pupil size and alpha power variations are not driven by a unitary and putatively noradrenergically governed pathway. Overall, the current study demonstrates that the dynamics of the neurophysiological indices of cognitive effort are multi-faceted and potentially modality-dependent under complex audio-visual task conditions.

References

Ai, L., & Ro, T. (2014). The phase of prestimulus alpha oscillations affects tactile perception. *Journal of Neurophysiology*, *111*(6), 1300–1307. https://doi.org/10.1152/jn.00125.2013

Ala, T. S., Graversen, C., Wendt, D., Alickovic, E., Whitme, W. M., & Lunner, T. (2020). An exploratory study of EEG alpha oscillation and pupil dilation in hearing-aid users during effortful listening to continuous speech. *PLoS ONE*, *15*(7 July), 1–15. https://doi.org/10.1371/journal.pone.0235782

Alhanbali, S., Dawes, P., Millman, R. E., & Munro, K. J. (2019). Measures of Listening Effort Are Multidimensional. *Ear and Hearing*, *40*(5), 1084–1097. https://doi.org/10.1097/AUD.0000000000000697

Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus-norepinephrine function: Adaptive gain and optimal performance. *Annual Review of Neuroscience*, *28*, 403–450. https://doi.org/10.1146/annurev.neuro.28.061604.135709

Bakeman, R. (2005). Recommended effect size statistic. *Behavior Research Methods*, *37*(3), 379–384.
https://link.springer.com/content/pdf/10.3758%2FBF03192707.pdf

Başar, E., Schürmann, M., Başar-Eroglu, C., & Karakaş, S. (1997). Alpha oscillations in brain functioning:
An integrative theory. *International Journal of Psychophysiology*, *26*(1–3), 5–29.
https://doi.org/10.1016/S0167-8760(97)00753-8

Bollimunta, A., Chen, Y., Schroeder, C. E., & Ding, M. (2008). Neuronal mechanisms of cortical alpha
oscillations in awake-behaving macaques. *Journal of Neuroscience*, *28*(40), 9976–9988.
https://doi.org/10.1523/JNEUROSCI.2699-08.2008

Bonnefond, M., & Jensen, O. (2012). Alpha oscillations serve to protect working memory maintenance
against anticipated distracters. *Current Biology*, *22*(20), 1969–1974.
https://doi.org/10.1016/j.cub.2012.08.029

Breton-Provencher, V., & Sur, M. (2019). Active control of arousal by a locus coeruleus GABAergic
circuit. *Nature Neuroscience*, *22*(2), 218–228. https://doi.org/10.1038/s41593-018-0305-z

Cavanagh, P., & Alvarez, G. A. (2005). Tracking multiple targets with multifocal attention. *Trends in*
Cognitive Sciences, *9*(7), 349–354. https://doi.org/10.1016/j.tics.2005.05.009

Cohen, J. (1988). Statistical power analysis for the behavioral sciences. Lawrence Erlbaum Associates.
Hillsdale, NJ, 20–26.

Dahl, M. J., Mather, M., Sander, M. C., & Werkle-Bergner, M. (2020). Noradrenergic responsiveness
supports selective attention across the adult lifespan. *Journal of Neuroscience*, *40*(22), 4372–
4390. https://doi.org/10.1523/JNEUROSCI.0398-19.2020

Dahl, M. J., Mather, M., & Werkle-Bergner, M. (2022). Noradrenergic modulation of rhythmic neural
activity shapes selective attention. *Trends in Cognitive Sciences*, *26*(1), 38–52.
https://doi.org/https://doi.org/10.1016/j.tics.2021.10.009

Desjardins, J. L., & Doherty, K. A. (2013). Age-related changes in listening effort for various types of
masker noises. *Ear and Hearing*, *34*(3), 261–272.
https://doi.org/10.1097/AUD.0b013e31826d0ba4

Dimitrijevic, A., Smith, M. L., Kadis, D. S., & Moore, D. R. (2017). Cortical alpha oscillations predict
speech intelligibility. *Frontiers in Human Neuroscience*, *11*.
https://doi.org/10.3389/fnhum.2017.00088

Erickson, M. A., Smith, D., Albrecht, M. A., & Silverstein, S. (2019). Alpha-band desynchronization
reflects memory-specific processes during visual change detection. *Psychophysiology*, *56*(11), 1–9.
https://doi.org/10.1111/psyp.13442

Fink, L., Simola, J., Tavano, A., Lange, E., Wallot, S., & Laeng, B. (2021). From pre-processing to
advanced dynamic modeling of pupil data. *PsyArXiv*, 1–56.

Foxe, J. J., & Snyder, A. C. (2011). The role of alpha-band brain oscillations as a sensory suppression mechanism during selective attention. *Frontiers in Psychology*, *2*(JUL), 1–13. https://doi.org/10.3389/fpsyg.2011.00154

Gagné, J. P., Besser, J., & Lemke, U. (2017). Behavioral assessment of listening effort using a dual-task paradigm: A review. *Trends in Hearing*, *21*, 1–25. https://doi.org/10.1177/2331216516687287

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Essen, D. C. Van. (2017). Europe PMC Funders Group Europe PMC Funders Author Manuscripts A multi-modal parcellation of human cerebral cortex. *Nature*, *536*(7615), 171–178. https://doi.org/10.1038/nature18933.A

Gosselin, P. A., & Gagné, J. (2014). Older Adults Expend More Listening Effort Than. *Journal of Speech, Language, and Hearing Research*, *54*(June 2011), 944–958. https://doi.org/10.1044/1092-4388(2010/10-0069)a

Gross, J., Kujala, J., Hämäläinen, M., Timmermann, L., Schnitzler, A., & Salmelin, R. (2001). Dynamic imaging of coherent sources: Studying neural interactions in the human brain. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(2), 694–699. https://doi.org/10.1073/pnas.98.2.694

Henry, M. J., Herrmann, B., Kunke, D., & Obleser, J. (2017). Aging affects the balance of neural entrainment and top-down neural modulation in the listening brain. *Nature Communications*, *8*. https://doi.org/10.1038/ncomms15801

Herrmann, Björn, Henry, M. J., Haegens, S., & Obleser, J. (2016). Temporal expectations and neural amplitude fluctuations in auditory cortex interactively influence perception. *NeuroImage*, *124*, 487–497. https://doi.org/10.1016/j.neuroimage.2015.09.019

Herrmann, Bjorn, & Johnsrude, I. S. (2018). Neural signatures of the processing of temporal patterns in sound. *Journal of Neuroscience*, *38*(24), 5466–5477. https://doi.org/10.1523/JNEUROSCI.0346-18.2018

Herrmann, Björn, & Johnsrude, I. S. (2018). Attentional State Modulates the Effect of an Irrelevant Stimulus Dimension on Perception. *Journal of Experimental Psychology: Human Perception and Performance*, *44*(1), 89–105. https://doi.org/10.1037/xhp0000432

Herrmann, Björn, & Johnsrude, I. S. (2020). A model of listening engagement (MoLE). *Hearing Research*, *397*. https://doi.org/10.1016/j.heares.2020.108016

Herrmann, Björn, Maess, B., Henry, M. J., Obleser, J., & Johnsrude, I. S. (2022). *Task-related temporal fluctuations in auditory attention generate neural signatures that change with age*.

Herrmann, Björn, Maess, B., & Johnsrude, I. S. (2018). Aging affects adaptation to sound-level statistics in human auditory cortex. *Journal of Neuroscience*, *38*(8), 1989–1999. https://doi.org/10.1523/JNEUROSCI.1489-17.2018

Holmes, C. J., Hoge, R., Collins, L., Woods, R., Toga, A. W., & Evans, A. C. (1998). Enhancement of MR
Images Using Registration for Signal Averaging. *Journal of Computer Assisted Tomography*, *22*(2).
https://journals.lww.com/jcat/Fulltext/1998/03000/Enhancement_of_MR_Images_Using_Registr
ation_for.32.aspx

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity: Gating
by inhibition. *Frontiers in Human Neuroscience*, *4*(November), 1–8.
https://doi.org/10.3389/fnhum.2010.00186

Joshi, S., & Gold, J. I. (2020). Pupil Size as a Window on Neural Substrates of Cognition. *Trends in*
Cognitive Sciences, *24*(6), 466–480. https://doi.org/10.1016/j.tics.2020.03.005

Joshi, S., Li, Y., Kalwani, R. M., & Gold, J. I. (2016). Relationships between Pupil Diameter and Neuronal
Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron*, *89*(1), 221–234.
https://doi.org/10.1016/j.neuron.2015.11.028

Kadem, M., Herrmann, B., Rodd, J. M., & Johnsrude, I. S. (2020). Pupil Dilation Is Sensitive to Semantic
Ambiguity and Acoustic Degradation. *Trends in Hearing*, *24*.
https://doi.org/10.1177/2331216520964068

Kahneman, D. (1973). *Attention and Effort* (Vol. 88, Issue 2). Prentice Hall.
https://doi.org/10.2307/1421603

Kahneman, D., & Beatty, J. (1966). Pupil Diameter and Load on Memory. *Science*, 1583–1586.

Keitel, A., & Gross, J. (2016). Individual Human Brain Areas Can Be Identified from Their Characteristic
Spectral Activation Fingerprints. *PLoS Biology*, *14*(6), 1–22.
https://doi.org/10.1371/journal.pbio.1002498

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing
hypothesis. In *Brain Research Reviews* (Vol. 53, Issue 1, pp. 63–88).
https://doi.org/10.1016/j.brainresrev.2006.06.003

Koelewijn, T., Zekveld, A. A., Festen, J. M., & Kramer, S. E. (2012). Pupil dilation uncovers extra listening
effort in the presence of a single-talker masker. *Ear and Hearing*, *33*(2), 291–300.
https://doi.org/10.1097/AUD.0b013e3182310019

Lavie, N. (1995). Perceptual Load as a Necessary Condition for Selective Attention. *Journal of*
Experimental Psychology: Human Perception and Performance, *21*(3), 451–468.
https://doi.org/10.1037/0096-1523.21.3.451

Linkenkaer-Hansen, K., Nikulin, V. V., Palva, S., Ilmoniemi, R. J., & Palva, J. M. (2004). Prestimulus
oscillations enhance psychophysical performance in humans. *Journal of Neuroscience*, *24*(45),
10186–10190. https://doi.org/10.1523/JNEUROSCI.2584-04.2004

Magosso, E., De Crescenzio, F., Ricci, G., Piastra, S., & Ursino, M. (2019). EEG alpha power is modulated
by attentional changes during cognitive tasks and virtual reality immersion. *Computational*

Intelligence and Neuroscience, *2019*. https://doi.org/10.1155/2019/7051079

Martin, J. T., Whittaker, A. H., & Johnston, S. J. (2020). Component processes in free-viewing visual
search: Insights from fixation-aligned pupillary response averaging. *Journal of Vision*, *20*(7), 1–24.
https://doi.org/10.1167/JOV.20.7.5

Mick, P., Kawachi, I., & Lin, F. R. (2014). The association between hearing loss and social isolation in
older adults. *Otolaryngology - Head and Neck Surgery (United States)*, *150*(3), 378–384.
https://doi.org/10.1177/0194599813518021

Miles, K., McMahon, C., Boisvert, I., Ibrahim, R., de Lissa, P., Graham, P., & Lyxell, B. (2017). Objective
Assessment of Listening Effort: Coregistration of Pupillometry and EEG. *Trends in Hearing*, *21*, 1–
13. https://doi.org/10.1177/2331216517706396

Mo, J., Schroeder, C. E., & Ding, M. (2011). Attentional modulation of alpha oscillations in macaque
inferotemporal cortex. *Journal of Neuroscience*, *31*(3), 878–882.
https://doi.org/10.1523/JNEUROSCI.5295-10.2011

Montefusco-Siegmund, R., Schwalm, M., Jubal, E. R., Devia, C., Egaña, J. I., & Maldonado, P. E. (2022).
Alpha EEG Activity and Pupil Diameter Coupling during Inactive Wakefulness in Humans. *ENeuro*,
9(2). https://doi.org/10.1523/ENEURO.0060-21.2022

Murphy, P. R., Robertson, I. H., Balsters, J. H., & O'connell, R. G. (2011). Pupillometry and P3 index the
locus coeruleus-noradrenergic arousal function in humans. *Psychophysiology*, *48*(11), 1532–1543.
https://doi.org/10.1111/j.1469-8986.2011.01226.x

Nolte, G. (2003). The magnetic lead field theorem in the quasi-static approximation and its use for
magnetoenchephalography forward calculation in realistic volume conductors. *Physics in
Medicine and Biology*, *48*(22), 3637–3652. https://doi.org/10.1088/0031-9155/48/22/002

Obleser, J., Wöstmann, M., Hellbernd, N., Wilsch, A., & Maess, B. (2012). Adverse listening conditions
and memory load drive a common alpha oscillatory network. *Journal of Neuroscience*, *32*(36),
12376–12383. https://doi.org/10.1523/JNEUROSCI.4908-11.2012

Ohlenforst, B., Wendt, D., Kramer, S. E., Naylor, G., Zekveld, A. A., & Lunner, T. (2018). Impact of SNR,
masker type and noise reduction processing on sentence recognition performance and listening
effort as indicated by the pupil dilation response. *Hearing Research*, *365*, 90–99.
https://doi.org/10.1016/j.heares.2018.05.003

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence
and Neuroscience*, *2011*. https://doi.org/10.1155/2011/156869

Oostenveld, R., Stegeman, D. F., Praamstra, P., & Van Oosterom, A. (2003). Brain symmetry and
topographic analysis of lateralized event-related potentials. *Clinical Neurophysiology*, *114*(7),
1194–1202. https://doi.org/10.1016/S1388-2457(03)00059-2

Palva, S., & Palva, J. M. (2011). Functional roles of alpha-band phase synchronization in local and large-
scale cortical networks. In *Frontiers in Psychology* (Vol. 2, Issue SEP).
https://doi.org/10.3389/fpsyg.2011.00204

Paul, B. T., Chen, J., Le, T., Lin, V., & Dimitrijevic, A. (2021). Cortical alpha oscillations in cochlear
implant users reflect subjective listening effort during speech-in-noise perception. *PLoS ONE*, *16*(7
July), 1–22. https://doi.org/10.1371/journal.pone.0254162

Petersen, E. B., Wöstmann, M., Obleser, J., Stenfelt, S., & Lunner, T. (2015). Hearing loss impacts neural
alpha oscillations under adverse listening conditions. *Frontiers in Psychology*, *6*(FEB), 1–11.
https://doi.org/10.3389/fpsyg.2015.00177

Pichora-Fuller, M. K., Kramer, S. E., Eckert, M. A., Edwards, B., Hornsby, B. W. Y., Humes, L. E., Lemke,
U., Lunner, T., Matthen, M., Mackersie, C. L., Naylor, G., Phillips, N. A., Richter, M., Rudner, M.,
Sommers, M. S., Tremblay, K. L., & Wingfield, A. (2016). Hearing impairment and cognitive energy:
The framework for understanding effortful listening (FUEL). *Ear and Hearing*, *37*, 5S-27S.
https://doi.org/10.1097/AUD.0000000000000312

Picou, E. M., & Ricketts, T. A. (2014). The effect of changing the secondary task in dual-task paradigms
for measuring listening effort. *Ear and Hearing*, *35*(6), 611–622.
https://doi.org/10.1097/AUD.0000000000000055

Porter, G., Troscianko, T., & Gilchrist, I. D. (2007). Effort during visual search and counting: Insights
from pupillometry. *Quarterly Journal of Experimental Psychology*, *60*(2), 211–229.
https://doi.org/10.1080/17470210600673818

Pylyshyn, Z. W., & Storm, R. W. (1988). Tracking multiple independent targets: Evidence for a parallel
tracking mechanism. *Spatial Vision*, *3*(3), 179–197.

Ritz, H., Wild, C., & Johnsrude, I. (2020). Parametric cognitive load reveals hidden costs in the neural
processing of perfectly intelligible degraded speech. *BioRxiv*, 2020.10.02.324509.
https://doi.org/10.1101/2020.10.02.324509

Roijendijk, L., Farquhar, J., Van Gerven, M., Jensen, O., & Gielen, S. (2013). Exploring the impact of
target eccentricity and task difficulty on covert visual spatial attention and its implications for
brain computer interfacing. *PLoS ONE*, *8*(12). https://doi.org/10.1371/journal.pone.0080489

Scholl, B. J. (2009). What have we learned about attention from multiple-object tracking (and vice
versa)? In *Computation, cognition, and Pylyshyn.* (pp. 49–77). MIT Press.

Shukla, A., Harper, M., Pedersen, E., Goman, A., Suen, J. J., Price, C., Applebaum, J., Hoyer, M., Lin, F.
R., & Reed, N. S. (2020). Hearing Loss, Loneliness, and Social Isolation: A Systematic Review.
Otolaryngology - Head and Neck Surgery (United States), *162*(5), 622–633.
https://doi.org/10.1177/0194599820910377

Stolte, M., Gollan, B., & Ansorge, U. (2020). Tracking visual search demands and memory load through

pupil dilation. *Journal of Vision*, *20*(6), 1–19. https://doi.org/10.1167/JOV.20.6.21

Strand, J. F., Brown, V. A., Merchant, M. B., Brown, H. E., & Smith, J. (2018). Measuring listening effort:
Convergent validity, sensitivity, and links with cognitive and personality measures. *Journal of*
Speech, Language, and Hearing Research, *61*(6), 1463–1486.
https://doi.org/10.1044/2018_JSLHR-H-17-0257

Sung, Y. K., Li, L., Blake, C., Betz, J., & Lin, F. R. (2016). Association of Hearing Loss and Loneliness in
Older Adults. *Journal of Aging and Health*, *28*(6), 979–994.
https://doi.org/10.1177/0898264315614570

Vazey, E. M., Moorman, D. E., & Aston-Jones, G. (2018). Phasic locus coeruleus activity regulates
cortical encoding of salience information. *Proceedings of the National Academy of Sciences of the*
United States of America, *115*(40), E9439–E9448. https://doi.org/10.1073/pnas.1803716115

Weisz, N., Hartmann, T., Müller, N., Lorenz, I., & Obleser, J. (2011). Alpha rhythms in audition:
Cognitive and clinical perspectives. *Frontiers in Psychology*, *2*(APR), 1–15.
https://doi.org/10.3389/fpsyg.2011.00073

Wendt, D., Dau, T., & Hjortkjær, J. (2016). Impact of background noise and sentence complexity on
processing demands during sentence comprehension. *Frontiers in Psychology*, *7*(MAR), 1–12.
https://doi.org/10.3389/fpsyg.2016.00345

Winn, M. B., Edwards, J. R., & Litovsky, R. Y. (2016). The Impact of Auditory Spectral Resolution on
Listening Effort Revealed by Pupil Dilation. In *Ear and hearing* (Vol. 36, Issue 4).
https://doi.org/10.1097/AUD.0000000000000145.The

Winn, M. B., Wendt, D., Koelewijn, T., & Kuchinsky, S. E. (2018). Best Practices and Advice for Using
Pupillometry to Measure Listening Effort: An Introduction for Those Who Want to Get Started.
Trends in Hearing, *22*, 1–32. https://doi.org/10.1177/2331216518800869

Winneke, A. H., Schulte, M., Vormann, M., & Latzel, M. (2020). Effect of Directional Microphone
Technology in Hearing Aids on Neural Correlates of Listening and Memory Effort: An
Electroencephalographic Study. *Trends in Hearing*, *24*.
https://doi.org/10.1177/2331216520948410

Wisniewski, M. G., Thompson, E. R., & Iyer, N. (2017). Theta- and alpha-power enhancements in the
electroencephalogram as an auditory delayed match-to-sample task becomes impossibly difficult.
Psychophysiology, *54*(12), 1916–1928. https://doi.org/10.1111/psyp.12968

Wisniewski, M. G., Zakrzewski, A. C., Bell, D. R., & Wheeler, M. (2021). EEG power spectral dynamics
associated with listening in adverse conditions. *Psychophysiology*, *58*(9), 1–19.
https://doi.org/10.1111/psyp.13877

Wöstmann, M., Herrmann, B., Wilsch, A., & Obleser, J. (2015). Neural alpha dynamics in younger and
older listeners reflect acoustic challenges and predictive benefits. *Journal of Neuroscience*, *35*(4),

1458–1467. https://doi.org/10.1523/JNEUROSCI.3250-14.2015

Wöstmann, M., Lim, S. J., & Obleser, J. (2017). The Human Neural Alpha Response to Speech is a Proxy
of Attentional Control. *Cerebral Cortex*, *27*(6), 3307–3317. https://doi.org/10.1093/cercor/bhx074

Wu, Y. H., Stangl, E., Zhang, X., Perkins, J., & Eilers, E. (2016). Psychometric Functions of Dual-Task
Paradigms for Measuring Listening Effort. *Ear and Hearing*, *37*(6), 660–670.
https://doi.org/10.1097/AUD.0000000000000335

Wutz, A., Zazio, A., & Weisz, N. (2020). Oscillatory bursts in parietal cortex reflect dynamic attention
between multiple objects and ensembles. *Journal of Neuroscience*, *40*(36), 6927–6937.
https://doi.org/10.1523/JNEUROSCI.0231-20.2020

Zekveld, A. A., & Kramer, S. E. (2014). Cognitive processing load across a wide range of listening
conditions: Insights from pupillometry. *Psychophysiology*, *51*(3), 277–284.
https://doi.org/10.1111/psyp.12151

Zekveld, A. A., Kramer, S. E., & Festen, J. M. (2010). Pupil response as an indication of effortful
listening: The influence of sentence intelligibility. *Ear and Hearing*, *31*(4), 480–490.
https://doi.org/10.1097/AUD.0b013e3181d4f251

Zhao, S., Bury, G., Milne, A., & Chait, M. (2019). Pupillometry as an Objective Measure of Sustained
Attention in Young and Older Listeners. *Trends in Hearing*, *23*.
https://doi.org/10.1177/2331216519887815

795