1   Plasma after both SARS-CoV-2 boosted vaccination and COVID-19 potently neutralizes BQ.1.1
2   and XBB.1
3   Running title: COVID-19 convalescent plasma neutralizes BQ.1.1.
4

5   David J Sullivan[1], Massimo Franchini[2], Jonathon W. Senefeld[3], Michael J. Joyner[3], Arturo
6   Casadevall[1], Daniele Focosi[4]
7
[1]Johns Hopkins Bloomberg School of Public Health and School of Medicine, Baltimore, MD
21218, USA; dsulliv7@jhmi.edu; acasade1@jhu.edu
[2]Division of Transfusion Medicine, Carlo Poma Hospital, 46100 Mantua, Italy;
massimo.franchini@asst-mantova.it
[3]Department of Anesthesiology & Perioperative Medicine, Mayo Clinic, Rochester, MN 55902,
USA joyner.michael@mayo.edu; senefeld.jonathon@mayo.edu
[4]North-Western Tuscany Blood Bank, Pisa University Hospital, 56124 Pisa, Italy.
daniele.focosi@gmail.com
16

Keywords: convalescent plasma; SARS-CoV-2; COVID-19; BQ.1.1; XBB; BF.7: virus
neutralization.
Abbreviations: VOC: variant of concern; VaxCCP, plasma from both vaccinated and COVID-
19 convalescent subjects.

Author contributions: DJS: data extraction, table curation and design, and writing original
draft; D.F.: conceptualization, data access and verification and manuscript revision; M.F. data
access and verification and manuscript revision; JWS data access and verification and
manuscript revision, MJJ and AC manuscript revision.
Data availability statement: all the data used for this manuscript are available in the
Supplementary Dataset file.
27

28 **Corresponding author**: David Sullivan, 615 N. Wolfe St Rm W4606, Baltimore MD 21205.
29 Tel 410 502 2522, fax 410 955 0105. E-mail dsulliv7@jhmi.edu

30
Word counts: Abstract 199; Text – 2087.

Abstract

Objectives

Recent 2022 SARS-CoV-2 Omicron variants, have acquired resistance to most neutralizing anti-
Spike monoclonal antibodies authorized, and the BQ.1.* sublineages are notably resistant to all
authorized monoclonal antibodies. Polyclonal antibodies from individuals both vaccinated and
recently recovered from Omicron COVID-19 (VaxCCP) could retain new Omicron neutralizing
activity.

Methods

Here we reviewed BQ.1.* virus neutralization data from 920 individual patient samples from 43
separate cohorts defined by boosted vaccinations with or without recent Omicron COVID-19, as
well as infection without vaccination.

Results

More than 90% of the plasma samples from individuals in the recently (within 6 months) boosted
VaxCCP study cohorts neutralized BQ.1.1, and BF.7 with 100% neutralization of WA-1, BA.4/5,
BA.4.6 and BA.2.75. The geometric mean of the geometric mean 50% neutralizing titers (GM
(GMT$_{50}$) were 314, 78 and 204 for BQ.1.1, XBB.1 and BF.7, respectively. Compared to VaxCCP,
plasma sampled from COVID-19 naïve subjects who also recently within 6 months received at
least a third vaccine dose had about half of the GM (GMT$_{50}$) for all viral variants.

Conclusions

Boosted VaxCCP characterized by either recent vaccine dose or infection event within 6 months
represents a robust, variant-resilient, passive immunotherapy against the new Omicron BQ.1.1,
XBB.1 and BF.7 variants.

Table of contents

75
76

Introduction

In immunocompromised (IC) patients both passive immunotherapies and small molecule antivirals are often necessary to treat COVID-19 or eliminate persistently high SARS-CoV-2 viral load. Chronic, persistent viral loads increase both transmission and mutation risk, and prevent administration of the required immunosuppressive/antineoplastic therapies[1]. Small molecule antivirals have not been formally validated for IC patients, who often have contraindications, and the convergent evolution of the Omicron variant of concern (VOC) has led to inefficacy of all the anti-Spike monoclonal antibodies (mAbs) authorized so far for both treatment or prevention, e.g. in the highly prevalent BQ.1.* sublineages[2]. The other rapidly growing XBB.* and BF.7 sublineages are also highly resistant to anti-Spike mAbs[3]. Polyclonal plasma from individuals who are both vaccinated and had COVID-19 (VaxCCP) has more than ten times the antibody levels capable of neutralizing pre-Omicron variants as well as Omicron variants BA.1 through BA.4/5[4, 5]. Polyclonal COVID-19 convalescent plasma (CCP) has thousands of distinct antibody specificities of different isotypes, including many capable of SARS-CoV-2 neutralization. High-titer pre-Omicron CCP contains Omicron neutralizing activity despite being collected before variant appearance[4, 5].

Given that CCP remains a recommended therapy for IC[1, 6, 7], we systematically reviewed recent primary research for neutralization results against BQ.1.1 by plasma collected from vaccinated subjects with or without COVID-19 or after recent Omicron infection alone.

Results

Ten articles were included (**Figure 1**) which contained virus neutralizations with WA-1, BQ.1.1, BA.4/5, BA.4.6, XBB.1 and BF.7, assessed with either live authentic SARS-CoV-2 or SARS-CoV-2 pseudovirus neutralization assays and represented data from 920 patients (**Supplementary Table 1**). Qu *et al*. in the USA reported on Spring and Summer 2022 breakthrough infections with BA.1 and BA.4/5 in two sampled cohorts with predominantly unvaccinated individuals, as well as a third cohort of healthcare workers after a single monovalent booster vaccination in the Fall of 2021[8] (**Table 1**). Zou *et al*. in the USA in the Summer and Fall of 2022 sampled individuals who had already received 3 mRNA BNT162b2 vaccinations with or without previous COVID-19, both before and about 4 weeks after a 4^{th} monovalent or bivalent vaccine booster vaccination[9]. Miller *et al*. also in the USA sampled both before the 3^{rd} vaccination dose and about 4 weeks after monovalent mRNA vaccination in the Fall of 2021, as well as with the 4th vaccine dose in the Summer or Fall of 2022, with either monovalent or bivalent booster vaccinations in Fall of 2022 in those with no documented COVID-19[10]. Cao *et al*. in China investigated BQ.1.1 neutralizations from plasma of 4 cohorts after 3 doses of CoronaVac (Fall 2021) without COVID-19 or 2-12 weeks after BA.1, BA.2 and BA.5 infection[3]. Planas *et al*. in France evaluated GMT_{50} in plasma from individuals both 4 and 16 weeks after a third monovalent mRNA vaccine dose in the Fall of 2021 as well as 12 and 32 weeks after vaccine breakthrough BA.1/2 or BA.5 infection[11]. Davis *et al* in the USA sampled after the 3^{rd} mRNA vaccine monovalent dose in the Fall of 2021 and also after either a 4th monovalent mRNA dose or a bivalent (wild-type + BA.4/5) vaccine dose in the Summer and Fall of 2022[12]. Kurhade *et al* in the USA also compared GMT_{50} after the 4^{th} monovalent vaccine dose or 3 mRNA doses with the 4^{th} the bivalent dose without COVID-19 and also after bivalent boost with recent COVID-19[13]. Wang *et al* in the USA compared GMT_{50} after three vaccine doses, the 4^{th} monovalent vaccine dose or 3 mRNA doses with the 4^{th} the bivalent dose without COVID-19, and also after 2-3 vaccine doses and recent

BA.2 breakthrough infection or 3-4 mRNA vaccine doses and recent BA.4/5 breakthrough
infection(14). Ito et al in Japan compared breakthrough infections after BA.2 and BA.5 after 2-3
doses of mRNA vaccines in the Spring and Summer of 2022(15). Akerman et al in Australia
characterized neutralizing antibodies in four groups 1) sampling one to three months after 3 doses
of mRNA vaccines with an Omicron infection in 2022; 2) sampling 3 months after 4 doses of
mRNA vaccine; 3) sampling 6 months after 3 doses of mRNA vaccine and 4) sampling 3-6 months
after last vaccine in a larger cohort who had the original WA-1 infection in early 2020 as well as
3 more doses of mRNA vaccine(16).
131
These diverse cohorts were assembled into 3 groups, 1) plasma after both 2-4 vaccine doses and
COVID-19 (VaxCCP); 2) plasma from subjects after administration of 3-4 vaccine doses (i.e.
boosted), but either self-reported as COVID-19-*naïve* or anti-nucleocapsid negative; and 3)
Omicron infection without vaccination (CCP) as well as participants sampled 6 to 11 months after
previous vaccine dose and before the booster vaccination. Boosted VaxCCP neutralized BQ.1.1,
XBB.1 and BF.7 with approximately 3 times the dilutional potency of the vaccine-only or 2-6
times CCP/pre-boost vaccination groups for all viral variants (**Table 2 and Figure 2**). Importantly,
while there was a 19-fold reduction in neutralization by boosted VaxCCP against BQ.1.1 compared
to WA-1, more than 90% of the boosted VaxCCP samples neutralized BQ.1.1 as well as XBB.1
and BF.7 (**Table 2 and Figure 2c**). Three cohorts within the boosted VaxCCP group were below
at 90% neutralization with one sampled late, 8 months after BA.1/2 breakthrough infection(11)
and the other two from a single study after BA.2 and BA.5 (**Supplementary Table 2 and 3**).
Except for the GMT (GMT_{50}) against XBB.1 at 78, the other viral variant neutralizations were in
the same range as pre-Alpha CCP neutralizing WA-1 (i.e., 311)(4). By comparison the large
randomized clinical trial which effectively reduced outpatient COVID-19 progression to
hospitalizations had a GMT_{50} of 60 for WA-1 with pre-Alpha CCP(17). Boosted vaccinations at
3-4 doses without COVID-19, showed GM (GMT_{50}) of 118 for BQ.1.1, with only 6 of 23 cohorts
over 90% neutralizations, for 79% overall (i.e. 326 of 414 individuals). Four separate
studies(8),(13),(12),(10) characterized BQ.1.1 virus neutralizations with plasma after the new
bivalent (wild-type + BA.4/5) mRNA vaccine booster in the Fall of 2022, with 88% (103 of 117
samples) neutralization activity within 4 weeks of bivalent booster (**Supplementary Table 3**).
153
Many studies performed virus neutralizations on samples drawn before the 3rd or 4th vaccine dose
which were 6 to 11 months after last vaccine dose. The GM (GMT_{50})'s for BQ.1.1 and BA.2.75
were about 6 times reduced compared to VaxCCP even though the fold reductions were similar
(**Figure 3**, **Table 2**). In agreement with lower GMT_{50} for neutralizations was the low percent
neutralizing BQ.1.1 (63%), XBB.1 (50%), and BF.7 (75%) at 6 to 11 months after vaccination
(**Figure 3, Table 2 and Supplementary Table 3**).
160
Five studies used the lentiviral pseudovirus assays, with diverse Spike proteins cloned in, while
the other four were live virus assays using different cell types (**Supplementary Table 1**). Notably,
Planas *et al* employed the IGROV-1 cell type for better growth of Omicron sublineages(11). While
the single study fold reductions (FR) and percent neutralizations normalize the results between
studies, the GMT_{50} can vary between studies even amongst the live authentic viral neutralization
studies (e.g., mNeonGreen™ reporter assays versus cytopathic effects)(9, 13). We sorted the live
authentic viral neutralizations from the pseudoviral neutralizations, plotting also the minimum and
maximums (**Supplementary Figures 1-3**). In general, the live authentic SARS-CoV-2

neutralization assays for VaxCCP appeared to have similar antibody neutralization levels, with the
single study by Cao et al(3) employing lentiviral pseudovirus with lower dilutional titers. In
contrast, the GMT_{50} achieved with pseudoviral assays in the boosted vaccinations without COVID-
19 appeared slightly higher than the ones achieved with authentic virus.
173
174 Discussion
175 The FDA deemed CCP safe and effective for both immunocompetent and IC COVID-19
176 outpatients(6, 7, 18), and further extended its authorized use in the IC patient population in
177 December 2021(7, 18), at a time when oral antiviral therapy promised a no transfusion outpatient
178 solution and many anti-Spike mAbs were still effective.
179
Up until the present, CCP remained a backup bridge for IC patients, durable against the changing
variants and as a salvage therapy in seronegative IC patients. With the recent advent of Omicron
XBB.* and BQ.1.* defeating the remaining anti-Spike mAbs, boosted VaxCCP, recently collected
within the last 6 months of either a vaccine dose or SARS-CoV-2 is likely to be the only viable
remaining passive antibody therapy in the 2022-23 Winter for IC patients who have failed to make
antibodies after vaccination and still require B-cell depleting drugs or immunosuppressive therapy.
In a literature review of CCP from diverse VOC waves as well as boosted vaccinees and VaxCCP
up to BA-1, VaxCCP showed higher neutralization titers against Omicron at levels above 300
dilutional GMT_{50}[4].
189
The accelerated evolution of SARS-CoV-2 VOCs has created the problem that the pharmaceutical
development of additional mAbs is not worth the effort and cost given their expected short useful
clinical life expectancy, so the anti-Spike mAb pipeline has remained stuck in 2022. High levels
of antibodies in donor plasma from both boosted vaccinations and COVID-19 convalescent plasma
(VaxCCP) neutralizing more than 93% of BQ.1.1and BF.7, with XBB.1 at 89%. Recently collected
plasma within a 6 months window from those boosted vaccinees without prior documented
COVID-19 had a 20-30% reduction in neutralization percent for BQ.1.1and XBB.1 with 10%
reduction for the others and a third of the GM (GMT_{50}) neutralizing antibody levels compared to
VaxCCP. In those vaccinated with last dose more than 6 months prior to sample collection, both
the neutralization percent and neutralizing antibody titers fell further, compared to the recently
boosted VaxCCP group. Four studies (Planas(11), Zou(9), Cao(3) and Kurhade(13)) had directly
comparative cohorts in the three groups which increases the robustness reduction in neutralizations
with the vaccine only or more than 6 months to last vaccine or infection event compared to
VaxCCP. The main limitation of our systematic review is the small number of studies reporting
virus neutralization with BQ.1.1 with most available as pre-preprints without peer-review yet.
However, we note that peer-review itself does not change GMT_{50} or neutralization numbers and
the authors of these papers have considerable expertise in the topic.
207
Boosted VaxCCP has full potential to replace anti-Spike mAbs for passive antibody therapy of IC
patients against recent Omicron sublineages, in the meanwhile polyclonal IgG formulations can
be manufactured. VaxCCP qualification in the real-world will likely remain constrained on high-
throughput serology, whose correlation with GMT_{50} is not perfect(19, 20). Nevertheless, the very
high prevalence (93%) of Omicron-neutralizing antibodies and the high GM (GMT_{50}) in recently
boosted VaxCCP reassure about its potency, and further confirm that exact donor-recipient VOC

matching is dispensable. Overall, our findings urge WHO to revise its guidelines and recommend
boosted VaxCCP for therapy of COVID-19 in IC patients.
216
217

Search strategy and selection criteria
On November 19, 2022 we initially searched PubMed, medRxiv and bioRxiv for manuscripts
reporting BQ.1.1 neutralization, using English language as a restriction. Search of bioRxiv with
same keywords now yields 17 records of which only 10 contained plasma viral neutralization data.
Search of medRxiv produced 3 records which did not have BQ.1.1 neutralizations. PubMed
retrieved 3 entries using ("BQ.1.1") and ("neutralization"), one of which was focused on anti-
Spike mAb alone(2) and the other 2 were duplicates from bioRxiv(8, 12). Articles underwent
evaluation for data extraction by two assessors (DS and DF) with disagreements resolved by third
assessor (AC). Articles lacking plasma BQ.1.1 virus neutralizations were excluded. The process
of study selection is represented in the PRISMA flow diagram (**Figure 1**).
228
The type of viral assay (live or pseudovirus), time interval to blood sample, GMT_{50}, minimum and
maximum neutralizing 50% dilutional titer for WA-1 (pre-Alpha wild-type) and Omicron
sublineages BQ.1.1, BA.4/5, BA.4.6, BA.2.75, XBB.1 and BF.7 and number out of total that
neutralized Omicron were abstracted from study text, graphs and tables. Two studies (Wang(14)
and Qu(8)) reported BQ.1 and those were separate cohorts in addition to BQ.1.1. Prism v. 9.4
(GraphPad Software, San Diego, CA, USA) was used for data analysis. While all manuscripts
included neutralization data against WA-1, BQ.1.1, BA.4/5 and BA.2.75, only a subset of
manuscripts included neutralization data for BA.4.6, XBB.1 and BF.7 which were assembled for
relevance to present circulating variants. Historic early Omicron partial neutralization data on
variants like BA.1 or BA.2 were excluded because of the full set data with BA.4/5 and BA.2.75.
239
240 Statistical significance between $\log_{10}$ transformed GMT_{50} was investigated using Tukey's test. The
241 multiple comparison test was a two-way ANOVA with Alpha 0.05 on log transformed GMT_{50}.
242 The log normal test was performed on WA-1, BQ.1.1, BA.4/5, BA.4.6, XBB.1 and BF.7 virus
243 GMT_{50}. Two studies(10, 11) reported the median titer rather than the GMT_{50}. Compiled data
244 abstracted from the published studies is available in the supplementary dataset.
245

Funding
This study was supported by the U.S. Department of Defense's Joint Program Executive Office
for Chemical, Biological, Radiological and Nuclear Defense (JPEO-CBRND), in collaboration
with the Defense Health Agency (DHA) (contract number: W911QY2090012) (DS), with
additional support from Bloomberg Philanthropies, State of Maryland, the National Institutes of
Health (NIH) National Institute of Allergy and Infectious Diseases 3R01AI152078-01S1 (DS,
AC), NIH National Center for Advancing Translational Sciences U24TR001609-S3 and
UL1TR003098.

Role of the funding source
The study sponsors did not contribute to the study design; the collection, analysis, and
interpretation of data; manuscript preparation, and the decision to submit the paper for publication.

The views expressed are those of the authors and should not be construed to represent the positions
of the U.S. Army or the Department of Defense. The data and opinions presented do not reflect
the view of the U.S. government.
261
Conflict of interest disclosure
DJS reports AliquantumRx Founder and Board member with stock options (macrolide for
malaria), Hemex Health malaria diagnostics consulting and royalties for malaria diagnostic test
control standards to Alere- all outside of submitted work. AC reports being part of the scientific
advisory board of SabTherapeutics and has received personal fees from Ortho Diagnostics,
outside of the submitted work. All other authors report no relevant disclosures.
268
269

References

1.	Senefeld JW, Franchini M, Mengoli C, Cruciani M, Zani M, Gorman EK, et al. COVID-19 convalescent plasma for the treatment of immunocompromised patients: a systematic review. medRxiv. 2022:2022.08.03.22278359.

2.	Arora P, Kempf A, Nehlmeier I, Schulz SR, Jack HM, Pohlmann S, et al. Omicron sublineage BQ.1.1 resistance to monoclonal antibodies. Lancet Infect Dis. 2022 Nov 18. PubMed PMID: 36410372. Epub 2022/11/22.

3.	Cao Y, Jian F, Wang J, Yu Y, Song W, Yisimayi A, et al. Imprinted SARS-CoV-2 humoral immunity induces convergent Omicron RBD evolution. bioRxiv. 2022:2022.09.15.507787.

4.	Sullivan DJ, Franchini M, Joyner MJ, Casadevall A, Focosi D. Analysis of anti-SARS-CoV-2 Omicron-neutralizing antibody titers in different vaccinated and unvaccinated convalescent plasma sources. Nature Communications. 2022 2022/10/29;13(1):6478.

5.	Li M, Beck EJ, Laeyendecker O, Eby YJ, Tobian AA, Caturegli P, et al. Convalescent plasma with a high level of virus-specific antibody effectively neutralizes SARS-CoV-2 variants of concern. Blood Adv. 2022:2022.03.01.22271662.

6.	Hinton DM. Convalescent Plasma EUA Letter of Authorization 06032021. In: FDA, editor. FDA in brief: FDA; 2021.

7.	O'Shaughnessy JA. Convalescent Plasma EUA Letter of Authorization 12282021 . FDA website2021.

8.	Qu P, Evans JP, Faraone J, Zheng Y-M, Carlin C, Anghelina M, et al. Distinct Neutralizing Antibody Escape of SARS-CoV-2 Omicron Subvariants BQ.1, BQ.1.1, BA.4.6, BF.7 and BA.2.75.2. bioRxiv. 2022:2022.10.19.512891.

9.	Zou J, Kurhade C, Patel S, Kitchin N, Tompkins K, Cutler M, et al. Improved Neutralization of Omicron BA.4/5, BA.4.6, BA.2.75.2, BQ.1.1, and XBB.1 with Bivalent BA.4/5 Vaccine. bioRxiv. 2022:2022.11.17.516898.

10.	Miller J, Hachmann NP, Collier A-rY, Lasrado N, Mazurek CR, Patio RC, et al. Substantial Neutralization Escape by the SARS-CoV-2 Omicron Variant BQ.1.1. bioRxiv. 2022:2022.11.01.514722.

11.	Planas D, Bruel T, Staropoli I, Guivel-Benhassine F, Porrot F, Maes P, et al. Resistance of Omicron subvariants BA.2.75.2, BA.4.6 and BQ.1.1 to neutralizing antibodies. bioRxiv. 2022:2022.11.17.516888.

12.	Davis-Gardner ME, Lai L, Wali B, Samaha H, Solis D, Lee M, et al. mRNA bivalent booster enhances neutralization against BA.2.75.2 and BQ.1.1. bioRxiv. 2022:2022.10.31.514636.

13.	Kurhade C, Zou J, Xia H, Liu M, Chang HC, Ren P, et al. Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1, and XBB.1 by 4 doses of parental mRNA vaccine or a BA.5-bivalent booster. bioRxiv. 2022:2022.10.31.514580.

14.	Wang Q, Iketani S, Li Z, Liu L, Guo Y, Huang Y, et al. Alarming antibody evasion properties of rising SARS-CoV-2 BQ and XBB subvariants. bioRxiv. 2022:2022.11.23.517532.

15.	Ito J, Suzuki R, Uriu K, Itakura Y, Zahradnik J, Deguchi S, et al. Convergent evolution of the SARS-CoV-2 Omicron subvariants leading to the emergence of BQ.1.1 variant. bioRxiv. 2022:2022.12.05.519085.

16.   Akerman A, Milogiannakis V, Jean T, Esneu C, Silva MR, Ison T, et al. Emergence and
antibody evasion of BQ and BA.2.75 SARS-CoV-2 sublineages in the face of maturing antibody
breadth at the population level. medRxiv. 2022:2022.12.06.22283000.
17.   Sullivan DJ, Gebo KA, Shoham S, Bloch EM, Lau B, Shenoy AG, et al. Early Outpatient
Treatment for Covid-19 with Convalescent Plasma. N Engl J Med. 2022 Mar 30. PubMed
PMID: 35353960. Pubmed Central PMCID: PMC9006786. Epub 2022/03/31.
18.   Villa C. EUA 26382 COVID-19 Convalescent Plasma. In: OBRR, editor. FDA
(https://www.fda.gov/media/155159/download)2021.
19.   Stone M, Grebe E, Sulaeman H, Di Germanio C, Dave H, Kelly K, et al. Evaluation of
Commercially Available High-Throughput SARS-CoV-2 Serologic Assays for Serosurveillance
and Related Applications. Emerg Infect Dis. 2022 Mar;28(3):672-83. PubMed PMID: 35202525.
Pubmed Central PMCID: PMC8888213. Epub 2022/02/25.
20.   Patel EU, Bloch EM, Clarke W, Hsieh YH, Boon D, Eby Y, et al. Comparative
Performance of Five Commercially Available Serologic Assays To Detect Antibodies to SARS-
CoV-2 and Identify Individuals with High Neutralizing Titers. J Clin Microbiol. 2021 Jan
21;59(2). PubMed PMID: 33139419. Epub 2020/11/04.
330

Table 1

Synopsis of included studies, reporting plasma sources, epoch of sampling, region, time since vaccination/infection to plasma sampling, and sample size. The cohorts were split into three groups-1) boosted vaccinations and recent COVID-19 (VaxCCP), 2) boosted vaccines only without documented COVID-19 (Vac only) and 3) infection alone or pre-boosted sampling before 3rd or 4th vaccine dose (Infection only or pre-boost)

Study	Vaccine and COVID-19 history at sample time	Group	Time period of plasma sampling	Geography	Sampling time mean or median (min-max)	Sample number
Cao(3)	3xCorVac+BA.1 inf	VaxCCP	Spring 2022	China	5-7 weeks post hosp admit (42 weeks avg)	50
Cao(3)	3xCorVac +BA.2 inf	VaxCCP	Summer 2022	China	3-11 weeks post hosp admit (8 weeks mean)	39
Cao(3)	3xCorVac +BA.5 inf	VaxCCP	Summer/Fall 2022	China	2-11 weeks (mean 5 weeks)	36
Zou(9)	4xBNT162b2+BTI	VaxCCP	Summer/Fall 2022	USA	4 weeks post dose	20
Planas(11)	mRNAvacx3+ BA.1/2 inf	VaxCCP	Spring/Fall 2022	France	32 weeks post BTI BA.1/2	13
Wang(14)	2-3xmRNAvac+ BA.2 BTI	VaxCCP	Spring/Fall 2022	USA	2-23 weeks (mean 6 wk (3over 90 days))	14
Wang(14)	3-4xmRNAvac+ BA.4/5 BTI	VaxCCP	Summer/Fall 2022	USA	2-8 weeks (mean 4 weeks)	20
Kurhade(13)	3xmRNAvac+bivalent+BTI	VaxCCP	Fall 2022	USA	4 weeks post with infection history	23
Ito(15)	2-3xmRNAvac+BA.2 BTI	VaxCCP	Spring 2022	Japan	2-8 weeks	14
Ito(15)	2-3xmRNAvac+BA.5 BTI	VaxCCP	Summer 2022	Japan	2-3 weeks	20
Zou(9)	3xBNT162b2+bivalent+BTI	VaxCCP	Summer/Fall 2022	USA	4 weeks post dose	19
Akerman(16)	3xmRNA +bivalent	VaxCCP	Fall 2022	Australia	4-12 weeks post BTI	29
Planas(11)	3xmRNAvac+ BA.1/2 inf	VaxCCP	Spring/Fall 2022	France	12 weeks post BTI BA.1/2	16
Planas(11)	3xmRNAvac+ BA.5 inf	VaxCCP	Fall 2022	France	8 weeks post BTI BA.5	15
Davis(12)	3xmRNAvac	Vac only	Fall 2021	USA	1-4 weeks post boost	12
Kurhade(13)	4xmRNAvac	Vac only	Summer 2022	USA	4-12 weeks	25
Cao(3)	3xCorVac	Vac only	Fall 2021	China	4 weeks	40
Zou(9)	4xBNT162b2	Vac only	Summer/Fall 2022	USA	4 weeks post dose	20
Planas(11)	3xmRNAvac	Vac only	Winter 2021/2022	France	16 weeks post 3rd dose	10
Wang(14)	3xmRNAvac	Vac only	Fall 2021	USA	2-12 weeks (mean 5 weeks)	14
Wang(14)	3xmRNAvac+monovalent	Vac only	Summer/Fall 2022	USA	3-4 weeks	19
Wang(14)	3xmRNAvac+bivalent	Vac only	Summer/Fall 2022	USA	3-4 weeks	21
Davis(12)	3xmRNAvac+monovalent	Vac only	Summer/Fall 2022	USA	10-15 weeks post boost	12
Akerman(16)	4xmRNA	Vac only	Fall 2022	Australia	12 weeks	23
Akerman(16)	3xmRNAvac after 2020 WA-1	Vac only	Summer/Fall 2022	Australia	3-6 months	47

Kurhade(13)	3xmRNAvac+bivalent	Vac only	Fall 2022	USA	4 weeks post	29	
Davis(12)	3xmRNAvac+bivalent	Vac only	Summer/Fall 2022	USA	2-6 weeks post booster (8 no vacc. 10 no infection)	12	
Qu(8)	3xmRNAvac	Vac only	Fall 2021	USA	2-13 weeks	15	
Zou(9)	3xBNT162b2+bivalent	Vac only	Summer/Fall 2022	USA	4 week post dose	18	
Planas(11)	3xmRNAvac	Vac only	Fall/Winter 2021	France	4 weeks post 3rd dose	18	
Miller(10)	3xBNT162b2	Vac only	Fall 2021	USA	2-4 weeks	16	
Miller(10)	3xmRNA+ monovalent	Vac only	Spring/Fall 2022	USA	2-9 weeks	18	
Miller(10)	3xmRNA +bivalent	Vac only	Fall 2022	USA	2-3 weeks	15	
Qu(8)	BA.4/5 inf (17-unvac)	Inf only	Summer 2022	USA	not stated	20	
Qu(8)	Hosp BA.1 (6-unvac;5-2xmRNAvac)	Inf only	Spring 2022	USA	1 week post hospitalization	15	
Zou(9)	3xBNT162b2+BTI	preboost with BNT162b	Summer/Fall 2022	USA	preboost with BNT162b (6-11 months post last dose)	20	
Zou(9)	3xBNT162b2 +BTI	preboost with bivalent	Summer/Fall 2022	USA	preboost with bivalent (6-11 months post last dose)	19	
Zou(9)	3xBNT162b2	preboost with bivalent	Summer/Fall 2022	USA	preboost with bivalent (6-11 months post last dose)	18	
Zou(9)	3xBNT162b2	preboost with BNT162b	Summer/Fall 2022	USA	preboost with BNT162b (6-11 months post last dose)	20	
Akerman(16)	3xmRNA	Preboost	Fall 2022	Australia	preboost (6 months)	47	
Miller(10)	2xBNT162b2	preboost with BNT162b	Fall 2021	USA	preboost (6-11 months post last dose)	16	
Miller(10)	3xmRNA	preboost with bivalent	Fall 2022	USA	preboost with bivalent (6-11 months post last dose)	15	
Miller(10)	3xmRNA	preboost with monovalent	Spring/Fall 2022	USA	preboost with monovalent (6-11 months post last dose)	18	

Table 2

GM (GMT_{50}) of plasma from three different sources against recent Omicron sublineages.

Neutralization virus	WA-1	BQ.1.1	BA.4/5	BA.4.6	BA.2.75	XBB.1	BF.7
Post COVID-19/vaccine (study cohorts)	12	16	14	7	9	9	4
GM (GMT_{50})	5876*	314	987	346	303	78	204
Fold reduction from WA-1	ref	19	6	17	19	75	32
Samples tested	294	237**	328	135	231	125	148
Samples neutralizing	285	221	326	135	230	111	146
Percent neutralizing	97	93	99	100	100	89	99
Boosted vaccine (study cohorts)	19	23	19	10	14	10	7
GM (GMT_{50})	3766	118	346	126	107	52	357
Fold reduction from WA-1	ref	32	11	30	35	72	12
Samples tested	384	414	384	206	261	217	158
Samples neutralizing	383	326	363	191	231	125	149
Percent neutralizing	100	79	95	93	89	58	94
Infection only or preboosted vaccine (study cohorts)	10	12	10	7	9	5	5
GM (GMT_{50})	870	47	96	103	57	23	110
Fold reduction from WA-1	ref	19	9	8	15	38	9
Samples tested	184	220	184	136	162	101	84
Samples neutralizing	182	139	144	104	104	50	63
Percent neutralizing	99	63	78	76	64	50	75

*Pre-Alpha CCP from 27 different studies had a GM (GMT_{50}) of 311 from 707 samples with 315 or 45% neutralizing omicron BA.1(4).

** percent neutralizations after CoronaVac and Omicron COVID-19 in the paper by Cao *et al* could not be retrieved from the manuscript. 237 samples from the 6 other cohorts were used for percent neutralization.

347 # Figure 1

348 PRISMA flowchart for the current study. Number of records identified from various sources,
349 excluded by manual screening, found eligible and included according to subgroup analyses.
350

 Figure 2

 Neutralizing GMT (GMT$_{50}$) against WA-1, BQ.1.1, BA.4/5, BA.4.6, BA.2.75, XBB, BF.7. A) post boosted
 vaccinations and COVID-19 and B) boosted vaccinated plasma without COVID-19. Geometric standard
 deviation for error bars, fold reduction (FR) below data, and number of studies above x-axis. Geomeans
 statistically significant in difference by multiple comparison in Tukey's test are indicated. C) The percent
 of total samples within a study which neutralized Omicron BQ.1.1 graphed in increasing percentages on
 left y axis with the total number of samples tested on the right y axis.

Figure 3

Geometric mean neutralizing titers (GMT$_{50}$) against WA-1, BQ.1.1, BA.4/5, BA.4.6, BA.2.75, XBB, BF.7 A) plasma Omicron infection alone or pre-boosted-6 to 11 months after last vaccine dose sampled in 2021 or 2022. Geometric standard deviation for error bars, fold reduction (FR) above data, and number of studies above x-axis. GM (GMT$_{50}$) statistically significant in difference by multiple comparison in Tukey's test are indicated. B). The percent of total samples within a study which neutralized Omicron BQ.1.1 graphed in increasing percentages on left y-axis with the total number of samples tested on the right y-axis.

Supplementary Table 1

Virus neutralization assays.

Reference	Assay	Virus	Replication-competent cells	neutralization threshold
Qu(8)	Pseudovirus	Lentiviral pseudovirus	HEK293T-ACE-2	80
Miller(10)	Pseudovirus	Lentiviral pseudovirus	HEK293T-ACE-2	20
Cao(3)	Pseudovirus	VSV pseudovirus	HEK293T-ACE-2	20
Wang(14)	Pseudovirus	VSV pseudovirus	VeroE6-	100
Ito(15)	Pseudovirus	Lentiviral pseudovirus	HEK293T-ACE-2	150
Akerman(16)	Live virus	Live authentic SARS-CoV-2	VeroE6-TMPRSS2	20
Davis(12)	Live virus	Live authentic SARS-CoV-2	VeroE6-TMPRSS2	20
Kurhade(13)	Live virus	mNeonGreen reporter USA-WA1/2020 SARS-CoV-2	Vero E6-TMPRSS2	20
Planas(11)	Live virus	Live authentic SARS-CoV-2	IGROV-1 or Vero E6-TMPRSS2	30
Zou(9)	Live virus	mNeonGreen reporter USA-WA1/2020 SARS-CoV-2	Vero E6-TMPRSS2	20

374 GMT_{50} of different plasma sources against BQ.1.1, BA.4/5, BA.4.6, BA.2.75, XBB.1 and BF.7 and fold-reductions (FR) compared to
375 WA-1.

papers	Vaccine and COVID-19 history at sample time	Group	#	WA-1 GMT_{50}	BQ.1.1 GMT_{50}	FR BQ.1.1	BA.4/5 GMT_{50}	FR BA.4/5	BA.4.6 GMT_{50}	FR BA.4.6	BA.2.75 GMT_{50}	FR BA.2.75	XBB.1 GMT_{50}	FR XBB	BF.7 GMT_{50}	FR BF.7
Cao(3)	3xCorVac+BA.1 BTI	VaxCCP	50	1557	27	58	107	15	23	68	197	8	23	68	70	22
Cao(3)	3xCorVac +BA.2 BTI	VaxCCP	39	1245	40	31	175	7	22	57	217	6	23	54	97	13
Cao(3)	3xCorVac +BA.5 BTI	VaxCCP	36	1136	77	15	508	2	27	42	145	8	27	42	208	5
Zou(9)	4xBNT162b2+BTI	VaxCCP	20	5120	132	39	629	8	587	9	265	19	99	52		
Planas(11)	3xmRNAvac+ BA.1/2 BTI	VaxCCP	13	8000	200	40	400	20	500	16	200	40				
Wang(14)	2-3xmRNAvac+ BA.2 BTI	VaxCCP	14	24970	849	29	3727	7					186	134		
Wang(14)	2-3xmRNAvac+ BA.2 BTI--BQ.1	VaxCCP	14		1250	20										
Wang(14)	3-4xmRNAvac+ BA.4/5 BTI	VaxCCP	20	20507	671	31	5541	4					214	96		
Wang(14)	3-4xmRNAvac+ BA.4/5 BTI--BQ.1	VaxCCP	20		1644	12										
Kurhade(13)	3xmRNAvac+bivalent+ BTI	VaxCCP	23	5776	267	22	1558	4	744	8	367	16	103	56	1223	5
Ito(15)	2-3xmRNAvac+BA.2 BTI	VaxCCP	14		400	7	800									
Ito(15)	2-3xmRNAvac+BA.5 BTI	VaxCCP	20		400	5	2100									
Zou(9)	3xBNT162b2+bivalent+ BTI	VaxCCP	19	4847	444	11	1377	4	1564	3	326	15	131	37		
Akerman(16)	3xmRNA +bivalent	VaxCCP	29	1748	88	20	468	4	289	6			139	13		
Planas(11)	3xmRNAvac+ BA.1/2 BTI	VaxCCP	16	25000	700	36	1000	25	2000	13	600	42				
Planas(11)	3xmRNAvac+ BA.5 BTI	VaxCCP	15	30000	3000	10	10000	3	9000	3	900	33				
Davis(12)	3xmRNAvac	Vac only	12	758	19	40	50	15			23	33				
Kurhade(13)	4xmRNAvac	Vac only	25	1533	22	70	95	16	62	25	26	59	15	102	69	22
Cao(3)	3xCorVac	Vac only	40	652	24	27	72	9	21	31	90	7	20	33	45	14
Zou(9)	4xBNT162b2	Vac only	20	1325	26	51	89	15	92	14	37	36	17	78		
Planas(11)	3xmRNAvac	Vac only	10	1500	40	38	60	25	60	25	10	150				
Wang(14)	3xmRNAvac	Vac only	14	7687	139	55	628	12					108	71		

Wang(14)	3xmRNAvac--BQ.1	Vac only	14		208	37										
Wang(14)	3xmRNAvac+monovalent	Vac only	19	21182	261	81	1540	14					137	155		
Wang(14)	3xmRNAvac+monovalent--BQ.1	Vac only	19		496	43										
Wang(14)	3xmRNAvac+bivalent	Vac only	21	13736	337	41	1688	8					162	85		
Wang(14)	3xmRNAvac+bivalent--BQ.1	Vac only	21		568	24										
Davis(12)	3xmRNAvac+monovalent	Vac only	12	1812	53	34	142	13			65	28				
Akerman(16)	4xmRNA	Vac only	23	1141	80	14	257	4	179	6			117	10		
Akerman(16)	3xmRNAvac after 2020 WA-1	Vac only	47	1075	70	15	212	5	110	10			54	20		
Kurhade(13)	3xmRNAvac+bivalent	Vac only	29	3620	73	50	298	12	183	20	98	37	35	103	305	12
Davis(12)	3xmRNAvac+bivalent	Vac only	12	2312	112	21	576	4			201	12				
Qu(8)	3xmRNAvac--BQ.1	Vac only	15		140	19										
Qu(8)	3xmRNAvac	Vac only	15	2616	114	23	300	9	246	11	589	4			238	11
Zou(9)	3xBNT162b2+bivalent	Vac only	18	2237	143	16	518	4	524	4	117	19	55	41		
Planas(11)	3xmRNAvac	Vac only	18	6000	200	30	300	20	300	20	60	100				
Miller(10)	3xBNT162b2	Vac only	16	45695	261	175	887	52			387	118			595	77
Miller(10)	3xmRNA+ monovalent	Vac only	18	21507	406	53	2829	8			745	29			2276	9
Miller(10)	3xmRNA +bivalent	Vac only	15	40515	508	80	3693	11			883	46			2399	17
Qu(8)	BA.4/5 inf (17-unvac)	Inf only	20	707	66	11	190	4	180	4	210	3			162	4
Qu(8)	BA.4/5 inf (17-unvac)--BQ.1	Inf only	20		68	10										
Qu(8)	Hosp BA.1 (6-unvac;5-2xmRNAvac)	Inf only	15	720	145	5	263	3	205	4	186	4			227	3
Qu(8)	Hosp BA.1 (6-unvac;5-2xmRNAvac)--BQ.1	Inf only	15		135	5										
Zou(9)	3xBNT162b2+BTI	Preboost with BNT162b	20	2516	60	42	226	11	283	9	126	20	55	46		
Zou(9)	3xBNT162b2 +BTI	Preboost with bivalent	19	1377	74	19	207	7	282	5	62	22	27	51		
Zou(9)	3xBNT162b2	Preboost with bivalent	18	**226**	**11**	**21**	**20**	**11**	**24**	**9**	**14**	**16**	**12**	**19**		
Zou(9)	3xBNT162b2	Preboost with BNT162b	20	**303**	**17**	**18**	**30**	**10**	**36**	**8**	**18**	**17**	**13**	**23**		
Akerman(16)	3xmRNA	Preboost	47	**203**	**34**	**6**	**63**	**3**	**50**	**4**			**26**	**8**		

Miller(10)	2xBNT162b2	Preboost with BNT162b	16	484	20	24	20	24			20	24			20	24
Miller(10)	3xmRNA	Preboost with bivalent	15	3633	45	81	211	17			33	110			131	28
Miller(10)	3xmRNA	Preboost with monovalent	18	5731	49	117	184	31			117	49			168	34

Supplementary Table 3

Neutralizing activity numbers (#) by study cohort.

papers	Vaccine and COVID-19 history at sample time	groups	WA-1 number	WA-1 neutraliz ing number	BQ.1 .1 numb er	BQ.1.1 neutraliz ing number	BA.4 /5 numb er	BA.4/5 neutraliz ing number	BA.4 .6 numb er	BA.4.6 neutraliz ing number	BA.2. 75 numb er	BA.2.75 neutraliz ing number	XBB .1 numb er	XBB.1 neutraliz ing number	BF.7 numb er	BF.7 neutraliz ing number
Cao(3)	3xCorVac+BA.1 BTI	VaxCCP	50	50			50	50			50	50			50	49
Cao(3)	3xCorVac +BA.2 BTI	VaxCCP	39	39			39	39			39	39			39	38
Cao(3)	3xCorVac +BA.5 BTI	VaxCCP	36	36			36	36			36	36			36	36
Zou(9)	4xBNT162b2+BTI	VaxCCP	20	20	20	20	20	20	20	20	20	20	20	19		
Planas(11)	3xmRNAvac+ BA.1/2 BTI	VaxCCP	13	13	13	11	13	12	13	13	13	11				
Wang(14)	2-3xmRNAvac+ BA.2 BTI	VaxCCP	14	14	14	13	14	14					14	8		
Wang(14)	2-3xmRNAvac+ BA.2 BTI--BQ.1	VaxCCP			14	13										
Wang(14)	3-4xmRNAvac+ BA.4/5 BTI	VaxCCP	20	20	20	18	20	20					20	14		
Wang(14)	3-4xmRNAvac+ BA.4/5 BTI--BQ.1	VaxCCP			20	20										
Kurhade(13)	3xmRNAvac+bivale nt+BTI	VaxCCP	23	23	23	22	23	23	23	23	23	23	23	22	23	23
Ito(15)	2- 3xmRNAvac+BA.2 BTI	VaxCCP			14	11	14	13								
Ito(15)	2- 3xmRNAvac+BA.5 BTI	VaxCCP			20	14	20	20								
Zou(9)	3xBNT162b2+bivale nt+BTI	VaxCCP	19	19	19	19	19	19	19	19	19	20	19	19		
Akerman (16)	3xmRNA +bivalent	VaxCCP	29	20	29	29	29	29	29	29			29	29		
Planas(11)	3xmRNAvac+ BA.1/2 BTI	VaxCCP	16	16	16	16	16	16	16	16	16	16				
Planas(11)	3xmRNAvac+ BA.5 BTI	VaxCCP	15	15	15	15	15	15	15	15	15	15				
Davis(12)	3xmRNAvac	Vac only	12	12	12	6	12	11			12	9				
Kurhade(13)	4xmRNAvac	Vac only	25	25	25	15	25	23	25	22	25	17	25	8	25	21
Cao(3)	3xCorVac	Vac only	40	40			40	39			40	40			40	37
Zou(9)	4xBNT162b2	Vac only	20	20	20	15	20	8	20	19	20	19	20	10		

Study	Vaccine	Type														
Planas(11)	3xmRNAvac	Vac only	10	10	10	6	10	7	10	5	10	3				
Wang(14)	3xmRNAvac	Vac only	14	14	14	5	14	14							14	3
Wang(14)	3xmRNAvac--BQ.1	Vac only			14	6										
Wang(14)	3xmRNAvac+monovalent	Vac only	19	19	19	15	19	19							19	8
Wang(14)	3xmRNAvac+monovalent--BQ.1	Vac only			19	16										
Wang(14)	3xmRNAvac+bivalent	Vac only	21	21	21	16	21	21							21	9
Wang(14)	3xmRNAvac+bivalent--BQ.1	Vac only			21	18										
Davis(12)	3xmRNAvac+monovalent	Vac only	12	12	12	9	12	12			12	10				
Akerman(16)	4xmRNA	Vac only	23	23	23	20	23	23	23	22					23	17
Akerman(16)	3xmRNAvac after 2020 WA-1	Vac only	47	46	47	34	47	45	47	43					47	32
Kurhade(13)	3xmRNAvac+bivalent	Vac only	29	29	29	26	29	29	29	28	29	28	29	20	29	28
Davis(12)	3xmRNAvac+bivalent	Vac only	12	12	12	10	12	12			12	11				
Qu(8)	3xmRNAvac--BQ.1	Vac only			15	14										
Qu(8)	3xmRNAvac	Vac only	15	15	15	12	15	15	15	15	15	14			15	14
Zou(9)	3xBNT162b2+bivalent	Vac only	18	18	19	18	18	18	19	19	19	18	19	18		
Planas(11)	3xmRNAvac	Vac only	18	18	18	16	18	18	18	18	18	13				
Miller(10)	3xBNT162b2	Vac only	16	16	16	16	16	16			16	16			16	16
Miller(10)	3xmRNA+ monovalent	Vac only	18	18	18	18	18	18			18	18			18	18
Miller(10)	3xmRNA +bivalent	Vac only	15	15	15	15	15	15			15	15			15	15
Qu(8)	BA.4/5 inf (17-unvac)	Inf only	20	20	20	12	20	15	20	14	20	15			20	14
Qu(8)	BA.4/5 inf (17-unvac)--BQ.1	Inf only			20	11										
Qu(8)	Hosp BA.1 (6-unvac;5-2xmRNAvac)	Inf only	15	14	15	13	15	11	15	12	15	8			15	14

Qu(8)	Hosp BA.1 (6-unvac;5-2xmRNAvac)--BQ.1	Inf only			15	14										
Zou(9)	3xBNT162b2+BTI	Preboost with BNT162b	20	20	20	16	20	19	20	19	20	17	20	17		
Zou(9)	3xBNT162b2 +BTI	Preboost with bivalent	19	19	19	16	19	19	19	19	19	17	19	12		
Zou(9)	**3xBNT162b2**	**Preboost with bivalent**	18	18	19	1	18	9	19	9	19	5	19	2		
Zou(9)	3xBNT162b2	**Preboost with BNT162b**	20	19	20	6	20	12	20	12	20	8	20	5		
Akerman (16)	**3xmRNA**	**Preboost**	23	23	23	16	23	21	23	19			23	14		
Miller(10)	2xBNT162b2	**Preboost with BNT162b**	16	16	16	6	16	5			16	7			16	4
Miller(10)	3xmRNA	**Preboost with bivalent**	15	15	15	13	15	15			15	9			15	14
Miller(10)	3xmRNA	**Preboost with monovalent**	18	18	18	15	18	18			18	18			18	17

Supplementary Figure 1

Plasma GMT$_{50}$ from post boosted vaccinations and COVID-19 sorted by study cohort with live virus assays on the left and pseudovirus on right, with individual sample minimum and maximum dilution titer also shown. A) WA-1; B) BQ.1.1; C) BA.2.75; D) BA.4/5.

Supplementary Figure 2

413 Plasma GMT$_{50}$ from boosted vaccinations only without COVID-19 sorted by study cohort with
414 live virus assays on the left and pseudovirus on right with with individual sample minimum and
415 maximum dilution titer also shown. A) WA-1; B) BQ.1.1; C) BA.2.75; and D) BA.4/5.

422 Supplementary Figure 3
423 Plasma GMT$_{50}$ from Omicron infection alone and also pre-boosted in 2021 or 2022 6 to 11 months
424 after last vaccine dose sampled sorted by study cohort with live virus assays on the left and
425 pseudovirus on right with minimum and maximum dilution titer also shown. A) WA-1; B) BQ.1.1;
426 C) BA.2.75; and D) BA.4/5.
427