Resilient CD8[+] T cells maintain a high cytotoxic capacity by balancing ROS via ME1 upregulation

Joanina K. Gicobi[1], Zhiming Mao[1], Grace DeFranco[1], Ying Li[3], Cindy Liu[2], Jacob B. Hirdler[2], Vianca V. Vianzon[1], Emilia R. Dellacecca[1], Michelle A. Hsu[1], Whitney Barham[1], Yohan Kim[2], Feven Abraha[4], William S. Harmsen[4], Yiyi Yan[5], Roxana S. Dronca[5], Mojun Zhu[6], Svetomir N. Markovic[6], Aaron S. Mansfield[6], Yi Lin[6], Xiaosheng Wu[6], Dawn Owen[7], Michael P. Grams[7], Jacob J. Orme[6], Fabrice Lucien[2], Hu Zeng[1, 8], Sean S. Park[7], Haidong Dong[1, 2, *]

[1] Department of Immunology, Mayo Clinic College of Medicine and Science, Rochester, MN
[2] Department of Urology, Mayo Clinic, Rochester, MN
[3] Department of Biomedical Statistics and Informatics, Mayo Clinic, Rochester, MN
[4] Department of Quantitative Health Sciences, Mayo Clinic, Rochester, MN
[5] Division of Medical Oncology, Mayo Clinic, Jacksonville, FL
[6] Division of Medical Oncology, Mayo Clinic, Rochester, MN
[7] Department of Radiation Oncology, Mayo Clinic, Rochester, MN
[8] Department of Rheumatology, Mayo Clinic, Rochester, MN

[*]Corresponding Author: dong.haidong@mayo.edu

Graphical abstract

Highlights

- CX3CR1[+] and low $\Delta\Psi$m identify functional resilient CD8[+] T cells.

- Resilient CD8[+] T cells are highly cytotoxic and have less ROS.

- Resilient CD8[+] T cells express more ME1 that can balance ROS.

- ME1 overexpression can promote cytotoxic function of CD8[+] T cells.

Abstract

Cytotoxic T lymphocytes (CTL) are indispensable in anti-tumor immunity. Although CTLs are prone to exhaustion in patients with advanced cancer, T cell resiliency explains the presence of tumor-reactive CTLs that are less exhausted, capable of cytolytic function, expand, and rebound in response to immunotherapy to reject metastatic malignances. However, the features of resilient T cells have not been clearly defined. In this report, we demonstrate peripheral CX3CR1[+] CD8[+] T cells with low mitochondrial membrane potential portray a functional resiliency in patients with large tumor burden as their CTL function rebounded quickly after radiation therapy. Furthermore, CX3CR1[+] CD8[+] T cell with low, but not high, mitochondrial membrane potential are highly cytotoxic, accumulate less reactive oxygen species (ROS) and express more Malic enzyme 1 (ME1). ME1 overexpression increases ATP production in a glycolysis-independent manner while concurrently curtailing excessive ROS in activated CD8[+] T cells; and expands CX3CR1[+]NKG7[+] effector CD8[+] T cells with enhanced cytotoxicity. Importantly, transfection of *ME1* mRNA promotes tumoricidal activity in CD8[+] T cells from patients with advanced cancers. Our study reveals a mechanism used by CTLs to balance excessive ROS via ME1 to maintain a metabolic and functional resiliency. Modification of ME1 expression in CTLs may be a novel method to improve the efficacy of cancer immunotherapy by preventing T cell exhaustion.

Introduction.

Immunotherapies (immune checkpoint inhibitors/ICI, vaccines, or T cell therapies) are dependent on the presence of endogenous cytotoxic T lymphocytes (CTLs) to respond to ICI or provide substrate for cell transfer therapy[1]. Moreover, radiation therapy has been used frequently to reduce the tumor burden and to create opportunity to synergize with immunotherapy [2]. To that end, the functionality of endogenous tumor-reactive CTLs plays a key role in predicting clinical response to combined radiation and immunotherapy[3,4].

Recently, we have proposed the term "resilient T cells" (Trs) to describe a functional state of tumor-reactive cytotoxic T cells that are capable of withstanding tumor burden and recovering quickly from stressful conditions such as harsh tumor microenvironment to respond to immunotherapy [5,6]. From there, a computational model of T cell resilience in multiple tumors has been established to predict whether a cancer immunotherapy will be effective in certain patients [7]. However, it is not clear whether Trs cells can be identified in circulation in patients with large tumor burden (5-10 cm or more in diameter). Of note, the identification and characterization of circulating Trs cells in these patients is pivotal because the presence of Trs cells is the base of combination immunotherapy to control an advanced cancer.

It has been reported that a successful ICI therapy expands tumor-reactive CD8[+] T cells with effector phenotype in peripheral blood, which have the potential to replace exhausted T cells inside tumor tissue [8-12]. To that end, we and others identified CX3CR1[+] CD8[+] T cells that are responsive to ICI therapy in both preclinical and clinical setting [11,13,14]. Because CX3CR1[+] CD8[+] T cells are highly cytotoxic, proliferative, and active in preclinical models and in the peripheral blood of patients who respond to ICI therapy [10,11,13], we hypothesize that CX3CR1[+] CD8[+] T cells are prototypes of Trs cells in circulation of patients with advanced cancers. A

further characterization of CX3CR1$^+$ CD8$^+$ T cells will provide insights into the biology of Trs cells in circulation.

In this report, we found that CX3CR1$^+$ CD8$^+$ T cells with low mitochondrial membrane potential ($\Delta\Psi$m) undergo a rapid cytotoxic response in patients with advanced cancer after radiation therapy. Upon further characterization of CD8$^+$ T cells with low $\Delta\Psi$m, we found that they are highly cytotoxic and accumulate less ROS (reactive oxygen species) but express more ME1(Malic enzyme 1) compared to T cells with high $\Delta\Psi$m. From there, we found ME1 acts as an anti-oxidant in CD8$^+$ T cells and expands CX3CR1$^+$ NKG7$^+$ effector CD8$^+$ T cells with enhanced cytotoxicity. Importantly, ME1 overexpression introduced by *ME1* mRNA also increases mitochondrial-linked ATP while curtailing excessive ROS, thus enhancing the tumoricidal activity in CD8$^+$ T cells from patients with advanced cancers. Our study suggests that resilient T cells balance excessive ROS upon antigen stimulation via upregulation of ME1 to maintain their metabolic fitness and retain a high cytotoxic capacity. Modification of ME1 expression in T cells may be a new method to avoid T cell exhaustion and improve the efficacy of cancer immunotherapy.

Results

CX3CR1 and low mitochondria membrane potential identify resilient CD8$^+$ T cells in patients with advanced cancers

To reduce the tumor burden in patients with very large (5-10 cm or more in diameter) or therapy-resistant tumors, patients were treated with a unique form of radiation therapy, spatially fractionated radiotherapy (SFRT) (**Fig. 1a**). SFRT is unique in that it intentionally delivers a heterogeneous dose distribution consisting of alternating regions of very high and low doses throughout the tumor[15]. SFRT can produce dramatic relief of severe symptoms, cause significant tumor regression, and achieve above average local control rates [16]. The biologic mechanisms responsible for the effectiveness of SFRT are still unknown, but the heterogeneous dose distribution is hypothesized to be responsible for modifying tumor-reactive immune response locally and systemically.

We and others previously identified circulating CX3CR1$^+$ CD8$^+$ T cells that are less exhausted and demonstrate high cytotoxic capability in patients with advanced cancers such as melanoma and lung cancers [10,11,13]. We hypothesized that a change in the functional state of CX3CR1$^+$ CD8$^+$ T cells may reflect an optimal response to a successful SFRT. Given the presence of large tumor burden in these patients in which compromised mitochondrial function has been linked to T cell exhaustion [17-22], we measured the CTL function of CX3CR1$^+$ CD8$^+$ T cells along with their mitochondrial membrane potential ($\Delta\Psi$m) to determine their mitochondria function. The CTL function of these T cells was measured by CD107a expression, which reflects degranulation and cytotoxicity. In lung cancer and sarcoma patients with large tumor burden, CX3CR1$^+$ CD8$^+$ T cells with low, but not high $\Delta\Psi$m showed improved CTL function one day after SFRT (**Fig. 1a-c, Extended Data Fig. 1a**). Similarly, one week after SFRT we found CD8$^+$ T cells with low $\Delta\Psi$m showed improved CTL function compared to cells with high $\Delta\Psi$m (**Fig. 1d**). In another cohort of patients with advanced melanoma that are resistant to ICI therapy, we found CX3CR1$^+$ CD8$^+$ T cells, but not CX3CR1$^-$ CD8$^+$ T cells, exhibit a low $\Delta\Psi$m phenotype in the peripheral

blood (**Fig. 1d**) prior to radiation therapy. We treated this cohort of patients with stereotactic body radiotherapy (SBRT) to reduce tumor burden. From there, we found patients with a higher frequency (>28%, cut-off line indicated in **Fig. 1e**) of CX3CR1$^+$ CD8$^+$ T cells with low $\Delta\Psi$m at baseline demonstrated superior overall survival (OS) in response to SBRT (**Fig. 1f**). On the other hand, we found the frequency of CX3CR1$^+$ CD8$^+$ T cells with low $\Delta\Psi$m gradually decreased in responders to ICI therapy as the tumor burden reduced but fluctuated in non-responders as the tumor burden persisted (**Extended Data Fig. 1b-c**). Taken together our clinical observations suggest that CX3CR1$^+$ CD8$^+$ T cells with low $\Delta\Psi$m are present in patients with large tumor burdens, and demonstrate a resiliency in their cytotoxic function after radiation therapy.

Resilient CD8$^+$ T cells are highly cytotoxic and less exhausted

Although murine CD8$^+$ T cells with low $\Delta\Psi$m demonstrates a higher antitumor activity [23], the functionality of human CD8$^+$ T cells with low or high $\Delta\Psi$m with respect to cytotoxicity has not been clearly defined. To understand whether the low $\Delta\Psi$m feature of CD8$^+$ T cells is associated with their functional state, we sorted CD8$^+$ T cells isolated from healthy donors into low or high $\Delta\Psi$m using TMRM staining as reported [23] (**Fig. 2a; Extended Data Fig. 2a**). First, we performed bulk RNA-seq analysis to compare the transcription differences between CD8$^+$ T cells with low and high $\Delta\Psi$m. We found CD8$^+$ T cells with low $\Delta\Psi$m expressed more genes that code for cytotoxic effector molecules such as *GZMB, PRF1*, and *NKG7* than CD8$^+$ T cells with high $\Delta\Psi$m (**Fig. 2b**). We then confirmed a higher expression of granzyme B in CD8$^+$ T cells with low $\Delta\Psi$m compared to CD8$^+$ T cells with high $\Delta\Psi$m using RT-PCR and flow cytometry (**Fig. 2c-d**). In a T cell-mediated tumor cytotoxicity assay following a brief T cell activation with anti-CD3/CD28, we found that CD8$^+$ T cells with low $\Delta\Psi$m demonstrated 1.5-fold higher cytolytic activity compared to CD8$^+$ T cells with high $\Delta\Psi$m in killing of two tumor cell lines (breast cancer and prostate cancer) (**Fig. 2e**). Also, we observed significantly increased degranulation in CD8$^+$ T cells with low $\Delta\Psi$m compared to CD8$^+$ T cells with high $\Delta\Psi$m upon T cell activation (**Fig. 2f**). In addition, gene set enrichment analysis (GSEA) showed CD8$^+$ T cells with low $\Delta\Psi$m are enriched for mediators of inflammatory responses including Interferon (IFN) and IL-2/STAT5 pathways (**Fig. 2g; Extended Data Fig. 2b**). To determine which phenotype of human CD8$^+$ T cells with low or high $\Delta\Psi$m would be, we found CD8$^+$ T cells with low $\Delta\Psi$m cells were enriched in both Temra-like (CCR7$^-$CD45RA$^+$) and Tnaive-like (CCR7$^+$CD45RA$^+$) subsets of CD8$^+$ T cells, indicating a limitation of using CCR7 and CD45RA to identify a functional state of resilient T cells (**Extended Data Fig. 2c**). Overall, our results suggest that human CD8$^+$ T cells with low $\Delta\Psi$m are highly cytotoxic and more responsive to inflammatory signals.

Given a high cytotoxic capability of CD8$^+$ T cells with low $\Delta\Psi$m, we considered whether they are prone to exhaustion upon activation. Interestingly, CD8$^+$ T cells with low $\Delta\Psi$m had lower expression of exhaustion markers such as TOX, a causative factor of T cell exhaustion[24] , and PD-1(**Fig. 3a, b**) compared to cells with high $\Delta\Psi$m. Interestingly, we found CD8$^+$ T cells with low $\Delta\Psi$m are enriched with TCF-1$^+$PD-1$^+$ stem-like cells that have been reported to be responsive to immunotherapy (**Fig. 3c**) [23]. To confirm this possibility, we compared the cytotoxicity of CD8$^+$ T cells with low or high $\Delta\Psi$m in co-culture with anti-PD-1 or anti-PD-L1 antibody. We found the cytotoxicity of CD8$^+$ T cells with low $\Delta\Psi$m increased in the presence of anti-PD-L1 or anti-PD-1 antibody compared to CD8$^+$ T cells with high $\Delta\Psi$m, although only the

anti-PD-L1 group reached a statistical significance (**Fig. 3d**). Additionally, a pathway analysis of our RNAseq data revealed upregulation of anti-PD-1/L1 cancer immunotherapy pathway in CD8[+] T cells with low ΔΨm cells compared to high ΔΨm cells (**Extended Data Fig. 3a**). Since transcriptional factors: Eomesodermin (Eomes) and T-bet regulate the expression of effector genes in CD8[+] T cells [25-28] and the ratio of Eomes/T-bet may determine the T cell exhaustion and immune pathology [29,30], we measured and compared Eomes and T-bet expression between CD8[+] T cells with low or high ΔΨm. Although Eomes and T-bet expression at mRNA level were significantly higher in CD8[+] T cells with low ΔΨm compared to high ΔΨm cells, the protein levels of Eomes and T-bet were comparable between CD8[+] T cells with low or high ΔΨm (**Fig. 3e-f**). These data suggest that although CD8[+] T cells with low ΔΨm seem to be programmed with a high capability of effector function that could lead to exhaustion (based on high Eomes transcription), their functional state may be regulated by other mechanisms to prevent them from entering exhaustion.

Resilient CD8[+] T cells have lower glycolysis and less ROS

As the levels of ΔΨm of CD8[+] T cells reflect the metabolism state of T cells[23,31], we compared the glycolysis (ECAR) and mitochondria respiration (OCR) between CD8[+] T cells with low ΔΨm or high ΔΨm. Effector CD8[+] T cells rapidly utilize glycolysis as a quick but inefficient means of ATP production to carry out their effector functions[32-34]. However, a recent report argues that activated human CD8[+] T cells increase their metabolic pathways in both glycolysis and oxidative phosphorylation [35]. We found CD8[+] T cells with low ΔΨm had lower glycolysis and glycolytic capacity than CD8[+] T cells with high ΔΨm (**Fig. 4a, b**), while their mitochondria respiration (OCR) was comparable between CD8[+] T cells with low ΔΨm or high ΔΨm (**Fig. 4c, d**). However, the OCR/ECAR ratio was higher in CD8[+] T cells with low ΔΨm compared to high ΔΨm cells (**Fig. 4f**), implying that the resilient CD8[+] T cells may utilize more mitochondrial respiration for energy needs instead of glycolysis. Accordingly, mitochondrial-linked ATP and spare respiratory capacity are comparable between CD8[+] T cells with low ΔΨm and high ΔΨm (**Fig. 4d, f**). Given the high cytotoxicity capacity of CD8[+] T cells with low ΔΨm, it is unexpected that they would exhibit low glycolysis. Since GLUT1 levels were comparable between CD8[+] T cells with low ΔΨm and high ΔΨm (**Fig. 4g**), it is possible that the lower glycolysis of CD8[+] T cells with low ΔΨm is not due to a lower intake of glucose but rather a preference for lower lactate production via glycolysis which could result into Warburg effects limiting their functionality[36]. Our data suggest that although CD8[+] T cells with low ΔΨm do not have increased glycolysis, but rather meet their energy requirements via mitochondrial-linked respiration.

In consistent with above observations, our GSEA analysis shows that reactive oxygen species (ROS) pathway was enriched in CD8[+] T cells with high ΔΨm compared to CD8[+] T cells with low ΔΨm (**Fig. 5a**). From there, we measured and compared ROS levels between CD8[+] T cells with low ΔΨm or high ΔΨm. We found both cytosolic ROS and mitochondrial ROS were lower in resting and activated CD8[+] T cells with low ΔΨm compared to CD8[+] T cells with high ΔΨm (**Fig. 5b-c**), and the lower levels of ROS were maintained up to 7 days during *in vitro* culture (**Fig. 5c**). To that end, we measured and compared the mass and numbers of mitochondria between CD8[+] T cells with low and high ΔΨm. Although there was a trend of lower mass and

numbers of mitochondria in CD8$^+$ T cells with low $\Delta\Psi$m (**Fig. 5d-e**), this difference did not reach statistical significance, suggesting other mechanism may contribute the lower levels of ROS beyond the volume of mitochondria.

Since excessive ROS signals result in DNA damage and functional exhaustion in T cells [22,37,38], the ability to maintain a lower ROS level in the context of a higher OCR/ECAR ratio could be a unique feature of resilient CD8$^+$ T cells with low $\Delta\Psi$m that enable them to tolerate harsh tumor microenvironment that would lead to increase ROS in T cells. To test whether this feature of CD8$^+$ T cells with low $\Delta\Psi$m could be retained in T cells in patients with large tumor burdens, we measured and compared the levels of ROS in CD8$^+$ T cells with low or high $\Delta\Psi$m isolated from patients with advanced lung cancers and sarcoma before and after SFRT. Interestingly, a lower level of ROS was consistently maintained in CD8$^+$ T cells with low $\Delta\Psi$m compared to cells with high $\Delta\Psi$m before and after SFRT (**Fig. 5f**, n=9). Similar findings were confirmed in another cohort of patients with advanced prostate cancer receiving SBRT (**Fig. 5f**, n=21). Taken together, our results suggest that CD8$^+$ T cells with low $\Delta\Psi$m selectively use mitochondria respiration to meet their energy needs but have a capacity of balancing ROS to avoid an accumulation of excessive ROS.

ME1 is upregulated in Resilient CD8$^+$ T cells

To determine how CD8$^+$ T cells with low $\Delta\Psi$m can maintain their ATP production via mitochondria respiration while keeping a lower ROS level, we performed RNA-seq analysis and found malic enzyme 1 (ME1) was among the most upregulated genes in the CD8$^+$ T cells with low $\Delta\Psi$m cells in both resting and activated state compared to CD8$^+$ T cells with high $\Delta\Psi$m (**Fig. 6a-b**). Since ME1 alone has the potential to regulate metabolism and ROS among these upregulated genes[39,40], we focused on ME1 and confirmed its higher expression in CD8$^+$ T cells with low $\Delta\Psi$m cells compared to high $\Delta\Psi$m cells via quantitative RT-PCR and Western blot in both resting and activated CD8$^+$ T cells (**Fig. 6c-d**).

To examine the consequence of ME1 expression in CD8$^+$ T cells with low or high $\Delta\Psi$m, we sorted CD8$^+$ T cells with low or high $\Delta\Psi$m from healthy donors followed with ME1 mRNA transfection and functional analysis (**Fig. 6e**). The overexpression of ME1 was confirmed by RT-PCR in CD8$^+$ T cells with low or high $\Delta\Psi$m, respectively (**Fig. 6f**). Interestingly, we found that ME1 overexpression reduced ROS levels only in CD8$^+$ T cells with high $\Delta\Psi$m but not in CD8$^+$ T cells with low $\Delta\Psi$m (**Fig. 6g**). Of note, although ME1 has the capability to reduce ROS levels in CD8$^+$ T cells, ME1 did not completely abolish ROS there, suggesting ME1 is a regulator of ROS rather than a scavenger of ROS in T cells. Since a basal level of ROS is also required for T cell activation[41], to determine whether ME1 overexpression would affect the functionality of CD8$^+$ T cells, we measured and compared the T cell-mediated cytotoxicity between CD8$^+$ T cells transfected with control or ME1 mRNA. The overexpression of ME1 in CD8$^+$ was confirmed via western blot (**Extended Data Fig. 4a-b**). Interestingly, we found ME1 overexpression significantly increased the cytotoxicity of CD8$^+$ T cells in killing tumor cells in vitro (**Fig. 6h**). Accordingly, we found ME1 overexpression significantly increased the frequency of highly cytotoxic CX3CR1$^+$NKG7$^+$ CD8$^+$ T cells [42] upon T cell activation (**Fig. 6i**). Our results suggest

that ME1 may not only regulate the level of ROS in CD8[+] T cells according to their mitochondria potential, but also enhance CTL function of CD8[+] T cells.

ME1 increases metabolic output of CD8[+] T cells without increasing ROS

To understand how ME1 enhance CTL function of CD8[+] T cells, we examined whether ME1 may regulate T cell metabolism upon activation, as ME1 is a cytosolic enzyme that converts malate into pyruvate and produces NAPDH[39,40]. Interestingly, we found CD8[+] T cells with ME1 overexpression had higher basal and maximal respiration as well as a higher ATP-linked respiration, but their spare respiratory capacity did not increase significantly (**Fig. 7a; Extended Data Fig. 4c**). On the other hand, we found ME1 overexpression did not affect the glycolysis and glycolytic capacity in CD8[+] T cells (**Fig. 7b**). Importantly, the increased mitochondria respiration induced by ME1 overexpression did not result in an elevated level of cytosolic ROS among total CD8[+] T cells at both activation and resting stage (**Fig. 7c; Extended Data Fig. 4d**).

To test whether ME1 may have a therapeutic potential in improving CTL function of PBMCs in patients with large tumor burdens, we introduced ME1 mRNA into PBMCs isolated from patients with advanced melanoma and prostate cancer. We found that ME1 overexpression increased the tumor killing ability of PBMCs in about 50% of patients with advanced melanoma or prostate cancer (**Fig. 7d**). From there, we found that ME1 overexpression reduced ROS only in CD8[+] T cells with low $\Delta\Psi$m but not in high $\Delta\Psi$m cells (**Fig. 7e**), suggesting CD8[+] T cells with low $\Delta\Psi$m in patients with large tumor burdens could be more sensitive to regulation mediated by ME1 overexpression. Taken together, our results imply that ME1 can promote CTL function via increasing the ATP production and mitochondrial activity in activated CD8[+] T cells to meet their energy requirements while concurrently curtailing excessive ROS to prevent T cell exhaustion (**Fig. 7f**).

Discussion

In this study, we characterized the features of resilient T cells in patients with advanced malignancies. We found CX3CR1[+] CD8[+] T cells with low $\Delta\Psi$m are endowed with functional resiliency and metabolic fitness that allow them to withstand large tumor burden, retain their cytotoxicity and be responsive to radiation therapy or ICI treatment. One potential underlying mechanism of this resiliency is the upregulation of ME1 that not only can balance excessive ROS in activated T cells but also can sufficiently enhance cytotoxic T cell response. Thus, our study reveals a mechanism by which a highly cytotoxic function can be maintained in resilient T cells in patients with large tumor burdens and provides a method to improve the efficacy of cancer immunotherapy and radiation therapy for patients with advanced cancer.

Since PD-1 signals dynamically influence mitochondrial functions resulting compromised function of effector T cells along with large tumor burdens[17-22], we measured the mitochondrial membrane potential ($\Delta\Psi$m) of CX3CR1[+] CD8[+] T cells to determine their mitochondria functional state as a start to understand T cell resilience. According to the level of $\Delta\Psi$m, we found CX3CR1[+] CD8[+] T cells with low $\Delta\Psi$m are more responsive to radiation therapy in patients with large tumor burdens. From there, we further characterized CD8[+] T cells with low

ΔΨm and found they are highly cytotoxic and produce less ROS. We found that ME1 (malic enzyme 1) is upregulated in CD8[+] T cells with low ΔΨm that can balance elevated levels of ROS in activated T cells and can promote tumoricidal activity of CD8[+] T cells in both healthy donors and patients with advanced cancers. To our knowledge, the role of ME1 in T cell function has not been clearly defined although ME1 have been identified as a key pro-tumor factor in cancer cells where ME1 helps tumor cells to survive in tumor microenvironments with high ROS [39,43]. ME1 is a NADP+ dependent cytosolic enzyme that is required for catalytic activity of a reaction that converts malate into pyruvate with NAPDH as a by-product[39,40]. This means that the catalytic function of ME1 in CD8[+] T cells with low ΔΨm may serve to provide pyruvate which feeds into either glycolytic or oxidative phosphorylation pathway for ATP production, and thus provide an optimal maintenance of metabolic fitness in T cells. Additionally, NADPH produced from the ME-1-mediated reaction can act as an anti-oxidant to prevent accumulation of excessive ROS, which may trigger T cell death [44] and T cell exhaustion [22,37,38]. Of note, although ME1 has the potential to reduce cytosolic ROS probably through NADPH, our data suggest ME1 is not a scavenger of cytosolic or mitochondrial ROS in T cells that needs a basal level of ROS for activation[41].

Large tumor burden has been significantly correlated with more T cells exhaustion and poorer response to ICI therapy [45,46]. However, the presence of resilient T cells that can recover their CTL function quickly after radiation therapy is the unexpected discovery in our study. Although the concept of resilient T cells in response to cancer immunotherapy has been conceived recently [5-7], no markers have been defined to identify them in patients with advanced diseases. To that end, our study suggest that CX3CR1[+] CD8[+] T cells with low ΔΨm could be a surrogate of resilient T cells, based on several features we identified in this report, such as (1) their CTL function (degranulation/CD107a expression) can be recovered quickly after tumor burden has been reduced by radiation therapy; (2) they maintain a low profile of glycolysis but produce optimal levels of ATP; (3) they are not prone to exhaustion due to lower expression of PD-1 and TOX; (4) they share a feature of stem-like T cells (TCF1[+]PD-1[+]) that can be responsive to ICI therapy[47-52], which is consistent with murine CD8[+] T cells with low ΔΨm [23]; (5) they maintain a relative lower level of ROS to avoid exhaustion but do not compromise T cell activation[41]; (6) they upregulate ME1 expression to balance excessive ROS and contribute to expansion of highly cytotoxic effector T cells. These unique features of resilient T cells not only can be used as T cell biomarker to predict or monitor patient responses to cancer immunotherapy or combination therapy, but also can be used to convey a resilient functional state to T cells in patients who may be lack of enough resilient T cells in responses to immunotherapy. To that end, our study suggests that introduction of ME1 mRNA into T cells could be used to improve the efficacy of cancer immunotherapy either in response to ICI therapy or in adoptive T cell therapy.

Several limitations in our studies should be addressed in future studies. Although our current studies are mainly focusing on the circulating CD8[+] T cells in patients with advanced diseases, it is warranted to investigate whether CD8[+] T cells within tumor tissues would share the features of resilient T cells. On the other hand, since there are no animal models to address T cell-specific role of ME1 in antitumor immunity, new animal models of ME1 are needed to test how a persistent or transient expression of ME1 in tumor-reactive T cells would contribute to either endogenous tumor control or in response to ICI therapy. Also, the efficacy and safety of transfer of ME1-modified T cells in treatment of established tumors need to be addressed in mouse tumor

models in vivo. Finally, one open question remains as why some CD8$^+$ T cells are endowed or maintain their low or high ΔΨm status. It is not clear to what degree the heterogeneity of ΔΨm within human circulating CD8$^+$ T cells would be related with a systemic antitumor immunity, as resilient T cells need to maintain a depolarized low ΔΨm to avoid the consequences associated with high ΔΨm such as increased ROS or exhaustion. By adapting to this unique feature, resilient T cells can protect the organism from metastatic cancer and chronic viral infections.

In summary, our study identified a mechanism by which resilient T cells can avoid exhaustion in patients with advanced cancers. Leveraging the knowledge of T cell resiliency may improve the efficacy of ICI therapy, CAR-T cell, or TCR-T cell therapy to control metastatic malignancies that need a robust systemic anti-tumor immunity.

FIGURES

Fig. 1: CX3CR1 and low ΔΨm identifies resilient CD8 T cells in patients with advanced cancers upon radiation therapy. a, A diagram of SFRT therapy and study design for **b-c**. **b,** Degranulation assay (measured by CD107a expression) of CX3CR1$^+$CD8$^+$ T cells with low or high ΔΨm. **c-d,** Summaries of % CD107a$^+$ cells among CX3CR1$^+$CD8$^+$ T cells with low or high ΔΨm from patients with advanced lung cancer and sarcoma (n=17) at baseline and 24 hrs post SFRT therapy (c), and one week post SFRT (d). Each line indicates one individual patient. **e,** Percentages of CX3CR1$^+$ and CX3CR1$^-$ CD8$^+$ T cells with low or high ΔΨm from patients with melanoma receiving SBRT at baseline. **f,** Survival probabilities at baseline prior to SBRT in patients (n=20) with advanced melanoma. Kaplan-Meier estimates were used to estimate survival curves and a univariate Cox model was used to determine the association of patients and disease variables with the risk of each outcome of interest. Statistical significance was determined by Student's Paired two-tailed t-test in **b-c**, and non-paired t-test in **d**, *P<0.05, **P<0.01, ****P<0.0001, ns: not significant.

Extended Data Fig. 1: Functionality and frequency of CX3CR1$^+$ CD8$^+$ T cells with low or high ΔΨm in response to radiation or ICI therapy. a, Degranulation analysis of CX3CR1$^+$ CD8$^+$ T cells with high ΔΨm from patients with advanced lung cancer and sarcoma (n=17) as measured by CD107a expression comparing baseline and 24 hrs post SFRT therapy. **b,** Frequency of CX3CR1$^+$CD8$^+$ T cells with low ΔΨm was measured in the peripheral blood of patients with advanced melanoma at baseline (BL) and follow up time points (N=42) in responders (**c**) and non-responders (**d**) to ICI therapy. Statistical significance was determined by Student's Paired two-tailed t-test in **a-b** *P<0.05, ns: not significant.

Fig. 2: Resilient CD8 T cells have high cytolytic capability. a, Experimental schematics. **b,** Heatmap of differentially expressed genes from bulk RNAseq of anti-CD3/CD28 activated CD8+ T cells sorted according to low and high ΔΨm (N=3). Framed area indicates genes code cytotoxic molecules**. c,** Relative GZMB mRNA expression in 5 donors. **d,** Expression of granzyme B protein in activated CD8$^+$ T cells with low or high ΔΨm as analyzed by flow cytometry for MFI (median fluorescent intensity) and percentage of positive cells. **e,** Cytotoxicity assay of CD8$^+$ T cells with low or high ΔΨm against tumor cells (MCF-7 breast and PC.3 prostate cancer) at a ratio of 1:20 (Tumor: Effector) in a 4-hour Calcein release assay (n=9-10). **f,** Degranulation assay of CX3CR1$^+$CD8$^+$ T cells with low or high ΔΨm as measured by CD107a expression. **g,** Hallmark Gene Set Enrichment Analysis (GSEA) of RNAseq data from three healthy donors using IPA analysis software. Statistical significance was determined by Student's paired (**f**) or non-paired (**c-e**) two-tailed t-test, *P<0.05, **P<0.01.

Extended Data Fig. 2: Sorting and phenotype analysis of CD8$^+$ T cells with low or high ΔΨm. a, Gating strategy that defines the low and high ΔΨm among CD8$^+$ T cells for sorting and analysis. **b,** Hallmark Gene Set Enrichment Analysis (GSEA) of RNAseq data from three healthy donors using IPA analysis software. **c, Phenotype of** CD8$^+$ T cell subsets as defined by CCR7 and CD45RA expression among low or high ΔΨm cells. Statistical significance was determined by Student's Paired two-tailed t-test in **c,** **P<0.01, ***P<0.001, ****P<0.0001.

Fig. 3: Resilient CD8 T cells are not prone to be exhausted. A-b, TOX (a) and PD-1 (b) expression by activated CD8$^+$ T cells with low and high ΔΨm as analyzed by flow cytometry in median fluorescent intensity (MFI) and frequency (%). **c,** Frequency of TCF-1$^+$ PD-1$^+$ among CD8$^+$ T cells with low or high ΔΨm at activated state. **d,** CD8$^+$ T cells with low or high ΔΨm were activated with anti-CD3/CD28 beads in the presence of anti-PD-1 (Pembro) or anti-PD-L1 (Atezo) for 72hrs followed with a cytotoxicity assay at a ratio of 1:20 (Tumor: Effector) against PC.3 tumor cells (n=7-13). **e-f,** Box plots of bulk RNA seq transcripts (**e**) and flow cytometry analysis (**f**) of Eomes and T-bet in CD8$^+$ T cells with low or high ΔΨm at activated state (n=6). Statistics significance was determined by Student's Paired (**a**) and non-paired (**b-d, f**) two-tailed t-test, *P<0.05, **P<0.01, ns: not significant.

Extended Data Fig. 3: PD-1/L1 pathway is upregulated in CD8$^+$ T cells with low ΔΨm cells. Hallmark Gene Set Enrichment Analysis (GSEA) of RNA-seq data from three healthy donors was further analyzed with IPA analysis software for pathway regulation. Star indicates the upregulation of PD-1/PD-L1 pathway in low ΔΨm cells.

Fig. 4: Resilient CD8$^+$ T cells have lower glycolytic capacity. a-b, Glycolysis, and glycolytic capacity of sorted CD8$^+$ T cells with low and high ΔΨm after activation with anti-CD3/CD28 beads as measured with extracellular acidification rate (ECAR) analysis and shown with a representative of Seahorse graph (**b,** n=6). **c,** Basal respiration, and spare respiratory capacity of sorted CD8$^+$ T cells with low and high ΔΨm after activation as measured with Mitochondrial Oxygen Consumption Rate (OCR) analysis. **d,** Representative graph of OCR as measured by MitoStress Test (n=6). **e,** OCR/ECAR ratio calculated from the basal respiration values(n=6). **f,** Mitochondrial ATP of activated CD8$^+$ T cells obtained from Mitochondrial Oxygen Consumption Rate (OCR) analysis(n=6). **g,** GLUT1 expression (MFI) in activated CD8$^+$ T cells with low and high ΔΨm (n=4). Statistical significance was determined by Student's paired two-tailed t-test; *P<0.05, ns: not significant.

Fig. 5: Resilient CD8 T cells have lower cytosolic and mitochondrial ROS. a, Hallmark GSEA analysis of ROS pathway genes from bulk RNA-seq data. **b,** Cytosolic ROS was measured by flow cytometry using CellROX Green and shown as MFI in both resting and activated CD8$^+$ T cells. **c,** Mitochondrial ROS was measured by flow cytometry using MitoSOX over one week in culture media. **d,** Mitochondrial mass was measured using MitoTracker Green with flow cytometry and shown as MFI. **e,** Number of mitochondria per cell was counted via Transmission electron microscopy (TEM) among 10-16 view fields. **f-g,** Cytosolic ROS in CD8$^+$ T cells with low or high ΔΨm were measured from the PBMCs isolated from patients with lung cancer or sarcoma receiving SFRT (**f,** n=9), and from patient patients with prostate cancer receiving SBRT (**g,** n=21). Statistical significance was determined with Student's Non-paired two-tailed t-test; **P<0.01, ***P<0.001, ****P<0.0001, ns: not significant.

Fig. 6: Resilient CD8 T cells express more ME1 to regulate ROS and promote CTL function. a, Volcano plots of bulk RNA-seq of CD8$^+$ T cells with low or high ΔΨm at resting and activated state (n=3) (ME1 in green circle). **b,** Box plots of ME1 expression analysis with bulk RNA-seq as in (**a**). **c,**

Relative ME1 mRNA expression measured with qRT-PCR. **d,** Western blot of ME1 expression as in (c) in two donors. **e,** Schematics of functional study of T cells with overexpression of ME1. **f,** ME1 overexpression measured by qRT-PCR in low and high $\Delta\Psi$m cells. **g,** ROS levels measured in CD8$^+$ T cells with low or high $\Delta\Psi$m isolated from healthy donors (n=3) after transfection with control or ME1 mRNA. **h,** Cytotoxicity assay of CD8$^+$ T cells isolated from healthy donors (n=6) after transfection with control or ME1 mRNA in killing tumor cells (MCF-7) at a ratio of 1:10 of tumor: effector cells. **i,** % CX3CR1$^+$ NKG7$^+$ among CD8$^+$ T cells after transfection with control or ME1 mRNA as measured by flow cytometry (n=4). Statistical significance was determined by Student's Paired two-tailed t-test; *P<0.05, **P<0.01, ns: not significant.

Fig. 7: ME1 increases glycolysis and oxidative phosphorylation, but not ROS, in CD8 T cells. a-b, Seahorse analysis of mitochondria respiration   and mitochondrial ATP production (a) and glycolysis (b) in CD8$^+$ T cells following transfection with control or ME1 mRNA. **c,** Cytosolic ROS in CD8$^+$ T cells following transfection with control or ME1 mRNA as measured by flow cytometry (MFI). **d,** Percent of increased cytotoxicity of PBMCs transfection with ME1 mRNA over PBMCs transfected with control mRNA to normalize different basal level of cytotoxicity of each patient. Cytotoxicity assay was performed at a ratio of 1:10 of tumor: effector cells. **e,** ROS levels (MFI) measured in CD8$^+$ T cells with low or high $\Delta\Psi$m among PBMCs of patients (n=5) with advanced prostate cancer after transfection with control or ME1 mRNA followed with T cell activation by anti-CD3 antibody for 48 hrs. **f,** Graphical illustration of the potential role of ME1 in resilient CD8$^+$ T cells. Statistical significance was determined by Student's Paired (a, b) or two-tailed t-test; non-parametric Mann-Whitney correction (d, e); *P<0.05, **P<0.01, ns: not significant.

Extended Data Fig. 4: ME1 expression and its regulation of metabolism in CD8$^+$ T cells. a-b, ME1 overexpression measured by Western blot (**a**) followed with quantification of ME1 proteins (**b**) in CD8$^+$ T cells after transfection with control or ME1 mRNA. **c,** Spare respiration capacity as measured via Seahorse assay. **d**, ROS MFI in resting CD8$^+$ T cells overexpressing ME1 vs. control mRNA. Statistical significance was determined by Student's unpaired (c, d) two-tailed t-test; non-parametric Mann-Whitney correction, ns: not significant.

METHODS

Specimen collection and T cells isolation

Peripheral blood Mononuclear Cells (PBMCs) were isolated from healthy donors or patients via centrifugation with lymphoprep (07851, StemCell Technologies) and SepMate conical tubes (85450, StemCell Technologies). CD8$^+$ T cells were then isolated using a magnet based CD8 T cell isolation kit (19053, StemCell Technologies) and used immediately for experiments. Some experiments used PBMCs (patient samples) stored in liquid nitrogen. After thawing, PBMCs would be incubated in CTL medium (RPMI 1640 complete medium, rhIL-2:10U/ml, rh IL-15:5ng/ml; rhIL-7:5ng/ml) at 37°C overnight for recovery before transfection.

Patient information. Peripheral blood for this study were collected after written consent was obtained from each participant. Clinical course, treatment information, and outcomes in patients treated with anti-PD-1/L1 therapy and radiation therapy at Mayo Clinic were retrospectively collected. Response to treatment was evaluated according to standard clinical practice guidelines

using RECIST [11]. Note that the response listed for anti-PD-1/PD-L1 therapy, (R = complete response, NR = progressive disease, SD= stable disease) was apparent (using RECIST) at the 12-week post-initiation of therapy time-point for all patients. For prostate cancer (receiving SFRT and SBRT), the response was evaluated based on the PSA levels and distance reoccurrence.

Study approval. Peripheral blood for this study from healthy people were acquired from anonymous donors who had consented for blood donation at the Blood Transfusion Center at Mayo Clinic. All donors and patients provided signed informed written consent; the study was approved by the Mayo Clinic Rochester IRB and was conducted according to Declaration of Helsinki principles.

TMRM Staining and Cell Sorting

$CD8^+$ T cells were washed once with 1x PBS and adjusted to a concentration of $1x10^6$ cells/ml and stained with 0.02μM final concentration of tetramethylrhodamine methyl ester (TMRM) or 2nM final concentration of Carbonyl Cyanide 3-chlorophenylhydrazone (CCCP) as control. The TMRM stained cells were incubated at 37°C for 30 minutes with intermittent shaking while the CCCP stained cells were incubated at 37°C for 5 minutes. The cells were then washed twice with 1x PBS and resuspended at 10-15 $x10^6$cells/ml of cell culture media for sorting. The cells were sorted with BD FACSMelody™ Cell Sorter using the 100-micron sort nozzle and on the PE channel (561nm excitation).

siRNA Transfection

Sorted CD8+ T cells with low and high ΔΨm were centrifuged at 200g for 10 min prior to nucleofection (4D nucleofector system, Lonza). Three- five million cells were combined with 200 pMol siRNA (siControl or siME1) in 20 μl P3 nucleofection media (Lonza) per well of the 16-well Nucleocuvette strips (X unit). Program FI-115 was used. Following nucleofection, cells were rested in warm RPMI (no FBS, no cytokines) for 4 hours before adding 10% FBS and 10IU/ml of IL-2, 5ng/ml of IL-7, and 5ng/ml of IL-15 into the culture for an overnight recovery followed with use in experiments.

mRNA Transfection

T cells were transfected with control mRNA or ME1 mRNA (220 μg/ml) using the P3 nucleofection kit (Lonza V4XP-3024) and program FI-115 on the 4D Nucleofector (Lonza). Following nucleofection, cells were rested in warm RPMI (no FBS, no cytokines) for 4 hours before adding 2x CTL medium (10% FBS and IL-2, IL-7, IL-15) for an overnight recovery followed with T cell activation and functional analysis.

Cytotoxicity Assay

Target tumor cells were first washed twice with HBSS, and then labeled with Calcein-AM (5 μM) for 30 min and incubation at 37°C in dark, shaking occasionally. The cells were then washed with HBSS twice and re-suspended at $1x10^5$/ml in CTL medium with no FBS. Pre-

activated CD8[+] T cells and target cells were mixed at 1:20 or 1:10 (target to effector ratio) in CTL media without FBS and seeded into 96-U bottom well plate at 200µl per well. Saponin (0.1%) or Triton X-100 (2%) was added to wells that contained tumor cells only to provide a value for "maximum calcein release," in each assay; tumor cell-only wells were included to measure spontaneous release of calcein. All experimental and control conditions were performed in triplicate wells. The plate was briefly centrifuged at 1000rpm for 30 sec, followed by incubated at 37°C for 4 hours. After 4hrs incubation, the plate was centrifuged at 2000 rpm for 5 min. 100µl of the 200µl supernatant was then removed from each well and added to a new, opaque (black) 96 well flat bottom plate (Thermo scientific, 237105). Calcein fluorescence was read using an automated fluorescence measurement system (BioTeK Synergy HTX multi-mode reader) with an excitation of 485/20 and an emission filter of 530/25 scanning for 1 sec per well. Percent cytotoxicity was then calculated using the following formula:
 % Cytotoxicity = 100 x [(Experimental well release-Spontaneous release well)/Maximum release well-Spontaneous release well].

Degranulation assay

T cells were adjusted at a concentration of $1x10^6$ cells/100µl CTL medium including Golgi-Stop (Biolegend, 420701) and Golgi-Plug (Biolegend, 420601) and incubated with 5µl of CD107a antibody (Biolegend, H4A3) and 5 µl of anti-CD3/CD28 beads. The cells were then briefly centrifuged at 300g for 1 minute and incubated at 37°C for 5 hours. After the incubation, the cells were stained with TMRM followed with surface antibody staining before flow cytometry analysis.

ROS detection

Cells were stained with 250 nM of CellROX (ThermoFisher Scientific, C10492 or 1 µM of MitoSOX (ThermoFisher Scientific, M36008) in complete media and incubated at 37°C for 45 minutes. From there, the cells were resuspended in FACS buffer (1x PBS, 2mM EDTA, and 3% FBS) at a concentration of $1 x 10^6$ cells/ 100µl followed by staining with antibodies for surface molecules for 20 minutes at room temperature in the dark. Cells were then washed once in FACS buffer, resuspended in 200µl, and analyzed on Bio-Rad ZE5 Cell Analyzer. Analysis was performed using FlowJo V10.

Flow Cytometry Analysis

Cells were adjusted to 0.5-1x 10^6 cells/ml with 1x PBS and stained with live/dead dye and incubated at 4°C for 30 minutes. The cells were then washed once with 1x PBS followed by staining for cell surface molecules. For intracellular molecule staining, cells were incubated with FoxP3 Fixation Buffer overnight at 4°C. After fixation, cells were washed once 1x permeabilization buffer, and then stained with antibodies for intracellular molecules such as ME1, TCF-1/7, TOX, Eomes, T-bet, Ki67, HIF1-α, NKG7, and Glut1 for 1 hour at 4°C in 100µl of 1x permeabilization buffer. Following intracellular staining, cells were washed once with 1x permeabilization buffer and then resuspended in 200µl FACS buffer for flow cytometry analysis, which was completed on a Cytoflex LX (Beckman Coulter) (DAQ Version V2.233, MCB

Version: V3.01) running CytExpert software. Flow cytometric analysis was performed using FlowJo V10.

Quantitative PCR

RNA was isolated using Qiagen RNeasy Plus Mini Kit (74134, Qiagen). RT-reaction was completed using SuperScript™ III Reverse Transcriptase (18080-400, Invitrogen). qRT-PCR analysis was completed using a Quant-Studio 3 Real-Time PCR System (Applied Biosystems) and TaqMan Fast Advanced Master Mix (444455, Applied Biosystems). Taqman assays included: Hs01120688-gl, Human ME1-FAM and Hs99999901-Sl, Human-18s-FAM, both from Applied Biosystems. All samples were run in doublet. Relative expression was calculated using the $2^{-\Delta Ct}$ method. Granzyme B (F 5-TACCATTGAGTTGTGCGTGGG-3, R 5-GCCATTGTTTCGTCCATAGGAGA-3, ME1 (F 5-GGGAGACCTTGGCTGTAATGG-3, R 5-TTCGGTTCCCACATCCAGAAT-3), UBE (F 5-GTACTCTTGTCCATCTGTTCTCTG-3; R 5-CCATTCCCGAGCTATTCTGTT-3). RNA input was normalized to 20ng/ul with 10ul input for the RT reaction. Post RT reaction cDNA was diluted 1:5 before amplification on the Quantstudio 3 in the following volume per well: 5µl cDNA template, 10µl SYBR green (Applied Biosystems), 3µl H_2O, 1µl 10 mM F/R primer.

Western Blots

Cell pellets were lysed in NP-40 buffer and concentrations measured via protein assay using BioRad reagent (#500-0006). Once diluted to equal concentrations, samples were combined with Laemmli sample buffer (BioRad) and run using a Mini-PROTEAN Electrophoresis and Transfer system (BioRad). Membranes were incubated with primary antibody overnight at 4°C (ME1, Beta-Actin, or GAPDH). Secondary antibodies were added the next morning for 2 hours (HRP-conjugated anti-mouse). Signal was visualized using SuperSignal West Pico PLUS Chemiluminescence Substrate Kit (34577, Thermo Fisher) on a Syngene G: Box Chemi XX6 system running GeneSys software (V1.6.1.0). Full images of all membranes are included in the supplementary source data files.

Transmission electron microscopy (TEM) analysis of mitochondria

Cells were fixed in Trump fixative for 1 hour at room temperature or at 4ºC overnight followed with a fixation for 1 hr in 1% osmium tetroxide. The samples were dehydrated, embedded in Spurrs resin, sectioned at 90 nm, and observed using a Joel 1400 electron microscope (Joel USA Inc.). For quantification, images of individual T cell in a single field of view downloaded into JPEG images and the number of mitochondria structures within the T cells were counted manually by two different readers.

Metabolic Assays

Seahorse Xfe96 Bioanalyser (Agilent) was used to determine OCR and ECAR. Sorted cells or total CD8$^+$ T cells were washed in XF Base media (Seahorse XF RPMI medium with 2 mM glutamine, 10 mM glucose, 1 mM sodium pyruvate, and 5 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid), pH 7.4 at 37 °C) for OCR or XF media with 2mM glutamine for

ECAR before being plated onto Seahorse cell culture plates coated with Cell-Tak (Corning #354240) at 1×10^5 cells per well. The cells were allowed to adhere to culture plate. The OCR was measured via Seahorse Mito Stress assay (Agilent), with addition of oligomycin (2 µM), carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone (FCCP; 1.2 µM) and Rotenone and Antimycin (1.0 µM)). The ECAR was measured with addition of 10mM glucose, 2 µM oligomycin, and 50 nM 2-deoxy-D-glucose (2-DG). Assay parameters were as follows: 3 min mix, no wait, 3 min measurement, repeated 3–4 times at basal and after each addition. SRC was calculated as oxygen consumption rate (OCR) at maximum rate (OCR_{Max}) − OCR in basal state (OCR_{Bas}). Mitochondrial ATP production was calculated by subtracting the minimum respiration rate after oligomycin injection from the basal respiration rate before oligomycin injection.

Mitochondria mass Staining

Cells at a concentration of 1×10^6 cell/ml were washed once with 1xPBS and incubated with 100 nM of MitoTracker Green FM dye for 30 minutes at 37 °C. The cells were then analyzed by flow cytometry.

Bulk RNA sequencing

RNA was isolated using Qiagen Rneasy Plus Mini Kit (74134, Qiagen). The bulk RNA paired-end sequencing reads were processed through the Mayo bioinformatics pipelines MAP-Rseq (v3.0) as we previously reported. Reads were aligned to human reference genomes (hg38). The RNA aligned reads were quantified for gene expression using the Subread package. Differences across groups were assessed using bioinformatics package edgeR 2.6.2 to identify differentially expressed genes. Such genes were reported with magnitude of change (log2 scale) and their level of significance (False Discovery Rate, FDR < 5%). For pathway analysis, T cells with low or high ΔΨm were randomized and put through Gene Set Enrichment Analysis (GSEA) as described in the user guide.

Statistical Analyses

Data were analyzed in GraphPad Prism (version 9) using the unpaired or paired, two-tailed *t*-test without correction for multiple comparisons, as indicated in figure legends.

Antibody List

Target	Host	Clone	Flour	Company	Catalogue #	Final conc./dilution
Human CX3CR1	Rat	2A9-1	APC-Cy7	Biolegend	341616	Flow @1:20
Human CX3CR1	Rat	2A9-1	BV-510	Biolegend	341622	Flow @1:20
Human Granzyme B	Mouse	CLB-GB11	PERCP	Novus	NBP1-50071PCP	Flow @1:100
Human PD1	Mouse	EH12.2H7	APC	Biolegend	329908	Flow @1:20

Human PD-1	Mouse	EH12.2H7	APC-Cy7	Biolegend	329922	Flow @1:20
Human PD-1	Mouse	EH12.1	BV510	BD Horizon	563076	Flow @1:20
Human TCF-1	Rabbit	C63D9	Pac-Blue	Cell Signaling	#9066	Flow @1:50
Human TCF-1	Rabbit	C63D9	PE	Cell Signaling	#14456	Flow @1:50
Human EOMES	Mouse	WD1928	FITC	eBioscience by ThermoFisher	11-4877-42	Flow @1:20
Human TOX	Recombinant	REA473	PE	Miltenyi Biotec	130-120-716	Flow @1:50
Human Tbet	Mouse	4B10	APC	Biolegend	644814	Flow @1:20
Human HIF-1α	Mouse	Mgc3	APC	eBioscience by ThermoFisher	17-7528-82	Flow @1:20
Human GLUT1	Mouse	202915	FITC	R&D Systems	FAB1418F	Flow @1:20
Human CCR7	Mouse	2-L1-A	BV 650	BD Horizon	566756	Flow @1:20
Human CCR7	Mouse	G043H7	FITC	Biolegend	353216	Flow @1:20
Human CD45RA	Mouse	HI100	BV 510	BD Horizon	563031	Flow @1:20
Human CD45RA	Mouse	HI100	APC	eBioscience by ThermoFisher	17-0458-42	Flow @1:20
Human CD45RA	Mouse	5H9	BUV496	Biolegend	741182	Flow @1:20
Human Ki67	Mouse	Ki-67	APC	Biolegend	350514	Flow @1:20
Human CD8	Mouse	RPA-T8	PE-Cy7	BD Horizon	557746	Flow @1:20
Human CD8	Mouse	SK1	BV 510	Biolegend	344732	Flow @1:20
Human CD8	Mouse	SK1	APC	Biolegend	344722	Flow @1:20
Human CD3	Mouse	UCHT1	BV 650	Biolegend	300468	Flow @1:20
Human CD3	Mouse	UCHT1	PECy5.5	Biolegend	300410	Flow @1:20
Human CD3	Mouse	OKT3	BV 421	Biolegend	317344	Flow @1:20
Human CD11a	Mouse	TS2/4	PerCP/Cy5.5	Biolegend	350614	Flow @1:20
Human CD11a	Mouse	HI111		Biolegend	301220	Flow @1:20
Human CD11a	Mouse	HI111	PE-Cy7 APC-Cy7	Biolegend	301236	Flow @1:20
Human ME1	Mouse		None	Santa Cruz Biotechnology	sc-365891	Western Blot @ 1:1000
Human ME1	Mouse		PE		sc-365891	Flow @1:50
Human CD107a	Mouse	H4A3	FITC	Biolegend	328606	Flow @1:20
Brefeldin A				Biolegend	420601	
Monensin				Biolegend	420701	
Mouse IgG	Mouse	Polyclonal	None			
Rabbit IgG	Rabbit	Polyclonal	None			

| Peroxidase affiniPure Donkey anti-mouse IgG (H+L) | | Polyclonal | | Jackson ImmunoResearch | 715-035-151 | Western blot @ 1:5000 |
| Beta-actin | Rabbit | | None | Cell Signaling | #4970S | Western blot @ 1:2000 |

Dyes

Reagent (Fluoro)	Supplier	Catalogue #	Final conc.
Ghost Dye UV 450 (UV405)	Tonbo biosciences	13-0868-T100	Flow @1:100
Ghost Dye V 450 (BV450)	Tonbo biosciences	13-0863-T100	Flow @1:100
CellROX Green Flow Kit	ThermoFisher Scientific	C10492	Flow @500nM/mL
MitoSOX Red	ThermoFisher Scientific	M36008	Flow @100nM/mL
MitoTracker Green FM	ThermoFisher Scientific	M7514	Flow @100nM/mL
CellTrace™ CFSE dye	ThermoFisher Scientific	C34554	Flow @ 2.5μM/mL
TMRM, perchlorate	Biotium	70017	Flow @0.02μM
TMRM	ThermoFisher Scientific	M20036	Flow @0.02μM

CD8+ isolation and nucleofection reagents

Reagent	Supplier	Catalogue #
Lymphoprep	StemCell Technologies	07851
SepMate Conical tubes	StemCell Technologies	85450
EasySep Human CD8$^+$ T cell Isolation Kit	StemCell Technologies	Q-263176
P3 Primary Cell 4D-Nucleofector X Kit S	Lonza	V4XP-3032
P3 Primary Cell 4D-Nucleofector X Kit L	Lonza	V4XP-3024

qPCR reagents

Reagent	Supplier	Catalogue #
RLT buffer	Qiagen	79216
Qiagen RNeasy Plus Mini Kit	Qiagen	74134
SuperScript™ III Reverse Transcriptase	ThermoFisher Scientific	18080-400
TaqMan Fast Advanced Master Mix	Applied Biosystems	444455

Seahorse reagents

Reagent	Supplier	Catalogue #
Seahorse XF Cell Mito Stress Test Kit	Agilent	103015-100
Seahorse XF Glycolysis Stress Test Kit	Agilent	103020-100
Seahorse XFe96 FluxPak mini	Agilent	102601-100
Seahorse XF Cell Energy Phenotype Test Kit	Agilent	103325-100

Seahorse XF RPMI medium pH 7.4, 500mL	Agilent	103576-100
Seahorse XF 100mM pyruvate solution, 50mL	Agilent	103578-100
Seahorse XF 200mM glutamine solution, 50mL	Agilent	103579-100
Seahorse XF 1.0 M glucose solution, 50mL	Agilent	103577-100
Corning Cell-Tak Cell and Tissue Adhesive, 1 mg	Corning	354240

Inhibitors

Reagent	Supplier	Catalogue #	Final conc.
Oligomycin (OCR)	Agilent	103015-100	2.0μM
FCCP	Agilent	103015-100	1.5μM
Rot/AA	Agilent	103015-100	1.0μM
Glucose	Agilent	103020-100	10mM
Oligomycin (ECAR)	Agilent	103020-100	2.0μM
2-DG	Agilent	103020-100	50nM

Primer List

Primer	Sequence	Usage Notes
Human Full Length ME1 F	GGGAGACCTTGGCTGTAATGG	
Human Full Length ME1 R	TTCGGTTCCCACATCCAGAAT	
Human Full Length GZMB F	TACCATTGAGTTGTGCGTGGG	
Human Full Length GZMB R	GCCATTGTTTCGTCCATAGGAGA	
Human Full Length UBE F	GTACTCTTGTCCATCTGTTCTCTG	
Human Full Length UBE R	CCATTCCCGAGCTATTCTGTT	

KD and mRNA Overexpression reagents:

Reagent	Company/Catalogue or sequence	Catalogue #	Usage Notes
ME1 siRNA SMARTpool: ON-TARGETplus	Dharmacon	L-009348-00-0020	
Control siRNA ON-TARGETplus Non-targeting pool	Dharmacon	D-001810-10-20	
ME1 mRNA for human studies	TriLink Biotechnologies	L-7007	
Control mRNA	TriLink Biotechnologies	L-7610	

Cell culture reagents:

Reagent	Company	Catalogue #	Final conc/Usage Notes
DMEM	Gibco	11885-084	Base media for MCF-7
RPMI	Corning	10-040-CV	Base media for PC.3
HEPES buffer	Corning	25-060-CI	Added 10 mL of the 1M solution to 500 mL of all cell culture media
FBS	Gibco	10437-028	Final concentration of 10% FBS used for all cell culture media
FoxP3/TF Fixation/Permeabilization concentrate and diluent	eBioscience by ThermoFisher	00-5521-00	
Pen/Strep	Cellgro	30-002-CI	Used 5mL of the 100X solution in 500 mL of all cell culture media
Recombinant human IL-2	Peprotech	200-02	10U/ml unless otherwise stated in methods.
Recombinant human IL-15	Peprotech	200-15	10ng/mL
Recombinant human IL-7	Peprotech	200-07	10ng/mL
Purified NA/LE Mouse Anti-human CD3 (Clone HIT3a)	BD Horizon	555336	
Purified NA/LE Mouse anti-human CD28 (Clone CD28.2)	BD Horizon	555725	

Acknowledgements
We thank all the patients who generously contributed samples and participated in
the study. We thank all members of the Dong laboratory for helpful discussions and critical
analysis of the manuscript. We also acknowledge the Department of Radiation Oncology and
Division of Medical Oncology at the Mayo Clinic for providing excellent patient care. This study
was supported in part by the generosity of Mayo Clinic benefactors. We acknowledge the
Genome Analysis Core at Mayo Clinic, including technologists Vernadette A. Simon and
Fariborz Rakhshan-Rohakhtar as well as supervisors Julie S. Lau, Samantha J. McDonough, and
Mark Mutawe. We are grateful for the electron microscopy imaging service provided by Bing
Huang and Scott I. Gamb; and Flow cytometry and Sorting services provided by Holly Lamb,
Colleen Moe, Han Yong Hwan, and Drew Kluge at the Mayo Clinic Microscope and Cell
Analysis Core. The staff at Mayo Clinic Blood transfusion center. BioRender was used to create

the graphics throughout the manuscript. We acknowledge support from the Mayo Clinic Center for Individualized Medicine's IMPRESS program and High-Definition Therapeutics program (H. Dong), the Mayo Clinic Center for Biomedical Discovery (H. Dong), the Mayo Clinic Cancer Center's David F. and Margaret T. Grohne Cancer Immunology and Immunotherapy program (H. Dong), NIH grant R21 CA251923 (A.S. Mansfield), NIH grant K12 CA 090628 (Y. Yan), NIH grant R01 AR077518 (Hu Zeng), NIH grant R21 CA197878 (H. Dong), NIH grant R01 AI095239 (H. Dong), NIH grant R01 CA200551 (H. Dong and S.S. Park), NIH grant R01 CA256927 (H. Dong), and NIH grant R01 200551 (S.S. Park and H. Dong). This project is partially funded by the Lawrence and Marilyn Matteson Award S.S. Park, D.O, and M.P. Grams), and Sydney Luckman Family Predoctoral Award (J.K.G.).

Author contributions

J.K.G. and H.D. conceived the project, designed experiments, and wrote the manuscript. J.K.G. performed most of the experiments. Z.M., C.L., J.B.H., V.V.V., E.R.D., M.A.H., W.B., and Y.H. processed patient blood samples, conducted some of the flow cytometry experiments, Western blot experiments and qRT-PCR. Y.L (Li). analyzed the bulk RNAseq data and J.K.G. ran the GSEA analysis. G.D. performed healthy donors' ME1 overexpression cytotoxicity assays and Western blots. F.A., S.H., and J.J.O. performed statistical analysis. F.L. and H.Z. advised on biochemical and metabolic assays and provided critical edits to the manuscript. X.W. provided guidance on messenger RNA design and construct. Y.Y., R.X.D., M.Z., S.N.M., A.S.M., Y.L.(Lin)., M. G., D.O. and S.S.P. provided clinical samples and interpretation of clinical data. All authors reviewed the manuscript.

References

1. Walsh, S.R.*, et al.* Endogenous T cells prevent tumor immune escape following adoptive T cell therapy. *The Journal of clinical investigation* **129**, 5400-5410 (2019).
2. Evans, J.D.*, et al.* Prospective Immunophenotyping of CD8(+) T Cells and Associated Clinical Outcomes of Patients With Oligometastatic Prostate Cancer Treated With Metastasis-Directed SBRT. *Int J Radiat Oncol Biol Phys* **103**, 229-240 (2019).
3. Zhang, H.*, et al.* Phase II Evaluation of Stereotactic Ablative Radiotherapy (SABR) and Immunity in (11)C-Choline-PET/CT-Identified Oligometastatic Castration-Resistant Prostate Cancer. *Clin Cancer Res* **27**, 6376-6383 (2021).
4. Yan, Y.*, et al.* CX3CR1 identifies PD-1 therapy-responsive CD8+ T cells that withstand chemotherapy during cancer chemoimmunotherapy. *JCI Insight* **3**(2018).
5. Gicobi, J.K., Barham, W. & Dong, H. Immune resilience in response to cancer therapy. *Cancer Immunology, Immunotherapy* **69**, 2165-2167 (2020).
6. Gicobi, J.K., Dellacecca, E.R. & Dong, H. Resilient T-cell responses in patients with advanced cancers. *Int J Hematol* (2022).
7. Zhang, Y.*, et al.* A T cell resilience model associated with response to immunotherapy in multiple tumor types. *Nat Med*, 1421–1431 (2022).
8. An, H.J., Chon, H.J. & Kim, C. Peripheral Blood-Based Biomarkers for Immune Checkpoint Inhibitors. *Int J Mol Sci* **22**, 9414 (2021).
9. Sade-Feldman, M.*, et al.* Defining T Cell States Associated with Response to Checkpoint Immunotherapy in Melanoma. *Cell* **175**, 998-1013.e1020 (2018).

10. Wu, T.D.*, et al.* Peripheral T cell expansion predicts tumour infiltration and clinical response. *Nature* **579**, 274-278 (2020).
11. Yan, Y.*, et al.* CX3CR1 identifies PD-1 therapy-responsive CD8+ T cells that withstand chemotherapy during cancer chemoimmunotherapy. *JCI Insight* **3**, e97828 (2018).
12. Yost, K.E.*, et al.* Clonal replacement of tumor-specific T cells following PD-1 blockade. *Nat Med* **25**, 1251-1259 (2019).
13. Yamauchi, T.*, et al.* T-cell CX3CR1 expression as a dynamic blood-based biomarker of response to immune checkpoint inhibitors. *Nat Commun* **12**, 1402 (2021).
14. Zander, R.*, et al.* CD4(+) T Cell Help Is Required for the Formation of a Cytolytic CD8(+) T Cell Subset that Protects against Chronic Infection and Cancer. *Immunity* **51**, 1028-1042.e1024 (2019).
15. Griffin, R.J.*, et al.* Understanding High-Dose, Ultra-High Dose Rate, and Spatially Fractionated Radiation Therapy. *Int J Radiat Oncol Biol Phys* **107**, 766-778 (2020).
16. Grams, M.P.*, et al.* VMAT Grid Therapy: A Widely Applicable Planning Approach. *Pract Radiat Oncol* **11**, e339-e347 (2021).
17. Li, C.*, et al.* A high OXPHOS CD8 T cell subset is predictive of immunotherapy resistance in melanoma patients. *J Exp Med* **219**, e20202084 (2022).
18. Li, C.*, et al.* The Transcription Factor Bhlhe40 Programs Mitochondrial Regulation of Resident CD8(+) T Cell Fitness and Functionality. *Immunity* **51**, 491-507.e497 (2019).
19. Nishida, M.*, et al.* Mitochondrial reactive oxygen species trigger metformin-dependent antitumor immunity via activation of Nrf2/mTORC1/p62 axis in tumor-infiltrating CD8T lymphocytes. *J Immunother Cancer* **9**, e002954 (2021).
20. Ogando, J.*, et al.* PD-1 signaling affects cristae morphology and leads to mitochondrial dysfunction in human CD8(+) T lymphocytes. *J Immunother Cancer* **7**, 151 (2019).
21. Simula, L.*, et al.* PD-1-induced T cell exhaustion is controlled by a Drp1-dependent mechanism. *Mol Oncol* **16**, 188-205 (2022).
22. Yu, Y.-R.*, et al.* Disturbed mitochondrial dynamics in CD8+ TILs reinforce T cell exhaustion. *Nature Immunology* **21**, 1540-1551 (2020).
23. Sukumar, M.*, et al.* Mitochondrial Membrane Potential Identifies Cells with Enhanced Stemness for Cellular Therapy. *Cell Metab* **23**, 63-76 (2016).
24. Khan, O.*, et al.* TOX transcriptionally and epigenetically programs CD8(+) T cell exhaustion. *Nature* **571**, 211-218 (2019).
25. Buggert, M.*, et al.* T-bet and Eomes are differentially linked to the exhausted phenotype of CD8+ T cells in HIV infection. *PLoS Pathog* **10**, e1004251 (2014).
26. Intlekofer, A.M.*, et al.* Effector and memory CD8+ T cell fate coupled by T-bet and eomesodermin. *Nat Immunol* **6**, 1236-1244 (2005).
27. Pearce, E.L.*, et al.* Control of effector CD8+ T cell function by the transcription factor Eomesodermin. *Science* **302**, 1041-1043 (2003).
28. Szabo, S.J.*, et al.* A novel transcription factor, T-bet, directs Th1 lineage commitment. *Cell* **100**, 655-669 (2000).
29. Llaó-Cid, L.*, et al.* EOMES is essential for antitumor activity of CD8(+) T cells in chronic lymphocytic leukemia. *Leukemia*, 3152–3162 (2021).
30. Raveney, B.J.E.*, et al.* Involvement of cytotoxic Eomes-expressing CD4[+] T cells in secondary progressive multiple sclerosis. *Proceedings of the National Academy of Sciences* **118**, e2021818118 (2021).

31.   Amitrano, A.M.*, et al.* Optical Control of CD8(+) T Cell Metabolism and Effector Functions. *Front Immunol* **12**, 666231 (2021).

32.   Maciolek, J.A., Pasternak, J.A. & Wilson, H.L. Metabolism of activated T lymphocytes. *Curr Opin Immunol* **27**, 60-74 (2014).

33.   Newsholme, E.A., Crabtree, B. & Ardawi, M.S. The role of high rates of glycolysis and glutamine utilization in rapidly dividing cells. *Biosci Rep* **5**, 393-400 (1985).

34.   Finlay, D. & Cantrell, D.A. Metabolism, migration and memory in cytotoxic T cells. *Nat Rev Immunol* **11**, 109-117 (2011).

35.   Ma, E.H.*, et al.* Metabolic Profiling Using Stable Isotope Tracing Reveals Distinct Patterns of Glucose Utilization by Physiologically Activated CD8(+) T Cells. *Immunity* **51**, 856-870 e855 (2019).

36.   Vander Heiden, M.G., Cantley, L.C. & Thompson, C.B. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* **324**, 1029-1033 (2009).

37.   Dan, L.*, et al.* The phosphatase PAC1 acts as a T cell suppressor and attenuates host antitumor immunity. *Nature Immunology* **21**, 287-297 (2020).

38.   Scharping, N.E.*, et al.* Mitochondrial stress induced by continuous stimulation under hypoxia rapidly drives T cell exhaustion. *Nature Immunology* **22**, 205-215 (2021).

39.   Simmen, F.A., Alhallak, I. & Simmen, R.C.M. Malic enzyme 1 (ME1) in the biology of cancer: it is not just intermediary metabolism. *J Mol Endocrinol* **65**, R77-R90 (2020).

40.   Lu, Y.X.*, et al.* ME1 Regulates NADPH Homeostasis to Promote Gastric Cancer Growth and Metastasis. *Cancer Res* **78**, 1972-1985 (2018).

41.   Sena, L.A.*, et al.* Mitochondria are required for antigen-specific T cell activation through reactive oxygen species signaling. *Immunity* **38**, 225-236 (2013).

42.   Wen, T.*, et al.* NKG7 Is a T-cell-Intrinsic Therapeutic Target for Improving Antitumor Cytotoxicity and Cancer Immunotherapy. *Cancer Immunol Res* **10**, 162-181 (2022).

43.   Liu, C.*, et al.* Malic Enzyme 1 Indicates Worse Prognosis in Breast Cancer and Promotes Metastasis by Manipulating Reactive Oxygen Species. *Onco Targets Ther* **13**, 8735-8747 (2020).

44.   Gorrini, C., Harris, I.S. & Mak, T.W. Modulation of oxidative stress as an anticancer strategy. *Nat Rev Drug Discov* **12**, 931-947 (2013).

45.   Meermeier, E.W.*, et al.* Tumor burden limits bispecific antibody efficacy through T cell exhaustion averted by concurrent cytotoxic therapy. *Blood Cancer Discov* **2**, 354-369 (2021).

46.   Zheng, L.*, et al.* Pan-cancer single-cell landscape of tumor-infiltrating T cells. *Science* **374**, abe6474 (2021).

47.   Fang, X.*, et al.* TCF-1(+) PD-1(+) CD8(+)T cells are associated with the response to PD-1 blockade in non-small cell lung cancer patients. *J Cancer Res Clin Oncol*, 2653–2660 (2021).

48.   Hudson, W.H.*, et al.* Proliferating Transitory T Cells with an Effector-like Transcriptional Signature Emerge from PD-1(+) Stem-like CD8(+) T Cells during Chronic Infection. *Immunity* **51**, 1043-1058.e1044 (2019).

49.   Jansen, C.S.*, et al.* An intra-tumoral niche maintains and differentiates stem-like CD8 T cells. *Nature* **576**, 465-470 (2019).

50.   Siddiqui, I.*, et al.* Intratumoral Tcf1[+]PD-1[+]CD8[+] T Cells with Stem-like Properties Promote Tumor Control in Response to Vaccination and Checkpoint Blockade Immunotherapy. *Immunity* **50**, 195-211.e110 (2019).

51.   Wen, S.*, et al.* TCF-1 maintains CD8(+) T cell stemness in tumor microenvironment. *J Leukoc Biol* **110**, 585-590 (2021).

52.   Zhang, J., Lyu, T., Cao, Y. & Feng, H. Role of TCF-1 in differentiation, exhaustion, and memory of CD8(+) T cells: A review. *Faseb j* **35**, e21549 (2021).

Figures

Fig.1

a

b Gated on CD8+

c Low ΔΨm CX3CR1+

d 1 wk post-SFRT

e

f

Extended data Fig. 1

High ΔΨm CX3CR1+

% CX3CR1+ Low ΔΨm trends in Responders

% CX3CR1+ Low ΔΨm trends in Non−Responders

Fig. 2

Extended data Fig. 2

a

b

IL-2 STAT5 signaling

c

Fig. 3

Extended data Fig. 3

Fig. 4

Fig. 6

Fig. 7

Extended data Fig. 4

a

Ctrl mRNA   ME1 mRNA

ME1

β-actin

b

Western Blot Relative Protein Levels

Relative ME1 Protein Levels

Healthy Donors

■ control mRNA
■ ME1 mRNA

Spare Respiratory Capacity

c

ns

OCR (pmol/min)

Control mRNA   ME1 mRNA

Resting CD8⁺ T cells

d

ns

ROS MFI

Control mRNA   ME1 mRNA