A primary microcephaly-associated *sas-6* mutation perturbs centrosome duplication,

dendrite morphogenesis, and ciliogenesis in *Caenorhabditis elegans*

Mary Bergwell[1*], Amy Smith[2*], Ellie Smith[3§], Carter Dierlam[3§], Ramon Duran[3], Erin Haastrup[3],

Rebekah Napier-Jameson[4], Rory Seidel[3], William Potter[3], Adam Norris[4], and Jyoti Iyer[3#]

* and § These authors contributed equally to this work

From [1]Oklahoma Medical Research Foundation, Oklahoma City, OK 73104, [2]Pfizer Inc., 875

Chesterfield Parkway West, Chesterfield, MO 63017, [3]Department of Chemistry and

Biochemistry, University of Tulsa, Tulsa, OK 74104, [4]Southern Methodist University, PO Box

750100, Dallas, TX 75275

Running Head: *sas-6(L69T)* affects centrioles and cilia

[#]To whom correspondence should be addressed:

Jyoti Iyer: [3]Department of Chemistry and Biochemistry, University of Tulsa, Tulsa, OK 74104;

jgi2708@utulsa.edu; Tel. (918) 631-2070.

Keywords: SAS-6, SASS6, rare disease, microcephaly, centrosome, centriole, cilia, *C. elegans*

Abstract

The human SASS6(I62T) missense mutation has been linked with the incidence of primary
microcephaly in a Pakistani family, although the mechanisms by which this mutation causes
disease remain unclear. The SASS6(I62T) mutation corresponds to SAS-6(L69T) in *C. elegans*.
Given that SAS-6 is highly conserved, we modeled this mutation in *C. elegans* and examined
sas-6(L69T) effect on centrosome duplication, ciliogenesis and dendrite morphogenesis. Our
studies revealed that all the above processes are perturbed by the *sas-6(L69T)* mutation.
Specifically, *C. elegans* carrying the *sas-6(L69T)* mutation exhibit an increased failure of
centrosome duplication in a sensitized genetic background. Further, worms carrying this
mutation also display shortened phasmid cilia, an abnormal phasmid cilia morphology, shorter
phasmid dendrites and chemotaxis defects. Our data show that the centrosome duplication
defects caused by this mutation are only uncovered in a sensitized genetic background,
indicating that these defects are mild. However, the ciliogenesis and dendritic defects caused by
this mutation are evident in an otherwise wild-type background, indicating that they are stronger
defects. Thus, our studies shed light on the novel mechanisms by which the *sas-6(L69T)*
mutation could contribute to the incidence of primary microcephaly in humans.

Introduction

Autosomal recessive primary microcephaly (MCPH) is a rare disease that causes a reduced
brain size in infants along with a host of other physical and intellectual disabilities (Zaqout et. al.
2017). According to the OMIM database, thirty microcephaly loci named MCPH1-MCPH30 have
been identified thus far (https://www.omim.org/entry/251200). The current consensus in the field
is that most mutations linked to the incidence of MCPH deplete the pool of neural progenitor
cells during brain development. This results in fewer neurons being produced during embryonic
development, thereby leading to a smaller brain size (Phan and Holland 2021). Although some
MCPH-associated mutations have been previously modeled in *in vivo* model systems to

understand how MCPH-associated mutations contribute to the disease pathology, there are still

many MCPH-linked mutations that remain uninvestigated (Phan and Holland 2021). One such

MCPH-associated mutation is a mutation within the *SASS6* gene (*sas-6* in *C. elegans*) (Khan et

al. 2014). In 2014, a study by Khan et al. identified a leucine to threonine missense mutation in

the human *SASS6* gene to be associated with the incidence of MCPH in a consanguineous

Pakistani family (Khan et al. 2014).

SAS-6 protein function is conserved amongst several species including the multicellular

eukaryotic nematode *C. elegans* and humans (Dammerman et al. 2004, Leidel et al. 2005). The

sas-6 gene was first identified to regulate centrosome duplication in *C. elegans* (Dammerman et

al. 2004, Leidel et al. 2005). Centrosomes consist of a pair of cylindrical centrioles: a mother

and a daughter centriole that are embedded in pericentriolar material. Major functions of these

organelles involve organizing a mitotic spindle and aiding in cilia assembly (Nigg and Holland,

2018). Proper cell division requires a precise duplication of centrosomes. There are two levels

of regulation of centrosome duplication. First, it is critical that centrioles duplicate only once per

cell cycle, and second, it is imperative that each mother centriole gives rise to a single daughter

centriole. Defects in centrosome duplication can impair proper chromosome segregation,

leading to genome instability and aneuploidy. Indeed, increasing centrosome numbers by

elevating the expression levels of the master centrosome duplication kinase Plk4 (ZYG-1 in *C.*

elegans) is sufficient to induce tumorigenesis in mice (Levine et al. 2017). Along with ZYG-1,

SAS-6 also plays a critical role in mediating centrosome duplication. SAS-6 is recruited to the

daughter centriole through its direct interaction with ZYG-1 and with another centriolar protein

SAS-5 (Lettman *et al.* 2013). These interactions are essential to initiate daughter centriole

assembly.

In addition to regulating centrosome biogenesis, SAS-6 function is also essential for cilia

assembly in *C. elegans* (Li et al. 2017). Cilia are microtubule-based organelles that are critical

for cell signaling and development (Drummond 2012). Sensory cilia are the sole type of cilia that

are present in *C. elegans.* These cilia are non-motile in nature and they play an essential role in

sensing the extracellular environment (Inglis et al. 2007). There are a total of sixty ciliated

sensory neurons in *C. elegans* hermaphrodites*.* The cilia are present at the dendritic ends of the

sensory neurons (Inglis et al. 2007). Although a previous study has shown that SAS-6 function

is critical for proper ciliogenesis (Li et al. 2017), the mechanism by which SAS-6 promotes

ciliogenesis is unknown. It has been especially difficult to study the role of SAS-6 in ciliogenesis

because *sas-6/SASS6* is an essential gene (Kamath et al. 2003, Blomen et al. 2015). Due to the

important role of SAS-6 in regulating centrosome duplication, most mutations in *sas-6* yield

embryonic lethality or sterility (O'Rourke et al. 2011, Song et al. 2011, The C. elegans Deletion

Mutant Consortium et al. 2012, Lettman *et al.* 2013).

Ciliogenesis defects have been proposed as an important underlying cause of

microcephaly (Alcantara et al. 2014, Gabriel et al. 2016, Zhang et al. 2019, Ding et al. 2019,

Sohayeb et al. 2020, Farooq et al. 2020). However, the effect of this *SASS6* mutation on

ciliogenesis has never been investigated. The effect of the MCPH-associated *SASS6* mutation

on centrosome duplication in its endogenous context has also not been determined thus far. It is

important to note that SAS-6 levels in the cell are tightly regulated. An increase or a decrease in

SAS-6 levels causes either centrosome overduplication or a failure of centrosome duplication,

respectively (Dammerman et al. 2004, Leidel et al. 2005). Therefore, it is critical to model the

MCPH-associated *SASS6* mutation in an endogenous context in a multicellular eukaryotic

animal model. However, to date such a study has not been reported.

In this study, we used CRISPR/Cas9 genome editing to re-create the MCPH-associated

sas-6(L69T) mutation in *C. elegans*. Our *in vivo* studies in the worm indicate that the *sas-*

6(L69T) mutation does not result in embryonic lethality when this is the sole mutation in *sas-6* in

an otherwise wild-type background. However, introducing this mutation into a tight temperature-

sensitive *zyg-1(it25)* strain that is partially compromised for centrosome duplication results in a

robust increase in embryonic lethality at the semi-permissive temperature. These data indicate

that this MCPH-associated mutation mildly perturbs centrosome duplication in *C. elegans.*

Surprisingly, while the centrosome duplication defects were only observed in a sensitized

genetic background, ciliogenesis and dendritic defects were observed even in a wild-type

background in *C. elegans* carrying the *sas-6(L69T)* mutation. Specifically, our studies reveal

that the *sas-6(L69T)* mutation causes a reduction of the phasmid cilia and phasmid dendrite

lengths and results in an abnormal phasmid cilia morphology. Thus, our studies indicate that the

MCPH-associated *sas-6(L69T)* mutation affects ciliogenesis and dendrite morphogenesis more

severely than centrosome duplication in *C. elegans.* In conclusion, our studies have uncovered

a novel role of the MCPH-associated *sas-6(L69T)* mutation in regulating ciliogenesis and

dendrite morphogenesis in the worm.

Materials and Methods

C. elegans **maintenance**

All *C. elegans* strains were grown at 20°C on OP50 *E. coli* bacteria seeded onto MYOB agar

plates, unless otherwise indicated (as in the case of different assays). **Supplementary Table**

S1 lists all the *C. elegans* strains that have been generated and used in this study.

CRISPR/Cas9 genome editing

Assembled ribonucleoprotein complexes were microinjected into *C. elegans* gonads to perform

CRISPR/Cas9 genome editing as described previously (Paix et al. 2015, Paix et al. 2016, Iyer et

al. 2019, Smith et al. 2020). Homozygosity of the *sas-6(L69T)* edit was confirmed by PCR

followed by restriction digestion with EcoRI and by DNA sequencing. **Supplementary Table S2**

lists the sequences of the screening primers, guide RNAs and repair templates that were used

to make the CRISPR strains that have been used in this study.

Brood count and embryonic viability assays

All brood count assays and embryonic viability assays were performed as described previously

(Xie *et al.* 2022). For both brood count as well as embryonic viability assays, L4 stage *C.*

elegans were plated at the respective indicated temperatures. For brood count assays, all the

progeny that were produced by each single worm over the worm's reproductive lifespan were

counted and quantified. For embryonic viability assays, the percentage of dead and live progeny

produced by each worm was determined by counting. The percentage of viable progeny was

then computed by dividing the number of live progeny by the total number of progeny and by

multiplying this ratio by 100.

Confocal live imaging of centrosome duplication

To monitor centrosome duplication in *C. elegans* embryos, live imaging was performed using an

Olympus Spinning-Disk confocal microscope mounted with a Yokagawa scanhead. The

cellSens software (Olympus America Inc., Waltham, MA) was used for image acquisition and

processing. To prepare the *C. elegans* strains for imaging, L4 stage *C. elegans* were plated at

21.8°C overnight. The next day, gravid adult worms were picked into 8 to 12 µl of egg buffer

(118 mM NaCl, 47 mM KCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$, 5 mM HEPES pH 7.4) on a square glass

coverslip and dissected near the vulva using a needle to release embryos. The coverslip was

carefully mounted on a 2% agarose pad (2% agarose LE in egg buffer). The edges of the

coverslip were sealed with molten Vaseline using a paintbrush. These slides were subjected to

time-lapse imaging by taking 1.25 µm Z-stacks over an interval of approximately 132 seconds.

Maximum intensity Z-projections were computed using the cellSens software (Olympus America

Inc.) and converted into an AVI format for making videos. The brightness and contrast were

adjusted for the control and *sas-6(L69T)* videos for better non-quantitative video visualization.

Genetic crosses

To introduce the *sas-6(L69T)* mutation into the *zyg-1(it25)* background, first, males were

generated from the *zyg-1(it25)* strain using alcohol treatment. Specifically, L4 stage

hermaphrodites from the *zyg-1(it25)* strain were incubated with 10% ethanol in M9 buffer for 30

minutes, washed thrice with M9 buffer and allowed to recover for an hour. After recovery, the

worms were plated onto MYOB plates and incubated at 20°C until males appeared. Following

this, *zyg-1(it25)* homozygous males were mated with *sas-6(L69T)* L4 hermaphrodites. The F1

progeny from this cross were screened for heterozygosity of the *sas-6(L69T)* mutation using

PCR and restriction digestion. Cross progeny were identified and screened for homozygosity of

both edits using PCR and restriction digestion (for homozygosity screening of the *sas-6(L69T)*

mutation) and 24°C temperature shifts (for homozygosity screening of the *zyg-1(it25)* mutation).

Homozygosity of fluorescent markers was confirmed by confocal fluorescence imaging.

For the cilia imaging experiments, PY6100 males *(oyIs59[osm-6p::gfp] III)* were crossed with

hermaphrodites from the IYR001 (*sas-6(luv1[sas-6::ha]) IV*) and IYR002 (*sas-6(luv2[sas-*

6[L69T])) IV) strains to produce strains expressing either SAS-6::HA (control) or SAS-

6(L69T)::HA along with a pan-ciliary GFP marker. These strains were named IYR026 and

IYR027, respectively. F1 progeny were screened for heterozygosity at both loci by screening F1

hermaphrodites for the presence of GFP fluorescence in the phasmid neurons. F2 homozygotes

for *sas-6::ha* were identified as described previously (Bergwell et al. 2019) by PCR and

digestion by NdeI followed by agarose gel electrophoresis. F2 homozygotes for *sas-6(L69T)::ha*

were identified by PCR and digestion by EcoRI followed by agarose gel electrophoresis.

***C. elegans* phasmid cilia and dendrite imaging**

C. elegans phasmid cilia and dendrites were imaged as described previously (Xie et al. 2022).

For the phasmid and dendrite imaging experiments, a new strain IYR040 (*oyIs59 [osm-6p::gfp]*

III; sas-6(luv40)) IV) was created by CRISPR/Cas9 editing of the IYR026 strain. Specifically, to

generate the IYR040 strain, the IYR026 strain (*oyIs59 [osm-6p::gfp] III; sas-6(luv1[sas-6::ha])*

IV) was CRISPR-edited using the same guide RNA and the repair template that was used to

introduce the leucine 69 to threonine (L69T) mutation, with the only exception that the IYR040

repair template lacked the L69T mutation. As a result, the IYR040 strain is genetically identical

to the IYR027 strain except that it is wild-type for SAS-6. This strain served as a control for all of

our phasmid cilia and dendrite imaging experiments. For the phasmid cilia and dendrite imaging

assays, L4 stage *C. elegans* from the IYR040 and IYR027 strains were grown at 16°C

overnight. The next day, young adult worms were anaesthetized using 10 mM Levamisole,

mounted on 2% agarose pads and imaged within 30 minutes using an Olympus IX83 Spinning

Disk confocal microscope coupled with a Yokogawa X1 Scan head (Olympus America Inc.).

0.25 μm Z-slices were taken from the top to the bottom of each phasmid neuron using the

cellSens software (Olympus America Inc.). All images were taken at the same laser intensity

and exposure time.

Phasmid cilia and dendrite length quantification

To quantify phasmid cilia and dendrite lengths, the raw .vsi files that were captured using the

cellSens software (Olympus America Inc.) were opened in ImageJ using the OlympusImageJ

Plugin. ImageJ (Abràmoff et al. 2004) was then used to convert the raw data into maximum

intensity Z-projections. The scale was set to 1 pixel = 110 nm. Since the expression of GFP in

the phasmid neurons varied from worm to worm, the brightness and contrast were adjusted

non-uniformly to delineate the beginning and end of each phasmid cilium to ensure an accurate

measurement of the phasmid cilia lengths. For the dendrite imaging, the single frame view was

used in cellSens (Olympus America, Inc.) to visualize the beginning and end of each phasmid

dendrite in 3D to ensure that the dendrite lengths were being properly measured. Dendrites and

cilia whose beginning or end could not be delineated were excluded from our analysis. To

measure cilia and dendrite lengths, a line was traced along the entire length of the cilium or the

dendrite using the Segmented Lines tool in ImageJ (Abràmoff et al. 2004). The lengths of the

shorter of the two phasmid dendrites and the longer of the two phasmid cilia were then

measured in micrometers using ImageJ (Abràmoff et al. 2004) and have been presented in

Figures 4D and **6C**.

Criteria used for quantifying phasmid cilia morphology defects

Phasmid cilia that did not exhibit the canonical phasmid cilia structure with two transition zones

and cilia projecting from the transition zones were quantified as having an abnormal phasmid

cilia morphology. Specifically, instances where the transition zones could not be clearly

distinguished from the ciliary axoneme or where we were unable to clearly identify the transition

zone due to additional transition zone-like GFP aggregates being present in the vicinity of the

transition zone were classified as being abnormal.

C. elegans **chemotaxis assay**

Olfaction assays were performed as previously described in Liang et al. 2022.

Briefly, *C. elegans* were maintained by standard techniques on NGM agar plates seeded with

OP50 *Escherichia coli* at 20°C. Worms were synchronized to L4 stage. Worms were washed off

the plates using M9 buffer, allowed to gravity settle, supernatant removed, and then washed a

further 4 times. Unseeded NGM plates (50 mM NaCl, 1.7% agar, 0.25% peptone, 1 mM $CaCl_2$,

0.013 mM Cholesterol, 1 mM $MgSO_4$, 25 mM KPO_4) were divided into four quadrants, two with

solvent and two with odorant. Ethanol was used as a solvent. A 1:500 dilution of the odorant

butanone was made in ethanol. The Chemotaxis Index (CI) was calculated using the formula CI

= (# animals in two odorant quadrants)/(# animals in any of the four quadrants). Four plates

were used per genotype per experiment. Assays were repeated 3 times.

Whole worm extracts for western blotting

Whole worm extracts for western blotting were prepared as described previously (Xie et al.

2022). Briefly, 100 gravid adults from each genotype were washed with M9 buffer twice. The M9

buffer was removed and the worm pellet was resuspended in 40 µl of 4X SDS PAGE sample

buffer, boiled for 10 minutes at 95°C and stored at -30°C until further use.

Western blotting

Western blotting was performed using the Bio-Rad wet-transfer method as described previously

(Xie et al. 2022). Briefly, for western blotting of whole worm extracts, 10 to 14 µl of extracts were

loaded onto each well of a 7.5% SDS-PAGE gel. The gels were run until proper band separation

was achieved and the protein was transferred onto a 0.2 µm nitrocellulose membrane using the

wet-transfer method (Bio-Rad Laboratories). The membranes were blocked with Odyssey

Blocking Buffer (LiCOR Biosciences, Inc.), probed with anti-HA (Cell Signaling, catalog C29F4

Rabbit mAb #3724) and anti-tubulin (Santa Cruz Biotechnology, Inc., catalog sc-32293)

antibodies overnight at 4°C. The membranes were washed thrice with 1X TBST (0.1% w/v

Tween-20, 50 mM Tris-Cl, pH 7.6, 150 mM NaCl) and incubated with IRDye® 800CW Donkey

anti-Rabbit IgG secondary antibody (LiCOR Biosciences, Inc. catalog # 926-32213) and IRDye®

680RD Goat anti-Mouse IgG secondary antibody (LiCOR Biosciences, Inc. catalog # 926-

68070) for 1 hour at room temperature. The membranes were washed again thrice with

1XTBST and imaged using the LI-COR Odyssey CLx imager (LI-COR Biosciences, Inc.). The

band intensities were adjusted equally for all the gel lanes for better data visualization and

presentation. **Supplementary Table S3** lists all the antibodies that have been used in this

study.

Western blot quantification and image presentation

The band intensities across the entire membrane were uniformly adjusted using the LiCOR

Image Studio software (LI-COR Biosciences, Inc.) and saved in a .tiff format for performing

quantitative analysis of the bands. Quantitative western blotting was performed using ImageJ

(Abràmoff et al. 2004). Briefly, the band intensities for SAS-6::HA were determined using

ImageJ and normalized with respect to the loading control Tubulin. ImageJ was also used to

rotate and crop representative gel images presented in **Figure S1** and in all other fluorescence

images that are presented in this paper**.**

Modeling SAS-6(L69T) using AlphaFold2

The N-terminal coiled-coil dimer construct of *C. elegans* centriolar protein SAS-6 was

crystallized by Hilbert et al. 2013. The ceSAS-constructs required for crystallization involved the

removal of the $\alpha 2$-$\beta 5$ loop ($\Delta 103$-130) which was reported to not affect the N-terminal domain

structure within a C_α rmsds of ~0.3 Angstroms. The crystallographic data was deposited in the

Research Collaboration for Structural Bioinformatics databank under accession number PDB ID

code 4G79.

The native SAS-6 structure (AF-062479-F1) available from the AlphaFold Protein

Structure Database (Jumper et al. 2021; https://AlphaFold.ebi.ac.uk/) has leucine at position 69

at the end of the β_4 sheet just prior to the asparagine-asparagine which initiates the α_1 helix

occurring at positions 70 to 82 (using the Hilbert et al. 2013 author numbering system). It

should be noted that the residue numbering system reported for ceSAS-6 in Uniprot (062479)

has two additional amino acids (glycine-serine) at the N-terminus and thus the sequence is

offset and the α_1 helix starts at position 72 in this numbering system. The reported per-residue

confidence score (pLDDT) for residue leucine 69 was 92.37 suggesting that the AlphaFold

model confidence level was very high for this residue.

The SAS-6(L69T) mutant protein structure containing the single leucine to threonine

substitution was modeled by loading the modified sequence into the Google Colaboratory

(Colab) services using ColabFold: AlphaFold2 using MMseqs2 as described by Mirdita et al.

2022, and available through the AlphaFold2.ipynb v2.1.0. with the suite of public notebooks

available through the deepmind/AlphaFold github. The 4G79 and the AlphaFold models were

uploaded into the program ChimeraX (https://www.cgl.ucsf.edu/chimerax/) developed from the

Resource for Biocomputing, Visualization, and Informatics for Visualization (RBVI), University of

California, San Francisco.

The Molecular Lipophilicity Potential (MLP) is a construct that spreads atomic values out

in 3D. The MLP coloring on the molecular surface is based on pyMLP

(https://www.rbvi.ucsf.edu/chimerax/docs/user/commands/mlp.html) as originally developed by

Broto et al. 1984 and Laguerre et al. 1997 and as modified

https://www.cgl.ucsf.edu/chimera/data/mlp-sep2016/index.html and was used without any color

scale modifications.

Statistical analysis

Graphpad Prism (GraphPad Software, Inc.) was used for performing statistical analysis of the

data. Unpaired two-tailed t-tests were performed to determine statistical significance of the data

for the brood counts, western blot assays, phasmid cilia lengths, phasmid dendrite lengths and

chemotaxis assays. The Fisher's exact test was performed to determine if the phasmid cilia

morphology defects were statistically significant. A p-value of less than 0.05 was considered to

attain statistical significance.

Results

An isoleucine to threonine substitution of SASS6 was reported to be linked with the occurrence

of MCPH in a Pakistani family (Khan et al. 2014). As shown previously by Khan et al., alignment

of the amino acid sequences of the SAS-6 protein of different species showed that the MCPH-

associated amino acid of human SASS6 is remarkably conserved among several species

including *C. elegans* (Khan et al. 2014). Notably, the *sas-6* gene was first identified in *C.*

elegans and was subsequently shown to function similarly in humans (Dammerman et al. 2004,

Leidel et al. 2005). Due to the high functional conservation of SAS-6, we determined that *C.*

elegans would be an appropriate eukaryotic model to thoroughly investigate the consequences

of this mutation. The human SASS6(I62T) mutation corresponds to SAS-6(L69T) in *C. elegans.*

The *sas-6(L69T)* mutation was successfully introduced into *C. elegans* by CRISPR/Cas9

editing

Using CRISPR/Cas9 genome editing, we generated a leucine to threonine substitution at

amino acid 69 of the *C. elegans* SAS-6 protein by editing the endogenous *sas-6* gene of *C.*

elegans. This strain was named IYR002 and the corresponding allele was named *sas-6(luv2).*

For ease of understanding, this strain has been referred to as the *sas-6(L69T)* strain and the

corresponding MCPH-associated mutation has been referred to as the *sas-6(L69T)* mutation

throughout this manuscript. The repair template used for introducing the *sas-6(L69T)* mutation

carried an EcoRI restriction site **(Figure 1A)**. As shown in **Figure 1B**, upon PCR amplification

and restriction digestion of the *sas-6* gene with EcoRI, several homozygous edited worms were

obtained in the F2 generation (lanes with black stars). The incorporation of the *sas-6(L69T)*

mutation was confirmed by DNA sequencing.

The microcephaly-associated *sas-6(L69T)* mutation does not affect brood size or

embryonic viability in a wild-type background

The function of the *sas-6* gene is essential for centrosome duplication (Dammerman et

al. 2004, Leidel et al. 2005). Proper centrosome duplication is required for accurate embryonic

divisions in *C. elegans.* Since the *sas-6(L69T)* mutation has a pathological relevance in

humans, we proposed that this mutation could impair SAS-6 function in regulating centrosome

duplication, thereby leading to defective centrosome duplication in *C. elegans.* This could, in

turn, affect *C. elegans* brood size and embryonic viability. Therefore, to investigate the effect of

the *sas-6(L69T)* mutation on centrosome duplication, brood counts and embryonic viability

assays were performed using *C. elegans* carrying this mutation. Surprisingly, the introduction of

the *sas-6(L69T)* mutation did not yield any major effect on either brood size or embryonic

viability at 16°C, 20°C and 25°C as compared with control worms (wild-type N2 worms with HA-

tagged endogenous *sas-6*) **(Figures 2A and 2B)**. These data indicate that the *sas-6(L69T)*

mutation likely has a mild effect on SAS-6 function in regulating centrosome duplication which

cannot be uncovered in an otherwise wild-type background.

The *sas-6(L69T)* mutation increases centrosome duplication failures and embryonic

lethality in a sensitized *zyg-1(it25)* background

ZYG-1 is a functional homolog of human Plk4 and a master kinase whose activity is

critical for proper centrosome duplication (O'Connell et al. 2001, Habedanck et al. 2005,

Bettencourt-Dias et al. 2005). Decreasing ZYG-1 protein levels causes a failure of centrosome

duplication while elevating them causes centrosome overduplication (O'Connell et al. 2001,

Peel et al. 2017, Wolf et al. 2018). The temperature-sensitive *zyg-1(it25)* allele has a proline to

leucine substitution within the ZYG-1 cryptic polo box which impairs ZYG-1 function at the

restrictive temperature, leading to a failure of proper centrosome duplication (Kemphues et al.

1988, Kemp et al. 2007). By controlling the temperature at which the *zyg-1(it25)* strain is grown,

the rate of centrosome duplication failure in this strain can be controlled. For example, at a

restrictive temperature of 24°C and above, the *zyg-1(it25)* strain produces 100% dead embryos

due to a complete failure of centrosome duplication (Kemp et al. 2007). However, at lower semi-

permissive temperatures (temperatures below 24°C), centrosome duplication still partially

occurs in the *zyg-1(it25)* strain, yielding varying levels of embryonic viability with a temperature

of 20°C yielding between 93.7% to 98.7% embryonic viability (Song et al. 2008). To uncover

mild defects in SAS-6 function in regulating centrosome duplication in the *sas-6(L69T)* mutant

worms, we first obtained a *zyg-1(it25)* strain expressing fluorescently-tagged centrosome

(mCherry::SPD-2) and chromosome (GFP::histone) markers (OC869 from the O'Connell lab) to

visualize centrosome duplication. This strain served as a control for our experiment and has

been referred to as the *zyg-1(it25)* strain in **Figure 3**. Using this strain as the background strain,

as shown in **Figure 3A,** we performed genetic crossing to combine the *sas-6(L69T)* and *zyg-*

1(it25) mutations. This strain has been referred to as the *zyg-1(it25);sas-6(L69T)* strain in

Figure 3. We reasoned that if the *sas-6(L69T)* mutation mildly affects SAS-6 function in

regulating centrosome duplication, combining it with the *zyg-1(it25)* mutation at the semi-

permissive temperature would yield higher levels of centrosome duplication failures and

increased embryonic lethality than what each mutant produces on its own.

Upon introducing the *sas-6(L69T)* mutation into the *zyg-1(it25)* strain expressing

fluorescent centrosome and chromosome markers, centrosome duplication in the *zyg-1(it25)*

and *zyg-1(it25);sas-6(L69T)* worms was assessed at the semi-permissive temperature of 21.8°C

(Figures 3B and 3C). Notably, introducing the *sas-6(L69T)* mutation greatly increased the

centrosome duplication failures exhibited by the *zyg-1(it25)* mutants at 21.8°C **(Figure 3C)**.

Specifically, in the *zyg-1(it25)* single mutant alone, about 83% of the centrosomes analyzed

duplicated normally at the semi-permissive temperature of 21.8°C (n=52). However,

interestingly, in the *zyg-1(it25);sas-6(L69T)* double mutant, proper centrosome duplication was

reduced to only about 28% (n=58). Since accurate centrosome duplication is required for

embryonic viability, we next assessed whether embryonic viability of the *zyg-1(it25)* strain was

also affected in the presence of the *sas-6(L69T)* mutation.

The embryonic viability of the *zyg-1(it25)* strain expressing fluorescent markers alone at

the semi-permissive temperature of 21.7°C was about 44% (number of worms=18; number of

progeny= 1103). However, introducing the *sas-6(L69T)* mutation into this background reduced

embryonic viability to only about 7% (number of worms=18; number of progeny= 912; unpaired

two-tailed t-test p<0.0001) **(Figure 3D)**. These data collectively indicate that the *sas-6(L69T)*

mutation mildly inhibits SAS-6 function in regulating centrosome duplication, thereby resulting in

a failure of proper centrosome duplication and increased embryonic lethality in the *zyg-1(it25)*

background.

Ciliogenesis is perturbed in *sas-6(L69T)* mutant *C. elegans*

Ciliogenesis defects have been linked with the incidence of microcephaly (Alcantara et

al. 2014, Gabriel et al. 2016, Wambach et al. 2018, Zhang et al. 2019, Ding et al. 2019,

Sohayeb et al. 2020, Farooq et al. 2020). Defective ciliogenesis is thought to result in cell cycle

arrest in neural progenitors during brain development. The loss of neural progenitors due to

either cell-cycle arrest or apoptosis has been proposed to be an underlying cause for the

incidence of MCPH (Phan and Holland 2021). The effect of the *sas-6(L69T)* mutation on

ciliogenesis is currently unknown. Since ciliogenesis defects have been previously linked with

MCPH, we questioned whether worms carrying the *sas-6(L69T)* mutation exhibit any

ciliogenesis defects. Preliminary dye-filling assays (performed according to the protocol by

Power et al. 2020) did not reveal any severe dye-filling defects between control (wild-type N2

worms with HA-tagged endogenous SAS-6) and *sas-6(L69T)* worms. Specifically, we did not

observe any worm that completely excluded DiI in their phasmid neurons in either control or

sas-6(L69T) worms upon dye-filling. Our preliminary experiments revealed that 100% of the

control (n=23) and *sas-6(L69T)* (n=25) worms exhibited dye-filling with DiI. Nevertheless,

although dye-filling assays are a quick and easy method to visualize cilia defects, a lack of a

dye-filling defect phenotype does not exclude the presence of ciliary defects.

To directly determine whether the *sas-6(L69T)* strain exhibits ciliary defects, we obtained

a PY6100 strain that expresses a pan-ciliary marker. This strain expresses a GFP

transcriptional reporter for OSM-6*,* a protein that is critical for proper ciliogenesis (Perkins et al.

1986, Collet et al. 1998). Thus, the introduction of *osm-6p::GFP* marks *C. elegans* ciliated

sensory neurons with soluble GFP driven from the *osm-6* promoter, allowing for easy

visualization of the cilia (Bayer et al. 2020, Xie et al. 2022).

We introduced the *sas-6(L69T)* mutation into the PY6100 strain (*oyIs59 [osm-6p::gfp] III*)

by genetic crossing. The resultant strain was named IYR027. The experimental design for

combining *osm-6p::GFP* with the *sas-6(L69T)* mutation is shown in **Figure 4A.** The introduction

of *osm-6p::GFP* into the *sas-6(L69T)* strain by genetic crossing was confirmed by both confocal

microscopy and by genotyping using PCR. For ease of understanding, this new IYR027 strain

has been referred to as the *osm-6p::GFP; sas-6(L69T)* strain throughout the paper and has

been used to examine the cilia and dendritic defects that arise as a consequence of the *sas-*

6(L69T) mutation in Figures 4, 5 and 6.

Since the *sas-6(L69T)* strain also has an HA tag on the mutant SAS-6 protein, to create

an appropriate control strain, we first crossed the PY6100 strain with a strain expressing SAS-

6::HA to account for the potential effect of the HA-tag on SAS-6 function in regulating

ciliogenesis. This strain was named IYR026. We then CRISPR-edited the IYR026 strain to

make it genetically identical to the *osm-6p::GFP; sas-6(L69T)* strain except for the fact that this

new strain lacked the leucine 69 to threonine mutation. For convenience, we refer to this control

strain (IYR040) as the *osm-6p::GFP* strain. This strain served as the control for all the cilia and

dendrite analysis conducted in Figures 4, 5 and 6.

To investigate whether *sas-6(L69T)* mutant worms exhibit defects in ciliogenesis, we

examined the cilia of the phasmid neurons which are present in the posterior region of the

worms **(Figure 4B)**. Intriguingly, upon measuring the length of the phasmid neuronal cilia, we

found a significant decrease in the average ciliary length in the *osm-6p::GFP; sas-6(L69T)*

mutant worms **(Figure 4C, bottom panel, white bracket)** as compared with control (*osm-*

6p::GFP) worms **(Figure 4C, top panel, white bracket)**. Specifically, control *osm-6p::GFP*

worms had an average phasmid cilia length of about 5.82 μm (n=54) while the *osm-6p::GFP;*

sas-6(L69T) worms had shorter cilia with an average phasmid cilia length of only 3.99 μm

(n=59) (unpaired two-tailed t-test, p<0.0001). These data indicate that the *sas-6(L69T)* mutation

indeed perturbs ciliogenesis in *C. elegans.*

Phasmid cilia morphology and dendrite length are defective in the *sas-6(L69T)* mutant

worms

While analyzing cilia length of *osm-6p::GFP; sas-6(L69T)* worms, we found that some of

the cilia from this strain exhibited a disfigured cilia morphology as compared with control worms.

Our analysis revealed that 100% of the phasmid cilia of control (*osm-6p::gfp)* worms (n=61)

showed a normal phasmid cilia morphology with two distinct transition zones and a ciliary

axoneme projecting from each transition zone **(Figure 5A, 5B, 5D)**. However, 10 out of 65

(15.4%) of the *osm-6p::GFP; sas-6(L69T)* phasmid cilia exhibited an abnormal morphology

(Figures 5A, 5C, 5D). This abnormal phasmid cilia morphology defect of the *osm-6p::GFP; sas-*

6(L69T) worms was found to be statistically significant via a Fisher's exact test (p=0.0014).

These data indicate that the *sas-6(L69T)* mutation perturbs phasmid cilia morphology in a

subset of the phasmid cilia.

In addition to the phasmid cilia morphology defect, the *osm-6p::GFP; sas-6(L69T)*

worms also displayed a dendritic extension defect. In general, the average lengths of the

phasmid dendrites of control *osm-6p::GFP* worms **(Figure 6A)** were shorter than those of *osm-*

6p::GFP; *sas-6(L69T)* worms **(Figure 6B)** . Specifically, phasmid dendrites analyzed from

control worms exhibited an average length of 37.4 µm (n=38). In contrast, the average length of

phasmid dendrites of *sas-6(L69T)* worms was 28.1 µm (n=42) **(Figure 6C).** This phasmid

dendrite length defect of the *osm-6p::GFP;sas-6(L69T)* worms was found to be statistically

significant via an unpaired 2-tailed t-test (p<0.0001).

Chemotaxis is defective in the *sas-6(L69T)* mutant worms

Since defects in ciliogenesis in the *C. elegans* sensory neurons often affect worm

behaviors, we investigated the effect of the *sas-6(L69T)* mutation on chemotaxis to the

chemoattractant butanone. The experimental scheme for the chemotaxis assays is shown in

Figure 7A. Our studies demonstrated that *C. elegans* carrying the *sas-6(L69T)* mutation

exhibited a reduced attraction to 1:500 butanone as compared with control worms (two-tailed

unpaired t-test, p<0.0001) **(Figures 7B, 7C)**. Specifically, our data showed that the average

chemotaxis index of control worms (*sas-6::ha*) was 0.4107 (n=2825). In comparison, the

average chemotaxis index of *sas-6(L69T)* worms was reduced to 0.3307 (n=1711) **(Figures 7B,**

7C). These data indicate that defective ciliogenesis in the *sas-6(L69T)* mutant worms likely

causes a defective *C. elegans* chemotaxis response to 1:500 butanone. Hence, SAS-6 function

in regulating ciliogenesis is likely to be important for mediating certain worm behaviors such as

chemotaxis.

Interestingly, chemotaxis towards butanone is predominantly controlled by the AWC

amphid sensory neurons (Bargmann et al. 1993). These data indicate that ciliogenesis may also

be perturbed in the amphid sensory neurons in *sas-6(L69T)* mutant worms. In the present study,

we have mainly focused our efforts on analyzing ciliogenesis in the *C. elegans* phasmid

neurons. Future studies should be directed towards determining whether ciliogenesis in the

amphid neurons is also affected in *C. elegans* carrying the *sas-6(L69T)* mutation.

SAS-6 protein levels are altered in the presence of the *sas-6(L69T)* mutation in a *zyg-

***1(it25)* background but not in a wild-type background**

Some of the other known MCPH-associated mutations in centrosome duplication genes

such as STIL and PLK4 have been reported to affect their protein levels (Martin et al. 2014,

Patwardhan et al. 2018). The effect of *sas-6(L69T)* mutation on SAS-6 protein levels is

unknown. Therefore, we questioned whether the *sas-6(L69T)* mutation has any effect on SAS-6

protein levels. To address this, western blotting was performed on whole worm lysates from

zyg-1(it25);sas-6::ha (control) vs *zyg-1(it25);sas-6(L69T)::ha* worms and from *sas-6::ha* (control)

vs *sas-6(L69T)::ha* worms. Western blot analysis indicated that SAS-6 levels decrease on

average by 15% in the presence of the *sas-6(L69T)* mutation in the *zyg-1(it25)* background

(n=4; p=0.0475, unpaired two-tailed t-test) **(Figure S1A, S1B)**. However, the decrease in SAS-6

protein levels in the presence of the *sas-6(L69T)* mutation in an otherwise wild-type background

was not found to be statistically significant (n=4; p=0.1398, unpaired two-tailed t-test) **(Figures**

S1C, S1D). Since we observe ciliogenesis defects in the presence of the *sas-6(L69T)* mutation

even in a wild-type background **(Figures 4C, 4D, 5B, 5C, 5D)**, we propose that decreased SAS-

6 levels are likely not sufficient to explain the effects of the *sas-6(L69T)* mutation.

Modeling SAS-6(L69T) using AlphaFold2 suggests that the leucine to threonine

substitution at amino acid 69 does not have a major effect on SAS-6 protein structure

The L69T mutation is located within the conserved PISA (Present In SAS-6) motif of

SAS-6 **(Figure 8A)**. Further, this amino acid does not have a known involvement in the

dimerization of SAS-6 coiled-coil region, in the interaction of SAS-6 N-terminal head domains, in

the interaction of SAS-6 with ZYG-1 or in the interaction of SAS-6 with SAS-5 (van Breugel et al.

2011, Kitagawa et al. 2011, Qiao et al. 2013, Hilbert *et al,* 2013, Lettman et al. 2013) **(Figure**

8A). To elucidate the possible mechanism by which the *sas-6(L69T)* mutation affects

centrosome duplication and ciliogenesis, we used AlphaFold2 to predict how the L69T mutation

could alter the structure of the SAS-6 protein **(Figures 8B, 8C)**. **Figures 8B and 8C** show the

comparison of the SAS-6 PDB model 4GFC from Hilbert et al. 2013 **(Figure 8B)** compared with

the AlphaFold2-derived SAS-6(L69T) model **(Figure 8C)** using the ChimeraX program. Our

analysis indicates that the overall structural differences due to the L69T mutation in SAS-6 are

minor. Therefore, we propose that centriole architecture may not be significantly affected in the

sas-6(L69T) mutant worms. This is supported by our data that *C. elegans* carrying homozygous

copies of the *sas-6(L69T)* mutation are close to 100% viable at all temperatures **(Figure 2B)**.

However, the ChimeraX hydrophobicity surface model suggests that a deep hydrophobic pocket

that is found within the SAS-6 protein is reduced due to the MCPH-associated L69T mutation

(Figures 8B, 8C, dark blue arrows). The function of this hydrophobic pocket within SAS-6 has

not yet been elucidated.

Discussion

Through our studies, we have re-created the MCPH-associated *sas-6(L69T)* mutation in *C.*

elegans by CRISPR to generate the first *C. elegans* model of a MCPH-associated mutation. A

previous study investigated the effects of the human SASS6(I62T) mutation (the human

equivalent of the SAS-6(L69T) mutation) on centrosome duplication in U2OS cells (Khan et al.

2014). In their studies, the authors noted an inhibition of centrosome duplication in U2OS cells

expressing the SASS6(I62T) protein from exogenous plasmid constructs carrying the

SASS6(I62T) mutation. The levels of SAS-6 in the cell must be tightly regulated for its proper

cellular function. Either overexpressing the SAS-6 protein or depleting it can affect proper

centrosome duplication (Dammermann et al. 2004, Leidel et al. 2005). Therefore, to investigate

the effect of the *sas-6(L69T)* mutation on centrosome duplication in an endogenous context, we

monitored centrosome duplication in our *sas-6(L69T)* CRISPR worm model. Interestingly, *C.*

elegans carrying this mutation appeared phenotypically wild-type and did not exhibit any defects

in embryonic viability or brood size **(Figure 2)**. *C. elegans* embryonic divisions are reliant on

proper centrosome duplication. Since we did not detect any embryonic lethality in the presence

of the *sas-6(L69T)* mutation **(Figure 2B)**, we concluded that centrosome duplication is likely to

be only modestly affected in the *sas-6(L69T)* mutant worms in a wild-type background.

To uncover the potential milder effects of this mutation on centrosome duplication that

may not be visible in a wild-type genetic background, we introduced this mutation in the

sensitized *zyg-1(it25)* background. Interestingly, the *sas-6(L69T)* mutation significantly

enhanced the centrosome duplication failures and embryonic lethality of *zyg-(it25)* worms

(Figures 3B, 3C, 3D). These data demonstrate that the MCPH-associated *sas-6(L69T)*

mutation does, in fact, reduce SAS-6 function in regulating centrosome duplication. However,

this effect can only be revealed in a sensitized genetic background. Thus, at least in the context

of this MCPH-associated mutation, centrosome duplication is only mildly affected by this

mutation. Future studies will involve elucidating the molecular mechanism by which the *sas-*

6(L69T) mutation inhibits centrosome duplication.

Our data that the *sas-6(L69T)* mutation likely has a mild effect on SAS-6 function in

regulating centrosome duplication are consistent with the observed phenotypic effects of the

SASS6(I62T) mutation in human MCPH patients. *SASS6*/*sas-6* is an essential gene that is

required for proper centrosome duplication and cell division (Kamath et al. 2003, Dammerman

et al. 2004, Leidel et al. 2005, Blomen et al. 2015). If the *SASS6(I62T)* mutation severely

inhibited SASS6 function in regulating centrosome duplication, it would be expected to result in

embryonic lethality. As a result, there would not be human patients living with homozygous

copies of the *SASS6(I62T)* mutation. However, the fact that there are *SASS(I62T)* homozygous

individuals in the population (Khan et al. 2014) indicates that this mutation does not severely

perturb SASS6 role in regulating centrosome duplication in humans. Therefore, our data in *C.*

elegans which demonstrate that embryonic viability is unaffected in the presence of this

mutation **(Figure 2B)** are in line with the observations in human MCPH patients carrying this

mutation.

The effect of the *SASS6(I62T)* mutation on ciliogenesis has not been investigated. Upon

introducing the *sas-6(L69T)* mutation into a pan-ciliary marker expressing *C. elegans* strain, we

determined that ciliogenesis is perturbed in the *sas-6(L69T)* mutant worms **(Figures 4C, 4D,**

5B, 5C, 5D). Specifically, the *sas-6(L69T)* worms have shorter phasmid cilia and a defective

phasmid cilia morphology **(Figures 4C, 4D, 5B, 5C, 5D)**. Ciliogenesis defects have been

associated with the incidence of microcephaly (Alcantara et al. 2014, Gabriel et al. 2016,

Wambach et al. 2018, Zhang et al. 2019, Ding et al. 2019, Sohayeb et al. 2020, Farooq et al.

2020). Interestingly, in a study by Wambach et al.*,* the authors cultured MCPH patient derived

fibroblasts and found that these fibroblasts had shorter primary cilia than those of fibroblasts

that were cultured from a healthy control individual (Wambach et al. 2018). In another study,

Shohayeb et al. determined that mouse embryonic fibroblasts that were derived from WDR62

mutant mice carrying CRISPR-edited MCPH patient mutations exhibited a shortened ciliary

length, like we report here (Shohayeb et al. 2019). Thus, there is precedence for shorter ciliary

length being associated with MCPH.

We find that in *C. elegans* carrying the *sas-6(L69T)* mutation, centrosome duplication

defects can only be detected in a sensitized *zyg-1(it25)* genetic background **(Figures 3B, 3C).**

However, the ciliogenesis and dendritic defects induced by this mutation can be uncovered

even in an otherwise wild-type background **(Figures 4C, 4D, 5B, 5C, 5D)**. These findings lead

us to speculate that perhaps, at least in the context of this specific MCPH-associated mutation,

the inhibition of proper ciliogenesis may be a more important contributor to the development of

MCPH than the inhibition of centrosome duplication. However, it is also possible that the

centrosome duplication and ciliogenesis defects induced by this mutation synergize to cause in

MCPH in human patients carrying this mutation. In the future, it will be interesting to determine if

the *SASS6(I62T)* mutation also affects ciliogenesis in human cell lines. It will also be interesting

to examine if other such MCPH-associated mutations likewise affect ciliogenesis in addition to

affecting centrosome duplication.

Since *sas-6* is an essential gene, many *sas-6* mutant alleles are difficult to maintain and

analyze (O'Rourke et al. 2011, Song et al. 2011, The C. elegans Deletion Mutant Consortium et

al. 2012, Lettman *et al.* 2013). This has made it tedious to dissect the function of SAS-6 in

ciliogenesis. A previous study expressed a *sas-6* sgRNA and Cas9 protein using a heat-shock

promoter to demonstrate that SAS-6 plays a critical role in *C. elegans* ciliogenesis (Li et al.

2017). However, the mechanistic investigation of these defects has been challenging due to the

high lethality and sterility of *sas-6* mutant alleles. Notably, our studies indicate that *sas-6(L69T)*

mutant worms appear phenotypically wild-type and yet exhibit significant ciliogenesis defects.

Therefore, we expect that our *sas-6(L69T)* CRISPR strain will be a useful tool to unravel the

mechanism by which SAS-6 regulates ciliogenesis. In the future, forward genetic screens can

be performed with the *sas-6(L69T)* allele to identify additional genes that may genetically

interact with *sas-6* to regulate ciliogenesis in the worm.

Our data indicate that *C. elegans* harboring the *sas-6(L69T)* mutation also exhibit

sensory neuronal defects such as a shortened dendrite length **(Figure 6)**. This is a surprising

finding since *C. elegans* centrioles have been found to degenerate during neuronal

differentiation (Li et al. 2017, Serwas et al. 2017, Nechipurenko et al. 2017). One possibility is

that SAS-6 functions independently of its role as a core centriolar protein in regulating dendrite

length in *C. elegans* sensory neurons. Another possibility is that SAS-6 function at the centrioles

may be required for proper phasmid dendrite extension through an unknown mechanism. In the

future, transmission electron microscopy analysis of dendrites and cilia from *sas-6(L69T)* mutant

worms should be performed to determine the nature of the dendritic and ciliary defects in these

mutant worms. Future studies will also involve analyzing *C. elegans* strains expressing a variety

of fluorescently-tagged markers (e.g. Intraflagellar transport markers, transition zone markers,

neuron-specific markers, etc.) to further dissect the mechanism by which the *sas-6(L69T)*

mutation affects ciliogenesis and dendrite extension.

Defective chemotaxis towards butanone was seen in *C. elegans* carrying the *sas-*

6(L69T) mutation **(Figure 7).** Future studies should also be directed towards examining what

other worm behaviors (if any) are altered in *C. elegans* carrying the *sas-6(L69T)* mutation. It is

important to note that in this study, we have only examined the effect of the *sas-6(L69T)*

mutation on ciliogenesis and behaviors in the *C. elegans* hermaphrodite. The effect of this

mutation on *C. elegans* males is unknown. Assays such as male mating assays could be carried

out in the future to determine if male-specific behaviors are affected by the *sas-6(L69T)*

mutation.

Finally, modeling the L69T mutation using AlphaFold2 did not reveal any significant

changes in the structure of the SAS-6 protein **(Figures 8B, 8C)**. This is not a surprising finding

since *sas-6* is an essential gene and therefore severely affecting SAS-6 structure may alter

centriole architecture and result in embryonic lethality. The fact that there are live individuals

within the population harboring the *SASS6(I62T)* mutation suggest that this mutation does not

significantly alter SASS6 protein structure **(Figures 8B, 8C)**. However, our AlphaFold2 analysis

revealed that a deep hydrophobic pocket within SAS-6 is altered by the *sas-6(L69T)* mutation

(Figures 8B, 8C, dark blue arrows). Presently, no known function has been ascribed to this

hydrophobic pocket in SAS-6. Based upon our data, we are tempted to speculate that this SAS-

6 hydrophobic pocket may mediate an as-yet unknown interaction of SAS-6 with a protein(s)

that may be important for ciliogenesis, dendrite morphogenesis and/or centrosome duplication.

As our *C. elegans* model of the *sas-6(L69T)* mutation recapitulates many of the effects

seen in the human MCPH patients carrying this mutation, we expect that *C. elegans* will be

more widely adopted as a model system to study the effects of additional MCPH-associated

human mutations in the future.

Figure 1. The MCPH-associated *sas-6(L69T)* mutation was successfully generated in *C.*

***elegans* by CRISPR/Cas9 editing** A) Schematic of the CRISPR repair template used for

inserting the leucine to threonine mutation at amino acid 69 of the *C. elegans* SAS-6 protein

(Note: Only 10 out of 45 bases of the 5' and 3' homology ends of the repair template are

shown). The teal highlights represent bases that are identical between the repair template and

the *sas-6* gene sequence. The magenta highlight indicates the bases that were altered to

introduce the leucine 69 to threonine mutation into the endogenous *sas-6* gene. The bases

highlighted in gray represent silent mutations that were introduced in the repair template to

prevent its cutting by the Cas9 enzyme. B) Agarose gel electrophoresis of PCR products

digested with EcoRI to screen for homozygous-edited *C. elegans* carrying the *sas-6(L69T)*

mutation. Black stars: Homozygous-edited worms; het: heterozygote; WT: wild-type.

Figure 2. The *sas-6(L69T)* mutation does not affect *C. elegans* brood size or embryonic

viability A) Brood sizes of wild-type and *sas-6(L69T)* mutant worms were determined at

temperatures of 16°C, 20°C and 25°C. Each circle or triangle represents the brood size of a

single worm. No statistically significant difference in brood size was observed between wild-type

and *sas-6(L69T)* worms at any of the assessed temperatures. n= number of worms whose

brood was analyzed for each strain. n.s: Not significant. Error bars represent the standard

deviation. The middle bar represents the mean. B) Embryonic viability of wild-type and *sas-*

6(L69T) worms was analyzed at temperatures of 16°C, 20°C and 25°C. Each circle or triangle

represents the embryonic viability of a single worm. The middle line represents the median

embryonic viability with a 95% confidence interval. No major change in embryonic viability was

observed between wild-type and *sas-6(L69T)* worms at any of the assessed temperatures. The

viability of over 2000 embryos was analyzed for each strain at every temperature.

Figure 3. The *sas-6(L69T)* mutation increases the centrosome duplication failures and the

embryonic lethality of *zyg-1(it25)* worms. A) Schematic showing the genetic cross that was

performed to combine the *sas-6(L69T)* mutation with the *zyg-1(it25)* mutation. B) Stills from

time-lapse imaging of *zyg-1(it25)* and *zyg-1(it25);sas-6(L69T)* strains expressing fluorescently-

tagged centrosome and DNA markers. Magenta: mCherry-SPD-2 (centrosome marker); Green:

GFP-histone (DNA marker). In a majority of *zyg-1(it25)* mutant embryos, centrosome duplication

proceeds normally at a semi-permissive temperature of 21.8°C, yielding two centrosomes in

each cell of a 2-cell *C. elegans* embryo (**Figure 3B, top panel**). Scale bar = 10 µm. On the

contrary, most of the *zyg-1(it25);sas-6(L69T)* mutant centrosomes fail to duplicate at 21.8°C

resulting in a single centrosome present in each cell of a 2-cell *C. elegans* embryo (**Figure 3B,**

bottom panel). C) Quantification of B). n=Number of centrosomes analyzed. Approximately

83% of the *zyg-1(it25)* centrosomes duplicate normally at 21.8°C. In contrast, only about 28% of

the *zyg-1(it25);sas-6(L69T)* centrosomes duplicate normally at this temperature. D)

Quantification of embryonic viability of *zyg-1(it25)* single and *zyg-1(it25);sas-6(L69T)* double

mutants at the semi-permissive temperature of 21.7°C. Average embryonic viability is reduced

from ~44% to ~7% in the presence of the *sas-6(L69T)* mutation. The viability of 1103 embryos

from *zyg-1(it25)* worms and 912 embryos from *zyg-1(it25);sas-6(L69T)* worms was analyzed.

Embryos from 18 worms were analyzed for each condition. Unpaired 2-tailed t-test p<0.0001.

Error bars represent the standard deviation and the middle bar represents the mean.

Figure 4. Phasmid cilia length is shorter in *sas-6(L69T)* mutant worms. A) A schematic of

the genetic crossing experiment that was performed to introduce the MCPH-associated *sas-*

6(L69T) mutation into the PY6100 pan-ciliary marker strain. B) Cartoon showing the anatomical

positioning of a *C. elegans* phasmid cilium and dendrite within the tail region of the worm. C) Z-

projection confocal images of phasmid neurons from *osm-6p::gfp* (top panel) and *osm-6p::gfp;*

sas-6(L69T) (bottom panel) worms highlighting the difference in phasmid cilia lengths between

the two genotypes. On average, control *osm-6p::gfp* worms have longer phasmid cilia (top

panel, white bracket) than *osm-6p::gfp; sas-6(L69T)* worms (bottom panel, white bracket). White

brackets denote the cilia. Scale bar = 10 µm. D) Quantification of phasmid cilia lengths of wild-

type (*osm-6p::gfp*) and *sas-6(L69T)* (*osm-6p::gfp; sas-6(L69T)*) worms. Each circle represents

the length of a single phasmid cilium. n=number of phasmid cilia analyzed for each genotype.

Phasmid cilia length was reduced from 5.82 µm in control worms (n=54) to 3.99 µm in *osm-*

6p::gfp; sas-6(L69T) worms (n=59). p<0.0001, unpaired 2-tailed t-test. The error bars represent

the standard deviation and the middle bar represents the mean.

Figure 5. Phasmid cilia morphology is defective in a subset of the *sas-6(L69T)* mutant

worms. A) Cartoon showing the organization of a typical phasmid neuron with a ciliary

axoneme (A) built on top the transition zone (TZ). The dendrite of the phasmid neuron connects

to the TZ. B) and C) Representative Z-projection confocal images of phasmid cilia morphology

in *osm-6p::gfp* and *osm-6p::gfp; sas-6(L69T)* worms, respectively. B) *osm-6p::gfp* worms have a

stereotypical phasmid cilia morphology with two prominent TZs at the base and ciliary

axonemes projecting from the TZs. Scale bar = 10 µm. C) A subset of the *osm-6p::gfp; sas-*

6(L69T) worms exhibit deformed phasmid cilia morphologies. Scale bar = 10 µm. D)

Quantification of B) and C). While all of the wild-type (*osm-6p::gfp)* phasmid neurons (n=61)

exhibit a normal phasmid cilia morphology, 15.4% of the analyzed *osm-6p::gfp; sas-6(L69T)*

phasmid neurons show a deformed cilia morphology (n=65). n=number of phasmid neurons

analyzed. p=0.0014, Fisher's exact test.

Figure 6. Dendrite length is shorter in *osm-6p::gfp;sas-6(L69T)* phasmid neurons. A)

Representative Z-projection confocal image of the average phasmid dendrite length in *osm-*

6p::gfp worms. Scale bar = 10 µm. B) Representative Z-projection confocal image of the

average phasmid dendrite length in *osm-6p::gfp;sas-6(L69T)* worms. Scale bar = 10 µm. C)

Quantification of A) and B). The average dendrite length of control (*osm-6p::GFP*) worms is 37.4

µm (n=38) while that of *osm-6p::gfp;sas-6(L69T)* worms is 28.1 µm (n=42). n= number of

phasmid dendrites analyzed. p<0.0001, unpaired 2-tailed t-test. The error bars represent the

standard deviation and the middle bar represents the mean.

Figure 7. The *sas-6(L69T)* mutation reduces *C. elegans* chemotaxis to 1:500 Butanone. A)

Schematic of chemotaxis assay experiments. L4 stage *C. elegans* were plated onto unseeded

NGM plates that had been divided into four quadrants, two with solvent (ethanol) and two with

odorant (1:500 butanone in ethanol). The worms were placed in the center of the plate and

allowed to move around for 1 hour at 25°C. After 1 hour, the number of worms in each quadrant

were counted and quantified. The formula C.I. = (# animals in two odorant quadrants)/(#

animals in any of the four quadrants) was used to compute the Chemotaxis Index (C.I.). Worms

on any line or within the origin were not counted. Four plates were used for each genotype per

experiment. Assays were repeated in triplicate. B) Quantification of data obtained from A). The

sas-6(L69T) mutants exhibit a decreased chemotaxis towards 1:500 butanone as compared

with control worms. n= Number of worms whose chemotaxis towards 1:500 butanone was

analyzed. p<0.0001, unpaired 2-tailed t-test. C) Summary of data from A) The average

chemotaxis index of *sas-6(L69T)* worms was 0.3307. This chemotaxis index was less than the

average chemotaxis index of control worms which was 0.4107. C.I.=Chemotaxis Index.

Figure 8. The L69T mutation does not significantly alter the overall structure of the *C.*

***elegans* SAS-6 protein.** A) Cartoon depicting the location of the L69T mutation within the *C.*

elegans SAS-6 protein oriented with respect to known SAS-6 protein domains and SAS-6 amino

acid residues that are important for the interaction of SAS-6 with either itself or with other known

interactors (van Breugel et al. 2011, Kitagawa et al. 2011, Qiao et al. 2013, Hilbert *et al,* 2013,

Lettman et al. 2013). B) AlphaFold structure prediction of the *C. elegans* SAS-6 protein (PDB

accession number: 4GFC) that was crystallized in Hilbert et al. 2013. C) The *C. elegans* SAS-

6(L69T) protein structure was modeled using AlphaFold2. The coloring for the models are

based on the ChimeraX default palette-option where lipophilicity ranges from -20 to 20 going

from cyan (least lipophilic) to golden (most lipophilic). Blue arrows: A deep hydrophobic pocket

within the mutant SAS-6(L69T) protein is decreased by the L69T mutation.

Supplementary Figures

Figure S1. SAS-6 protein levels are slightly reduced in the presence of the *sas-6(L69T)*

mutation in a *zyg-1(it25)* background but are unaffected in a wild-type background. A) A

representative western blot image of an HA blot showing SAS-6 levels in *zyg-1(it25);sas-6::ha*

and *zyg-1(iti25);sas-6(L69T)::ha* worms. Tubulin served as a loading control. SAS-6 protein

levels are slightly decreased in the presence of the *sas-6(L69T)* mutation in a *zyg-1(it25)*

background B) Quantification of A) from four independent experiments. On average, SAS-6

protein levels are reduced by 15% in the presence of the *sas-6(L69T)* mutation in the *zyg-1(it25)*

background. P=0.0475, unpaired two-tailed t-test. C) A representative western blot image of a

HA blot showing SAS-6 levels in *sas-6::ha* and *sas-6(L69T)::ha* worms. Tubulin was used as a

loading control. Although SAS-6 protein levels tend to decrease in the presence of the *sas-*

6(L69T) mutation in a wild-type background, this decrease is not statistically significant

(p=0.1398, unpaired two-tailed t-test).

Data Availability Statement

719   All the strains made and used in this study will be made available upon request.

720   **Supplementary Table S1** lists all the strains made in this study.

Acknowledgments

We would like to thank the Sengupta lab for providing the PY6100 strain that was used for cilia

analysis. We would like to thank the O'Connell lab for providing the OC14 and OC869 strains

that have been used in this study. The wild-type N2 Bristol strain was provided by the CGC,

which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). We would

like to acknowledge Peter Nunan and Mariah Moreland for their technical assistance. Adam

Norris received funding from NIH R35GM133461. Finally, we would like to thank Drs. Steve

Caplan and Nina Peel for their critical comments on this manuscript.

Funding

The authors would like to acknowledge the National Institute of General Medical Sciences

(NIGMS) of the National Institutes of Health (NIH) for grant support 5P20GM103636-09,

University of Tulsa (TU) Start-up funding, the TU Faculty Summer Development Fellowship and

Shark Tank Grant awarded to Jyoti Iyer. Amy Smith, Mary Bergwell, Ellie Smith, Carter Dierlam,

Ramon Duran and Rory Seidel received funding and research support from the TU Chemistry

Summer Undergraduate Research Program and/or from the Tulsa Undergraduate Research

Challenge program. Ellie Smith and Carter Dierlam also received financial support from the

NIGMS grant 5P20GM103636-09 awarded to Jyoti Iyer.

Conflict of Interest

The authors declare no competing interests.

References

Abràmoff MD, Magalhães PJ, Ram SJ. Image processing with ImageJ. Biophotonics

international. 2004;11(7):36-42.

Alcantara D, O'Driscoll M. Congenital microcephaly. Am J Med Genet C Semin Med

Genet. 2014;166(2):124-139.

Arribere, JA, Bell, RT, Fu, BX, Artiles, KL, Hartman, PS, Fire, AZ. Efficient marker-free recovery

of custom genetic modifications with CRISPR/Cas9 in Caenorhabditis elegans. Genet.

2014;198(3),837-846.

Bargmann CI, Hartwieg E, Horvitz HR. Odorant-selective genes and neurons mediate

olfaction in C. elegans. Cell. 1993;74(3):515-27.

Bayer EA, Sun H, Rafi I, Hobert O. Temporal, spatial, sexual and environmental

regulation of the master regulator of sexual differentiation in C. elegans. Curr Biol.

2020;30(18):3604-16.

Bergwell M, Smith A, Lakin H, Slay R, Iyer J. Generation of sas-6:: ha by CRISPR/Cas9

editing. MicroPubl Biol. 2019;2019.

Bettencourt-Dias M, Rodrigues-Martins A, Carpenter L, Riparbelli M, Lehmann L, Gatt

MK, Carmo N, Balloux F, Callaini G, Glover DM. SAK/PLK4 is required for centriole

duplication and flagella development. Curr Biol. 2005;15(24):2199-207.

Blomen VA, Májek P, Jae LT, Bigenzahn JW, Nieuwenhuis J, Staring J, Sacco R, van

Diemen FR, Olk N, Stukalov A, et al. Gene essentiality and synthetic lethality in haploid

760   human cells. Science. 2015;350(6264):1092-6.

761   Broto P, Moreau G, Vandycke C. Molecular structures: perception, autocorrelation descriptor

762   and sar studies: system of atomic contributions for the calculation of the n-octanol/water

763   partition coefficients. Eur J Med Chem. 1984;19(1):71-8.

764   Collet J, Spike CA, Lundquist EA, Shaw JE, Herman RK. Analysis of osm-6, a gene that

765   affects sensory cilium structure and sensory neuron function in Caenorhabditis elegans.

766   Genetics. 1998 Jan 1;148(1):187-200.

767   Dammermann A, Müller-Reichert T, Pelletier L, Habermann B, Desai A, Oegema K.

768   Centriole assembly requires both centriolar and pericentriolar material proteins. Dev

769   Cell. 2004;7(6):815-29.

770   Ding W, Wu Q, Sun L, Pan NC, Wang X. Cenpj regulates cilia disassembly and

771   neurogenesis in the developing mouse cortex. J Neurosci. 2019;39(11):1994-2010.

772   Drummond IA. Cilia functions in development. Curr Opin Cell Biol. 2012;24(1):24-30.

773   Farooq M, Lindbæk L, Krogh N, Doganli C, Keller C, Mönnich M, Gonçalves AB,

774   Sakthivel S, Mang Y, Fatima A, et al. RRP7A links primary microcephaly to dysfunction

775   of ribosome biogenesis, resorption of primary cilia, and neurogenesis. Nat Commun.

776   2020;11(1):1-6.

777   Gabriel E, Wason A, Ramani A, Gooi LM, Keller P, Pozniakovsky A, Poser I, Noack F,

778   Telugu NS, Calegari F, et al. CPAP promotes timely cilium disassembly to maintain

779   neural progenitor pool. EMBO J. 2016;35(8):803-19.

780   Habedanck R, Stierhof YD, Wilkinson CJ, Nigg EA. The Polo kinase Plk4 functions in

781   centriole duplication. Nat Cell Biol. 2005;7(11):1140-6.

782   Hilbert M, Erat MC, Hachet V, Guichard P, Blank ID, Flückiger I, Slater L, Lowe ED,

783  GN, Steinmetz MO, et al. Caenorhabditis elegans centriolar protein SAS-6 forms a spiral

784   that is consistent with imparting a ninefold symmetry. Proc Natl Acad Sci U S A.

785   2013;110(28):11373-8.

786 Inglis PN, Ou G, Leroux MR, Scholey JM. The sensory cilia of Caenorhabditis elegans.

787   WormBook, ed. The C. elegans Research Community. Wormbook. 2007;1-22.

788 Iyer J, DeVaul N, Hansen T, Nebenfuehr B. Using microinjection to generate genetically

789   modified Caenorhabditis elegans by CRISPR/Cas9 Editing. In Microinjection. Humana

790   Press, New York, NY. 2019;431-457.

791 Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates

792   R, Žídek A, Potapenko A, et al. Highly accurate protein structure prediction with

793   AlphaFold. Nature. 2021 Aug;596(7873):583-9.

794 Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, Kanapin A, Le Bot N,

795   Moreno S, Sohrmann M, et al. Systematic functional analysis of the Caenorhabditis

796   elegans genome using RNAi. Nature. 2003;421(6920):231-7.

797 Kemp CA, Song MH, Addepalli MK, Hunter G, O'Connell K. Suppressors of zyg-1 define

798   regulators of centrosome duplication and nuclear association in Caenorhabditis elegans.

799   Genetics. 2007;176(1):95-113.

800 Kemphues KJ, Kusch M, Wolf N. Maternal-effect lethal mutations on linkage group II of

801   Caenorhabditis elegans. Genetics. 1988;120(4):977-86.

802 Khan MA, Rupp VM, Orpinell M, Hussain MS, Altmüller J, Steinmetz MO, Enzinger C,

803   Thiele H, Höhne W, Nürnberg G, Baig SM. A missense mutation in the PISA domain of

804   HsSAS-6 causes autosomal recessive primary microcephaly in a large consanguineous

805   Pakistani family. Hum Mol Genet. 2014;23(22):5940-9.

806 Kitagawa D, Vakonakis I, Olieric N, Hilbert M, Keller D, Olieric V, Bortfeld M, Erat MC, Flückiger

807   I, Gönczy P, et al. Structural basis of the 9-fold symmetry of centrioles. Cell.

808   2011;144(3):364-75.

Laguerre ML, Saux M, Dubost JP, Carpy A. MLPP: a program for the calculation of molecular

lipophilicity potential in proteins. Pharm. Sci. 1997;3.5–6: 217–222.

Leidel S, Delattre M, Cerutti L, Baumer K, Gönczy P. SAS-6 defines a protein family

required for centrosome duplication in C. elegans and in human cells. Nat Cell Biol.

2005;7(2):115-25.

Lettman MM, Wong YL, Viscardi V, Niessen S, Chen SH, Shiau AK, Zhou H, Desai A,

Oegema K. Direct binding of SAS-6 to ZYG-1 recruits SAS-6 to the mother

centriole for cartwheel assembly. Dev cell. 2013;25(3):284-98.

Levine MS, Bakker B, Boeckx B, Moyett J, Lu J, Vitre B, Spierings DC, Lansdorp PM,

Cleveland DW, Lambrechts D, et al. Centrosome amplification is sufficient to promote

spontaneous tumorigenesis in mammals. Dev cell. 2017;40(3):313-22.

Li W, Yi P, Zhu Z, Zhang X, Li W, Ou G. Centriole translocation and degeneration

during ciliogenesis in Caenorhabditis elegans neurons. EMBO J. 2017;36(17):2553-66.

Liang X, Calovich-Benne C, Norris A. Sensory neuron transcriptomes reveal complex

neuron-specific function and regulation of mec-2/Stomatin splicing. Nucleic Acids Res.

2022;50(5):2401-16.

Martin CA, Ahmad I, Klingseisen A, Hussain MS, Bicknell LS, Leitch A, Nürnberg G,

Toliat MR, Murray JE, Hunt D, et al. Mutations in PLK4, encoding a master regulator of

centriole biogenesis, cause microcephaly, growth failure and retinopathy. Nat Genet.

2014;46(12):1283-92.

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making

protein folding accessible to all. Nat Methods. 2022; 30:1-4.

Nechipurenko IV, Berciu C, Sengupta P, Nicastro D. Centriolar remodeling underlies

basal body maturation during ciliogenesis in Caenorhabditis elegans. Elife.

2017;6:e25686.

Nigg EA, Holland AJ. Once and only once: mechanisms of centriole duplication and

their deregulation in disease. Nat Rev Mol Cell Biol. 2018;19(5):297-312.

O'Rourke SM, Yochem J, Connolly AA, Price MH, Carter L, Lowry JB, Turnbull DW,

Kamps-Hughes N, Stiffler N, Miller MR, et al. Rapid mapping and identification of

mutations in Caenorhabditis elegans by restriction site-associated DNA mapping and

genomic interval pull-down sequencing. Genetics. 2011;189(3):767-78.

O'Connell KF, Caron C, Kopish KR, Hurd DD, Kemphues KJ, Li Y, White JG. The C.

elegans zyg-1 gene encodes a regulator of centrosome duplication with distinct maternal

and paternal roles in the embryo. Cell. 2001;105(4):547-58.

Paix A, Folkmann A, Rasoloson D, Seydoux G. High efficiency, homology-directed

genome editing in Caenorhabditis elegans using CRISPR-Cas9 ribonucleoprotein

complexes. Genetics. 2015;201(1):47-54.

Paix A, Schmidt H, Seydoux G. Cas9-assisted recombineering in C. elegans: genome

editing using in vivo assembly of linear DNAs. Nucleic Acids Res. 2016;44(15):e128.

Patwardhan D, Mani S, Passemard S, Gressens P, El Ghouzzi V. STIL balancing

primary microcephaly and cancer. Cell Death Dis. 2018;9(2):1-1.

Peel N, Iyer J, Naik A, Dougherty MP, Decker M, O'Connell KF. Protein phosphatase 1

down regulates ZYG-1 levels to limit centriole duplication. PLoS Genet.

2017;13(1):e1006543.

Perkins LA, Hedgecock EM, Thomson JN, Culotti JG. Mutant sensory cilia in the

nematode Caenorhabditis elegans. Dev Biol. 1986;117(2):456-87.

Phan TP, Holland AJ. Time is of the essence: the molecular mechanisms of primary

microcephaly. Genes Dev. 2021;35(23-24):1551-78.

Piano F, Schetter AJ, Mangone M, Stein L, Kemphues KJ. RNAi analysis of genes expressed in

the ovary of Caenorhabditis elegans. Curr Biol. 2000;10(24):1619-22.

Power KM, Akella JS, Gu A, Walsh JD, Bellotti S, Morash M, Zhang W, Ramadan YH,

Ross N, Golden A, et al. Mutation of NEKL-4/NEK10 and TTLL genes suppress neuronal

ciliary degeneration caused by loss of CCPP-1 deglutamylase function. PLoS Genet.

2020;16(10): e1009052.

Qiao R, Cabral G, Lettman MM, Dammermann A, Dong G. SAS-6 coiled-coil structure and

interaction with SAS-5 suggest a regulatory mechanism in C. elegans centriole

assembly. EMBO J. 2012;31(22):4334-47.

Serwas D, Su TY, Roessler M, Wang S, Dammermann A. Centrioles initiate cilia

assembly but are dispensable for maturation and maintenance in C. elegans. J Cell Biol.

2017;216(6):1659-71.

Shohayeb B, Ho U, Yeap YY, Parton RG, Millard SS, Xu Z, Piper M, Ng DC. The

association of microcephaly protein WDR62 with CPAP/IFT88 is required for cilia

formation and neocortical development. Hum Mol Genet. 2020;29(2):248-63.

Smith A, Bergwell M, Smith E, Mathew D, Iyer J. CRISPR/Cas9 Editing of the C.

elegans rbm-3.2 Gene using the dpy-10 Co-CRISPR Screening Marker and Assembled

Ribonucleoprotein Complexes. J Vis Exp. 2020;(166):e62001.

Song MH, Aravind L, Müller-Reichert T, O'Connell KF. The conserved protein SZY-20

opposes the Plk4-related kinase ZYG-1 to limit centrosome size. Dev Cell.

2008;15(6):901-12.

Song MH, Liu Y, Anderson DE, Jahng WJ, O'Connell KF. Protein phosphatase 2A-SUR-

6/B55 regulates centriole duplication in C. elegans by controlling the levels of

centriole assembly factors. Dev cell. 2011;20(4):563-71.

The C. elegans Deletion Mutant Consortium, et al. Large-scale screening for targeted

knockouts in the Caenorhabditis elegans genome. G3 (Bethesda). 2012;2(11):1415-25.

van Breugel M, Hirono M, Andreeva A, Yanagisawa HA, Yamaguchi S, Nakazawa Y, Morgner

N, Petrovich M, Ebong IO, Robinson CV, et al. Structures of SAS-6 suggest its

organization in centrioles. Science. 2011;331(6021):1196-9.

Wambach JA, Wegner DJ, Yang P, Shinawi M, Baldridge D, Betleja E, Shimony JS, Spencer D,

Hackett BP, Andrews MV, et al. Functional characterization of biallelic RTTN variants

identified in an infant with microcephaly, simplified gyral pattern, pontocerebellar

hypoplasia, and seizures. Pediatric research. 2018;84(3):435-41.

Wilsch-Bräuninger M, Huttner WB. Primary Cilia and Centrosomes in Neocortex

Development. Front Neurosci. 2021;15.

Wolf B, Balestra FR, Spahr A, Gönczy P. ZYG-1 promotes limited centriole amplification

in the C. elegans seam lineage. Dev Biol. 2018;434(2):221-30.

Xie S, Dierlam C, Smith E, Duran R, Williams A, Davis A, Mathew D, Naslavsky N, Iyer

J, Caplan S. The retromer complex regulates C. elegans development and mammalian

ciliogenesis. J Cell Sci. 2022;135(10):jcs259396.

Xu YS, Su ZY, Sun L, Yang XM, Sun SY, Ji XF, Ji YZ, Zhang SQ, Tian J, Wu QC.

Overexpression of the PLK4 Gene as a Novel Strategy for the Treatment of Autosomal

Recessive Microcephaly by Improving Centrosomal Dysfunction. J Mol Neurosci.

2021;71(12):2618-27.

Zaqout S, Morris-Rosendahl D, Kaindl AM. Autosomal recessive primary microcephaly

(MCPH): an update. Neuropediatrics. 2017;48(03):135-42.

Zhang W, Yang SL, Yang M, Herrlinger S, Shao Q, Collar JL, Fierro E, Shi Y, Liu A, Lu

H, et al. Modeling microcephaly with cerebral organoids reveals a WDR62–CEP170–

KIF2A pathway promoting cilium disassembly in neural progenitors. Nat Commun.

2019;10(1):1-4.

Figure 1

A

ECoRI site

CGATGATTTT GAATTCCTCTTTTTCAGAGACAAACG AACAACGAGA

TTG → ACG
Leu → Thr

B

Figure 2

A

n=31   n=23   n=12   n=12   n=21   n=16

B

n=40   n=23   n=16   n=16   n=36   n=21

Figure 3

A

zyg-1(it25) ✕ *sas-6(L69T)* → *zyg-1(it25);sas-6(L69T)*

Low embryonic lethality   No embryonic lethality   High embryonic lethality?

B

zyg-1(it25)

zyg-1(it25); sas-6(L69T)

C

% Normal centrosome duplication

n=52
82.7

n=58
27.6

zyg-1(it25)   *zyg-1(it25);sas-6(L69T)*

D

**** p<0.0001

Embryonic viability (%)

44.36%

7.14%

zyg-1(it25)   *zyg-1(it25);sas-6(L69T)*
n=18   n=18

Figure 4

A

osm-6p::gfp   X   *sas-6(L69T)*

F2 generation

osm-6p::gfp; sas-6(L69T)

B

C. elegans tail

Phasmid cilium

Phasmid dendrite

C

osm-6p::gfp

osm-6p::gfp; sas-6(L69T)

D

p<0.0001

****

5.82

3.99

Phasmid Cilia length (µm)

control
n=54

sas-6(L69T)
n=59

Figure 5

A

Transition zone (TZ)

Ciliary
Axoneme (A)

Dendrite

B

C

osm-6p::gfp

osm-6p::gfp; sas-6(L69T)

D

Phasmid cilia morphology defects

p=0.0014
**

Number of phasmid cilia

0%

15.4%

control
n=61

sas-6(L69T)
n=65

Normal cilia
Abnormal cilia

Figure 6

A

osm-6p::gfp

B

osm-6p::gfp;
sas-6(L69T)

C

Figure 7

A

Chemotaxis Assay

Chemotaxis index = (# animals in two odorant quadrants)/ (# animals in any of the four quadrants)

B

1:500 Butanone Response

C

Figure 8

A

SAS-5 interaction
E286,E287

N-terminal
Dimerization
I154

ZYG-1
interaction

L69*

46   98

C. elegans
SAS-6

PISA

1   Head domain   191   Coiled-coil   406 Tail 492

Residues involved in coiled-coil
dimer interactions: L36, S58,
F61, E62, L64, L172, I173, H175,
L176, L177, C180, A184

Residues involved in N-terminal
dimer interactions: D82, D83,
T84, V85, F90, V93, T152, P153,
I154, S155, K156, I159, F160

B

SAS-6 PDB 4GFC

C

SAS-6(L69T) modeled
using AlphaFold2

Figure S1