A primary microcephaly-associated *sas-6* mutation perturbs centrosome duplication and ciliogenesis in *Caenorhabditis elegans*

Mary Bergwell[1*], Amy Smith[2*], Ellie Smith[3], Ramon Duran[3], Erin Haastrup[3], Rebekah Napier-Jameson[4], Carter Dierlam[3], Rory Seidel[3], William Potter[3], Adam Norris[4], and Jyoti Iyer[3#]

* These authors contributed equally to this work

From [1]Oklahoma Medical Research Foundation, Oklahoma City, OK 73104, [2]Pfizer Inc., 875 Chesterfield Parkway West, Chesterfield, MO 63017, [3]Department of Chemistry and Biochemistry, University of Tulsa, Tulsa, OK 74104, [4]Southern Methodist University, PO Box 750100, Dallas, TX 75275

Running Head: *sas-6(L69T)* affects centrioles and cilia

[#]To whom correspondence should be addressed:

Jyoti Iyer: [3]Department of Chemistry and Biochemistry, University of Tulsa, Tulsa, OK 74104;

jgi2708@utulsa.edu; Tel. (918) 631-2070.

Keywords: SAS-6, SASS6, rare disease, microcephaly, centrosome, centriole, cilia, *C. elegans*

Abstract

A missense mutation within the human *SASS6* gene has been linked with the incidence of primary microcephaly in a Pakistani family. However, the mechanism by which this mutation causes disease is still unclear. SASS6 protein function is conserved between humans and *C. elegans.* Therefore, in the present study, we used *C. elegans* to model this primary microcephaly associated mutation to determine which molecular pathways are affected by this mutation in *C. elegans*. The human primary microcephaly-associated mutation corresponds to SAS-6(L69T) in *C. elegans*. Our studies revealed that both centrosome duplication and ciliogenesis are perturbed by the *sas-6(L69T)* mutation. Specifically, *C. elegans* carrying the *sas-6(L69T)* mutation exhibit an increased failure of centrosome duplication in a sensitized genetic background. Further, worms carrying this mutation also display shortened phasmid cilia, an abnormal phasmid cilia morphology, defective phasmid dendrite morphology and chemotaxis defects. Our data show that the centrosome duplication defects caused by this mutation are only uncovered in a sensitized genetic background, indicating that these defects are mild. However, the ciliogenesis defects caused by this mutation are evident in an otherwise wild-type background, indicating that they are stronger defects. Based on our findings, we propose that the ciliogenesis defects induced by this mutation are likely a more important contributing factor to the development of primary microcephaly in humans carrying this mutation than centrosome duplication defects. Thus, our studies shed light on a novel mechanism by which the *sas-6(L69T)* mutation could lead to the incidence of primary microcephaly in humans.

Introduction

Autosomal recessive primary microcephaly (MCPH) is a rare disease that causes a reduced brain size in infants along with a host of other physical and intellectual disabilities (Zaqout *et. al.*

2017). According to the OMIM database, twenty-nine microcephaly loci named MCPH1-MCPH29 have been identified thus far (https://www.omim.org/entry/251200). The current consensus in the field is that most mutations linked to the incidence of MCPH deplete the pool of neural progenitor cells during brain development. This results in fewer neurons being produced during embryonic development, thereby leading to a smaller brain size (Phan and Holland 2021). Although some MCPH-associated mutations have been previously modeled in *in vivo* model systems to understand how MCPH-associated mutations contribute to the disease pathology, there are still many MCPH-linked mutations that remain uninvestigated (Phan and Holland 2021). One such MCPH-associated mutation is a missense mutation within the *SASS6* gene (*sas-6* in *C. elegans*) (Khan *et al.* 2014).

SAS-6 protein function is conserved amongst several species including the multicellular eukaryotic nematode *C. elegans* and humans (Dammerman *et al.* 2004, Leidel *et al.* 2005). The *sas-6* gene was first identified to regulate centrosome duplication in *C. elegans* (Dammerman *et al.* 2004, Leidel *et al.* 2005). Centrosomes consist of a pair of cylindrical centrioles: a mother and a daughter centriole that are surrounded by pericentriolar material. Major functions of these organelles involve organizing a mitotic spindle and aiding in cilia assembly (Nigg and Holland, 2018). Proper cell division requires a precise duplication of centrosomes. There are two levels of regulation of centrosome duplication. First, it is critical that centrioles duplicate only once per cell cycle, and second, it is imperative that each mother centriole gives rise to a single daughter centriole. Defects in centrosome duplication can impair proper chromosome segregation, leading to genome instability and aneuploidy. Indeed, increasing centrosome numbers is sufficient to induce tumorigenesis in mice (Levine *et al.* 2017).

In addition to regulating centrosome biogenesis, SAS-6 function is also essential for cilia assembly in *C. elegans* (Li *et al.* 2017). Cilia are microtubule-based organelles that are critical for cell signaling and development (Drummond 2012). Primary cilia are the sole type of cilia that

are present in *C. elegans.* There are a total of sixty ciliated sensory neurons in *C. elegans* hermaphrodites*.* The cilia are present at the dendritic tips of the sensory neurons (Inglis *et al.* 2007). Although a previous study has shown that SAS-6 function is critical for proper ciliogenesis (Li *et al.* 2017), the mechanism by which SAS-6 promotes ciliogenesis is unknown. It has been especially difficult to study the role of SAS-6 in ciliogenesis because *sas-6* is an essential gene (Kamath *et al.* 2003, Blomen *et al.* 2015). Due to the important role of SAS-6 in regulating centrosome duplication, most mutations in *sas-6* yield embryonic lethality or sterility (Piano *et al.* 2000, Kamath *et al.* 2003, O'Rourke *et al.* 2011, Song *et al.* 2011, The C. elegans Deletion Mutant Consortium *et al.* 2012, Lettman *et al.* 2013).

In 2014, a study by Khan *et al.* identified a missense mutation in the human *SASS6* gene to be associated with the incidence of MCPH in a consanguineous Pakistani family (Khan *et al.* 2014). Ciliogenesis defects have been proposed as an important underlying cause of microcephaly (Alcantara *et al.* 2014, Gabriel *et al.* 2016, Zhang *et al.* 2019, Ding *et al.* 2019, Sohayeb *et al.* 2020, Farooq *et al.* 2020). However, the effect of this *SASS6* mutation on ciliogenesis has never been investigated. The effect of the MCPH-associated *SASS6* mutation on centrosome duplication in its endogenous context has also not been determined thus far. It is important to note that, SAS-6 levels in the cell are tightly regulated. An increase or a decrease in SAS-6 levels causes either centrosome overduplication or a failure of centrosome duplication, respectively (Dammerman *et al.* 2004, Leidel *et al.* 2005). Therefore, it is critical to model the MCPH-associated *SASS6* mutation in an endogenous context in a multicellular eukaryotic animal model. However, to date such a study has not been reported.

In this study, we used CRISPR/Cas9 genome editing to re-create the MCPH-associated *sas-6(L69T)* mutation in *C. elegans*. Our *in vivo* studies in the worm indicate that the *sas-6(L69T)* mutation does not result in embryonic lethality in a wild-type *C. elegans* background. However, introducing this mutation into a *zyg-1(it25)* strain that is partially compromised for

centrosome duplication results in a robust increase in embryonic lethality at the semi-permissive temperature. These data indicate that this MCPH-associated mutation mildly perturbs centrosome duplication in *C. elegans.* Surprisingly, while the centrosome duplication defects were only observed in a sensitized genetic background, ciliogenesis defects were observed even in a wild-type background in *C. elegans* carrying this mutation. Specifically, our studies revealed that the *sas-6(L69T)* mutation causes a reduction of the phasmid cilia length and results in an abnormal phasmid cilia and phasmid dendrite morphology. Thus, our studies indicate that the MCPH-associated *sas-6(L69T)* mutation affects ciliogenesis more severely than centrosome duplication in *C. elegans.* Our studies have also uncovered a novel role of the MCPH-associated *sas-6(L69T)* mutation in regulating ciliogenesis in the worm.

Materials and Methods

C. elegans maintenance

All *C. elegans* strains were grown at 20°C on OP50 *E. coli* bacteria seeded onto MYOB agar plates, unless otherwise indicated (as in the case of different assays). Supplementary table S1 lists all the *C. elegans* stains that have been generated and used in this study.

CRISPR/Cas9 genome editing

Assembled ribonucleoprotein complexes were microinjected into *C. elegans* gonads to perform CRISPR/Cas9 genome editing as described previously (Paix *et al.* 2015, Paix *et al.* 2016, Iyer *et al.* 2019, Smith *et al.* 2020). Homozygosity of the *sas-6(L69T)* edit was confirmed by PCR followed by restriction digestion with EcoRI and by DNA sequencing. Supplementary table S2 lists the sequences of the guide RNAs and repair templates that were used to make the CRISPR strains that have been used in this study.

Brood count and embryonic viability assays

All brood count assays and embryonic viability assays were performed as described previously (Xie *et al.* 2022). For both brood count as well as embryonic viability assays, L4 stage *C. elegans* were plated at the respective indicated temperatures. For brood count assays, all the progeny that were produced by each single worm over the worm's reproductive lifespan were counted and quantified. For embryonic viability assays, the percentage of dead and live progeny produced by each worm was determined by counting. The percentage of viable progeny was then computed by dividing the number of live progeny by the total number of progeny and by multiplying this ratio by 100.

Confocal live imaging

To monitor centrosome duplication in *C. elegans* embryos, live imaging was performed using an Olympus Spinning-Disk confocal microscope mounted with a Yokagawa scanhead. The cellSens software was used for image acquisition and processing. To prepare the *C. elegans* strains for imaging, L4 stage *C. elegans* were plated at 21.8°C overnight. The next day, gravid adult worms were picked into 8 to 12 µl of egg buffer (118 mM NaCl, 47 mM KCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$, 5 mM HEPES pH 7.4) on a square glass coverslip and dissected near the vulva using a needle to release embryos. The coverslip was carefully mounted on a 2% agarose pad (2% agarose LE in molecular biology grade water). The edges of the coverslip were sealed with molten Vaseline using a paintbrush. These slides were subjected to time-lapse imaging by taking 1.25 µm Z-stacks over an interval of approximately 132 seconds. Maximum intensity Z-projections were computed using the cellSens software and converted into an AVI format for making videos. The brightness and contrast were adjusted for the control and *sas-6(L69T)* videos for better non-quantitative video visualization.

Genetic crosses

To introduce the *sas-6(L69T)* mutation into the *zyg-1(it25)* background, first, males were generated from the *zyg-1(it25)* strain using alcohol treatment. Specifically, L4 stage

hermaphrodites from the *zyg-1(it25)* strain were incubated with 10% ethanol for 30 minutes, washed thrice with M9 buffer and allowed to recover for an hour. After recovery, the worms were plated onto MYOB plates and incubated at 20°C until males appeared. Following this, *zyg-1(it25)* homozygous males were mated with *sas-6(L69T)* L4 hermaphrodites. The F1 progeny from this cross were screened for heterozygosity of the *sas-6(L69T)* mutation using PCR and restriction digestion. Cross progeny were identified and screened for homozygosity of both edits using PCR and restriction digestion (for homozygosity screening of the *sas-6(L69T)* mutation) and 24°C temperature shifts (for homozygosity screening of the *zyg-1(it25)* mutation). For the cilia imaging experiments, PY6100 males *(oyIs59 Is[osm-6p::gfp] III)* were crossed with hermaphrodites from the IYR001 (*sas-6(luv1) IV*) and IYR002 (*sas-6(luv2) IV*) strains to produce strains expressing either SAS-6::HA (control) or SAS-6(L69T)::HA along with a pan-ciliary GFP marker. F1 progeny were screened for heterozygosity at both loci by screening F1 hermaphrodites for the presence of GFP fluorescence in the phasmid neurons. F2 homozygotes for *sas-6::ha* were identified as described previously (Bergwell *et al.* 2019) by PCR and digestion by NdeI followed by agarose gel electrophoresis. F2 homozygotes for *sas-6(L69T)::ha* were identified by PCR and digestion by EcoRI followed by agarose gel electrophoresis.

C. elegans phasmid cilia and dendrite imaging

C. elegans phasmid cilia were imaged as described previously (Xie *et al.* 2022). Briefly, L4 stage *C. elegans* from IYR026 (*oyIs59 Is[osm-6p::gfp] III; sas-6(luv1) IV*) and IYR027 (*oyIs59 Is[osm-6p::gfp] III; sas-6(luv2) IV*) strains were grown at 16°C overnight . The next day, young adult worms were anaesthetized using 10 mM Levamisole, mounted on 2% agarose pads and imaged within 30 minutes using an Olympus IX83 Spinning Disk confocal microscope coupled with a Yokogawa X1 Scan head (Olympus America Inc., Waltham, MA). 0.25 µm Z-slices were taken from the top to the bottom of each phasmid neuron using the cellSens software (Olympus America Inc., Waltham, MA) and maximum-intensity Z-projections were used for the cilia length,

cilia morphology and dendrite morphology analysis. All images were taken at the same laser intensity and exposure time.

Phasmid cilia length quantification

Phasmid cilia lengths were quantified as described previously (Xie *et al*. 2022). Briefly, the maximum intensity Z-projections of the phasmid cilia were used for measuring cilia lengths using the cellSens software (Olympus America Inc.). The shorter of the two phasmid cilia was used for quantifying phasmid cilia length since their lengths were more consistent than the lengths of the longer phasmid cilia. Images in which we were unable to delineate at which position the cilia began or ended from the Z-projections were excluded from our analysis.

Criteria used for quantifying phasmid cilia and dendrite morphology defects

Phasmid cilia that did not exhibit the canonical phasmid cilia structure with two basal bodies and cilia projecting from the basal bodies were quantified as having an abnormal phasmid cilia morphology. Specifically, instances where the basal bodies could not be clearly distinguished from the ciliary axoneme were classified as being abnormal. In the case of phasmid dendrite defects, dendrites that exhibited an uninterrupted filling of the dendrite with GFP were classified as having a normal dendrite morphology. On the other hand, phasmid dendrites that exhibited a discontinuous GFP fluorescence within the dendrite with aggregates of GFP evident along the dendrites were classified as having an abnormal dendrite morphology.

***C. elegans* chemotaxis assay**

Olfaction assays were performed as previously described in Liang *et al*. 2022.

Briefly, *C. elegans* were maintained by standard techniques on NGM agar plates seeded with OP50 *Escherichia coli* at 20°C. Worms were synchronized to L4 stage. Worms were washed off the plates using M9 buffer, allowed to gravity settle, supernatant removed, and then washed a further 4 times. Unseeded NGM plates (50 mM NaCl, 1.7% agar, 0.25% peptone, 1 mM $CaCl_2$, 0.013 mM Cholesterol, 1 mM $MgSO_4$, 25 mM KPO_4) were divided into four quadrants, two with solvent and two with odorant. Ethanol was used as a solvent. A 1:500 dilution of the odorant

butanone was made in ethanol. The Chemotaxis Index (CI) was calculated using the formula CI = (# animals in two odorant quadrants)/(# animals in any of the four quadrants). Four plates were used per genotype per experiment. Assays were repeated 3 times.

Whole worm extracts for western blotting

Whole worm extracts for western blotting were prepared as described previously (Xie *et al.* 2022). Briefly, 100 gravid adults from each genotype were washed with M9 buffer twice. The M9 buffer was removed and the worm pellet was resuspended in 40 µl of 4X SDS PAGE sample buffer, boiled for 10 minutes at 95°C and stored at -30°C until further use.

Western blotting

Western blotting was performed using the Bio-Rad wet-transfer method as described previously (Xie *et al.* 2022). Briefly, for western blotting of whole worm extracts, 10 to 14 µl of extracts were loaded onto each well of a 7.5% SDS-PAGE gel. The gels were run until proper band separation was achieved and the protein was transferred onto a 0.2 µm nitrocellulose membrane using the wet-transfer method (Bio-Rad Laboratories). The membranes were blocked with Odyssey Blocking Buffer (LiCOR Biosciences, Inc.), probed with anti-HA (Cell Signaling, catalog C29F4 Rabbit mAb #3724) and anti-tubulin (Santa Cruz Biotechnology, Inc., catalog sc-32293) antibodies overnight at 4°C. The membranes were washed thrice with 1X TBST (0.1% w/v Tween-20, 50 mM Tris-Cl, pH 7.6, 150 mM NaCl) and incubated with IRDye® 800CW Donkey anti-Rabbit IgG secondary antibody (LiCOR Biosciences, Inc. catalog # 926-32213) and IRDye® 680RD Goat anti-Mouse IgG secondary antibody (LiCOR Biosciences, Inc. catalog # 926-68070) for 1 hour at room temperature. The membranes were washed again thrice with 1XTBST and imaged using the LI-COR Odyssey CLx imager (LI-COR Biosciences, Inc.). The band intensities were adjusted equally for all the gel lanes for better data visualization and presentation. Supplementary table S3 lists all the antibodies that have been used in this study.

Western blot quantification and presentation

The band intensities across the entire membrane were uniformly adjusted using the LiCOR Image Studio software (LI-COR Biosciences, Inc.) and saved in a .tiff format for performing quantitative analysis of the bands. Quantitative western blotting was performed using ImageJ (Abràmoff *et al.* 2004). Briefly, the band intensities for SAS-6::HA were determined using ImageJ and normalized with respect to the loading control Tubulin. ImageJ was also used to rotate and crop representative gel images presented in **Figure S1.**

Modeling SAS-6(L69T) using Alphafold2

The N-terminal coiled-coil dimer construct of *C. elegans* centriolar protein SAS-6 was crystallized by Hilbert *et al.* 2013. The ceSAS-constructs required for crystallization involved the removal of the α2-β5 loop (Δ103-130) which was reported to not affect the N-terminal domain structure within a C_α rmsds of ~0.3 Angstroms. The crystallographic data was deposited in the Research Collaboration for Structural Bioinformatics databank under accession number PDB ID code 4G79.

The native SAS-6 structure (AF-062479-F1) available from the AlphaFold Protein Structure Database (Jumper *et al.* 2021; https://AlphaFold.ebi.ac.uk/) has leucine at position 69 at the end of the β_4 sheet just prior to the asparagine-asparagine which initiates the α_1 helix occurring at positions 70 to 82 (using the Hilbert *et al.* 2013 author numbering system). It should be noted that the residue numbering system reported for ceSAS-6 in Uniprot (062479) has two additional amino acids (glycine-serine) at the N-terminus and thus the sequence is offset and the α_1 helix starts at position 72 in this numbering system. The reported per-residue confidence score (pLDDT) for residue leucine 69 was 92.37 suggesting that the AlphaFold model confidence level was very high for this residue.

The SAS-6(L69T) mutant protein structure was modeled by loading the modified sequence into the Google Colaboratory (Colab) services using ColabFold: AlphaFold2 using

MMseqs2 as described by Mirdita *et al.* 2022, and available through the AlphaFold2.ipynb

v2.1.0. with the suite of public notebooks available through the deepmind/AlphaFold github.

The 4G79 and the AlphaFold models were uploaded into the program ChimeraX

(https://www.cgl.ucsf.edu/chimerax/) developed from the Resource for Biocomputing,

Visualization, and Informatics for Visualization (RBVI), University of California, San Francisco.

The Molecular Lipophilicity Potential (MLP) is a construct that spreads atomic values out

in 3D. The MLP coloring on the molecular surface is based on pyMLP

(https://www.rbvi.ucsf.edu/chimerax/docs/user/commands/mlp.html) as originally developed by

Broto *et al.* 1984 and Laguerre *et al.* 1997 and as modified

https://www.cgl.ucsf.edu/chimera/data/mlp-sep2016/index.html and was used without any color

scale modifications.

Statistical analysis

Graphpad Prism (GraphPad Software, Inc.) was used for performing statistical analysis of the

data. Unpaired two-tailed t-tests were performed to determine statistical significance of the data

for the brood counts, western blot assays, phasmid cilia lengths and chemotaxis assays. The

Fisher's exact test was performed to determine if the phasmid cilia dendritic defects and

phasmid cilia morphology defects were statistically significant. A p-value of less than 0.05 was

considered attain statistical significance.

Results

An isoleucine to threonine substitution of SASS6 was reported to be linked with the occurrence

of MCPH in a Pakistani family (Khan *et al.* 2014). As shown previously by Khan *et al.*, alignment

of the amino acid sequences of the SAS-6 protein of different species showed that the MCPH-

associated amino acid of human SASS6 is remarkably conserved among several species

including *C. elegans* (Khan *et al.* 2014). Notably, the *sas-6* gene was first identified in *C.

elegans* and was subsequently shown to function similarly in humans (Dammerman *et al.* 2004,

Leidel *et al.* 2005). Due to the high functional conservation of SAS-6, we determined that *C. elegans* would be an appropriate eukaryotic model to thoroughly investigate the consequences of this mutation. The human SASS6(I62T) mutation corresponds to SAS-6(L69T) in *C. elegans.*

The *sas-6(L69T)* mutation was successfully introduced into *C. elegans* by CRISPR/Cas9 editing

Using CRISPR/Cas9 genome editing, we generated a leucine to threonine substitution at amino acid 69 of the *C. elegans* SAS-6 protein by editing the endogenous *sas-6* gene of *C. elegans.* This strain was named IYR002 and the corresponding allele was named *sas-6(luv2).* For ease of understanding, this strain has been referred to as the *sas-6(L69T)* strain and the corresponding MCPH-associated mutation has been referred to as the *sas-6(L69T)* mutation throughout this manuscript. The repair template used for introducing the *sas-6(L69T)* mutation carried an EcoRI restriction site **(Figure 1A)**. As shown in **Figure 1B**, upon PCR amplification and restriction digestion of the *sas-6* gene with EcoRI, several homozygous edited worms were obtained in the F2 generation. The incorporation of the *sas-6(L69T)* mutation was confirmed by DNA sequencing **(Figures 1C, 1D)**.

The microcephaly-associated *sas-6(L69T)* mutation does not affect brood size or embryonic viability in a wild-type background

The function of the *sas-6* gene is essential for centrosome duplication (Dammerman *et al.* 2004, Leidel *et al.* 2005). Proper centrosome duplication is required for accurate embryonic divisions in *C. elegans.* Since the *sas-6(L69T)* mutation has a pathological relevance in humans, we proposed that this mutation could impair SAS-6 function in regulating centrosome duplication, thereby leading to defective centrosome duplication in *C. elegans.* This could, in turn, affect *C. elegans* brood size and embryonic viability. Therefore, to investigate the effect of the *sas-6(L69T)* mutation on centrosome duplication, brood counts and embryonic viability

assays were performed using *C. elegans* carrying this mutation. Surprisingly, the introduction of the *sas-6(L69T)* mutation did not yield any major effect on either brood size or embryonic viability at 16°C, 20°C and 25°C **(Figures 2A and 2B)**. These data indicate that the *sas-6(L69T)* mutation likely has a mild effect on SAS-6 function in regulating centrosome duplication which cannot be uncovered in an otherwise wild-type background.

The *sas-6(L69T)* mutation increases centrosome duplication failures and embryonic lethality in a sensitized *zyg-1(it25)* background

ZYG-1 is a functional homolog of human Plk4 and a master kinase whose activity is critical for proper centrosome duplication (O'Connell *et al.* 2001, Habedanck *et al.* 2005, Bettencourt-Dias *et al.* 2005). Decreasing ZYG-1 protein levels causes a failure of centrosome duplication while elevating them causes centrosome overduplication (O'Connell *et al.* 2001, Peel *et al.* 2017, Wolf *et al.* 2018). The temperature-sensitive *zyg-1(it25)* allele has a proline to leucine substitution within the ZYG-1 cryptic polo box which impairs ZYG-1 function at the restrictive temperature, leading to a failure of proper centrosome duplication (Kemphues *et al.* 1988, Kemp *et al.* 2007). By controlling the temperature at which the *zyg-1(it25)* strain is grown, the rate of centrosome duplication failure in this strain can be controlled. For example, at restrictive temperatures of 24°C and above, the *zyg-1(it25)* strain produces 100% dead eggs due to a complete failure of centrosome duplication (Kemp *et al.* 2007). However, at lower semi-permissive temperatures (temperatures below 24°C), centrosome duplication still partially occurs in the *zyg-1(it25)* strain, yielding varying levels of embryonic viability with a temperature of 20°C yielding between 93.7% to 98.7% embryonic viability (Song *et al.* 2008). To uncover mild defects in SAS-6 function in regulating centrosome duplication in the *sas-6(L69T)* mutant worms, the *sas-6(L69T)* mutation was introduced into a *zyg-1(it25)* strain expressing fluorescently tagged centrosome and chromosome markers to visualize centrosome duplication. We reasoned that if the *sas-6(L69T)* mutation mildly affects SAS-6 function in regulating

centrosome duplication, combining it with the *zyg-1(it25)* mutation at the semi-permissive temperature would yield higher levels of centrosome duplication failures and increased embryonic lethality than what each mutant produces on its own. The schematic and rationale for introducing the *sas-6(L69T)* mutation into the *zyg-1(it25)* strain is shown in **Figure 3A**.

Upon introducing the *sas-6(L69T)* mutation into a *zyg-1(it25)* strain expressing fluorescent centrosome and chromosome markers, centrosome duplication in the *zyg-1(it25)* and *zyg-1(it25);sas-6(L69T)* worms was assessed at the semi-permissive temperature of 21.8°C **(Figures 3B and 3C)**. Notably, introducing the *sas-6(L69T)* mutation greatly increased the centrosome duplication failures exhibited by the *zyg-1(it25)* mutants at 21.8°C **(Figure 3C)**. Specifically, in the *zyg-1(it25)* single mutant alone, about 83% of the centrosomes analyzed duplicated normally at the semi-permissive temperature of 21.8°C (n=52). However, interestingly, in the *sas-6(L69T);zyg-1(it25)* double mutant, proper centrosome duplication was reduced to only about 28% (n=58). Since accurate centrosome duplication is required for embryonic viability, we next assessed whether embryonic viability of the *zyg-1(it25)* strain was also affected in the presence of the *sas-6(L69T)* mutation.

The embryonic viability of the *zyg-1(it25)* strain expressing fluorescent centrosome and DNA markers alone at the semi-permissive temperature of 21.7°C was about 44% (number of worms=18; number of progeny= 1103). However, introducing the *sas-6(L69T)* mutation into this background reduced embryonic viability to only about 7% (number of worms=18; number of progeny= 912; unpaired two-tailed t-test p<0.0001) **(Figure 3D)**. These data collectively indicate that the *sas-6(L69T)* mutation mildly inhibits SAS-6 function in regulating centrosome duplication, thereby resulting in a failure of proper centrosome duplication and increased embryonic lethality in the *zyg-1(it25)* background.

Ciliogenesis is perturbed in *sas-6(L69T)* mutant *C. elegans*

Ciliogenesis defects have been linked with the incidence of microcephaly (Alcantara *et al.* 2014, Gabriel *et al.* 2016, Wambach *et al.* 2018, Zhang *et al.* 2019, Ding *et al.* 2019, Sohayeb *et al.* 2020, Farooq *et al.* 2020). Defective ciliogenesis is thought to result in cell cycle arrest in neural progenitors during brain development. The loss of neural progenitors due to either cell-cycle or apoptosis is proposed to be an underlying cause for the incidence of MCPH (Phan and Holland 2021). The effect of the *sas-6(L69T)* mutation on ciliogenesis is currently unknown. Since ciliogenesis defects have been previously linked with MCPH, we questioned whether worms carrying the *sas-6(L69T)* mutation exhibit any ciliogenesis defects. Preliminary dye-filling assays did not reveal any severe dye-filling defects between control and *sas-6(L69T)* worms. Specifically, we did not observe any worm that completely excluded DiI in their phasmid neurons in either control or *sas-6(L69T)* worms upon dye-filling. Our preliminary experiments revealed that 100% of the control (n=23) and *sas-6(L69T)* (n=25) worms exhibited dye-filling with DiI. Nevertheless, although dye-filling assays are a quick and easy method to visualize cilia defects, a lack of a dye-filling defect phenotype does not preclude the presence of ciliary defects.

To directly determine whether the *sas-6(L69T)* strain exhibits ciliary defects, we obtained a PY6100 strain that expresses a pan-ciliary marker. This strain expresses a GFP transcriptional reporter for OSM-6, a protein that is critical for proper ciliogenesis (Perkins *et al.* 1986, Collet *et al.* 1998). Thus, the introduction of *osm-6p::GFP* marks *C. elegans* ciliated neurons with GFP, allowing for easy visualization of the cilia (Bayer *et al.* 2020, Xie *et al.* 2022).

Since the *sas-6(L69T)* strain also has an HA tag on the mutant SAS-6 protein, we first crossed the PY6100 strain with a stain expressing SAS-6::HA to account for the potential effect of the HA-tag on SAS-6 function in regulating ciliogenesis. For convenience, we refer to this control strain as the *osm-6p::GFP* strain. We introduced the *sas-6(L69T)* mutation into the PY6100 strain by genetic crossing. The experimental design for introducing the *osm-6p::GFP*

into the *sas-6(L69T)* strain is shown in **Figure 4A.** The introduction of *osm-6p::GFP* into the *sas-6(L69T)* strain by genetic crossing was confirmed by both confocal microscopy and by genotyping using PCR.

To investigate whether *sas-6(L69T)* mutant worms exhibit defects in ciliogenesis, we examined the cilia of the phasmid neurons which are present in the posterior region of the worms. Intriguingly, upon measuring the length of the phasmid neuronal cilia, we found a statistically significant decrease in ciliary length in the *sas-6(L69T)* mutant worms (unpaired two-tailed t-test, p=0.0047). Specifically, control worms had a phasmid cilia length of about 5.8 μm (n=26) while the *sas-6(L69T)* worms had shorter cilia with a phasmid cilia length of only 4.71 μm (n=35). These data indicate that the *sas-6(L69T)* mutation indeed perturbs ciliogenesis in *C. elegans.*

Phasmid cilia and dendrite morphology are defective in the *sas-6(L69T)* mutant worms

While analyzing cilia length of *sas-6(L69T)* worms, we found that many of our cilia images for this strain were unusable due to exhibiting a disfigured cilia morphology as compared with control worms. Our analysis revealed that 100% of the phasmid cilia of control worms (n=63) showed a normal phasmid cilia morphology with two distinct basal bodies and a ciliary axoneme projecting from each basal body **(Figure 5A, 5C)**. However, 15 out of 66 (22.7%) of the *sas-6(L69T)* phasmid cilia exhibited an abnormal morphology **(Figures 5B, 5C).** This abnormal phasmid cilia morphology defect of the *sas-6(L69T)* worms was found to be statistically significant via a Fisher's exact test (p<0.0001). These data indicate that the *sas-6(L69T)* mutation perturbs phasmid cilia morphology in a subset of the phasmid cilia.

In addition to the phasmid cilia morphology defect, the *sas-6(L69T)* worms also displayed an abnormal phasmid dendrite morphology. Specifically, 60 out of 63 phasmid dendrites (95.2%) analyzed from control worms showed an even GFP fluorescence throughout

the dendrites **(Figures 6A, 6D).** However, 30 out of 66 (45.5%) of the phasmid dendrites analyzed from the *sas-6(L69T)* worms exhibited an uneven GFP fluorescence **(Figures 6B, 6C, 6D)** with aggregates of GFP present either throughout the dendrite or in certain regions of the dendrite. This abnormal phasmid dendrite morphology defect of the *sas-6(L69T)* worms was found to be statistically significant via a Fisher's exact test (p<0.0001). On the basis of these data, we propose that the *sas-6(L69T)* mutation likely causes structural defects in some of the phasmid dendrites. Electron microscopy analysis of these phasmid dendrites should be performed in the future to determine if this is indeed the case.

Chemotaxis is defective in the *sas-6(L69T)* mutant worms

Since defects in ciliogenesis in the *C. elegans* sensory neurons often affect worm behaviors, we investigated the effect of the *sas-6(L69T)* mutation on chemotaxis to the chemoattractant butanone. The experimental scheme for the chemotaxis assays is shown in **Figure 7A**. Our studies demonstrated that *C. elegans* carrying the *sas-6(L69T)* mutation exhibited a reduced attraction to 1:500 butanone as compared with control worms (two-tailed unpaired t-test, p<0.0001) **(Figures 7B, 7C)**. Specifically, our data showed that the average chemotaxis index of control worms was 0.4107. In comparison, the average chemotaxis index of *sas-6(L69T)* worms was reduced to 0.3307 **(Figure 7C)**. These data indicate that defective ciliogenesis in the *sas-6(L69T)* mutant worms likely causes a defective *C. elegans* chemotaxis response to 1:500 butanone. Hence, SAS-6 function in regulating ciliogenesis is likely to be important for mediating certain worm behaviors such as chemotaxis.

SAS-6 protein levels are altered in the presence of the *sas-6(L69T)* mutation in a *zyg-1(it25)* background but not in a wild-type background

Some of the other known MCPH-associated mutations in centrosome duplication genes such as STIL and PLK4 have been reported to affect their protein levels (Martin *et al.* 2014,

Patwardhan *et al.* 2018, Xu *et al.* 2021). The effect of *sas-6(L69T)* mutation on SAS-6 protein levels is unknown. Therefore, we questioned whether the *sas-6(L69T)* mutation has any effect on SAS-6 protein levels. To address this, western blotting was performed on whole worm lysates from *sas-6::ha* (control) vs *sas-6(L69T)::ha* worms and from *zyg-1(it25);sas-6::ha* (control) vs *zyg-1(it25);sas-6(L69T)::ha* worms. Western blot analysis indicated that SAS-6 levels decrease on average by 15% in the presence of the *sas-6(L69T)* mutation in the *zyg-1(it25)* background (n=4; p=0.0475, unpaired two-tailed t-test) **(Figure S1A, S1B)**. However, the decrease in SAS-6 protein levels in the presence of the *sas-6(L69T)* mutation in an otherwise wild-type background was not found to be statistically significant (n=4; p=0.1398, unpaired two-tailed t-test) **(Figures S1C, S1D)**. Since we observe ciliogenesis defects in the presence of the *sas-6(L69T)* mutation even in a wild-type background **(Figures 4B, 4C, 4D, 5A, 5B, 5C)**, we propose that decreased SAS-6 levels are likely not sufficient to explain the effects of the *sas-6(L69T)* mutation.

Modeling SAS-6(L69T) using AlphaFold suggests that the leucine to threonine substitution at amino acid 69 does not have a major effect on SAS-6 protein structure

The L69T mutation is located within the conserved PISA motif of SAS-6 **(Figure 8A)**. Further, this amino acid does not have a known involvement in the dimerization of SAS-6 coiled-coil region, in the oligomerization of SAS-6 N-terminal head domains, in the interaction of SAS-6 with ZYG-1 or in the interaction of SAS-6 with SAS-5 (van Breugel *et al.* 2011, Kitagawa *et al.* 2011, Qiao *et al.* 2013, Hilbert *et al,* 2013, Lettman *et al.* 2013) **(Figure 8A)**. To elucidate the possible mechanism by which the *sas-6(L69T)* mutation affects centrosome duplication and ciliogenesis, we used AlphaFold to predict how the L69T mutation could alter the structure of the SAS-6 protein **(Figures 8B, 8C)**. Figures 8B and 8C show the comparison of the SAS-6 PDB model 4GFC from Hilbert *et al.* 2013 **(Figure 8B)** compared with the AlphaFold-derived SAS-6(L69T) model **(Figure 8C)** using the ChimeraX program. Our analysis indicates that the

overall structural differences due to the L69T mutation in SAS-6 are minor. Therefore, we propose that centriole architecture may not be significantly affected in the *sas-6(L69T)* mutant worms. This is supported by our data that wild-type *C. elegans* carrying homozygous copies of the *sas-6(L69T)* mutation are close to 100% viable at all temperatures **(Figure 2B)**. However, the ChimeraX hydrophobicity surface model suggests that a deep hydrophobic pocket that is found within the SAS-6 protein is reduced due to the MCPH-associated L69T mutation **(Figure 8B)**. The function of this hydrophobic pocket within SAS-6 has not yet been elucidated.

Discussion

Through our studies, we have re-created the MCPH-associated *sas-6(L69T)* mutation in *C. elegans* by CRISPR to generate the first *C. elegans* model of a MCPH-associated mutation. A previous study investigated the effects of the human SASS6(I62T) mutation (the human equivalent of the *sas-6(L69T)* mutation) on centrosome duplication in U2OS cells (Khan *et al.* 2014). In their studies, the authors noted an inhibition of centrosome duplication in U2OS cells expressing the SASS6(I62T) protein from exogenous plasmid constructs carrying the SASS6(I62T) mutation. The levels of SAS-6 in the cell must be tightly regulated for its proper cellular function. Either overexpressing the SAS-6 protein or depleting it can affect proper centrosome duplication (Dammermann *et al.* 2004, Leidel *et al.* 2005). Therefore, to investigate the effect of the *sas-6(L69T)* mutation on centrosome duplication in an endogenous context, we monitored centrosome duplication in our *sas-6(L69T)* CRISPR worm model. Interestingly, *C. elegans* carrying this mutation appeared phenotypically wild-type and did not exhibit any defects in embryonic viability or brood size **(Figure 2)**. *C. elegans* embryonic divisions are reliant on proper centrosome duplication. Since we did not detect any embryonic lethality in the presence of the *sas-6(L69T)* mutation **(Figure 2B)**, we concluded that centrosome duplication is likely to be relatively unaffected in the *sas-6(L69T)* mutant worms in a wild-type background.

To uncover the potential milder effects of this mutation on centrosome duplication that may not be visible in a wild-type genetic background, we introduced this mutation in the sensitized *zyg-1(it25)* background. Interestingly, the *sas-6(L69T)* mutation significantly enhanced the centrosome duplication failures and embryonic lethality of *zyg-(it25)* worms **(Figures 3B, 3C, 3D)**. These data demonstrate that the MCPH-associated *sas-6(L69T)* mutation does, in fact, reduce SAS-6 function in regulating centrosome duplication. However, this effect can only be revealed in a sensitized genetic background. Thus, at least in the context of this MCPH-associated mutation, centrosome duplication is only mildly affected by this mutation. Future studies will involve elucidating the molecular mechanism by which the *sas-6(L69T)* mutation inhibits centrosome duplication.

Our data that the *sas-6(L69T)* mutation likely has a mild effect on SAS-6 function in regulating centrosome duplication are consistent with the observed phenotypic effects of the *SASS6(I62T)* mutation in human MCPH patients. SASS6 is an essential gene that is required for proper centrosome duplication and cell division (Kamath *et al.* 2003, Dammerman *et al.* 2004, Leidel *et al.* 2005, Blomen *et al.* 2015). If the *SASS6(I62T)* mutation severely inhibited SASS6 function in regulating centrosome duplication, it would be expected to result in embryonic lethality. As a result, there would not be human patients living with homozygous copies of the *SASS6(I62T)* mutation. However, the fact that there are *SASS(I62T)* homozygous individuals in the population (Khan *et al.* 2014) indicates that this mutation does not severely perturb SASS6 role in regulating centrosome duplication in humans. Therefore, our data in *C. elegans* which demonstrate that embryonic viability is unaffected in the presence of this mutation **(Figure 2B)** are in line with the observations in human MCPH patients carrying this mutation.

The effect of the *SASS6(I62T)* mutation on ciliogenesis has never been investigated. Upon introducing the *sas-6(L69T)* mutation into a pan-ciliary marker expressing *C. elegans*

strain, we determined that ciliogenesis is perturbed in the *sas-6(L69T)* mutant worms **(Figures 4B, 4C, 4D, 5A, 5B, 5C)**. Specifically, the *sas-6(L69T)* worms have shorter phasmid cilia and a defective phasmid cilia morphology **(Figures 4B, 4C, 4D, 5A, 5B, 5C)**. Ciliogenesis defects have been associated with the incidence of microcephaly (Alcantara *et al.* 2014, Gabriel *et al.* 2016, Wambach *et al.* 2018, Zhang *et al.* 2019, Ding *et al.* 2019, Sohayeb *et al.* 2020, Farooq *et al.* 2020). Interestingly, in a study by Wambach *et al.,* the authors cultured MCPH patient derived fibroblasts and found that these fibroblasts had shorter primary cilia than those of fibroblasts that were cultured from a healthy control individual (Wambach *et al.* 2018). In another study, Shohayeb *et al.* determined that mouse embryonic fibroblasts that were derived from WDR62 mutant mice carrying CRISPR-edited MCPH patient mutations exhibited a shortened ciliary length, like we report here (Shohayeb *et al.* 2019). Thus, there is precedence for shorter ciliary length being associated with MCPH.

We find that in *C. elegans* carrying the *sas-6(L69T)* mutation, centrosome duplication defects can only be detected in a sensitized *zyg-1(it25)* genetic background **(Figures 3B, 3C).** However, the ciliogenesis defects induced by this mutation can be uncovered even in a wild-type background **(Figures 4B, 4C, 4D, 5A, 5B, 5C)**. These findings lead us to speculate that perhaps, at least in the context of this specific MCPH-associated mutation, the inhibition of proper ciliogenesis may be a more important contributor to the development of MCPH than the inhibition of centrosome duplication. However, it is also possible that the centrosome duplication and ciliogenesis defects induced by this mutation synergize to cause in MCPH in human patients carrying this mutation. In the future, it will be interesting to determine if the *SASS6(I62T)* mutation also affects ciliogenesis in human cell lines. It will also be interesting to determine if other such MCPH-associated mutations likewise affect ciliogenesis in addition to affecting centrosome duplication.

Since *sas-6* is an essential gene, many *sas-6* mutant alleles are difficult to maintain and analyze (O'Rourke *et al.* 2011, Song *et al.* 2011, The C. elegans Deletion Mutant Consortium *et al.* 2012, Lettman *et al.* 2013). This has made it tedious to dissect the function of SAS-6 in ciliogenesis. A previous study expressed a *sas-6* sgRNA and Cas9 protein using a heat-shock promoter to demonstrate that SAS-6 plays a critical role in *C. elegans* ciliogenesis (Li *et al.* 2017). However, the mechanistic investigation of these defects has been challenging due to the high lethality of *sas-6* mutant alleles. Notably, our studies indicate that *sas-6(L69T)* mutant worms appear phenotypically wild-type and yet exhibit significant ciliogenesis defects. Therefore, we expect that our *sas-6(L69T)* CRISPR strain will be a useful tool to unravel the mechanism by which SAS-6 regulates ciliogenesis. In the future, forward genetic screens can be performed with the *sas-6(L69T)* allele to identify additional genes that may genetically interact with SAS-6 to regulate ciliogenesis in the worm.

Our data indicate that *C. elegans* harboring the *sas-6(L69T)* mutation also exhibit sensory neuronal defects such as defective dendrite morphology **(Figure 6)**. This is a surprising finding since *C. elegans* centrioles degenerate early during neuronal differentiation (Serwas *et al.* 2017, Nechipurenko *et al.* 2017). One possibility is that SAS-6 functions independently of its role as a core centriolar protein in regulating dendrite morphology in *C. elegans* sensory neurons. Another possibility is that proper basal body assembly may be required for proper phasmid neuron dendrite morphology through an unknown mechanism. In the future, transmission electron microscopy analysis of dendrites and cilia from *sas-6(L69T)* mutant worms should be performed to determine the nature of the dendritic and ciliary defects in these mutant worms. Future studies will also involve analyzing *C. elegans* strains expressing a variety of fluorescently-tagged ciliary markers (e.g. Intraflagellar transport markers, transition zone markers, basal body markers, etc.) to further dissect the mechanism by which the *sas-6(L69T)* mutation affects ciliogenesis.

Defective chemotaxis towards butanone was seen in *C. elegans* carrying the *sas-6(L69T)* mutation **(Figure 7).** Chemotaxis towards butanone is controlled by the AWC amphid sensory neuron (Bargmann *et al.* 1993). These data indicate that ciliogenesis may also be perturbed in the amphid sensory neurons in *sas-6(L69T)* mutant worms. In the present study, we have mainly focused our efforts in analyzing ciliogenesis in the *C. elegans* phasmid neurons. Future studies should be directed towards determining whether ciliogenesis in the amphid neurons is also affected in *C. elegans* carrying the *sas-6(L69T)* mutation.

Finally, modeling the L69T mutation using AlphaFold did not reveal any significant changes in the structure of the SAS-6 protein **(Figure 8B, 8C)**. This is not a surprising finding since *sas-6* is an essential gene and therefore severely affecting SAS-6 structure may result in embryonic lethality. The fact that there are live individuals within the population harboring the *SASS6(I62T)* mutation suggest that this mutation does not significantly alter SASS6 protein structure. However, our AlphaFold analysis revealed that a deep hydrophobic pocket within SAS-6 is altered by the *sas-6(L69T)* mutation. Presently, no known function has been ascribed to this hydrophobic pocket in SAS-6. Based upon our data, we are tempted to speculate that this SAS-6 hydrophobic pocket may mediate an as-yet unknown interaction of SAS-6 with a protein(s) that may be important for ciliogenesis and/or centrosome duplication.

As our *C. elegans* model of the *sas-6(L69T)* mutation recapitulates many of the effects seen in the human MCPH patients carrying this mutation, we expect that *C. elegans* will be more widely adopted as a model system to study the effects of additional MCPH-associated human mutations in the future.

Figure 1. The MCPH-associated *sas-6(L69T)* mutation was successfully generated in *C. elegans* by CRISPR/Cas9 editing A) Schematic of the CRISPR repair template used for inserting the leucine to threonine mutation at amino acid 69 of the *C. elegans* SAS-6 protein. B) Agarose gel electrophoresis of PCR products digested with EcoRI to screen for homozygous-

edited *C. elegans* carrying the *sas-6(L69T)* mutation. Red stars: Homozygous-edited worms; het: heterozygote; WT: wild-type. C) Protein alignment of sequenced SAS-6(L69T) homozygotes and wild-type SAS-6 protein using CLC sequence viewer showing that the leucine to threonine mutation was successfully introduced in the homozygous-edited worm line (Red arrow). D) Sequencing peaks from an identified *sas-6(L69T)* homozygous worm line demonstrating that the original codon UUG coding for leucine in wild-type worms was successfully changed to the codon ACG that codes for threonine.

Figure 2. The *sas-6(L69T)* mutation does not affect *C. elegans* brood size or embryonic viability A) Brood sizes of wild-type and *sas-6(L69T)* mutant worms were determined at temperatures of 16°C, 20°C and 25°C. No statistically significant difference in brood size was observed between wild-type and *sas-6(L69T)* worms at any of the assessed temperatures. n= number of worms whose brood was analyzed for each strain. n.s: Not significant. Error bars represent the standard deviation. The middle bar represents the mean. B) Embryonic viability of wild-type and *sas-6(L69T)* worms was analyzed at temperatures of 16°C, 20°C and 25°C. No major change in embryonic viability was observed between wild-type and *sas-6(L69T)* worms at any of the assessed temperatures. The viability of over 2000 embryos was analyzed for each strain at every temperature.

Figure 3. The *sas-6(L69T)* mutation increases the centrosome duplication failures and the embryonic lethality of *zyg-1(it25)* worms. A) Schematic showing the genetic cross that was performed to combine the *sas-6(L69T)* mutation with the *zyg-1(it25)* mutation. B) Stills from time-lapse imaging of *zyg-1(it25)* and *zyg-1(it25);sas-6(L69T)* strains expressing fluorescently-tagged centrosome and DNA markers. Red: mCherry-SPD-2 (centrosome marker); Green: GFP-histone (DNA marker). In a majority of *zyg-1(it25)* mutant embryos, centrosome duplication proceeds normally at a semi-permissive temperature of 21.8°C, yielding two centrosomes in each cell of a 2-cell *C. elegans* embryo (**Figure 3B, top panel**). On the contrary, most of the

zyg-1(it25);sas-6(L69T) mutant centrosomes fail to duplicate at 21.8°C resulting in a single centrosome present in each cell of a 2-cell *C. elegans* embryo (**Figure 3B, bottom panel**). C) Quantification of B). n=Number of centrosomes analyzed. Approximately 83% of the *zyg-1(it25)* centrosomes duplicate normally at 21.7°C. In contrast, only about 28% of the *zyg-1(it25);sas-6(L69T)* centrosomes duplicate normally at this temperature. D) Quantification of embryonic viability of *zyg-1(it25)* single and *zyg-1(it25);sas-6(L69T)* double mutants at the semi-permissive temperature of 21.7°C. Average embryonic viability is reduced from ~44% to ~7% in the presence of the *sas-6(L69T)* mutation. The viability of 1103 embryos from *zyg-1(it25)* worms and 912 embryos from *zyg-1(it25);sas-6(L69T)* worms was analyzed. Embryos from 18 worms were analyzed for each condition. Unpaired 2-tailed t-test p<0.0001. Error bars represent the standard deviation.

Figure 4. Phasmid cilia length is shorter in *sas-6(L69T)* mutant worms. A) A schematic of the genetic crossing experiment that was performed to introduce the MCPH-associated *sas-6(L69T)* mutation into the PY6100 pan-ciliary marker strain. B) and C): Z-projection confocal images of phasmid neurons from *osm-6p::gfp* and *osm-6p::gfp;sas-6(L69T)* worms highlighting the difference in phasmid cilia lengths between the two genotypes. On average, control *osm-6p::gfp* worms B) have longer phasmid cilia than *osm-6p::gfp;sas-6(L69T)* worms C). Red brackets denote the cilia. Scale bar: 10 μm D) Quantification of phasmid cilia lengths of wild-type (*osm-6p::gfp*) and *sas-6(L69T)* (*osm-6p::gfp;sas-6(L69T)*)) worms. Each circle represents the length of a single phasmid cilium. n=number of phasmid cilia analyzed for each genotype. Phasmid cilia length was reduced from 5.80 μm in wild-type worms to 4.71 μm in *sas-6(L69T)* worms. p=0.0047, unpaired 2-tailed t-test.

Figure 5. Phasmid cilia morphology is defective in a subset of the *sas-6(L69T)* mutant worms. A) and B) Representative Z-projection confocal images of phasmid cilia morphology in *osm-6p::gfp* and *osm-6p::gfp;sas-6(L69T)* worms, respectively. A) *osm-6p::gfp* worms have a

stereotypical phasmid cilia morphology with two prominent basal bodies at the base and ciliary axonemes projecting from the basal bodies. Scale bar=10 µm. B) Some *osm-6p::gfp;sas-6(L69T)* worms exhibit severely deformed phasmid cilia morphologies. Scale bar=10 µm. C) Quantification of A) and B). While all of the wild-type (*osm-6p::gfp)* phasmid neurons exhibit a normal phasmid cilia morphology, 22.7% of the analyzed *sas-6(L69T)* (*osm-6p::gfp;sas-6(L69T))* phasmid neurons show a deformed cilia morphology. n=number of phasmid neurons analyzed. p<0.0001, Fisher's exact test.

Figure 6. Dendrite morphology is defective in some of the *sas-6(L69T)* phasmid neurons.
A) Representative Z-projection confocal image of phasmid dendrite morphology in *osm-6p::gfp* worms. In *osm-6p::gfp* worms, an uninterrupted filling of the dendrite with GFP is observed in a majority of the phasmid dendrites. Scale bar=10 µm. B) and C): Representative Z-projection confocal images of phasmid dendrite morphologies in *osm-6p::gfp;sas-6(L69T)* worms. B) Some of the phasmid dendrites of worms carrying the *sas-6(L69T)* mutation display GFP aggregates along the length of the dendrite. Scale bar=10 µm. C) Some *osm-6p::gfp;sas-6(L69T)* worms exhibit severely defective phasmid dendrite morphologies with some parts of the dendrite almost completely excluding GFP fluorescence. Scale bar=10 µm. D) Quantification of A), B) and C). While 95.2% of the wild-type (*osm-6p::gfp*) phasmid neurons exhibit a phasmid dendrite morphology with GFP completely filling the phasmid dendrite, 45.5% of the analyzed *sas-6(L69T)* (*osm-6p::gfp;sas-6(L69T))* phasmid neurons show aggregates of GFP and/or an uneven filling of the phasmid dendrites with GFP. n=number of phasmid neurons analyzed. p<0.0001, Fisher's exact test.

Figure 7. The *sas-6(L69T)* mutation reduces *C. elegans* chemotaxis to 1:500 Butanone. A) Schematic of chemotaxis assay experiments. L4 stage *C. elegans* were plated onto unseeded NGM plates that had been divided into four quadrants, two with solvent and two with odorant. Ethanol was used as a solvent. A 1:500 dilution of the odorant butanone was made in ethanol.

The worms were placed in the center of the plate and allowed to move around for 1 hour at 25°C. After 1 hour, the number of worms in each quadrant were counted and quantified. The Chemotaxis Index (C.I.) was calculated using the formula C.I. = (# animals in two odorant quadrants)/(# animals in any of the four quadrants). Worms on any line or within the origin were not counted. Four plates were used per genotype per experiment. Assays were repeated 3 times. B) Quantification of data obtained from A). The *sas-6(L69T)* mutants exhibit a decreased chemotaxis towards 1:500 Butanone as compared with control worms. n= Number of worms whose chemotaxis towards 1:500 butanone was analyzed. p<0.0001, unpaired 2-tailed t-test. B) Summary of data from A) The average chemotaxis index of *sas-6(L69T)* worms was 0.3307. This chemotaxis index was less than the average chemotaxis index of control worms which was 0.4107. C.I.=Chemotaxis Index.

Figure 8. The L69T mutation does not significantly alter the overall structure of the *C. elegans* SAS-6 protein. A) Cartoon depicting the location of the L69T mutation within the *C. elegans* SAS-6 protein oriented with respect to known SAS-6 protein domains and SAS-6 amino acid residues that are important for the interaction of SAS-6 with either itself or with other known interactors. B) Alphafold structure prediction of the *C. elegans* SAS-6 protein (PDB accession number: 4GFC) that was crystallized in Hilbert *et al.* 2013. C) The *C. elegans* SAS-6(L69T) protein structure was modeled using Alphafold2. The coloring for the models are based on the ChimeraX default palette-option where lipophilicity ranges from -20 to 20 going from cyan (least lipophilic) to golden (most lipophilic). Red arrows: A deep hydrophobic pocket within the mutant SAS-6(L69T) protein is decreased by the leucine to threonine mutation.

Supplementary Figures

Figure S1. SAS-6 protein levels are slightly reduced in the presence of the *sas-6(L69T)* mutation in a *zyg-1(it25)* background but are unaffected in a wild-type background. A) A representative western blot image of an HA blot showing SAS-6 levels in *zyg-1(it25);sas-6::ha*

and *zyg-1(iti25);sas-6(L69T)::ha* worms. Tubulin served as a loading control. SAS-6 protein levels are slightly decreased in the presence of the *sas-6(L69T)* mutation in a *zyg-1(it25)* background B) Quantification of A) from four independent experiments. On average, SAS-6 protein levels are reduced by 15% in the presence of the *sas-6(L69T)* mutation in the *zyg-1(it25)* background. p=0.0475, unpaired two-tailed t-test. C) A representative western blot image of a HA blot showing SAS-6 levels in *sas-6::ha* and *sas-6(L69T)::ha* worms. Tubulin was used as a loading control. Although SAS-6 protein levels tend to decrease in the presence of the *sas-6(L69T)* mutation in a wild-type background, this decrease is not statistically significant (p=0.1398, unpaired two-tailed t-test).

Data Availability Statement

All the strains made and used in this study will be made available upon request. Table S1 lists all the strains made in this study.

Acknowledgments

We would like to thank the Sengupta lab for providing the PY6100 strain that was used for cilia analysis. We would like to thank the O'Connell lab for providing the OC14 and OC779 strains that have been used in this study. The wild-type N2 Bristol strain was provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). We would like to acknowledge Peter Nunan for his technical assistance. Adam Norris received funding from NIH R35GM133461. Finally, we would like to thank Dr. Steve Caplan for critical comments and discussions regarding our manuscript.

Funding

The authors would like to acknowledge the National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health (NIH) for grant support 5P20GM103636, University of Tulsa (TU) Start-up funding, the TU Faculty Summer Development Fellowship and Shark

Tank Grant awarded to Jyoti Iyer. Amy Smith, Mary Bergwell, Ramon Duran, Ellie Smith and Rory Seidel received funding and research support from the TU Chemistry Summer Undergraduate Research Program and/or from the Tulsa Undergraduate Research Challenge program.

Conflict of Interest

The authors declare no competing interests.

References

Abràmoff MD, Magalhães PJ, Ram SJ. Image processing with ImageJ. Biophotonics international. 2004;11(7):36-42.

Alcantara D, O'Driscoll M. Congenital microcephaly. Am J Med Genet C Semin Med Genet. 2014;166(2):124-139.

Bargmann CI, Hartwieg E, Horvitz HR. Odorant-selective genes and neurons mediate olfaction in C. elegans. Cell. 1993;74(3):515-27.

Bayer EA, Sun H, Rafi I, Hobert O. Temporal, spatial, sexual and environmental regulation of the master regulator of sexual differentiation in C. elegans. Curr Biol. 2020;30(18):3604-16.

Bergwell M, Smith A, Lakin H, Slay R, Iyer J. Generation of sas-6:: ha by CRISPR/Cas9 editing. MicroPubl Biol. 2019;2019.

Bettencourt-Dias M, Rodrigues-Martins A, Carpenter L, Riparbelli M, Lehmann L, Gatt MK, Carmo N, Balloux F, Callaini G, Glover DM. SAK/PLK4 is required for centriole duplication and flagella development. Curr Biol. 2005;15(24):2199-207.

Blomen VA, Májek P, Jae LT, Bigenzahn JW, Nieuwenhuis J, Staring J, Sacco R, van Diemen FR, Olk N, Stukalov A, *et al.* Gene essentiality and synthetic lethality in haploid human cells. Science. 2015;350(6264):1092-6.

Broto P, Moreau G, Vandycke C. Molecular structures: perception, autocorrelation descriptor and sar studies: system of atomic contributions for the calculation of the n-octanol/water partition coefficients. Eur J Med Chem. 1984;19(1):71-8.

Collet J, Spike CA, Lundquist EA, Shaw JE, Herman RK. Analysis of osm-6, a gene that affects sensory cilium structure and sensory neuron function in Caenorhabditis elegans. Genetics. 1998 Jan 1;148(1):187-200.

Dammermann A, Müller-Reichert T, Pelletier L, Habermann B, Desai A, Oegema K. Centriole assembly requires both centriolar and pericentriolar material proteins. Dev Cell. 2004;7(6):815-29.

Ding W, Wu Q, Sun L, Pan NC, Wang X. Cenpj regulates cilia disassembly and neurogenesis in the developing mouse cortex. J Neurosci. 2019 ;39(11):1994-2010.

Drummond IA. Cilia functions in development. Curr Opin Cell Biol. 2012;24(1):24-30.

Farooq M, Lindbæk L, Krogh N, Doganli C, Keller C, Mönnich M, Gonçalves AB, Sakthivel S, Mang Y, Fatima A, *et al*. RRP7A links primary microcephaly to dysfunction of ribosome biogenesis, resorption of primary cilia, and neurogenesis. Nat Commun. 2020;11(1):1-6.

Gabriel E, Wason A, Ramani A, Gooi LM, Keller P, Pozniakovsky A, Poser I, Noack F, Telugu NS, Calegari F, *et al*. CPAP promotes timely cilium disassembly to maintain neural progenitor pool. EMBO J. 2016;35(8):803-19.

Habedanck R, Stierhof YD, Wilkinson CJ, Nigg EA. The Polo kinase Plk4 functions in centriole duplication. Nat Cell Biol. 2005;7(11):1140-6.

Hilbert M, Erat MC, Hachet V, Guichard P, Blank ID, Flückiger I, Slater L, Lowe ED, GN, Steinmetz MO, *et al*. Caenorhabditis elegans centriolar protein SAS-6 forms a spiral that is consistent with imparting a ninefold symmetry. Proc Natl Acad Sci U S A. 2013;110(28):11373-8.

Inglis PN, Ou G, Leroux MR, Scholey JM. The sensory cilia of Caenorhabditis elegans. WormBook, ed. The C. elegans Research Community. Wormbook. 2007;1-22.

Iyer J, DeVaul N, Hansen T, Nebenfuehr B. Using microinjection to generate genetically modified Caenorhabditis elegans by CRISPR/Cas9 Editing. In Microinjection. Humana Press, New York, NY. 2019;431-457.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, *et al*. Highly accurate protein structure prediction with AlphaFold. Nature. 2021 Aug;596(7873):583-9.

Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, Kanapin A, Le Bot N, Moreno S, Sohrmann M, *et al*. Systematic functional analysis of the Caenorhabditis elegans genome using RNAi. Nature. 2003;421(6920):231-7.

Kemp CA, Song MH, Addepalli MK, Hunter G, O'Connell K. Suppressors of zyg-1 define regulators of centrosome duplication and nuclear association in Caenorhabditis elegans. Genetics. 2007;176(1):95-113.

Kemphues KJ, Kusch M, Wolf N. Maternal-effect lethal mutations on linkage group II of Caenorhabditis elegans. Genetics. 1988;120(4):977-86.

Khan MA, Rupp VM, Orpinell M, Hussain MS, Altmüller J, Steinmetz MO, Enzinger C, Thiele H, Höhne W, Nürnberg G, Baig SM. A missense mutation in the PISA domain of HsSAS-6 causes autosomal recessive primary microcephaly in a large consanguineous Pakistani family. Hum Mol Genet. 2014;23(22):5940-9.

Kitagawa D, Vakonakis I, Olieric N, Hilbert M, Keller D, Olieric V, Bortfeld M, Erat MC, Flückiger I, Gönczy P, *et al*. Structural basis of the 9-fold symmetry of centrioles. Cell. 2011;144(3):364-75.

Laguerre ML, Saux M, Dubost JP, Carpy A. MLPP: a program for the calculation of molecular lipophilicity potential in proteins. Pharm. Sci. 1997;3.5–6: 217–222.

Leidel S, Delattre M, Cerutti L, Baumer K, Gönczy P. SAS-6 defines a protein family

required for centrosome duplication in C. elegans and in human cells. Nat Cell Biol. 2005;7(2):115-25.

Lettman MM, Wong YL, Viscardi V, Niessen S, Chen SH, Shiau AK, Zhou H, Desai A, Oegema K. Direct binding of SAS-6 to ZYG-1 recruits SAS-6 to the mother centriole for cartwheel assembly. Dev cell. 2013;25(3):284-98.

Levine MS, Bakker B, Boeckx B, Moyett J, Lu J, Vitre B, Spierings DC, Lansdorp PM, Cleveland DW, Lambrechts D, *et al*. Centrosome amplification is sufficient to promote spontaneous tumorigenesis in mammals. Dev cell. 2017;40(3):313-22.

Li W, Yi P, Zhu Z, Zhang X, Li W, Ou G. Centriole translocation and degeneration during ciliogenesis in Caenorhabditis elegans neurons. EMBO J. 2017;36(17):2553-66.

Liang X, Calovich-Benne C, Norris A. Sensory neuron transcriptomes reveal complex neuron-specific function and regulation of mec-2/Stomatin splicing. Nucleic Acids Res. 2022;50(5):2401-16.

Martin CA, Ahmad I, Klingseisen A, Hussain MS, Bicknell LS, Leitch A, Nürnberg G, Toliat MR, Murray JE, Hunt D, *et al*. Mutations in PLK4, encoding a master regulator of centriole biogenesis, cause microcephaly, growth failure and retinopathy. Nat Genet. 2014;46(12):1283-92.

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making protein folding accessible to all. Nat Methods. 2022; 30:1-4.

Nechipurenko IV, Berciu C, Sengupta P, Nicastro D. Centriolar remodeling underlies basal body maturation during ciliogenesis in Caenorhabditis elegans. Elife. 2017;6:e25686.

Nigg EA, Holland AJ. Once and only once: mechanisms of centriole duplication and their deregulation in disease. Nat Rev Mol Cell Biol. 2018;19(5):297-312.

O'Rourke SM, Yochem J, Connolly AA, Price MH, Carter L, Lowry JB, Turnbull DW,

Kamps-Hughes N, Stiffler N, Miller MR, *et al*. Rapid mapping and identification of
mutations in Caenorhabditis elegans by restriction site-associated DNA mapping and
genomic interval pull-down sequencing. Genetics. 2011;189(3):767-78.

O'Connell KF, Caron C, Kopish KR, Hurd DD, Kemphues KJ, Li Y, White JG. The C.
elegans zyg-1 gene encodes a regulator of centrosome duplication with distinct maternal
and paternal roles in the embryo. Cell. 2001;105(4):547-58.

Paix A, Folkmann A, Rasoloson D, Seydoux G. High efficiency, homology-directed
genome editing in Caenorhabditis elegans using CRISPR-Cas9 ribonucleoprotein
complexes. Genetics. 2015;201(1):47-54.

Paix A, Schmidt H, Seydoux G. Cas9-assisted recombineering in C. elegans: genome
editing using in vivo assembly of linear DNAs. Nucleic Acids Res. 2016;44(15):e128.

Patwardhan D, Mani S, Passemard S, Gressens P, El Ghouzzi V. STIL balancing
primary microcephaly and cancer. Cell Death Dis. 2018;9(2):1-1.

Peel N, Iyer J, Naik A, Dougherty MP, Decker M, O'Connell KF. Protein phosphatase 1
down regulates ZYG-1 levels to limit centriole duplication. PLoS Genet.
2017;13(1):e1006543.

Perkins LA, Hedgecock EM, Thomson JN, Culotti JG. Mutant sensory cilia in the
nematode Caenorhabditis elegans. Dev Biol. 1986;117(2):456-87.

Phan TP, Holland AJ. Time is of the essence: the molecular mechanisms of primary
microcephaly. Genes Dev. 2021;35(23-24):1551-78.

Piano F, Schetter AJ, Mangone M, Stein L, Kemphues KJ. RNAi analysis of genes expressed in
the ovary of Caenorhabditis elegans. Curr Biol. 2000;10(24):1619-22.

Qiao R, Cabral G, Lettman MM, Dammermann A, Dong G. SAS-6 coiled-coil structure and
interaction with SAS-5 suggest a regulatory mechanism in C. elegans centriole
assembly. EMBO J. 2012;31(22):4334-47.

Serwas D, Su TY, Roessler M, Wang S, Dammermann A. Centrioles initiate cilia

assembly but are dispensable for maturation and maintenance in C. elegans. J Cell Biol. 2017;216(6):1659-71.

Shohayeb B, Ho U, Yeap YY, Parton RG, Millard SS, Xu Z, Piper M, Ng DC. The association of microcephaly protein WDR62 with CPAP/IFT88 is required for cilia formation and neocortical development. Hum Mol Genet. 2020;29(2):248-63.

Smith A, Bergwell M, Smith E, Mathew D, Iyer J. CRISPR/Cas9 Editing of the C. elegans rbm-3.2 Gene using the dpy-10 Co-CRISPR Screening Marker and Assembled Ribonucleoprotein Complexes. J Vis Exp. 2020;(166):e62001.

Song MH, Aravind L, Müller-Reichert T, O'Connell KF. The conserved protein SZY-20 opposes the Plk4-related kinase ZYG-1 to limit centrosome size. Dev Cell. 2008;15(6):901-12.

Song MH, Liu Y, Anderson DE, Jahng WJ, O'Connell KF. Protein phosphatase 2A-SUR-6/B55 regulates centriole duplication in C. elegans by controlling the levels of centriole assembly factors. Dev cell. 2011;20(4):563-71.

The C. elegans Deletion Mutant Consortium, *et al*. Large-scale screening for targeted knockouts in the Caenorhabditis elegans genome. G3 (Bethesda). 2012;2(11):1415-25.

van Breugel M, Hirono M, Andreeva A, Yanagisawa HA, Yamaguchi S, Nakazawa Y, Morgner N, Petrovich M, Ebong IO, Robinson CV, *et al*. Structures of SAS-6 suggest its organization in centrioles. Science. 2011;331(6021):1196-9.

Wambach JA, Wegner DJ, Yang P, Shinawi M, Baldridge D, Betleja E, Shimony JS, Spencer D, Hackett BP, Andrews MV, *et al*. Functional characterization of biallelic RTTN variants identified in an infant with microcephaly, simplified gyral pattern, pontocerebellar hypoplasia, and seizures. Pediatric research. 2018;84(3):435-41.

Wilsch-Bräuninger M, Huttner WB. Primary Cilia and Centrosomes in Neocortex Development. Front Neurosci. 2021;15.

Wolf B, Balestra FR, Spahr A, Gönczy P. ZYG-1 promotes limited centriole amplification

in the C. elegans seam lineage. Dev Biol. 2018;434(2):221-30.

Xie S, Dierlam C, Smith E, Duran R, Williams A, Davis A, Mathew D, Naslavsky N, Iyer
J, Caplan S. The retromer complex regulates C. elegans development and mammalian
ciliogenesis. J Cell Sci. 2022;135(10):jcs259396.

Xu YS, Su ZY, Sun L, Yang XM, Sun SY, Ji XF, Ji YZ, Zhang SQ, Tian J, Wu QC.
Overexpression of the PLK4 Gene as a Novel Strategy for the Treatment of Autosomal
Recessive Microcephaly by Improving Centrosomal Dysfunction. J Mol Neurosci.
2021;71(12):2618-27.

Zaqout S, Morris-Rosendahl D, Kaindl AM. Autosomal recessive primary microcephaly
(MCPH): an update. Neuropediatrics. 2017;48(03):135-42.

Zhang W, Yang SL, Yang M, Herrlinger S, Shao Q, Collar JL, Fierro E, Shi Y, Liu A, Lu
H, *et al.* Modeling microcephaly with cerebral organoids reveals a WDR62–CEP170–
KIF2A pathway promoting cilium disassembly in neural progenitors. Nat Commun.
2019;10(1):1-4.

Figure 1

A

TTG → ACG
Leu → Thr

B

C

D

TTG → ACG
Leu → Thr

Figure 2

A

n=31   n=23   n=12   n=12   n=21   n=16

B

n=40   n=23   n=16   n=16   n=36   n=21

Figure 3

A

zyg-1(it25)   *sas-6(L69T)*   *zyg-1(it25);sas-6(L69T)*

Low embryonic   No embryonic   High embryonic
lethality   lethality   lethality?

B

zyg-1(it25)

zyg-1(it25);sas-6(L69T)

C

n=52
82.7

n=58
27.6

% Normal centrosome duplication

zyg-1(it25)   *zyg-1(it25);sas-6(L69T)*

D

**** p<0.0001

44.36%

7.14%

Embryonic viability (%)

zyg-1(it25)   *zyg-1(it25);sas-6(L69T)*
n=18   n=18

Figure 4

A

osm-6p::gfp   X   *sas-6(L69T)*

F2 generation

osm-6p::gfp; sas-6(L69T)

B   **C**   **D**

osm-6p::gfp   *osm-6p::gfp; sas-6(L69T)*

Figure 5

A

B

osm-6p::gfp

osm-6p::gfp; sas-6(L69T)

C

Phasmid cilia morphology defects

p<0.0001
****

0%   22.7%

Number of ciliated neurons

Normal cilia
Abnormal cilia

wlld-type   *sas-6(L69T)*
n=63   n=66

Figure 6

A

B

C

osm-6p::gfp   osm-6p::gfp; sas-6(L69T)   osm-6p::gfp; sas-6(L69T)

D

Figure 7

A

Chemotaxis Assay

Chemotaxis index = (# animals in two odorant quadrants)/ (# animals in any of the four quadrants)

B

C

1:500 Butanone Response

Figure 8

A

SAS-5 interaction
E286,E287

N-terminal
Oligomerization
I154

ZYG-1
interaction

L69*

46   98

C. elegans
SAS-6

1   Head domain   191   Coiled-coil   406   Tail   492

PISA

Residues involved in coiled-coil
dimer interactions: L36, S58,
F61, E62, L64, L172, I173, H175,
L176, L177, C180, A184

Residues involved in N-terminal
dimer interactions: D82, D83,
T84, V85, F90, V93, T152, P153,
I154, S155, K156, I159, F160

B

C

SAS-6 PDB 4GFC

SAS-6(L69T) modeled using
alphafold

Figure S1

A

B

C

D

Table S1

STRAIN	GENOTYPE	METHOD	RESOURCE
N2	WT	N/A	CGC
PY6100	*oyIs59 Is[osm-6p::gfp] III*	N/A	Piali Sengupta Lab
OC14	*zyg-1(it25) II*	N/A	Kevin O'Connell Lab
			Kemphues and Wolf, 1988
OC779	*bsSi15 [pKO109: spd-2p::spd-2::*	N/A	Kevin O'Connell Lab
	mCherry::spd-2 3'-utr, unc-119(+)] I;		
	bsSi30[pCW9: ZC504.3p::sfgfp::		
	his-58::ZC504.3 3'-utr, unc-119(+)] II		
IYR001	*sas-6(luv1[sas-6::ha]) IV*	CRISPR	Bergwell *et al.* 2019
IYR002	*sas-6(luv2[(sas-6(L69T)::ha)] IV*	CRISPR	This study
IYR003	*zyg-1(it25) II; sas-6(luv1) IV*	Crossed IYR001	This study
		with OC14	
IYR004	*zyg-1(it25) II; sas-6(luv2) IV*	Crossed IYR002	This study
		with OC14	
IYR005	*bsSi15 [pKO109: spd-2p::spd-2::*	Crossed IYR001	This study
	mCherry::spd-2 3'-utr, unc-119(+)] I;	with OC779	
	bsSi30[pCW9: ZC504.3p::sfgfp::		
	his-58::ZC504.3 3'-utr, unc-119(+)] II;		
	zyg-1(it25) II; sas-6(luv1) IV		
IYR006	*bsSi15 [pKO109: spd-2p::spd-2::*	Crossed IYR002	This study
	mCherry::spd-2 3'-utr, unc-119(+)] I;	with OC779	
	bsSi30[pCW9: ZC504.3p::sfgfp::		
	his-58::ZC504.3 3'-utr, unc-119(+)] II;		
	zyg-1(it25) II; sas-6(luv2) IV		

STRAIN	GENOTYPE	METHOD	RESOURCE
IYR026	*oyIs59 Is[osm-6p::gfp] III; sas-6(luv1) IV*	Crossed IYR001 with PY6100	This study
IYR027	*oyIs59 Is[osm-6p::gfp] III; sas-6(luv2) IV*	Crossed IYR002 with PY6100	This study

Table S2

Gene	Type of oligo	Sequence (5' to 3')
dpy-10 (Paix *et al.* 2015)	co-CRISPR crRNA	GCUACCAUAGGCACCACGAG
sas-6 (Bergwell *et al.* 2019)	C-terminal HA-tag crRNA	AUUUUAUCGUUGAGCGGGUG
sas-6	L69T crRNA	UGAUUUUGAAUUUCUAUUCU
dpy-10 (Arribere *et al.* 2014)	*dyp-10(cn64)* repair oligo	CACTTGAACTTCAATACGGC AAGATGAGAATGACTGGAAA CCGTACCGCATGCGGTGCCTA TGGTAGCGGAGCTTCACATGG CTTCAGACCAACAGCCTAT
sas-6 (Bergwell *et al.* 2019)	HA repair oligo	TATTTTCAAGTAAAGGACAA GAAAAAATCAATAAAAAGA TTTTAAGCGTAATCTGGAACA TCATATGGGTAACGCTGTGCC GGCGGAGTGTTCGTCACACTT GAACCAGTAGTCTCGTCGGC GATTAGTTGA
sas-6	L69T repair oligo	AAAGGAGCTAAAATTCGAAAT CAGCCGCTCCGATGATTTTGA ATTCCTCTTTTCAGAGACAACG AACAACGAGAAATATCAGATTT TGGCTCGAGATCACGATTTAA CA

Table S3

Antibodies	Host	Manufacturer	Catalog#	Application	Dilution
Alpha tubulin	Mouse	Santa Cruz Biotechnology	sc-32293	IB	1:200
HA	Rabbit	Cell Signaling	3724S	IB	1:1000
Mouse IRDye 680 RD	Goat	LI-COR	926-68070	IB	1:14000
Rabbit IRDye 800CW	Donkey	LI-COR	926-32213	IB	1:14000