Centromere-associated retroelement evolution in *Drosophila melanogaster* reveals an

underlying conflict

Lucas W. Hemmer[1,2*], Sherif Negm[1,3], Xuewen Geng[1], Cécile Courret[1], Beatriz Navarro-

Domínguez[1,4], Iain Speece[1], Xiaolu Wei[1], Eddyson Altidor[1], James Chaffer[1], John S. Sproul[1,5],

Amanda M. Larracuente[1*]

Affiliations:

[1] Department of Biology, University of Rochester, Rochester, NY, USA

[2] Invaio Sciences, Cambridge, MA, USA

[3] Department of Human Genetics, University of Chicago, Chicago, IL, USA

[4] Departamento de Genética, Facultad de Ciencias, Universidad de Granada, Granada, 18071,

Spain

[5] Department of Biology, University of Nebraska Omaha, Omaha, NE, USA

[*]**Correspondence:** alarracu@bio.rochester.edu, lhemmer08@gmail.com

17   **ABSTRACT**

Centromeres are chromosomal regions essential for coordinating chromosome segregation during

cell division. While centromeres are defined by the presence of a centromere-specific histone H3

variant called CENP-A rather than a particular DNA sequence, they are typically embedded in

repeat-dense and heterochromatic chromosomal genome regions. In many species, centromeres

are associated with transposable elements but it is unclear if these elements are selfish and target

centromeres for insertion or if they play a role in centromere specification and function. Here we

use *Drosophila melanogaste*r as a model to understand the evolution of centromere-associated

transposable elements. *G2/Jockey-3* is a non-LTR retroelement in the *Jockey* clade and the only

sequence shared by all centromeres. We study the evolution of *G2/Jockey-3* using short and long

read population genomic data to infer insertion polymorphisms across the genome. We combine

estimates of the age, frequency, and location of insertions to infer the evolutionary processes

shaping *G2/Jockey-3* and its association with the centromeres. We find that *G2/Jockey-3* is an

active retroelement that is targeted by the piRNA pathway. Our results suggest that *G2-Jockey-3*

is highly enriched in centromeres at least in part due to an insertion bias. We do not detect any

signature of positive selection on any *G2/Jockey-3* insertions that would suggest than individual

insertions are favored by natural selection. Instead, we infer that most insertions are neutral or

weakly deleterious both inside and outside of the centromeres. Therefore, *G2/Jockey-3* evolution

is consistent with it being a selfish genetic element that targets centromeres. We suspect targeting

centromeres for insertion helps active retroelements like *G2/Jockey-3* escape host defenses, as the

unique centromeric chromatin may prevent targeting by the host silencing machinery. On the other

hand, centromeric TEs insertions may be tolerated or even beneficial if they also contribute to the

right transcriptional and chromatin environment. Thus, we suspect centromere-associated

retroelements like *G2/Jockey-3* reflect a balance between conflict and cooperation at the

centromeres.

INTRODUCTION

Centromeres are important chromosomal structures for kinetochore assembly and proper chromosome segregation during cell division. While centromeres are usually in highly repetitive regions, the role of DNA sequences in centromere function or establishment remains unclear (reviewed in [1]). Across many organisms, centromeres are associated with satellite DNAs (satDNAs), large blocks of tandemly repeated DNA sequences (reviewed in [2, 3]). Centromere-associated tandem repeats are capable of initiating centromere assembly in fission yeast [4] and humans [5]. However, centromeres are defined epigenetically by CENP-A, a specific variant of histone H3 [6-11] and the centromere domains consist of unique chromatin with H3 nucleosomes interspersed with CENP-A nucleosomes [12-15]. Repeat sequences are not always required for centromere formation, as neocentromeres can form outside of centromeric DNA in the absence of satDNA sequences [16-18] and some species have centromeres devoid of repeats (*e.g.* satellite-free centromeres in the Equus genus [19]). Neocentromeres have the potential to replace older centromeres with implications for karyotype evolution and speciation [20-22].

Centromeres and pericentromeric heterochromatin are also enriched in transposable elements (TEs)— selfish genetic elements that mobilize in genomes. Evidence for specific TE-centromere associations has emerged in the last two decades, including in grasses [23, 24], rice [25, 26], maize [27], marsupials [28, 29], bats [30], gibbons [31, 32], and humans [18, 33, 34]. However, it is unclear how, or if, TEs contribute to centromere biology for most species [35]. Centromeres may offer a 'safe haven' to TEs that jump there and for the genome [36, 37], as centromere insertions are unlikely to disrupt centromere function [35]. At the same time, centromeric TEs may contribute

to important aspects of centromere evolution and biology. Homology between some TEs and

centromeric tandem repeats [3, 38] suggests that TEs can contribute to centromere evolution by

giving rise to new centromeric satDNAs [39]. TEs may also play roles in centromere function.

These roles may be indirect, such as maintaining the constitutive pericentromeric heterochromatin,

which is important for centromere function [40-42]. However, more direct roles are also possible.

Some transcription of centromeric regions is necessary for recruitment of CENP-A and

establishing centromere location [43-47]. Centromeric transcripts may also facilitate recruitment

of kinetochores to centromeric DNA [48, 49]. Given that TEs contain sequences important for

promoting transcription [50], their presence in centromeres may contribute to centromere

specification and function [35]. In most cases it is unclear whether TEs are a part of the CENP-A

domain corresponding to functional centromeres or just associated with adjacent, repeat-rich

regions of chromosomes in which functional centromeres are often embedded. Advances in single

molecule long-read sequencing technology now allows for detailed study of centromere

organization at the sequence level to provide insights into centromere structure, composition, and

evolution [51-56].

Drosophila melanogaster is an excellent model organism to study centromere evolution with its

compact genome and expansive genetic toolkit [57]. A recent study of centromere organization in

D. melanogaster revealed that CENP-A is bound to TE-rich islands of complex DNA embedded

within large arrays of satellite repeats [53]. The composition of each centromere island is unique.

The only sequence element shared among all centromeres is a single non-LTR retroelement in the

Jockey superfamily, *G2/Jockey-3*. However, *G2/Jockey-3* is not exclusively located in

centromeres; about 40% of all insertions are found outside of the centromere islands [53].

Polymorphic *G2/Jockey-3* insertions [58] and transcripts from published RNAseq datasets [59-61]

suggest this element may still be active in *D. melanogaster* [53]. *G2/Jockey-3* is also found at the

centromeres in *D. simulans*, a close relative of *D. melanogaster,* suggesting that its centromere

association may be broadly conserved [53].

It is unclear whether *G2/Jockey-3* is an important centromere component in *D. melanogaster* or if

they are instead purely selfish TEs. If *G2/Jockey-3* are important for centromere specification or

function, we may detect evidence for positive natural selection on centromeric copies.

Alternatively, *G2/Jockey-3* may be a selfish element that targets insertions in centromeres but are

otherwise selectively neutral or weakly deleterious. These hypotheses are not necessarily mutually

exclusive. Here we investigated the age and frequency of *G2/Jockey-3* insertions across *D.*

melanogaster genomes to determine if its evolution is consistent with a role in centromere function

or underlying selfish behavior. Our analyses suggest that *G2/Jockey-3* insertions are largely neutral

or weakly deleterious both inside and outside centromeres, suggesting that these are selfish genetic

elements. Our results also suggest that *G2/Jockey-3* has an insertion bias toward centromeres.

Although there is no evidence for positive selection on individual *G2/Jockey-3* centromeric

insertions, insertion of active TEs like *G2/Jockey-3* may still contribute to transcription and the

unique chromatin environment at centromeres. We propose that there may be a balance between

conflict and cooperation between active TEs like *G2/Jockey-3* and centromeres.

RESULTS

1. A new consensus sequence for G2/Jockey-3

Prior to analyzing the distribution of *G2/Jockey-3* in populations, we first confirmed the consensus

annotation. We analyzed all *G2/Jockey-3* insertions in our *D. melanogaster* genome assembly [53]

and found two lines of evidence that the annotation needed updating: 1) 25 of the 167 *G2/Jockey-*

3 sequences in the genome shared up to 92 bp of upstream sequence at the 5'-end of the annotation;

and 2) our reanalysis of PROseq data [62] suggests that *G2/Jockey-3* transcription starts 92 bp

upstream of the published start site (Figure S1A). This upstream region is also enriched for GO

terms for RNA polymerase II transcription factors and other transcription factor activity (Figure

S1B). We therefore extended the original *G2/Jockey-3* consensus sequence by 92 bases (File S1).

G2/Jockey-3 is a part of the *Jockey* superfamily of non-LTR retroelements, which typically contain

two open reading frames (ORFs) (Figure S1A). While ORF1 is generally less conserved [63],

ORF2 is well-conserved across *Jockey* elements and encodes for a reverse transcriptase (RT) and

an endonuclease. NCBI ORF finder detected both ORFs within *G2/Jockey-3*. ORF1 is predicted

to produce a 427 amino-acid protein approximately 186 bp downstream of the 5' end. The NCBI

Conserved Domains identified a PRE_C2GC super family (cl11655, E-value 9.22E-18) within

ORF1 and while the function is unknown, it is associated with Cys2His2 type zinc fingers found

in putative GAG proteins in other arthropod species. The predicted ORF2 overlaps with the last

four bp of ORF1 and contains a domain corresponding to endonuclease (EEP Superfamily

Accession cl00490, E-value 7.21E-18) and RT (RT_like Superfamily Accession cd01650, E-value

4.62E-56). *G2/Jockey-3* ends in a poly(A) signal.

2. Overview of data sets analyzed

We studied the evolutionary dynamics of *G2/Jockey-3* and other TEs by investigating the patterns

of insertion polymorphisms both in a single panmictic population, and in individuals from a wide

geographic range (Figure 1, Table S1). We conducted a survey of TE dynamics and genetic

variation in 135 samples from the *Drosophila melanogaster* Genetic Reference Panel (DGRP)

collected from a single population in North Carolina with whole genome Illumina sequence data

[64, 65] and a subset of long read data from the same population for validation [66] (see

Supplemental Text). For the samples from a wide geographic range, we used both long- and short-

read data from 13 founder lines of the *Drosophila* Synthetic Population Resource (DSPR) and

Oregon-R [67]. We also analyzed long-read PacBio data from five Global Diversity Lines (GDL)

samples from different continents [68]. We generated *de novo* genome assemblies of each line to

determine the organization of the centromere islands originally described in the reference genome

for comparison (Table S2) and predicted insertions in short read data with the McClintock pipeline

[69]. We assessed the reliability of our TE insertion predictions by looking for concordance

between short and long read predictions (see Supplemental Text) and through PCR validation of a

subset of insertions. We confirmed 64% (67 / 105) of the *G2/Jockey-3* insertions detected by

McClintock are found in the long-read samples we examined (File S2). PCR confirmed 80% (28 /

35) of *G2/Jockey-3* insertions predicted in the GDL lines, and ~67% (12 / 18, File S3) of the

centromere insertions. Based on our assessment, we limit our analysis to short read datasets with

>20X centromere coverage (Table S3) and TEs predicted by more than one TE caller implemented

in McClintock (Supplemental Text).

Figure 1: The summary of data and analyses using one source or a combination of sources

in this paper. A heterochromatin-enriched assembly was originally used to discover centromere

islands in Chang et al. [53] and identify reference TE insertions. We used short-reads from the

DGRP to identify reference and *de novo* insertions to study TE dynamics within a single population

from North Carolina (map source: R package usmaps https://cran.r-

project.org/web/packages/usmap/). We generated genome assemblies for 18 samples from global

populations to uncover the centromere island sequences and *de novo* elements (map source:

www.outline-world-map.com). We combined the reference sequences with population frequencies

to infer selection with an age-of-allele test [70]. We also combined reference TE sequences with

de novo elements uncovered in our assemblies for phylogenetic analysis. The fly illustrations are

modified images from Wikipedia Commons

(https://commons.wikimedia.org/wiki/File:Drosophila-drawing.svg).

3. G2/Jockey-3 is active in Drosophila melanogaster populations

Previous studies of TE polymorphism in *D. melanogaster* are consistent with *G2/Jockey-3*

(previously annotated as *G2*) being recently active [58, 71, 72]. To determine if *G2/Jockey-3* is

active in the DGRP, we detected TE insertion polymorphisms with the McClintock pipeline [69]

and classified each insertion based on their presence (a "reference" insertion) or absence (a non-

reference or "*de novo*" insertion, Figure 2) in our reference genome [53, 73]. *G2/Jockey-3* showed

approximately five *de novo* insertions per line (File S4-S5), which is considerably fewer than the

most highly active *Drosophila* TEs. For example, we estimate *Hobo* and *Jockey-1* have

approximately 35 *de novo* insertions per line (consistent with [58]). However, the fraction of *de*

novo insertions as percentage of all insertions (39% or 733 / 1124 for *G2/Jockey3*) is comparable

to other highly active elements [72] such as *Doc* (25%; 2944 / 3826), *jockey* (24% ;1308 / 5448),

I element (27%; 658 / 2364), and *Copia* (88% ;1083 / 1226) (File S4).

We detected insertions in the centromeres, here defined as the centromere islands identified in

Chang et al. [53]. Of the 18 different TE families detected within centromeres, *G2/Jockey-3* is the

most common. Within centromeres, *G2/Jockey-3* has the highest percentage of *de novo* (32%;

15/47) and reference (72%; 28 / 39, Figure 2) TE insertions. Only *G2/Jockey-3* and *Doc* have more

than 10 unique insertions (File S4). Other *de novo* centromere insertions include active families

such as the *F* element, *R1* and *Roo*.

Many active TEs are targeted by small, 24-30-nt long Piwi-interacting RNAs (piRNA) in the

germline [74, 75]. These piRNAs target TEs for both transcriptional and post-transcriptional

silencing. In the *D. melanogaster* germline, piRNAs are produced from precursor transcripts that

originate from discrete loci in the genome that contain clusters of nested TEs [76] called piRNA

clusters [77]. Subsequent amplification of the piRNA signal comes from the piRNA-guided

cleavage of active TEs through the 'ping-pong' cycle [77, 78] generating a signature of 10-nt

overlap between sense and antisense piRNAs. Consistent with *G2/Jockey-3* being a recently active

element, we detect insertions of *G2/Jockey-3* in the *38C1* (2 copies) and *38C2* (9 copies) piRNA

clusters, and *G2/Jockey-3*-derived piRNAs in the germline. *G2/Jockey-3* is ranked in the top 4%

and top 18% of TE-derived piRNAs in testes and ovaries, respectively (Table S4). The resulting

G2/Jockey-3 piRNAs show signatures of ping-pong in ovaries (Z-score = 7.11; data from [60, 79,

80]) but not in testes (Z-score = 0.15; data from [81-83]), consistent with other testis piRNAs.

Taken together, these results suggest that *G2/Jockey-3* is an active TE enriched in centromere

islands.

Figure 2: The distribution and frequency of reference and _de novo G2/Jockey-3_ insertions along the chromosomes and within the centromeres in _D. melanogaster_. The height of the bars corresponds to frequency of an insertion detected in the DGRP samples and the color denotes whether an insertion is found in the reference genome or _de novo_. Each A-D subpanels depict a chromosome (2,3,4,X) from the heterochromatin-enriched assembly of _D. melanogaster_ genome [53, 73]. Corresponding centromere sequences are zoomed in as highlighted (X chromosome centromere = CenX, _etc_).

4. Young G2/Jockey3 insertions are enriched in centromeres

To understand if the observed enrichment of *G2/Jockey3* at centromeres is due to insertion bias,

we studied insertion frequencies across the genome. We reasoned that an enrichment of putatively

young *G2/Jockey-3* at centromeres may reflect an insertion preference. Young recently inserted

TEs are expected to occur at lower frequencies than older TEs. We thus examined the distribution

of rare insertions (<10% population frequency) of *G2/Jockey-3* and other TE families in the DGRP.

We found that rare *G2/Jockey-3* insertions are significantly enriched in centromeres compared to

other regions relative to their size (12 insertions; FET, *P* < 1E-3, Figure 3). *R1* elements are the

only other TE enriched in centromeres, but only relative to euchromatin (FET, *P* = 0.014, Figure

3). *G2/Jockey-3* is therefore unique in being the only TE that is significantly enriched in

centromeres over the heterochromatin and euchromatin (Figure 3). Among rare insertions,

G2/Jockey-3 is by far the most enriched TE in centromeres (39% of all rare insertions).

Figure 3: Young *G2/Jockey-3* insertions are highly enriched in centromeres relative to

heterochromatin and euchromatin. The distribution of rare insertions (< 10% of the population)

per Mb between the centromere islands, heterochromatin, and euchromatin regions for several TE

families in DGRP lines. Significant differences are noted as: * *P* < 0.05, *** *P* < 0.001.

We confirmed these patterns using an orthogonal approach with long-read genome assemblies of

flies from a wide geographic distribution (*i.e.,* founder DSPR lines, GDL lines, and our reference

genome; Table S1). Here we identified putatively young *G2/Jockey-3* insertions based on sequence

divergence instead of frequency. We classified young insertions as those with < 1% divergence

from the consensus sequence, or with < 2% divergence from the consensus and found within a

single genome assembly. Insertions classified as young were found in less than 50% of the

assemblies, had fewer unique substitutions, and were more likely to have a complete ORF than

older fragments (Figure 4, FET *P* = 3.93E-12, File S6, File S7). Consistent with our short-read

DGRP findings (Figure 3), we found significant relative enrichment in centromeres (Figure 5, FET

P = 8.28E-5; Y chromosome excluded *P* = 0.0142), even with some of the centromere island

sequences missing in the long-read assemblies (Table S2).

These results suggest that *G2/Jockey-3* shows at least some insertion preference for centromeres.

As insertion bias can be associated with a target site sequence, we searched for motif enrichment

at insertions sites. Although we found an enrichment of AT-rich simple motifs like "TAGTTTT"

and rDNA-associated sequences (*e.g.* 18S and non-transcribed spacers; Figure S2) within 100

bases of predicted insertion sites, these sequences are generally abundant in repeat-dense regions

of the genome and not specifically enriched at *G2/Jockey-3* insertions (File S6, File S8; MEME

E-value = 3.8E-16).

Figure 4: Young *G2/Jockey-3* insertions share a common origin and are enriched in centromeres. A maximum likelihood phylogenetic tree with all non-redundant *G2/Jockey-3* sequences larger than 10% of the full-length (400 bp) in the reference genome [53] and long-read assemblies of founder DSPR and GDL lines, with the *D. simulans* consensus as the outgroup. Black circles denote "complete" *G2/Jockey-3* elements containing a full ORF2; gray circles are incomplete elements. The columns and colors correspond to characteristics of each insertion including chromatin region (centromere or not), relative age, and presence on Y chromosome. Young insertions, determined using an orthogonal approach to the Figure 3 analysis, are those with < 1% divergence from the consensus sequence or < 2% and found within a single genome assembly.

5. Skew toward high frequency variants of G2/Jockey-3 in the centromere

To test if *G2/Jockey3* insertions are retained at centromeres by natural selection we studied their

insertion frequencies. Because most TE insertions are deleterious in the euchromatin, they are

found at low population frequencies [84]. In contrast, TEs in heterochromatin are allowed to reach

higher population frequencies because recombination is infrequent, and natural selection is less

efficacious (reviewed in [85-87]. To infer differences in the strength and direction of natural

selection in different genomic regions, we examined the frequency spectra of a subset of TE

families in centromeres, heterochromatin, and euchromatin (Figure 5, Figure S3). If natural

selection favors centromeric insertions, we should observe a skew toward high frequency variants

in centromeres compared to heterochromatin. We compared the frequency spectra of two TE

families which had five or more insertions in the centromeres: *G2/Jockey-3* and *Doc*. As expected,

both elements showed an enrichment for rare insertions in the euchromatin, consistent with TEs

being generally deleterious in gene-rich regions (Figure 5; FET *P* < 0.001). Similarly, we observed

a   slight   skew   toward   higher   frequency   *G2/Jockey-3*   and   *Doc*   insertions   (*e.g.*

common/intermediate) in heterochromatin and centromeres compared to euchromatin (Figure 5,

FET *P* < 0.001), suggesting relaxed constraints or a reduced efficacy of natural selection outside

of euchromatic regions. Other TE families found in centromeres are also enriched for low

frequency insertions in the euchromatin (Figure S3). However, we did not observe differences

between the frequency spectra of the euchromatin and centromeres, suggesting most TEs may be

deleterious in these regions (Figure S3).

The frequency spectra of *G2/Jockey3* does not appear significantly different between centromeres

and heterochromatin. However, *G2/Jockey-3* has more "common" insertions in centromeres

relative to other TEs, including *Doc* (Figure 5; Figure S3). This could be due to relaxed constraints

allowing *G2/Jockey-3* insertions to rise to higher frequencies over time or positive selection for

some *G2/Jockey-3* insertions in centromeres.

Figure 5: *G2/Jockey-3* has higher frequency variants in centromeres. TE frequencies were

binned into rare (less than 10% of all samples), intermediate (10-50% of all samples), and common

(> 50% of all samples). The frequency distributions of the four TE families with > 10 unique

insertions in the centromeres were compared among the centromeres, heterochromatin, and

euchromatin. Fisher's Exact Tests between regions, Bonferroni adjusted two-tailed P-value * $P <$

0.05; *** $P < 0.001$; ns is not significant.

6. G2/Jockey-3 ages suggest a historical burst of activity

While young *G2/Jockey-3* insertions appear to be enriched at centromeres (Figure 3), centromeric copies are generally older—within a range of 2-7% divergence from the consensus—than those in heterochromatin (MWU, $P < 0.001$, Figure 6A), but not significantly different than euchromatic copies (MWU, $P > 0.5$; Figure 6A). This pattern suggests that most centromeric copies of *G2/Jockey-3* resulted from a historical burst of activity. This burst of activity is consistent with general patterns observed at other non-LTR retroelements [72, 88]. Other TE families show a similar age distribution between heterochromatin and the centromeres (Figure S4). These copies are also older and highly fragmented. In general, both the lack of recombination and lack of strong purifying selection against TE insertions in centromeres and heterochromatin may allow these insertions to persist over longer evolutionary time periods [89].

We can combine information about the TE's age and frequency to infer how, and if, its history is consistent with natural selection. In the absence of selection, the frequency of a TE insertion should correlate with its age: *e.g.,* older TEs have more time to drift to higher population frequencies than young TEs and thus accumulate more unique mutations. Natural selection can cause deviations from expected frequencies based on the age. While negative selection might cause TEs to have lower population frequency than expected given their age, positive selection might increase their frequencies. We therefore used an age-of-allele test [70] to detect whether *G2/Jockey-3* frequencies deviate from our expectations. We estimated age as the number of unique substitutions for each TE copy based on the reference genome and we estimated frequency from the 28 high-coverage DGRP samples, controlling for demographic history [70]. For this test, we need to

compare insertions spanning a range of ages to identify individual insertion frequencies that

deviate from expectations [70]. We thus examine *G2/Jockey-3* and other TE families that have

centromere insertions. We find that *G2/Jockey-3* (Figure 6B, File S9)*, like the majority of TEs,

segregates either at the expected frequencies or lower than expected, given their age (Figure S6,

File S9). This suggests that many *G2/Jockey-3* copies, both within and outside the centromere, are

either neutral or weakly deleterious. Therefore *G2/Jockey-3* shares similar selective pressures as

other presumably weakly deleterious TE families in the genome.

Interestingly, we observed an enrichment of copies with zero unique substitutions for the

heterochromatin and CenY insertions (Figure S5). However, we suspect this pattern arises from

duplications rather than independent *G2/Jockey-3* insertion events (see below). Genomic

duplication events are common in heterochromatin in the Y chromosome [73, 76, 90].

In sum, while we do not see evidence for strong recent positive selection on any individual

G2/Jockey-3 insertions in the centromeres, there may be selection on centromere haplotypes that

differ in their composition or structure.

Figure 6: *G2/Jockey-3* copies are either as frequent or less frequent than expected given their age. A) The number of *G2/Jockey-3* insertions grouped by divergence from the consensus sequence split between the euchromatin, heterochromatin, and centromeres. *** $P < 0.001$, ns is not significant. B) The difference between observed and expected frequency of *G2/Jockey-3* copies in the DGRP. The *P*-values are calculated as the sum of the probabilities that an insertion is less frequent than expected (if observed - expected < 0) or more frequent than expected (observed - expected > 0). * $P < 0.1$, a stringent threshold given the reduced power to detect deviations from neutrality as stated in Blumenstiel et al. [70].

7. Centromere island haplotype structure not likely driven by G2/Jockey-3 insertions

To detect signatures of natural selection on centromere haplotypes and any associations with

G2/Jockey-3 copies, we examined patterns of nucleotide variation. Centromeres and

pericentromeric regions in *D. melanogaster* do not recombine [91, 92] therefore the effects of

linked selection in non-recombining regions should lead to low levels of nucleotide diversity. We

thus expect to see reduced genetic variation in the centromeres and high levels of linkage

disequilibrium. We quantified genetic variation in the centromere islands for DGRP samples with

high centromere coverage ($> 20X$, see Methods for more details). As expected, nucleotide diversity

is greatly reduced in every centromere compared to the major chromosome arms (MWU $P < 2.0E$-

07, Figure S7, Table S5) except for chromosome 4, which does not recombine, and its centromere

and chromosome arms are similar (MWU $p = 0.56$, Figure S8, Table S5). We find that Tajima's

D, a summary of the site frequency spectrum, is also typically lower in centromere islands (Figure

S8), indicating a skew toward rare alleles in centromeres. The centromere island haplotype

structures (Figures S9-S11) indicate that the majority of centromeric SNPs are randomly

distributed with respect to the *G2/Jockey-3* sequences (FET $P > 0.46$). Linked selection anywhere

in or near centromeres can reduce genetic diversity, skew frequency spectra, and create haplotype

structure. While centromere island haplotype structure is consistent with a history of linked

selection, we do not see evidence that *G2/Jockey-3* insertions themselves are necessarily the targets

of selection.

8. G2/Jockey-3 copy number amplification on the Y chromosome is driven by duplication

The Y chromosome centromere is unique in that it has the largest centromere island containing

nearly half of all *G2/Jockey-3* fragments, but no identifiable satellite sequences [53]. We cannot

use the genomic re-sequencing data to study polymorphic insertions on the Y chromosome, as the

data are either female-only or mixed sex and thus lower coverage. However, our high-quality

reference assembly provides an opportunity to infer the organization and evolutionary history of

Y-linked *G2/Jockey-3*. We inferred a phylogenetic tree of all Y-linked copies of *G2/Jockey-3*.

Based on this phylogeny, we classified centromeric Y-linked *G2/Jockey-3* copies into four broad

classes (Figure S12). Class 1 *G2/Jockey-3* correspond to young and mostly long insertions, class

2 to short (< 160 bp) 3' end fragments, class 3 to old internal fragments missing the 5' and 3'

terminal ends, and class 4 to generally long and old 3' end fragments divided into two subclasses

(4 and 4A) based on position (Figure 9, Figure S12). We mapped these classifications onto our Y

chromosome assembly and estimated pairwise divergence between all *G2/Jockey-3* copies in the

Y centromere (Figure 7) and elsewhere on the Y chromosome (Figure S13). Each class seems to

undergo independent duplication events, however at a larger scale we do not identify higher order

repeat domains (Figure 7). This structure differs from other known centromeres (*e.g.* primates),

which are organized into higher order repeats, the youngest repeats are found in the center and the

older, more divergent repeats, are at the outer edges of the array [51, 52]. By contrast, in the *D.*

melanogaster Y centromere we observe that regions enriched with young *G2/Jockey-3* copies

punctuate regions rich with older copies (Figure 9), suggesting that recent, local rearrangements

affect centromere organization.

Figure 7: *G2/Jockey-3* evolution in the Y chromosome centromere is shaped by expansion via

duplication, gene conversion and rearrangements. The colored pairwise divergences between

individual *G2/Jockey-3* copies on the Y centromere are shown on top with their relative positions

on the line graph below. White spaces represent *G2/Jockey-3* copies which do not overlap, so

pairwise divergence cannot be calculated. Below the graph is a visual representation of the broad

categories of repeats on the Y chromosome centromere, labeled by color with length and direction

illustrated with arrowed lines. The repeats appear arranged into roughly four regions with

overlapping borders based on the types of repeats and pairwise divergence. The classes of

G2/Jockey-3 are as follows: Class 1 younger insertions, Class 2 short (< 160 bp) 3' end fragments,

Class 3 older internal fragments missing the 5' and 3' terminal ends, Class 4 generally longer and

older 3' end fragments. Class 4 contains a subclade (4A) with highly similar sequences almost

exclusively from the single region of CenY.

We estimated gene conversion rates of *G2/Jockey-3* on the Y centromere with respect to their

classes (Supplemental Text). Our estimates for gene conversion are on the order of 1E-6 per site

per generation (Table S6), ~10E3 times higher than the other chromosomes in *Drosophila* [91-93]

and an order of magnitude less than previously estimated gene conversion rates between duplicated

gene families on the Y chromosome [73]. Moreover, we do not see complete sequence

homogenization within classes. We observe similar patterns of duplication and expansion outside

the centromere on the Y chromosome (Figure S13), as 32 of 47 non-centromere Y-linked copies

appear to be duplicated; they are of similar fragment size, orientation, and have few no zero unique

substitutions (File S6). Therefore, we conclude that *G2/Jockey-3* evolution on the Y chromosome

is driven by duplication of historical insertions with some gene conversion among copies.

Population genomic data from the Y chromosome will help distinguish between types of

recombination affecting Y centromere structure.

Outside of the Y chromosome, putative duplications involving *G2/Jockey-3* are less common

(Figure S13). Many copies of *G2/Jockey-3* within the centromeres are fragmented and therefore

share no homologous sequences for pairwise calculations (Figure S13). Pairwise divergence

between centromeric *G2/Jockey-3* fragments is elevated in comparison to fragments in both the

euchromatin and non-centromeric heterochromatin (Figure S14, MWU, *P* < 0.01), and lowest in

non-centromeric heterochromatin, especially on the Y chromosome. Therefore, the majority of

G2/Jockey-3 within the non-Y centromeres appears to be the result of recurrent past insertions.

426   CONCLUSIONS

G2/Jockey-3 is a unique TE in that it is the only sequence shared among centromeres of *D.*

melanogaster [53]. Here we show that *G2/Jockey-3* is a recently active non-LTR retroelement that

may have an insertion preference for centromeres (Figures 3-4). While the centromere enrichment

and skew toward high frequency centromere copies are expected for insertions favored by natural

selection, we do not find evidence for positive selection on individual *G2/Jockey-3* insertions.

Instead, most *G2/Jockey-3* insertions both inside and outside of the centromeres are at the expected

frequency or significantly rarer than expected given their age, suggesting they may be neutral or

slightly deleterious and accumulating due to relaxed constraints against centromeric TE insertions

(Figure 6B). The majority of *G2/Jockey-3* copies within the centromeres appear to be the result of

recurrent past insertions, except for on the Y chromosome. Some Y-linked copies are likely a result

of duplications rather than *de novo* insertions, similar to patterns reported in other species (*e.g.*

[28]).

We propose that most *G2/Jockey-3* insertions in the centromeres resulted from a burst of activity,

which is consistent with the evolutionary history of other non-LTR retroelements [88]. When a TE

first invades, it can proliferate rapidly until it comes under the surveillance of germline defense

mechanisms like the piRNA pathway for example, by inserting into a piRNA cluster [94-96].

These dynamics are consistent with the invasion and control [94, 97] of the *P* element in *D.*

melanogaster [98] and its close relative, *D. simulans* [99]. Once targeted by the piRNA pathway

for transcriptional and posttranscriptional silencing, the TE may eventually become dormant or go

extinct. Here we show that *G2/Jockey-3* had a burst of activity and is now inside piRNA clusters

and actively targeted by the piRNA pathway (Table S4).

Therefore, the *G2/Jockey-3* insertion polymorphism patterns and piRNA observations are most

consistent with some underlying selfish behavior typical among TE families, except that this

element can target centromeres. Although we are unsure of the precise targeting mechanism, we

suspect the target is the centromere chromatin rather than any particular DNA sequence. Such

centromere-specific targeting was reported for LTR retroelements in other species such as maize

[100, 101] and *Arabidopsis* [102, 103]*.* For example, when the *Arabidopsis lyrata Tal-1 Copia-*

type retroelement [103] is introduced into *A. thaliana,* the element targets the centromeres despite

divergence in centromeric DNA sequences between these species [102].

TEs may accumulate in heterochromatic regions where they experience relaxed selection

compared to euchromatic insertions, which have potential to disrupt functional elements, risk

ectopic exchange and resulting genomic rearrangements [85-87], and spread silent chromatin into

nearby functional elements [85-87, 104, 105]. However, centromeres are not typical

heterochromatin [106]. Here we propose that centromeres offer a unique environment for TEs

(Figure 8A). First, insertions are unlikely to be strongly disfavored, as the function of centromeres

is unlikely to be disrupted by a single insertion event. Second, centromeric insertions may escape

host silencing mechanisms. Small RNAs target TEs to limit potentially harmful TE activity by

inducing H3K9 methylation and thus transcriptional silencing (reviewed in [107-109]). However,

CENP-A structure and posttranslational modifications differ from H3 [106], and the host silencing

machinery is not known to target CENP-A. Therefore, centromeres may offer a 'safe haven' for

some TEs not just because they can avoid removal by purifying selection, but also because TEs

may temporarily escape targeting by small RNAs (*e.g.* piRNAs, siRNAs) and alternative host

defenses (*e.g.* DNA methylation in pericentromeres [110] hypomethylated DNA in humans [52,

111]).

While most TEs are considered to be parasitic elements that are neutral or mildly deleterious [112,

113], some TE insertions, including those in non-recombining and/or gene poor heterochromatic

environments, might contribute to host function [114]. For example, TEs can adopt commensal

and/or mutualistic strategies to propagate without direct confrontation by cooperating with their

host (reviewed in [115]). One extreme outcome is TE domestication, where the elements no longer

pose a threat to the host and instead now perform some important genome function. One often

cited example is at *Drosophila* telomeres, where three non-LTR retroelements in the *Jockey* clade

(*HeT-A*, *TAHRE*, and *TART* collectively referred to as HTT) have specialized roles [116, 117]. The

transposition activity of HTT is limited to telomeres, where they directly contribute to telomere

length and integrity [116-119]. However, detailed study of the evolutionary genetic history of the

retroelements across species suggests instead that there is still an underlying conflict between the

HTT and the genome, as there is rapid turnover of HTT between species [120]. Alternatively,

given the specialization of HTT, the dynamic and unstable telomeric environment may necessitate

rapid HTT evolution [121].

While centromeric TEs do not currently show evidence for domestication in *D. melanogaster*, they

may be tolerated, or even beneficial for three reasons. First, TEs may contribute to centromere

transcription (reviewed in [35]), as some transcription at centromeres is important for CENP-A

assembly and function [14, 46, 122-126] and centromere function [14, 45, 127]. Second, TEs might

contribute to the unique chromatin at centromeres. H3 nucleosomes are typically interspersed with

CENP-A within centromere domains [12, 14] [15], including a limited amount of H3K9me2/3

[13]. We propose that insertion of active TEs like *G2/Jockey-3* may contribute to this unique

CENP-A/H3 chromatin at the centromeres (Figure 8A). While landing in CENP-A nucleosomes

may allow the TE to temporarily escape silencing, if an active TE lands in an H3 nucleosome

region of the centromere, it may be targeted for silencing through H3K9 methylation (*e.g.* piRNA-

or siRNA-mediated silencing) and thus contribute to H3K9me2/3 domains within the centromere.

Third, TEs may help mitigate other conflicts that happen at centromeres. Some centromeric

sequences have the capacity to be selfish and bias their transmission through female meiosis in a

process known as meiotic drive [128]. Driving centromeres generally have expansions of satellite

DNA sequences that can recruit more kinetochore proteins and cause biased segregation toward

the egg pole [129, 130] (Figure 8B). Meiotic drive at the centromeres can lead to rapid evolution

of both centromere DNA and proteins that work to restore fair segregation [131, 132]. There are

parallel pathways to suppress meiotic drive involving rebalancing kinetochore proteins and

spreading heterochromatin to limit the CENP-A domain [133]. While centromere drive and

suppression models predict that the flanking heterochromatin domains are the main participants in

the conflict [133], the H3K9me2/3 domains within the centromeres might also contribute. Active

centromeric TEs that are targeted by small RNA pathways, like *G2/Jockey-3,* could lead to

spreading of H3K9me2/3 heterochromatin from within the centromere (Figure 8C). In this way,

the centromeric TEs and 'internal' centromeric H3K9 domains may also be a substrate for

suppressors of centromere drive to stop the unchecked spread of CENP-A caused by meiotic

drivers.

Even if TEs cooperate with centromeres, this may be an uneasy alliance as there could still be an

underlying conflict. For example, the invasion of H3K9me2/3 heterochromatin in centromeres is

potentially dangerous, as it may evict CENP-A and silence a centromere [5]. Therefore, the

centromeric chromatin, and the relative abundance of CENP-A and H3 nucleosomes, may mediate

the balance between conflict and cooperation between active TEs and centromeres.

In many species across the tree of life, centromeres are associated with TEs [18, 23-34, 53]. The

connection between TEs and centromeres is supported not just by TE sequences embedded in and

around present-day centromeres, but also in the origins of some proteins that function in

centromere biology (*e.g.* similarity between centromere protein CENP-B and Pogo family

transposons [134]). This relationship may represent a delicate balance between conflict and

cooperation between the centromeres and the repetitive selfish genetic elements that occupy these

genome regions. We expect this balance to vary depending on genomic context, level of TE

activity, and presence and frequency of centromere meiotic drive. If TEs contribute positively to

centromere function, many types of elements may be sufficient to provide the right 'environment'.

In this case, we would not necessarily expect to see a signature of selection on any one insertion

or type of insertion. While *G2/Jockey-3* is currently the principal element occupying the

centromere niche in *D. melanogaster*, it may be replaced over time by another type of repeat. In

our model, any repeat that can be transcribed and can recruit H3K9me2/3 (*e.g.* is targeted by small

RNAs) can contribute to centromere function. Detecting the mode, tempo, and target of selection

in non-recombining and dynamic environments like the centromeres is difficult. Exploring the

evolution of centromere composition in closely related species will provide additional insights

necessary to test these hypotheses.

Figure 8. Model for the accumulation of centromere-associated retroelements. Centromeres

have unique chromatin with interspersed CENP-A and H3 domains, including some H3K9me2/3

[12-15]. The diagrams represent nucleosomes, either CENP-A (blue) or H3 (green), with

H3K9me2/3 modifications indicated with a black dot. A. Active TEs like *G2/Jockey-3* jump in

centromeric chromatin (magenta triangles and lines in nucleosomes) and could contribute to

transcription (black wavy line) or be silenced by host machinery (orange hexagon). Centromeres

may offer a safe haven for TE insertion because insertions there are unlikely to be strongly

disfavored and because insertions in CENP-A nucleosomes may escape host silencing machinery

(*e.g.* piRNA pathway). Centromeric TE insertions might be tolerated or even favored by the host

if they contribute to centromere transcription or a low level of H3K9me2/3 inside centromeres. B.

Selfish centromeric DNA sequences can expand their copy number and bias their transmission to

the next generation through female meiosis [128]. C. One way to suppress cheating centromeres

is by spreading heterochromatin to limit the CENP-A domain [133]. TE insertions in H3

nucleosomes within the centromeres may be targeted for silencing by host silencing machinery

(green circle with magenta lines and black circle). These H3K9me2/3 nucleosomes within

centromeres may help create a more efficient response to driving centromeres by merging

spreading pericentric heterochromatin with internal H3K9me2/3.

559   **MATERIALS AND METHODS**

G2/Jockey-3 re-annotation and expression analysis

Chang et. al. [53] generated the *G2/Jockey-3* consensus sequence after a phylogenetic analysis

confirmed previously annotated *G2* and *Jockey-3* were the same element [53]. We annotated a

heterochromatin-enriched long-read genome assembly of *D. melanogaster* [73] with

Repeatmasker v4.1.0 using the previous *G2/Jockey-3* consensus sequence along with a *Drosophila*

repeat library [135]. We extracted *G2/Jockey-3* and the surrounding sequences using BEDTools

v2.29.2 [136], aligned the sequences to the previous consensus sequence, manually inspected the

alignment, and generated the new consensus sequence with Geneious v8.1.6 [137] (File S1). We

identified Open Reading Frames (ORFs) with NCBI ORFfinder

(https://www.ncbi.nlm.nih.gov/orffinder/) and screened the ORF candidates by identifying

conserved domains common in non-LTR retrotransposons with the NCBI Conserved Domains

search tool (https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi). We defined the 5' end by

identifying transcription along the length of *G2/Jockey-3* by mapping publicly-available single-

end Precision nuclear Run-On Sequencing (PROseq) downloaded from NCBI SRA under

BioProject PRJNA451204 and accessions SRR7050623 - SRR7050630 [62]. We removed

adapters and reads < 16bp using cutadapt v2.1. Reads were reverse complemented using the fastX

toolkit v0.0.13 (fastx_reverse_complement). We mapped reads to the heterochromatin-enriched

genome assembly of *D. melanogaster* using bowtie2 v2.3.5.1 (--no-discordant --dovetail --very-

sensitive options; [138]). We extracted reads > 16 bp in length mapping to *G2/Jockey-3*

annotations in the genome with samtools v1.7 [139] and remapped the reads to the *G2/Jockey-3*

consensus with bowtie2 v2.3.5.1 [138]. We separated reads based on their strand using samtools

(v1.7). We calculated read depth at each position along the *G2/Jockey-3* consensus and normalized

coverage by the total number of genome mapped reads (RPM) using bedtools genomecov [136].

We used alignment depth of all *G2/Jockey-3* genomic fragments to identify where transcription

occurs along the length of *G2/Jockey-3*. We submitted the sequence in 5' end motifs under

transcription peaks for a GoMo analysis [140] to identify motifs as possible promoters.

To detect piRNAs corresponding to *G2/Jockey-3,* we trimmed the small RNA-seq reads for

adaptors and excluded the reads that align to rRNA or tRNA. We aligned the remaining reads to

the heterochromatin-enriched genome assembly [53, 73] using Bowtie [141], and then used a

customized python script as described in Wei et al. [83] to count reads that mapped to each TE

feature (htseq_bam_count_proportional.py;

https://github.com/LarracuenteLab/Dmelanogaster_satDNA_regulation; last accessed 11/2022,

https://doi.org/10.5061/dryad.hdr7sqvj3). We reported the RPM (Reads Per Million) values in a

summary table. We calculated the 10nt overlap Z-score of piRNAs using piPipes [142].

Short-read analysis

We detected TE insertions from publicly available FASTQ files from 135 Drosophila

melanogaster Genetic Reference Panel (DGRP) lines from BioProject PRJNA36679 [64, 65]

downloaded from the European Nucleotide Archive last accessed September 20, 2019, 13 of the

founder lines of the Drosophila Synthetic Population Resource (DSPR) BioProject

PRJNA156883[143], and 85 samples from the Global Diversity Lines BioProject PRJNA268111

(Table S3)[144]. We detected TE insertions in each sample with the McClintock pipeline [69] (last

downloaded and updated, August 20, 2020) combining several split-read and discordant read

mapping programs including the following: ngs_te_mapper, popoolationTE, popoolationTE2,

Retroseq, TE-locate, and TEMP (scripts and files for running McClintock at

https://github.com/LarracuenteLab/Dmel_Jockey-3_Evolution). We concatenated the outputs to

calculate the number of TEs detected by each program and sample and analyzed the outputs with

Python and R scripts available at https://github.com/LarracuenteLab/Dmel_Jockey-3_Evolution.

We repeated our analyses of TEs with and without insertions in known piRNA clusters and

removed fragments of Gypsy6_I_Dmoj and Gypsy1_I_Dmoj which resemble satellite sequences

in *D. melanogaster*. Depth of each short-read sample was calculated from the sorted BAM outputs

from the McClintock pipeline using Mosdepth [145] (Table S3). We retained samples which had

a centromere coverage greater than 20X for our analyses (see Supplemental Text for more

information).

We used the same short-read FASTQ files from the DGRP to quantify genetic diversity between

the centromere islands and the rest of the genome using a best practices GATK pipeline v4.1.9.0

to detect variant SNPs. We mapped reads from high coverage samples to our reference genome

with BWA v1.7.15 and samtools 1.7 to generate sorted BAM files and used GATK

MarkDuplicates, BaseRecalibrator, ApplyBQSR, AnalyzeCovariates, and HaplotypeCaller to call

and filter SNPs for each sample into a VCF file. The VCFs for all samples were concatenated with

GenotypeGVCFs and relicabrated with VariantRecalibrator and ApplyVQSR. We quality filtered

SNPs in the resulting GVCF file with the following criteria; QD < 2.0, FS > 60.0, MQ < 40.0,

MQRankSum < -12.5, and ReadPosRankSum < -8.0. We calculated π and Tajima's *D* with the

SimplifyVCF_Basic.v2.pl and Windows_Basic.v2.pl Perl scripts on the filtered SNPs (available

at https://github.com/bnavarrodominguez/SD-Population-genomics, [146]. We further analyzed

variants within the centromeres with VCFtools [147] and the vcfR R package [148]. Additional

details of our analysis with short-read data and the results can be found in the Supplemental Text

section.

Long-read analysis

We assembled genomes for several *D. melanogaster* lines following the heterochromatin-enriched

assembly protocol from Chang & Larracuente [73]. We downloaded raw PacBio reads for five of

the Global Diversity Lines (GDL) from the NBCI BioProject PRJNA451471 [68], eight

Drosophila Synthetic Population Resources (DSPR) founder lines and Oregon-R with raw Pacbio

reads from BioProject PRJNA418342 [67], and five DGRP lines from Bioproject PRJNA559813

[66]. We assembled draft genomes using all reads with Canu v1.7.1 [149] with the relaxed

corrected   error   rate   and   the   parameters   "genomeSize=160m   useGrid=false

stopOnReadQuality=false corMinCoverage=0 corOutCoverage=100 correctedErrorRate=0.100".

We then mapped PacBio reads to the reference genome with minimap2 (downloaded August 28,

2018) with the "-x map-bp" parameter setting [150]. We then used unmapped reads and reads

mapping to heterochromatic regions to generate an additional "heterochromatin-enriched"

assembly. We merged the two assemblies into a single assembly with quickmerge [151] and

polished the assemblies with Racon v1.3.1 [152] by mapping the PacBio reads three times to the

assembly using minimap2 [150]. We also polished the GDL assemblies twice with Illumina reads

from BioProject PRJNA268111 [144], the DSPR assemblies with reads from

https://wfitch.bio.uci.edu/~dspr/index.html (last accessed September 6, 2021 [153]), BioProject

PRJNA252961 for Oregon-R, and the DGRP assemblies with reads from BioProject PRJNA36679

[64, 65]. We annotated all assemblies with Repeatmasker v4.1.0 and our *Drosophila* TE library

with the updated *G2/Jockey-3* consensus sequence [135] (File S10). We identified centromere

island sequences in assemblies using a combination of methods to find sequences corresponding

to the centromere islands identified in our reference genome [53]. These include aligning

assemblies to the centromere islands with MUMMER v.3.23 nucmer [154], identification of

similar sequences with BLAST 2.10.0+ [155], and identifying contigs enriched in *G2/Jockey-3* or

satellite DNA repeats enriched around the centromere islands [53]. We also used MEME [156] to

detect motifs enriched in the flanking regions of young insertions in our assemblies. Additional

details of our analysis with long-read data, the results, and the MEME motif analysis are in the

Supplemental Text.

Insertion polymorphism validation

We validated TE insertion predictions in three ways: 1) we used short reads to confirm TE

predictions in long read assemblies (see Supplemental Text); 2) we used long read assemblies to

confirm TE predictions in short read data (see Supplemental Text); and 3) we selected a subset to

validate with PCR (File S3). We used the results from these analyses to establish criteria for

including a predicted insertion in our analyses. For PCR, we selected a subset of 59 *G2/Jockey-3*

copies to confirm in four GDL lines; B59, I23, T29A, and ZH26. We included copies from each

chromosome (X, 2, 3, 4), each chromatin region (euchromatin, heterochromatin, and centromere),

and both reference and non-reference copies. We were able to detect 33 copies including 13 copies

in the centromere and seven *de novo* insertions (File S3).

G2/Jockey-3 sequence and phylogenetic analysis in D. melanogaster

We extracted all *G2/Jockey-3* sequences from the annotated assemblies using BEDTools v2.29.2

[136] and classified the insertions as reference or non-reference (*de novo*). We classified insertions

as homologous to those in our reference if they were both fragments of similar size (+/- 10 bp) and

within +100 bp of the same location in the reference genome. We identified corresponding

locations by aligning the assemblies to the reference genome with nucmer within MUMMER

v.3.23 [154]. Insertions were classified as *de novo* if they differed in fragment size length and

position of reference insertions, found outside the reference fragment position range, and located

within breaks in the nucmer alignment to the reference genome.

We performed phylogenetic analyses for candidate *G2/Jockey-3* sequences identified in our

genome assemblies. We used *G2/Jockey-3* sequences from the reference and the non-reference

insertions discovered in the DSPR and GDL assemblies. If a non-reference insertion was found in

multiple assemblies, we randomly selected an assembly from which to extract that insertion to

include in our analysis. Prior to phylogenetic analysis, we filtered sequences to exclude fragments

shorter than 400 bp or ~10% of the length of a full *G2/Jockey-3* element. We aligned the sequences

with MAFFT [157] with the *D. simulans G2/Jockey-3* consensus as the outgroup and inferred a

maximum likelihood phylogenetic tree with RAxML v8.2.11 [158]. To study the evolution of

G2/Jockey-3 on the Y chromosome, we inferred a maximum likelihood phylogeny of all fragments

exclusive to the Y chromosome regardless of size. We determined the relative age of individual

insertions using divergence from the consensus from the Repeatmasker output and the number of

substitutions unique to a fragment divided by fragment length. "Young" insertions were either <

1% diverged from the consensus sequence or *de novo* insertions exclusive to a single genome

assembly and < 2% diverged from the consensus sequence. We calculated pairwise divergence

between fragments with MEGA X [159] using the Kimura 2-parameter model assuming uniform

rates among all sites. We estimated gene conversion rates between elements using the same method

outlined from Chang & Larracuente [73] based on Ohta's α equation [160] and Rozen's equation

[161]. We used the following equation for our calculation of gene conversion:

706
$$c = \frac{2u(n-1)}{\pi}$$

We determined the rate at which a number of sequences (*n*) homogenize each other per generation

(*c*) based on the point mutation rate (*u*) and divergence among the sequences (*π*). We grouped

reference *G2/Jockey-3* sequences located within the same contig except for insertions on the Y

centromere. Fragments from the Y centromere were grouped based on the phylogenetically-

defined classes. We aligned the sequences with MAFFT [157] and calculated divergence with

MEGA X [159]. We normalized *c* based on gene conversion tract length of a single event (c_g)

based on previously estimated 400 bp tract length of a single event in *D. melanogaster* [92, 93]:

714
$$c_g = \frac{c}{400 \; x \; 2}$$

Age-of-allele test

Natural selection, transposition, and frequency of insertion influence TE evolution within

populations. We used an age-of-allele test [70] to determine departure from neutral evolution of

individual insertions based on population frequency, age, and demographic history in *D.*

melanogaster. The age-of-allele test determines the probability distribution of an insertion

frequency within a population under neutrality conditioned on the age of the insertion. The age is

inferred by the number of unique substitutions within a TE sequence accumulated since the time

of the insertion. To determine the number of unique substitutions, we aligned insertions with

MAFFT [157] for *G2/Jockey-3*, *BS*, *G5*, *Doc*, and *Doc2* detected as reference insertions in the

McClintock output. Unique substitutions were quantified from each individual insertion using

Biostrings [162]. We calculated the probability distribution of the frequency of each unique

insertion with the age-of-allele test script from Blumenstiel et al. [70] in Mathematica v12.0.0.0.

The probability distribution is also conditioned on population demography of North American *D.*

melanogaster populations outlined in the original paper [70]. North American *D. melanogaster*

populations experienced two bottlenecks in their migration from Africa [163, 164] that led to non-

equilibrium TE evolution [70]. We compared the probability distribution to the frequency of

insertions detected in the 28 DGRP samples with >25X average coverage of the centromere

islands. We excluded insertions in the Y chromosome given the high rate of duplication. The

expected frequency of each insertion in the population is the probability of observing an insertion

as frequent or more frequent within the population multiplied by the observed number of insertions

in our population sample. We calculated significant deviations from neutrality as probability of

observing as many or fewer, or as many or greater, copies < 0.1.

DATA AVAILABILITY STATEMENT

All scripts for data analysis and data for figure generation are available on Github at https://github.com/LarracuenteLab/Dmel_Jockey-3_Evolution. The genome assemblies, genome annotations, and large datasets including the McClintock outputs will be available on Dryad (DOI forthcoming).

ACKNOWLEDGEMENTS

This work was funded by the National Science Foundation (NSF MCB 1844693) and University of Rochester funds to AML. We thank the Mellone lab, Dr. Justin Blumenstiel, and the Larracuente lab for helpful discussion. We also thank the University of Rochester Center for Integrated Resource Computing for access to computing cluster resources.

REFERENCES

1.	Allshire RC, Karpen GH: **Epigenetic regulation of centromeric chromatin: old dogs, new tricks?** *Nat Rev Genet* 2008, **9:**923-937.
2.	Garrido-Ramos MA: **Satellite DNA: An Evolving Topic.** *Genes* 2017, **8**.
3.	Hartley G, O'Neill RJ: **Centromere Repeats: Hidden Gems of the Genome.** *Genes* 2019, **10**.
4.	Pidoux AL, Allshire RC: **Kinetochore and heterochromatin domains of the fission yeast centromere.** *Chromosome Res* 2004, **12:**521-534.
5.	Ohzeki J-I, Bergmann JH, Kouprina N, Noskov VN, Nakano M, Kimura H, Earnshaw WC, Larionov V, Masumoto H: **Breaking the HAC Barrier: histone H3K9 acetyl/methyl balance regulates CENP-A assembly.** *EMBO J* 2012, **31:**2391-2402.
6.	Choo KH: **Centromerization.** *Trends Cell Biol* 2000, **10:**182-188.
7.	Henikoff S, Dalal Y: **Centromeric chromatin: what makes it unique?** *Curr Opin Genet Dev* 2005, **15:**177-184.

8.   Henikoff S, Furuyama T: **Epigenetic inheritance of centromeres.** *Cold Spring Harb*
Symp Quant Biol 2010, **75:**51-60.

9.   Black BE, Cleveland DW: **Epigenetic centromere propagation and the nature of**
CENP-a nucleosomes. *Cell* 2011, **144:**471-479.

10.   Mendiburo MJ, Padeken J, Fülöp S, Schepers A, Heun P: **Drosophila CENH3 is**
sufficient for centromere formation. *Science* 2011, **334:**686-690.

11.   McKinley KL, Cheeseman IM: **The molecular basis for centromere identity and**
function. *Nat Rev Mol Cell Biol* 2016, **17:**16-29.

12.   Blower MD, Sullivan BA, Karpen GH: **Conserved Organization of Centromeric**
Chromatin in Flies and Humans. *Developmental Cell* 2002, **2:**319-330.

13.   Ribeiro SA, Vagnarelli P, Dong Y, Hori T, McEwen BF, Fukagawa T, Flors C, Earnshaw
WC: **A super-resolution map of the vertebrate kinetochore.** *Proc Natl Acad Sci U S A*
2010, **107:**10484-10489.

14.   Bergmann JH, Rodríguez MG, Martins NMC, Kimura H, Kelly DA, Masumoto H,
Larionov V, Jansen LET, Earnshaw WC: **Epigenetic engineering shows H3K4me2 is**
required for HJURP targeting and CENP-A assembly on a synthetic human
kinetochore. *EMBO J* 2011, **30:**328-340.

15.   Martins NMC, Bergmann JH, Shono N, Kimura H, Larionov V, Masumoto H, Earnshaw
WC: **Epigenetic engineering shows that a human centromere resists silencing**
mediated by H3K27me3/K9me3. *Mol Biol Cell* 2016, **27:**177-196.

16.   Lo AW, Magliano DJ, Sibson MC, Kalitsis P, Craig JM, Choo KH: **A novel chromatin**
immunoprecipitation and array (CIA) analysis identifies a 460-kb CENP-A-binding
neocentromere DNA. *Genome Res* 2001, **11:**448-457.

17.   Alonso A, Fritz B, Hasson D, Abrusan G, Cheung F, Yoda K, Radlwimmer B, Ladurner
AG, Warburton PE: **Co-localization of CENP-C and CENP-H to discontinuous**
domains of CENP-A chromatin at human neocentromeres. *Genome Biol* 2007,
8:R148.

18.   Hasson D, Panchenko T, Salimian KJ, Salman MU, Sekulic N, Alonso A, Warburton PE,
Black BE: **The octamer is the major form of CENP-A nucleosomes at human**
centromeres. *Nat Struct Mol Biol* 2013, **20:**687-695.

19.   Nergadze SG, Piras FM, Gamba R, Corbo M, Cerutti F, McCarter JGW, Cappelletti E,
Gozzo F, Harman RM, Antczak DF, et al: **Birth, evolution, and transmission of**
satellite-free mammalian centromeric domains. *Genome Res* 2018, **28:**789-799.

20.   Kasai F, Garcia C, Arruga MV, Ferguson-Smith MA: **Chromosome homology between**
chicken (*Gallus gallus domesticus*) and the red-legged partridge (Alectoris rufa);
evidence of the occurrence of a neocentromere during evolution. *Cytogenet Genome*
Res 2003, **102:**326-330.

21.   Marshall OJ, Chueh AC, Wong LH, Choo KHA: **Neocentromeres: new insights into**
centromere structure, disease development, and karyotype evolution. *Am J Hum*
Genet 2008, **82:**261-282.

22.   Rocchi M, Archidiacono N, Schempp W, Capozzi O, Stanyon R: **Centromere**
repositioning in mammals. *Heredity* 2012, **108:**59-67.

23.   Miller JT, Dong F, Jackson SA, Song J, Jiang J: **Retrotransposon-related DNA**
sequences in the centromeres of grass chromosomes. *Genetics* 1998, **150:**1615-1623.

24.   Presting GG, Malysheva L, Fuchs J, Schubert I: **A Ty3/gypsy retrotransposon-like**
sequence localizes to the centromeric regions of cereal chromosomes. *Plant J* 1998,
16:721-728.

25.   Cheng Z, Dong F, Langdon T, Ouyang S, Buell CR, Gu M, Blattner FR, Jiang J:
Functional rice centromeres are marked by a satellite repeat and a centromere-
specific retrotransposon. *Plant Cell* 2002, **14:**1691-1704.

26.   Nagaki K, Cheng Z, Ouyang S, Talbert PB, Kim M, Jones KM, Henikoff S, Buell CR,
Jiang J: **Sequencing of a rice centromere uncovers active genes.** *Nat Genet* 2004,
36:138-145.

27.   Nagaki K, Song J, Stupar RM, Parokonny AS, Yuan Q, Ouyang S, Liu J, Hsiao J, Jones
KM, Dawe RK, et al: **Molecular and cytological analyses of large tracks of**
centromeric DNA reveal the structure and evolutionary dynamics of maize
centromeres. *Genetics* 2003, **163:**759-770.

28.   Ferreri GC, Brown JD, Obergfell C, Jue N, Finn CE, O'Neill MJ, O'Neill RJ: **Recent**
amplification of the kangaroo endogenous retrovirus, KERV, limited to the
centromere. *J Virol* 2011, **85:**4761-4771.

29.   Ferreri GC, Marzelli M, Rens W, O'Neill RJ: **A centromere-specific retroviral element**
associated with breaks of synteny in macropodine marsupials. *Cytogenet Genome Res*
2004, **107:**115-118.

30.   de Sotero-Caio CG, Cabral-de-Mello DC, Calixto MdS, Valente GT, Martins C, Loreto
V, de Souza MJ, Santos N: **Centromeric enrichment of LINE-1 retrotransposons and**
its significance for the chromosome evolution of Phyllostomid bats. *Chromosome Res*
2017, **25:**313-325.

31.   Carbone L, Harris RA, Gnerre S, Veeramah KR, Lorente-Galdos B, Huddleston J, Meyer
TJ, Herrero J, Roos C, Aken B, et al: **Gibbon genome and the fast karyotype evolution**
of small apes. *Nature* 2014, **513:**195-201.

32.   Carbone L, Harris RA, Mootnick AR, Milosavljevic A, Martin DIK, Rocchi M, Capozzi
O, Archidiacono N, Konkel MK, Walker JA, et al: **Centromere remodeling in *Hoolock***
***leuconedys* (Hylobatidae) by a new transposable element unique to the gibbons.**
Genome Biol Evol 2012, **4:**648-658.

33.   Contreras-Galindo R, Kaplan MH, He S, Contreras-Galindo AC, Gonzalez-Hernandez
MJ, Kappes F, Dube D, Chan SM, Robinson D, Meng F, et al: **HIV infection reveals**
widespread expansion of novel centromeric human endogenous retroviruses.
Genome Res 2013, **23:**1505-1513.

34.   Zahn J, Kaplan MH, Fischer S, Dai M, Meng F, Saha AK, Cervantes P, Chan SM, Dube
D, Omenn GS, et al: **Expansion of a novel endogenous retrovirus throughout the**
pericentromeres of modern humans. *Genome Biol* 2015, **16:**74.

35.   Klein SJ, O'Neill RJ: **Transposable elements: genome innovation, chromosome**
diversity, and centromere conflict. *Chromosome Res* 2018, **26:**5-23.

36.   Birchler JA, Presting GG: **Retrotransposon insertion targeting: a mechanism for**
homogenization of centromere sequences on nonhomologous chromosomes. *Genes*
Dev 2012, **26:**638-640.

37.   Sultana T, Zamborlini A, Cristofari G, Lesage P: **Integration site selection by**
retroviruses and transposable elements in eukaryotes. *Nature Reviews Genetics* 2017,
18:292-308.

38. Wong LH, Choo KHA: **Evolutionary dynamics of transposable elements at the**
centromere. *Trends Genet* 2004, **20:**611-616.
39. Sharma A, Wolfgruber TK, Presting GG: **Tandem repeats derived from centromeric**
retrotransposons. *BMC Genomics* 2013, **14**.
40. Bourc'his D, Bestor TH: **Meiotic catastrophe and retrotransposon reactivation in**
male germ cells lacking Dnmt3L. *Nature* 2004, **431:**96-99.
41. Kato M, Takashima K, Kakutani T: **Epigenetic control of CACTA transposon mobility**
in Arabidopsis thaliana. *Genetics* 2004, **168:**961-969.
42. Ariumi Y: **Guardian of the Human Genome: Host Defense Mechanisms against**
LINE-1 Retrotransposition. *Front Chem* 2016, **4:**28.
43. Chan FL, Wong LH: **Transcription in the maintenance of centromere chromatin**
identity. *Nucleic Acids Res* 2012, **40:**11178-11188.
44. Wong LH, Brettingham-Moore KH, Chan L, Quach JM, Anderson MA, Northrop EL,
Hannan R, Saffery R, Shaw ML, Williams E, Choo KHA: **Centromere RNA is a key**
component for the assembly of nucleoproteins at the nucleolus and centromere.
Genome Res 2007, **17:**1146-1160.
45. Quénet D, Dalal Y: **A long non-coding RNA is required for targeting centromeric**
protein A to the human centromere. *Elife* 2014, **3:**e03254.
46. Catania S, Pidoux AL, Allshire RC: **Sequence features and transcriptional stalling**
within centromere DNA promote establishment of CENP-A chromatin. *PLoS Genet*
2015, **11:**e1004986.
47. Chen C-C, Bowers S, Lipinszki Z, Palladino J, Trusiak S, Bettini E, Rosin L, Przewloka
MR, Glover DM, O'Neill RJ, Mellone BG: **Establishment of Centromeric Chromatin**
by the CENP-A Assembly Factor CAL1 Requires FACT-Mediated Transcription.
Dev Cell 2015, **34:**73-84.
48. Hall LE, Mitchell SE, O'Neill RJ: **Pericentric and centromeric transcription: a**
perfect balance required. *Chromosome Research* 2012, **20:**535-546.
49. Talbert PB, Henikoff S: **Transcribing Centromeres: Noncoding RNAs and**
Kinetochore Assembly. *Trends in Genetics* 2018, **34:**587-599.
50. Chuong EB, Elde NC, Feschotte C: **Regulatory activities of transposable elements:**
from conflicts to benefits. *Nat Rev Genet* 2017, **18:**71-86.
51. Jain M, Olsen HE, Turner DJ, Stoddart D, Bulazel KV, Paten B, Haussler D, Willard HF,
Akeson M, Miga KH: **Linear assembly of a human centromere on the Y**
chromosome. *Nature Biotechnology* 2018, **36:**321-323.
52. Logsdon GA, Vollger MR, Hsieh P, Mao Y, Liskovykh MA, Koren S, Nurk S, Mercuri
L, Dishuck PC, Rhie A, et al: **The structure, function and evolution of a complete**
human chromosome 8. *Nature* 2021, **593:**101-107.
53. Chang C-H, Chavan A, Palladino J, Wei X, Martins NMC, Santinello B, Chen C-C,
Erceg J, Beliveau BJ, Wu C-T, et al: **Islands of retroelements are major components**
of Drosophila centromeres. *PLoS Biol* 2019, **17:**e3000241.
54. Belser C, Baurens F-C, Noel B, Martin G, Cruaud C, Istace B, Yahiaoui N, Labadie K,
Hřibová E, Doležel J, et al: **Telomere-to-telomere gapless chromosomes of banana**
using nanopore sequencing. *Commun Biol* 2021, **4:**1047.
55. Wang B, Yang X, Jia Y, Xu Y, Jia P, Dang N, Wang S, Xu T, Zhao X, Gao S, et al:
High-quality Arabidopsis thaliana Genome Assembly with Nanopore and HiFi Long
Reads. *Genomics Proteomics Bioinformatics* 2021.

56. Naish M, Alonge M, Wlodzimierz P, Tock AJ, Abramson BW, Schmucker A,
Mandakova T, Jamge B, Lambing C, Kuo P, et al: **The genetic and epigenetic**
landscape of the Arabidopsis centromeres. *Science* 2021, **374:**eabi7489.
57. Hales KG, Korey CA, Larracuente AM, Roberts DM: **Genetics on the Fly: A Primer on**
the Drosophila Model System. *Genetics* 2015, **201:**815-842.
58. Rahman R, Chirn G-W, Kanodia A, Sytnikova YA, Brembs B, Bergman CM, Lau NC:
Unique transposon landscapes are pervasive across Drosophila melanogaster
genomes. *Nucleic Acids Res* 2015, **43:**10655-10672.
59. Gerstein MB, Rozowsky J, Yan K-K, Wang D, Cheng C, Brown JB, Davis CA, Hillier L,
Sisu C, Li JJ, et al: **Comparative analysis of the transcriptome across distant species.**
Nature 2014, **512:**445-448.
60. Andersen PR, Tirian L, Vunjak M, Brennecke J: **A heterochromatin-dependent**
transcription machinery drives piRNA expression. *Nature* 2017, **549:**54-59.
61. Laktionov PP, Maksimov DA, Romanov SE, Antoshina PA, Posukh OV, White-Cooper
H, Koryakov DE, Belyakin SN: **Genome-wide analysis of gene regulation mechanisms**
during Drosophila spermatogenesis. *Epigenetics Chromatin* 2018, **11:**14.
62. Tettey TT, Gao X, Shao W, Li H, Story BA, Chitsazan AD, Glaser RL, Goode ZH,
Seidel CW, Conaway RC, et al: **A Role for FACT in RNA Polymerase II Promoter-**
Proximal Pausing. *Cell Rep* 2019, **27:**3770-3779.e3777.
63. Metcalfe CJ, Casane D: **Modular organization and reticulate evolution of the ORF1**
of Jockey superfamily transposable elements. *Mobile DNA* 2014, **5:**19.
64. Huang W, Massouras A, Inoue Y, Peiffer J, Ràmia M, Tarone AM, Turlapati L, Zichner
T, Zhu D, Lyman RF, et al: **Natural variation in genome architecture among 205**
Drosophila melanogaster **Genetic Reference Panel lines.** *Genome Res* 2014, **24:**1193-
1208.
65. Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y,
Magwire MM, Cridland JM, et al: **The** *Drosophila melanogaster* **Genetic Reference**
Panel. *Nature* 2012, **482:**173-178.
66. Rech GE, Radío S, Guirao-Rico S, Aguilera L, Horvath V, Green L, Lindstadt H,
Jamilloux V, Quesneville H, González J: **Population-scale long-read sequencing**
uncovers transposable elements associated with gene expression variation and
adaptive signatures in Drosophila. *Nat Commun* 2022, **13:**1948.
67. Chakraborty M, Emerson JJ, Macdonald SJ, Long AD: **Structural variants exhibit**
widespread allelic heterogeneity and shape variation in complex traits. *Nat Commun*
2019, **10:**4872.
68. Long E, Evans C, Chaston J, Udall JA: **Genomic Structural Variations Within Five**
Continental Populations of *Drosophila melanogaster.* *G3: Genes|Genomes|Genetics*
2018, **8:**3247-3253.
69. Nelson MG, Linheiro RS, Bergman CM: **McClintock: An Integrated Pipeline for**
Detecting Transposable Element Insertions in Whole-Genome Shotgun Sequencing
Data. *G3* 2017, **7:**2763-2778.
70. Blumenstiel JP, Chen X, He M, Bergman CM: **An age-of-allele test of neutrality for**
transposable element insertions. *Genetics* 2014, **196:**523-538.
71. Kofler R, Betancourt AJ, Schlötterer C: **Sequencing of pooled DNA samples (Pool-Seq)**
uncovers complex dynamics of transposable element insertions in *Drosophila*
melanogaster. *PLoS Genet* 2012, **8:**e1002487.

72. Kofler R, Nolte V, Schlötterer C: **Tempo and Mode of Transposable Element Activity**
in Drosophila. *PLoS Genet* 2015, **11:**e1005406.

73. Chang C-H, Larracuente AM: **Heterochromatin-Enriched Assemblies Reveal the**
Sequence and Organization of the Y Chromosome. *Genetics* 2019, **211:**333-348.

74. Ghildiyal M, Zamore PD: **Small silencing RNAs: an expanding universe.** *Nat Rev*
Genet 2009, **10:**94-108.

75. Ernst C, Odom DT, Kutter C: **The emergence of piRNAs against transposon invasion**
to preserve mammalian genome integrity. *Nature Communications* 2017, **8**.

76. Bergman CM, Quesneville H, Anxolabéhère D, Ashburner M: **Recurrent insertion and**
duplication generate networks of transposable element sequences in the *Drosophila*
melanogaster **genome.** *Genome Biol* 2006, **7:**R112.

77. Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ:
Discrete small RNA-generating loci as master regulators of transposon activity in
Drosophila. *Cell* 2007, **128:**1089-1103.

78. Gunawardane LS, Saito K, Nishida KM, Miyoshi K, Kawamura Y, Nagami T, Siomi H,
Siomi MC: **A slicer-mediated mechanism for repeat-associated siRNA 5' end**
formation in Drosophila. *Science* 2007, **315:**1587-1590.

79. Mohn F, Sienski G, Handler D, Brennecke J: **The rhino-deadlock-cutoff complex**
licenses noncanonical transcription of dual-strand piRNA clusters in Drosophila.
Cell 2014, **157:**1364-1379.

80. Parhad SS, Tu S, Weng Z, Theurkauf WE: **Adaptive Evolution Leads to Cross-Species**
Incompatibility in the piRNA Transposon Silencing Machinery. *Dev Cell* 2017,
43:60-70.e65.

81. Rozhkov NV, Aravin AA, Zelentsova ES, Schostak NG, Sachidanandam R, McCombie
WR, Hannon GJ, Evgen'ev MB: **Small RNA-based silencing strategies for transposons**
in the process of invading Drosophila species. *RNA* 2010, **16:**1634-1645.

82. Quénerch'du E, Anand A, Kai T: **The piRNA pathway is developmentally regulated**
during spermatogenesis in Drosophila. *RNA* 2016, **22:**1044-1054.

83. Wei X, Eickbush DG, Speece I, Larracuente AM: **Heterochromatin-dependent**
transcription of satellite DNAs in the female germline. *Elife* 2021, **10**.

84. Petrov DA, Fiston-Lavier A-S, Lipatov M, Lenkov K, González J: **Population genomics**
of transposable elements in *Drosophila melanogaster***.** *Mol Biol Evol* 2011, **28:**1633-
1644.

85. Montgomery E, Charlesworth B, Langley CH: **A test for the role of natural selection in**
the stabilization of transposable element copy number in a population of *Drosophila*
*melanogaster***.** *Genetical Research* 1987, **49:**31-41.

86. Charlesworth B, Campos JL: **The relations between recombination rate and patterns**
of molecular variation and evolution in Drosophila. *Annu Rev Genet* 2014, **48:**383-
403.

87. Kent TV, Uzunović J, Wright SI: **Coevolution between transposable elements and**
recombination. *Philos Trans R Soc Lond B Biol Sci* 2017, **372**.

88. Bergman CM, Bensasson D: **Recent LTR retrotransposon insertion contrasts with**
waves of non-LTR insertion since speciation in *Drosophila melanogaster***.** *Proc Natl*
Acad Sci U S A 2007, **104:**11340-11345.

89. Blumenstiel JP, Hartl DL, Lozovsky ER: **Patterns of insertion and deletion in**
contrasting chromatin domains. *Mol Biol Evol* 2002, **19:**2211-2225.

90. Chang CH, Gregory LE, Gordon KE, Meiklejohn CD, Larracuente AM: **Unique structure and positive selection promote the rapid divergence of Drosophila Y chromosomes.** *Elife* 2022, **11**.

91. Comeron JM, Ratnappan R, Bailin S: **The many landscapes of recombination in Drosophila melanogaster.** *PLoS Genet* 2012, **8:**e1002905.

92. Miller DE, Smith CB, Kazemi NY, Cockrell AJ, Arvanitakis AV, Blumenstiel JP, Jaspersen SL, Hawley RS: **Whole-Genome Analysis of Individual Meiotic Events in** *Drosophila melanogaster* **Reveals That Noncrossover Gene Conversions Are Insensitive to Interference and the Centromere Effect.** *Genetics* 2016, **203:**159-171.

93. Miller DE, Takeo S, Nandanan K, Paulson A, Gogol MM, Noll AC, Perera AG, Walton KN, Gilliland WD, Li H, et al: **A Whole-Chromosome Analysis of Meiotic Recombination in** *Drosophila melanogaster.* *G3* 2012, **2:**249-260.

94. Khurana JS, Wang J, Xu J, Koppetsch BS, Thomson TC, Nowosielska A, Li C, Zamore PD, Weng Z, Theurkauf WE: **Adaptation to P Element Transposon Invasion in** *Drosophila melanogaster.* *Cell* 2011, **147:**1551-1563.

95. Kofler R: **Dynamics of Transposable Element Invasions with piRNA Clusters.** *Mol Biol Evol* 2019, **36:**1457-1472.

96. Said I, McGurk MP, Clark AG, Barbash DA: **Patterns of piRNA Regulation in Drosophila Revealed through Transposable Element Clade Inference.** *Mol Biol Evol* 2022, **39**.

97. Kofler R, Senti K-A, Nolte V, Tobler R, Schlötterer C: **Molecular dissection of a natural transposable element invasion.** *Genome Res* 2018, **28:**824-835.

98. Anxolabéhère D, Kidwell MG, Periquet G: **Molecular characteristics of diverse populations are consistent with the hypothesis of a recent invasion of** *Drosophila melanogaster* **by mobile P elements.** *Mol Biol Evol* 1988, **5:**252-269.

99. Hill T, Schlotterer C, Betancourt AJ: **Hybrid Dysgenesis in** *Drosophila simulans* **Associated with a Rapid Invasion of the P-Element.** *PLoS Genet* 2016, **12:**e1005920.

100. Wolfgruber TK, Sharma A, Schneider KL, Albert PS, Koo D-H, Shi J, Gao Z, Han F, Lee H, Xu R, et al: **Maize centromere structure and evolution: sequence analysis of centromeres 2 and 5 reveals dynamic Loci shaped primarily by retrotransposons.** *PLoS Genet* 2009, **5:**e1000743.

101. Schneider KL, Xie Z, Wolfgruber TK, Presting GG: **Inbreeding drives maize centromere evolution.** *Proc Natl Acad Sci U S A* 2016, **113:**E987-996.

102. Tsukahara S, Kawabe A, Kobayashi A, Ito T, Aizu T, Shin-i T, Toyoda A, Fujiyama A, Tarutani Y, Kakutani T: **Centromere-targeted de novo integrations of an LTR retrotransposon of** *Arabidopsis lyrata.* *Genes Dev* 2012, **26:**705-713.

103. Tsukahara S, Kobayashi A, Kawabe A, Mathieu O, Miura A, Kakutani T: **Bursts of retrotransposition reproduced in Arabidopsis.** *Nature* 2009, **461:**423-426.

104. Hollister JD, Gaut BS: **Epigenetic silencing of transposable elements: a trade-off between reduced transposition and deleterious effects on neighboring gene expression.** *Genome Res* 2009, **19:**1419-1428.

105. Lee YCG: **The Role of piRNA-Mediated Epigenetic Silencing in the Population Dynamics of Transposable Elements in** *Drosophila melanogaster.* *PLoS Genet* 2015, **11:**e1005269.

106. Srivastava S, Foltz DR: **Posttranslational modifications of CENP-A: marks of distinction.** *Chromosoma* 2018, **127:**279-290.

107. Czech B, Munafò M, Ciabrelli F, Eastwood EL, Fabry MH, Kneuss E, Hannon GJ: **piRNA-Guided Genome Defense: From Biogenesis to Silencing.** *Annu Rev Genet* 2018, **52:**131-157.

108. Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD: **PIWI-interacting RNAs: small RNAs with big functions.** *Nat Rev Genet* 2019, **20:**89-108.

109. Onishi R, Yamanaka S, Siomi MC: **piRNA- and siRNA-mediated transcriptional repression in Drosophila, mice, and yeast: new insights and biodiversity.** *EMBO Rep* 2021, **22:**e53062.

110. Scelfo A, Fachinetti D: **Keeping the Centromere under Control: A Promising Role for DNA Methylation.** *Cells* 2019, **8**.

111. Altemose N, Logsdon GA, Bzikadze AV, Sidhwani P, Langley SA, Caldas GV, Hoyt SJ, Uralsky L, Ryabov FD, Shew CJ, et al: **Complete genomic and epigenetic maps of human centromeres.** *Science* 2022, **376:**eabl4178.

112. Doolittle WF, Ford Doolittle W, Sapienza C: **Selfish genes, the phenotype paradigm and genome evolution.** *Nature* 1980, **284:**601-603.

113. Orgel LE, Crick FH: **Selfish DNA: the ultimate parasite.** *Nature* 1980, **284:**604-607.

114. Dimitri P, Junakovic N: **Revising the selfish DNA hypothesis: new evidence on accumulation of transposable elements in heterochromatin.** *Trends Genet* 1999, **15:**123-124.

115. Cosby RL, Chang N-C, Feschotte C: **Host-transposon interactions: conflict, cooperation, and cooption.** *Genes Dev* 2019, **33:**1098-1116.

116. Danilevskaya ON, Slot F, Traverse KL, Hogan NC, Pardue ML: **Drosophila telomere transposon HeT-A produces a transcript with tightly bound protein.** *Proc Natl Acad Sci U S A* 1994, **91:**6679-6682.

117. Mason JM, Biessmann H: **The unusual telomeres of Drosophila.** *Trends Genet* 1995, **11:**58-62.

118. Pardue M-L, DeBaryshe PG: **Drosophila telomeres: A variation on the telomerase theme.** *Fly* 2008, **2:**101-110.

119. Pardue ML, Debaryshe PG: **Drosophila telomere transposons: genetically active elements in heterochromatin.** *Genetica* 2000, **109:**45-52.

120. Saint-Leandre B, Nguyen SC, Levine MT: **Diversification and collapse of a telomere elongation mechanism.** *Genome Res* 2019, **29:**920-931.

121. McGurk MP, Dion-Côté A-M, Barbash DA: **Rapid evolution at the Drosophila telomere: transposable element dynamics at an intrinsically unstable locus.** *Genetics* 2021, **217**.

122. Moazed D, Bühler M, Buker SM, Colmenares SU, Gerace EL, Gerber SA, Hong EJE, Motamedi MR, Verdel A, Villén J, Gygi SP: **Studies on the mechanism of RNAi-dependent heterochromatin assembly.** *Cold Spring Harb Symp Quant Biol* 2006, **71:**461-471.

123. Choi ES, Strålfors A, Castillo AG, Durand-Dubief M, Ekwall K, Allshire RC: **Identification of noncoding transcripts from within CENP-A chromatin at fission yeast centromeres.** *J Biol Chem* 2011, **286:**23600-23607.

124. Chan FL, Marshall OJ, Saffery R, Kim BW, Earle E, Choo KHA, Wong LH: **Active transcription and essential role of RNA polymerase II at the centromere during mitosis.** *Proc Natl Acad Sci U S A* 2012, **109:**1979-1984.

125. Rošić S, Köhler F, Erhardt S: **Repetitive centromeric satellite RNA is essential for**
kinetochore formation and cell division. *J Cell Biol* 2014, **207:**335-349.
126. Bobkov GOM, Gilbert N, Heun P: **Centromere transcription allows CENP-A to**
transit from chromatin association to stable incorporation. *J Cell Biol* 2018,
217:1957-1972.
127. Ohkuni K, Kitagawa K: **Endogenous transcription at the centromere facilitates**
centromere activity in budding yeast. *Curr Biol* 2011, **21:**1695-1703.
128. Sandler L, Novitski E: **Meiotic Drive as an Evolutionary Force.** *The American*
Naturalist 1957, **91:**105-110.
129. Akera T, Chmátal L, Trimm E, Yang K, Aonbangkhen C, Chenoweth DM, Janke C,
Schultz RM, Lampson MA: **Spindle asymmetry drives non-Mendelian chromosome**
segregation. *Science* 2017, **358:**668-672.
130. Akera T, Trimm E, Lampson MA: **Molecular Strategies of Meiotic Cheating by Selfish**
Centromeres. *Cell* 2019, **178:**1132-1144.e1110.
131. Henikoff S, Ahmad K, Malik HS: **The centromere paradox: stable inheritance with**
rapidly evolving DNA. *Science* 2001, **293:**1098-1102.
132. Malik HS, Henikoff S: **Adaptive Evolution of Cid, a Centromere-Specific Histone in**
Drosophila. *Genetics* 2001, **157:**1293-1298.
133. Kumon T, Ma J, Akins RB, Stefanik D, Nordgren CE, Kim J, Levine MT, Lampson MA:
Parallel pathways for recruiting effector proteins determine centromere drive and
suppression. *Cell* 2021, **184:**4904-4918.e4911.
134. Kipling D, Warburton PE: **Centromeres, CENP-B and Tigger too.** *Trends Genet* 1997,
13:141-145.
135. Khost DE, Eickbush DG, Larracuente AM: **Single-molecule sequencing resolves the**
detailed structure of complex satellite DNA loci in *Drosophila melanogaster*. *Genome*
Res 2017, **27:**709-721.
136. Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic**
features. *Bioinformatics* 2010, **26:**841-842.
137. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper
A, Markowitz S, Duran C, et al: **Geneious Basic: an integrated and extendable**
desktop software platform for the organization and analysis of sequence data.
Bioinformatics 2012, **28:**1647-1649.
138. Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2.** *Nat Methods*
2012, **9:**357-359.
139. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R, Genome Project Data Processing S: **The Sequence Alignment/Map format**
and SAMtools. *Bioinformatics* 2009, **25:**2078-2079.
140. Buske FA, Bodén M, Bauer DC, Bailey TL: **Assigning roles to DNA regulatory motifs**
using comparative genomics. *Bioinformatics* 2010, **26:**860-866.
141. Langmead B: **Aligning short sequencing reads with Bowtie.** *Curr Protoc*
Bioinformatics 2010, **Chapter 11:**Unit 11.17.
142. Han BW, Wang W, Zamore PD, Weng Z: **piPipes: a set of pipelines for piRNA and**
transposon analysis via small RNA-seq, RNA-seq, degradome- and CAGE-seq,
ChIP-seq and genomic DNA sequencing. *Bioinformatics* 2015, **31:**593-595.
143. Long AD, Macdonald SJ, King EG: **Dissecting complex traits using the Drosophila**
Synthetic Population Resource. *Trends Genet* 2014, **30:**488-495.

144.   Grenier JK, Roman Arguello J, Moreira MC, Gottipati S, Mohammed J, Hackett SR,
Boughton R, Greenberg AJ, Clark AG: **Global Diversity Lines–A Five-Continent**
Reference Panel of Sequenced *Drosophila melanogaster* Strains. *G3:*
Genes|Genomes|Genetics 2015, **5:**593-603.
145.   Pedersen BS, Quinlan AR: **Mosdepth: quick coverage calculation for genomes and**
exomes. *Bioinformatics* 2018, **34:**867-868.
146.   Navarro-Dominguez B, Chang C-H, Brand CL, Muirhead CA, Presgraves DC,
Larracuente AM: **Epistatic selection on a selfish supergene - drive, recombination,**
and genetic load. *Elife* 2022, **11**.
147.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,
Lunter G, Marth GT, Sherry ST, et al: **The variant call format and VCFtools.**
Bioinformatics 2011, **27:**2156-2158.
148.   Knaus BJ, Grünwald NJ: **vcfr: a package to manipulate and visualize variant call**
format data in R. *Mol Ecol Resour* 2017, **17:**44-53.
149.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM: **Canu: scalable**
and accurate long-read assembly via adaptive -mer weighting and repeat
separation. *Genome Res* 2017, **27:**722-736.
150.   Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics* 2018,
34:3094-3100.
151.   Chakraborty M, Baldwin-Brown JG, Long AD, Emerson JJ: **Contiguous and accurate**
de novo assembly of metazoan genomes with modest long read coverage. *Nucleic*
Acids Res 2016, **44:**e147.
152.   Vaser R, Sović I, Nagarajan N, Šikić M: **Fast and accurate de novo genome assembly**
from long uncorrected reads. *Genome Res* 2017, **27:**737-746.
153.   King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, Long AD,
Macdonald SJ: **Genetic dissection of a model complex trait using the Drosophila**
Synthetic Population Resource. *Genome Res* 2012, **22:**1558-1566.
154.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL:
Versatile and open software for comparing large genomes. *Genome Biol* 2004, **5:**R12.
155.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search**
tool. *J Mol Biol* 1990, **215:**403-410.
156.   Bailey TL, Johnson J, Grant CE, Noble WS: **The MEME Suite.** *Nucleic Acids Research*
2015, **43:**W39-W49.
157.   Katoh K, Standley DM: **MAFFT multiple sequence alignment software version 7:**
improvements in performance and usability. *Mol Biol Evol* 2013, **30:**772-780.
158.   Stamatakis A: **RAxML version 8: a tool for phylogenetic analysis and post-analysis**
of large phylogenies. *Bioinformatics* 2014, **30:**1312-1313.
159.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K: **MEGA X: Molecular Evolutionary**
Genetics Analysis across Computing Platforms. *Mol Biol Evol* 2018, **35:**1547-1549.
160.   Ohta T: **Allelic and nonallelic homology of a supergene family.** *Proc Natl Acad Sci U S*
A 1982, **79:**3251-3254.
161.   Rozen S, Skaletsky H, Marszalek JD, Minx PJ, Cordum HS, Waterston RH, Wilson RK,
Page DC: **Abundant gene conversion between arms of palindromes in human and**
ape Y chromosomes. *Nature* 2003, **423:**873-876.
162.   **Biostrings: Efficient manipulation of biological strings.**
[https://bioconductor.org/packages/Biostrings]

163.   Duchen P, Zivkovic D, Hutter S, Stephan W, Laurent S: **Demographic inference reveals**
1183   **African and European admixture in the North American *Drosophila melanogaster***
1184   **population.** *Genetics* 2013, **193:**291-301.
1185   164.   Li H, Stephan W: **Inferring the demographic history and rate of adaptive**
1186   **substitution in Drosophila.** *PLoS Genet* 2006, **2:**e166.
1187