Centromere-associated retroelement evolution in *Drosophila melanogaster* reveals an

underlying conflict

Lucas W. Hemmer[1,2*], Sherif Negm[1,3], Xuewen Geng[1], Cécile Courret[1], Beatriz Navarro-

Domínguez[1,4], Iain Speece[1], Xiaolu Wei[1], Eddyson Altidor[1], James Chaffer[1], John S. Sproul[1,5],

Amanda M. Larracuente[1*]

Affiliations:

[1] Department of Biology, University of Rochester, Rochester, NY, USA

[2] Invaio Sciences, Cambridge, MA, USA

[3] Department of Human Genetics, University of Chicago, Chicago, IL, USA

[4] Departamento de Genética, Facultad de Ciencias, Universidad de Granada, Granada, 18071,

Spain

[5] Department of Biology, University of Nebraska Omaha, Omaha, NE, USA

ABSTRACT

Centromeres are chromosomal regions essential for coordinating chromosome segregation during cell division. While centromeres are defined by the presence of a centromere-specific histone H3 variant rather than a particular DNA sequence, they are typically embedded in repeat-dense chromosomal genome regions. In many species, centromeres are associated with transposable elements, but it is unclear if these elements are selfish or if they play a role in centromere specification or function. Here we use *Drosophila melanogaster* as a model to understand the evolution of centromere-associated transposable elements. *G2/Jockey-3* is a non-LTR retroelement in the *Jockey* clade and the only sequence shared by all centromeres. We study the evolution of *G2/Jockey-3* using short and long read population genomic data to infer insertion polymorphisms across the genome. We combine estimates of the age, frequency, and location of insertions to infer the evolutionary processes shaping *G2/Jockey-3* and its association with the centromeres. We find that *G2/Jockey-3* is an active retroelement targeted by the piRNA pathway that is enriched in centromeres at least in part due to an insertion bias. We do not detect signatures of positive selection on any *G2/Jockey-3* insertions that would suggest than individual copies are favored by natural selection. Instead, we infer that most insertions are neutral or weakly deleterious both inside and outside of the centromeres. Therefore, *G2/Jockey-3* evolution is consistent with it being a selfish genetic element that targets centromeres. We propose that targeting centromeres helps active retroelements escape host defenses, as the unique centromeric chromatin may prevent targeting by the host silencing machinery. At the same time, centromeric TEs insertions may be tolerated or even beneficial if they also contribute to the transcriptional and chromatin environment. Thus, we suspect centromere-associated retroelements like *G2/Jockey-3* reflect a balance between conflict and cooperation at the centromeres.

INTRODUCTION

Centromeres are important chromosomal structures for kinetochore assembly and proper chromosome segregation during cell division. While centromeres are usually in highly repetitive regions, the role of DNA sequences in centromere function or establishment remains unclear (reviewed in [1]). Across many organisms, centromeres are associated with satellite DNAs (satDNAs), large blocks of tandemly repeated DNA sequences (reviewed in [2, 3]). Centromere-associated tandem repeats are capable of initiating centromere assembly in fission yeast [4] and humans [5]. However, centromeres are not defined by any particular DNA sequences. They are instead defined epigenetically by CENP-A, a specific variant of histone H3 [6-11]. Centromere domains consist of unique chromatin where H3 nucleosomes are interspersed with CENP-A nucleosomes [12-15]. Repeat sequences are not always required for centromere formation, as neocentromeres can form outside of centromeric DNA in the absence of satDNA sequences [16-18] and some species have centromeres devoid of repeats (*e.g.* satellite-free centromeres in the Equus genus [19]). Neocentromeres have the potential to replace older centromeres with implications for karyotype evolution and speciation [20-22].

Centromeres and pericentromeric heterochromatin are also enriched in transposable elements (TEs)— selfish genetic elements that mobilize in genomes. Evidence for specific TE-centromere associations has emerged in the last two decades, including in grasses [23, 24], rice [25, 26], maize [27], marsupials [28, 29], bats [30], gibbons [31, 32], and humans [18, 33, 34]. However, it is unclear how, or if, TEs contribute to centromere biology for most species [35]. Centromeres may offer a 'safe haven' to TEs that jump there and for the genome [36, 37], as centromere

70 insertions are unlikely to disrupt centromere or gene function [35]. At the same time, centromeric

71 TEs may contribute to important aspects of centromere evolution and biology. Homology

72 between some TEs and centromeric tandem repeats [3, 38] suggests that TEs can contribute to

73 centromere evolution by giving rise to new centromeric satDNAs [39]. TEs may also play roles

74 in centromere function. These roles may be indirect, such as maintaining the constitutive

75 pericentromeric heterochromatin, which is important for centromere function [40-42]. However,

76 more direct roles are also possible. Some transcription of centromeric regions is necessary for

77 recruitment of CENP-A and establishing centromere location [43-47]. Centromeric transcripts

78 may also facilitate recruitment of kinetochores to centromeric DNA [48, 49]. Given that TEs

79 contain sequences important for promoting transcription [50], their presence in centromeres may

80 contribute to centromere specification and function [35]. In most cases it is unclear whether TEs

81 are a part of the CENP-A domain corresponding to functional centromeres or just associated with

82 adjacent, repeat-rich regions of chromosomes where functional centromeres are often embedded.

83 Advances in single molecule long-read sequencing technology now allows for detailed study of

84 centromere organization at the sequence level to provide insights into centromere structure,

85 composition, and evolution [51-56].

87 With its compact genome and expansive genetic toolkit, *Drosophila melanogaster* is an excellent

88 model organism to study centromere evolution [57]. A recent study of centromere organization

89 in *D. melanogaster* revealed that CENP-A is bound to TE-rich islands of complex DNA

90 embedded within large arrays of satellite repeats [53]. The composition of each centromere

91 island is unique. The only sequence element shared among all centromeres is a single non-LTR

92 retroelement in the *Jockey* superfamily, *G2/Jockey-3*. However, *G2/Jockey-3* is not exclusively

located in centromeres; about 40% of all insertions are found outside of the centromere islands

[53]. Polymorphic *G2/Jockey-3* insertions [58] and transcripts from published RNAseq datasets

[59-61] suggest this element may still be active in *D. melanogaster* [53]. *G2/Jockey-3* is also

found at the centromeres in *D. simulans*, a close relative of *D. melanogaster,* suggesting that its

centromere association may be broadly conserved [53].

It is unclear whether *G2/Jockey-3* is an important centromere component in *D. melanogaster* or

if they are instead purely selfish TEs. If *G2/Jockey-3* are important for centromere specification

or function, we may detect evidence for positive natural selection on centromeric copies.

Alternatively, *G2/Jockey-3* may be a selfish element that targets insertions in centromeres but are

otherwise selectively neutral or weakly deleterious. These hypotheses are not necessarily

mutually exclusive. Here we investigated the age and frequency of *G2/Jockey-3* insertions across

D. melanogaster genomes to determine if its evolution is consistent with a role in centromere

function or underlying selfish behavior. Our analyses suggest that *G2/Jockey-3* insertions are

largely neutral or weakly deleterious both inside and outside centromeres, suggesting that they

are selfish genetic elements. Our results also suggest that *G2/Jockey-3* has had a weak insertion

bias toward centromeres. Although there is no evidence for positive selection on individual

G2/Jockey-3 centromeric insertions, insertion of active TEs like *G2/Jockey-3* may still contribute

to transcription and the unique chromatin environment at centromeres. We propose that there

may be a balance between conflict and cooperation between active TEs like *G2/Jockey-3* and

centromeres.

RESULTS

1. A new consensus sequence for G2/Jockey-3

Prior to analyzing the distribution of *G2/Jockey-3* insertion polymorphisms in *D. melanogaster*

populations, we first confirmed the consensus annotation. We analyzed all *G2/Jockey-3*

insertions in our *D. melanogaster* genome assembly [53] and found two lines of evidence that the

annotation needed updating: 1) 25 of the 167 *G2/Jockey-3* sequences in the genome shared up to

92 bp of upstream sequence at the 5'-end of the annotation; and 2) our reanalysis of PROseq data

[62] suggests that *G2/Jockey-3* transcription starts 92 bp upstream of the published start site

(Figure S1A). This upstream region is also enriched for GO terms for RNA polymerase II

transcription factors and other transcription factor activity (Figure S1B). We therefore extended

the original *G2/Jockey-3* consensus sequence by 92 bases (File S1).

G2/Jockey-3 is a part of the *Jockey* superfamily of non-LTR retroelements, which typically

contain two open reading frames (ORFs) (Figure S1A). While ORF1 is generally less conserved

[63], ORF2 is well-conserved across *Jockey* elements and encodes for a reverse transcriptase

(RT) and an endonuclease. NCBI ORF finder detected both ORFs within *G2/Jockey-3*. ORF1 is

predicted to produce a 427 amino-acid protein approximately 186 bp downstream of the 5' end.

The NCBI Conserved Domains identified a PRE_C2GC super family (cl11655, E-value 9.22E-

18) within ORF1 and while the function is unknown, it is associated with Cys2His2 type zinc

fingers found in putative GAG proteins in other arthropod species. The predicted ORF2 overlaps

with the last four bp of ORF1 and contains a domain corresponding to endonuclease (EEP

Superfamily Accession cl00490, E-value 7.21E-18) and RT (RT_like Superfamily Accession

cd01650, E-value 4.62E-56). *G2/Jockey-3* ends in a poly(A) signal.

2. Overview of data sets analyzed

We studied the evolutionary dynamics of *G2/Jockey-3* and other TEs by investigating the

patterns of insertion polymorphisms both in a single panmictic population, and in individuals

from a wide geographic range (Figure 1, Table S1). We conducted a survey of TE dynamics and

genetic variation in 135 samples from the *Drosophila melanogaster* Genetic Reference Panel

(DGRP) collected from a single population in North Carolina with whole genome Illumina

sequence data [64, 65] and a subset of long read data from the same population for validation

[66] (see Supplemental Text). For the samples from a wide geographic range, we used both long-

and short-read data from 13 founder lines of the *Drosophila* Synthetic Population Resource

(DSPR) and Oregon-R [67]. We also analyzed long-read PacBio data from five Global Diversity

Lines (GDL) samples from different continents [68]. We generated *de novo* genome assemblies

of each line to determine the organization of the centromere islands originally described in the

reference genome for comparison (Table S2) and predicted insertions in short read data with the

McClintock pipeline [69]. We assessed the reliability of our TE insertion predictions by looking

for concordance between short and long read predictions (see Supplemental Text) and through

PCR validation of a subset of insertions. We confirmed 64% (67 / 105) of the *G2/Jockey-3*

insertions detected by McClintock are found in the long-read samples we examined (File S2).

PCR confirmed 80% (28 / 35) of *G2/Jockey-3* insertions predicted in the GDL lines, and ~67%

(12 / 18, File S3) of the centromere insertions. Based on our assessment, we limit our analysis to

short read datasets with >20X centromere coverage (Table S3) and TEs predicted by more than

one TE caller implemented in McClintock (Supplemental Text).

Figure 1: The summary of data and analyses using one source or a combination of sources

in this paper. A heterochromatin-enriched assembly was originally used to discover centromere

islands in Chang et al. [53] and identify reference TE insertions. We used short reads from the

DGRP to identify reference and *de novo* insertions to study TE dynamics within a single

population from North Carolina (map source: R package usmaps https://cran.r-

project.org/web/packages/usmap/). We generated genome assemblies for 18 samples from global

populations to uncover the centromere island sequences and *de novo* elements (map source:

www.outline-world-map.com). We combined the reference sequences with population

frequencies to infer selection with an age-of-allele test [70]. We also combined reference TE

sequences with *de novo* elements uncovered in our assemblies for phylogenetic analysis. The fly

illustrations are modified images from Wikipedia Commons

(https://commons.wikimedia.org/wiki/File:Drosophila-drawing.svg).

3. G2/Jockey-3 is active in Drosophila melanogaster populations

Previous studies of TE polymorphism in *D. melanogaster* are consistent with *G2/Jockey-3*

(previously annotated as *G2*) being recently active [58, 71, 72]. To determine if *G2/Jockey-3* is

active in the DGRP, we detected TE insertion polymorphisms with the McClintock pipeline [69]

and classified each insertion based on their presence (a "reference" insertion) or absence (a non-

reference or "*de novo*" insertion, Figure 2) in our reference genome [53, 73]. *G2/Jockey-3*

showed approximately five *de novo* insertions per line (File S4-S5), which is considerably fewer

than the most highly active *Drosophila* TEs. For example, we estimate *Hobo* and *Jockey-1* have

approximately 35 *de novo* insertions per line (consistent with [58]). However, the fraction of *de*

novo insertions as percentage of all insertions (39% or 733 / 1124 for *G2/Jockey3*) is comparable

to other highly active elements [72] such as *Doc* (25%; 2944 / 3826), *Jockey* (24% ;1308 /

5448), *I* element (27%; 658 / 2364), and *Copia* (88% ;1083 / 1226) (File S4).

We detected insertions in the centromeres, here defined as the centromere islands identified in

Chang et al. [53]. Of the 18 different TE families detected within centromeres, *G2/Jockey-3* is

the most common. Within centromeres, *G2/Jockey-3* has the highest percentage of *de novo*

(32%; 15/47) and reference (72%; 28 / 39, Figure 2) TE insertions. Only *G2/Jockey-3* and *Doc*

have more than 10 unique insertions (File S4). Other *de novo* centromere insertions include

active families such as the *F* element, *R1* and *Roo*.

Many active TEs are targeted by small, 24-30-nt long Piwi-interacting RNAs (piRNA) in the

germline [74, 75]. These piRNAs target TEs for both transcriptional and post-transcriptional

silencing. In the *D. melanogaster* germline, piRNAs are produced from precursor transcripts that

originate from discrete loci in the genome that contain clusters of nested TEs [76] called piRNA

clusters [77]. Subsequent amplification of the piRNA signal comes from the piRNA-guided

cleavage of active TEs through the 'ping-pong' cycle [77, 78] generating a signature of 10-nt

overlap between sense and antisense piRNAs. Consistent with *G2/Jockey-3* being a recently

active element, we detect insertions of *G2/Jockey-3* in the *38C1* (2 copies) and *38C2* (9 copies)

piRNA clusters, and *G2/Jockey-3*-derived piRNAs in the germline. *G2/Jockey-3* is ranked in the

top 4% and top 18% of TE-derived piRNAs in testes and ovaries, respectively (Table S4). The

resulting *G2/Jockey-3* piRNAs show signatures of ping-pong in ovaries (Z-score = 7.11; data

from [60, 79, 80]) but not in testes (Z-score = 0.15; data from [81-83]), consistent with other

testis piRNAs. Taken together, these results suggest that *G2/Jockey-3* is an active TE enriched in

centromere islands.

Figure 2: The distribution and frequency of *G2/Jockey-3* insertions in *D. melanogaster*. The

height of the bars corresponds to frequency of an insertion detected in the DGRP samples and the

color denotes whether an insertion is found in the reference genome or is *de novo*. Each A-D

subpanels depict a chromosome (Ch 2, Ch 3, Ch 4, Ch X) from the heterochromatin-enriched

assembly of *D. melanogaster* genome [53, 73], with a window zoomed in on the corresponding

centromere (*e.g.,* X chromosome centromere = CenX, *etc*).

4. Young G2/Jockey3 insertions are enriched in centromeres

To understand if the observed enrichment of *G2/Jockey3* at centromeres is due to insertion bias,

we studied insertion frequencies across the genome. We reasoned that an enrichment of

putatively young *G2/Jockey-3* at centromeres may reflect an insertion preference. Young,

recently inserted TEs are expected to occur at lower frequencies than older TEs. We thus

examined the distribution of rare insertions (<10% population frequency) of *G2/Jockey-3* and

other TE families in the DGRP. We found that rare *G2/Jockey-3* insertions are significantly

enriched in centromeres compared to other regions relative to their size (19 centromere

insertions; Permutation tests: n_{perm} = 10,000, *P* < 0.0001; *Z*-score = 26.736 for whole genome; *P*

< 0.0001; *Z*-score = 23.895, excluding heterochromatin windows*; P* < 0.0001; *Z*-score = 17.354,

excluding euchromatin windows; Figure 3). *R1* elements are the only other TE enriched in

centromeres (4 centromere insertions; Permutation tests: n_{perm} = 10,000, *P* < 0.0001; *Z*-score =

6.892 for whole genome; *P* < 0.0001; *Z*-score = 5.92, excluding heterochromatin windows*; P* =

0.0187; *Z*-score = 2.605, excluding euchromatin windows; Figure 3), but to a lesser degree than

G2/Jockey-3. Among rare insertions, *G2/Jockey-3* is by far the most enriched TE in centromeres

(39% of all rare insertions).

Figure 3: Young *G2/Jockey-3* insertions are highly enriched in centromeres relative to

heterochromatin and euchromatin. The distribution of rare insertions (< 10% of the

population) per Mb between the centromere islands, heterochromatin, and euchromatin regions

for several TE families in DGRP lines. Significant differences are noted as: * $P < 0.05$, *** $P <$

0.0001 in the permutation test.

We confirmed these patterns using an orthogonal approach with long-read genome assemblies of

flies from a wide geographic distribution (*i.e.,* founder DSPR lines, GDL lines, and our reference

genome; Table S1). Here we identified putatively young *G2/Jockey-3* insertions based on

sequence divergence instead of frequency. We classified young insertions as those with < 1%

divergence from the consensus sequence, or with < 2% divergence from the consensus and found

within a single genome assembly. Insertions classified as young were found in less than 50% of

the assemblies, had fewer unique substitutions, and were more likely to have a complete ORF

than older fragments (Figure 4, FET P = 3.93E-12, File S6, File S7). Consistent with our short-

read DGRP findings (Figure 3), we found significant relative enrichment in centromeres

(Permutation tests n_{perm} = 10,000: P < 0.0001; Z-score = 15.633 for whole genome; P < 0.0001;

Z-score = 7.505 excluding euchromatin windows), even with some of the centromere island

sequences missing in the long-read assemblies (Table S2).

These results suggest that *G2/Jockey-3* has had at least some insertion preference for

centromeres. As insertion bias can be associated with a target site sequence, we searched for

motif enrichment at insertions sites. Although we found an enrichment of AT-rich simple motifs

like "TAGTTTT" and rDNA-associated sequences (*e.g.* 18S and non-transcribed spacers; Figure

S2) within 100 bases of predicted insertion sites, these sequences are generally abundant in

repeat-dense regions of the genome and not specifically enriched at *G2/Jockey-3* insertions (File

S6, File S8; MEME *E*-value = 3.8E-16).

Figure 4: Young *G2/Jockey-3* insertions share a common origin. A maximum likelihood

phylogenetic tree with all non-redundant *G2/Jockey-3* sequences larger than 10% of the full-

length (400 bp) in the reference genome [53] and long-read assemblies of founder DSPR and

GDL lines, with the *D. simulans* consensus as the outgroup. Black circles denote "complete"

G2/Jockey-3 elements containing a full ORF2; gray circles are incomplete elements. The

columns and colors correspond to characteristics of each insertion including chromatin region

(centromere or not), relative age, and presence on Y chromosome. Young insertions, determined

using an orthogonal approach to the Figure 3 analysis, are those with < 1% divergence from the

consensus sequence or < 2% and found within a single genome assembly.

5. Skew toward high frequency copies of G2/Jockey-3 in the centromere

To test if *G2/Jockey3* insertions are retained at centromeres by natural selection, we studied their

frequencies. Because most TE insertions are deleterious in the euchromatin, they are found at

low population frequencies [84]. In contrast, TEs in heterochromatin are allowed to reach higher

population frequencies because recombination is infrequent, and natural selection is less

efficacious (reviewed in [85-87]). To infer differences in the strength and direction of natural

selection in different genomic regions, we examined the frequency spectra of a subset of TE

families in centromeres, heterochromatin, and euchromatin (Figure 5, Figure S3). If natural

selection favors centromeric insertions, we should observe a skew toward high frequency

variants in centromeres compared to heterochromatin. We compared the frequency spectra of

two TE families which had five or more insertions in the centromeres: *G2/Jockey-3* and *Doc*. As

expected, both elements showed an enrichment for rare insertions in the euchromatin, consistent

with TEs being generally deleterious in gene-rich regions (Figure 5; FET *P* < 0.001). Similarly,

we observed a slight skew toward higher frequency *G2/Jockey-3* and *Doc* insertions (*e.g.*

common/intermediate) in heterochromatin and centromeres compared to euchromatin (Figure 5,

FET *P* < 0.001), suggesting relaxed constraints or a reduced efficacy of natural selection outside

of euchromatic regions. Other TE families found in centromeres are also enriched for low

frequency insertions in the euchromatin (Figure S3). However, we did not observe differences

between the frequency spectra of the euchromatin and centromeres, suggesting most TEs may be

deleterious in these regions (Figure S3).

The frequency spectra of *G2/Jockey3* does not appear significantly different between

centromeres and heterochromatin. However, *G2/Jockey-3* has more "common" insertions in

centromeres relative to other TEs, including *Doc* (Figure 5; Figure S3). This could be due to

relaxed constraints allowing *G2/Jockey-3* insertions to rise to higher frequencies over time or

positive selection for some *G2/Jockey-3* insertions in centromeres.

Figure 5: *G2/Jockey-3* has higher frequency variants in centromeres. TE frequencies were

binned into rare (less than 10% of all samples), intermediate (10-50% of all samples), and

common (> 50% of all samples). The frequency distributions of the four TE families with > 10

unique insertions in the centromeres were compared among the centromeres, heterochromatin,

and euchromatin. Fisher's Exact Tests between regions, Bonferroni adjusted two-tailed P-value *

P < 0.05; *** *P* < 0.001; ns is not significant.

6. G2/Jockey-3 ages suggest a historical burst of activity

Centromeric *G2/Jockey-3* copies are generally older—within a range of 2-7% divergence from

the consensus—than those in heterochromatin (MWU, $P < 0.001$, Figure 6A), but not

significantly different than euchromatic copies (MWU, $P > 0.5$; Figure 6A). This pattern

suggests that most centromeric copies of *G2/Jockey-3* resulted from a historical burst of activity.

This burst of activity is consistent with general patterns observed at other non-LTR retroelements

[72, 88]. Other TE families show a similar age distribution between heterochromatin and the

centromeres (Figure S4). These copies are also older and highly fragmented. In general, both the

lack of recombination and lack of strong purifying selection against TE insertions in centromeres

and heterochromatin may allow these insertions to persist over longer evolutionary time periods

[89].

We can combine information about the TE's age and frequency to infer if, and how, its history is

consistent with natural selection. In the absence of selection, the frequency of a TE insertion

should correlate with its age: *e.g.,* older TEs have more time to drift to higher population

frequencies than young TEs and thus accumulate more unique mutations. Natural selection can

cause deviations from expected frequencies based on the age. While negative selection might

cause TEs to have lower population frequency than expected given their age, positive selection

might increase their frequencies. We therefore used an age-of-allele test [70] to detect whether

G2/Jockey-3 frequencies deviate from our expectations. We estimated age as the number of

unique substitutions for each TE copy based on the reference genome and we estimated

frequency from the 28 high-coverage DGRP samples, controlling for demographic history [70].

For this test, we need to compare insertions spanning a range of ages to identify individual

insertion frequencies that deviate from expectations [70]. We thus examine *G2/Jockey-3* and

other TE families that have centromere insertions. We find that *G2/Jockey-3* (Figure 6B, File

S9)*,* like the majority of TEs, segregates either at the expected frequencies or lower than

expected, given their age (Figure S6, File S9). This suggests that many *G2/Jockey-3* copies, both

within and outside the centromeres, are either neutral or weakly deleterious. Therefore

G2/Jockey-3 shares similar selective pressures as other presumably weakly deleterious TE

families in the genome.

Interestingly, we observed an enrichment of copies with zero unique substitutions for the

heterochromatin and CenY insertions (Figure S5). However, we suspect this pattern arises from

duplications rather than independent *G2/Jockey-3* insertion events (see below). Genomic

duplication events are common in heterochromatin in the Y chromosome [73, 76, 90].

While we do not see evidence for strong recent positive selection on any individual *G2/Jockey-3*

insertions in the centromeres, there may be selection on centromere haplotypes that differ in their

composition or structure.

Figure 6: *G2/Jockey-3* copies are either as frequent or less frequent than expected given their age. A) The number of *G2/Jockey-3* insertions grouped by divergence from the consensus sequence split between the euchromatin, heterochromatin, and centromeres. Two-tailed Mann Whitney U tests *** $P < 0.001$, ns is not significant. B) The difference between observed and expected frequency of *G2/Jockey-3* copies in the DGRP. The *P*-values are calculated as the sum of the probabilities that an insertion is less frequent than expected (if observed - expected < 0) or more frequent than expected (observed - expected > 0). * $P < 0.1$, a stringent threshold given the reduced power to detect deviations from neutrality as stated in Blumenstiel et al. [70].

7. Centromere island haplotype structure not likely driven by G2/Jockey-3 insertions

To detect signatures of natural selection on centromere haplotypes and any associations with

G2/Jockey-3 copies, we examined patterns of nucleotide variation. Centromeres and

pericentromeric regions in *D. melanogaster* generally do not recombine by crossing over [91, 92]

therefore the effects of linked selection in non-recombining regions should lead to low levels of

nucleotide diversity. We thus expect to see reduced genetic variation in the centromeres and high

levels of linkage disequilibrium. We quantified genetic variation in the centromere islands for

DGRP samples with high centromere coverage (> 20X, see Methods). As expected, nucleotide

diversity is greatly reduced in every centromere compared to the major chromosome arms

(MWU $P < 2.0E-07$; Figure S7, Table S5) except for chromosome 4, which does not recombine,

and its centromere and chromosome arms are similar (MWU $p = 0.56$; Figure S8, Table S5). We

find that Tajima's *D*, a summary of the site frequency spectrum, is also typically lower in

centromere islands (Figure S8), indicating a skew toward rare alleles in centromeres. The

centromere island haplotype structures (Figures S9-S11) indicate that the majority of centromeric

SNPs are randomly distributed with respect to the *G2/Jockey-3* sequences (FET $P > 0.46$).

Linked selection anywhere in or near centromeres can reduce genetic diversity, skew frequency

spectra, and create haplotype structure. While centromere island haplotype structure is consistent

with a history of linked selection, we do not detect evidence that *G2/Jockey-3* insertions

themselves are necessarily the targets of selection.

8. G2/Jockey-3 copy number amplification on the Y chromosome is driven by duplication

The Y chromosome centromere is unique in that it has the largest centromere island containing

nearly half of all *G2/Jockey-3* fragments, but no identifiable satellite sequences [53]. We cannot

use the genomic re-sequencing data to study polymorphic insertions on the Y chromosome, as

the data are either female-only or mixed sex and thus lower coverage. However, our high-quality

reference assembly provides an opportunity to infer the organization and evolutionary history of

Y-linked *G2/Jockey-3*. We inferred a phylogenetic tree of all Y-linked copies of *G2/Jockey-3*.

Based on this phylogeny, we classified centromeric Y-linked *G2/Jockey-3* copies into four broad

classes (Figure S12). Class 1 *G2/Jockey-3* correspond to young and mostly long insertions, class

2 to short (< 160 bp) 3' end fragments, class 3 to old internal fragments missing the 5' and 3'

terminal ends, and class 4 to generally long and old 3' end fragments divided into two subclasses

(4 and 4A) based on position (Figure 9, Figure S12). We mapped these classifications onto our Y

chromosome assembly and estimated pairwise divergence between all *G2/Jockey-3* copies in the

Y centromere (Figure 7) and elsewhere on the Y chromosome (Figure S13). Each class seems to

undergo independent duplication events, however at a larger scale we do not identify higher

order repeat domains (Figure 7). This structure differs from other known centromeres (*e.g.*

primates), which are organized into higher order repeats, the youngest repeats are found in the

center and the older, more divergent repeats, are at the outer edges of the array [51, 52]. By

contrast, in the *D. melanogaster* Y centromere we observe that regions enriched with young

G2/Jockey-3 copies punctuate regions rich with older copies (Figure 9), suggesting that recent,

local rearrangements affect centromere organization.

Figure 7: *G2/Jockey-3* **evolution in the Y chromosome centromere is shaped by expansion**

via duplication, gene conversion and rearrangements. The colored pairwise divergences

between individual *G2/Jockey-3* copies on the Y centromere are shown on top with their relative

positions on the line graph below. White spaces represent *G2/Jockey-3* copies which do not

overlap, so pairwise divergence cannot be calculated. Below the graph is a visual representation

of the broad categories of repeats on the Y chromosome centromere, labeled by color with length

and direction illustrated with arrowed lines. The repeats appear arranged into roughly four

regions with overlapping borders based on the types of repeats and pairwise divergence. The

classes of *G2/Jockey-3* are as follows: Class 1 younger insertions, Class 2 short (< 160 bp) 3' end

fragments, Class 3 older internal fragments missing the 5' and 3' terminal ends, Class 4

generally longer and older 3' end fragments. Class 4 contains a subclade (4A) with highly similar

sequences almost exclusively from the single region of CenY.

We estimated gene conversion rates of *G2/Jockey-3* on the Y centromere with respect to their

classes (Supplemental Text). Our estimates for gene conversion are on the order of 1E-6 per site

per generation (Table S6), ~10E3 times higher than the other chromosomes in *Drosophila* [91-

93] and an order of magnitude less than previously estimated gene conversion rates between

duplicated gene families on the Y chromosome [73]. Moreover, we do not see complete

sequence homogenization within classes. We observe similar patterns of duplication and

expansion outside the centromere on the Y chromosome (Figure S13), as 32 of 47 non-

centromere Y-linked copies appear to be duplicated; they are of similar fragment size,

orientation, and have few no zero unique substitutions (File S6). Therefore, we conclude that

G2/Jockey-3 evolution on the Y chromosome is driven by duplication of historical insertions

with some gene conversion among copies. Population genomic data from the Y chromosome will

help distinguish between types of recombination affecting Y centromere structure.

Outside of the Y chromosome, putative duplications involving *G2/Jockey-3* are less common

(Figure S13). Many copies of *G2/Jockey-3* within the centromeres are fragmented and therefore

share no homologous sequences for pairwise calculations (Figure S13). Pairwise divergence

between centromeric *G2/Jockey-3* fragments is elevated in comparison to fragments in both the

euchromatin and non-centromeric heterochromatin (Figure S14, MWU, $P < 0.01$), and lowest in

non-centromeric heterochromatin, especially on the Y chromosome. Therefore, the majority of

G2/Jockey-3 within the non-Y centromeres appears to be the result of recurrent past insertions.

CONCLUSIONS

G2/Jockey-3 is a unique TE in that it is the only sequence shared among centromeres of *D.*

melanogaster [53]. Here we show that *G2/Jockey-3* is a recently active non-LTR retroelement

that may have had an insertion preference for centromeres, given their relatively small size

compared to the rest of the genome (Figures 3). While the centromere enrichment and skew

toward high frequency centromere copies are expected for insertions favored by natural

selection, we do not find evidence for positive selection on individual *G2/Jockey-3* insertions.

Instead, most *G2/Jockey-3* insertions both inside and outside of the centromeres are at the

expected frequency or significantly rarer than expected given their age, suggesting they may be

neutral or slightly deleterious and accumulating due to relaxed constraints against centromeric

TE insertions (Figure 6B). The majority of *G2/Jockey-3* copies within the centromeres appear to

be the result of recurrent past insertions, except for on the Y chromosome. Some Y-linked copies

are likely a result of duplications rather than *de novo* insertions, similar to patterns reported in

other species (*e.g.* [28]).

We propose that most *G2/Jockey-3* insertions in the centromeres resulted from a burst of activity,

which is consistent with the evolutionary history of other non-LTR retroelements [88]. When a

TE first invades, it can proliferate rapidly until it comes under the surveillance of germline

defense mechanisms like the piRNA pathway, for example by inserting into a piRNA cluster

[94-96]. These dynamics are consistent with the invasion and control [94, 97] of the *P* element in

D. melanogaster [98] and its close relative, *D. simulans* [99]. Once targeted by the piRNA

pathway for transcriptional and posttranscriptional silencing, the TE may eventually become

dormant or go extinct. Here we show that *G2/Jockey-3* had a burst of activity and is now inside

piRNA clusters and actively targeted by the piRNA pathway (Table S4).

The *G2/Jockey-3* insertion polymorphism patterns and piRNA observations are most consistent

with some underlying selfish behavior typical among TE families, except that this element can

target centromeres. Although we are unsure of the precise targeting mechanism, we suspect the

target is the centromere chromatin rather than any particular DNA sequence. Such centromere

targeting was reported for LTR retroelements in other species such as maize [100, 101] and

Arabidopsis [102, 103]. For example, when the *Arabidopsis lyrata Tal-1 Copia*-type

retroelement [103] is introduced into *A. thaliana,* the element targets the centromeres despite

divergence in centromeric DNA sequences between these species [102].

TEs may accumulate in heterochromatic regions where they experience relaxed selection

compared to euchromatic insertions, which have potential to disrupt functional elements, risk

ectopic exchange and resulting genomic rearrangements [85, 86, 104], and spread silent

chromatin or DNA methylation into nearby functional elements [*e.g.*, 105-108]. However,

centromeres are not typical heterochromatin [109]. Here we propose that centromeres offer a

unique environment for TEs (Figure 8A). First, insertions are unlikely to be strongly disfavored,

as the function of centromeres is unlikely to be disrupted by a single insertion event. Second,

centromeric insertions may escape host silencing mechanisms. Small RNAs target TEs to limit

potentially harmful TE activity by inducing H3K9 methylation and thus transcriptional silencing

(reviewed in [110-112]). However, CENP-A structure and posttranslational modifications differ

from H3 [109], and the host silencing machinery is not known to target CENP-A. Therefore,

centromeres may offer a 'safe haven' for some TEs not just because they can avoid removal by

purifying selection, but also because TEs may temporarily escape targeting by small RNAs (*e.g.*

piRNAs, siRNAs) and alternative host defenses (*e.g.* DNA methylation in pericentromeres [113],

hypomethylated DNA in humans [52, 114]).

While most TEs are considered to be parasitic elements that are neutral or mildly deleterious

[115, 116], some TE insertions, including those in non-recombining and/or gene poor

heterochromatic environments, might contribute to host function [117]. For example, TEs can

adopt commensal and/or mutualistic strategies to propagate without direct confrontation by

cooperating with their host (reviewed in [118]). One extreme outcome is TE domestication,

where the elements no longer pose a threat to the host and instead now perform some important

genome function. One often cited example is at *Drosophila* telomeres, where three non-LTR

retroelements in the *Jockey* clade (*HeT-A*, *TAHRE*, and *TART* collectively referred to as HTT)

have specialized roles [119, 120]. The transposition activity of HTT is limited to telomeres,

where they directly contribute to telomere length and integrity [119-122]. However, detailed

study of the evolutionary genetic history of the retroelements across species suggests instead that

there is still an underlying conflict between the HTT and the genome, as there is rapid turnover

of HTT between species [123]. Alternatively, given the specialization of HTT, the dynamic and

unstable telomeric environment may necessitate rapid HTT evolution [124].

While centromeric TEs do not currently show evidence for domestication in *D. melanogaster*,

they may be tolerated, or even beneficial for three reasons. First, TEs may contribute to

centromere transcription (reviewed in [35]), as some transcription at centromeres is important for

CENP-A assembly and function [14, 46, 125-129] and centromere function [14, 45, 130].

Second, TEs might contribute to the unique chromatin at centromeres. H3 nucleosomes are

typically interspersed with CENP-A within centromere domains [12, 14] [15], including a

limited amount of H3K9me2/3 (*e.g.*, [13]). We propose that insertion of active TEs like

G2/Jockey-3 may contribute to this unique CENP-A/H3 chromatin at the centromeres (Figure

8A). While landing in CENP-A nucleosomes may allow the TE to temporarily escape silencing,

if an active TE lands in an H3 nucleosome region of the centromere, it may be targeted for

silencing through H3K9 methylation (*e.g.* piRNA- or siRNA-mediated silencing) and thus

contribute to H3K9me2/3 domains within the centromere. Third, TEs may help mitigate other

conflicts that happen at centromeres. Some centromeric sequences have the capacity to be selfish

and bias their transmission through female meiosis in a process known as meiotic drive [131].

Driving centromeres generally have expansions of satellite DNA sequences that can recruit more

kinetochore proteins and cause biased segregation toward the egg pole [132, 133] (Figure 8B).

Meiotic drive at the centromeres can lead to rapid evolution of both centromere DNA and

proteins that work to restore fair segregation [134, 135]. There are parallel pathways to suppress

meiotic drive involving rebalancing kinetochore proteins and spreading heterochromatin to limit

the CENP-A domain [136]. While centromere drive and suppression models predict that the

flanking heterochromatin domains are the main participants in the conflict [136], the

H3K9me2/3 domains within the centromeres might also contribute. Active centromeric TEs that

are targeted by small RNA pathways, like *G2/Jockey-3,* could lead to spreading of H3K9me2/3

heterochromatin from within the centromere (Figure 8C). In this way, the centromeric TEs and

'internal' centromeric H3K9 domains may also be a substrate for suppressors of centromere

drive to stop the unchecked spread of CENP-A caused by meiotic drivers.

Even if TEs cooperate with centromeres, this may be an uneasy alliance, as there could still be an

underlying conflict. For example, the invasion of H3K9me2/3 heterochromatin in centromeres is

potentially dangerous, as it may evict too much CENP-A and silence a centromere [5].

Therefore, the centromeric chromatin, and the relative abundance of CENP-A and H3

nucleosomes, may mediate the balance between conflict and cooperation between active TEs and

centromeres.

In many species across the tree of life, centromeres are associated with TEs [18, 23-34, 53]. The

connection between TEs and centromeres is supported not just by TE sequences embedded in

and around present-day centromeres, but also in the origins of some proteins that function in

centromere biology (*e.g.* similarity between centromere protein CENP-B and Pogo family

transposons [137]). This relationship may represent a delicate balance between conflict and

cooperation between the centromeres and the repetitive selfish genetic elements that occupy

these genome regions. We expect this balance to vary depending on genomic context, level of

TE activity, and presence and frequency of centromere meiotic drive. If TEs contribute

positively to centromere function, many types of elements may be sufficient to provide the right

'environment'. In this case, we would not necessarily expect to see a signature of selection on

any one insertion or type of insertion. While *G2/Jockey-3* is currently the principal element

occupying the centromere niche in *D. melanogaster*, it may be replaced over time by another

type of repeat. In our model, any repeat that can be transcribed and can recruit H3K9me2/3 (*e.g.*

is targeted by small RNAs) can contribute to centromere function. Detecting the mode, tempo,

and target of selection in non-recombining and dynamic environments like the centromeres is

difficult. Exploring the evolution of centromere composition in closely related species will

provide additional insights necessary to test these hypotheses.

Figure 8. Model for the accumulation of centromere-associated retroelements. Centromeres

have unique chromatin with interspersed CENP-A and H3 domains, including some H3K9me2/3

[12-15]. The diagrams represent nucleosomes, either CENP-A (blue) or H3 (green), with

H3K9me2/3 modifications indicated with a black dot. The nucleosome distributions are not

drawn to scale. A. Active TEs like *G2/Jockey-3* jump in centromeric chromatin (magenta

triangles and lines in nucleosomes) and could contribute to transcription (black wavy line) or be

silenced by host machinery (orange hexagon). Centromeres may offer a safe haven for TE

insertion because insertions there are unlikely to be strongly disfavored and because insertions in

CENP-A nucleosomes may escape host silencing machinery (*e.g.* piRNA pathway). Centromeric

TE insertions might be tolerated or even favored by the host if they contribute to centromere

transcription or a low level of H3K9me2/3 inside centromeres. B. Selfish centromeric DNA

sequences can expand their copy number and bias their transmission to the next generation

through female meiosis [131]. C. One way to suppress cheating centromeres is by spreading

heterochromatin to limit the CENP-A domain [136]. TE insertions in H3 nucleosomes within the

centromeres may be targeted for silencing by host silencing machinery (green circle with

magenta lines and black circle). These H3K9me2/3 nucleosomes within centromeres may help

create a more efficient response to driving centromeres by merging spreading pericentric

heterochromatin with internal H3K9me2/3 domains.

614   **MATERIALS AND METHODS**

G2/Jockey-3 re-annotation and expression analysis

Chang et. al. [53] generated the *G2/Jockey-3* consensus sequence after a phylogenetic analysis

confirmed previously annotated *G2* and *Jockey-3* were the same element [53]. We annotated a

heterochromatin-enriched long-read genome assembly of *D. melanogaster* [73] with

Repeatmasker v4.1.0 using the previous *G2/Jockey-3* consensus sequence along with a

Drosophila repeat library [138]. We extracted *G2/Jockey-3* and the surrounding sequences using

BEDTools v2.29.2 [139], aligned the sequences to the previous consensus sequence, manually

inspected the alignment, and generated the new consensus sequence with Geneious v8.1.6 [140]

(File S1). We identified Open Reading Frames (ORFs) with NCBI ORFfinder

(https://www.ncbi.nlm.nih.gov/orffinder/) and screened the ORF candidates by identifying

conserved domains common in non-LTR retrotransposons with the NCBI Conserved Domains

search tool (https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi). We defined the 5' end by

identifying transcription along the length of *G2/Jockey-3* by mapping publicly-available single-

end Precision nuclear Run-On Sequencing (PROseq) downloaded from NCBI SRA under

BioProject PRJNA451204 and accessions SRR7050623 - SRR7050630 [62]. We removed

adapters and reads < 16bp using cutadapt v2.1. Reads were reverse complemented using the

fastX toolkit v0.0.13 (fastx_reverse_complement). We mapped reads to the heterochromatin-

enriched genome assembly of *D. melanogaster* using bowtie2 v2.3.5.1 (--no-discordant --

dovetail --very-sensitive options; [141]). We extracted reads > 16 bp in length mapping to

G2/Jockey-3 annotations in the genome with samtools v1.7 [142] and remapped the reads to the

G2/Jockey-3 consensus with bowtie2 v2.3.5.1 [141]. We separated reads based on their strand

using samtools (v1.7). We calculated read depth at each position along the *G2/Jockey-3*

consensus and normalized coverage by the total number of genome mapped reads (RPM) using

bedtools genomecov [139]. We used alignment depth of all *G2/Jockey-3* genomic fragments to

identify where transcription occurs along the length of *G2/Jockey-3*. We submitted the sequence

in 5' end motifs under transcription peaks for a GoMo analysis [143] to identify motifs as

possible promoters.

To detect piRNAs corresponding to *G2/Jockey-3,* we trimmed the small RNA-seq reads for

adaptors and excluded the reads that align to rRNA or tRNA. We aligned the remaining reads to

the heterochromatin-enriched genome assembly [53, 73] using Bowtie [144], and then used a

customized python script as described in Wei et al. [83] to count reads that mapped to each TE

feature (htseq_bam_count_proportional.py;

https://github.com/LarracuenteLab/Dmelanogaster_satDNA_regulation; last accessed 11/2022,

https://doi.org/10.5061/dryad.hdr7sqvj3). We reported the RPM (Reads Per Million) values in a

summary table. We calculated the 10nt overlap Z-score of piRNAs using piPipes [145].

Short-read analysis

We detected TE insertions from publicly available FASTQ files from 135 *Drosophila*

melanogaster Genetic Reference Panel (DGRP) lines from BioProject PRJNA36679 [64, 65]

downloaded from the European Nucleotide Archive last accessed September 20, 2019, 13 of the

founder lines of the Drosophila Synthetic Population Resource (DSPR) BioProject

PRJNA156883[146], and 85 samples from the Global Diversity Lines BioProject PRJNA268111

(Table S3)[147]. We detected TE insertions in each sample with the McClintock pipeline [69]

(last downloaded and updated, August 20, 2020) combining several split-read and discordant

read mapping programs including the following: ngs_te_mapper, popoolationTE,

popoolationTE2, Retroseq, TE-locate, and TEMP (scripts and files for running McClintock at

https://github.com/LarracuenteLab/Dmel_Jockey-3_Evolution). We concatenated the outputs to

calculate the number of TEs detected by each program and sample and analyzed the outputs with

Python and R scripts available at https://github.com/LarracuenteLab/Dmel_Jockey-3_Evolution.

We repeated our analyses of TEs with and without insertions in known piRNA clusters and

removed fragments of Gypsy6_I_Dmoj and Gypsy1_I_Dmoj which resemble satellite sequences

in *D. melanogaster*. Depth of each short-read sample was calculated from the sorted BAM

outputs from the McClintock pipeline using Mosdepth [148] (Table S3). We retained samples

which had a centromere coverage greater than 20X for our analyses (see Supplemental Text for

more information).

We used the same short-read FASTQ files from the DGRP to quantify genetic diversity between

the centromere islands and the rest of the genome using a best practices GATK pipeline v4.1.9.0

to detect variant SNPs. We mapped reads from high coverage samples to our reference genome

with BWA v1.7.15 and samtools 1.7 to generate sorted BAM files and used GATK

MarkDuplicates, BaseRecalibrator, ApplyBQSR, AnalyzeCovariates, and HaplotypeCaller to

call and filter SNPs for each sample into a VCF file. The VCFs for all samples were

concatenated with GenotypeGVCFs and relicabrated with VariantRecalibrator and ApplyVQSR.

We quality filtered SNPs in the resulting GVCF file with the following criteria; QD < 2.0, FS >

60.0, MQ < 40.0, MQRankSum < -12.5, and ReadPosRankSum < -8.0. We calculated π and

Tajima's D with the SimplifyVCF_Basic.v2.pl and Windows_Basic.v2.pl Perl scripts on the

filtered SNPs (available at https://github.com/bnavarrodominguez/SD-Population-genomics,

[149]). We further analyzed variants within the centromeres with VCFtools [150] and the vcfR R

package [151]. Additional details of our analysis with short-read data and the results can be

found in the Supplemental Text section.

Long-read analysis

We assembled genomes for several *D. melanogaster* lines following the heterochromatin-

enriched assembly protocol from Chang & Larracuente [73]. We downloaded raw PacBio reads

for five of the Global Diversity Lines (GDL) from the NBCI BioProject PRJNA451471 [68],

eight Drosophila Synthetic Population Resources (DSPR) founder lines and Oregon-R with raw

Pacbio reads from BioProject PRJNA418342 [67], and five DGRP lines from Bioproject

PRJNA559813 [66]. We assembled draft genomes using all reads with Canu v1.7.1 [152] with

the relaxed corrected error rate and the parameters "genomeSize=160m useGrid=false

stopOnReadQuality=false corMinCoverage=0 corOutCoverage=100 correctedErrorRate=0.100".

We then mapped PacBio reads to the reference genome with minimap2 (downloaded August 28,

2018) with the "-x map-bp" parameter setting [153]. We then used unmapped reads and reads

mapping to heterochromatic regions to generate an additional "heterochromatin-enriched"

assembly. We merged the two assemblies into a single assembly with quickmerge [154] and

polished the assemblies with Racon v1.3.1 [155] by mapping the PacBio reads three times to the

assembly using minimap2 [153]. We also polished the GDL assemblies twice with Illumina

reads from BioProject PRJNA268111 [147], the DSPR assemblies with reads from

https://wfitch.bio.uci.edu/~dspr/index.html (last accessed September 6, 2021 [156]), BioProject

PRJNA252961 for Oregon-R, and the DGRP assemblies with reads from BioProject

PRJNA36679 [64, 65]. We annotated all assemblies with Repeatmasker v4.1.0 and our

Drosophila TE library with the updated *G2/Jockey-3* consensus sequence [138] (File S10). We

identified centromere island sequences in assemblies using a combination of methods to find

sequences corresponding to the centromere islands identified in our reference genome [53].

These include aligning assemblies to the centromere islands with MUMMER v.3.23 nucmer

[157], identification of similar sequences with BLAST 2.10.0+ [158], and identifying contigs

enriched in *G2/Jockey-3* or satellite DNA repeats enriched around the centromere islands [53].

We also used MEME [159] to detect motifs enriched in the flanking regions of young insertions

in our assemblies. Additional details of our analysis with long-read data, the results, and the

MEME motif analysis are in the Supplemental Text.

Insertion polymorphism validation

We validated TE insertion predictions in three ways: 1) we used short reads to confirm TE

predictions in long read assemblies (see Supplemental Text); 2) we used long read assemblies to

confirm TE predictions in short read data (see Supplemental Text); and 3) we selected a subset to

validate with PCR (File S3). We used the results from these analyses to establish criteria for

including a predicted insertion in our analyses. For PCR, we selected a subset of 59 *G2/Jockey-3*

copies to confirm in four GDL lines; B59, I23, T29A, and ZH26. We included copies from each

chromosome (X, 2, 3, 4), each chromatin region (euchromatin, heterochromatin, and

centromere), and both reference and non-reference copies. We were able to detect 33 copies

including 13 copies in the centromere and seven *de novo* insertions (File S3).

We used permutation tests to test for significant differences in density of recent *G2/Jockey-3*

insertions between genome regions. For these tests, the genome was divided into 90-kb windows

(average size of a centromere, excluding the Y chromosome and 2nd chromosome) and

categorized as centromere, heterochromatin, or euchromatin. We estimated *G2/Jockey-3* counts

in each window based on final insertion polymorphisms detected by short-read and long-read

datasets separately. We excluded the Y chromosome from these analyses. To test for enrichment

of *G2/Jockey-3* counts within categories, we permuted the datasets 10,000 times using regioneR

1.18.1 ([160] ntimes=10000, randomize.function=resampleRegions,

evaluate.function=numOverlaps). Results were similar for permutation tests with the whole

genome, excluding heterochromatin windows, and excluding euchromatin windows.

G2/Jockey-3 sequence and phylogenetic analysis in D. melanogaster

We extracted all *G2/Jockey-3* sequences from the annotated assemblies using BEDTools v2.29.2

[139] and classified the insertions as reference or non-reference (*de novo*). We classified

insertions as homologous to those in our reference if they were both fragments of similar size

(+/- 10 bp) and within +100 bp of the same location in the reference genome. We identified

corresponding locations by aligning the assemblies to the reference genome with nucmer within

MUMMER v.3.23 [157]. Insertions were classified as *de novo* if they differed in fragment size

length and position of reference insertions, found outside the reference fragment position range,

and located within breaks in the nucmer alignment to the reference genome.

We performed phylogenetic analyses for candidate *G2/Jockey-3* sequences identified in our

genome assemblies. We used *G2/Jockey-3* sequences from the reference and the non-reference

insertions discovered in the DSPR and GDL assemblies. If a non-reference insertion was found

in multiple assemblies, we randomly selected an assembly from which to extract that insertion to

include in our analysis. Prior to phylogenetic analysis, we filtered sequences to exclude

fragments shorter than 400 bp or ~10% of the length of a full *G2/Jockey-3* element. We aligned

the sequences with MAFFT [161] with the *D. simulans G2/Jockey-3* consensus as the outgroup

and inferred a maximum likelihood phylogenetic tree with RAxML v8.2.11 [162]. To study the

evolution of *G2/Jockey-3* on the Y chromosome, we inferred a maximum likelihood phylogeny

of all fragments exclusive to the Y chromosome regardless of size. We determined the relative

age of individual insertions using divergence from the consensus from the Repeatmasker output

and the number of substitutions unique to a fragment divided by fragment length. "Young"

insertions were either < 1% diverged from the consensus sequence or *de novo* insertions

exclusive to a single genome assembly and < 2% diverged from the consensus sequence. We

calculated pairwise divergence between fragments with MEGA X [163] using the Kimura 2-

parameter model assuming uniform rates among all sites. We estimated gene conversion rates

between elements using the same method outlined from Chang & Larracuente [73] based on

Ohta's ɑ equation [164] and Rozen's equation [165]. We used the following equation for our

calculation of gene conversion:

772
$$c = \frac{2u(n-1)}{\pi}$$

We determined the rate at which a number of sequences (*n*) homogenize each other per

generation (*c*) based on the point mutation rate (*u*) and divergence among the sequences (*π*). We

grouped reference *G2/Jockey-3* sequences located within the same contig except for insertions on

the Y centromere. Fragments from the Y centromere were grouped based on the

phylogenetically-defined classes. We aligned the sequences with MAFFT [161] and calculated

divergence with MEGA X [163]. We normalized *c* based on gene conversion tract length of a

single event (*c_g*) based on previously estimated 400 bp tract length of a single event in *D.*

melanogaster [92, 93]:

781
$$c_g = \frac{c}{400 \; x \; 2}$$

Age-of-allele test

Natural selection, transposition, and frequency of insertion influence TE evolution within

populations. We used an age-of-allele test [70] to determine departure from neutral evolution of

individual insertions based on population frequency, age, and demographic history in *D.*

melanogaster. The age-of-allele test determines the probability distribution of an insertion

frequency within a population under neutrality conditioned on the age of the insertion. The age is

inferred by the number of unique substitutions within a TE sequence accumulated since the time

of the insertion. To determine the number of unique substitutions, we aligned insertions with

MAFFT [161] for *G2/Jockey-3*, *BS*, *G5*, *Doc*, and *Doc2* detected as reference insertions in the

McClintock output. Unique substitutions were quantified from each individual insertion using

Biostrings [166]. We calculated the probability distribution of the frequency of each unique

insertion with the age-of-allele test script from Blumenstiel et al. [70] in Mathematica v12.0.0.0.

The probability distribution is also conditioned on population demography of North American *D.*

melanogaster populations outlined in the original paper [70]. North American *D. melanogaster*

populations experienced two bottlenecks in their migration from Africa [167, 168] that led to

non-equilibrium TE evolution [70]. We compared the probability distribution to the frequency of

insertions detected in the 28 DGRP samples with >25X average coverage of the centromere

islands. We excluded insertions in the Y chromosome given the high rate of duplication. The

expected frequency of each insertion in the population is the probability of observing an insertion

as frequent or more frequent within the population multiplied by the observed number of

insertions in our population sample. We calculated significant deviations from neutrality as

probability of observing as many or fewer, or as many or greater, copies < 0.1.

COMPETING INTERESTS STATEMENT

The authors have no competing interests.

DATA AVAILABILITY STATEMENT

All scripts for data analysis and data for figure generation are available on Github at

https://github.com/LarracuenteLab/Dmel_Jockey-3_Evolution. The genome assemblies, genome

annotations, and large datasets including the McClintock outputs will be available on Dryad

(DOI forthcoming).

ACKNOWLEDGEMENTS

This work was funded by the National Science Foundation (NSF MCB 1844693) and University

of Rochester funds to AML. We thank the Mellone lab, Dr. Justin Blumenstiel, and the

Larracuente lab for helpful discussion. We also thank the University of Rochester Center for

Integrated Resource Computing for access to computing cluster resources.

REFERENCES

1. Allshire RC, Karpen GH: **Epigenetic regulation of centromeric chromatin: old dogs,**
new tricks? *Nat Rev Genet* 2008, **9:**923-937.
2. Garrido-Ramos MA: **Satellite DNA: An Evolving Topic.** *Genes* 2017, **8**.
3. Hartley G, O'Neill RJ: **Centromere Repeats: Hidden Gems of the Genome.** *Genes*
2019, **10**.
4. Pidoux AL, Allshire RC: **Kinetochore and heterochromatin domains of the fission**
yeast centromere. *Chromosome Res* 2004, **12:**521-534.
5. Ohzeki J-I, Bergmann JH, Kouprina N, Noskov VN, Nakano M, Kimura H, Earnshaw
WC, Larionov V, Masumoto H: **Breaking the HAC Barrier: histone H3K9**
acetyl/methyl balance regulates CENP-A assembly. *EMBO J* 2012, **31:**2391-2402.

6.   Choo KH: **Centromerization.** *Trends Cell Biol* 2000, **10:**182-188.

7.   Henikoff S, Dalal Y: **Centromeric chromatin: what makes it unique?** *Curr Opin*
Genet Dev 2005, **15:**177-184.

8.   Henikoff S, Furuyama T: **Epigenetic inheritance of centromeres.** *Cold Spring Harb*
Symp Quant Biol 2010, **75:**51-60.

9.   Black BE, Cleveland DW: **Epigenetic centromere propagation and the nature of**
CENP-a nucleosomes. *Cell* 2011, **144:**471-479.

10.   Mendiburo MJ, Padeken J, Fülöp S, Schepers A, Heun P: **Drosophila CENH3 is**
sufficient for centromere formation. *Science* 2011, **334:**686-690.

11.   McKinley KL, Cheeseman IM: **The molecular basis for centromere identity and**
function. *Nat Rev Mol Cell Biol* 2016, **17:**16-29.

12.   Blower MD, Sullivan BA, Karpen GH: **Conserved Organization of Centromeric**
Chromatin in Flies and Humans. *Developmental Cell* 2002, **2:**319-330.

13.   Ribeiro SA, Vagnarelli P, Dong Y, Hori T, McEwen BF, Fukagawa T, Flors C, Earnshaw
WC: **A super-resolution map of the vertebrate kinetochore.** *Proc Natl Acad Sci U S A*
2010, **107:**10484-10489.

14.   Bergmann JH, Rodríguez MG, Martins NMC, Kimura H, Kelly DA, Masumoto H,
Larionov V, Jansen LET, Earnshaw WC: **Epigenetic engineering shows H3K4me2 is**
required for HJURP targeting and CENP-A assembly on a synthetic human
kinetochore. *EMBO J* 2011, **30:**328-340.

15.   Martins NMC, Bergmann JH, Shono N, Kimura H, Larionov V, Masumoto H, Earnshaw
WC: **Epigenetic engineering shows that a human centromere resists silencing**
mediated by H3K27me3/K9me3. *Mol Biol Cell* 2016, **27:**177-196.

16.   Lo AW, Magliano DJ, Sibson MC, Kalitsis P, Craig JM, Choo KH: **A novel chromatin**
immunoprecipitation and array (CIA) analysis identifies a 460-kb CENP-A-binding
neocentromere DNA. *Genome Res* 2001, **11:**448-457.

17.   Alonso A, Fritz B, Hasson D, Abrusan G, Cheung F, Yoda K, Radlwimmer B, Ladurner
AG, Warburton PE: **Co-localization of CENP-C and CENP-H to discontinuous**
domains of CENP-A chromatin at human neocentromeres. *Genome Biol* 2007,
8:R148.

18.   Hasson D, Panchenko T, Salimian KJ, Salman MU, Sekulic N, Alonso A, Warburton PE,
Black BE: **The octamer is the major form of CENP-A nucleosomes at human**
centromeres. *Nat Struct Mol Biol* 2013, **20:**687-695.

19.   Nergadze SG, Piras FM, Gamba R, Corbo M, Cerutti F, McCarter JGW, Cappelletti E,
Gozzo F, Harman RM, Antczak DF, et al: **Birth, evolution, and transmission of**
satellite-free mammalian centromeric domains. *Genome Res* 2018, **28:**789-799.

20.   Kasai F, Garcia C, Arruga MV, Ferguson-Smith MA: **Chromosome homology between**
chicken (*Gallus gallus domesticus*) and the red-legged partridge (*Alectoris rufa*);
evidence of the occurrence of a neocentromere during evolution. *Cytogenet Genome*
Res 2003, **102:**326-330.

21.   Marshall OJ, Chueh AC, Wong LH, Choo KHA: **Neocentromeres: new insights into**
centromere structure, disease development, and karyotype evolution. *Am J Hum*
Genet 2008, **82:**261-282.

22.   Rocchi M, Archidiacono N, Schempp W, Capozzi O, Stanyon R: **Centromere**
repositioning in mammals. *Heredity* 2012, **108:**59-67.

23.   Miller JT, Dong F, Jackson SA, Song J, Jiang J: **Retrotransposon-related DNA**
sequences in the centromeres of grass chromosomes. *Genetics* 1998, **150:**1615-1623.

24.   Presting GG, Malysheva L, Fuchs J, Schubert I: **A Ty3/gypsy retrotransposon-like**
sequence localizes to the centromeric regions of cereal chromosomes. *Plant J* 1998,
16:721-728.

25.   Cheng Z, Dong F, Langdon T, Ouyang S, Buell CR, Gu M, Blattner FR, Jiang J:
Functional rice centromeres are marked by a satellite repeat and a centromere-
specific retrotransposon. *Plant Cell* 2002, **14:**1691-1704.

26.   Nagaki K, Cheng Z, Ouyang S, Talbert PB, Kim M, Jones KM, Henikoff S, Buell CR,
Jiang J: **Sequencing of a rice centromere uncovers active genes.** *Nat Genet* 2004,
36:138-145.

27.   Nagaki K, Song J, Stupar RM, Parokonny AS, Yuan Q, Ouyang S, Liu J, Hsiao J, Jones
KM, Dawe RK, et al: **Molecular and cytological analyses of large tracks of**
centromeric DNA reveal the structure and evolutionary dynamics of maize
centromeres. *Genetics* 2003, **163:**759-770.

28.   Ferreri GC, Brown JD, Obergfell C, Jue N, Finn CE, O'Neill MJ, O'Neill RJ: **Recent**
amplification of the kangaroo endogenous retrovirus, KERV, limited to the
centromere. *J Virol* 2011, **85:**4761-4771.

29.   Ferreri GC, Marzelli M, Rens W, O'Neill RJ: **A centromere-specific retroviral element**
associated with breaks of synteny in macropodine marsupials. *Cytogenet Genome Res*
2004, **107:**115-118.

30.   de Sotero-Caio CG, Cabral-de-Mello DC, Calixto MdS, Valente GT, Martins C, Loreto
V, de Souza MJ, Santos N: **Centromeric enrichment of LINE-1 retrotransposons and**
its significance for the chromosome evolution of Phyllostomid bats. *Chromosome Res*
2017, **25:**313-325.

31.   Carbone L, Harris RA, Gnerre S, Veeramah KR, Lorente-Galdos B, Huddleston J, Meyer
TJ, Herrero J, Roos C, Aken B, et al: **Gibbon genome and the fast karyotype evolution**
of small apes. *Nature* 2014, **513:**195-201.

32.   Carbone L, Harris RA, Mootnick AR, Milosavljevic A, Martin DIK, Rocchi M, Capozzi
O, Archidiacono N, Konkel MK, Walker JA, et al: **Centromere remodeling in** *Hoolock*
leuconedys **(Hylobatidae) by a new transposable element unique to the gibbons.**
Genome Biol Evol 2012, **4:**648-658.

33.   Contreras-Galindo R, Kaplan MH, He S, Contreras-Galindo AC, Gonzalez-Hernandez
MJ, Kappes F, Dube D, Chan SM, Robinson D, Meng F, et al: **HIV infection reveals**
widespread expansion of novel centromeric human endogenous retroviruses.
Genome Res 2013, **23:**1505-1513.

34.   Zahn J, Kaplan MH, Fischer S, Dai M, Meng F, Saha AK, Cervantes P, Chan SM, Dube
D, Omenn GS, et al: **Expansion of a novel endogenous retrovirus throughout the**
pericentromeres of modern humans. *Genome Biol* 2015, **16:**74.

35.   Klein SJ, O'Neill RJ: **Transposable elements: genome innovation, chromosome**
diversity, and centromere conflict. *Chromosome Res* 2018, **26:**5-23.

36.   Birchler JA, Presting GG: **Retrotransposon insertion targeting: a mechanism for**
homogenization of centromere sequences on nonhomologous chromosomes. *Genes*
Dev 2012, **26:**638-640.

37. Sultana T, Zamborlini A, Cristofari G, Lesage P: **Integration site selection by**
retroviruses and transposable elements in eukaryotes. *Nature Reviews Genetics* 2017,
18:292-308.
38. Wong LH, Choo KHA: **Evolutionary dynamics of transposable elements at the**
centromere. *Trends Genet* 2004, **20:**611-616.
39. Sharma A, Wolfgruber TK, Presting GG: **Tandem repeats derived from centromeric**
retrotransposons. *BMC Genomics* 2013, **14**.
40. Bourc'his D, Bestor TH: **Meiotic catastrophe and retrotransposon reactivation in**
male germ cells lacking Dnmt3L. *Nature* 2004, **431:**96-99.
41. Kato M, Takashima K, Kakutani T: **Epigenetic control of CACTA transposon mobility**
in Arabidopsis thaliana. *Genetics* 2004, **168:**961-969.
42. Ariumi Y: **Guardian of the Human Genome: Host Defense Mechanisms against**
LINE-1 Retrotransposition. *Front Chem* 2016, **4:**28.
43. Chan FL, Wong LH: **Transcription in the maintenance of centromere chromatin**
identity. *Nucleic Acids Res* 2012, **40:**11178-11188.
44. Wong LH, Brettingham-Moore KH, Chan L, Quach JM, Anderson MA, Northrop EL,
Hannan R, Saffery R, Shaw ML, Williams E, Choo KHA: **Centromere RNA is a key**
component for the assembly of nucleoproteins at the nucleolus and centromere.
Genome Res 2007, **17:**1146-1160.
45. Quénet D, Dalal Y: **A long non-coding RNA is required for targeting centromeric**
protein A to the human centromere. *Elife* 2014, **3:**e03254.
46. Catania S, Pidoux AL, Allshire RC: **Sequence features and transcriptional stalling**
within centromere DNA promote establishment of CENP-A chromatin. *PLoS Genet*
2015, **11:**e1004986.
47. Chen C-C, Bowers S, Lipinszki Z, Palladino J, Trusiak S, Bettini E, Rosin L, Przewloka
MR, Glover DM, O'Neill RJ, Mellone BG: **Establishment of Centromeric Chromatin**
by the CENP-A Assembly Factor CAL1 Requires FACT-Mediated Transcription.
Dev Cell 2015, **34:**73-84.
48. Hall LE, Mitchell SE, O'Neill RJ: **Pericentric and centromeric transcription: a**
perfect balance required. *Chromosome Research* 2012, **20:**535-546.
49. Talbert PB, Henikoff S: **Transcribing Centromeres: Noncoding RNAs and**
Kinetochore Assembly. *Trends in Genetics* 2018, **34:**587-599.
50. Chuong EB, Elde NC, Feschotte C: **Regulatory activities of transposable elements:**
from conflicts to benefits. *Nat Rev Genet* 2017, **18:**71-86.
51. Jain M, Olsen HE, Turner DJ, Stoddart D, Bulazel KV, Paten B, Haussler D, Willard HF,
Akeson M, Miga KH: **Linear assembly of a human centromere on the Y**
chromosome. *Nature Biotechnology* 2018, **36:**321-323.
52. Logsdon GA, Vollger MR, Hsieh P, Mao Y, Liskovykh MA, Koren S, Nurk S, Mercuri
L, Dishuck PC, Rhie A, et al: **The structure, function and evolution of a complete**
human chromosome 8. *Nature* 2021, **593:**101-107.
53. Chang C-H, Chavan A, Palladino J, Wei X, Martins NMC, Santinello B, Chen C-C,
Erceg J, Beliveau BJ, Wu C-T, et al: **Islands of retroelements are major components**
of Drosophila centromeres. *PLoS Biol* 2019, **17:**e3000241.
54. Belser C, Baurens F-C, Noel B, Martin G, Cruaud C, Istace B, Yahiaoui N, Labadie K,
Hřibová E, Doležel J, et al: **Telomere-to-telomere gapless chromosomes of banana**
using nanopore sequencing. *Commun Biol* 2021, **4:**1047.

55. Wang B, Yang X, Jia Y, Xu Y, Jia P, Dang N, Wang S, Xu T, Zhao X, Gao S, et al:
High-quality Arabidopsis thaliana Genome Assembly with Nanopore and HiFi Long
Reads. *Genomics Proteomics Bioinformatics* 2021.

56. Naish M, Alonge M, Wlodzimierz P, Tock AJ, Abramson BW, Schmucker A,
Mandakova T, Jamge B, Lambing C, Kuo P, et al: **The genetic and epigenetic**
landscape of the Arabidopsis centromeres. *Science* 2021, **374**:eabi7489.

57. Hales KG, Korey CA, Larracuente AM, Roberts DM: **Genetics on the Fly: A Primer on**
the Drosophila Model System. *Genetics* 2015, **201**:815-842.

58. Rahman R, Chirn G-W, Kanodia A, Sytnikova YA, Brembs B, Bergman CM, Lau NC:
Unique transposon landscapes are pervasive across *Drosophila melanogaster*
genomes. *Nucleic Acids Res* 2015, **43**:10655-10672.

59. Gerstein MB, Rozowsky J, Yan K-K, Wang D, Cheng C, Brown JB, Davis CA, Hillier L,
Sisu C, Li JJ, et al: **Comparative analysis of the transcriptome across distant species.**
Nature 2014, **512**:445-448.

60. Andersen PR, Tirian L, Vunjak M, Brennecke J: **A heterochromatin-dependent**
transcription machinery drives piRNA expression. *Nature* 2017, **549**:54-59.

61. Laktionov PP, Maksimov DA, Romanov SE, Antoshina PA, Posukh OV, White-Cooper
H, Koryakov DE, Belyakin SN: **Genome-wide analysis of gene regulation mechanisms**
during Drosophila spermatogenesis. *Epigenetics Chromatin* 2018, **11**:14.

62. Tettey TT, Gao X, Shao W, Li H, Story BA, Chitsazan AD, Glaser RL, Goode ZH,
Seidel CW, Conaway RC, et al: **A Role for FACT in RNA Polymerase II Promoter-**
Proximal Pausing. *Cell Rep* 2019, **27**:3770-3779.e3777.

63. Metcalfe CJ, Casane D: **Modular organization and reticulate evolution of the ORF1**
of Jockey superfamily transposable elements. *Mobile DNA* 2014, **5**:19.

64. Huang W, Massouras A, Inoue Y, Peiffer J, Ràmia M, Tarone AM, Turlapati L, Zichner
T, Zhu D, Lyman RF, et al: **Natural variation in genome architecture among 205**
***Drosophila melanogaster* Genetic Reference Panel lines.** *Genome Res* 2014, **24**:1193-
1208.

65. Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y,
Magwire MM, Cridland JM, et al: **The *Drosophila melanogaster* Genetic Reference**
Panel. *Nature* 2012, **482**:173-178.

66. Rech GE, Radío S, Guirao-Rico S, Aguilera L, Horvath V, Green L, Lindstadt H,
Jamilloux V, Quesneville H, González J: **Population-scale long-read sequencing**
uncovers transposable elements associated with gene expression variation and
adaptive signatures in Drosophila. *Nat Commun* 2022, **13**:1948.

67. Chakraborty M, Emerson JJ, Macdonald SJ, Long AD: **Structural variants exhibit**
widespread allelic heterogeneity and shape variation in complex traits. *Nat Commun*
2019, **10**:4872.

68. Long E, Evans C, Chaston J, Udall JA: **Genomic Structural Variations Within Five**
Continental Populations of *Drosophila melanogaster*. *G3: Genes|Genomes|Genetics*
2018, **8**:3247-3253.

69. Nelson MG, Linheiro RS, Bergman CM: **McClintock: An Integrated Pipeline for**
Detecting Transposable Element Insertions in Whole-Genome Shotgun Sequencing
Data. *G3* 2017, **7**:2763-2778.

70. Blumenstiel JP, Chen X, He M, Bergman CM: **An age-of-allele test of neutrality for**
transposable element insertions. *Genetics* 2014, **196**:523-538.

71.   Kofler R, Betancourt AJ, Schlötterer C: **Sequencing of pooled DNA samples (Pool-Seq)**
1018   **uncovers complex dynamics of transposable element insertions in *Drosophila***
1019   ***melanogaster.*** *PLoS Genet* 2012, **8:**e1002487.
1020   72.   Kofler R, Nolte V, Schlötterer C: **Tempo and Mode of Transposable Element Activity**
1021   **in Drosophila.** *PLoS Genet* 2015, **11:**e1005406.
1022   73.   Chang C-H, Larracuente AM: **Heterochromatin-Enriched Assemblies Reveal the**
1023   **Sequence and Organization of the *Drosophila melanogaster* Y Chromosome.**
1024   *Genetics* 2019, **211:**333-348.
1025   74.   Ghildiyal M, Zamore PD: **Small silencing RNAs: an expanding universe.** *Nat Rev*
1026   *Genet* 2009, **10:**94-108.
1027   75.   Ernst C, Odom DT, Kutter C: **The emergence of piRNAs against transposon invasion**
1028   **to preserve mammalian genome integrity.** *Nature Communications* 2017, **8**.
1029   76.   Bergman CM, Quesneville H, Anxolabéhère D, Ashburner M: **Recurrent insertion and**
1030   **duplication generate networks of transposable element sequences in the *Drosophila***
1031   ***melanogaster* genome.** *Genome Biol* 2006, **7:**R112.
1032   77.   Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ:
1033   **Discrete small RNA-generating loci as master regulators of transposon activity in**
1034   **Drosophila.** *Cell* 2007, **128:**1089-1103.
1035   78.   Gunawardane LS, Saito K, Nishida KM, Miyoshi K, Kawamura Y, Nagami T, Siomi H,
1036   Siomi MC: **A slicer-mediated mechanism for repeat-associated siRNA 5' end**
1037   **formation in Drosophila.** *Science* 2007, **315:**1587-1590.
1038   79.   Mohn F, Sienski G, Handler D, Brennecke J: **The rhino-deadlock-cutoff complex**
1039   **licenses noncanonical transcription of dual-strand piRNA clusters in Drosophila.**
1040   *Cell* 2014, **157:**1364-1379.
1041   80.   Parhad SS, Tu S, Weng Z, Theurkauf WE: **Adaptive Evolution Leads to Cross-Species**
1042   **Incompatibility in the piRNA Transposon Silencing Machinery.** *Dev Cell* 2017,
1043   **43:**60-70.e65.
1044   81.   Rozhkov NV, Aravin AA, Zelentsova ES, Schostak NG, Sachidanandam R, McCombie
1045   WR, Hannon GJ, Evgen'ev MB: **Small RNA-based silencing strategies for transposons**
1046   **in the process of invading Drosophila species.** *RNA* 2010, **16:**1634-1645.
1047   82.   Quénerch'du E, Anand A, Kai T: **The piRNA pathway is developmentally regulated**
1048   **during spermatogenesis in Drosophila.** *RNA* 2016, **22:**1044-1054.
1049   83.   Wei X, Eickbush DG, Speece I, Larracuente AM: **Heterochromatin-dependent**
1050   **transcription of satellite DNAs in the *Drosophila melanogaster* female germline.** *Elife*
1051   2021, **10**.
1052   84.   Petrov DA, Fiston-Lavier A-S, Lipatov M, Lenkov K, González J: **Population genomics**
1053   **of transposable elements in *Drosophila melanogaster.*** *Mol Biol Evol* 2011, **28:**1633-
1054   1644.
1055   85.   Montgomery E, Charlesworth B, Langley CH: **A test for the role of natural selection in**
1056   **the stabilization of transposable element copy number in a population of *Drosophila***
1057   ***melanogaster.*** *Genetical Research* 1987, **49:**31-41.
1058   86.   Charlesworth B, Campos JL: **The relations between recombination rate and patterns**
1059   **of molecular variation and evolution in Drosophila.** *Annu Rev Genet* 2014, **48:**383-
1060   403.
1061   87.   Kent TV, Uzunović J, Wright SI: **Coevolution between transposable elements and**
1062   **recombination.** *Philos Trans R Soc Lond B Biol Sci* 2017, **372**.

88. Bergman CM, Bensasson D: **Recent LTR retrotransposon insertion contrasts with waves of non-LTR insertion since speciation in *Drosophila melanogaster*.** *Proc Natl Acad Sci U S A* 2007, **104:**11340-11345.

89. Blumenstiel JP, Hartl DL, Lozovsky ER: **Patterns of insertion and deletion in contrasting chromatin domains.** *Mol Biol Evol* 2002, **19:**2211-2225.

90. Chang CH, Gregory LE, Gordon KE, Meiklejohn CD, Larracuente AM: **Unique structure and positive selection promote the rapid divergence of Drosophila Y chromosomes.** *Elife* 2022, **11**.

91. Comeron JM, Ratnappan R, Bailin S: **The many landscapes of recombination in *Drosophila melanogaster*.** *PLoS Genet* 2012, **8:**e1002905.

92. Miller DE, Smith CB, Kazemi NY, Cockrell AJ, Arvanitakis AV, Blumenstiel JP, Jaspersen SL, Hawley RS: **Whole-Genome Analysis of Individual Meiotic Events in *Drosophila melanogaster* Reveals That Noncrossover Gene Conversions Are Insensitive to Interference and the Centromere Effect.** *Genetics* 2016, **203:**159-171.

93. Miller DE, Takeo S, Nandanan K, Paulson A, Gogol MM, Noll AC, Perera AG, Walton KN, Gilliland WD, Li H, et al: **A Whole-Chromosome Analysis of Meiotic Recombination in *Drosophila melanogaster*.** *G3* 2012, **2:**249-260.

94. Khurana JS, Wang J, Xu J, Koppetsch BS, Thomson TC, Nowosielska A, Li C, Zamore PD, Weng Z, Theurkauf WE: **Adaptation to P Element Transposon Invasion in *Drosophila melanogaster*.** *Cell* 2011, **147:**1551-1563.

95. Kofler R: **Dynamics of Transposable Element Invasions with piRNA Clusters.** *Mol Biol Evol* 2019, **36:**1457-1472.

96. Said I, McGurk MP, Clark AG, Barbash DA: **Patterns of piRNA Regulation in Drosophila Revealed through Transposable Element Clade Inference.** *Mol Biol Evol* 2022, **39**.

97. Kofler R, Senti K-A, Nolte V, Tobler R, Schlötterer C: **Molecular dissection of a natural transposable element invasion.** *Genome Res* 2018, **28:**824-835.

98. Anxolabéhère D, Kidwell MG, Periquet G: **Molecular characteristics of diverse populations are consistent with the hypothesis of a recent invasion of *Drosophila melanogaster* by mobile P elements.** *Mol Biol Evol* 1988, **5:**252-269.

99. Hill T, Schlotterer C, Betancourt AJ: **Hybrid Dysgenesis in *Drosophila simulans* Associated with a Rapid Invasion of the P-Element.** *PLoS Genet* 2016, **12:**e1005920.

100. Wolfgruber TK, Sharma A, Schneider KL, Albert PS, Koo D-H, Shi J, Gao Z, Han F, Lee H, Xu R, et al: **Maize centromere structure and evolution: sequence analysis of centromeres 2 and 5 reveals dynamic Loci shaped primarily by retrotransposons.** *PLoS Genet* 2009, **5:**e1000743.

101. Schneider KL, Xie Z, Wolfgruber TK, Presting GG: **Inbreeding drives maize centromere evolution.** *Proc Natl Acad Sci U S A* 2016, **113:**E987-996.

102. Tsukahara S, Kawabe A, Kobayashi A, Ito T, Aizu T, Shin-i T, Toyoda A, Fujiyama A, Tarutani Y, Kakutani T: **Centromere-targeted de novo integrations of an LTR retrotransposon of *Arabidopsis lyrata*.** *Genes Dev* 2012, **26:**705-713.

103. Tsukahara S, Kobayashi A, Kawabe A, Mathieu O, Miura A, Kakutani T: **Bursts of retrotransposition reproduced in Arabidopsis.** *Nature* 2009, **461:**423-426.

104. Langley CH, Montgomery E, Hudson R, Kaplan N, Charlesworth B: **On the role of unequal exchange in the containment of transposable element copy number.** *Genet Res* 1988, **52:**223-235.

105.   Hollister JD, Gaut BS: **Epigenetic silencing of transposable elements: a trade-off**
between reduced transposition and deleterious effects on neighboring gene
expression. *Genome Res* 2009, **19:**1419-1428.
106.   Lee YCG: **The Role of piRNA-Mediated Epigenetic Silencing in the Population**
Dynamics of Transposable Elements in *Drosophila melanogaster. PLoS Genet* 2015,
11:e1005269.
107.   Ahmed I, Sarazin A, Bowler C, Colot V, Quesneville H: **Genome-wide evidence for**
local DNA methylation spreading from small RNA-targeted sequences in
Arabidopsis. *Nucleic Acids Res* 2011, **39:**6919-6931.
108.   Cridland JM, Thornton KR, Long AD: **Gene expression variation in** *Drosophila*
melanogaster **due to rare transposable element insertion alleles of large effect.**
Genetics 2015, **199:**85-93.
109.   Srivastava S, Foltz DR: **Posttranslational modifications of CENP-A: marks of**
distinction. *Chromosoma* 2018, **127:**279-290.
110.   Czech B, Munafò M, Ciabrelli F, Eastwood EL, Fabry MH, Kneuss E, Hannon GJ:
piRNA-Guided Genome Defense: From Biogenesis to Silencing. *Annu Rev Genet*
2018, **52:**131-157.
111.   Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD: **PIWI-interacting RNAs:**
small RNAs with big functions. *Nat Rev Genet* 2019, **20:**89-108.
112.   Onishi R, Yamanaka S, Siomi MC: **piRNA- and siRNA-mediated transcriptional**
repression in Drosophila, mice, and yeast: new insights and biodiversity. *EMBO Rep*
2021, **22:**e53062.
113.   Scelfo A, Fachinetti D: **Keeping the Centromere under Control: A Promising Role**
for DNA Methylation. *Cells* 2019, **8**.
114.   Altemose N, Logsdon GA, Bzikadze AV, Sidhwani P, Langley SA, Caldas GV, Hoyt SJ,
Uralsky L, Ryabov FD, Shew CJ, et al: **Complete genomic and epigenetic maps of**
human centromeres. *Science* 2022, **376:**eabl4178.
115.   Doolittle WF, Ford Doolittle W, Sapienza C: **Selfish genes, the phenotype paradigm**
and genome evolution. *Nature* 1980, **284:**601-603.
116.   Orgel LE, Crick FH: **Selfish DNA: the ultimate parasite.** *Nature* 1980, **284:**604-607.
117.   Dimitri P, Junakovic N: **Revising the selfish DNA hypothesis: new evidence on**
accumulation of transposable elements in heterochromatin. *Trends Genet* 1999,
15:123-124.
118.   Cosby RL, Chang N-C, Feschotte C: **Host-transposon interactions: conflict,**
cooperation, and cooption. *Genes Dev* 2019, **33:**1098-1116.
119.   Danilevskaya ON, Slot F, Traverse KL, Hogan NC, Pardue ML: **Drosophila telomere**
transposon HeT-A produces a transcript with tightly bound protein. *Proc Natl Acad*
Sci U S A 1994, **91:**6679-6682.
120.   Mason JM, Biessmann H: **The unusual telomeres of Drosophila.** *Trends Genet* 1995,
11:58-62.
121.   Pardue M-L, DeBaryshe PG: **Drosophila telomeres: A variation on the telomerase**
theme. *Fly* 2008, **2:**101-110.
122.   Pardue ML, Debaryshe PG: **Drosophila telomere transposons: genetically active**
elements in heterochromatin. *Genetica* 2000, **109:**45-52.
123.   Saint-Leandre B, Nguyen SC, Levine MT: **Diversification and collapse of a telomere**
elongation mechanism. *Genome Res* 2019, **29:**920-931.

124. McGurk MP, Dion-Côté A-M, Barbash DA: **Rapid evolution at the Drosophila**
telomere: transposable element dynamics at an intrinsically unstable locus. *Genetics*
2021, **217**.
125. Moazed D, Bühler M, Buker SM, Colmenares SU, Gerace EL, Gerber SA, Hong EJE,
Motamedi MR, Verdel A, Villén J, Gygi SP: **Studies on the mechanism of RNAi-**
dependent heterochromatin assembly. *Cold Spring Harb Symp Quant Biol* 2006,
71:461-471.
126. Choi ES, Strålfors A, Castillo AG, Durand-Dubief M, Ekwall K, Allshire RC:
Identification of noncoding transcripts from within CENP-A chromatin at fission
yeast centromeres. *J Biol Chem* 2011, **286:**23600-23607.
127. Chan FL, Marshall OJ, Saffery R, Kim BW, Earle E, Choo KHA, Wong LH: **Active**
transcription and essential role of RNA polymerase II at the centromere during
mitosis. *Proc Natl Acad Sci U S A* 2012, **109:**1979-1984.
128. Rošić S, Köhler F, Erhardt S: **Repetitive centromeric satellite RNA is essential for**
kinetochore formation and cell division. *J Cell Biol* 2014, **207:**335-349.
129. Bobkov GOM, Gilbert N, Heun P: **Centromere transcription allows CENP-A to**
transit from chromatin association to stable incorporation. *J Cell Biol* 2018,
217:1957-1972.
130. Ohkuni K, Kitagawa K: **Endogenous transcription at the centromere facilitates**
centromere activity in budding yeast. *Curr Biol* 2011, **21:**1695-1703.
131. Sandler L, Novitski E: **Meiotic Drive as an Evolutionary Force.** *The American*
Naturalist 1957, **91:**105-110.
132. Akera T, Chmátal L, Trimm E, Yang K, Aonbangkhen C, Chenoweth DM, Janke C,
Schultz RM, Lampson MA: **Spindle asymmetry drives non-Mendelian chromosome**
segregation. *Science* 2017, **358:**668-672.
133. Akera T, Trimm E, Lampson MA: **Molecular Strategies of Meiotic Cheating by Selfish**
Centromeres. *Cell* 2019, **178:**1132-1144.e1110.
134. Henikoff S, Ahmad K, Malik HS: **The centromere paradox: stable inheritance with**
rapidly evolving DNA. *Science* 2001, **293:**1098-1102.
135. Malik HS, Henikoff S: **Adaptive Evolution of Cid, a Centromere-Specific Histone in**
Drosophila. *Genetics* 2001, **157:**1293-1298.
136. Kumon T, Ma J, Akins RB, Stefanik D, Nordgren CE, Kim J, Levine MT, Lampson MA:
Parallel pathways for recruiting effector proteins determine centromere drive and
suppression. *Cell* 2021, **184:**4904-4918.e4911.
137. Kipling D, Warburton PE: **Centromeres, CENP-B and Tigger too.** *Trends Genet* 1997,
13:141-145.
138. Khost DE, Eickbush DG, Larracuente AM: **Single-molecule sequencing resolves the**
detailed structure of complex satellite DNA loci in *Drosophila melanogaster.* *Genome*
Res 2017, **27:**709-721.
139. Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic**
features. *Bioinformatics* 2010, **26:**841-842.
140. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper
A, Markowitz S, Duran C, et al: **Geneious Basic: an integrated and extendable**
desktop software platform for the organization and analysis of sequence data.
Bioinformatics 2012, **28:**1647-1649.

141.   Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2.** *Nat Methods*
1201   2012, **9:**357-359.
1202   142.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
1203   Durbin R, Genome Project Data Processing S: **The Sequence Alignment/Map format**
1204   **and SAMtools.** *Bioinformatics* 2009, **25:**2078-2079.
1205   143.   Buske FA, Bodén M, Bauer DC, Bailey TL: **Assigning roles to DNA regulatory motifs**
1206   **using comparative genomics.** *Bioinformatics* 2010, **26:**860-866.
1207   144.   Langmead B: **Aligning short sequencing reads with Bowtie.** *Curr Protoc*
1208   *Bioinformatics* 2010, **Chapter 11:**Unit 11.17.
1209   145.   Han BW, Wang W, Zamore PD, Weng Z: **piPipes: a set of pipelines for piRNA and**
1210   **transposon analysis via small RNA-seq, RNA-seq, degradome- and CAGE-seq,**
1211   **ChIP-seq and genomic DNA sequencing.** *Bioinformatics* 2015, **31:**593-595.
1212   146.   Long AD, Macdonald SJ, King EG: **Dissecting complex traits using the Drosophila**
1213   **Synthetic Population Resource.** *Trends Genet* 2014, **30:**488-495.
1214   147.   Grenier JK, Roman Arguello J, Moreira MC, Gottipati S, Mohammed J, Hackett SR,
1215   Boughton R, Greenberg AJ, Clark AG: **Global Diversity Lines–A Five-Continent**
1216   **Reference Panel of Sequenced Drosophila melanogaster Strains.** *G3:*
1217   *Genes|Genomes|Genetics* 2015, **5:**593-603.
1218   148.   Pedersen BS, Quinlan AR: **Mosdepth: quick coverage calculation for genomes and**
1219   **exomes.** *Bioinformatics* 2018, **34:**867-868.
1220   149.   Navarro-Dominguez B, Chang C-H, Brand CL, Muirhead CA, Presgraves DC,
1221   Larracuente AM: **Epistatic selection on a selfish supergene - drive, recombination,**
1222   **and genetic load.** *Elife* 2022, **11**.
1223   150.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,
1224   Lunter G, Marth GT, Sherry ST, et al: **The variant call format and VCFtools.**
1225   *Bioinformatics* 2011, **27:**2156-2158.
1226   151.   Knaus BJ, Grünwald NJ: **vcfr: a package to manipulate and visualize variant call**
1227   **format data in R.** *Mol Ecol Resour* 2017, **17:**44-53.
1228   152.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM: **Canu: scalable**
1229   **and accurate long-read assembly via adaptive -mer weighting and repeat**
1230   **separation.** *Genome Res* 2017, **27:**722-736.
1231   153.   Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics* 2018,
1232   **34:**3094-3100.
1233   154.   Chakraborty M, Baldwin-Brown JG, Long AD, Emerson JJ: **Contiguous and accurate**
1234   **de novo assembly of metazoan genomes with modest long read coverage.** *Nucleic*
1235   *Acids Res* 2016, **44:**e147.
1236   155.   Vaser R, Sović I, Nagarajan N, Šikić M: **Fast and accurate de novo genome assembly**
1237   **from long uncorrected reads.** *Genome Res* 2017, **27:**737-746.
1238   156.   King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, Long AD,
1239   Macdonald SJ: **Genetic dissection of a model complex trait using the Drosophila**
1240   **Synthetic Population Resource.** *Genome Res* 2012, **22:**1558-1566.
1241   157.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL:
1242   **Versatile and open software for comparing large genomes.** *Genome Biol* 2004, **5:**R12.
1243   158.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search**
1244   **tool.** *J Mol Biol* 1990, **215:**403-410.

159. Bailey TL, Johnson J, Grant CE, Noble WS: **The MEME Suite.** *Nucleic Acids Research*
2015, **43:**W39-W49.
160. Gel B, Diez-Villanueva A, Serra E, Buschbeck M, Peinado MA, Malinverni R:
regioneR: an R/Bioconductor package for the association analysis of genomic
regions based on permutation tests. *Bioinformatics* 2016, **32:**289-291.
161. Katoh K, Standley DM: **MAFFT multiple sequence alignment software version 7:**
improvements in performance and usability. *Mol Biol Evol* 2013, **30:**772-780.
162. Stamatakis A: **RAxML version 8: a tool for phylogenetic analysis and post-analysis**
of large phylogenies. *Bioinformatics* 2014, **30:**1312-1313.
163. Kumar S, Stecher G, Li M, Knyaz C, Tamura K: **MEGA X: Molecular Evolutionary**
Genetics Analysis across Computing Platforms. *Mol Biol Evol* 2018, **35:**1547-1549.
164. Ohta T: **Allelic and nonallelic homology of a supergene family.** *Proc Natl Acad Sci U S*
A 1982, **79:**3251-3254.
165. Rozen S, Skaletsky H, Marszalek JD, Minx PJ, Cordum HS, Waterston RH, Wilson RK,
Page DC: **Abundant gene conversion between arms of palindromes in human and**
ape Y chromosomes. *Nature* 2003, **423:**873-876.
166. **Biostrings: Efficient manipulation of biological strings.**
[https://bioconductor.org/packages/Biostrings]
167. Duchen P, Zivkovic D, Hutter S, Stephan W, Laurent S: **Demographic inference reveals**
African and European admixture in the North American Drosophila melanogaster
population. *Genetics* 2013, **193:**291-301.
168. Li H, Stephan W: **Inferring the demographic history and rate of adaptive**
substitution in Drosophila. *PLoS Genet* 2006, **2:**e166.