Separating the control of moving and holding in post-stroke arm paresis

Alkis M. Hadjiosif[1,2,3], Kahori Kita[1,4], Scott T. Albert[5,6], Robert A. Scheidt[7], Reza Shadmehr[5], and John W. Krakauer[1,8,9,10]

Abstract

Moving and holding still have been posited to be under separate control regimes for both eye and arm movements. The paretic arm after stroke exhibits different abnormalities during rest vs. movement, providing an opportunity to ask whether control of these behaviors is independently affected in stroke. Here, we quantified resting postural abnormalities in stroke patients by measuring their biases in force production as they held their hand still in various locations in a planar workspace, and then assessed the influence of these resting force biases on reaching in the same workspace. We found that patients had marked resting force biases at each location, even when the arm was supported. However, these biases did not transfer to arm-supported planar reaching movements: not during initial acceleration, not in response to mid-trajectory perturbations, and not during deceleration to a stop. Rather, the abnormal resting forces only appeared to switch on after a movement had fully stopped. These findings suggest that moving and holding are functionally separable modes of control. At the same time, we found that the resting biases mirrored characteristics of abnormal synergies during movement: they markedly decreased when arm support was provided; they were higher in more distal positions which require breaking out of flexion; and they scaled with the Fugl-Meyer score for the upper extremity (a measure of intrusion of abnormal synergies during active movement). These three shared features suggest a common mechanism for resting biases and abnormal synergies, which appears to be a contradiction given the functional separation of moving and holding observed in the same patients. To resolve this paradox, we propose a conceptual model that predicts a breakdown in the functional separation between reaching and holding when patients move in the absence of weight support. This conceptual model posits that synergies are the behavioral manifestation of a spillover of posture into movement. Mapping these functional systems onto anatomical and physiological details of lesioned substrate after stroke may provide implementation-level insight into how normal arm motor control is assembled.

Author affiliations:

1 Department of Neurology, Johns Hopkins University, Baltimore MD

2 John A. Paulson School of Engineering and Applied Sciences, Harvard University,
Cambridge MA

3 Center for Neurotechnology and Neurorecovery, Massachusetts General Hospital, Boston
MA

4 Kyoto University School of Medicine, Kyoto Japan

5 Department of Biomedical Engineering, Johns Hopkins University, Baltimore MD

6 University of North Carolina School of Medicine, Chapel Hill NC

7 Department of Biomedical Engineering, Marquette University, Milwaukee WI

8 Department of Neuroscience, Johns Hopkins University, Baltimore MD

9 Department of Physical Medicine and Rehabilitation, Johns Hopkins University, Baltimore
MD

10 Santa Fe Institute, Santa Fe, NM

Correspondence to: Alkis Hadjiosif

101 Merrimac Street, Suite 317C

Boston, MA 02114

alkis@seas.harvard.edu; ahadjiosif@mgh.harvard.edu

Running title: Moving vs Holding After Stroke

Keywords: reaching; holding; hemiparesis; stroke; motor control

Abbreviations: ARAT = Action Research Arm Test; CCW = counter-clockwise; CST =
corticospinal tract; CW = clockwise; FM-UE = Fugl-Meyer score for the Upper Extremity; RST
= reticulospinal tract

Introduction

A longstanding idea in motor control is that moving and holding still (abbreviated to *holding* throughout this paper) rely upon separate control regimes. Such separation has been shown in the oculomotor system by the classic work of David Robinson (Robinson, 1970) who demonstrated that that while the controllers for moving and holding are neurally distinct, the hold controller integrates the motor commands that are generated by the move controller. It has been suggested that a separation between these two modes of control may extend to other effectors (Shadmehr, 2017). In particular, substantial behavioral (Ghez et al., 2007) and physiological (Kurtzer et al., 2005; Shalit et al., 2012) evidence supports the idea that a similar dissociation may govern the control of reaching and holding for the arm (Jayasinghe et al., 2022). Moreover, behavioral evidence suggests that commands that are generated during reaching are integrated to produce a component of the commands that are generated during subsequent holding (Albert et al., 2020). This has raised the possibility that like the control architecture of the oculomotor system, a holding controller may lie downstream of the reaching controller.

Patients with hemiparesis after stroke typically exhibit dissociable deficits in the control of reaching and holding within and across effectors (Garland et al., 2009; Levin, 1996; Mani et al., 2013; Schaefer et al., 2009; Trombly, 1992; Zackowski et al., 2004), making hemiparesis a potent model for understanding the interaction, or dissociation, between these two modes of motor control. The magnitude of reaching and holding abnormalities can be relatively large, making it easier to measure and compare them.

Here, we focused on one particular aspect of holding: resting at a position. One of the most common and readily recognizable motor symptoms after stroke is abnormal resting posture (Twitchell, 1951). For example, the typical hemiparetic arm posture consists of flexion at the fingers, wrist, and elbow (Carr and Kenney, 1992) (Figure 1A). Prominent approaches for the treatment of hemiparesis have been based on the idea that abnormal resting posture has a direct deleterious effect upon movement control; they advocate for adjusting overall posture in ways that may minimize such effects (Bobath, 1982; Carr and Kenney, 1994, 1992).

Our main aim was to determine whether resting abnormalities bleed over into active movements in the post-stroke arm. This is of great interest because the ability to separate moving from holding may be precisely what is lost after stroke, with postural abnormalities contaminating voluntary movement. If this is indeed the case, it might lend credence to the idea in rehabilitation that treating resting abnormalities can benefit voluntary movement.

In Experiment 1, we assessed resting postural abnormalities by measuring resting postural
force biases in patients with stroke using a planar workspace. We investigated how these
force biases varied with arm position, presence of arm support, and overall motor impairment.
In Experiment 2, we proceeded to assess patients' control of reaching and holding in the same
workspace. We separately investigated effects on initial reach and bringing the reach to a
stop, as these two might be separately controlled (Ghez et al., 2007; Scheidt and Ghez, 2007).
We also investigated active holding control *after* the movement was over, by examining
responses to perturbations that attempted to move the arm off the target, in order to confirm
that the same controller is engaged for both passive (as in Experiment 1) and active holding
at the same position.

Materials and methods

Participants and Ethics Statement

Sixteen stroke patients (age: 58.5±17.8 [mean±standard deviation], nine female) and nine
age-range matched healthy control participants (age: 62.6±15.2, six female) were recruited
for this study. Table 1 shows details for each patient whereas Table 2 shows summary
demographics and assessment metrics for patients and controls. Procedures were approved
by the Johns Hopkins Institutional Review Board, and participants provided written informed
consent in accordance with the Declaration of Helsinki.

Eligibility criteria

108 We recruited patients with hemiparesis due to stroke. To be eligible for the study, patients had
109 to be adult, exhibit some movement with the affected arm, and be able to provide informed
110 consent and understand the tasks involved. Exclusion criteria were marked cognitive
111 impairment (assessed based on the Montreal Cognitive Assessment, MoCA, cutoff of 20);
112 severe aphasia or ideomotor apraxia, neglect or hemianopia; and orthopedic or pain issues.

Task Details

Participants were seated in a chair and grasped the handle of a robotic arm; the handle moved
either passively (by itself, Experiment 1) or actively (by the participant, Experiment 2) on the
horizontal plane. In Experiment 2 and in two out of four conditions of Experiment 1,
participants' lower arm was supported using a custom-made air-sled (Figure 1C). Above the
plane was a screen which blocked direct vision of the arm; on this opaque screen we

continuously projected a cursor indicating hand position (diameter: 3 mm), as well as the
currently active target (diameter: 10 mm). Handle position was recorded at 200Hz, whereas
subject-produced forces on the handle were recorded using a 6-axis force transducer.
Experiments typically began with the paretic arm (see specific details below); for healthy
controls, an arm was randomly assigned as primary, with its schedule matching that of the
paretic arm in stroke patients. We decided to prioritize the paretic arm, since the primary
analyses in this paper involve comparing different metrics taken from the paretic arm. We note
that this absence of paretic/non-paretic counterbalancing could potentially have a limiting
effect on interpreting paretic vs. non-paretic comparisons; these comparisons, however, are
secondary, and we see no reason why any ordering effects could have more than a minimal
effect on them.

Figure 1: Framework and experiment setup. (***A***) *A patient exhibiting a typical flexor posture at rest.*
Dashed arrows indicate elements of the posture: shoulder depression, arm adduction/internal rotation,
elbow flexion. If one were to physically constrain the hand in a position away from the resting posture,
the torques involved in each component of the abnormal resting posture translate to a force on the hand
(sketched as a blue arrow); we thus designed an experiment to measure the resting force bias on the
hand, as a marker of the overall postural abnormality. The goal was to compare resting postural force
*biases to active movement control in the same area (**B**). (**C**) Experiment setup. The participant holds*
the handle of the robotic arm; reach targets and cursor position are projected on a screen on top; for
*arm support, the participant's arm is strapped on an armrest (**c**) connected to an air sled (**a**) which rests*
*on the table. Air is provided through tube labeled (**b**). (**D**) Top-down view of setup, illustrating the*
different hand positions where resting postural forces were measured in Experiment 1 (open circles).
Also shown are the five target positions used in the reaching and holding task for Experiment 2 (filled
*red circles). The gray box indicates the workspace depicted in Figure 2. (**E**) Examples of measurements*
of resting force biases. Each panel shows the evolution of resting forces during the 5s holding period
for one participant (same participants as in Figure 2), taken at positions close to midline and distal from

the body, under the same condition (paretic arm, arm support given). Solid lines indicate the force on the x-axis (positive values indicate forces towards the left), whereas dashed lines indicate the force on the y-axis (positive values indicate forces towards the body). The shaded area indicates the time window over which forces were averaged to estimate the resting bias, illustrating how resting biases were relatively stable by the 2s mark. Note that the third panel includes one trial (blue) which was rejected following visual inspection as described in Materials and Methods – Data Exclusion Criteria, due to instability in force production and movement during the hold period.

Experiment 1: Measuring resting postural abnormalities

Following a previous paradigm (Kanade-Mehta et al., 2023; Laczko et al., 2017; Simo et al., 2013), Experiment 1 assessed resting postural forces by having the robot passively move participants to different positions in a 2D workspace and hold them still in each position while it measured the forces they inadvertently exerted. Participants were instructed to maintain grasp on the robotic handle but otherwise rest and not resist the robot's motion as it slowly (5s movement time) moved from one position to the next and held them still (for an additional 5s). The array of positions (see Figure 1) could vary from one participant to the next, with each position visited three times for each block. During the passive moving and holding, a 3 mm white cursor indicating handle position and a 10 mm yellow disk indicating the destination of the passive movement were displayed. Though not essential from the participant's point of view, this allowed the experimenter to monitor the status of the experiment.

Each participant completed 4 blocks, two with each arm and in each arm support condition (air sled, no air sled); the typical order was {paretic, no support} -> {non-paretic, no support} -> {paretic, air sled support} -> {non-paretic, air sled support}; however in four individuals we completed the two paretic blocks first as their hand had to be secured to the handle (with self-adherent wrap) for a stable grasp.

Each block typically took less than 10 minutes to complete, with Experiment 1 lasting about 40 minutes including breaks.

Experiment 2: Assessing reaching control

Using the same workspace as Experiment 1, Experiment 2 assessed motor control in a reaching task. Participants made 10-cm point-to-point reaches across an array of 5 targets (diameter: 10mm) within the workspace located both distal and proximal to the body (Figure 3A), sampling 8 different movement directions. A white cursor (diameter: 3mm), indicating hand position, was visible throughout the experiment. Participants were instructed to try and stop at each target within a 600-800ms window after movement onset. At the end of movement, feedback was provided to indicate whether they were too fast (time <600ms, target

turning red), too slow (time>800ms, target turning blue), or within the right time range (target

"exploding" with a chirping sound).

The experiment was divided into blocks of 96 movements each (12 in each of the 8 movement

directions). It began with three blocks with the paretic arm, followed by three more with the

non-paretic arm, two more with the paretic arm, and ending with two blocks with the non-

paretic arm. The first block with each arm was a familiarization block. Arm support (air sled)

was provided throughout the experiment, and breaks were given between blocks.

Except for two participants, who performed Experiment 1 and Experiment 2 on different days

due to limitations in their schedule, the entirety of each session – consisting of Experiment 1,

Experiment 2 and standard assessments - took place on the same day and typically lasted

about 3.5 hours with breaks given between the blocks as necessary.

Most trials (two-thirds) consisted of unperturbed movements to assess reaching control. In the

remaining third of reaches, a 12N, 70 ms bell-shaped force pulse that was 12N at its peak was

applied by the robot lateral to the ideal movement direction (i.e. the direction formed between

the center of the start position and the center of the target) after participants reached 2cm

away from the starting position (Fine and Thoroughman, 2006; Smith and Shadmehr, 2005).

On half of these trials the perturbation was oriented leftwards with respect to the movement

(counterclockwise pulse) and the other half rightwards (clockwise pulse).

A fraction of trials in each block (20/96) imposed a perturbation after movement in order to

assess active holding control. For these trials, the holding time at the target was extended by

5 to 7 seconds, during which participants were instructed to hold still on the target (to remind

them, the word "HOLD" were shown close to the target). During this extended hold period, a

6N force was gradually imposed over 2s in one of four different directions (45˚, 135˚, 225˚,

315˚), held constant for a pseudorandom time interval uniformly ranging from 3 to 5 seconds,

and then abruptly released. We refer to this as "release perturbation" throughout the paper.

Each block presented each position/release perturbation direction combination exactly once.

Thus, in summary, each 96-movement block consisted of 64 unperturbed movements and 32

movements perturbed with a force pulse (16 clockwise, and 16 counter-clockwise). For 20 out

of the 96 movements in each block, the hold period was extended to test the hold perturbation

(4 trials for each of the 5 target locations, each one of the 4 trials testing one perturbation

direction as shown in Figure 7C).

A different perturbation was applied to movement (pulse) as compared to holding (release) so

as to maximize disruption for each type of control, rather than to enable direct comparison

between them. For holding control, instead of an abruptly-imposed force similar to the pulse used to probe movement control, we used a perturbation in which the arm first held against a 6N force which was then abruptly released. The rationale was that this force preloading would further engage active holding control and thus be better suited to examine it. In line with this, recent work that examined static hold under both types of perturbation – one in which a load is abruptly imposed vs. one in which a load is abruptly removed as we did here – found that the latter was more difficult and hence a better assay for unmasking a holding abnormality (Lowrey et al., 2019).

Data Analysis

Analysis was performed using MATLAB (Mathworks, Natick MA). For Experiment 1, we averaged resting postural forces for the last 3s of the 5s passive holding period for each trial. We excluded the first 2s in order to avoid measuring potential increases in muscle tone that may arise due to the velocity by which the robot passively brought the arm to each measurement position (i.e., avoiding potential effects of spasticity). Recent work using a similar task to examine resting force biases, found that the effects of the velocity at which the robot brought the arm to a position are present for 2s after the end of the passive movement, but then dissipate (Kanade-Mehta et al., 2023). To obtain a measure of the average resting postural force at each position for each individual and condition, we further averaged forces across the three visits to the same position. For comparisons, we focused on resting postural forces on the 5 positions shown in Figure 1D; these forces were obtained directly (when the exact positions were sampled for the individual) or through interpolation (7/16 patients and 3/9 controls).

For Experiment 2, movement onset was defined as the moment in which participants' velocity from the starting position exceeded 3.5 cm/s, movement end was defined as the moment the participant was within the target and moving at a speed of less than 3.5 cm/s. Initial reaching angle was calculated between hand position at movement onset and 150ms later; endpoint reaching angle was calculated between the position of the hand when it crossed within 2cm of the target and 150ms later. In pulse trials, settling time was defined as the time taken from pulse onset to the first moment absolute lateral velocity dipped below 2cm/s and remained so for at least 100ms (or the movement ended). In release perturbation trials, settling time was defined as the time taken from perturbation onset (release of holding force) to the moment when velocity dipped below 2cm/s (and remained below that amount for at least 100ms) and the distance from the target was less than 2cm.

Data exclusion criteria

In Experiment 1, some trials (7.5%) were flagged as erroneous after visual inspection of force and movement profiles. Erroneous here refers to trials where forces appeared unstable and/or there was movement during the robot hold period (please see Figure 1E illustrating an example one such trial, blue curves on the third panel). We excluded these trials from our main analyses. To ensure that this exclusion did not bias our findings, we repeated our analyses including these trials and obtained similar results (Figures S5-E, S6-E, S7-E).

In Experiment 2, we excluded as outliers movements in which initial movement direction (150ms after movement onset) was ≥90° away from target direction. This excluded 0.95% of patients' movements and 0.33% of controls' movements.

Moreover, in some release perturbation trials, patients took a long time to reach the stabilization criterion described in the previous section; mistakenly, our setup limited its recording time to only the first 2s after force release. The exact time to stabilization thus could not be measured for these particular trials, so they had to be excluded from analysis. Though only 13.2±3.3% (mean±SEM) of paretic stabilization trials were thus excluded in the patient population (1.4±0.4% in their non-paretic side, 2.3±2.0% in controls), there were three patients for which excluded trials were 25% or more of all paretic trials. To ensure there are no systematic effects of this issue, we repeated the analysis of Figure 7F (a) by excluding these three patients altogether or (b) by assigning a value of 2.0 seconds to the affected trials. In both cases, we obtained results similar to our main analysis (Figure S7-5).

Stability of resting posture bias measurements in Experiment 1

Figure 1E shows a few examples of the within-trial evolution of force bias measurements during the 5s hold period, illustrating the high within-trial temporal stability in our dataset – particularly after the 2s mark, which begins the measurement window. To systematically assess the stability of our estimates of resting force biases *across* different trials, we estimated the variance attributed to different measurements at the same location (in contrast to the variance attributed to different locations, conditions, and patients). We found that this measurement variance was only 9.0% of the total variance for resting bias magnitude.

Statistical comparisons

In Experiment 1, we used an ANOVA to investigate any effect of conditions {Position (distal targets [furthest three positions] vs. proximal targets [closest two positions]), Support (with/without air sled), FM-UE (continuous)} and their interactions.

In Experiment 2, we used paired t-tests for the within-subject comparisons of outcome
variables (For unperturbed reaching: initial reaching angle, endpoint reaching angle; for in-
movement pulse perturbations: maximum deviation in the pulse direction, settling time; for
holding release perturbations: maximum deviation in the perturbation direction; time to
stabilization; path to stabilization) against the corresponding resting biases measured in
Experiment 1. We performed two types of comparisons:

First, our main analyses took, for each individual, the two instances for which the resting
biases would have the strongest opposing effects. For variables related to unperturbed
reaching, we would take the movement direction for which the resting biases would be oriented
the strongest CCW vs. the movement direction for which the resting biases would be oriented
the strongest CW (two out of eight directions). The corresponding trials (32 for each of the two
instances per participant) were averaged for each individual. For variables related to
responses to pulse perturbation, we would take the two movement directions for which the
resting bias at start would most oppose vs. most align with the pulse. The corresponding trials
(8 for each of the two instances) were averaged for each individual. For variables related to
responses to the release perturbation, we took the two perturbation direction/hold position
combinations for which the resting bias on that position would most oppose vs. most align with
the perturbation. The corresponding trials (4 for each of the two instances) were averaged for
each individual.

Second, we used linear regression to calculate, for each individual, the sensitivities of outcome
variables from Experiment 2 to the corresponding resting biases measured in Experiment 1
using data from all trials (Figures S5-2, S6-2, and S7-2), rather than the most extreme
instances. We then tested whether the average of these per-individual sensitivities was
significantly greater than zero using a paired t-test.

We used the circular statistics toolbox CircStat (Berens, 2009) to estimate the circular
mean±SEM for the direction of resting biases shown in Figure 3B, right.

Fugl-Meyer assessments

305   Assessments were separately scored by AMH and KK with scores subsequently averaged
306   (hence some scores having decimal values). For cases of substantial score differences (3
307   points or more) scores were again reviewed by both raters together.

Results

Patients exhibited resting force biases across the whole workspace

We first assessed post-stroke resting postural abnormalities across a 2D workspace. In Experiment 1, participants grasped the handle of a robotic arm, which passively moved their hand to a series of positions that sampled the workspace in front of them (the setup and example workspace are shown in Figure 1C,D). Participants were instructed to maintain grasp, but otherwise relax their arm and not resist the actions of the robotic arm. Once the participant's arm was passively moved to a given position, the robotic arm would hold still for 5 seconds, enabling us to measure the forces involuntarily exerted by the participant as their hand was held at that location. Each experimental block consisted of three visits to 15-25 positions sampled for each participant, in random order. Participants completed four different experimental blocks: two with each arm, with or without arm weight support (provided by an air sled, Figure 1C).

Patients displayed abnormal postural force biases when the robot held the hand still at various locations across the workspace. Figure 2 shows examples of this for three patients and a typical healthy control participant. The non-paretic arm produced little to no postural force biases, whereas the paretic arm produced substantial postural force biases, particularly when the hand was held in a more distal position. The postural force biases were strongest when patients had to support the weight of the arm against gravity. Moreover, the patient with the highest degree of impairment (top left subplot of Figure 2, as assessed using the Fugl-Meyer score for the Upper Extremity, FM-UE (Fugl-Meyer et al., 1975)) exhibited the strongest resting postural abnormalities.

To aggregate results across all participants, we focused on abnormal resting postural force biases at five specific hand locations, illustrated by the red dots in Figure 2. These locations were chosen as they were within the sampled workspace of all participants (as this workspace could differ from one participant to the next due to different participants' segment lengths), represented positions both distal and proximal to the body, and, importantly, they were also the movement targets used in Experiment 2. Figure 3A shows subject-averaged resting postural forces at each of these five positions and Figure 3B shows the corresponding force magnitudes and directions averaged across subjects and the five positions. There were two key observations: first, postural force biases were stronger at locations further away from the body (average magnitude – paretic side – without arm support: 6.5±1.3N for the three more distal targets vs. 3.5±0.5N for the two more proximal targets, $t_{15}=3.5$, p=0.0030; with arm

support: 3.3±0.5N for the three more distal targets vs. 1.9±0.3N for the two more proximal targets, t_{15}=3.7, p=0.0022) and tended to point towards it (average direction – paretic side: -93.3±6.0° unsupported vs. -136.5±21.8° supported, with -90° directed towards the body); second, postural force biases were roughly halved in magnitude when arm support was provided (average magnitude – paretic side: 5.3±0.9N unsupported vs. 2.8±0.4N supported, t_{15}=4.0, p = 0.0013).

Figure 2: Examples of resting postural force biases. *Shown are three stroke patients and one healthy control. Arrows indicate magnitude and direction of abnormal resting postural forces as measured at the hand at each location. Isoclines and the corresponding color levels provide a visual representation of where these biases would tend to direct the hand (from red towards blue). These isoclines represent different levels of the (spatial) integral of the posture bias-field (with zero referring to the isocline passing through the center position). Postural bias force vectors cross these isoclines perpendicularly. Please see Supplementary materials on how this visual aid was constructed. The red dots are the reach targets, with the center location circled (used in Experiment 2). Note how abnormalities in the paretic side are considerably stronger when arm support is removed. FM-UE: Fugl-Meyer score for the Upper Extremity (0-66).*

Examining the influence of resting postural force biases on active motor control

In Experiment 2, we sought to investigate whether resting postural force biases influence active reaching and holding in the same workspace. Specifically, we measured their effects upon initial reach direction (Figure 4, blue) and approach direction when bringing the reach to a stop (Figure 4, red), as it has been suggested that these two components are separately controlled (Ghez et al., 2007; Scheidt and Ghez, 2007); see also (Karst and Hasan, 1991; Lestienne, 1979). We also asked whether the same controller is engaged while the arm is passively (Experiment 1) vs. actively (Experiment 2) held at the same position. For this purpose, we examined active holding control *after* the movement was over (Figure 4, black), using perturbations that attempted to push the arm off the target.

Figure 3: Average resting postural force biases. (A) *Average resting postural forces for the paretic (red) and non-paretic (cyan) arms of patients, as well as control participants (gray), illustrating how abnormal forces in the paretic arm are stronger in more distal targets and attenuated when arm support is provided (lighter shades). To average across left- and right-hemiparetic patients, left-arm forces were flipped left to right.* ***(B)*** *corresponding average resting postural force magnitudes and resting postural force directions (0 indicating the 3 o'clock direction, with negative values indicating clockwise directions). Errorbars indicate mean±SEM (circular mean±SEM in the case of movement directions).*

Figure 4 Three aspects of active motor control that we tested in Experiment 2. *We separately examined the early part of the reaching movement (blue) and the late part, when the arm was coming to a stop (red). This was done by studying both unperturbed movements at different stages and movements that were perturbed with brief force pulses. In addition, we examined active holding control after the movement was over (black), using perturbations that tried to move the arm away from the held point. Shown is an example of trajectory and (absolute) velocity profiles from the reaching and coming-to-a-stop parts of a trial (left) and active holding against a perturbation after the trial was over (right).*

Resting postural force biases did not affect the control of active reaching in the same workspace

We began by examining the control of active reaching. Participants made 10-cm point-to-point reaching movements within an array of five positions (filled circles in Figure 1D), for a total of 8 different movement directions (Figure 5A). Arm support was provided by the air sled (Figure 1C). Patients' movements were generally impaired, taking more time and traveling a longer path to reach the target than controls (Time to target: 1.59±0.12s for patients' paretic side vs. 0.82±0.03s for controls, p = 0.000063; Path traveled to target 13.4±0.6cm for patients' paretic side vs. 10.8±0.2cm for controls, p = 0.0071, Fig. 5B,C).

We then examined whether patients' movements were affected by the resting postural biases measured in Experiment 1 in the same workspace and under the same arm-support condition. Would there, for example, be a difference when moving the arm through a high-postural bias

area vs. a low-postural bias area? We examined effects of resting postural biases upon the

initial reach vs. the approach to a hold position, as separate mechanisms may be involved in

the control of each phase of movement (Ghez et al., 2007; Hannaford and Stark, 1985; Karst

and Hasan, 1991; Sainburg et al., 1999; Scheidt and Ghez, 2007); resting postural biases

might affect one phase but not the other. Specifically, we examined: (a) whether the direction

of resting postural forces at the start position influenced trajectory deviations near the

beginning of the movement (Figure 5D) or (b) whether direction of resting postural forces at

the end position influenced trajectory deviations near the end of the movement (Figure 5G,

after the participant reached within 2cm of the target).

To investigate within-subject effects, we selected, for each participant, the two movement

directions for which the corresponding postural forces had strongest opposing effects – i.e.

the most rightwards (CW) vs. most leftwards (CCW) with respect to movement direction. The

idea behind picking the most extreme values for each individual was to maximize our

sensitivity in detecting potential effects of resting postural forces upon active movement. Our

analysis found no significant differences in directional biases between these two conditions:

while the selected start-point force biases differed considerably between the most CCW and

CW instances (1.9±0.4N vs. -2.1±0.4N, correspondingly [negative signs indicating CW forces],

t_{15}=5.68, p=0.000044), the corresponding initial angular deviations did not (0.4±2.1° for the

most CCW vs. -1.6±2.2° for the most CW postural force, t_{15}=0.56, p=0.58, Figure 5D-F).

Similarly, while the selected end-point force biases also differed considerably (2.1±0.4N vs. -

1.9±0.4N, for the most CCW vs. CW instances, t_{15}=5.68, p=0.000044), the corresponding

endpoint angular deviations did not (3.6±2.3° for the most CCW vs. 4.9±2.6° for the most CW

postural force, t_{15}=0.50, p=0.62, Figure 5G-I). Figure S5-1 shows the same analyses but for

non-paretic and control data, illustrating that, like in the paretic data, a lack of evidence that

resting posture biases affect active movement.

While selecting the most extreme instances in terms of resting posture biases for each

individual provides the greatest bandwidth to detect effects of such biases upon active

movement, it also does not use the majority of the data (the 6 out of 8 movement directions

corresponding to intermediate strengths of posture biases). We thus conducted an additional

analysis which estimated the sensitivity (slope) of initial or endpoint deviations against the

corresponding resting biases using the entirety of the data. In line with our main analysis, we

found that patients' slopes were not significant in either case (Initial reaching angle vs. posture

bias at start: 1.5±1.4°/N, p = 0.30; endpoint reaching angle vs. posture bias at endpoint: -

1.0±0.8°/N, p = 0.25, Figure S5-2), further illustrating the lack of compelling evidence that

postural abnormalities directly affect active movement.

Figure 5: Abnormal resting postural force biases do not interact with active reaching. (A) *Target array for Experiment 2 (movement task), illustrating the 5 start/end points of reaches and the 8 movement directions. **(B)** Example outwards trajectories (unperturbed trials) for a patient (cyan: non-paretic side; red: paretic side) and a healthy control (gray). **(C)** Subject-averaged reach performance based on either time (top) or path length to target (bottom) indicates impaired reaching control in patients' paretic side. **(D-F)** Within-subject analysis of whether resting postural forces at movement start bias early movement towards their direction. **(D)** Sketch illustrating the concept behind this analysis. Assuming a movement from a start position subject to a strong rightwards resting bias (F_{start}), will that translate to a corresponding rightwards movement bias which can be expressed as the early reach angle θ_{start}? **(E)** For each individual, we selected the target direction for which the counter-clockwise*

(CCW) component of F_{start} was the strongest (red) vs. the target direction for which the clockwise (CW)
component was the strongest (blue). The left panel shows this selection for an example participant:
postural forces at start position were projected lateral to the movement direction, allowing us to select
movement directions for which the lateral component was directed the strongest CCW or CW. The right
*panel shows the magnitude of these selected components across all patients. (**F**) Left: Corresponding*
movement trajectories (rotated so start position is at the bottom and end position at the top) for the
directions selected for the same example participant. Right: Average initial angular deviations, θ_{start},
across the selected directions for each participant. Note the lack of difference between the instances
for which the CCW vs. the CW component of F_{start} was the strongest, and thus no evidence that F_{start}
*impinges upon the movement. (**G-I**) same as D-F but for endpoint resting postural forces, F_{end} and*
endpoint deviations, θ_{end}. Errorbars indicate SEM.

A potential limitation in our data and analysis above is that unperturbed reaching movements
may provide limited sensitivity in detecting effects of resting postural force biases, as any such
effects may be largely compensated by a predetermined motor plan. To address this, we
probed the effect of the biases to responses to less predictable, externally applied mid-
movement force perturbations. In 1/3 of reaching movements in Experiment 2, chosen
randomly, we imposed a 70ms bell-shaped force pulse which was 12N at its peak and acted
lateral to the movement (Smith and Shadmehr, 2005). These pulses were imposed 2cm into
the reach. Half of these pulses were clockwise (CW, blue in Figure 6A) and the other half were
counter-clockwise (CCW, red in Figure 6A). We first verified that these pulses had a clear
effect upon movement: perturbed movements took longer to complete in both patients (paretic
movement time: 1.72±0.13s vs. 1.59±0.12s, t_{15} =5.31, p = 0.000088) and controls (movement
time: 0.88±0.03s vs. 0.82±0.03s, t_8 =5.56, p = 0.00053). Patients generally had impaired
response to these pulses compared to their non-paretic side and healthy controls, deviating
further (maximum lateral deviation – paretic: 2.5±0.1cm vs. non-paretic: 2.1±0.1cm [p=
0.0008] and controls: 2.1±0.1cm [p = 0.0244] - average of CW and CCW pulses) and taking a
longer time to stabilize in the pulse direction (i.e. settling time; paretic: 0.61±0.03s vs. non-
paretic: 0.47±0.01s [p=0.0027] and controls: 0.50±0.01s [p = 0.028]). This is illustrated in
Figure 6B,C.

We then investigated whether resting postural forces played any role in patients' response to
the pulse perturbation. We reasoned that, should resting postural forces play a role, they would
tend to decrease the effect of the pulse if they were in the opposite direction to it, and increase
the effect of the pulse if they were in the same direction. This is because force pulses acting
against the gradient of the postural bias field would act to stretch the already active muscles,
further increasing postural resistance; whereas force pulses acting along the gradient of the
17

postural bias field would act to shorten the same active muscles, reducing postural resistance.

We thus compared the lateral component of resting postural forces in the start position against

(a) the maximum lateral deviation in the direction of each pulse and (b) the time taken to

stabilize in the pulse direction (settling time). In a manner similar to the previous section, we

selected, for each individual, the directions where the starting lateral postural force was most

Figure 6: Responses to pulse perturbations during movement are not affected by resting

postural force biases. (A) Examples of perturbed (red: perturbed with CCW pulse; blue: perturbed

with CW pulse) and unperturbed (gray) outward trajectories - same individuals as in Figure 3B. (B)

Lateral velocity (positive: CCW to movement) before and after pulse onset, and corresponding

responses from controls (gray), illustrating how patients, in response to the pulse, take longer time to

settle and tend to experience larger lateral deviations compared to controls. (C) Summary performance

measures for patients and controls, indicating impaired performance with the paretic side: settling time

(left) and maximum lateral deviation on pulse direction (right). (D) Within-individual analysis: here, for

each individual, we selected the movements for which the starting-position resting postural force would

be either the strongest CCW or CW (left); we then examined the corresponding settling time (middle)

and maximum lateral deviations (right). We find no effects of the most CCW vs. most CW resting

postural forces in either case: there is no evidence for either reduced settling time or increased

maximum lateral deviation for instances where pulse and resting bias are most opposing (open circles)

compared to the instances where pulse and resting bias are most aligned (filled circles).

CCW vs. most CW, and compared the corresponding deviations. We found no clear effect:
when resting postural forces were the most opposed to the pulse (vs. most aligned with it)
there were no clear differences in deviation along the pulse direction, for neither the CCW
pulses (0.97±0.17 vs. 1.30±0.14cm, t_{15} = 1.48, p = 0.16), the CW pulses (1.27±0.16
vs.1.50±0.20cm, t_{15} =0.87, p = 0.40), or with both types of pulses pooled together (1.12±0.12
vs. 1.40±0.11cm for the most-opposed vs. most-aligned pulse/bias instances, t_{15} =1.42, p =
0.18). Similarly, we found no clear differences in settling time between the instances where
resting postural forces were the most opposed to the pulse vs. most aligned with it, for neither
the CCW pulses (0.65±0.07 vs. 0.59±0.07s, t_{15} = 0.58, p = 0.57), the CW pulses (0.64±0.05
vs. 0.61±0.05s, t_{15} =0.40, p = 0.70), or with both types of pulses pooled together (0.64±0.05
vs. 0.60±0.05s, t_{15} =0.74, p = 0.47). For both metrics, we got similar findings when examining
non-paretic and control data (Figure S6-1). In line with the above analyses, we also found no
effects when we instead estimated the sensitivity of either outcome variable against the
corresponding resting biases using data from all trials (for CCW and CW pulses combined,
maximum lateral deviation along the pulse direction 0.07±0.06 cm/N, p=0.21; settling time:
0.00±0.02 s/N, p = 0.92, Figure S6-2).

In summary, we found no evidence for an interaction between resting postural force biases
and the ability to respond to perturbations that were applied during movement.

Resting postural force biases emerged during active holding at the end of movement

Finally, in Experiment 2, we also investigated the relationship between resting postural force
biases and active holding. In 20 out of 96 movements in each block, participants had to hold
steady on the target for an additional 5 to 7 seconds (Figure 7A). During this time, the robot
imposed a 6N force in one of four directions (45˚, 135˚, 225˚, 315˚, as shown in Figure 7B).
This force was gradually applied over two seconds, held at a 6N level for 3-5 seconds, and
then abruptly released, acting to displace the arm in the direction opposite to the original force,
as illustrated in the examples in Figure 7B.

Patients showed impaired capacity to resist and recover from this perturbation (the abrupt
release of the imposed force). The time to stabilization for the paretic side (0.94±0.05s) was
longer compared to the non-paretic side (0.79±0.03s, p = 0.024) and controls (0.78±0.06s,
though this was statistically marginal, p = 0.061) as shown in Figure 7E, left. Moreover,
patients traveled a longer path to stabilization (7.7±0.7cm for the paretic side vs. 5.9±0.3cm

for the non-paretic side [p = 0.012] and 5.4±0.5cm for controls [p = 0.026], Figure 7E, middle)

and they deviated more in the direction of the perturbation (3.3±0.2cm for the paretic side vs.

Figure 7: Responses to release perturbations during holding and their relationship to resting

postural forces. (A) Time course of the perturbation. (B) Example responses (all for the same position

in the workspace) from two patients (top row) and two controls (bottom row). (C) Corresponding

imposed force directions, the abrupt removal of which perturbs the movement in the opposite direction

(compare with B). (D) Examples of tangential velocity profiles after the sudden release to the imposed

hand force, averaged for all trials at the same position for each participant. Dashed line indicates the

2cm/s threshold used to assess time to stabilize. Left; example patient (paretic side); Right; example
*control. Colors correspond to different directions of the imposed hand force. (**E**) Summary of*
performance metrics after the perturbation for the paretic and non-paretic side of patients and healthy
*controls. (**F**) Within-subject analysis of the relationship between resting postural forces in the direction*
of the perturbation vs. performance against the perturbation. For each individual, we selected the two
position/perturbation direction combinations for which resting postural forces were either the most
opposed (green) to the perturbation or the most aligned (blue) with it. From left to right: forces in selected
position/perturbation direction combinations; corresponding path traveled to stabilization;
corresponding time to stabilization; corresponding maximum deviation. Note how the most-opposed
resting bias for each patient is equal and opposite to the their most-aligned resting bias. This is because
the same resting bias, when projected along the direction of two oppositely-directed perturbations
(illustrated in C), it would oppose one with the same magnitude it would align with the other. This
analysis suggests that resisting postural perturbations and restoring hand position after the perturbation
was indeed easier when resting postural forces opposed, rather than were aligned with, the
perturbation. Gray dots indicate individual data; colored dots and errorbars indicate mean±SEM.

2.6±0.1cm for the non-paretic side [p = 0.012] and 2.3±0.2 for controls [p = 0.0078], Figure
7E, right).

To investigate whether resting postural force biases affected the control of actively holding
across workspace locations, we performed a within-individual analysis analogous to the one
we used for unperturbed reaches and moving perturbations. We first projected resting postural
forces upon the directions of the release perturbation to assess the component of the resting
postural force that opposed or aligned with the release perturbation. We then selected, for
each participant, the two position/perturbation direction combinations for which these forces
were either the most opposed (Figure 7F, green) to the perturbation or the most aligned
(Figure 7F, blue) with it. For these selected position/perturbation direction combinations, we
compared capacity to resist and recover from the perturbation, and found that this capacity
was indeed better when the resting postural force was in a direction that opposed the
perturbation (path traveled to stabilization: 3.7±1.0cm vs. 6.3±0.9cm, t_{15}=2.8, p=0.014; time to
stabilization: 0.6±0.1s vs. 0.9±0.1s, t_{15}=3.8, p = 0.0017; maximum deviation: 1.7±0.4cm vs.
3.0±0.4cm, t_{15}=4.6, p = 0.00036, corresponding to instances with the most opposed vs. the
most aligned resting postural force). We did not observe significant differences when we
performed the same analyses for non-paretic and control data (Figure S7-1).

We then proceeded, in a secondary analysis, to estimate the sensitivity of active holding
control to resting postural biases for each individual using data from all trials (Figure S7-2).

Specifically, we estimated the sensitivity of all three outcome variables (path to stabilization, time to stabilization, and maximum deviation) against the resistive component of the resting bias for each trial. Consistent with our main analysis in the previous paragraph, we found significant sensitivities in all cases (path to stabilization: sensitivity of -0.69 ± 0.30 cm/N, $p = 0.034$; time to stabilization: sensitivity of -0.050 ± 0.009 s/N, $p = 0.000089$; maximum deviation: -0.30 ± 0.12 cm/N, $p= 0.027$). The negative slopes indicate reductions in these metrics – i.e. better performance – as the corresponding resting bias is more resistive. In short, our perturbations revealed that resting flexor biases "switched on" after movement was over, providing evidence for separate control between moving and holding.

Relating resting postural force biases to the Fugl-Meyer scale for the Upper Extremity

The observation, from Experiment 1, that resting postural force biases are reduced by external arm support bears parallels to the same effect of arm support upon abnormal synergies active during movement (Sukal et al., 2007). Here, "synergies" refer to abnormal co-activation patterns across joints that manifest as the patient tries to move – for example, the elbow involuntarily flexing as the patient tries to abduct their shoulder (Brunnstrom, 1966; Twitchell, 1951). Yet, Experiment 2 found no relationship between resting postural force biases and active movement control. To further investigate this apparent paradox, we examined the relationship between resting postural force biases and abnormal synergies in further detail. To assess the level of synergy abnormalities for each patient, we measured their Fugl-Meyer scores for the upper extremity (FM-UE), a scale which was designed to capture abnormal muscle synergy after stroke (Brunnstrom, 1966; Fugl-Meyer et al., 1975) and closely corresponds to EMG-based synergy measures (Bourbonnais et al., 1989).

We entered patients' resting postural bias magnitudes into a mixed-effects ANOVA with FM-UE (continuous), Proximity (distal: the three locations furthest from the body, proximal: the two locations closest to the body, Figure 8, top left) and Support (with air sled, without air sled) as factors, evaluating for main effects and interactions. All factors showed significant main effects, with resting postural force magnitudes decreasing with FM-UE ($R^2 = 0.27$, p 2×10^{-8}), target proximity ($\eta^2 = 0.13$, $p = 0.00003$), and weight support ($\eta^2 = 0.13$, $p = 0.00004$). Significant interactions were observed between FM-UE and both Support ($p=0.0062$) and Proximity ($p=0.0034$), with Support and Proximity becoming more important for lower FM-UE scores (i.e. higher overall motor impairment / higher synergy intrusion) as illustrated in Figure 8.

In summary, we made three key observations with regard to abnormal resting postural force biases. First, like abnormal synergies, they were exaggerated when active arm support was required. Second, they were more pronounced in more distal positions where the elbow was more extended; if patients were to actively reach to the same locations, they would have faced increased intrusion of flexor synergy. Third, they scaled with the synergy-based FM-UE. These observations suggest a common mechanism behind resting postural force biases and abnormal synergies.

Figure 8: Relationship between resting force biases and abnormal synergies. *Across-patient relationships of FM-UE (/66, higher scores indicating lower impairment) and resting postural force magnitudes, for distal (green) and proximal (blue) target positions, with (left) and without support (right). Note the strong effects of arm support, proximity, and FM-UE. Lines indicate linear fits; shading indicates 95% confidence interval for each fit.*

Discussion

We assessed abnormal resting posture in stroke patients by measuring the resting force biases they involuntarily exerted while their arm was held at different points within a planar workspace (Experiment 1). We found that these resting postural force biases were strongest in more distal positions of the arm, generally pulled the arm toward a flexed position, and were significantly reduced, but nevertheless remained present, with arm weight support. We then proceeded to assess reaching and holding control in the same workspace in the presence of arm support (Experiment 2) and examined whether resting postural forces could partially account for deficits in the motor control of reaching and holding in the same patients. Remarkably, resting postural force biases did not have a detectable effect upon any

component of active reaching, and only emerged during the control of holding after the movement ended. The results suggest a dissociation between the control of movement and posture. At the same time, assessing patients' impairment using the FM-UE, a metric designed to measure for abnormal synergies during 3D arm movements, revealed a strong association between resting force biases and abnormal movement synergies; which raises the possibility that the observed dissociation of movement and posture control for planar weight-supported movements may break down for unsupported 3D arm movements. This dissociation raises interesting questions about both the neural architecture responsible for these two forms of control, and how to approach rehabilitation of the post-stroke arm.

Dissociation between reaching and holding

Previous research provides evidence for separate control of reaching and holding in the healthy arm – for a review, see (Shadmehr, 2017) and (Jayasinghe et al., 2022). For example, following visuomotor rotation training specific to the outward phase of an out-and-back movement, participants did not transfer this rotation to holding after a point-to-point reach in the same direction: after the movement was over, and visual feedback was removed, participants' held position drifted from the rotated movement endpoint towards the baseline hold position (Scheidt and Ghez, 2007). There is also neurophysiological evidence for separate control of moving and holding. A previous study found populations of neurons in macaque M1 that represent mechanical loads during posture or movement but not both (Kurtzer et al., 2005), whereas in another study cortical neurons and spinal interneurons coded parameters related to either movement or posture maintenance (Shalit et al., 2012).

The dissociation reported here between the control of reaching and holding in the post-stroke arm is consistent with a recently proposed hybrid model, which posits distinct controllers for reaching and for bringing the arm to a stop (Jayasinghe et al., 2022). Our finding that resting posture control does not interact either with the initial reach or bringing the arm to a stop extends this idea, suggesting *three* distinct controllers: one for the initial reach; another for bringing the arm to a stop; and another for control of holding at the endpoint after movement is over. It should be noted, however, that having distinct neural circuits for reaching and holding does not rule out interactions between them. For example, we recently demonstrated how arm holding control reflects the integration of motor commands driving the preceding active movement that led to the hold position, in both healthy participants and patients with hemiparesis (Albert et al., 2020). However, in that paper, we did not claim that this integration is the only source of holding control. Indeed, in Experiment 1 of the current study, we used

668   passive movement to bring the arm to each probed position, which means that the postural
669   biases could not be the result of integration of motor commands.

Potential origins of abnormal resting flexor biases

Experiment 1 revealed several parallels between resting postural force biases and abnormal
synergies: a propensity for flexion, mitigation by arm weight support and, a significant
correlation between patients' resting postural force biases and their FM-UE scores. The FM-
UE is a measure designed to assess post-stroke abnormal synergies during active movement.
These similarities raise the possibility that post-stroke resting postural biases and movement
synergies share a generative mechanism. Consistent with this possibility, a study found that
externally-imposed elbow flexion led to (involuntary) shoulder flexion and external adduction
in stroke patients but not healthy controls, suggesting abnormal synergy patterns do not
require active voluntary movement to be expressed (Sangani et al., 2007) – also see (Schmit
and Rymer, 2001).

It has been proposed that abnormal resting posture after brain injury can be ascribed to an
extrapyramidal system (i.e., other than the corticospinal tract, CST) (Denny-Brown, 1964).
Consistent with this suggestion, multiple lines of evidence from animal models point towards
reticulospinal tract (RST) involvement in postural control (Deliagina et al., 2007; Lacquaniti et
al., 1997; Takakusaki, 2017). Lesions of the monkey ponto-medullary reticular formation
resulted in abnormal postures; notably, these were characterized by trunk and limb flexion,
elevated shoulders, and arms held close to the body (Lawrence and Kuypers, 1968) mirroring
the flexor posture pattern seen after stroke. Several studies in cats also suggest a postural
role for the reticulospinal tract. Reticular formation neurons respond to vestibular inputs such
as head tilts or whole-body tilts, in line with a role in adjusting posture against gravity (Bolton
et al., 1992; Matsuyama and Drew, 2000; Pompeiano et al., 1984). Electrical and chemical
stimulation of the reticular formation leads to modulations in muscle tone (Takakusaki et al.,
2016), and reticulospinal neurons display tonic activity patterns that are related to postural
adjustments that precede a movement rather than to the movement itself (Schepens and
Drew, 2004).

There is also recent evidence that the RST is the descending system responsible for the
generation of large forces during voluntary movement (Glover and Baker, 2022; Tapia et al.,
2022). Thus, the accumulated evidence suggests that the RST could control posture and large
force production in the upper limb. Upregulation of the RST has also been implicated in the
generation of abnormal movement synergies after stroke (McPherson et al., 2018). For

example, the ipsilateral RST can facilitate flexors but suppress extensors, a pattern mirroring

the flexor synergy (Davidson et al., 2007; Davidson and Buford, 2004).

There appears to be a contradiction, however, between the idea that abnormal postures share

a common mechanism with abnormal movement synergies, and our finding that there was no

evidence for intrusion of resting biases into any phase of active reaching. A potential

explanation for this apparent contradiction is that our experiments were conducted on a 2D

surface with weight support. In contrast, the FM-UE scale, a synergy-based measure, is

performed by patients in 3D without weight support. It has been shown that for 3D reaching,

patients with chronic stroke express intrusive flexor synergies (Zackowski et al., 2004). Weight

support – as was used in Experiment 2 – reduces intrusion of flexor synergies for planar

movements (Sukal et al., 2007) and improves planar kinematics (Beer et al., 2007, 2004).

Earlier work, using a weight-supported planar task, found similar degrees of reaching

abnormality whether movements were made in or out of synergy (Levin, 1996); more recently,

we showed that reaching dexterity can be dissociated from synergy intrusion when arm

support is provided (Hadjiosif et al., 2022). Thus, while full weight support reduces both resting

flexor biases and movement-related flexor synergies, this reduction is more complete for

synergies compared to resting biases. This is not inconsistent with positing a shared substrate

for abnormalities at rest and during movement, but this substrate might play a greater role in

holding than in moving, and so the consequences of damage to it may be more apparent in a

task that stresses holding over movement.

What would this framework look like? We posit that the motor system has separable functional

modes for moving vs. holding and that this is accomplished by differentially weighting the

contributions of descending systems that are operative in both modes (Figure 9). The CST is

weighted more towards fast and fractionated control during movement, whereas the RST, in

contrast, is weighted more towards slower postural control and generation of large isometric

forces (such as vertical forces for arm support, or horizontal forces for holding the arm still

against a background load like in our posture/release perturbation trials). In healthy

individuals, these two modes are kept in balance by the CST, which has a moderating

influence on the RST (Schepens and Drew, 2006) (Figure 9, left). In this framework, the CST

is the controller during movement and the modulator during holding.

Post-stroke damage to the CST reduces the moderating influence the CST has upon the RST

(Figure 9, right); influence that is likely further compromised by upregulation of the RST

through plasticity mechanisms (García-Alías et al., 2015; Zaaimi et al., 2012). This conceptual

model can explain our results as follows: As the CST is the dominant system during movement

it can still modulate the RST in this mode, especially with weight support, as this reduces RST

drive, with the consequence that resting biases do not markedly contaminate active
movement. Conversely, the RST is the dominant system for postural control, and can
overcome weakened

Figure 9: An architecture for the separable control of reaching and holding and spillover effects
in stroke.

CST modulation, which leads to the resting biases we observed. Thus, weight support allows
a weakened CST to keep moving protected from holding, but it cannot prevent abnormal
holding itself. The interesting implication of this conceptual model is that synergies are in fact
postural abnormalities that spill over into active movement when the CST can no longer
modulate the increased RST activation that occurs when weight support is removed (i.e.
resting biases may influence active reaching in absence of weight support). Supporting this
idea, a study found increased ipsilateral activity (which primarily represents activation via the
descending ipsilateral RST (Zaaimi et al., 2012)) when the paretic arm had reduced support
compared to full support (McPherson et al., 2018).

A side question is how the control of decelerating to a stop – which we show here is distinct
from the control of holding after the movement is over – fits within such a scheme. A recent
saccade study found that deceleration may be controlled through the cerebellum (Sedaghat-
Nejad et al., 2022). In reaching tasks in the mouse, activity in the interpositus nucleus scaled
with limb deceleration (Becker and Person, 2019) whereas disruption of the pontine nuclei did
not impair movement initiation as much as kinematic variables related to bringing the
movement to a stop (Guo et al., 2021).

Increased recruitment of the RST has also been implicated in power grip (Baker and Perez,
2017; Tazoe and Perez, 2017). This may be relevant to our study, as our task had participants
actively grasp the handle of the robot for both Experiments 1 and 2. If increased RST
recruitment indeed explains abnormal resting postural biases as we discussed in the previous
section, a stronger grasp would in turn increase the strength of these biases even further.
Moreover, the intermixing of free-reaching and perturbation trials in Experiment 2 could have
led to increased uncertainty in environmental dynamics; uncertainty can lead to adjustments
such as even stronger grip (Hadjiosif and Smith, 2015). Yet, despite this potential additional
RST recruitment, we found that resting biases did not affect active reaching and only switched
on after the reach was over; detected when we a applied a perturbing hand force. This result
lends further support for separate controllers for reaching vs. holding.

Another potential source for the resting postural force biases we observe could be abnormally
low thresholds (Levin and Feldman, 1994; Powers et al., 1989) or abnormally high gains
(Thilmann et al., 1991) of the stretch reflex. Passively extending the elbow, even at low speeds
– something Experiment 1 did – can lead to increased muscle activity which may persist after
the end of movement (Kanade-Mehta et al., 2023; Levin et al., 2000). This velocity-dependent
increase in muscle tone – spasticity – could potentially explain some of the resting force biases
measured in Experiment 1. However, in an earlier task where the elbow was passively
extended at different velocities, the resulting tonic biceps EMG terminated around the time
that the applied elbow extension was completed (Thilmann et al., 1991). And, in recent work,
(Kanade-Mehta et al., 2023) used a paradigm similar to Experiment 1 and found that, while
resting force biases may initially reflect the velocity at which the arm (passively) approached
each test position, this velocity dependence dissipated 2s after the end of passive movement.
In our study, resting biases were measured beginning 2s *after* the end of movement, which
makes a central rather than a reflex mechanism a more likely culprit.

Conclusions

Our examination of the interplay between abnormalities in moving and holding in patients making planar reaching movements suggests the existence of two functional modes of control, likely constructed out of differing configurations of the CST and the RST. The components of the paretic syndrome – loss of dexterity, weakness, abnormal resting posture, and intrusive synergies provide insight into how a normal movement is assembled by descending systems. To the degree that hemiparesis is a consequence of the CST losing, to varying degrees, both its direct control over motor neurons and its modulatory influence on the RST, then physiological and behavioral interventions that augment the residual CST may have a plurality of benefits. In support of this, in a recent study of epidural stimulation of the cervical spinal cord in two chronic stroke patients (the rationale behind the stimulation being to amplify residual CST commands) multiple hemiparetic components –strength, dexterity, synergy abnormalities – improved simultaneously (Powell et al., 2023).

Acknowledgements

We would like to thank Stuart Baker and Maurice Smith for helpful discussions about this manuscript.

Funding

AMH and KK were supported by the Sheikh Khalifa Stroke Institute.

Competing interests

The authors report no competing interests.

ID	Age (5y range)	Sex	Time since stroke	Handed-ness	Paretic arm	FM-UE (/66)	ARAT (/57)
S001	76-80	M	2y	Right	Left	57.5	57
S002	51-55	M	6y	Right	Left	40	47.5
S003	66-70	F	7y	Right	Right	34.5	19
S004	26-30	F	5y	Right	Left	55.5	43.5
S005	76-80	M	13mo	Right	Right	43.5	34
S007	51-55	F	2mo	Left	Right	63	57
S008	51-55	F	14mo	Right	Left	41	25
S009	56-60	F	5y	Right	Left	22	3
S010	66-70	M	5y	Right	Left	20	12
S011	41-45	F	20mo	Right	Right	64	57
S012	46-50	M	6y	Right	Left	18.5	6.5
S013	66-70	M	9y	Right	Left	14	8
S014	41-45	F	16mo	Right	Left	40	39.5
S015	61-65	F	10y	Right	Left	22	4.5
S016	36-40	F	21mo	Amb.	Right	62.5	57
S017	46-50	M	3mo	Right	Left	15	3

804 **Table 1: Patient characteristics.** *FM-UE: Fugl-Meyer Assessment for the Upper Extremity; ARAT:*
805 *Action Research Arm Test.*

	Stroke patients	Controls
N	16	9
Age	58.5±17.8	62.6±15.2
Gender	7M/9F	3M/6F
Paretic side	11L/5R	n/a
FM-UE	38.3±18.2	66.0±0.0
ARAT	29.6±21.8	57.0±0.0
MoCA	24.9±3.1	28.1±1.6
Time since stroke	[2mo,10y]	n/a

807 **Table 2: Summary of patient and control characteristics.** *FM-UE: Fugl-Meyer Assessment for the*

808 *Upper Extremity (/66); ARAT: Action Research Arm Test (/57). MoCA: Montreal Cognitive Assessment*

809 *(/30).). Here, ± indicates standard deviation.*

References

Albert ST, Hadjiosif AM, Jang J, Zimnik AJ, Soteropoulos DS, Baker SN, Churchland MM,
Krakauer JW, Shadmehr R. 2020. Postural control of arm and fingers through integration
of movement commands. *Elife* **9**:e52507.

Baker SN, Perez MA. 2017. Reticulospinal contributions to gross hand function after human
spinal cord injury. *Journal of Neuroscience* **37**:9778–9784.

Becker MI, Person AL. 2019. Cerebellar control of reach kinematics for endpoint precision.
Neuron **103**:335–348.

Beer RF, Dewald JP, Dawson ML, Rymer WZ. 2004. Target-dependent differences between
free and constrained arm movements in chronic hemiparesis. *Experimental Brain*
Research **156**:458–470.

Beer RF, Ellis MD, Holubar BG, Dewald JP. 2007. Impact of gravity loading on post-stroke
reaching and its relationship to weakness. *Muscle & Nerve: Official Journal of the*
American Association of Electrodiagnostic Medicine **36**:242–250.

Berens P. 2009. CircStat: a MATLAB toolbox for circular statistics. *Journal of statistical*
software **31**:1–21.

Bobath B. 1982. Adult Hemiplegia: Evaluation and Treatment, 2nd edn. London: Spottiswoode
Ballantyne. *Physiotherapy in Disorders of the Brain*.

Bolton P, Goto T, Schor R, Wilson V, Yamagata Y, Yates B. 1992. Response of
pontomedullary reticulospinal neurons to vestibular stimuli in vertical planes. Role in
vertical vestibulospinal reflexes of the decerebrate cat. *Journal of neurophysiology*
67:639–647.

Bourbonnais D, VANDEN NOVEN S, Carey KM, Rymer WZ. 1989. Abnormal spatial patterns
of elbow muscle activation in hemiparetic human subjects. *Brain* **112**:85–102.

Brunnstrom S. 1966. Motor testing procedures in hemiplegia: based on sequential recovery
stages. *Physical therapy* **46**:357–375.

Carr E, Kenney F. 1994. Observing seated posture after stroke: a reliability study. *Clinical*
rehabilitation **8**:329–333.

Carr E, Kenney F. 1992. Positioning of the stroke patient: a review of the literature.
International journal of nursing studies **29**:355–369.

Davidson AG, Buford JA. 2004. Motor outputs from the primate reticular formation to shoulder
muscles as revealed by stimulus-triggered averaging. *Journal of neurophysiology* **92**:83–
95.

Davidson AG, Schieber MH, Buford JA. 2007. Bilateral spike-triggered average effects in arm
and shoulder muscles from the monkey pontomedullary reticular formation. *Journal of*
Neuroscience **27**:8053–8058.

Deliagina TG, Zelenin PV, Beloozerova IN, Orlovsky GN. 2007. Nervous mechanisms
controlling body posture. *Physiology & behavior* **92**:148–154.

Denny-Brown D. 1964. The extrapyramidal system and postural mechanisms. *Clinical*
Pharmacology & Therapeutics **5**:812–827.

Fine MS, Thoroughman KA. 2006. Motor Adaptation to Single Force Pulses: Sensitive to
Direction but Insensitive to Within-Movement Pulse Placement and Magnitude. *Journal of*
Neurophysiology **96**:710–720. doi:10.1152/jn.00215.2006

Fugl-Meyer AR, Jääskö L, Leyman I, Olsson S, Steglind S. 1975. The post-stroke hemiplegic
patient. 1. a method for evaluation of physical performance. *Scandinavian journal of*
rehabilitation medicine **7**:13–31.

García-Alías G, Truong K, Shah PK, Roy RR, Edgerton VR. 2015. Plasticity of subcortical
pathways promote recovery of skilled hand function in rats after corticospinal and
rubrospinal tract injuries. *Experimental Neurology* **266**:112–119.
doi:10.1016/j.expneurol.2015.01.009

Garland SJ, Gray VL, Knorr S. 2009. Muscle activation patterns and postural control following
stroke. *Motor Control* **13**:387–411.

Ghez C, Scheidt R, Heijink H. 2007. Different Learned Coordinate Frames for Planning
Trajectories and Final Positions in Reaching. *Journal of Neurophysiology* **98**:3614–3626.
doi:10.1152/jn.00652.2007

Glover IS, Baker SN. 2022. Both Corticospinal and Reticulospinal Tracts Control Force of
Contraction. *J Neurosci* **42**:3150. doi:10.1523/JNEUROSCI.0627-21.2022

Guo J-Z, Sauerbrei BA, Cohen JD, Mischiati M, Graves AR, Pisanello F, Branson KM,
Hantman AW. 2021. Disrupting cortico-cerebellar communication impairs dexterity. *Elife*
10:e65906.

Hadjiosif AM, Branscheidt M, Anaya MA, Runnalls KD, Keller J, Bastian AJ, Celnik PA,
Krakauer JW. 2022. Dissociation between abnormal motor synergies and impaired
reaching dexterity after stroke. *Journal of neurophysiology* **127**:856–868.

Hadjiosif AM, Smith MA. 2015. Flexible Control of Safety Margins for Action Based on
Environmental Variability. *J Neurosci* **35**:9106–9121. doi:10.1523/JNEUROSCI.1883-
14.2015

Hannaford B, Stark L. 1985. Roles of the elements of the triphasic control signal. *Experimental*
Neurology **90**:619–634. doi:10.1016/0014-4886(85)90160-8

Jayasinghe SA, Scheidt RA, Sainburg RL. 2022. Neural Control of Stopping and Stabilizing
the Arm. *Frontiers in Integrative Neuroscience* **16**.

Kanade-Mehta P, Bengtson M, Stoeckmann T, McGuire J, Ghez C, Scheidt RA. 2023. Spatial
mapping of posture-dependent resistance to passive displacement of the hypertonic arm
post-stroke. *Journal of NeuroEngineering and Rehabilitation* **20**:163.

Karst G, Hasan Z. 1991. Timing and magnitude of electromyographic activity for two-joint arm
movements in different directions. *Journal of Neurophysiology* **66**:1594–1604.

Kurtzer I, Herter TM, Scott SH. 2005. Random change in cortical load representation suggests
distinct control of posture and movement. *Nature neuroscience* **8**:498–504.

Lacquaniti F, Maioli C, Borghese N, Bianchi L. 1997. Posture and movement: coordination
and control. *Archives italiennes de biologie* **135**:353–367.

Laczko J, Scheidt RA, Simo LS, Piovesan D. 2017. Inter-joint coordination deficits revealed in
the decomposition of endpoint jerk during goal-directed arm movement after stroke. *IEEE*
Transactions on Neural Systems and Rehabilitation Engineering **25**:798–810.

Lawrence DG, Kuypers HG. 1968. The functional organization of the motor system in the
monkey: II. The effects of lesions of the descending brain-stem pathways. *Brain* **91**:15–
36.

Lestienne F. 1979. Effects of inertial load and velocity on the braking process of voluntary limb
movements. *Experimental Brain Research* **35**:407–418.

Levin MF. 1996. Interjoint coordination during pointing movements is disrupted in spastic
hemiparesis. *Brain* **119**:281–293.

Levin MF, Feldman AG. 1994. The role of stretch reflex threshold regulation in normal and
impaired motor control. *Brain research* **657**:23–30.

Levin MF, Selles RW, Verheul MH, Meijer OG. 2000. Deficits in the coordination of agonist
and antagonist muscles in stroke patients: implications for normal motor control. *Brain*
research **853**:352–369.

Lowrey CR, Bourke TC, Bagg SD, Dukelow SP, Scott SH. 2019. A postural unloading task to
assess fast corrective responses in the upper limb following stroke. *Journal of*
NeuroEngineering and Rehabilitation **16**:1–17.

Mani S, Mutha PK, Przybyla A, Haaland KY, Good DC, Sainburg RL. 2013. Contralesional
motor deficits after unilateral stroke reflect hemisphere-specific control mechanisms. *Brain*
136:1288–1303.

Matsuyama K, Drew T. 2000. Vestibulospinal and reticulospinal neuronal activity during
locomotion in the intact cat. II. Walking on an inclined plane. *Journal of neurophysiology*
84:2257–2276.

McPherson JG, Chen A, Ellis MD, Yao J, Heckman C, Dewald JP. 2018. Progressive
recruitment of contralesional cortico-reticulospinal pathways drives motor impairment post
stroke. *The Journal of physiology* **596**:1211–1225.

Pompeiano O, Manzoni D, Srivastava U, Stampacchia G. 1984. Convergence and interaction
of neck and macular vestibular inputs on reticulospinal neurons. *Neuroscience* **12**:111–
128.

Powell MP, Verma N, Sorensen E, Carranza E, Boos A, Fields DP, Roy S, Ensel S, Barra B,
Balzer J. 2023. Epidural stimulation of the cervical spinal cord for post-stroke upper-limb
paresis. *Nature medicine* **29**:689–699.

Powers R, Campbell D, Rymer W. 1989. Stretch reflex dynamics in spastic elbow flexor
muscles. *Annals of neurology* **25**:32–42.

Robinson D. 1970. Oculomotor unit behavior in the monkey. *Journal of neurophysiology*
33:393–403.

Sainburg R, Ghez C, Kalakanis D. 1999. Intersegmental dynamics are controlled by sequential
anticipatory, error correction, and postural mechanisms. *Journal of neurophysiology*
81:1045–1056.

Sangani SG, Starsky AJ, Mcguire JR, Schmit BD. 2007. Multijoint reflexes of the stroke arm:
neural coupling of the elbow and shoulder. *Muscle & nerve* **36**:694–703.

Schaefer SY, Haaland KY, Sainburg RL. 2009. Hemispheric specialization and functional
impact of ipsilesional deficits in movement coordination and accuracy. *Neuropsychologia*
47:2953–2966.

Scheidt RA, Ghez C. 2007. Separate adaptive mechanisms for controlling trajectory and final
position in reaching. *Journal of neurophysiology* **98**:3600–3613.

Schepens B, Drew T. 2006. Descending signals from the pontomedullary reticular formation
are bilateral, asymmetric, and gated during reaching movements in the cat. *Journal of*
neurophysiology **96**:2229–2252.

Schepens B, Drew T. 2004. Independent and convergent signals from the pontomedullary
reticular formation contribute to the control of posture and movement during reaching in
the cat. *Journal of neurophysiology* **92**:2217–2238.

Schmit BD, Rymer WZ. 2001. Identification of static and dynamic components of reflex
sensitivity in spastic elbow flexors using a muscle activation model. *Annals of biomedical*
engineering **29**:330–339.

Sedaghat-Nejad E, Pi JS, Hage P, Fakharian MA, Shadmehr R. 2022. Synchronous spiking
of cerebellar Purkinje cells during control of movements. *Proceedings of the National*
Academy of Sciences **119**:e2118954119.

Shadmehr R. 2017. Distinct neural circuits for control of movement vs. holding still. *Journal of*
neurophysiology **117**:1431–1460.

Shalit U, Zinger N, Joshua M, Prut Y. 2012. Descending systems translate transient cortical
commands into a sustained muscle activation signal. *Cerebral cortex* **22**:1904–1914.

Simo LS, Piovesan, D., Botzer, L., Bengston M, Ghez, C. P., Scheidt, R.A. 2013. Arm
kinematics during blind and visually guided movements in hemiparetic stroke
survivorsNeuroscience Meeting Planner. Presented at the Society for Neuroscience. San
Diego, CA: Society for Neuroscience. p. Program No. 652.06/VV4 2013.

Smith MA, Shadmehr R. 2005. Intact ability to learn internal models of arm dynamics in
Huntington's disease but not cerebellar degeneration. *Journal of neurophysiology*
93:2809–2821.

Sukal TM, Ellis MD, Dewald JP. 2007. Shoulder abduction-induced reductions in reaching
work area following hemiparetic stroke: neuroscientific implications. *Experimental brain*
research **183**:215–223.

Takakusaki K. 2017. Functional neuroanatomy for posture and gait control. *Journal of*
movement disorders **10**:1.

Takakusaki K, Chiba R, Nozu T, Okumura T. 2016. Brainstem control of locomotion and
muscle tone with special reference to the role of the mesopontine tegmentum and
medullary reticulospinal systems. *Journal of neural transmission* **123**:695–729.

Tapia JA, Tohyama T, Poll A, Baker SN. 2022. The Existence of the StartReact Effect Implies
Reticulospinal, Not Corticospinal, Inputs Dominate Drive to Motoneurons during Voluntary
Movement. *J Neurosci* **42**:7634. doi:10.1523/JNEUROSCI.2473-21.2022

Tazoe T, Perez MA. 2017. Cortical and reticular contributions to human precision and power
grip. *The Journal of physiology* **595**:2715–2730.

Thilmann A, Fellows S, Garms E. 1991. The mechanism of spastic muscle hypertonus:
variation in reflex gain over the time course of spasticity. *Brain* **114**:233–244.

Trombly CA. 1992. Deficits of reaching in subjects with left hemiparesis: a pilot study. *The*
American Journal of Occupational Therapy **46**:887–897.

Twitchell TE. 1951. The restoration of motor function following hemiplegia in man. *Brain*
74:443–480.

Zaaimi B, Edgley SA, Soteropoulos DS, Baker SN. 2012. Changes in descending motor
pathway connectivity after corticospinal tract lesion in macaque monkey. *Brain* **135**:2277–
2289.

Zackowski KM, Dromerick A, Sahrmann S, Thach W, Bastian AJ. 2004. How do strength,
sensation, spasticity and joint individuation relate to the reaching deficits of people with
chronic hemiparesis? *Brain* **127**:1035–1046.