scRNAseq of healthy and irradiated mouse parotid glands highlights crosstalk between immune and secretory cells during chronic injury

Brenna A. Rheinheimer[2], Mary C. Pasquale[1], NIDCD/NIDCR Genomics and Computational Biology Core[3], Kirsten H. Limesand[2], Matthew P. Hoffman[1], Alejandro M Chibly[1*]

[1]Matrix and Morphogenesis Section, National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, MD 20892.

[2]Nutritional Sciences Department, University of Arizona, Tucson, AZ. 85721

[3]Genomics and Computational Biology Core, National Institute on Deafness and Other Communication Disorders, 35A Convent Drive, Room 1F-103, Bethesda, Maryland 20892.

Running title. Single cell RNAseq analysis of parotid glands
Key words. scRNAseq, salivary gland, parotid, PG, irradiation

*Corresponding author:
Alejandro M. Chibly
Matrix and Morphogenesis Section, NIDCR, NIH, Bethesda, MD 20892, USA
Email: martinez-chibly.agustin@gene.com
Current address: Reverse Translation, Oncology Bioinformatics, Genentech, Inc. 1 DNA Way, South San Francisco, CA. 94080

HIGHLIGHTS

- We generated a scRNAseq dataset of chronic irradiation injury in parotid glands

- A newly identified *Etv1+* epithelial population may be acinar precursors

- Ntrk2 and Erbb3 are highly specific Etv1+ cell receptors that may mediate cell-cell communication with myoepithelial cells

- CD8+ T-cells and secretory acinar cells have the greatest transcriptional changes post-IR

SUMMARY

Translational frameworks to understand the chronic loss of salivary dysfunction that follows after clinical irradiation, and the development of regenerative therapies remain an unmet clinical need. Understanding the transcriptional landscape long after irradiation treatment that results in chronic salivary hypofunction will help identify injury mechanisms and develop regenerative therapies to address this need. Advances in single cell (sc)RNAseq have made it possible to identify previously uncharacterized cell types within tissues and to uncover gene regulatory networks that mediate cell-cell communication and drive specific cell states. scRNAseq studies have been performed for virtually all major tissues including salivary glands; however, there are currently no scRNAseq studies evaluating the long-term chronic effects of irradiation on salivary glands. Here, we present scRNAseq from control and irradiated murine parotid glands collected 10 months post-irradiation. We identify a population of epithelial cells in the gland defined by expression of *Etv1*, which may be an acinar cell precursor. These Etv1+ cells also express *Ntrk2* and *Erbb3* and thus may respond to myoepithelial cell-derived growth factor ligands. Furthermore, our data suggests that CD4+CD8+ T-cells and secretory cells are the most transcriptionally affected during chronic injury with radiation, suggesting active immune

involvement during chronic injury post-irradiation. Thus, our study provides a resource to

understand the transcriptional landscape in a chronic post-irradiation microenvironment and

identifies cell-specific pathways that may be targeted to repair chronic damage.

Introduction

Of the three major pairs of salivary glands (SGs): the parotid (PG), submandibular

(SMG), and sublingual (SLG), the human PG is the largest and produces the largest volume of

saliva, particularly in response to gustatory simulation. In mice, the PG is smaller than the SMG

but also contributes to the majority of stimulated saliva [1]. In addition, the PG is the most

sensitive to irradiation (IR) damage, a therapeutic treatment for head and neck cancer that often

results in permanent salivary hypofunction. In terms of understanding salivary gland biology,

most studies have focused on the SMG both in the context of development and response to

injury; however, each gland has unique functions and transcriptional profile [2]. Here we set out to

investigate the effects of irradiation damage to PGs in mice using single cell (sc)RNAseq.

The PG is primarily comprised of serous acinar cells which produce large volumes of

watery serous saliva that is transported through the ductal system into the oral cavity to aid in

digestion and protection of mucosal surfaces. Despite advances in radiotherapy, it is estimated

that ~40% of head and neck cancer patients suffer from the chronic consequences of salivary

gland damage months to years after the completion of radiotherapy, even with newer modalities

such as intensity-modulated radiation treatment (IMRT) that reduces exposure to non-tumor

tissues [3-7]. Animal studies show that the acute effects of radiotherapy in the PG occur in the days

and weeks following initial treatment and are likely a result of high levels of acinar cell death [8],

DNA damage [9], dysregulated calcium signaling and ROS generation [10], inflammatory responses

[11], and alterations to the nerves and vasculature, whereas the chronic effects arise months to

years after initial treatment (Reviewed in [12]). Chronic loss of function is often attributed to

fibrosis and the inability of acinar regeneration to occur, and preclinical studies suggest that

persistent acinar cell proliferation, vascular damage, and parenchymal cell loss may be

contributing factors [13-16]. In a similar manner, patients with Sjogren's syndrome, an autoimmune

disease that damages the acinar cells of salivary and lacrimal glands, life-long consequences

include dental caries, reduced taste and smell, malnutrition, mucositis, and increased risk for oral

infections leading to a significant decrease in quality of life [17]. Therefore, translational

frameworks to understand chronic glandular dysfunction following IR therapy along with the

development of regenerative therapies remains an unmet need.

The development of scRNAseq has made it possible to identify previously

uncharacterized cell types within a tissue and to uncover and gene regulatory networks and

mechanisms regulating cell-cell communication and specific cell states [18-21]. To date, there have

been scRNAseq studies performed for virtually all major tissues, including atlas-level scRNAseq

datasets such as the Tabula Muris [22] or the Tabula Sapiens [23] which integrate data from multiple

organs in mouse and human, respectively. There are also numerous scRNAseq studies on

disease-specific models, which are important to understand the cellular mechanisms involved

that could be targeted for repair or regeneration. In SGs, scRNAseq studies have focused on

understanding homeostasis and development [24-28], as well as particular disease states, such cancer

[29], Sjogren syndrome [30,31], and COVID-19 infection [25].

In this study, we use scRNAseq to characterize the adult mouse PG and compare the

transcriptional landscape 10 months after IR damage to explore chronic dysfunction post-

irradiation. Due to the complex heterogeneity of the SGs, distinguishing cell-type compositional

differences and their specific and direct contribution to the loss of saliva following radiation

therapy is complex, and single-cell transcriptomics will begin to resolve this issue.

This dataset allows for discovery and exploratory research into the mechanisms and

cellular processes driving PG dysfunction post-IR in a model of fractionated IR with limited

acinar cell loss. Our work has been validated by immunofluorescence staining to confirm the

presence of selected markers in specific cell populations, confirming the potential to reveal

meaningful biological insights. It is noteworthy that scRNAseq of in vivo models of chronic IR

injury has only been performed in liver [32], lung [33], and skin [34], and data is only publicly available

for lung and skin. Thus, our study will also be an essential resource to better understand cell-

specific responses to IR in general.

Results

Generation of a single-cell resource of healthy and irradiated mouse parotid gland

Using the 10X Genomics platform, we generated 2 individual scRNAseq libraries of

healthy and IR mouse PG collected 10-months post-irradiation (Figure 1A). Mice received 6 Gy

IR/day to the head and neck region on five consecutive days, for a total dose of 30 Gy. This

mouse model of IR damage to SGs results in chronic loss of saliva with partial loss of epithelial

cells [35]. Control and IR PG samples were bioinformatically integrated with SEURAT v4 and

clustered following SEURAT's standard workflow [36,37]. The optimal resolution for clustering

was determined using clustree package [38] and the resulting 17 cell clusters were annotated based

on their gene expression profile (Figure 1B, S1A-B) and a previously generated atlas of SMG

development which provided cell type specific markers [26]. Stromal and myoepithelial cells

(MECs) clustered together with endothelial cells likely due to the low number of cells recovered

for these populations. Thus, they were manually annotated based on expression of a combination

of stromal (*Col1a12* and *Vim*) and myoepithelial (*Krt14* and *Acta2*) markers which were highly

specific (Figure S1C-D). We did not identify discrete clusters of basal duct cells (*Krt14+Krt5+*)

or peripheral nerves presumably due to limitations in the dissociation technique, which has been

previously reported for adult SG tissue dissociation [26].

The identified populations included acinar cells (*Amy1+*), intercalated duct (*Dcpp1-3+*),

striated duct (*Fxyd2+, Klk1+*), MECs (*Acta2+Krt14+*), stromal (*Col1a1+Vim+*), endothelial

(*Pecam1+*), and 9 distinct immune populations including B-cells (*Cd79a+* and Immunoglobulin

genes), five subtypes of T-cells (*CD4+; CD8+; CD4+CD8+; FoxP3+; Cxcr6+*), macrophages

(*Adgre1+*), dendritic cells (*S100a8/9+*), and natural killer cells (*Gzma+Nkg7+*). We also

identified a previously uncharacterized epithelial population defined by high expression of *Etv1*

and *Krt8* and moderate expression of *Amy1* (Figure 1B-C, S1B).

Etv1 expression delineates a secretory subpopulation in acinar and duct compartments

The two most striking observations from our initial clustering analysis are the

identification of an *Etv1+* epithelial population and the prominence of multiple resident immune

cell types after IR. *Etv1* has been described as one of the top transcription factors expressed in

the salivary glands [39]. To characterize this *Etv1+* cluster, and to generate gene expression

profiles of individual cell populations in healthy adult parotid glands, we performed differential

expression analysis with SEURAT in the annotated control sample (Figure 1C). Genes enriched

in a given cluster are herein referred to as cell-defining genes and were sometimes expressed

elsewhere at lower levels. The complete gene list is included in Supplementary File 1.

The expression of Amy1 in *Etv1+* cells suggested an acinar-like phenotype. When

comparing the gene expression profile of major epithelial populations, 38% of acinar-defining

genes (30 of 79) were enriched in *Etv1+* cells (Figure 2A-B). Both cell types expressed serous

secretory markers such as amylase (*Amy1*), parotid secretory protein (*Bpifa2*), prolactin induced

protein (*Pip*), and carbonic anhydrase 6 (*Car6*), but their expression was significantly higher in

acinar cells, while *Etv1+* cells had higher expression of *Krt8, Krt18,* and *Phlda1* (Figure 2C).

When compared to duct populations, *Etv1+* cells expressed 38% (19 genes) of intercalated duct

(ID)-defining genes (Figure S2A) and only 9.3% of striated duct (SD)-defining genes (Figure

2B, S2B), suggesting that *Etv1+* cells are transcriptionally similar to both acinar and ID

populations. Accordingly, Etv1 protein was detected by immunofluorescence in a subset of duct

and acinar cells. Duct cells showed strong nuclear and cytoplasmic Etv1+ signal while it was

predominantly nuclear in NKCC1+ acinar cells (Figure 2D).

Next, we performed functional analysis of all acinar and *Etv1+* cell-defining genes using

STITCH (search tool for interactions of chemicals, http://stitch.embl.de/), which integrates

information about interactions from metabolic and KEGG pathways, crystal structures, binding

experiments, and drug-target relationships. [40]. As expected, KEGG pathway analysis on acinar

genes showed salivary secretion as one of the top pathways (Figure S2D). In contrast, in *Etv1+*

cells the top functions and pathways were associated with organ development and activation of

Rap1, TNF, and ErbB signaling pathways (Figure 2E, S2C), suggesting that the *Etv1+*

population has distinct functions despite their transcriptional similarities to acinar cells.

Acinar and Etv1+ cells communicate with MECs and stromal cells via Erbb3 and Ntrk2

Given that cellular functions are often initiated by ligand-receptor interactions that trigger

signaling cascades, we next used two bioinformatic approaches to predict putative cell-cell

interactions: First, we used differential expression analysis for each cluster and cross-referenced

the resulting cell-defining genes with a previously published database of curated ligand-receptor

pairs [41]. For reproducibility of this approach, we used R-scripted code which is available as

supplementary material. In this database, a ligand is defined as any molecule that interacts with

known receptors and thus intracellular components such as *Hras* are included. As a

complementary approach, we used CellChat, which infers ligand-receptor pairs and associated

pathways based on a manually-curated list of literature-supported interactions grouped into 229

signaling pathways [42]. Both approaches were consistent and showed that acinar and duct cells

had the lowest number of enriched ligand and receptor genes compared to all other cell types

while MECs had the highest number across epithelial populations (Figure S3A-C).

Differential expression analysis identified 9 ligand and 5 receptor genes among the *Etv1+*

cell-defining genes, as well as 5 ligands and 2 receptors in acinar cells (Figure 3A). The

identified receptor genes enriched in *Etv1+* cells included *Ghr*, *Dddr1*, *St14*, *Erbb3,* and *Epha5*,

which were highly specific to this population (Figure 3B, left panel). On the other hand, the

ligands found in *Etv1+* cells were also enriched in other cell types, with the exception of *Col7a1,*

which was highly specific (Figure 3B, right panel). A distinct set of ligands and receptors were

enriched in acinar cells, including the receptor genes *Ntrk2* and *Kcnn4*, and the ligands *P4hb,*

Nucb2, Agt, Tcn2, and *Pip.* All of the resulting putative interactions from our differential gene

expression analysis are shown in Supplementary File 2. Interactions between acinar and *Etv1+*

cells with all other cell types are summarized as chord plots in Supplementary Figure S3D-E. All

interactions predicted by CellChat are available in Supplementary File 3. Based simply on the

total number of possible pairs (without accounting for the level of expression of individual

genes), the strongest outgoing interactions from *Etv1+* cell ligands were predicted to occur with

receptors in endothelial cells, whereas *Etv1+* cell receptors corresponded to ligands from

myoepithelial and stromal cells. In contrast, the corresponding pairs for acinar cell ligands were

expressed primarily in T-cells (Supplementary Figure S3D). These findings were largely

corroborated by CellChat (Figue 3D), which does take into account the level of gene expression

as well as the proportion of cells expressing a given ligand-receptor pair in a cluster.

CellChat analysis determined that Etv1+ cells were a source of ligands for

Thrombospondin (THBS) and EphrinA (EPHA) signaling pathways, and had receptors for

Carcinoembryonic antigen cell adhesion molecule (CEACAM) and Neuregulin (NRG) ligands,

whereas acinar cells were receptive to Neurotrophin (NT) signaling (Figure 3E). These results

revealed notable interactions between myoepithelial and Etv1+ cells via the *Erbb3* receptor and

two of its ligands, Neuregulin1 (*Nrg1*) and *Nrg2*, and between myoepithelial and acinar cells via

the neurotrophin receptor *Ntrk2* and its ligands, Neurotrophin 3 (*Ntf3*) and *5* (*Ntf5*) (Figure 3E-F,

Supplementary Figure S3D-E). *Ntrk2* was also expressed in *Etv1+*, myoepithelial and stromal

cells in our scRNAseq data but immunofluorescence staining confirmed enrichment of the

receptor in acinar cells of mouse parotid gland (Figure 3F). The cellular functions of *Ntrk2* in

acinar cells are currently unknown and thus further mechanistic studies are warranted.

CD8+CD4+ T-cells and acinar cells have the greatest transcriptional response to IR

The model of SG IR used in this study is based on a fractionated dosing schedule of 6Gy

x 5 consecutive days, which leads to significant loss of saliva [35] but it does not result in extensive

loss of acinar cells and development of fibrosis (Figure 4A) as reported by Ferreira *et al.* While

this model shows a milder phenotype compared to alternative models using a single 15Gy dose

in distinct mouse strains [10,43-45], it has been previously used to demonstrate the therapeutic

potential of adenovirus-based Neurturin gene transfer in the SMG to prevent the loss of saliva

caused by IR. Given that we did not perform multiple technical replicates of each treatment,

potential changes in cell proportions are reported as trends. In general, B cells and T cells were

the most affected (Figure 4A-B). We observed a 33 % relative decrease in the proportion of B

cells, a 39 % increase in CD4+ T cells, and a 195% increase in CD4+CD8+ T cells. A 22 %

decrease in the proportion of acinar cells was also noted.

Differential expression analysis with SEURAT was performed between control and

irradiated cell types. The complete list of differentially expressed genes (DEGs) is shown in

Supplementary file 4. CD4+CD8+ T-cells had the highest number of dysregulated genes (~70)

post-IR across all identified cell populations, followed by acinar cells (Figure 4C). We did not

detect DEGs in MEC and stromal populations post-IR, and only 1 gene was differentially

expressed in IR endothelial cells. The lack of DEGs in MECs is likely explained because of the

low number of MECs analyzed (Figure 4B). Stromal and endothelial cells also did not show

significant changes in gene expression, but they were well-represented in our dataset; thus, cell

numbers alone are not likely to account for the lack of DEGs post-IR in these populations.

Instead, the lack of DEGs may reflect the fact that this model of IR damage is not highly fibrotic

(Figure 4A). Alternatively, it's possible that endothelial and stromal populations may have

recovered a year after IR damage.

The top upregulated genes in acinar cells post-IR included *Actb, Tmsb4x*, and *Pfn1* which

are involved in actin polymerization (Figure 4D). The genes *Gm42418, Hba-a1*, and *Smr3a* were

the only downregulated genes in acinar cells and they were also downregulated in most other cell

types (Figure S4A, Supplementary file 4), suggesting a global response to IR rather than an

acinar-specific one. In CD4+CD8+ T-cells, the top upregulated genes post-IR were *Jun, Fos,*

Ltb, *Klf2*, and *Klf6*, and the most downregulated genes were *Ctla2a, Tcp11l2, Crip1, Ramp3,* and

Tubb4b (Figure 4D). In general, DEGs in acinar cells were associated with regulation of

transepithelial transport, electron transport, apoptosis, and translation processes according to

gene ontology analysis via The Gene Ontology Consortium [46], while DEGs in CD4+CD8+ T-

cells were associated with V(D)J recombination, lymphocyte differentiation, apoptosis,

axonogenesis, and ERK signaling pathway (Figure S4B).

When we cross-referenced the IR DEGs against the database of ligand-receptor pairs

from Ramilowsky *et al.*, only a handful of ligands and receptors were represented (Figure 4E-F),

and only a few of these had a corresponding pair (Figure 4G-H, Supplementary File 5). Most

differentially expressed pairs were found between immune populations, MECs, stromal, and

endothelial cells. In acinar and CD4+CD8+ T-cells, which were the most transcriptionally

affected, we identified 5 ligands (*Ptma, Hsp90aa1, Ltb, Hspa1a,* and *Hras*) and 5 receptor genes

(*Rpsa*, *Cd53, Ramp3, Cd28*, and *Ifngr1*) differentially expressed post-IR (Figure 4E-F).

However, these DEGs were expressed across multiple clusters and were not defining for any

individual population. For instance, *Hsp90aa1* was downregulated in all immune populations

except NK cells and macrophages, and both *Hspa1a* and *Hras* were downregulated in NK cells

(Figure 4E). Similarly, *Rpsa* was upregulated in acinar cells while *Ifngr1* was downregulated in

CD4+CD8+ T-cells post-IR (Figure 4F). Putative pairs were found for *Rpsa* (Ribosomal protein

SA (*Rpsa*), also known as Laminin receptor 1)*, Ifngr1* (Interferon Gamma Receptor 1)*,

Hsp90aa1* (Heatshock protein 90 Alpha Family Class A Member 1)*, Ltb* (Lymphotoxin Beta),

and *Hras* oncogene (Supplementary File 5).

This analysis suggested multiple signaling alterations including interactions with acinar

cells via *Lamb2-Rpsa* and between NK and CD8+ cells with CD4+CD8+ T-cells via *Ifng-Ifngr1*

(Supplementary File 5, Figure 4G-H). Paracrine signaling via *Hsp90aa1* from immune cells to

Egfr expressed in myoepithelial, stromal, and endothelial cells was potentially reduced, while

Ltb interaction with *Tnfrsf1a* and *Cd40* expressed by macrophages, endothelial cells, dendritic

cells, and B-cells was potentially increased.

Neurotrophin, neuregulin, ECM, and immune signaling are the main altered pathways in

Acinar and Etv1+ cells post-IR

Given that too few ligands and receptors were differentially expressed, we next used

CellChat, which is sensitive to expression patterns in ligands and receptors themselves, as well as

their cofactors, and weighs the size of a given cluster and the proportion of cells within a cluster

expressing a gene. CellChat predicted similar ligand-receptor interactions in the IR-PG compared

to those from the control glands (Supplementary File 3). There was an increase in the number of

interactions post-IR from 3128 in the control to 3191 in IR PG, although the predicted interaction

strength was slightly reduced (Figure 5A). Interactions with *Rpsa*, *Hsp90aa1*, or *Hras* could not

be confirmed since these are not included in CellChat's database of ligand-receptor pairs.

CellChat predicted that MECs, stromal, and endothelial cells had the highest number of

differential interactions while B-cells and T-cells had the greatest difference in interaction

strengths (Figure 5B, Figure S5A). A summary of the number of differential interactions per cell

type is provided in Supplementary File 6. Here, we focus specifically on acinar and Etv1+ cells.

CellChat predicted altered signaling from MECs, stromal, and endothelial cells to both

acinar and Etv1+ cells (Figure 5C-D). Generally, this seemed to be predominantly mediated by

fewer interactions between *Cd44* and *Sdc4* with multiple collagens and laminins (Figure 5E). We

also saw a shift in Neurotrophin signaling interactions post-IR, which were decreased between

acinar cells and MECs or Stromal cells, but increased with endothelial cells (Figure 5E); this

shift also involved changes in the ligand *Ntf5* and the appearance of a more significant

contribution from *Ngf-Ntrk1* interactions (Figure S5B). Disruption of neurotrophin signaling was

also recently reported in irradiated human salivary glands [47]. Altered signaling in Etv1+ cells

also involved loss of *Igf1-Igf1r* interactions with stromal and endothelial cells, and increased

interactions via *Nrg2-Erbb3* with stromal cells (Figure 5E). A key difference between the two

cell types was their differential interactions post-IR with immune populations. In acinar cells,

there were increased interactions with T-cells and B-cells (Figure 5C, Figure 5F), but the

strength of the interactions was generally lower (Figure S5C). In contrast, Etv1+ cells showed

generally stronger interactions with T-cells, with the exception of FoxP3+ cells, which had fewer

interactions (Figure 5D, Figure 5G, Figure S5C).

These results combined suggest that this model of IR injury shows alterations in cell

adhesion interactions with the extracellular matrix (i.e. via Cd44, Sdc4, collagen and laminin), as

well as changes in neurotrophin (Ntrk2, Ntf5, Ntf3), neuregulin (Nrg1, Nrg2, Erbb3), and IGF

(Igf1, Igf1r) pathways. Furthermore, they suggest active involvement of immune cells,

particularly T-cells, in mediating cellular responses post-IR, although the specific mechanisms

involved are still unclear. Further mechanistic studies are encouraged to determine the functional

relevance of these predicted interactions.

Discussion

We generated a scRNAseq resource of adult PG that includes a chronic IR injury model.

This resource allowed us to identify a discrete cell cluster of secretory cells defined by *Etv1*

expression, and to predict putative ligand-receptor interactions that mediate key signaling

pathways between secretory cell types and their microenvironment during homeostasis and post-

injury.

The near exclusivity of *Etv1* expression in a single cluster is intriguing but it is not known

whether it represents a cell-type-specific marker or a cell state. Their mixed histological

localization suggests the latter. We saw expression of *Etv1* in both acinar and duct

compartments, particularly close to the intercalated duct, similar to observations made by Song

et al. in adult mouse SMG [48]. *Etv1* is one of the top transcription factors in the salivary glands [39]

and it was previously reported to be enriched in putative salivary stem cells defined by

expression of Lin–CD24+c-Kit+Sca1+ [49]. *Etv1* was also recently associated with the

development of the acinar epithelium in the mouse SMG [26]. These observations may be

suggestive of *Etv1* being involved in an intermediate state between intercalated duct and acinar

cells. Indeed, experiments in rodents suggest that intercalated duct cells may harbor stem cells

that can differentiate into acinar cells or other duct cells [50-52]. *Etv1+* cells showed enrichment of

Erbb3 expression, which was predicted to mediate signaling via neuregulin ligands derived from

MECs and stromal cells. Erbb3 signaling is critical for SG development and plays a crucial role

in organogenesis. Branching morphogenesis in the mouse SMG depends on intraepithelial

signaling mediated by ErbB2, ErbB3, and neuregulin 1 (Nrg1) [53]. Nrg1-null embryos show

reduced innervation and defective branching morphogenesis [54,55]. Thus, it is plausible that *Etv1+*

(Erbb3+) cells in the adult parotid gland could be involved in either replenishment of the

epithelium or wound healing, and may function as a proacinar population in the PG.

Our finding that the neurotrophic receptor *Ntrk2* is enriched in acinar cells is interesting

because of the precedent of using neurotrophic factors such as neurturin to preserve function in

irradiated SGs [35,56]. Moreover, we recently reported that in humans, IR chronically dysregulates

the neurotrophin signaling pathway in both PG and SMG and is associated with functional and

morphological abnormalities [47]. Ligand-receptor analysis predicts that stromal cells and MECs

communicate with *Ntrk2*-expressing acinar cells via *Ntf5* and *Ntf3*, respectively. Considering the

localization of MECs surrounding acinar cells, it is likely that both juxtracrine and paracrine

signaling takes place. The function of the Ntrk2 receptor in salivary acinar cells is not known but

the gene is also highly expressed in Neurogenin 3-positive (Ngn3+) endocrine progenitors in the

pancreas [57] and its activation regulates Ngn3+ cell fate commitment. Neurotrophin receptors are

also mutated or upregulated in a variety of cancers, suggesting a role in proliferation and

differentiation. In the SMG, *Ntrk2* is expressed in serous acinar cells but not in seromucous

acinar cells [26], indicating that *Ntrk2* signaling may be important for the serous acinar phenotype,

which is predominant in the PG. Furthermore, we recently identified that *NTRK2* is highly

upregulated in MECs of irradiated human SGs along with other neurotrophin receptors and

stimulation of neurotrophin signaling *in vitro* promoted myoepithelial differentiation[47]. In the

lacrimal gland, neurotrophins are expressed in acini while neurotrophin receptors are expressed

by MECs [58], suggesting that neurotrophin signaling may mediate intercellular communication

between acinar cells and MECs in other exocrine tissues. Moreover, given that Ntrk2 is

expressed on the cell surface, it may also provide a viable strategy to FACS-sort acinar cells

from parotid gland to investigate expansion or differentiation of acinar cells *in vitro*. The latter

application would likely require a combination of markers since *Ntrk2* is also expressed in

Etv1+, myoepithelial and stromal cells.

The IR model used in our study has been previously used to demonstrate the protective

effect of neurturin gene-transfer to prevent loss of saliva in the IR-SMG [35]. In contrast to more

severe models of IR injury [10,43,45,59,60], this model does not result in extensive loss of acinar cells

and is only mildly fibrotic; thus, it is ideal for understanding transcriptional changes in acinar

cells before they are lost. Indeed, acinar cells had the second largest number of DEGs post-IR in

this model. Surprisingly, CD4+CD8+ cells showed the highest number of DEGs post-IR,

suggesting that chronic damage post-IR may be sustained by immunologic mechanisms. There is

growing evidence of immune-epithelial interactions in the regulation of tissue homeostasis and

wound healing responses with macrophages and regulatory T-cells (T_{regs}; FoxP3+) garnering the

most attention [61]. Through Notch-mediated signaling, mammary gland stem cells induced

resident macrophages to produce Wnt ligands ultimately leading to mammary stem cell

proliferation [62]. Depletion of T_{regs} in the intestine leads to a reduction in LGR5+ stem cells [63].

Given the extensive ligand-receptor interactions between *Etv1+* cells and immune cells, it is

interesting to speculate a functional role of *Etv1+* cells in directing the localization and activation

of resident immune populations. In the epidermis, distinct cellular populations around the hair

follicle produce distinct chemokines to direct innate immune cell populations [64]. For example,

the interaction between *Etv1+* and *FoxP3+* cells via *Cdh1-Itgae* (E-cadherin and integrin-α-E)

may represent the physical tethering of this sub-population of T-cells to the salivary epithelium

under homeostasis [65], and it was increased post-IR (Figure 5G). It's interesting to note that

radiation treatment led to a 1.5-fold increase in T_{regs} without a concomitant change in *Etv1+* cells

or macrophages. Given the extensive role macrophages and *FoxP3+* cells serve in injury and

regeneration models, more work is required to unravel the impact of these T_{regs} -epithelial

interactions population during SG dysfunction.

Radiation treatment also resulted in the greatest increase in CD4+CD8+ populations and

the most DEGs observed in the CD4+CD8+ cells (Figure 4). Clinical evaluation of SMG by

immunohistochemistry following radiotherapy has revealed increased T-cells (CD3+, CD4+ or

CD8+) in the periacinar area and B cell (CD20+) nodules in the periductal area [66]. The DEGs in

the CD4+CD8+ population suggest an imbalance in immune regulation following irradiation.

Increases in KLF2 in IR PGs may represent a shift in T-cell populations as KLF2 is highly

expressed in naïve and memory T-cells and downregulated by TCR activation and cytokine

stimulation in effector T-cells [67]. Additionally, high levels of KLF2 inhibit T-cell proliferation

and clonal expansion [67]. KLF6 also inhibits cell proliferation and is co-regulated with KLF2 in

MCF-7 cells [68]. Thus, high levels of KLF2 and KLF6 coupled with a lack of cytokines and

chemokines on the DEGs suggest that the increase in CD4+CD8+ T-cells may represent a naïve

population; however further kinetic analysis is required. This is also supported by a decrease in

Ctla2a, which encodes for a cysteine protease that serves an immunosuppressive function in

retinal pigment epithelium [69,70] and promotes the conversion of CD4+ T cells to Treg cells via

Transforming Growth Factor Beta (TGFβ) signaling [71]. Lymphotoxin-β (LT-β), encoded by *Ltb*,

is a TNF family member cytokine that has been predominantly studied in development and

organization of lymphoid tissues [72]. LT-β can mediate both regeneration and chronic tissue injury

in epithelial organs via nuclear factor-κB (NF-κB) pathway [73,74]. Blocking the LT-β receptor

suppresses immune responses by modulating trafficking mechanisms and disrupts the

progression of T1DM in NOD mice [72]. It is interesting to speculate whether the increased LT-β

interactions with *Tnfrsf1a* or *CD40* prevent the clearance of immune populations or maintenance

of naïve T cells. *Ltb* is induced following oxidative stress [75] and has been proposed to enable

communication between lymphocytes and stromal cells [73], findings that are corroborated by this

work predicting increased interactions with stromal and immune cell populations post-IR (Figure

5).

Limitations of the study

A caveat of this study is the lack of isolation of basal ducts and peripheral nerve cells during PG

dissociation, which were not represented. Similar limitations have been reported in other

scRNAseq studies working with adult tissues, which could potentially be overcome using single

nuclei RNAseq analysis. Furthermore, although multiple biological replicates were used, they

were pooled together during dissociation prior to sequencing, thus, cell proportion changes

should be considered with caution.

Acknowledgments

The authors thank the support from Dr. Daniel Martin, Dr. Robert Morell, and Dr. Erich Boger

from the Genomics and computational biology core (GCBC) at NIDCR for contributing to

library preparation and sequencing. This work used the NIDCR Veterinary Resources Core (ZIC

DE000740-05) and computational resources of the NIH HPC Biowulf cluster

(http://hpc.nih.gov). The GCBC funds were from the NIDCD Division of Intramural

Research/NIH (DC000086 to the GCBC). The study was supported by the Intramural Research

Program of the National Institute of Dental and Craniofacial Research, NIH.

Author Contributions

Conceptualization, writing and editing, A.M.C, B.R, K.H.L; Methodology, A.M.C., B.R.,

M.C.P., GCBC; Software, A.M.C, GCBC; Resources, M.P.H, K.H.L., A.M.C; Visualization,

A.M.C., B.R., M.C.P; Data curation, project administration, and supervision, A.M.C.

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Figure 1. scRNAseq analysis of control and irradiated PG

A) Single cell suspensions from 1-year-old control and irradiated PG from 2 C3H female mice were used to build scRNAseq libraries. Representative UMAP plots are colored by treatment group or cell type. Clusters were annotated based on the expression of known markers.

B) Balloon Plot with top 5 DEGs per cluster sorted by fold change. Statistical analysis performed using SEURAT package in R. Color is relative to scaled gene expression and size of the dot represents the percentage of cells expressing the gene.

C) Representative UMAP plots showing expression of Etv1 and Amy1

Figure 2. Characterization of acinar and Etv1+ cells

A) UMAP plot highlighting acinar, Etv1+, and duct populations with a representative heatmap of their gene expression profiles.

B) Venn diagram of cell-defining genes in acinar and Etv1+ clusters showing the number of unique and overlapping cell-defining genes. Representative genes from each group are shown. The bar graph shows the percentage of overlap between cell-defining genes in acinar and duct populations with Etv1+ cells.

C) Balloon plot showing expression of the 30 genes overlapping between acinar and Etv1+ cells. Genes marked with an asterisk are differentially expressed between Etv1+ and acinar cells (p<0.05, Wilcoxon rank sum test (SEURAT)). Color is relative to scaled gene expression and size of the dot represents the percentage of cells within a cluster expressing the gene.

D) Immunofluorescence staining of PG from 1 year-old C3H mice stained for Etv1 (Red), NKCC1 (green) and DAPI (blue). The area delineated by the yellow dotted line is magnified to the right for visualization. Scale bar = 50um.

E) Results from STITCH analysis showing top biological processes and KEGG pathways associated with defining-genes from Etv1+ cells.

Figure 3. Ligand-receptor analysis of Etv1+ and acinar cells

A) Bar graphs with number of identified ligands and receptors among cell-defining genes from epithelial populations.

B) Balloon plots of expression of ligands and receptors enriched in Etv1+ cells.

C) Balloon plots of expression of ligands and receptors enriched in acinar cells.

D) Chord plots summarizing putative ligand-receptor interactions with Acinar and Etv1+ cells, as predicted by CellChat. The arrows point to the cell expressing the receptors and are colored based on the source of the ligand. The thickness of the arrow is relative to the number of putative pairs identified.

E) Heatmap shows the relative importance of each cell group based on CellChat-computed network centrality measures of NRG, NT, THBS, EPHA, and CEACAM signaling networks.

F) Relative contribution of each ligand-receptor pair to the overall communication network of NRG and NT signaling pathways, which is the ratio of the total communication probability of the inferred network of each ligand-receptor pair to that of the signaling pathway.

G) Immunofluorescence staining for Smooth muscle actin (SMA, Red), NTRK2 (green) and Parotid Secretory Protein (PSP, blue). The area delineated by the yellow dotted line is magnified to the right for visualization. Scale bar = 50um.

Figure 4. Cell-specific IR-induced DEGs

A) Masson's trichrome staining of control and IR-PG collected 10-months post-IR. Areas in blue represent fibrotic/collagenous tissue.

B) Cell numbers and proportions in scRNAseq datasets from control and irradiated PG.

C) Bar graph showing number of DEGs post-IR in individual cell populations. DE analysis was performed with SEURAT's default Wilcoxon test (p<0.05).

D) Violin plots of top 5 (if present) up and downregulated genes in acinar and CD4+CD8+ T-cells. Red and blue arrows denote upregulated and downregulated genes, respectively.

E-F) Violin plots of differentially expressed ligands and receptors from SEURAT's DE analysis.

G-H) Chord plots of ligand-receptor interactions with IR-induced DE receptors and ligands.

Figure 5. Dysregulated ligand-receptor pairs post-IR

A) Number of ligand-receptor pairs and interaction strength in control and IR-PG determined by CellChat.

B) Differential number of possible interactions between acinar and Etv1+ cells with all other populations in IR-PG compared to control. Red (positive values) and blue (negative values) in

the color bar indicate higher number of predicted interactions in IR-PG and controls, respectively.

C-D) Differential ligand-receptor interactions in Acinar and Etv1+ cells between IR-PG and control, as predicted by CellChat. The arrows point to the cell expressing the receptors. Red arrows indicate increased number of interactions and blue arrows represent a decrease. The thickness of the arrow is relative to the change in number of interactions between control and IR-PG.

E) Comparison of the significant ligand-receptor pairs between control and IR-PG, which contribute to the signaling from MECs, stromal and endothelial cells to acinar and Etv1+ cells. Dot color reflects communication probabilities and dot size represents computed p-values. Empty space means the communication probability is zero. p-values are computed from one-sided permutation test.

F-G) Comparison of the significant ligand-receptor pairs between control and IR-PG, which contribute to the signaling from acinar and Etv1+ cells to immune populations. Dot color reflects communication probabilities and dot size represents computed p-values. Empty space means the communication probability is zero. p-values are computed from one-sided permutation test.

STAR METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Alejandro Chibly (martinez-chibly.agustin@gene.com)

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

The single-cell RNAseq libraries were deposited in GEO under accession number GSE223516. The code used for analysis is available in github: https://github.com/chiblya/scRNAseq_PG. All package versions are reported in the gitgub repository. Ready-to- use Seurat objects are also available via figshare: 10.6084/m9.figshare.20406219

EXPERIMENTAL MODEL AND SUBJECT DETAILS

C3H mice and irradiation (IR) treatment

C3H female mice were used for the study and were housed at the NIDCR Veterinary Resource Core in accordance with IACUC guidelines. At 6-10 weeks of age, mice received fractionated IR treatment at 5×6 Gy (6 Gy/day for 5 days). Only head and neck area was irradiated by placing each animal in a specially built Lucite jig so the animal could be immobilized without the use of anesthetics. IR treatment was delivered with by an X-Rad 320ix system. The mice were housed in a climate- and light-controlled environment, and allowed free access to food and water for 10 months post-IR before scRNA-seq analysis.

METHODS

Single-cell Dissociation. Parotid glands from 2 female mice per treatment were dissociated in a 15ml gentleMACS C tube with 5ml of digestion enzyme using the human tumor dissociation kit (#130-095-929, Miltenyi Biotech, Auburn CA) in RPMI 1640 w/L-Glutamine (Cell applications, Inc, USA). Cell dissociation was performed in a Miltenyi gentleMACS Octo Dissociator using the manufacturer's preset 37C_h_TDK_2 program. Following dissociation, 5ml of RPMI media were added to the dissociated cells and centrifuged at 1100 rpm for 10 min. Cells were resuspended in RPMI 1640 w/L-Glutamine with 5% PenStrep (Gibco, USA) and washed twice with RPMI. Cells were passed through 70 µm filters between centrifugation steps. Single-cell dissociation was confirmed by microscopic examination and cell concentration determined with a Cellometer (Nexcelom Biosciences). Cell concentration was adjusted to $5x10^5 - 1x10^6$ cells/ml prior to analysis with a 10X genomics Next GEM Chromium controller.

Library prep and sequencing: Single-cell RNA-seq library preparation was performed at the NIDCR Genomics and Computational Biology Core using a Chromium Single Cell v3 method (10X Genomics) following the manufacturer's protocol. Pooled single-cell RNA-seq libraries

were sequenced on a NextSeq500 sequencer (Illumina). Cell Ranger Single-Cell Software Suite (10X Genomics) was used for demultiplexing, barcode assignment, and unique molecular identifier (UMI) quantification using the mm10 reference genome (Genome Reference Consortium Mouse Build 38) for read alignment.

Computational analysis: Cell Ranger files were imported to SEURAT v3 using R & R Studio software and processed for clustering following their default pipeline. As a quality control measure, cells with fewer than 200 genes were not included in subsequent analyses, and those with >5% of UMIs mapping to mitochondrial genes were defined as non-viable or apoptotic and were also excluded. These metrics were based on our previous scRNAseq analysis of murine SMG [26]. Normalization and scaling were performed following SEURAT's default pipeline. Data from control and irradiated glands were bioinformatically integrated prior to assigning cell annotations. 'Clustree' package was used to determine an optimal resolution for clustering and the resulting clusters were annotated based on the expression of known cell type markers. Cell-defining genes were determined using the 'FindAllMarkers' function which uses a Wilcoxon Rank Sum statistical test for analysis. Only genes with adjusted p-values <0.05 were considered as cell-defining genes. To identify DEGs between treatments, each population was compared individually using the 'FindMarkers' function from SEURAT package.

Ligand-receptor analysis

A database of curated ligand-receptor pairs was downloaded from Ramikowski *et al.* (2015). We used scripted code in R to automate the search for ligand and receptor genes within our dataset and leverage that information against the curated database. Additionally, we used CellChat v.1.6.1 (Jin, S. *et al.,* 2021) to predict significant interactions and their associated pathways. Plots were generated using the 'circlize' and complexHeatmap packages in R. The code is available in https://github.com/chiblya/scRNAseq_PG.

Immunohistochemistry: PGs were fixed in 4% paraformaldehyde overnight at 4°C and dehydrated with 70% Ethanol prior to paraffin embedding. 5µm sections were deparaffinized with xylene substitute for 10 minutes and rehydrated with reverse ethanol gradient for 5 minutes each. Heat induced antigen retrieval was performed using a microwave maintaining sub-boiling

temperature for 10 minutes in a pH 6.0 Citrate Buffer (#21545, EDM Millipore, Darmstadt, Germany) . Sections were washed for 5 minutes with 0.1% Tween20 (Quality Biological, Inc) in PBS 1X (PBST). M.O.M.® (Mouse on Mouse) Immunodetection Kit (Vector Laboratories, Burlingame, CA) was used to block non-specific sites for 1 hour at room temperature followed by overnight incubation with primary antibodies at 4^oC. Tissue sections were washed 3 times for 5 minutes each with PBST and incubated in secondary antibodies and nuclear stain (Hoechst (Thermo Fisher Scientific, Marietta, OH)) at room temperature for 1 hour. Coverslips were mounted with Fluoro-Gel (Electron Microscopy Sciences, Hatfield, PA), and imaging was performed with a Nikon A1R confocal system.

Stitch analysis: Etv1+ cell defining genes from control parotid sample (Supplementary File 1) were directly imported into STITCH (http://stitch.embl.de/). For reproducibility, analysis was performed selecting a minimum interaction score of 0.7 and limited to less than 10 interactions.

References

1. Maruyama, C.L., Monroe, M.M., Hunt, J.P., Buchmann, L., and Baker, O.J. (2019). Comparing human and mouse salivary glands: A practice guide for salivary researchers. Oral Dis *25*, 403-415. 10.1111/odi.12840.

2. Gao, X., Oei, M.S., Ovitt, C.E., Sincan, M., and Melvin, J.E. (2018). Transcriptional profiling reveals gland-specific differential expression in the three major salivary glands of the adult mouse. Physiological Genomics *50*, 263-271. 10.1152/physiolgenomics.00124.2017.

3. Jensen, S.B., Pedersen, A.M., Vissink, A., Andersen, E., Brown, C.G., Davies, A.N., Dutilh, J., Fulton, J.S., Jankovic, L., Lopes, N.N., et al. (2010). A systematic review of salivary gland hypofunction and xerostomia induced by cancer therapies: prevalence, severity and impact on quality of life. Support Care Cancer *18*, 1039-1060. 10.1007/s00520-010-0827-8.

4. Tasaka, S., Jingu, K., Takahashi, N., Umezawa, R., Yamamoto, T., Ishikawa, Y., Takeda, K., Suzuki, Y., and Kadoya, N. (2021). The Long-Term Recovery of Parotid Glands in Nasopharyngeal Carcinoma Treated by Intensity-Modulated Radiotherapy. Front Oncol *11*, 665837. 10.3389/fonc.2021.665837.

5. Jensen, S.B., Vissink, A., Limesand, K.H., and Reyland, M.E. (2019). Salivary Gland Hypofunction and Xerostomia in Head and Neck Radiation Patients. J Natl Cancer Inst Monogr *2019*. 10.1093/jncimonographs/lgz016.

6. Eisbruch, A., Dawson, L.A., Kim, H.M., Bradford, C.R., Terrell, J.E., Chepeha, D.B., Teknos, T.N., Anzai, Y., Marsh, L.H., Martel, M.K., et al. (1999). Conformal and intensity modulated irradiation of head and neck cancer: the potential for improved target irradiation, salivary gland function, and quality of life. Acta Otorhinolaryngol Belg *53*, 271-275.

7. Henson, B.S., Eisbruch, A., D'Hondt, E., and Ship, J.A. (1999). Two-year longitudinal study of parotid salivary flow rates in head and neck cancer patients receiving unilateral neck parotid-sparing radiotherapy treatment. Oral Oncology *35*, 234-241. https://doi.org/10.1016/S1368-8375(98)00104-3.

8.   Avila, J.L., Grundmann, O., Burd, R., and Limesand, K.H. (2009). Radiation-induced salivary gland dysfunction results from p53-dependent apoptosis. Int J Radiat Oncol Biol Phys *73*, 523-529. 10.1016/j.ijrobp.2008.09.036.

9.   Meyer, S., Chibly, A.M., Burd, R., and Limesand, K.H. (2017). Insulin-Like Growth Factor-1-Mediated DNA Repair in Irradiated Salivary Glands Is Sirtuin-1 Dependent. J Dent Res *96*, 225-232. 10.1177/0022034516677529.

10.   Liu, X., Cotrim, A., Teos, L., Zheng, C., Swaim, W., Mitchell, J., Mori, Y., and Ambudkar, I. (2013). Loss of TRPM2 function protects against irradiation-induced salivary gland dysfunction. Nat Commun *4*, 1515. 10.1038/ncomms2526.

11.   Gilman, K.E., Camden, J.M., Klein, R.R., Zhang, Q., Weisman, G.A., and Limesand, K.H. (2019). P2X7 receptor deletion suppresses gamma-radiation-induced hyposalivation. Am J Physiol Regul Integr Comp Physiol *316*, R687-R696. 10.1152/ajpregu.00192.2018.

12.   Jasmer, K.J., Gilman, K.E., Munoz Forti, K., Weisman, G.A., and Limesand, K.H. (2020). Radiation-Induced Salivary Gland Dysfunction: Mechanisms, Therapeutics and Future Directions. J Clin Med *9*. 10.3390/jcm9124095.

13.   Grundmann, O., Mitchell, G.C., and Limesand, K.H. (2009). Sensitivity of Salivary Glands to Radiation: from Animal Models to Therapies. Journal of Dental Research *88*, 894-903. 10.1177/0022034509343143.

14.   Dirix, P., Nuyts, S., and Van den Bogaert, W. (2006). Radiation-induced xerostomia in patients with head and neck cancer: a literature review. Cancer *107*, 2525-2534. 10.1002/cncr.22302.

15.   Radfar, L., and Sirois, D.A. (2003). Structural and functional injury in minipig salivary glands following fractionated exposure to 70 Gy of ionizing radiation: an animal model for human radiation-induced salivary gland injury. Oral Surg Oral Med Oral Pathol Oral Radiol Endod *96*, 267-274. 10.1016/s1079-2104(03)00369-x.

16.   Li, Y., Taylor, J.M., Ten Haken, R.K., and Eisbruch, A. (2007). The impact of dose on parotid salivary recovery in head and neck cancer patients treated with radiation therapy. Int J Radiat Oncol Biol Phys *67*, 660-669. 10.1016/j.ijrobp.2006.09.021.

17.   Vissink, A., Mitchell, J.B., Baum, B.J., Limesand, K.H., Jensen, S.B., Fox, P.C., Elting, L.S., Langendijk, J.A., Coppes, R.P., and Reyland, M.E. (2010). Clinical management of salivary gland hypofunction and xerostomia in head-and-neck cancer patients: successes and barriers. Int J Radiat Oncol Biol Phys *78*, 983-991. 10.1016/j.ijrobp.2010.06.052.

18. Grün, D., and van Oudenaarden, A. (2015). Design and Analysis of Single-Cell Sequencing Experiments. Cell *163*, 799-810. https://doi.org/10.1016/j.cell.2015.10.039.

19. Kolodziejczyk, Aleksandra A., Kim, J.K., Svensson, V., Marioni, John C., and Teichmann, Sarah A. (2015). The Technology and Biology of Single-Cell RNA Sequencing. Molecular Cell *58*, 610-620. https://doi.org/10.1016/j.molcel.2015.04.005.

20. Trapnell, C. (2015). Defining cell types and states with single-cell genomics. Genome Res *25*, 1491-1498. 10.1101/gr.190595.115.

21. Wang, Y., and Navin, Nicholas E. (2015). Advances and Applications of Single-Cell Sequencing Technologies. Molecular Cell *58*, 598-609. https://doi.org/10.1016/j.molcel.2015.05.005.

22. Tabula Muris, C., Overall, c., Logistical, c., Organ, c., processing, Library, p., sequencing, Computational data, a., Cell type, a., Writing, g., et al. (2018). Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. Nature *562*, 367-372. 10.1038/s41586-018-0590-4.

23. Tabula Sapiens, C., Jones, R.C., Karkanias, J., Krasnow, M.A., Pisco, A.O., Quake, S.R., Salzman, J., Yosef, N., Bulthaup, B., Brown, P., et al. (2022). The Tabula Sapiens: A multiple-organ, single-cell transcriptomic atlas of humans. Science *376*, eabl4896. 10.1126/science.abl4896.

24. Chen, M., Lin, W., Gan, J., Lu, W., Wang, M., Wang, X., Yi, J., and Zhao, Z. (2022). Transcriptomic Mapping of Human Parotid Gland at Single-Cell Resolution. J Dent Res *101*, 972-982. 10.1177/00220345221076069.

25. Huang, N., Perez, P., Kato, T., Mikami, Y., Okuda, K., Gilmore, R.C., Conde, C.D., Gasmi, B., Stein, S., Beach, M., et al. (2021). SARS-CoV-2 infection of the oral cavity and saliva. Nat Med *27*, 892-903. 10.1038/s41591-021-01296-8.

26. Hauser, B.R., Aure, M.H., Kelly, M.C., Hoffman, M.P., and Chibly, A.M. (2020). Generation of a Single-Cell RNAseq Atlas of Murine Salivary Gland Development. iScience *23*, 101838. https://doi.org/10.1016/j.isci.2020.101838.

27. Oyelakin, A., Song, E.A.C., Min, S., Bard, J.E., Kann, J.V., Horeth, E., Smalley, K., Kramer, J.M., Sinha, S., and Romano, R.A. (2019). Transcriptomic and Single-Cell Analysis of the Murine Parotid Gland. J Dent Res *98*, 1539-1547. 10.1177/0022034519882355.

28. Sekiguchi, R., Martin, D., Genomics, Computational Biology, C., and Yamada, K.M. (2020). Single-Cell RNA-seq Identifies Cell Diversity in Embryonic Salivary Glands. J Dent Res *99*, 69-78. 10.1177/0022034519883888.

29. Praktiknjo, S.D., Obermayer, B., Zhu, Q., Fang, L., Liu, H., Quinn, H., Stoeckius, M., Kocks, C., Birchmeier, W., and Rajewsky, N. (2020). Tracing tumorigenesis in a solid tumor model at single-cell resolution. Nat Commun *11*, 991. 10.1038/s41467-020-14777-0.

30. Horeth, E., Oyelakin, A., Song, E.C., Che, M., Bard, J., Min, S., Kiripolsky, J., Kramer, J.M., Sinha, S., and Romano, R.A. (2021). Transcriptomic and Single-Cell Analysis Reveals Regulatory Networks and Cellular Heterogeneity in Mouse Primary Sjogren's Syndrome Salivary Glands. Front Immunol *12*, 729040. 10.3389/fimmu.2021.729040.

31. Hong, X., Meng, S., Tang, D., Wang, T., Ding, L., Yu, H., Li, H., Liu, D., Dai, Y., and Yang, M. (2020). Single-Cell RNA Sequencing Reveals the Expansion of Cytotoxic CD4(+) T Lymphocytes and a Landscape of Immune Cells in Primary Sjogren's Syndrome. Front Immunol *11*, 594658. 10.3389/fimmu.2020.594658.

32. Xu, Y., Feng, S., Peng, Q., Zhu, W., Zu, Q., Yao, X., Zhang, Q., Cao, J., and Jiao, Y. (2021). Single-cell RNA sequencing reveals the cell landscape of a radiation-induced liver injury mouse model. Radiation Medicine and Protection *2*, 181-183. 10.1016/j.radmp.2021.11.001.

33. Mukherjee, A., Epperly, M.W., Shields, D., Hou, W., Fisher, R., Hamade, D., Wang, H., Saiful Huq, M., Bao, R., Tabib, T., et al. (2021). Ionizing irradiation-induced Fgr in senescent cells mediates fibrosis. Cell Death Discov *7*, 349. 10.1038/s41420-021-00741-4.

34. Paldor, M., Levkovitch-Siany, O., Eidelshtein, D., Adar, R., Enk, C.D., Marmary, Y., Elgavish, S., Nevo, Y., Benyamini, H., Plaschkes, I., et al. (2022). Single-cell transcriptomics reveals a senescence-associated IL-6/CCR6 axis driving radiodermatitis. EMBO Mol Med, e15653. 10.15252/emmm.202115653.

35. Ferreira, J.N.A., Zheng, C., Lombaert, I.M.A., Goldsmith, C.M., Cotrim, A.P., Symonds, J.M., Patel, V.N., and Hoffman, M.P. (2018). Neurturin Gene Therapy Protects Parasympathetic Function to Prevent Irradiation-Induced Murine Salivary Gland Hypofunction. Mol Ther Methods Clin Dev *9*, 172-180. 10.1016/j.omtm.2018.02.008.

36. Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*, 1888-1902.e1821. https://doi.org/10.1016/j.cell.2019.05.031.

37. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*, 3573-3587 e3529. 10.1016/j.cell.2021.04.048.

38. Zappia, L., and Oshlack, A. (2018). Clustering trees: a visualization for evaluating clusterings at multiple resolutions. GigaScience *7*. 10.1093/gigascience/giy083.

39. Michael, D.G., Pranzatelli, T.J.F., Warner, B.M., Yin, H., and Chiorini, J.A. (2019). Integrated Epigenetic Mapping of Human and Mouse Salivary Gene Regulation. J Dent Res *98*, 209-217. 10.1177/0022034518806518.

40. Kuhn, M., von Mering, C., Campillos, M., Jensen, L.J., and Bork, P. (2008). STITCH: interaction networks of chemicals and proteins. Nucleic Acids Res *36*, D684-688. 10.1093/nar/gkm795.

41. Ramilowski, J.A., Goldberg, T., Harshbarger, J., Kloppmann, E., Lizio, M., Satagopam, V.P., Itoh, M., Kawaji, H., Carninci, P., Rost, B., and Forrest, A.R. (2015). A draft network of ligand-receptor-mediated multicellular signalling in human. Nat Commun *6*, 7866. 10.1038/ncomms8866.

42. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.H., Myung, P., Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat. Nat Commun *12*, 1088. 10.1038/s41467-021-21246-9.

43. Teos, L.Y., Zheng, C.Y., Liu, X., Swaim, W.D., Goldsmith, C.M., Cotrim, A.P., Baum, B.J., and Ambudkar, I.S. (2016). Adenovirus-mediated hAQP1 expression in irradiated mouse salivary glands causes recovery of saliva secretion by enhancing acinar cell volume decrease. Gene Therapy *23*, 572-579. 10.1038/gt.2016.29.

44. Zheng, C., Cotrim, A.P., Rowzee, A., Swaim, W., Sowers, A., Mitchell, J.B., and Baum, B.J. (2011). Prevention of radiation-induced salivary hypofunction following hKGF gene delivery to murine submandibular glands. Clin Cancer Res *17*, 2842-2851. 10.1158/1078-0432.CCR-10-2982.

45. Lombaert, I.M., Brunsting, J.F., Wierenga, P.K., Kampinga, H.H., de Haan, G., and Coppes, R.P. (2008). Keratinocyte growth factor prevents radiation damage to salivary glands by expansion of the stem/progenitor pool. Stem Cells *26*, 2595-2601. 10.1634/stemcells.2007-1034.

46. The Gene Ontology Consortium (2019). The Gene Ontology Resource: 20 years and still GOing strong. Nucleic Acids Research *47*, D330-D338. 10.1093/nar/gky1055.

47.   Chibly, A.M., Patel V. N., Aure M. H., Pasquale M. C., NIDCD/NIDCR Genomics and Computational Biology Core, Martin G. E., Ghannam M., Andrade J., Denegre N., Simpson C., Goldstein D. P., Liu F., Lombaert I. M. A., Hoffman, M. P. (2023). Neurotrophin signaling is a central mechanism of salivary dysfunction after irradiation that disrupts myoepithelial cells. NPJ Regenerative Medicine (Accepted, In press).

48.   Song, E.A.C., Smalley, K., Oyelakin, A., Horeth, E., Che, M., Wrynn, T., Osinski, J., Romano, R.A., and Sinha, S. (2022). Genetic Study of Elf5 and Ehf in the Mouse Salivary Gland. J Dent Res, 220345221130258. 10.1177/00220345221130258.

49.   Xiao, N., Lin, Y., Cao, H., Sirjani, D., Giaccia, A.J., Koong, A.C., Kong, C.S., Diehn, M., and Le, Q.T. (2014). Neurotrophic factor GDNF promotes survival of salivary stem cells. J Clin Invest *124*, 3364-3377. 10.1172/JCI74096.

50.   Katsumata, O., Sato, Y., Sakai, Y., and Yamashina, S. (2009). Intercalated duct cells in the rat parotid gland may behave as tissue stem cells. Anat Sci Int *84*, 148-154. 10.1007/s12565-009-0019-0.

51.   Denny, P.C., Liu, P., and Denny, P.A. (1999). Evidence of a phenotypically determined ductal cell lineage in mouse salivary glands. The Anatomical Record *256*, 84-90. 10.1002/(sici)1097-0185(19990901)256:1<84::Aid-ar11>3.0.Co;2-s.

52.   Denny, P.C., Ball, W.D., and Redman, R.S. (1997). Salivary glands: a paradigm for diversity of gland development. Crit Rev Oral Biol Med *8*, 51-75. 10.1177/10454411970080010301.

53.   Miyazaki, Y., Nakanishi, Y., and Hieda, Y. (2004). Tissue interaction mediated by neuregulin-1 and ErbB receptors regulates epithelial morphogenesis of mouse embryonic submandibular gland. Dev Dyn *230*, 591-596. 10.1002/dvdy.20078.

54.   Nedvetsky, Pavel I., Emmerson, E., Finley, Jennifer K., Ettinger, A., Cruz-Pacheco, N., Prochazka, J., Haddox, Candace L., Northrup, E., Hodges, C., Mostov, Keith E., et al. (2014). Parasympathetic Innervation Regulates Tubulogenesis in the Developing Salivary Gland. Developmental Cell *30*, 449-462. https://doi.org/10.1016/j.devcel.2014.06.012.

55.   Mattingly, A., Finley, J.K., and Knox, S.M. (2015). Salivary gland development and disease. Wiley Interdiscip Rev Dev Biol *4*, 573-590. 10.1002/wdev.194.

56.   Lombaert, I.M.A., Patel, V.N., Jones, C.E., Villier, D.C., Canada, A.E., Moore, M.R., Berenstein, E., Zheng, C., Goldsmith, C.M., Chorini, J.A., et al. (2020). CERE-120 Prevents Irradiation-Induced Hypofunction and Restores Immune Homeostasis in Porcine Salivary Glands. Molecular Therapy - Methods & Clinical Development *18*, 839-855. https://doi.org/10.1016/j.omtm.2020.07.016.

57. Shamblott, M.J., O'Driscoll, M.L., Gomez, D.L., and McGuire, D.L. (2016). Neurogenin 3 is regulated by neurotrophic tyrosine kinase receptor type 2 (TRKB) signaling in the adult human exocrine pancreas. Cell Communication and Signaling *14*, 23. 10.1186/s12964-016-0146-x.

58. Ghinelli, E., Johansson, J., Ríos, J.D., Chen, L.-L., Zoukhri, D., Hodges, R.R., and Dartt, D.A. (2003). Presence and Localization of Neurotrophins and Neurotrophin Receptors in Rat Lacrimal Gland. Investigative Ophthalmology & Visual Science *44*, 3352-3357. 10.1167/iovs.03-0037.

59. Lombaert, I.M., Brunsting, J.F., Wierenga, P.K., Faber, H., Stokman, M.A., Kok, T., Visser, W.H., Kampinga, H.H., de Haan, G., and Coppes, R.P. (2008). Rescue of salivary gland function after stem cell transplantation in irradiated glands. PLoS One *3*, e2063. 10.1371/journal.pone.0002063.

60. Ninche, N., Kwak, M., and Ghazizadeh, S. (2020). Diverse epithelial cell populations contribute to the regeneration of secretory units in injured salivary glands. Development *147*. 10.1242/dev.192807.

61. Naik, S., Larsen, S.B., Cowley, C.J., and Fuchs, E. (2018). Two to Tango: Dialog between Immunity and Stem Cells in Health and Disease. Cell *175*, 908-920. https://doi.org/10.1016/j.cell.2018.08.071.

62. Chakrabarti, R., Celià-Terrassa, T., Kumar, S., Hang, X., Wei, Y., Choudhury, A., Hwang, J., Peng, J., Nixon, B., Grady, J.J., et al. (2018). Notch ligand Dll1 mediates cross-talk between mammary stem cells and the macrophageal niche. Science *360*, eaan4153. 10.1126/science.aan4153.

63. Biton, M., Haber, A.L., Rogel, N., Burgin, G., Beyaz, S., Schnell, A., Ashenberg, O., Su, C.-W., Smillie, C., Shekhar, K., et al. (2018). T Helper Cell Cytokines Modulate Intestinal Stem Cell Renewal and Differentiation. Cell *175*, 1307-1320.e1322. https://doi.org/10.1016/j.cell.2018.10.008.

64. Mansfield, K., and Naik, S. (2020). Unraveling Immune-Epithelial Interactions in Skin Homeostasis and Injury. Yale J Biol Med *93*, 133-143.

65. Agace, W.W., Higgins, J.M., Sadasivan, B., Brenner, M.B., and Parker, C.M. (2000). T-lymphocyte-epithelial-cell interactions: integrin alpha(E)(CD103)beta(7), LEEP-CAM and chemokines. Curr Opin Cell Biol *12*, 563-568. 10.1016/s0955-0674(00)00132-0.

66. Teymoortash, A., Simolka, N., Schrader, C., Tiemann, M., and Werner, J.A. (2005). Lymphocyte subsets in irradiation-induced sialadenitis of the submandibular gland. Histopathology *47*, 493-500. 10.1111/j.1365-2559.2005.02256.x.

67. Preston, G.C., Feijoo-Carnero, C., Schurch, N., Cowling, V.H., and Cantrell, D.A. (2013). The Impact of KLF2 Modulation on the Transcriptional Program and Function of CD8 T Cells. PLOS ONE *8*, e77537. 10.1371/journal.pone.0077537.

68. Ebert, R., Zeck, S., Meissner-Weigl, J., Klotz, B., Rachner, T.D., Benad, P., Klein-Hitpass, L., Rudert, M., Hofbauer, L.C., and Jakob, F. (2012). Krüppel-like factors KLF2 and 6 and Ki-67 are direct targets of zoledronic acid in MCF-7 cells. Bone *50*, 723-732. https://doi.org/10.1016/j.bone.2011.11.025.

69. Sugita, S., Horie, S., Nakamura, O., Maruyama, K., Takase, H., Usui, Y., Takeuchi, M., Ishidoh, K., Koike, M., Uchiyama, Y., et al. (2009). Acquisition of T Regulatory Function in Cathepsin L-Inhibited T Cells by Eye-Derived CTLA-2α during Inflammatory Conditions. The Journal of Immunology *183*, 5013. 10.4049/jimmunol.0901623.

70. Sugita, S., Horie, S., Nakamura, O., Futagami, Y., Takase, H., Keino, H., Aburatani, H., Katunuma, N., Ishidoh, K., Yamamoto, Y., and Mochizuki, M. (2008). Retinal Pigment Epithelium-Derived CTLA-2α Induces TGFβ-Producing T Regulatory Cells. The Journal of Immunology *181*, 7525. 10.4049/jimmunol.181.11.7525.

71. Sugita, S., Yamada, Y., Horie, S., Nakamura, O., Ishidoh, K., Yamamoto, Y., Yamagami, S., and Mochizuki, M. (2011). Induction of T Regulatory Cells by Cytotoxic T-Lymphocyte Antigen-2α on Corneal Endothelial Cells. Investigative Ophthalmology & Visual Science *52*, 2598-2605. 10.1167/iovs.10-6322.

72. McCarthy, D.D., Summers-Deluca, L., Vu, F., Chiu, S., Gao, Y., and Gommerman, J.L. (2006). The lymphotoxin pathway. Immunologic Research *35*, 41-53. 10.1385/IR:35:1:41.

73. Wolf, M.J., Seleznik, G.M., Zeller, N., and Heikenwalder, M. (2010). The unexpected role of lymphotoxin β receptor signaling in carcinogenesis: from lymphoid tissue formation to liver and prostate cancer development. Oncogene *29*, 5006-5018. 10.1038/onc.2010.260.

74. Tumanov, A.V., Koroleva, E.P., Christiansen, P.A., Khan, M.A., Ruddy, M.J., Burnette, B., Papa, S., Franzoso, G., Nedospasov, S.A., Fu, Y.X., and Anders, R.A. (2009). T Cell-Derived Lymphotoxin Regulates Liver Regeneration. Gastroenterology *136*, 694-704.e694. https://doi.org/10.1053/j.gastro.2008.09.015.

75. Wong, G.H.W. (1995). Protective roles of cytokines against radiation: Induction of mitochondrial MnSOD. Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease *1271*, 205-209. https://doi.org/10.1016/0925-4439(95)00029-4.

A)

Single cell RNAseq from control and irradiated adult mouse PG

1-year-old old healthy PG

1-year-old old irradiated PG
5 x 6Gy (10 months post-IR)

- Acinar
- Etv1+
- Intercalated duct
- Striated duct
- Myoepithelial
- Stromal
- Endothelial
- B-cells
- T-cells CD4+
- T-cells CD4+CD8+
- T-cells CD8+
- T-cells FoxP3+
- T-cells Cxcr6+
- Macrophages
- DCs
- NK cells

B   Control PG

Percent Expressed
- 25
- 50
- 75
- 100

Average Expression
2.5
2.0
1.5
1.0
0.5
0.0

C

Etv1

Amy1

A

B

Acinar
Bpifa2^{hi}Amy1^{hi}

Etv1+

49 | 30 | 191

Prb1
Prpmp5
Aqp5
Tcn2
Lrrc26

Amy1
Bpifa2
Pip
Car6
Krt8

Etv1
Col7a1
Gm42892
Pdk4
Pnmal2

% of epithelial-defining genes enriched in Etv1+ cells

C

D Healthy 1 year-old PG
Etv1 NKCC1 DAPI

Duct

E

GO Biological Processes			
Pathway ID	Pathway description	Gene count	FDR
GO.0048519	Negative regulation of biological process	68	6.27E-08
GO.0009888	Tissue development	40	8.90E-08
GO.0048513	Organ development	53	1.59E-07
GO.0048523	Negative regulation of cellular process	63	1.59E-07
GO.0031325	Positive regulation of metabolic process	51	3.08E-07
KEGG pathways			
Pathway ID	Pathway description	Gene count	FDR
4015	Rap1 signaling pathway	13	4.92E-06
5210	Colorectal cancer	8	6.08E-06
4668	TNF signaling pathway	8	0.000203
5161	Hepatitis B	9	0.000203
4012	ErbB signaling pathway	7	0.000291

Figure S1: Annotation strategy

A) Unsupervised clustering of integrated control and irradiated mouse parotid gland (n=1 per treatment)
B) Balloon plot of top cluster-defining genes. Color is relative to scaled gene expression and size of the dot represents the percentage of cells within a cluster expressing the gene
C) UMAP highlighting cells that express the stromal markers Col1a1 and Vim
D) UMAP highlighting myoepithelial cells that express Krt14 and Acta2

A Intercalated duct   Etv1+

| 31 | 19 | 202 |

Dcpp2   *Dcpp1*   *Etv1*
Dcpp3   *Bpifa2*   *Chka*
Wfdc18   *Ehf*   *Gm42892*
Bglap3   *Krt18*   *Neat1*
Pglyrp   *Nupr1*   *Errfi1*

B Striated duct   Etv1+

| 164 | 17 | 204 |

Klk1   *Krt18*   *Etv1*
Fxyd2   *Krt8*   *Chka*
Crisp3   *Atf3*   *Gm42892*
Wfdc2   *Mt1*   *Neat1*
Ldhb   *Plet1*   *Phlda*

C

% of Etv1+ cell-defining genes enriched in other cells

Acinar / ID / SD

D

GO Biological Processes (Acinar cells)			
Pathway ID	**Pathway description**	**Gene count**	**FDR**
GO.0002018	renin-angiotensin regulation of aldosterone production	3	0.000359
GO.0003081	regulation of systemic arterial blood pressure by renin-angiotensin	5	0.000359
GO.0003073	regulation of systemic arterial blood pressure	6	0.00163
GO.0002035	brain renin-angiotensin system	3	0.00227
GO.0050708	regulation of protein secretion	9	0.00476
KEGG pathways (Acinar cells)			
Pathway ID	**Pathway description**	**Gene count**	**FDR**
4614	Renin-angiotensin system	5	2.21E-06
4150	mTOR signaling pathway	5	0.000499
4970	Salivary secretion	5	0.000867

Figure S2.

A) Venn diagram comparing defining genes for *Etv1+* and ID populations. The numbers in the left and right panels indicates the number of unique genes in the corresponding population whereas the number in the central panel reflects the overlap between the two populations.

B) Venn diagram comparing defining genes for *Etv1+* and SD populations. The numbers in the left and right panels indicates the number of unique genes in the corresponding population whereas the number in the central panel reflects the overlap between the two populations.

C) Bar graph with percentage of Etv1+ defining genes enriched in other epithelial cells.

D) Results from STITCH analysis showing top biological processes and KEGG pathways associated with defining-genes from acinar cells.

Figure S3. Ligand-receptor analysis of Etv1+ and acinar cells

A-B) Bar graphs with number of identified ligands and receptors among cell-defining genes from all populations.

C) Heatmap of possible interactions between any two cell populations. Red (positive values) in the color bar indicate higher number of predicted interactions.

D-E) Chord plot summarizing putative ligand-receptor interactions with Etv1+ cell ligands. The arrows point to the cell expressing the corresponding receptors and are color-coded based on the source of the ligand. The thickness of the arrow is relative to the number of putative pairs identified between Etv1 cells and the cell type pointed by the arrow. Representative ligand-receptor pairs are shown beside the chord plots.

A

B

GO biological process complete (IR-induced DEGs in acinar cells)	FDR	Genes
regulation of transepithelial transport (GO:0150111)	1.73E-02	Actb, Actg1
electron transport coupled proton transport (GO:0015990)	2.26E-02	mt-Cyb, Mtnd4
energy coupled proton transmembrane transport, against electrochemical gradient (GO:0015988)	2.23E-02	mt-Cyb, Mtnd4
positive regulation of intrinsic apoptotic signaling pathway by p53 class mediator (GO:1902255)	5.35E-04	Ubb, Rpl11, Rps7, Pppia, Rps3, Tpt1, Rack1
regulation of endodeoxyribonuclease activity (GO:0032071)	3.75E-02	Npm1, Rps3
cytoplasmic translation (GO:0002181)	6.15E-15	Rps and Rpl genes
positive regulation of signal transduction by p53 class mediator (GO:1901798)	1.67E-04	Ubb, Rpl11, Rps7, Ddx5

GO biological process complete (IR-induced DEGs in CD4+CD8+ T –cells)	FDR	Genes
T cell receptor V(D)J recombination (GO:0033153)	3.39E-02	Bcl11b, Tcf7
chaperone-mediated protein complex assembly (GO:0051131)	1.45E-02	Hspd1, Hsp90aa1, Hsp90ab1
lymphocyte differentiation (GO:0030098)	5.47E-03	Zfp36l2, Tcf7, Ptpn6, Rhoh, Cd28, Klf6, Gpr183
leukocyte differentiation (GO:0002521)	1.43E-03	Zfp36l2, Tcf7, Junb, Rhoh, Cd28, Klf6, Gpr183, Bcl11b, Jun,
apoptotic signaling pathway (GO:0097190)	3.99E-02	Shisa5, Prkca, Ppp1r15a, Jun, Crip1, Cd28
axonogenesis (GO:0007409)	9.62E-03	Hsp90aa1, Prkca, Emb, Actb, Bcl11b, Evl, Rhoh, Hsp90ab1
regulation of ERK1 and ERK2 cascade (GO:0070372)	4.48E-02	Prkca, Jun, Dusp1, Ptpn6, Gpr183, Ramp3

Figure S4. Cell-specific IR-induced DEGs

A) Violin plots of top 10 (if present) up and downregulated genes in epithelial populations. Red and blue arrows denote upregulated and downregulated genes, respectively.

B) Representative output from gene ontology analysis with IR-induced DEGs in acinar and CD4+CD8+ T-cells showing dysregulated processes and their associated genes.

Figure S5. Dysregulated ligand-receptor pairs post-IR

A) 2D representation of incoming vs outgoing interaction strengths for all cell types relative to each other.

B) Heatmap shows the relative importance of each cell group in IR-PG based on CellChat-computed network centrality measures of NRG, NT, THBS, EPHA, and CEACAM signaling networks. Relative contribution of each ligand-receptor pair to the overall communication network of NRG and NT signaling pathways, which is the ratio of the total communication probability of the inferred network of each ligand-receptor pair to that of the signaling pathway.

C) Differential interactions strength between acinar and Etv1+ populations with all other cell types in IR-PG compared to control. Red (positive values) and blue (negative values) in the color bar indicate higher number of predicted interactions in IR-PG and controls, respectively