Parametric information about eye movements is sent to the ears

Stephanie N Lovich[1-4]* Cynthia D King[1-4]*, David LK Murphy[1,3,4]*, Rachel Landrum[1-4], Christopher A Shera[7], Jennifer M Groh[1-6]

*These authors contributed equally to this work and are listed in reverse chronological order of their period of most intense involvement.

[1]Department of Psychology and Neuroscience
[2]Department of Neurobiology
[3]Center for Cognitive Neuroscience
[4]Duke Institute for Brain Sciences
[5]Department of Computer Science
[6]Department of Biomedical Engineering
Duke University

[7]Dept. of Otolaryngology
University of Southern California

Acknowledgments

We are grateful to Dr. Matthew Cooper, Dr. Kurtis Gruters, Jesse Herche, Dr. David

Kaylie, Dr. Jeff Mohl, Dr. Shawn Willett, Meredith Schmehl, Dr. Jonathan Siegel,

Chadbourne Smith, Dr. David Smith, Justine Shih, Chloe Weiser, Tingan Zhu, for

discussions and other assistance concerning this project. This work was supported by

NIH (NIDCD) grant DC017532 to JMG.

Abstract

Eye movements alter the relationship between the visual and auditory spatial scenes. Signals related to eye movements affect neural pathways from the ear through auditory cortex and beyond, but how these signals contribute to computing the locations of sounds with respect to the visual scene is poorly understood. Here, we evaluated the information contained in eye movement-related eardrum oscillations (EMREOs), pressure changes recorded in the ear canal that occur in conjunction with simultaneous eye movements. We show that EMREOs contain parametric information about horizontal and vertical eye displacement as well as initial/final eye position with respect to the head. The parametric information in the horizontal and vertical directions can be modelled as combining linearly, allowing accurate prediction of the EMREOs associated with oblique (diagonal) eye movements. Target location can also be inferred from the EMREO signals recorded during eye movements to those targets. We hypothesize that the (currently unknown) mechanism underlying EMREOs could impose a two-dimensional eye-movement related transfer function on any incoming sound, permitting subsequent processing stages to compute the positions of sounds in relation to the visual scene.

Significance Statement

When the eyes move, the alignment between the visual and auditory scenes changes. We are not perceptually aware of these shifts -- which indicates that the brain must incorporate accurate information about eye movements into auditory and visual processing. Here we show that the small sounds generated within the ear by the brain contain accurate information about contemporaneous eye movements in the spatial domain: the direction and amplitude of the eye movements could be inferred from these small sounds. The underlying mechanism(s) likely involve(s) the ear's various motor structures, and could facilitate the translation of incoming auditory signals into a frame of reference anchored to the direction of the eyes and hence the visual scene.

Introduction

Every time we move our eyes to localize multisensory stimuli, our retinae move in relation to our ears. These movements shift the alignment of the visual scene (as detected by the retinal surface) with respect to the auditory scene (as detected based on timing, intensity, and frequency differences in relation to the head and ears). Precise information about each eye movement is therefore needed to connect the brain's views of visual and auditory space to one another (e.g. e.g. Groh and Sparks, 1992; Boucher et al., 2001; Metzger et al., 2004). Most previous work about how eye movement information is incorporated into auditory processing has focused on cortical and subcortical brain structures (Jay and Sparks, 1984, 1987b, a; Russo and Bruce, 1994; Hartline et al., 1995; Stricanne et al., 1996; Cohen and Andersen, 2000; Groh et al., 2001; Zella et al., 2001; Werner-Reiss et al., 2003; Fu et al., 2004; Populin et al., 2004; Zwiers et al., 2004; Mullette-Gillman et al., 2005; Porter et al., 2006; Mullette-Gillman et al., 2009; Maier and Groh, 2010; Bulkin and Groh, 2012a, b; Lee and Groh, 2012; Caruso et al., 2021), but the recent discovery of eye-movement related eardrum oscillations (EMREOs) (Gruters et al., 2018) suggests that the process might manifest much earlier in the auditory periphery. EMREOs can be thought of as a biomarker of underlying efferent information impacting the internal structures of the ear in association with eye movements. What information this efferent signal contains is currently uncertain.

We reasoned that if this efferent signal is to play a role in linking auditory and visual space across eye movements, EMREOs should be *parametrically* related to the associated eye movement. Specifically, EMREOs should vary in a regular and predictable fashion with both horizontal and vertical displacements of the eyes, and some form of information regarding the initial position of the eyes should also be present. These properties are required if the efferent signal underlying EMREOs is to play a role in linking hearing and vision. Notably, this parametric relationship is not required of alternative possible roles, such as synchronizing visual and auditory processing in time or enhanced attentional processing of sounds regardless of their spatial location (Barczak et al., 2019; O'Connell et al., 2020; Köhler et al., 2021; Köhler and Weisz, 2023; Leszczynski et al., 2023).

Accordingly, we evaluated the parametric spatial properties of EMREOs in

human participants by varying the starting and ending positions of visually-guided

saccades in two dimensions. We find that EMREOs do in fact vary parametrically

depending on the saccade parameters in both horizontal and vertical dimensions and as

a function of both initial eye position in the orbits and the change in eye position relative

to that initial position. EMREOs associated with oblique saccades can be predicted by

the linear combination of the EMREOs associated with strictly horizontal and vertical

saccades. Furthermore, an estimate of target location can be decoded from EMREOs

alone – i.e. where subjects looked in space can be roughly determined from their

observed EMREOs.

These findings suggest that the eye-movement information needed to accomplish

a coordinate transformation of incoming sounds into a visual reference frame is fully

available in the most peripheral part of the auditory system. While the precise mechanism

that creates EMREOs remains unknown, we propose that the underlying mechanisms

might introduce a transfer function to the sound transduction process that serves to adjust

the gain, latency, and/or spectral dependence of responses in the cochlea. In principle,

this could provide later stages of auditory processing access to an eye-centered signal

of sound location for registration with the eye-centered visual scene (Groh and Sparks,

1992), Indeed, recent work has shown that changes in muscular tension on the ossicular

chain would be expected to affect gain and latency of sound transmission through the

middle ear, thus supporting the plausibility of this hypothesis (Gallagher et al., 2021; Cho

et al., 2023).

Figure 1. Events of the tasks in time and space. a. Task events across time. Each trial
began with the onset of an initial "fixation" cue (black trace). Participants made saccades
to the fixation point, then maintained fixation for a minimum of 200 ms. The fixation point
was then turned off and a new "target" was turned on (green trace). Participants
saccaded to this target and fixated for another 200 ms, at which point the target turned
red indicating that the trial was over. The ear-canal recordings were analyzed in
conjunction with the fixation-point-to-target saccade. b-d. Spatial layouts of fixation
points and targets for the three task designs used in this study. Points in space that were
used as both a fixation and a target across different trials are half-green, half-black.

Figure 2. *EMREOs recorded during the five-origin grid task. Each panel shows the grand average EMREO signal generated when saccades were made to that location on the screen (average of N=10 subjects' individual left ear averages). For example, the top right panel shows microphone recordings during saccades to the top right (contralateral) target location, and the color and line styles of each trace in that panel correspond to saccades from different initial fixation points. e.g. the red traces originated from the rightward fixation, the blue from the leftward fixation etc as indicated by the legend and boxes of the same color and line style. Both magnitude and phase vary as a function of initial eye position and target location, with contralateral responses being larger than ipsilateral. Phase reversal occurs based on the location of the target with respect to the initial fixation position, as can be seen for the central target location (central panel), where the EMREOs evoked for saccades from the rightward fixation (red traces) show an opposite phase relationship as those evoked for saccades from*

116 *the leftward fixation (blue traces). Corresponding grand averages for right ear data are*
117 *shown in Supplementary Figure 2. These data are presented in Z-units; the peak-*
118 *equivalent sound levels for 18 degree horizontal targets are roughly 55-56 dB SPL; see*
119 *Supplementary Figure 1 for the mean and distributions across the subject population*
120 *(range ~49-64 dB SPL).*

Results

We used earbud microphones to record internally-generated oscillations in the
ear canals of human subjects (n=10 for each task) while they performed eye-movement
tasks involving various visual fixation and target configurations (Figure 1). No external
sounds were presented in any task. The events of the tasks in time are shown in Figure
1a. At the beginning of each trial, subjects fixated a visual fixation point for a minimum
of 200 ms and then made a saccade to a second target, which they then fixated for
another 200 ms. Any trials with micro- or corrective saccades during the 200 ms prior to
or following main fixation-point-to-target saccade were discarded, to ensure a stable
baseline ear-canal recording could be established without intrusions by other eye
movements. Ten subjects were tested in the single origin and horizontal/vertical tasks,
and ten were tested in the five-origin grid task. Four subjects participated in both
groups, so that 16 subjects (8 female, 8 male) were tested overall. Female-male ratios
were equal in both subgroups.

We first tested subjects (N=10) on a task involving variation in both initial fixation
position and target locations varying along both horizontal and vertical dimensions – the
"five-origin grid task". Subjects fixated on an initial fixation light located either straight
ahead, 9° left or right, or 6° up or down, and then made a saccade to a target located
within the array of possible target locations spanning +/- 18° horizontally and +/- 12°
vertically as shown in Figure 1b. Results of this task are shown in Figure 2. Each panel
shows the average microphone signal recorded in the left ear canal (averaged across
all subjects) associated with saccades to a target at that location – e.g. the top right
panel shows all saccades to the top right target location. The color and line styles of the
waveforms correspond to the five initial fixation positions from which the saccades could
originate in space.

The first overall observation from this figure is that the *magnitude* of the
waveform of the EMREO depends on both the *horizontal* and *vertical* dimensions.   In

the horizontal dimension, EMREOs are larger for more contralateral target locations:

compare the column on the right (contralateral) to the column on the left (ipsilateral).

The pattern is reversed for right ear canal recordings (Supplementary Figure 2). In the

vertical dimension, EMREOs are larger for higher vs lower targets in both left and right

ears (compare top row to bottom row in Figure 2/Supplementary Figure 2).

The second overall observation from this figure is that the *phase* of the EMREO

waveform depends on the horizontal location of the target *with respect to* the fixation

position. Specifically, the first deflection after saccade onset is a peak for the most

ipsilateral targets (left-most column) and trough for the most contralateral targets (right-

most column). But *where* this pattern reverses depends on the initial fixation position.

Specifically, consider the red vs blue traces in the middle column of the figure, which

correspond to targets along the vertical meridian. Red traces involve saccades to these

targets from the fixation position on the right, and thus involve leftward (ipsiversive)

saccades. The red traces in this column begin with a peak followed by a trough. In

contrast, the blue traces involve saccades to these targets from the fixation position on

the left, i.e. rightward or contraversive saccades. The blue traces begin with a trough

followed by a peak. The pattern is particularly evident in the central panel (see arrows).

The phase reversal as a function of the combination of target location and initial

eye position suggests that the EMREO waveforms might align *better* when plotted in an

eye-centered frame of reference. Figure 3 demonstrates that this is indeed the case:

the data from Figure 2 is re-plotted as a function of target location *relative* to the initial

fixation position. The eight panels around the center represent the traces for the subset

of targets that can be fully analyzed in an eye-centered frame, i.e. the targets

immediately left, right, up, down, and diagonal relative to the five fixation locations. By

plotting the data based on the relative location of the targets to the origins, the

waveforms are better aligned, showing no obvious phase reversals.

Figure 3. *Replotting the grand average EMREOs as a function of* relative *target location shows better, but not perfect, correspondence of the EMREOs across different fixation positions. The data shown are a subset of those shown in Figure 2, but here each panel location corresponds to a particular target location defined <u>relative</u> to the associated fixation position. The color/linestyle indicates the associated relative fixation position. For example, the waveforms in the upper right panel all involved 9º rightward and 6º upward saccades; the red trace in that panel indicates those that originated from the 9º right fixation; the blue those from the 9º left fixation etc. Only relative target locations that existed for all 5 fixation positions are plotted, as indicated by the inset. Corresponding right ear data are shown in Supplementary Figure 3.*

Although the waveforms are better aligned when plotted based on target location

relative to initial eye position, some variation related to that fixation position is still

evident in the traces. That is, in each panel, the EMREO waveforms with different

colors/line styles (corresponding to different fixation positions) do not necessarily

superimpose perfectly. This suggests that a model that incorporates both relative target

191   position and original fixation position, in both horizontal and vertical dimensions, is

192   needed to account for the findings. Furthermore, a statistical accounting of these effects

193   is needed. Accordingly, we fit the data to the following regression equation:

195            $Mic(t) = B_H(t)\ H + B_{\Delta H}(t)\ \Delta H + B_V(t)\ V + B_{\Delta V}(t)\ \Delta V + C(t)$ Eq 1

where H and V correspond to the initial horizontal and vertical eye position and ΔH and

ΔV correspond to the respective changes in position associated with that trial. The

slope coefficients B_H, $B_{\Delta H}$, B_V, $B_{\Delta V}$ are time-varying and reflect the dependence of the

microphone signal on the respective eye position/movement parameters. The term $C(t)$

contributes a time-varying "constant" independent of eye movement metric and can be

thought of as the best fitting average oscillation across all initial eye positions and

changes in eye position. We used the measured values of eye position/change in eye

position for this analysis rather than the associated fixation and target locations so as to

incorporate trial-by-trial variability in fixation and saccade accuracy.

This model is a conservative one, assessing whether a linear relationship

between the microphone signal and the relevant eye position/movement variables can

provide a satisfactory fit to the data. As such, it provides a lower bound but does not

preclude that higher quality fits could be achieved via non-linear modeling. This

approach is similar to the general linear models applied to fMRI data (e.g. (Monti, 2011)

and differs chiefly in that we make no assumptions about the underlying temporal profile

of the signal (such as a hemodynamic response function) but allow the temporal pattern

to emerge in the time-varying fits of the coefficients.

Figure 4 shows the average of these time-varying values of the slope coefficients

across subjects (blue = left ear; red = right ear) and provides information about the

contribution of these various eye movement parameters to the EMREO signal.   A

strong, consistent dependence on horizontal eye displacement is observed, consistent

with our previous report (Figure 4a) (Gruters et al., 2018). This component is oscillatory

and begins slightly before the onset of the eye movement, inverting in phase for left vs

right ears. The thickened parts of the line indicate periods of time when this coefficient

differed significantly from 0 with 95% confidence (Shaded areas are +/-SEM). There is

also an oscillatory and binaurally phase-inverting signal related to the initial position of

222 the eyes in the horizontal dimension (Figure 4b). This signal is smaller and more

223 variable across subjects.

224   In the vertical dimension, the effect of vertical saccade amplitude is in phase for

225 both the left and right ears; it exhibits an oscillatory pattern, although not obviously

226 sinusoidal like the one observed for the horizontal saccade amplitude. Initial position of

227 the eyes in the vertical dimension exerts a variable effect across participants such that it

228 is not particularly evident in this grand average analysis; this may be related to poorer

229 abilities to localize sounds in the vertical vs. horizontal dimensions (Hebrank and

230 Wright, 1974a, b; Middlebrooks and Green, 1991; Macpherson and Sabin, 2013).

a Horizontal change-in-eye-position
$B_{\Delta H}(t)$

b Horizontal initial eye position
$B_H(t)$

c Vertical change-in-eye-position
$B_{\Delta V}(t)$

d Vertical initial eye position
$B_V(t)$

e Constant
$C(t)$

Approx saccade offset times

Time re: saccade onset (ms)

5-Origin Grid Task Regression Model

$Mic(t) = B_H(t)\,H + B_{\Delta H}(t)\Delta H + B_V(t)\,V + B_{\Delta V}(t)\Delta V + C(t)$

Grand average N=10
+/- SE ≠ 0 p<0.05

Right ears

Left ears

Figure 4. *Regression analysis of EMREOs shows contributions from multiple aspects of eye movement: horizontal and vertical change-in-eye-position **(a, c)**, horizontal initial eye position **(b)**, as well as a constant component that was consistent*

across saccades (e). The contribution of vertical initial eye position was weaker (d).
The regression involved modeling the microphone signal at each time point, and each
panel shows the time varying values of the coefficients associated with the different
aspects of the eye movement (horizontal vs. vertical, change-in-position and initial
position). The regressions were fit to individual subjects' microphone recordings, and
plotted here as grand averages of these regression coefficients across the N=10
subjects tested in the 5-origin grid task. Microphone signals were z-scored in reference
to baseline variability during a period -150 to 120 ms prior to saccade onset. Results are
presented in units of standard deviation (panel e) or standard deviation per degree
(panels a-d). Shaded areas represent +/-SEM.

Finally, there is a constant term that is similar in the two ears and peaks later with

respect to saccade onset than is the case for the other coefficients (Figure 4e). As

noted above, this constant term can be thought of as encapsulating the average

EMREO waveform that occurs when pooling across all the eye movements in the

dataset, regardless of their initial positions or horizontal or vertical components.

We next investigated whether the fits obtained in one task match those obtained

in a different task. We reasoned that if the information contained in the EMREO signal

reflects the eye movement itself, then task context should not matter. Furthermore, the

regression model should provide a good way to accomplish this comparison since it

does not require that the exact same locations and eye movements be tested across

tasks.

To test these questions, we collected data using two simplified tasks, the single-

origin-grid task (with a single initial fixation in the center, Figure 1c) and the

horizontal/vertical task (with fixation-target pairs on the horizontal and vertical

meridians, generating purely horizontal or vertical saccades, Figure 1d). Ten subjects

(four of whom also completed the 5-origin grid task) completed both the single-origin

grid task and the horizontal/vertical saccade. We fit the results from these tasks using

the same regression procedure but omitting the initial fixation position terms, i.e.:

Mic(t) = B$_{\Delta H}$(t) ΔH + B$_{\Delta V}$(t) ΔV + C(t)   *Eq 2*

As shown in Figure 5, both tasks yield similar values of the regression

coefficients for horizontal change-in-position (*B$_{\Delta H}$(t)*) and the constant term (*C(t)*) (grand

average across the population, black vs. green traces).   The vertical change-in-position

term ($B_{\Delta V}(t)$) was slightly more variable but also quite consistent across tasks.

Figure 5. *Different tasks generate similar regression coefficient curves. Grand*
average of the regression results for the single-origin grid (black lines) and
horizontal/vertical (green lines) tasks. The lines and shading represent the average and
standard error of the coefficient values across the same 10 subjects for the two tasks.
See Supplementary Figure 4 for corresponding findings among the individual subjects.

Given the consistency of the regression coefficient values between the single-

origin grid and horizontal/vertical tasks (and see (Lovich et al., 2023) for similar results

involving spontaneous vs. task-guided eye movements), we surmised that it should be

possible to use the *coefficient* values from one task to predict the EMREO *waveforms* in

the other. Specifically, we used the time-varying regression values from purely

horizontal and purely vertical saccades in the horizontal/vertical task to predict the

observed waveforms from oblique saccades in the single origin grid task. This method

can be used to evaluate the quality of the regression-based EMREO prediction not only

for target locations tested in both tasks, i.e. the horizontal and vertical meridians, but

also for oblique targets tested only in the grid task.

The black traces in Figure 6 show the grand average microphone signals

associated with each target in the single-origin grid task. The location of each trace

corresponds to the physical location of the associated target in the grid task (similar to

Figure 2). The superimposed predicted wave forms (red traces) were generated from

the $B_{\Delta H}(t)$, $B_{\Delta V}(t)$, and $C(t)$ regression coefficients fit to only the horizontal/vertical data,

then evaluated at each target location and moment in time to produce predicted curves

for each of the locations tested in the grid task.

Figure 6. *Regression coefficients fit to microphone recordings from the*
horizontal/vertical-saccade task can be used to predict the waveforms observed in the
grid task and their corresponding target locations. Combined results for all N=10
participants' left ears. The black traces indicate the grand average of all the individual
participants' mean microphone signals during the single-origin grid task, with the
shading indicating +/- the standard error across participants. The red traces show an
estimate of the EMREO at that target location based only on regression coefficients
measured from the horizontal/vertical task. Black values in parentheses are the actual
horizontal and vertical coordinates for each target in the grid task. Corresponding red
values indicate the inferred target location based on solving a multivariate regression
which fits the observed grid task microphone signals in a time window (-5 to 70 ms with

respect to saccade onset) to the observed regression weights from the
horizontal/vertical task for each target location. The averages of these values in the
horizontal and vertical dimensions are shown across the top and right sides. See
Figure 7 for additional plots of the inferred vs actual target values, and Supplementary
Figure 5 for corresponding right-ear data.

Overall, there is good correspondence between the predicted EMREO

oscillations and the observed EMREO from actual microphone recordings, including the

oblique target locations that were not tested in the horizontal/vertical task. This

illustrates two things: 1) the EMREO is reproducible across task contexts, and 2) the

horizontal and vertical change-in-position contributions interact in a reasonably

independent way, so that the EMREO signal observed for a combined horizontal-

vertical saccade can be predicted as the sum of the signals observed for purely

horizontal and purely vertical saccades with the corresponding component amplitudes.

Given that it is possible to predict the microphone signal from one task context to

another, it should also be possible to decode the target location and its associated eye

movement from just the simultaneously-recorded microphone signal.   To do this, we

again used the weights from the horizontal/vertical task data for the regression

equation:

$\text{Mic}(t) = B_{\Delta H}(t)\, \Delta H + B_{\Delta V}(t)\, \Delta V + C(t)$ Eq 2

Specifically, we used the Mic(t) values observed in the single-origin grid task to solve

this system of multivariate linear equations across the time window -5 to 70 ms with

respect to the saccade (a time period in which the EMREO appears particularly

consistent and substantial in magnitude) to generate the "read out" values of ΔH and ΔV

associated with each target's actual ΔH and ΔV. We conducted this analysis on the left

ear and right ear data separately. The left ear results of this analysis are seen in each of

the individual panels of Figure 6; the black values (e.g. -18, 12) indicate the actual

horizontal and vertical locations of the target, and the associated red values indicate the

inferred location of the target. Across the top of the figure, the numbers indicate the

average inferred horizontal location, and down the right side, the numbers indicate the

average inferred vertical location. These results indicate that, on average, the targets

can be read out in the proper order, but the spatial scale is compressed: the average

read-out values for the +/-18 degree horizontal targets are +/- ~11-12 degrees, and the

averages for the vertical +/- 12 degree targets are +/- ~6-7 degrees. Similar findings

applied to the right ear data (Supplementary Figure 5).

Plots of these target readouts in both horizontal and vertical dimensions for both

ears are shown in Figure 7a-f. Figure 7a shows the inferred location of the target (red

dots) connected to the actual location of the target (black dots) using the data from

Figure 6, i.e the left ear readout, and Figure 7b-c show regressions of these target

readouts as a function of the horizontal and vertical locations. Figure 7d-f show the

corresponding results for the right ears. Altogether, these figures illustrate that the

readout accuracy is better in the horizontal than in the vertical dimensions.

Quantitatively, the r^2 values for the horizontal dimension were 0.89 (LE) and 0.91 (RE),

and the corresponding values for the vertical dimension were 0.61 (LE) and 0.67 (RE).

Slopes were also closer to a value of 1 for the horizontal dimension (0.71, LE; 0.77, RE)

than for the vertical dimension (0.51, LE, 0.51, RE).

Given that it is known that the brain uses binaural computations for

reconstructing auditory space, we wondered whether the accuracy of this read-out could

be improved by combining signals recorded in each ear simultaneously. We first

considered a binaural difference computation, subtracting the right ear microphone

recordings from the left, thus eliminating the part of the signal that is common between

the two ears. Figure 7g shows the results. Generally, the horizontal dimension is well

ordered whereas the vertical dimension continues to show considerable shuffling. This

can also be seen in Figure 7h and 7i, which show the relationship between the inferred

target location and the true target location, plotted on the horizontal and vertical

dimension, respectively. The correlation between inferred and actual target is higher in

the horizontal dimension (r^2 0.95) than the vertical dimension (r^2 0.41), which is actually

worse than the monaural readouts. This makes sense because the binaural difference

computation serves to diminish the contribution from aspects of the signal that are in

phase across the two ears, such as the dependence on vertical change in eye position.

We then reasoned that improvement in the vertical readout could be achieved by

instead averaging (rather than subtracting) the signals across the two ears, and indeed

this is so. Averaging across the two ears produces an improved vertical readout (r^2

0.73, Figure 7k). A hybrid readout operation in which the horizontal location is

computed from the binaural difference, and the vertical location is computed from the

binaural average, produces a modest improvement in the overall reconstruction of

target location (Figure 7j).

We next considered how well this readout operation performed at the level of

individual subjects. Results for each subject are shown in Supplementary Figure 6 a-j,

and a population summary is shown in Supplementary Figure 6k. The relationship

between the inferred location and the actual location was statistically significant

($p<0.05$) for all 10 subjects in the horizontal dimension, and for 7 of 10 subjects in the

vertical dimension, confirming that we can predict eye position and movement in the

horizontal dimension from individual ear recordings within each subject.

Finally, we evaluated the error when reading out the target location of individual

trials. The preceding analyses show the results when the read-out operation is

performed on the average waveform observed across trials for a given target location.

The same readout can also be computed for each individual trial. When conducted on

all individual trials, the resulting scatter can be computed as the average standard

deviation observed across target locations and subjects. We found that the average

scatter or standard deviation was 19.1 degrees in the horizontal dimension and 23.8

degrees in the vertical dimension. For the horizontal dimension, this corresponds

roughly half of the range of space tested (+/- 18 degrees), whereas in the vertical

dimension this corresponds to nearly the whole range (+/- 12 degrees).

Overall, these results parallel human sound localization which relies on a

binaural difference computation in the horizontal dimension (and is more accurate in

that dimension), vs. potentially monaural or averaged spectral cues for the vertical

dimension (which is less accurate) (Blauert, 1997; Groh, 2014).   (Indeed, horizontal

and vertical sound localization show different patterns of dependence on the loudness

of the target sound relative to background noise, further supporting that these

operations are accomplished via distinct mechanisms. (Ege et al., 2018).)

Figure 7. *Multiple ways of reading out target location from the ear canal recordings. As in Figure 6 and Supplementary Figure 5, the relationship between EMREOs and eye movements was quantitatively modelled using Eq 2 and the ear canal data recorded in the horizontal/vertical task. Inferred grid task target location was "read out" by solving equation (2) for ΔH and ΔV using the coefficients as fit from the horizontal/vertical task and the microphone values as observed in the single-origin grid task; see main text for details. **a.** Inferred target location (red) compared to actual target location (black), based on the left ear (same data as in Figure 6). **b.** Horizontal component of the read-out target vs the actual horizontal component (left ear microphone signals). **c.** Same as **(b)** but for the vertical component. **d-f.** Same as A-C but for the right ear. **g-i,** Same as a-c and d-f but computed using the binaural difference between the microphone signals (left ear – right ear). **j., k.** A hybrid read-out model **(j)** using binaural difference in the horizontal dimension **(h)** and binaural average in the vertical dimension **(k)**. Related findings at the individual subject level are provided in Supplementary Figure 6.*

Discussion

Sound locations are inferred from head-centered differences in sound arrival time, intensity, and spectral content, but visual stimulus locations are inferred from eye-centered retinal locations (Blauert, 1997; Groh, 2014). Information about eye movements with respect to the head/ears is critical for connecting the visual and auditory scenes to one another (Groh and Sparks, 1992). This insight has motivated a number of previous neurophysiological studies in various brain areas in monkeys and

cats, all of which showed that changes in eye position affected the auditory response

properties of at least some neurons in the brain area studied (Inferior colliculus: (Groh

et al., 2001; Zwiers et al., 2004; Porter et al., 2006; Bulkin and Groh, 2012a, b) ;

auditory cortex: (Werner-Reiss et al., 2003; Fu et al., 2004; Maier and Groh, 2010) ;

superior colliculus: (Jay and Sparks, 1984, 1987a, b; Hartline et al., 1995; Zella et al.,

2001; Populin et al., 2004; Lee and Groh, 2012) ; frontal eye fields: (Russo and Bruce,

1994; Caruso et al., 2019) ; intraparietal cortex: (Stricanne et al., 1996; Cohen and

Andersen, 2000; Mullette-Gillman et al., 2005, 2009)).

These findings raised the question of where signals related to eye movements

first appear in the auditory processing stream. The discovery of EMREOs (Gruters et

al., 2018) (Abbasi et al., 2023; King et al., 2023b; Lovich et al., 2023) introduced the

intriguing possibility that the computational process leading to visual-auditory integration

might be manifested in the most peripheral part of the auditory system. Here we show

that the signals present in the ear exhibit the properties necessary for playing a role in

this process: these signals carry information about the horizontal and vertical

components of eye movements, and display signatures related to both change-in-eye-

position and the absolute position of the eyes in the orbits. Because of the parametric

information present in the EMREO signal, we were able to predict EMREOs in one task

from the EMREOs recorded in another, and even predict the target of eye movements

from the simultaneous EMREO recording. These predictions were accomplished using

strictly linear methods, a conservative approach providing a lower bound on what can

be deduced from these signals. Improvements in the "readout" may come from

exploration of more powerful non-linear techniques and/or other refinements such as

tailoring the time window used for the readout (here, a generous -5 to 70 ms) or stricter

criteria for the exclusion of trials contaminated by noise (see "*Methods: Trial exclusion*

criteria"). It should be noted that this read-out presumes knowledge of when the

saccade starts and that performance would be substantially poorer if conducted in a

continuous fashion across time.

Our present observations raise two key questions: what causes EMREOs and

how do those mechanisms impact hearing/auditory processing? The proximate cause of

EMREOs is likely to be one or more of the known types of motor elements in the ear[1]:

the middle ear muscles (stapedius and tensor tympani), which modulate the motion of

the ossicles (Mendelson, 1957; Hung and Dallos, 1972; Gelfand, 1984), and the outer

hair cells, which modulate the motion of the basilar membrane (Brownell et al., 1985).

One or more of these elements may be driven by descending brain signals originating

from within the oculomotor pathway and entering the auditory pathway somewhere

along the descending stream that ultimately reaches the ear via the 5th (tensor tympani),

7th (stapedius muscle), and/or 8th nerves (outer hair cells)   (see refs: Galambos, 1956;

Liberman and Guinan, 1998; Cooper and Guinan, 2006; Guinan, 2006; Mukerji et al.,

2010; Guinan, 2014) for reviews). Efforts are currently underway in our laboratory to

identify the specific EMREO generators/modulators (Schlebusch et al., 2019, 2020)

(King et al., 2023a).

Uncovering the underlying mechanism should shed light on another question.

Does the temporal pattern of the observed EMREO signal reflect the time course and

nature of that underlying mechanism's impact on auditory processing? It is not clear

how an oscillatory signal like the one observed here might contribute to hearing.

However, it is also not clear that the underlying mechanism is, in fact, oscillatory.

Microphones can only detect signals with oscillatory energy in the range of sensitivity of

the microphone. It is possible that the observed oscillations reflect ringing associated

with a change in some mechanical property of the transduction system, and that change

could have a non-oscillatory temporal profile (Figure 8a). Of particular interest would be

a ramp-to-step profile in which aspects of the middle or inner ear shift from one state to

another during the course of a saccade and hold steady at the new state during the

subsequent fixation period. This kind of temporal profile would match the time course of

the saccade itself.

---

[1] We note that EMREOs are unlikely to be due to the actual sound of the eyes moving in the orbits. Our original study, Gruters et al (2018) showed that when microphone recordings are aligned on saccade offset (as opposed to onset, as we did here), EMREOs continue for at least several 10's of ms after the eyes have stopped moving. We also have unpublished observations in patients with various hearing abnormalities; EMREOs are altered in such patients despite normal eye movements.

a Temporal profiles of relevant events and signals

Candidate corollary discharge signals

Saccade: ramp-and-hold

Burst

Tonic

Burst-Tonic

? Underlying mechanism

Observed EMREO: oscillatory

b Working conceptual model

Eye movement command

Retinal location

Copy of eye movement command

Sound localization cues modulated by eye movement command

Synthesized VA object

Figure 8. Temporal profiles of relevant signals and working conceptual model for how EMREOs might relate to our ability to link visual and auditory stimuli in space. ***a.*** *Temporal profiles of signals. The EMREO is oscillatory whereas the eye movement to which it is synchronized involves a ramp-and-hold temporal profile. Candidate source neural signals in the brain might exhibit a ramp-and-hold (tonic) pattern, suggesting a ramp-and-hold-like underlying effect on an as-yet-unknown peripheral mechanism, or could derive from other known temporal profiles including bursts of activity time-locked to saccades.* ***b.*** *Working conceptual model.   The brain causes the eyes to move by sending a command to the eye muscles. Each eye movement shifts the location of visual stimuli on the retinal surface. A copy, possibly a highly transformed one, of this eye movement command is sent to the ear, altering ear mechanics in some unknown way. When a sound occurs, the ascending signal to the brain will depend on the combination of its location in head-centered space (based on the physical values of*

binaural timing and level differences and spectral cues) and aspects of recent eye
movements and fixation position. This hybrid signal could then be read-out by the brain.

Available eye movement control signals in the oculomotor system include those
that follow this ramp-and-hold temporal profile, or tonic activity that is proportional to eye
position throughout periods of both movement and fixation. In addition to such tonic
signals, oculomotor areas also contain neurons that exhibit burst patterns, or elevated
discharge in association with the saccade itself, as well as combinations of burst and
tonic patterns (for reviews, see Fuchs et al., 1985; Takahashi and Shinoda, 2018).   It
remains to be seen which of these signals or signal combinations might be sent to the
auditory periphery and where they might come from. The paramedian pontine reticular
formation (PPRF) is a strong candidate for a source, having been implicated in
providing corollary discharge signals of eye movements in visual experiments (Sparks
et al., 1987) (see also Guthrie et al., 1983), and containing each of these basic temporal
signal profiles (Fuchs et al., 1985; Takahashi and Shinoda, 2018).   Regardless of the
source and nature of the descending corollary discharge signal, the oscillations
observed here should be thought of as possibly constituting a biomarker for an
underlying, currently unknown, mechanism, rather than necessarily the effect itself.

Despite these critical unknowns, it is useful to articulate a working conceptual
model of how EMREOs might facilitate visual and auditory integration (Figure 8b). The
general notion is that, by sending a copy of each eye movement command to the motor
elements of the auditory periphery, the brain keeps the ear informed about the current
orientation of the eyes. If, as noted above, these descending oculomotor signals cause
a ramp-to-step change in the state of tension of components within the EMREO
pathway, time-locked to the eye movement and lasting for the duration of each fixation
period, they would effectively change the transduction mechanism in an eye
position/eye movement dependent fashion. In turn, these changes could affect the
latency, gain, or frequency-filtering properties of the response to sound. Indeed,
intriguing findings from Puria and colleagues (Cho et al., 2023) have recently indicated
that the tension applied by the middle ear muscles likely affects all three of these
aspects of sound transmission throughout the middle ear. In short, the signal sent to
the brain in response to an incoming sound could ultimately reflect a mixture of the

physical cues related to the location of the sound itself - the ITD/ILD/spectral cues - and

eye position/movement information.

Most neurophysiological studies report signals consistent with a hybrid code in

which information about sound location is blended in a complex fashion with information

about eye position and movement, both within and across neurons (Jay and Sparks,

1984, 1987a; Groh et al., 2001; Porter et al., 2006; Maier and Groh, 2010; Lee and

Groh, 2012; Caruso et al., 2019). Computational modeling confirms that, in principle,

these complex signals can be "read out" to produce a signal of sound location with

respect to the eyes (Groh et al., 2001).   However, substantive differences do remain

between the observations here and such neural studies, chiefly in that the neural

investigations have focused primarily on periods of steady fixation. A more complete

characterization of neural signals time-locked to saccades is therefore needed (Porter et

al., 2007; Bulkin and Groh, 2012a).

Note that this working model differs from a spatial attention mechanism in which

the brain might direct the ears to "listen" selectively to a particular location in space.

Rather, under our working model, the response to sounds from any location would be

impacted by peripheral eye movement/position dependence in a consistent fashion

across all sound locations.   However, such a system could well work in concert with

top-down attention, which has previously been shown to impact outer hair cells even

when participants are required to fixate and not make eye movements (Delano et al.,

2007; Harkrider and Bowers, 2009; Srinivasan et al., 2012; Srinivasan et al., 2014;

Walsh et al., 2014; Wittekindt et al., 2014; Walsh et al., 2015).

Another question concerns whether EMREOs might actually impair sound

localization, specifically for brief sounds presented during an eye movement. We think

the answer to this is no.   Boucher et al (Boucher et al., 2001) reported that

perisaccadic sound localization is quite accurate, which suggests that EMREOs (or their

underlying mechanism) do not impair perception. This is an important insight because

given the rate at which eye movements occur - about 3/sec – and with each associated

EMREO signal lasting 100 ms or longer (due to extending past the end of saccades, as

explored by Gruters, Murphy et al. 2018 and (King et al., 2023b)), it would be highly

problematic if sounds could not be accurately detected or localized when they occur in

conjunction with saccades.   If there is indeed a step-ramp system underlying the

observed oscillations, then transduction of all sounds will be affected, regardless of

when they occur with respect to saccades. Indeed, recent work supports the view that

sound detection is unaffected by saccades (Bröhl and Kayser, 2023).

All this being said, a role for EMREOs in computing the spatial location of sounds

with respect to the visual scene does not preclude other roles. Specifically, they could

also play a role in synchronizing sampling in the temporal domain (e.g. (Barczak et al.,

2019; O'Connell et al., 2020; Köhler et al., 2021; Köhler and Weisz, 2023; Leszczynski

et al., 2023)). Such a possibility could account for the significant constant term C(t) of

the regression analysis (Equation 1, Figure 4). This temporally precise non-spatial

component could play a role in resetting or refreshing of auditory processing across

time or coordinating with the refreshing of the visual image on the retina.

Overall, how brain-controlled mechanisms adjust the signaling properties of

peripheral sensory structures is critical for understanding sensory processing as a

whole. Auditory signals are known to adjust the sensitivity of the visual system via

sound-triggered pupil dilation (Bala and Takahashi, 2000), indicating that

communication between these two senses is likely to be a two-way street. The

functional impact of such communication at low-level stages is yet to be fully explored

and may have implications for how individuals compensate when the information from

one sensory system is inadequate, either due to natural situations such as noisy sound

environments or occluded visual ones, or due to physiological impairments in one or

more sensory systems.

Methods

<u>General</u>

Healthy human subjects that were 18 years of age or older with no known

hearing deficits or visual impairments beyond corrected vision were recruited from the

surrounding campus community (N=16; 8 female, 8 male; female-male ratio was also

equal in subgroups tested on different tasks). If subjects were unable to perform the

saccade task without vision correction, they were excluded from the study. All study

procedures involving subjects were approved by the Duke Institutional Review Board,

and all subjects received monetary compensation for their participation.

Acoustic signals in both ear canals were measured simultaneously with Etymotic

ER10B+ microphone systems coupled with ER2 earphones (Etymotic Research, Elk

Grove Village, IL) to allow calibrations of the microphones (however, note that no

auditory stimuli were used during any of the saccade tasks in the current study). A low-

latency audio interface (Focusrite Scarlett 2i2, Focusrite Audio Engineering Ltd., High

Wycombe, UK) was used for audio capture and playback through the Etymotic

hardware at a sampling rate of 48kHz. Eye tracking was performed with an Eyelink

1000 system sampling at 1000Hz. Stimulus presentation and data acquisition were

controlled through custom scripts and elements of The Psychophysics Toolbox in

MATLAB, with visual stimuli presented on a large LED monitor.

In all experiments, eye position and microphone data were recorded while

subjects performed silent, visually-guided saccade tasks. Experimental sessions were

carried out in a darkened, acoustically isolated chamber made anechoic with the

addition of acoustic damping wall tiles. Subjects were seated 70 cm from the screen,

and a chin rest was used to maintain head position and minimize movement.

Experimental sessions were subdivided into multiple runs, approximately 5 minutes

each. This provided subjects with the opportunity to take a brief break from the

experiment if needed to maintain alertness or to address any possible discomfort from

maintaining their posture. Each run typically consisted of approximately 125 trials and

fixation positions and saccade targets were presented in pseudorandom order.

Before each experimental session, the eye-tracking system was calibrated using

the calibration routine provided with the Eyelink system to register raw eye-tracking data

to gaze locations on the stimulus presentation screen. If the subject requested an

adjustment to the chin rest or left the recording chamber for a break, the eye-tracking

calibration was repeated. Before each run, the microphone system was calibrated to

ensure that each microphone had a frequency response that was similar to the pre-

recorded frequency response of the microphones when placed in a volume that

approximated the size and geometry of the human ear canal - a 3ml syringe cut to

accept the Etymotic earpieces. The syringe stopper was pulled to 1.25 cm^3 to

approximate the volume of the average adult human ear canal. A small amount of

gauze (.25cm^3) was added to the volume to emulate the attenuation caused by the soft

tissue of the ear canal. The calibration routine played tones from 10 to 1600Hz, at a

constant system output amplitude. As the purpose of this calibration was to compare

microphone function in a control volume with that in an earpiece just placed in a subject,

the weighting of the tones was not specifically calibrated. If the input-output results of

the same tone sequences were consistent between ears and matched the overall shape

of the syringe calibration curves, microphone placement was considered successful.

No sounds were delivered during periods of experimental data collection.

<u>Task descriptions</u>

All tasks followed the same stimulus timing sequence: initial fixation points were

displayed on screen for 750ms and then removed as the saccade targets were

presented for 750ms (Figure 1A). Fixation and target locations were indicated by green

dots. Subjects were instructed to fixate on the initial fixation locations until targets were

presented on the screen, then to saccade to the targets and fixate on the targets until

they changed from green to red for the last 100ms of the target presentation (the color

cue was intended to help subjects maintain fixation through the end of the target

presentation). Inter-trial-intervals were jittered 350±150ms. This was done to minimize

the potential impact of non-saccade related noise signals that may be periodic (i.e.

heartbeat, external acoustic and electromagnetic sources).

In the five-origin grid task (Figure 1B), participants performed saccades to

multiple targets from five different initial eye positions in a plus-shaped configuration at -

9º, 0º, and +9º horizontally and at -6 º, 0º, and 6 º of elevation as shown. Twenty five

saccade targets ranged from -18º to +18º in 9º degree increments horizontally and from

-12º to +12 º in 6 º increments vertically.

In the horizontal/vertical task (Figure 1D), participants performed saccades to

targets along the vertical and horizontal axes from a central fixation. Vertical targets

ranged from -12º to +12º in 3º increments and horizontal targets ranged from -18º to

+18º in 3º increments.

In the single-origin grid task (Figure 1C), participants made saccades to 24

distinct targets of varying vertical and horizontal placement combinations from a central

fixation. Horizontal location components ranged from -18º to +18º in 9º increments and

vertical location components ranged from -12º to +12º in 6º increments.

<u>Preprocessing analysis</u>

Saccade-microphone synchronization:

Microphone data was synchronized to the onset the saccade from the fixation point to

the target. This was defined based on the third derivative of eye position, or jerk. The

first peak in the jerk represents the moment when the change in the eye acceleration is

greatest. Prior to each differentiation, a lowpass discrete filter with a 7ms window was

used to smooth the data and reduce the effects of noise and quantization error. This

filter was normalized, such that its output to a constant series of values equaled those

values.

<u>Trial exclusion criteria:</u>

Trials were excluded based on saccade performance and the quality of microphone

recordings. Exclusion criteria used for eye tracking: 1) if subjects made a sequence of

two or more saccades to achieve the target; 2) if the eye tracking signal dropped out

during the trial (e.g. due to blinks); 3) if the eye movement was slow and drifting, rather

than a saccade; 4) if the saccade curved by more than 4.5º (subtended angle); or 5)

subjects failed to maintain 200ms of fixation before and after the saccade. 6) If eye

tracking dropped samples that prevented the calculation of the saccade onset time.   On

average these saccade-related exclusion criteria resulted in the exclusion of ~15% of

the trials.

Prior to any further analysis, microphone data was downsampled from 48 kHz to

2 kHz sampling rate to reduce processing time given that the previously observed eye-

movement related signals of interest are well below 100 Hz (Gruters et al., 2018).

Exclusion based on noise in the microphone recordings was minimal. Within each block

of trials, the mean and standard deviation of the RMS values for each trial was

calculated. Individual trials were excluded if the microphone signal on that trial

contained any individual values that were more than 10 standard deviations away from

that mean. This typically resulted in the exclusion of a further ~1.4% of the trials, after

application of the eye position screen described above. On average, about 1200 trials

were included for each subject and task condition. See Supplementary Table for a full

breakdown across subjects and tasks.

<u>Z scoring</u>

To facilitate comparison across subjects, sessions, and experiments, all microphone

data reported in this study was z-scored within blocks and prior to the application of the

exclusion criteria described above. The mean and standard deviation of the

microphone values in a window -150 to -120 ms prior to saccade onset were used as

the normalization baseline period.

<u>EMREO intensity estimates.</u>

In addition to the Z scoring, we also calculated the peak-equivalent sound pressure

levels associated with EMREOs. EMREOs occur in a frequency range (~40 Hz) for

which the gain of the ER10B+ recording microphone is not flat, so standard conversions

from voltage-to-pressure do not apply. Rather, we used a Bruel & Kjaer sound level

meter (Model 2245) which has a roughly flat frequency response 20 Hz to 20 kHz (Z-

weighting). Using this meter, we measured the sound pressure levels associated with

40 Hz pure tones generated by Etymotic ER-2 transducers coupled in the earbud with

the ER10B+ microphone, which recorded these tones simultaneously. These

measurements were made in a short length of tygon tubing simulating the approximate

volume and sound-absorption properties of the human ear. This procedure provided a

voltage-to-sound-pressure-level mapping in the 40 Hz frequency range. We then used

these values to estimate the peak-equivalent sound pressure level of the EMREO signal

for selected targets in individual subjects (Supplementary Figure 1). By this technique,

EMREOs ranged in intensity from ~49 to 64 dB SPL, with a mean of 56 dB SPL. These

values closely corresponding with estimates using either the same (King et al., 2023b)

or a different method (Gruters et al., 2018).

<u>Regression Analyses</u>

Regression was used to assess how EMREOs vary with both eye position and

eye movement. Linear methods were used as a conservative approach providing a

lower bound on the ability to account for the variation in the microphone signal on the

basis of eye movement parameters.

The microphone signal at each moment in time *Mic(t)* was fit as follows:

$Mic(t) = B_H(t)\ H + B_{\Delta H}(t)\ \Delta H + B_V(t)\ V + B_{\Delta V}(t)\ \Delta V + C(t)$ *(1)*

where *H* and *V* correspond to the initial horizontal and vertical eye position and *ΔH* and

ΔV correspond to the respective changes in position associated with that trial. The

slope coefficients $B_H,\ B_{\Delta H},\ B_V,\ B_{\Delta V}$ are time-varying and reflect the dependence of the

microphone signal on the respective eye movement parameters. The term *C(t)*

contributes a time-varying "constant" independent of eye movement metrics, and can be

thought of as the best fitting average oscillation across all eye positions and

displacements.

The term *C(t)* was included for all regressions, but other parameters were

omitted when not relevant. Specifically, for the single-origin grid tasks and horizontal-

vertical tasks, the model used vertical and horizontal saccade displacement ($B_{\Delta H}(t)\ \Delta H$,

$B_{\Delta V}(t)\ \Delta V)$ as regression variables but not $B_H(t)\ H$ or $B_V(t)\ V$ as initial position did not

vary for those tasks. The analysis produced values for the intercept and variable

weights (or slopes), their 95% confidence intervals, R^2, and p-value for each time point.

For most analyses, the measured eye positions/changes-in-eye positions were

used as the independent variables, so as to incorporate any variability due to scatter in

fixation or saccade endpoint. For the target readout analysis described in Figures 6 and

7, the horizontal and vertical positions of the targets, rather than the associated eye

movements, were used.

<u>Data Availability</u>

The data and analysis code for this paper will be made available on publication on Duke

University's Research Data Repository. (citation to be added).

References

Abbasi H, King CD, Lovich S, Röder B, Groh JM, Bruns P (2023) Audiovisual temporal recalibration modulates eye movement-related eardrum oscillations. International Multisensory Research Forum.

Bala AD, Takahashi TT (2000) Pupillary dilation response as an indicator of auditory discrimination in the barn owl. J Comp Physiol [A] 186:425-434.

Barczak A, Haegens S, Ross DA, McGinnis T, Lakatos P, Schroeder CEJCr (2019) Dynamic modulation of cortical excitability during visual active sensing. 27:3447-3459. e3443.

Blauert J (1997) Spatial hearing. Cambridge, MA: MIT Press.

Boucher L, Groh JM, Hughes HC (2001) Afferent delays and the mislocalization of perisaccadic stimuli. Vision Research 41:2631-2644.

Bröhl F, Kayser C (2023) Detection of spatially-localized sounds is robust to saccades and concurrent eye movement-related eardrum oscillations (EMREOs). BioRxiv:2023.2004.2017.537161.

Brownell WE, Bader CR, Bertrand D, de Ribaupierre Y (1985) Evoked mechanical responses of isolated cochlear outer hair cells. Science 227:194-196.

Bulkin DA, Groh JM (2012a) Distribution of eye position information in the monkey inferior colliculus. J Neurophysiol 107:785-795.

Bulkin DA, Groh JM (2012b) Distribution of visual and saccade related information in the monkey inferior colliculus. Front Neural Circuits 6:61.

Caruso VC, Pages DS, Sommer MA, Groh JM (2019) Compensating for a shifting world: A quantitative comparison of the reference frame of visual and auditory signals across three multimodal brain areas. bioRxiv:669333.

Caruso VC, Pages DS, Sommer MA, Groh JM (2021) Compensating for a shifting world: evolving reference frames of visual and auditory signals across three multimodal brain areas. J Neurophysiol 126:82-94.

Cho NH, Ravicz ME, Puria S (2023) Human middle-ear muscle pulls change tympanic-membrane shape and low-frequency middle-ear transmission magnitudes and delays. Hear Res 430:108721.

Cohen YE, Andersen RA (2000) Reaches to sounds encoded in an eye-centered reference frame. Neuron 27:647-652.

Cooper NP, Guinan JJ, Jr. (2006) Efferent-mediated control of basilar membrane motion. J Physiol 576:49-54.

Delano PH, Elgueda D, Hamame CM, Robles L (2007) Selective attention to visual stimuli reduces cochlear sensitivity in chinchillas. J Neurosci 27:4146-4153.

Ege R, Opstal AJV, Van Wanrooij MM (2018) Accuracy-Precision Trade-off in Human Sound Localisation. Scientific reports 8:16399.

Fu KM, Shah AS, O'Connell MN, McGinnis T, Eckholdt H, Lakatos P, Smiley J, Schroeder CE (2004) Timing and laminar profile of eye-position effects on auditory responses in primate auditory cortex. J Neurophysiol 92:3522-3531.

Fuchs AF, Kaneko CR, Scudder CA (1985) Brainstem control of saccadic eye movements. Annu Rev Neurosci 8:307-337.

Galambos R (1956) Suppression of auditory nerve activity by stimulation of efferent fibers to cochlea. J Neurophysiol 19:424-437.

Gallagher L, Diop M, Olson ES (2021) Time-domain and frequency-domain effects of tensor tympani contraction on middle ear sound transmission in gerbil. Hear Res 405:108231.

Gelfand SA (1984) The contralateral acoustic reflex. . In: The acoustic reflex: Basic principles and clinical applications pp 137-186. New York, NY: Academic Press.

Groh JM (2014) Making space: how the brain knows where things are. Cambridge, MA: Harvard University Press.

Groh JM, Sparks DL (1992) Two models for transforming auditory signals from head-centered to eye- centered coordinates. Biological Cybernetics 67:291-302.

Groh JM, Trause AS, Underhill AM, Clark KR, Inati S (2001) Eye position influences auditory responses in primate inferior colliculus. Neuron 29:509-518.

Gruters KG, Murphy DLK, Jenson CD, Smith DW, Shera CA, Groh JM (2018) The eardrums move when the eyes move: A multisensory effect on the mechanics of hearing. Proc Natl Acad Sci U S A 115 E1309-E1318.

Guinan JJ, Jr. (2006) Olivocochlear efferents: anatomy, physiology, function, and the measurement of efferent effects in humans. Ear Hear 27:589-607.

Guinan JJ, Jr. (2014) Cochlear mechanics, otacoustic emisssions, and medial olivocochlear efferents: Twenty years of advances and controversies along with areas ripe for new work. In: Perspectives on Auditory Research, pp 229-246. New York: Springer.

Guthrie BL, Porter JD, Sparks DL (1983) Corollary discharge provides accurate eye position information to the oculomotor system. Science 221:1193-1195.

Harkrider AW, Bowers CD (2009) Evidence for a cortically mediated release from inhibition in the human cochlea. J Am Acad Audiol 20:208-215.

Hartline PH, Vimal RL, King AJ, Kurylo DD, Northmore DP (1995) Effects of eye position on auditory localization and neural representation of space in superior colliculus of cats. Exp Brain Res 104:402-408.

Hebrank J, Wright D (1974a) Are two ears necessary for localization of sound sources on the median plane? J Acoust Soc Am 56:935-938.

Hebrank J, Wright D (1974b) Spectral cues used in the localization of sound sources on the median plane. J Acoust Soc Am 56:1829-1834.

Hung IJ, Dallos P (1972) Study of the acoustic reflex in human beings. I. Dynamic characteristics. J Acoust Soc Am 52:1168-1180.

Jay MF, Sparks DL (1984) Auditory receptive fields in primate superior colliculus shift with changes in eye position. Nature 309:345-347.

Jay MF, Sparks DL (1987a) Sensorimotor integration in the primate superior colliculus. II. Coordinates of auditory signals. J Neurophysiol 57:35-55.

Jay MF, Sparks DL (1987b) Sensorimotor integration in the primate superior colliculus. I. Motor convergence. J Neurophysiol 57:22-34.

King C, Lovich S, Kaylie D, Shera C, Groh J (2023a) Measuring the impact of auditory system impairments on eye-movement-related eardrum oscillations (EMREOs). . Association for Research in Otolaryngology Abstracts Orlando, FL, February 2023

King CD, Lovich SN, Murphy DL, Landrum R, Kaylie D, Shera CA, Groh JM (2023b) Individual similarities and differences in eye-movement-related eardrum oscillations (EMREOs). BioRxiv:2023.2003.2009.531896.

Köhler MHA, Weisz N (2023) Cochlear Theta Activity Oscillates in Phase Opposition during Interaural Attention. J Cogn Neurosci 35:588-602.

Köhler MHA, Demarchi G, Weisz N (2021) Cochlear activity in silent cue-target intervals shows a theta-rhythmic pattern and is correlated to attentional alpha and theta modulations. BMC biology 19:48.

Lee J, Groh JM (2012) Auditory signals evolve from hybrid- to eye-centered coordinates in the primate superior colliculus. J Neurophysiol 108:227-242.

Leszczynski M, Bickel S, Nentwich M, Russ BE, Parra L, Lakatos P, Mehta A, Schroeder CE (2023) Saccadic modulation of neural excitability in auditory areas of the neocortex. Curr Biol 33:1185-1195 e1186.

Liberman MC, Guinan JJ, Jr. (1998) Feedback control of the auditory periphery: anti-masking effects of middle ear muscles vs. olivocochlear efferents. J Commun Disord 31:471-482; quiz 483; 553.

Lovich SN, King CD, Murphy DLK, Abbasi H, Bruns P, Shera CA, Groh JM (2023) Conserved features of eye movement related eardrum oscillations (EMREOs) across humans and monkeys. Philosophical transactions of the Royal Society of London Series B, Biological sciences In press.

Macpherson EA, Sabin AT (2013) Vertical-plane sound localization with distorted spectral cues. Hear Res 306:76-92.

Maier JX, Groh JM (2010) Comparison of gain-like properties of eye position signals in inferior colliculus versus auditory cortex of primates. Frontiers in Integrative Neuroscience 4:121-132.

Mendelson ES (1957) A sensitive method for registration of human intratympanic muscle reflexes. J Appl Physiol 11:499-502.

Metzger RR, Mullette-Gillman OA, Underhill AM, Cohen YE, Groh JM (2004) Auditory saccades from different eye positions in the monkey: implications for coordinate transformations. J Neurophysiol 92:2622-2627.

Middlebrooks JC, Green DM (1991) Sound localization by human listeners. Annu Rev Psychol 42:135-159.

Monti MM (2011) Statistical Analysis of fMRI Time-Series: A Critical Review of the GLM Approach. Frontiers in human neuroscience 5:28.

Mukerji S, Windsor AM, Lee DJ (2010) Auditory brainstem circuits that mediate the middle ear muscle reflex. Trends Amplif 14:170-191.

Mullette-Gillman OA, Cohen YE, Groh JM (2005) Eye-centered, head-centered, and complex coding of visual and auditory targets in the intraparietal sulcus. J Neurophysiol 94:2331-2352.

Mullette-Gillman OA, Cohen YE, Groh JM (2009) Motor-related signals in the intraparietal cortex encode locations in a hybrid, rather than eye-centered, reference frame. Cerebral Cortex 19:1761-1775.

O'Connell MN, Barczak A, McGinnis T, Mackin K, Mowery T, Schroeder CE, Lakatos PJI (2020) The role of motor and environmental visual rhythms in structuring auditory cortical excitability. 23:101374.

Populin LC, Tollin DJ, Yin TC (2004) Effect of eye position on saccades and neuronal responses to acoustic stimuli in the superior colliculus of the behaving cat. J Neurophysiol 92:2151-2167.

Porter KK, Metzger RR, Groh JM (2006) Representation of eye position in primate inferior colliculus. J Neurophysiol 95:1826-1842.

Porter KK, Metzger RR, Groh JM (2007) Visual- and saccade-related signals in the primate inferior colliculus. Proceedings of the National Academy of Sciences of the United States of America 104:17855-17860.

Russo GS, Bruce CJ (1994) Frontal eye field activity preceding aurally guided saccades. J Neurophysiol 71(3):1250-1253.

Schlebusch SN, Cooper MW, Kaylie DM, King CD, Murphy DLK, Shera CA, Groh JM (2019) Changes in saccade-related eardrum oscillations after surgical denervation of the stapedius muscle. Soc Neurosci Abstr.

Schlebusch SN, Cooper MW, Kaylie DM, King CD, Murphy DLK, Shera CA, Groh JM (2020) Changes in saccade-related eardrum oscillations after surgical denervation of the stapedius muscle. . Association for Research in Otolaryngology Abstracts.

Sparks DL, Mays LE, Porter JD (1987) Eye movements induced by pontine stimulation: interaction with visually triggered saccades. J Neurophysiol 58:300-318.

Srinivasan S, Keil A, Stratis K, Woodruff Carr KL, Smith DW (2012) Effects of cross-modal selective attention on the sensory periphery: cochlear sensitivity is altered by selective attention. Neuroscience 223:325-332.

Srinivasan S, Keil A, Stratis K, Osborne AF, Cerwonka C, Wong J, Rieger BL, Polcz V, Smith DW (2014) Interaural attention modulates outer hair cell function. Eur J Neurosci 40:3785-3792.

Stricanne B, Andersen RA, Mazzoni P (1996) Eye-centered, head-centered, and intermediate coding of remembered sound locations in area LIP. J Neurophysiol 76:2071-2076.

Takahashi M, Shinoda Y (2018) Brain Stem Neural Circuits of Horizontal and Vertical Saccade Systems and their Frame of Reference. Neuroscience 392:281-328.

Walsh KP, Pasanen EG, McFadden D (2014) Selective attention reduces physiological noise in the external ear canals of humans. I: auditory attention. Hear Res 312:143-159.

Walsh KP, Pasanen EG, McFadden D (2015) Changes in otoacoustic emissions during selective auditory and visual attention. J Acoust Soc Am 137:2737-2757.

Werner-Reiss U, Kelly KA, Trause AS, Underhill AM, Groh JM (2003) Eye position affects activity in primary auditory cortex of primates. Current Biology 13:554-562.

Wittekindt A, Kaiser J, Abel C (2014) Attentional modulation of the inner ear: a combined otoacoustic emission and EEG study. J Neurosci 34:9995-10002.

Zella JC, Brugge JF, Schnupp JW (2001) Passive eye displacement alters auditory spatial receptive fields of cat superior colliculus neurons. Nat Neurosci 4:1167-1169.

Zwiers MP, Versnel H, Van Opstal AJ (2004) Involvement of monkey inferior colliculus in spatial hearing. J Neurosci 24:4145-4156.

Supplementary Materials

Supplementary Table – summary of included data

Supplementary Figure 1

Supplementary Figure 2 – supplement to Figure 2

Supplementary Figure 3 – supplement to Figure 3

Supplementary Figure 4 – supplement to Figure 5

Supplementary Figure 5 – supplement to Figure 6

Supplementary Figure 6 – supplement to Figure 7

Five Origin Grid Task

Subjects	Number of days	Number of runs	Number of trials	N trials included after saccade screens	N trials included after saccade and microphone noise screens	Overall % included	Overall % excluded due to microphone noise screens
S11	5	17	2095	1905	1833	87.5%	3.4%
S13	4	16	1958	1608	1593	81.4%	0.8%
S45	4	16	1960	1687	1676	85.5%	0.6%
S71	5	25	3055	2618	2606	85.3%	0.4%
S56	2	18	2205	1745	1717	77.9%	1.3%
S66	4	16	1970	1498	1470	74.6%	1.4%
S75	3	15	1857	1377	1361	73.3%	0.9%
S77	7	32	3969	3358	3325	83.8%	0.8%
S78	4	26	3115	2363	2357	75.7%	0.2%
S80	3	10	1233	1060	1041	84.4%	1.5%
Average across subjects	**4.1**		**2341.7**		**1897.9**	**80.9%**	**1.1%**

Single Origin Grid Task

Subjects	Number of days	Number of runs	Number of trials	N trials included after saccade screens	N trials included after saccade and microphone noise screens	Overall % included	Overall % excluded due to microphone noise screens
S05	2	10	1403	1327	1299	92.6%	2.0%
S11	3	12	1615	1476	1416	87.7%	3.7%
S13	2	8	1064	906	891	83.7%	1.4%
S15	2	8	1217	1048	1037	85.2%	0.9%
S17	2	9	1182	1057	1028	87.0%	2.5%
S24	1	7	490	395	391	79.8%	0.8%
S45	1	9	667	455	443	66.4%	1.8%
S52	2	9	1116	1020	1002	89.8%	1.6%
S71	2	10	1208	1094	1090	90.2%	0.3%
S73	2	10	1172	1005	996	85.0%	0.8%
Average across subjects	**1.9**		**1113.4**		**959.3**	**84.7%**	**1.6%**

Horizontal-Vertical Task

Subjects	Number of days	Number of runs	Number of trials	N trials included after saccade screens	N trials included after saccade and microphone noise screens	Overall % included	Overall % excluded due to microphone noise screens
S05	1	4	488	455	445	91.2%	2.0%
S11	3	14	1491	1346	1279	85.8%	4.5%
S13	2	12	1165	976	968	83.1%	0.7%
S15	2	11	1094	844	825	75.4%	1.7%
S17	1	4	445	399	394	88.5%	1.1%
S24	1	5	509	436	434	85.3%	0.4%
S45	3	18	1307	1082	1054	80.6%	2.1%
S52	1	4	439	411	406	92.5%	1.1%
S71	2	8	747	649	647	86.6%	0.3%
S73	3	10	986	760	757	76.8%	0.3%
Average across subjects	**1.9**		**867.1**		**720.9**	**84.6%**	**1.4%**

Average across subjects and tasks	2.6	1440.7		1192.7	83.4%	1.4%

Supplementary Table. The numbers of days, blocks, and trials recorded and included/excluded based on saccade parameters and/or microphone noise across the different tasks.

Supplementary Figure 1. *Sound level of EMREOs associated with the 18 degree contralateral or ipsilateral target locations for individual subjects tested in the five-origin grid task. The origin for these data was the (0, -6) origin which corresponds to the conditions tested in our earlier study (Gruters et al., 2018). See* "Methods: EMREO intensity estimates" *for details.*

Supplementary Figure 2*. Grand average EMREOs recorded during the 5-origin grid task in right ears of ten subjects. Same format as Figure 2: Each panel is the grand average EMREO signal (average of the individual subject averages) that is generated when a saccade is made to that location on the screen e.g. the top right*

panel involves saccades to the top right target location. The color and line styles of each trace correspond to the initial fixation point as indicated by boxes of the same color and line style; e.g. all red oscillations are generated during a simultaneous saccade that originated from the right fixation point.

Supplementary Figure 3. *Grand average EMREOs as a function of target location with respect to the fixation position, for N=10 right ears. Same format as Figure 3. The data shown are the same as (a subset of) those shown in Supplementary Figure 2, but here each panel location corresponds to a particular target location defined with respect to the fixation point. The color/linestyle indicate the associated fixation position, as in Supplementary Figure 2.*

single origin grid task
horiz/vert task
p<0.05

a S05 LE

b S11 LE

c S13 LE

d S15 LE

e S17 LE

f S24 LE

g S45 LE

h S52 LE

i S71 LE

j S73 LE

Horiz. eye displacement
Mic SD/deg

Time re: saccade onset (ms)

Compare fits   -5 to 70 ms

k Horiz. eye displacement

Number of ears

Left ears
Right ears

Cross-task comparison R²

l Vert. eye displacement

m Constant

Supplementary Figure 4. Cross-task comparison for individual subjects and at the population level. A-J. Regression coefficients for horizontal eye displacement in the single-origin grid task (black) vs. the horizontal-vertical task (green) for individual subject left ears, similar to Figure 4A. K-L. All three coefficient values (horizontal eye displacement, vertical eye displacement, and constant terms) were then (separately) regressed against each other for the two tasks during the time window -5 to 70 ms (grey boxes in panels A-J), and the distribution of R^2 values across subjects and ears are shown as indicated. There is excellent cross task correspondence for the horizontal eye displacement signal at the individual level and somewhat more variable correspondence for the vertical eye displacement and constant terms, consistent with the population results shown in Figure 5.

Supplementary Figure 5. Same as Figure 6, but for right ear data. See main text for details.

Supplementary Figure 6. Target readout results for individual subjects following the population results shown in Figure 7J. The readout was computed based on the difference between the two ears for the horizontal dimension and the average between the two ears for the vertical dimension. Panels A-J show the results for each subject. Panel K shows the distribution of slope values relating the actual vs. read-out target locations in the horizontal and vertical dimensions (green: horizontal; brown: vertical). All values were statistically significant p<0.05 in the horizontal dimension and most were significant in the vertical dimension (darker shade of brown). The horizontal values are comparable to the population values indicated in Figure 7H, and the vertical are comparable to Figure 7K. Panel L shows the average standard deviations of the difference between the read-out location and the actual target when the readout is computed from individual trial waveforms.