1
2

Conformational changes in the essential *E. coli* septal cell wall synthesis complex suggest an activation mechanism

Brooke M. Britton[1*], Remy A. Yovanno[1*], Sara F. Costa[2], Joshua McCausland[1], Albert Y. Lau[1],
Jie Xiao[1#], Zach Hensel[2#]

1. Department of Biophysics and Biophysical Chemistry, Johns Hopkins School of Medicine, 725
N. Wolfe St, Baltimore, MD, 21205, U.S.A.

2. ITQB NOVA, Universidade NOVA de Lisboa, Lisbon, Av. da República, 2780-157 Oeiras,
Portugal.

[*]: These two authors contributed equally

[#]: Correspondences should be addressed to xiao@jhmi.edu and zach.hensel@itqb.unl.pt

ABSTRACT

The bacterial divisome, a macromolecular machine that is composed of more than thirty proteins

in *E. coli*, orchestrates the essential process of cell wall constriction during cell division. Novel

antimicrobial strategies can target protein-protein interactions within the divisome and will benefit

from insights into divisome structure and dynamics. In this work, we combined structure prediction,

molecular dynamics simulation, single-molecule imaging, and mutagenesis to construct a model

of the core complex of the *E. coli* divisome composed of the essential septal cell wall synthase

complex formed by FtsW and FtsI, and its regulators FtsQ, FtsL, FtsB, and FtsN. We observed

extensive interactions in four key regions in the periplasmic domains of the complex. FtsQ, FtsL,

and FtsB scaffold FtsI in an extended conformation with the FtsI transpeptidase domain lifted

away from the membrane through interactions among the C-terminal domains. FtsN binds

between FtsI and FtsL in a region rich in residues with superfission (activating) and dominant

negative (inhibitory) mutations. Mutagenesis experiments *in cellulo* and *in silico* revealed that the

essential domain of FtsN functions as a tether to tie FtsI and FtsL together, impacting interactions

between the anchor-loop of FtsI and the putative catalytic region of FtsW, suggesting a

mechanism of how FtsN activates the cell wall synthesis activities of FtsW and FtsI.

INTRODUCTION

Bacterial cell division is an essential process and a frequent target for novel antibiotics [1-
3]. The cytoskeletal protein FtsZ defines the future division site [4] and recruits more than thirty
proteins, many of them cell-wall enzymes and regulators. Collectively termed the divisome [5,6],
these proteins orchestrate cell-wall constriction during cell division. In *E. coli*, the essential septal
peptidoglycan (sPG) polymerase FtsW and its cognate transpeptidase FtsI form the sPG
synthase complex FtsWI and cooperate to synthesize new septal cell wall [7,8]. FtsWI activities are
regulated by a conserved subcomplex of transmembrane proteins FtsQ, FtsL, and FtsB (FtsQLB)
[9-12], and by FtsN in γ-proteobacteria [13]. However, how the septal cell wall synthesis activities of
FtsWI are regulated by FtsQLB and FtsN remains unclear.

In *E. coli*, FtsN has been regarded as the trigger of septum synthesis, as its arrival at the
division site coincides with the initiation of cell-wall constriction [13]. FtsN and FtsN-like proteins are
conserved in γ-proteobacteria and contain SPOR domains that bind denuded glycans [14]. In *E. coli,*
an essential, periplasmic segment of FtsN (FtsN^E) is sufficient to initiate constriction in the
absence of full-length FtsN [13,15,16]. Our recent single-molecule studies show that FtsN^E is part of
a processive sPG synthesis complex with active FtsWI and required to maintain the processivity
of the FtsWI complex [17,18]. Whether FtsQLB is also part of this complex and, if so, how it
associates with FtsWI, is unknown. *In vitro* experiments using purified proteins found that *E. coli*
FtsQLB inhibits FtsI activity [19], while *Pseudomonas aeruginosa* FtsQLB enhances FtsW activity [9].
In neither case did the addition of FtsN impact FtsI or FtsW activity. *In vivo* experiments showed
that mutations in FtsL, FtsW, and FtsI can be either dominant negative (DN) or superfission (SF,
bypassing FtsN and/or complementing DN mutations) [9,10,15,20,21]. These seemingly contradictory
observations suggest that FtsWI can transition between "on" and "off" states depending on the
environment and/or genetic mutations, and that FtsQLB and FtsN may play important roles in
shifting FtsWI between the two states [10,11,20]. However, it is unclear what conformations of FtsWI
correspond to "on" and "off" states and how FtsN shifts FtsWI between the two states.

A recent review synthesized evidence to date and used structure predictions to propose
a model of allosteric regulation of FtsWI in which multiple interactions cooperatively stabilize an
active conformation [22]. This work drew on structure predictions of complexes FtsWI, FtsQLB, and
FtsLWI using AlphaFold2 (AF2) [23]. Structure prediction was also recently applied to other
divisome interactions [20,24]. While AF2 makes predictions that are largely consistent with available
structural data and can accurately predict protein-protein interfaces [25-27], its predictions can fail to

distinguish between different states of the same complex [28] or describe the consequences of point

mutations [29].

In this work, we provided experimental evidence to show that FtsQLB forms a complex

with active FtsWI *in vivo* and conducted structure predictions for *E. coli* FtsWI, FtsWI in complex

with FtsQLB (FtsQLBWI), and for FtsQLBWI in complex with FtsN[E] (FtsQLBWIN). We subjected

these predicted structures to all-atom molecular dynamics (MD) simulation to observe dynamics

of protein-protein interfaces on the microsecond timescale. We observed extensive interactions

in the transmembrane and periplasmic domains of the FtsQLBWI and FtsQLBWIN complexes.

Next, we carried out mutagenesis and single-molecule tracking experiments to investigate these

observed interactions. Further MD simulations using DN and SF point mutations revealed critical

conformational changes at FtsQLB-FtsI protein-protein interfaces and near the putative catalytic

region of FtsW. Collectively, our results support a model in which FtsQLB scaffolds FtsWI in an

extended conformation poised for activation, with FtsN[E] functioning as a tether between FtsI and

FtsL to impact the interactions between the anchor-loop of FtsI and the catalytic region of FtsW,

suggesting an activation mechanism of FtsWI by FtsN.

RESULTS

Single-molecule tracking suggests that FtsQLB remains in complex with FtsWI on

both FtsZ and sPG tracks

Previously, using single-molecule tracking, we showed that FtsWI exhibits two

directionally moving subpopulations that reflect their activities in sPG synthesis [17,30]: a fast-moving

subpopulation (~30 nm/s) driven by FtsZ treadmilling dynamics and inactive in sPG synthesis (on

the "Z track") and a slow-moving subpopulation (~8 nm/s) independent of FtsZ treadmilling and

driven by active sPG synthesis (on the "sPG track"). We later observed that processive FtsN

exhibits only a slow-moving population, indicating association with FtsWI in an sPG synthesis

complex on the sPG track [18]. To investigate whether FtsQLB is part of the processive, active

complex on the sPG track and/or part of the inactive complex on the Z track, we ectopically

expressed a Halo-FtsB fusion protein that complemented an FtsB depletion strain (**Fig. S1A**,

Table S1, S2). We sparsely labeled Halo-FtsB with a fluorescent dye, JF646, and performed

single-molecule tracking as we previously described [17].

We observed that a large percentage of single Halo-FtsB molecules exhibited directional

motion (53.7% $\pm$ 2.7%, $\mu \pm$ s.e.m., n = 379 segments) (**Fig. 1B**, top, **Movie S1**, and **Movie S2**).

The velocity distribution of directional segments of Halo-FtsB single-molecule trajectories was

best described as the sum of fast- and slow-moving subpopulations (v_{fast} = 32.9 $\pm$ 5.2 nm/s, v_{slow}

= 8.2 $\pm$ 1.7 nm/s, p_{fast} = 38% $\pm$ 13%, μ $\pm$ s.e.m., *n* = 179 segments, **Fig. 1B**, bottom, **S1B**, and

Table S3). In the presence of fosfomycin, a drug that inhibits the synthesis of FtsW substrate

Lipid II, Halo-FtsB shifted to the fast-moving subpopulation (v_{fast} = 28.5 $\pm$ 2.9 nm/s, v_{slow} = 8.5$\pm$

3.0 nm/s, p_{fast} = 70% $\pm$ 10%, μ $\pm$ s.e.m., *n* = 218 segments, **Fig. 1B**, **S1C**, and **Table S3**). The

existence of two subpopulations of FtsB and the response to fosfomycin (**Fig. 1B**, **S1B**, **S1C**, and

Table S3) mirrored observations for FtsW in our previous study [17], but differed from those of FtsN,

which exhibited only slow motion [18]. Since FtsQ, FtsL, and FtsB form a stable heterotrimer [31],

these observations suggest that FtsQLB is in complex with both fast-moving, inactive FtsWI on

the Z track [17] and slow-moving, active FtsWI on the sPG track, with the latter also including FtsN

[18]. As such, FtsWI in complex with FtsQLB may be able to adopt both active and inactive states,

which are regulated by FtsN.

An FtsQLBWI model describes an extended conformation of FtsI and extensive

protein-protein interfaces

To gain insight into how the conformation of FtsWI can adopt both active and inactive

states for sPG synthesis in the FtsQLBWI complex, we used the ColabFold [32] implementation of

AlphaFold2 (AF2) [23] to predict the atomic structure of FtsQLBWI. In the prediction, we did not

utilize template coordinates, avoiding explicit dependence on homologous published structures.

The predicted structure of the complex showed an extended conformation of FtsI supported by

interactions between the membrane-distal C-terminal domains of FtsQ, FtsL, FtsB, and FtsI, as

well as interactions between transmembrane helices (TMH) of FtsL, FtsB, FtsW, and FtsI. These

interactions were predicted with high local confidence (pLDDT) except for extreme terminal

residues that have no predicted interactions (**Fig. S2A**). Furthermore, predicted DockQ values [33]

for protein-protein interfaces with individual subunits ranged from 0.64 to 0.73 (**Table S4**),

indicating accurate predictions of protein complexes [27]. Finally, predictions for homologous

complexes of FtsQLBWI in other diderm (gram negative) and monoderm (gram positive) species

were largely similar, with notable differences at less conserved C-terminal domains, such as the

different interactions reported for PASTA domains in FtsI homologs *Bacillus subtilis* Pbp2B [34] and

Streptococcus pneumonia Pbp2X [35] (**Fig. S2B**). These results suggest that AF2 can accurately

predict protein-protein interfaces of *E. coli* FtsQLBWI and homologous complexes, but it remains

essential to verify that predictions are consistent with experimental data and predictive in new

experiments.

Next, we built a system of FtsQLBWI in a lipid bilayer (simulation details in **Table S5**) and
performed a 1-µs all-atom molecular dynamics (MD) simulation to investigate the stability of
predicted protein-protein interfaces and conformational dynamics of the complex. We excluded
cytoplasmic, N-terminal regions of FtsQ (residues 1–19), FtsW (1–45), and FtsI (1–18) on the
basis of having very low pLDDT and/or lacking high-confidence predicted protein-protein
interactions. The MD simulation revealed a dynamic complex, with the FtsI transpeptidase
(TPase) domain tilting backward and the head domain wrapping around the FtsL helix (**Fig. 1C,**
S3 and **Movie S3**). Unless otherwise noted, we describe and depict structures after 1 µs of MD
hereafter. Throughout the complex, we observed extensive interactions reported in previous
experimental studies, such as leucine-zipper-like interactions between FtsL and FtsB helices [36]
(**Fig. S4A**), and interactions between the same FtsW and FtsI transmembrane helices observed
for the paralog RodA-PBP2 [37,38] (**Fig. S4B**). The resulting complex is also consistent with crystal
structures of *E. coli* FtsI (**Fig. S5A**) [39] and a partial FtsQ-FtsB complex (**Fig. S5B**) [40]. Finally, our
model of FtsQLBWI and its dynamics during 1 µs of MD are largely consistent with a recently
available CryoEM model of the orthologous complex in *Pseudomonas aeruginosa* [41], which we
discuss further in detail below.

To describe observed new interfaces in the context of previously defined domains, we
define four periplasmic interaction regions in FtsQLBWI (**Fig. 1C**, dashed boxes). These include
a "Truss" region that links FtsQ to FtsI via an extended β sheet, a "Hub" region that contains a
dense interaction network on both sides of FtsL and FtsB helices, a "Lid" region in which the
anchor domain of FtsI interacts with FtsW ECL4 (extracellular loop 4 containing the putative
catalytic residue FtsW^{D297} [42]), and a "Pivot" region where the first N-terminal periplasmic helix of
FtsI interacts with FtsW ECL4. FtsI is involved in all four regions, contacting FtsL and FtsB in the
membrane-distal periplasmic space and FtsW in the membrane, but essentially has no interaction
with FtsQ. These four regions are rich in residues that, when mutated, give rise to superfission
(SF) or dominant negative (DN) phenotypes, suggesting that they modulate FtsWI activity. While
our analysis below focuses on a subset of residues and protein-protein interfaces with well-
established phenotypes, we provide trajectories of top-ranked inter-residue hydrogen bonding
frequencies for each interface pair in the complex (**Supplementary Data 1**).

A collapsed structural model of FtsWI in the absence of FtsQLB reveals critical

interactions in the Pivot region

Our FtsQLBWI model and FtsW, FtsN, and FtsB single-molecule tracking results suggest
that FtsQLB forms a complex with FtsWI on the Z track that is poised for further activation by FtsN

on the sPG track. To investigate this possibility, we performed an MD simulation of an FtsWI

system without FtsQLB. In this simulation we observed flexibility in FtsI periplasmic domains in

the absence of FtsQLB. FtsI rotated about the Pivot region and collapsed onto the membrane,

where it remained until the end of the 1-µs simulation (**Fig. 1D** and **Movie S4**). While 1 µs is

insufficient to sample all conformations of FtsI in equilibrium, a collapsed FtsI conformation is

consistent with various flexible conformations of the paralogous *Thermus thermophilus* PBP2 in

the elongation complex with RodA in a previous cryo-EM study [38]. Notably, FtsI tilts in the opposite

direction relative to FtsW in comparison to a RodA-PBP2 crystal structure, with *E. coli* FtsI missing

a small loop that mediates RodA-PBP2 interaction in *T. thermophilus* PBP2 (**Fig. S5C**).

In the FtsQLBWI structure, the Pivot region consists of an FtsI helix (FtsI$^{D51–S61}$) on top of

the second short helix (FtsW$^{S260–G274}$) of FtsW ECL4 and is maintained by hydrogen bonding

between FtsI^{R60}-FtsW^{Q266} (**Fig. S6A**) and hydrophobic contact between FtsI^{L53} and FtsW^{M269} (**Fig.**

1E and **S7A**). These contacts were broken in FtsWI in the absence of FtsQLB, resulting in

separation of FtsI^{L53} and FtsW^{M269} and of FtsI^{R60} and FtsW^{Q266} in the FtsWI trajectory (**Fig. 1F**,

S6B, and **S7B**). Previous work showed that mutations to charged residues in this interface

(FtsI^{G57D}, FtsW^{M269K}) lead to the failure of cell wall constriction, while replacing with a hydrophobic

residue (FtsW^{M269I}) produced a SF variant of FtsW [20,21,43]. As the Pivot region conformation is

impacted by distal interactions with FtsQLB and contains both DN and SF mutations, it is likely

key for regulating FtsWI activity.

Truss region interactions stabilize an extended conformation of FtsI

Extreme C-terminal segments of FtsQLBI form an extended β-sheet

To identify what specific interactions between FtsI and FtsQLB support the extended

conformation of FtsI periplasmic domains, we examined the Truss region. The Truss region

consists of the C-terminal domains of FtsQLB and FtsI, with a striking feature that each one of

the four proteins contributes one β-strand to extend an FtsQ β-sheet (**Fig. 1C** and **2A**). The β-

sheet formed between FtsQ$^{A252–A257}$ (β12) and FtsB$^{T83–P89}$ (β1) has been previously reported [44]

(**Fig. S5B**), but that between FtsL$^{N116–Q120}$ (β1) and FtsI$^{E575–I578}$ (β16) has not been observed

experimentally and was only recently identified by structure prediction [45]. FtsL indirectly interacts

with FtsQ via FtsB within this β-sheet. The β-strand of FtsL$^{N116–Q120}$ maintains β dihedral angles

throughout the 1-µs simulation, while FtsI$^{E575–I578}$, which terminates the β-sheet, exhibits greater

flexibility (**Fig. S8A**, **B**). Previously, it was reported that the FtsI C-terminus (FtsI$^{I578–S588}$) was

cleaved, although it was unclear whether the cleavage occurred post-translationally, or during

sample processing [46,47]. We performed an additional 200-ns MD simulation of FtsQLBWI$^{\Delta I578–S588}$

and observed that this truncation did not disrupt β-strand formation of FtsL$^{N116–Q120}$ or FtsI$^{E575–V577}$

(**Fig. S8C**, **D**). In addition to the β-sheet linking these four proteins, two important contacts were

observed between FtsI^{R559} and FtsL^{E115}, and between FtsI^{R239} and FtsQ^{D249}, which may strengthen

C-terminal interactions between FtsI and FtsQLB (**Fig. S9**, **S10**, and **S11**).

β-strands deletions of FtsL and FtsI lead to varied degrees of defective cell division

To verify the importance of the β-sheet formed between FtsL and FtsI, we constructed

three FtsL mutants with varied degrees of C-terminal truncations (FtsL$^{\Delta 6}$, FtsL$^{\Delta 11}$, and FtsL$^{\Delta 16}$). All

three lack the C-terminal β-strand, and FtsL$^{\Delta 11}$ and FtsL$^{\Delta 16}$ additionally lack residues such as

FtsL^{E115} that interacts with FtsI (**Fig. 2B** and **S12**). FtsL$^{\Delta 6}$ and FtsL$^{\Delta 11}$ cells were filamentous,

indicating a division defect, but complemented an FtsL depletion strain at the highest induction

level tested (100 μM IPTG). FtsL$^{\Delta 16}$ cells remained filamentous and failed to rescue cell division

under the same condition (**Fig. 2C**, **S13A**, **B**, and **Table S6**). Fluorescently labeled FtsL$^{\Delta 6}$ and

FtsL$^{\Delta 11}$ (mVenus-FtsL) fusions showed clear, but significantly reduced, midcell localization, while

mVenus-FtsL$^{\Delta 16}$ cells exhibited only diffusive, cytoplasmic fluorescence (**Fig. 2D** and **S13C**, **Table**

S7, Table S8). These different localization patterns were not due to mVenus-FtsL truncations'

differential expression levels or their abilities to support cell division (**Fig. S13D, E**). These results

suggest that the interactions of the C-terminal β-strand of FtsL with FtsB and FtsI are important

for the recruitment of the complex, while further truncations impact essential divisome functions.

They are consistent with a previous observation that a C-terminal truncation mutant, FtsL$^{\Delta 114–121}$

(or FtsI$^{\Delta 7}$), complemented FtsL depletion, but was defective in co-immunoprecipitating with FtsQ

[48] and produced wrinkled colonies [49].

Next, to investigate the role of β-strand interaction between FtsL and FtsI, we constructed

two FtsI C-terminal truncations, FtsI$^{\Delta N579–S588}$ (or FtsI$^{\Delta 10}$), where only the disordered C-terminus

after the predicted β-strand was deleted, and FtsI$^{\Delta E575–S588}$ (or FtsI$^{\Delta 14}$), where the β-strand and the

C-terminus were both deleted (**Fig. 2B**, **S12**, and **Table S9**). We note that reported post-

translational cleavage of FtsI between FtsI$^{I578–S588}$ would be FtsI$^{\Delta 11}$ following this nomenclature,

removing one residue with a hydrophobic sidechain from the predicted β-strand that interacts with

FtsL. Both FtsI$^{\Delta 10}$ and FtsI$^{\Delta 14}$ complemented FtsI depletion (**Fig. S14A**). FtsI$^{\Delta 10}$ cells were longer

($l = 5.5 \pm 1.3$, $\mu \pm$ s.e.m. calculated from two biological replicates for n = 446 total cells) than wild-

type cells ($l = 3.9 \pm 0.04$, $\mu \pm$ s.e.m., n = 1215 cells) at the minimum expression level (1 μM IPTG).

However, extending the truncation in FtsI$^{\Delta14}$ corrected this defect, resulting in cell lengths

marginally shorter than wild-type (l = 3.3 ± 0.03, µ ± s.e.m., n = 487 cells) at the minimum

expression level (**Fig. 2E**, **S14B**, and **Table S9**). Western blot showed that FtsI$^{\Delta10}$ expression was

lower than that for either FtsI or FtsI$^{\Delta14}$ at equivalent induction levels, and that FtsI$^{\Delta14}$ expression

was marginally higher than that of FtsI (**Fig. S14C, D, Table S10**). Little change was observed in

midcell localization of FtsI across all conditions (**Fig. S14E, Table S11**). Together, these

observations indicate that interactions between C-terminal β-strands of FtsI and FtsL may

modulate FtsI stability and/or FtsWI activity.

Hub region interactions reveal inhibitory and activating interactions between FtsL and FtsI

The Hub region lies beneath the Truss region, encompassing the previously identified

CCD interface (Constriction Control Domain, containing multiple SF residues [9,10,50]) on one side

of FtsL and FtsB helices (**Fig. 3A** and **S15**), and the AWI interface (Activation of FtsW and FtsI,

containing multiple DN residues [51]) on the other side (**Fig. 3B**). As such, the Hub region may be

a key regulatory region for the activities of FtsWI. We analyzed predicted interactions in the Hub

region that were maintained during the final 500 ns of a 1-µs simulation of the FtsQLBWI system,

focusing on hydrogen bonds (as defined in **Methods**) and hydrophobic contacts monitored by

computing the distance between the side chain geometry center of each residue in a given pair.

The most prominent feature in the CCD interface of the Hub region is a network of residues

with polar sidechains extending through FtsQLB to FtsI. As shown in **Fig. 3A**, contacts between

FtsQ^{R196}-FtsB^{E69}, FtsQ^{R196}-FtsB^{E56} and FtsQ^{R213}-FtsB^{E56} connect FtsQ to FtsB. FtsB is connected

to FtsL through FtsB^{E56}-FtsB^{R70} and FtsB^{R70}-FtsL^{E88}. Finally, FtsL connects to FtsI through FtsL^{R82}-

FtsI^{S85} (**Fig. 3A** and **S16**). Mutations of residues in this region such as FtsB^{E56A}, FtsL^{E88K}, FtsL^{N89S},

FtsL^{G92D}, and FtsL^{H94Y} have been reported to be SF variants; cells expressing these variants are

shorter than wild-type cells and able to survive in the absence of FtsN [10,15]. Therefore, it is possible

that these interactions maintain FtsI in an inactive conformation, and that abolishing these

inhibitory interactions activates FtsI.

Interactions in the AWI interface are mainly between one face of the FtsL helix and two β-

strands at the neck of the FtsI head domain (**Fig. 3B**). A few hydrophobic residues, FtsI^{Y168}, FtsI^{V84},

and FtsI^{V86}, pack closely with hydrophobic residues FtsL^{I85} and FtsL^{L86}. Additionally, hydrogen

bonds were observed between FtsL^{R82}-FtsI^{S85} and between FtsL^{N83}-FtsI^{P87} (**Fig. S16**, hydrogen

bond frequencies for all protein-protein interfaces in the complex are available in **Supplementary**

Data 1). Previously both FtsL[R82E] and FtsL[L86F] were identified as DN mutants [10], suggesting that

these interactions may be required for maintaining the active conformation of FtsI.

MD simulation of the FtsI[R167S] SF variant to investigate an active conformation of FtsI

We reasoned that if predicted inhibitory interactions in the CCD region and activating

interactions in the AWI region play important roles in modulating FtsI activity, we may observe

corresponding conformational changes in the Hub region when some of these residues are

mutated. To examine interactions in the AWI region, we performed a 1-μs MD simulation of a SF

variant, FtsI[R167S] [17] (**Fig. S17A**). FtsI[R167] is next to hydrophobic residues FtsI[Y168], FtsI[V84], and FtsI[V86]

that interact with FtsL[I85] and FtsL[L86] (**Fig. 3B**). In the FtsI[R167S] simulation, the two β-strands at the

neck of the FtsI head domain pack closer to FtsL than in the WT (FtsQLBWI) simulation, as

measured by the sidechain geometry center distances between FtsI[V84] and FtsL[I85] (**Fig. 3C**, **D**,

and **S18A**, **B**; $d = 6.4 \pm 0.6$ Å for WT and 5.4 ± 0.5 Å for FtsI[R167S] complex). Consistent with this

measurement, the solvent-accessible surface area of FtsL residues with hydrophobic sidechains

in the helix linking the Lid and Hub regions was reduced from 534 Å^2 to 501 Å^2.

Interestingly, we observed drastic conformational changes in the turn connecting the two

FtsL helices immediately above the AWI region of the Hub. As shown in **Fig. 3E**, this

conformational change corresponds to a transition in the distribution of dihedral angles for FtsL[G92]

and with FtsL[H94] shifting from α-helical to β-strand dihedral angles, while the WT complex retains

α-helical dihedral angles. Note that both FtsL[H94Y] and FtsL[G92D] are SF variants, suggesting that

the conformation of this region plays an important role in FtsWI regulation. Taken together, the

FtsI[R167S] SF complex simulation suggests that replacing the charged arginine residue with

uncharged serine in FtsI[R167S] strengthens hydrophobic interactions between FtsI and FtsL in the

AWI region and results a conformational change in FtsL helices. These changes may correspond

to an activated conformation of FtsWI, leading to the SF phenotype of FtsI[R167S].

FtsB remains associated with FtsWI in the FtsI[R167S] SF background

In our previous single-molecule tracking experiments, we observed that FtsWI shifts to the

slow-moving, active population on the sPG track in the FtsB[E56A], FtsI[R167S], and FtsW[E289G] SF

backgrounds [17]. We asked whether activation of FtsWI was due to the dissociation of FtsQLB

from FtsWI on the Z track to relieve an inhibitory effect as previously proposed [19], or if FtsQLB

remains in complex with active FtsWI on the sPG track to maintain its activities as recently

suggested by an *in vitro* study [9]. To distinguish between these two possibilities, we performed

single-molecule tracking of Halo-FtsB in the FtsI[R167S] SF background under a rich growth condition,

which was previously used to assess the effect on FtsW. As shown in **Fig. 3F**, we observed that

Halo-FtsB was best fit as a single, slow-moving, active population (v_{slow} = 8.0 ± 0.4 nm/s, μ ±

s.e.m., *n* = 112 segments, **Table S3**), just as we previously observed for FtsW. This result is

consistent with the expectation that FtsI[R167S] enhances the formation of the active FtsQLBWI

complex on the sPG synthesis track, and that FtsQLB remains in the complex with activated FtsWI,

as we previously described for FtsN [18].

Lid region reveals important interactions between the FtsI anchor domain and

292 **FtsW ECL4**

Position of FtsI anchor domain may modulate FtsW activity

The Lid region is located at the periplasmic face of the inner membrane and involves

interactions between a polar patch on the back of FtsL and FtsB [24], a loop in the FtsI anchor

domain (FtsI anchor-loop) and FtsW ECL4 (**Fig. 4A**, **S19**, and **S20**). Specifically, hydrogen bonds

between FtsB[K23]-FtsI[E219], FtsB[D35]-FtsL[R67], FtsL[R67]-FtsI[D220], and FtsL[E68]-FtsI[R207] persist during the

final 500 ns of FtsQLBWI simulation and fix the orientation of the FtsI anchor-loop with respect to

the FtsL and FtsB helices (**Fig. 4A** and **S19**), while hydrogen bonding between FtsL[R61], FtsW[N283],

and FtsW[S284] positions ECL4 directly beneath the anchor-loop (**Fig. 4B** and **S20**). These

interactions allow FtsI[Y214] in the FtsI anchor-loop to hydrogen bond with FtsW[E289], positioning

FtsW ECL4 beside a central pore containing the putative catalytic residue FtsW[D297] [20] (**Fig. 4B, C**,

and **Fig. S20**). Previous studies showed that FtsW[E289G] is a SF variant while FtsL[R61E] and FtsL[R67E]

are DN variants [24], indicating that altering the interactions in the Lid region can indeed render a

constitutively active or inactive complex.

MD simulations of Lid region DN and SF complexes reveal drastically different

interactions

Observing that Lid region interactions in FtsQLBWI are rich in SF and DN residues, we

hypothesize that local conformational changes resulting from these mutations could shed light on

the FtsW activation mechanism. We simulated the effects of introducing the FtsW[E289G] SF variant

and the FtsL[R61E] DN variant into FtsQLBWI. Both complexes adopt similar global structures after

1 μs of MD compared to WT FtsQLBWI (**Fig. S17B, C**), but differ in local positioning of the FtsI

anchor-loop relative to FtsW ECL4 (**Fig. 4D, E**). In the SF FtsW[E289G] complex, the FtsI anchor-

loop is rotated sideways toward FtsL and FtsW TMH1, opening a central cavity of FtsW (**Fig. 4D**).

This conformation is stabilized by backbone hydrogen bonding between FtsI[R213] and FtsW[L288] in

addition to maintaining contacts with FtsL (**Fig. S20**). In contrast, in the DN FtsL[R61E] complex, the

317 FtsI anchor-loop is directly above the FtsW cavity (**Fig. 4E**). This interaction is stabilized by

318 contacts between FtsI^{R216} and FtsW^{E289} and persists through ~72% of the last 500 ns of simulation

319 of the FtsL^{R61E} DN complex (**Fig. S21**). As we described above, the FtsI anchor-loop is

320 coordinated by its interactions with the polar patch on one side of FtsL helix and with FtsW ECL4,

321 which are reduced in the FtsL^{R61E} variant (**Fig. S17C**). Loss of the interaction between FtsL^{R61E}

322 and FtsI^{E219} is also accompanied by conformational changes in FtsI; the relative orientation of FtsI

323 and FtsW helices in the Pivot region changes, and the β-sheet in the anchor domain tilts towards

324 FtsL (**Fig. S17C**). We also observed that in the SF FtsW^{E289G} simulation FtsW TM7 tilts outward

325 and expose a large extracytoplasmic cavity (**Fig. 4D** and **S17B**) as reported for the RodA-PBP2

326 complex [38]. In simulations of both FtsWI in the absence of FtsQLB and FtsQLBWI with FtsL^{R61E},

327 FtsW^{Y379} in ECL5 flipped from its predicted position to fit between FtsW TM1 and TM2. This flip

328 was only observed in simulations of these inactive complexes, suggesting that FtsW-FtsL

329 interactions may influence conformation near the FtsW active site.

330 **Binding of FtsN^E reduces inhibitory interactions and strengthens activating**

331 **interactions in the FtsQLBWI complex**

332 Previous studies have shown that the effects of SF variants are independent of FtsN, *i.e.*

333 SF variants bypass the need for FtsN binding. Therefore, we reason that binding of FtsN to

334 FtsQLBWI could switch the complex into an active conformation similar to what we observed for

335 SF variants. To investigate the role of FtsN binding, we predicted the structure of FtsQLBWI with

336 the addition of FtsN$^{K58–V108}$*,* which encompasses FtsN^E (FtsN$^{L75–Q93}$ [15]) and a sufficient number of

337 surrounding periplasmic residues required to obtain a predicted structure. As with other

338 complexes, we performed 1 μs of MD to investigate interactions and dynamics of the predicted

339 complex of FtsQLBWI and FtsN$^{K58–V108}$ (FtsQLBWIN, **Movie S5**).

340 While termini of FtsN$^{K58–V108}$ did not form stable interactions on this timescale, the FtsN^E

341 region is bookended by two interfaces predicted with high confidence and stability in MD (**Fig.**

342 **5A**). First, FtsN$^{L75–P79}$ forms a polyproline II helix that binds the FtsI head domain (**Fig. S22A**).

343 This region consists of residues with high polyproline II helix propensity [52], which is a property

344 conserved for FtsN (**Fig. S22B**) in taxonomic families within the previously identified subgroups

345 characterized by a four-amino-acid deletion in RpoB [53,54]. However, we did not identify this motif

346 for FtsN outside of these subgroups (*e.g.* in *Pseudomonadaceae*) and predicted complexes with

347 both *P. aeruginosa* FtsI and PBP3x did not include this interaction (**Fig. S22C**), suggesting that

348 FtsN interactions and functions may vary between γ-proteobacteria. This observation is consistent

with the conditional essentiality of FtsN and the inability to detect an effect of FtsN on FtsQLBWI

or reconstitute FtsQLBWIN *in vitro* for *P. aeruginosa* [9] [41].

The second interaction region, FtsN$^{W83–L89}$, contains conserved residues FtsN^{W83}, FtsN^{Y85},

and FtsN^{L89}, which are also the most sensitive residues to alanine scanning mutagenesis [15]. This

region interacts with both the AWI interface of FtsL (FtsL^{L86}, FtsL^{A90}) and at the neck of the FtsI

head domain (FtsI^{V86} and FtsI^{Y168}), largely through hydrophobic interactions and also including

FtsB^{L53} (**Fig. 5A**, right). Importantly, the FtsL and FtsI residues bound by FtsN overlap with those

impacted in the FtsI^{R167S} SF variant (FtsL^{L86}, FtsI^{V86}, and FtsI^{Y168}), suggesting a shared mechanism

of FtsQLBWI activation. Furthermore, this FtsN-binding interface includes both FtsL^{A90} in the

hydrophobic interface and an FtsL^{E87}-FtsN^{R84} interaction, suggesting that DN variants FtsL^{E87K}

and FtsL^{A90E} are defective in FtsN^{E} binding. The solvent-accessible surface area of FtsL residues

with hydrophobic sidechains in this region was reduced from 534 Å^2 in WT to 383 Å^2, greater than

that in the FtsI^{R167S} simulation.

To validate the prediction of FtsN binding specificity, we additionally predicted structures

of FtsN$^{K58–V108}$ together with only FtsQLB or FtsWI. Despite relatively low prediction confidence,

the predicted location of FtsN$^{W83–L89}$ binding did not change (**Fig. S22D**). We also note that

predictions of full-length FtsN are possible for *E. coli* and homologous complexes such as for

Aliivibrio fischeri, but that we failed to identify other high-confidence binding interfaces (**Fig. S22E**).

Little global change was observed in simulations of FtsQLBWIN compared to FtsQLBWI

(**Fig. S17D**). However, MD revealed that the presence of FtsN^{E} triggered extensive local

conformational changes in all four interaction regions in FtsQLBWI. In the Truss region, the

FtsL^{E115}-FtsI^{R559} interaction was broken (**Fig. S9A** and **S10E**), which impacts interactions between

the FtsL β-strand and adjacent FtsB and FtsI β-strands. Several interactions involving polar

residues at the CCD interface (FtsL^{E98}-FtsI^{R246} and FtsL^{N89}-FtsI^{P170}) in the Hub region were nearly

completely abolished (**Fig. S16**), resulting in conformational change for FtsL^{H94} and FtsL^{G92} similar

to that observed for FtsQLBWI^{R167S} (**Fig. 5B**). In the Lid region, charged interactions between the

polar patch of FtsL and the back of the FtsI anchor domain (FtsL^{R67}-FtsI^{D225}, **Fig. 5C** and **S23**) are

present; FtsW^{E289} in ECL4 switches from interacting with FtsI^{Y214} to FtsI^{R213} in the anchor-loop,

resulting in conformation changes in the FtsI anchor-loop and FtsW ECL4 that alter the catalytic

cavity on the periplasmic face of FtsW relative to the conformer observed for FtsQLBWI (**Fig. 5D**).

This conformational change, coincident with loss of FtsW^{E289}-FtsI^{Y214} interaction, is similar to that

observed with the FtsW^{E289G} SF mutation (**Fig. 4D**, **S22F**). However, the same conformational

change did not occur in the FtsI^{R167S} simulation within 1 μs and it remains unclear if this specific

conformational change is necessary or sufficient for FtsW activation (**Fig. S22G**). In the

FtsQLBWIN simulation, we also observed that FtsW^{K370} flipped to form a salt bridge with FtsW^{D297}

within the first 200 ns that persisted to the end of the simulation (**Fig. 5D, Movie S5**). This was

also observed in FtsI^{R167S} and FtsW^{E289G} SF simulations, but not in the wild-type FtsQLBWI

simulation. The consistency between FtsN^E-induced conformational changes and those that

appeared in simulations of SF complexes with FtsI^{E289G} or FtsI^{R167S}, but not the wild-type

FtsQLBWI or the DN FtsL^{R61E} complex, suggests that FtsN^E binding in the Hub region has

allosteric effects on FtsW and/or FtsI activity in the FtsQLBWI complex.

Defining long-range interaction paths of FtsQLB complexes

Analyses described above allow us to identify critical local interactions that are modulated

by FtsN^E binding and may contribute to the activation of FtsWI, but do not reveal how local

interactions trigger distal conformational changes in the complex. To investigate this, we

compared long-range interaction pathways for wild-type and two active (FtsI^{R167S} or FtsN^E)

complexes, as well as for the FtsI^{R61E} DN complex. We quantified correlations between pairs of

residues using a dynamical network model [55]. Highly correlated residues in a simulation trajectory

of a complex reflect their coordinated motions, allowing us to construct long-range interaction

pathways. Because FtsQ is primarily connected to the rest of the complex through its interactions

with FtsB in the C-terminal Truss region, and because we wish to identify long range interactions

that affect the activities of FtsW and FtsI, we computed optimal paths through the Truss and Hub

regions between FtsQ and putative catalytic residues FtsW^{D297} [20] and FtsI^{S307} (**Fig. S24**). These

paths reveal dominant routes of communication between the proteins. The optimal path ensemble

between FtsQ and FtsW^{D297} for the DN FtsL^{R61E} simulation exhibits greater path density through

FtsB, while the other simulation systems exhibit greater path density through FtsL (**Fig. 5E and**

Fig. S24A). Interestingly, active FtsI^{R167S} and especially FtsN^E simulations included more FtsI

residues in optimal paths connecting FtsI to FtsW, which is consistent with our observation that

FtsN^E binds between FtsL and the FtsI head domain in the Hub region to enhance this interaction.

The optimal path ensemble between FtsQ and FtsI^{S307} extends through SF residues including

FtsB^{E56}, FtsL^{E88}, and FtsL^{H94}, and varied across different complexes (**Fig. S24B**).

Comparison with a cryo-EM model of the orthologous complex in *Pseudomonas aeruginosa*

The recently publication of the CryoEM structure of the FtsQLBWI complex in

Pseudomonas aeruginosa [41] makes it possible to address concerns about the physiological

relevance of molecular dynamics of predicted divisome structures in *E. coli.* We found that the

global conformations of the *E. coli* FtsQLBWI complex was remarkably similar to the *P.*

aeruginosa complex (**Fig. S25A**). The *P. aeruginosa* cryo-EM structure was reported to exhibit a

tilt in its FtsI TPase domain, relative to an AF2-predicted structure. We observed a similar tilt of

FtsI early in all simulations and measured an angle of only 3° between the *P. aeruginosa* cryo-

EM structure and the *E. coli* FtsQLBWI complex after 1-μs MD. Rapid tilt of the TPase domain

towards the membrane was observed within the first 200 ns of MD for simulations of all *E. coli*

complexes (**Fig. S25B**). Furthermore, *Käshammer et al.* described a change in the interface

between the FtsI anchor domain and FtsL, where FtsL-interacting residues in the FtsI anchor

domain shift from residues FtsI[203-206] in the AF2 prediction to residues FtsI[208–212] in the cryo-EM

structure. We observed a similar conformational change as an interface is formed between FtsI[223–228]

(equivalent to *P. aeruginosa* FtsI[209–215]) and FtsL[69–75] upon the addition of FtsN[E] (**Fig. S25C**).

Unlike in the *P. aeruginosa* structure, we did not observe a direct interaction between the FtsI

head and anchor domains. Instead, the conformer after 1-μs MD for the *E. coli* complex with FtsN

suggests that FtsN bridges the head and anchor domains. This may reflect differing roles and

interactions for FtsN in *E. coli* and *P. aeruginosa* divisome.

Discussion

In this work, we first used single-molecule tracking to determine that *E. coli* FtsQLB forms

a complex with both inactive FtsWI on the Z track and active FtsWI on the sPG track (**Fig. 1A**, **B**).

Building upon this result, we used structure prediction and molecular dynamic simulations to

model the structures of FtsQLBWI (**Fig. 1C**), FtsWI (**Fig. 1D**), and FtsQLBWIN (**Fig. 5A**). We then

examined the structural models by a set of mutations both experimentally (**Fig. 2C–E** and **3F**) and

computationally (**Fig. 3C–F**, **4C**, **5D–E**). Combining these structural models with observations

from MD and in biological experiments, we propose a regulatory mechanism in which FtsWI

activities depend on FtsQLB and are further activated by FtsN in *E. coli.*

In the absence of FtsQLB, FtsWI adopts flexible conformations with low GTase and TPase

activities because FtsWI lacks interactions in the Pivot and Lid regions that stabilize FtsWI

conformations required for processive sPG strand polymerization and crosslinking. The addition

of FtsQLB stabilizes an extended conformation of FtsI through the Truss region. This

conformation supports higher FtsWI activities compared to the background level of FtsWI alone,

and could be sufficient in species that are not dependent on FtsN. The binding of FtsN[E] both

weakens inhibitory interactions in the CCD interface and enhances activating interactions in the

AWI interface of the Hub region, collectively resulting in stronger association of the FtsI head

domain and FtsL. These changes promote conformations in which interactions between the FtsI

anchor-loop and FtsW ECL4 are modified to open the catalytic pore of FtsW. We propose that

this conformational change corresponds to the highest activities of the complex because it

remodels the catalytic cavity of FtsW to facilitate processive PG polymerization and to allow the

growing PG strand to reach the TPase domain of FtsI.

Further, we hypothesize that binding of the nascent PG strand to FtsI may be coupled with

a transient conformational change to an extended conformation compatible with PG crosslinking

as proposed for the RodA-PBP2 elongasome complex [38] and observed in our predicted structures

(**Fig. S3**). This mechanism differs in specific details from those proposed in recent work that also

drew from structure prediction [22,41,45]. Despite their differences, each of these hypotheses involves

allostery linking periplasmic interactions to distal active sites.

Roles of FtsQLB in FtsWI activation

The structural models of various complexes allowed us to clarify the seemingly

contradictory roles of FtsQLB in activating FtsWI. Using a thioester substrate of FtsI, Boes *et al.*,

found that adding purified *E. coli* FtsQ moderately inhibited FtsI TPase activity, while adding FtsL

and FtsB had little effect [19]. Marmont *et al.*, found that purified *P. aeruginosa* FtsL and FtsB

enhanced FtsW GTase activity as well as FtsQLB [9]. In neither case did the addition of FtsN impact

FtsI or FtsW activity. Our model of FtsQLBWI shows that FtsQ has few interactions with members

of the complex other than FtsB. Thus, a role of FtsQ is to stabilize the global conformation of the

complex via these interactions without directly impacting the activity of FtsW or FtsI. The major

role of FtsL and FtsB is to scaffold FtsWI in a conformation poised for activation by clamping both

the membrane-distal TPase domain of FtsI and membrane-embedded FtsW. Truncations of β-

strands in FtsL$^{\Delta 6}$, FtsL$^{\Delta 11}$, FtsL$^{\Delta 16}$, and FtsI$^{\Delta 10}$ were not lethal, but resulted in filamentous

phenotypes indicating defects in cell division. Stacked β-strands between FtsL, FtsB, and FtsQ

are consistent with the observation that C-terminal truncation of FtsL$^{\Delta 100-121}$ (or FtsL$^{\Delta 21}$) both

abolishes FtsL interaction with FtsQ and also results in a reduced level of full length FtsB [48]. A

mild cell division defect was observed for FtsI$^{\Delta 10}$ but not for FtsI$^{\Delta 14}$, which is likely caused by a

lower expression level of FtsI$^{\Delta 10}$ than FtsI$^{\Delta 14}$ under the same induction levels (**Fig. S14**). These

observations suggest a complex role for C-terminal FtsL-FtsI interaction in the Truss region and

are consistent with the observation that deletion of FtsI$^{G571-V577}$ appears to be fully functional [56].

Note that our FtsB single-molecule tracking data indicates that FtsB is associated with FtsWI on

both the Z-track and the sPG-track in *E. coli*, and hence whether an FtsWI complex alone is

physiologically relevant is unclear.

The proposed activation mechanism is consistent with previous studies

The proposed activation mechanism is remarkably consistent with what was previously

deduced by genetic studies: following recruitment by FtsQ, the activating signal goes from FtsB

to FtsL, then to FtsI, and finally to FtsW [10]. We suggest that the final activation step includes

remodeling the central cavity of FtsW to facilitate processive PG polymerization and crosslinking.

This step can be achieved by multiple means: SF variants in the Pivot (such as FtsW[M269I]), Hub

(such as FtsB[E56A], FtsL[H94Y], and FtsI[R167S]), and Lid (such as FtsI[K211I]) regions can correctly position

the FtsI anchor domain away from the FtsW pore, or directly remodel the structure defined by the

FtsI anchor-loop and FtsW ECL4 to open a channel in FtsW (such as FtsW[E289G]). However, short

circuiting this pathway removes potential points of regulation needed to coordinate cell wall

synthesis in space and time and in response to environmental conditions.

This activation mechanism is also consistent with previous mutagenesis studies, as our

results provide unprecedented details in potential conformational changes caused by previously

identified SF and DN mutants. For example, previously it was observed that substitutions

removing the negative charge of FtsW[E289] do not affect the function of FtsW while only FtsW[E289G]

is superfission [20]. This observation can be explained by the fact that the loss of the side chain of

FtsW[E289] and/or the flexibility of FtsW[G289], but not necessarily specific interactions, removes the

capping of the FtsW cavity and activates FtsW GTase activity. The observation that SF variant

FtsW[E289G] or FtsW[M269I], but not overexpression of FtsN, can rescue the double-DN mutant

FtsL[L86F/E87K] [10] is also consistent with our proposed activation mechanism and results. Since the

double-DN mutant FtsL[L86F/E87K] loses its ability to bind to FtsN to trigger conformational changes

that increase FtsWI activity via allosteric paths extending from the Hub through the Lid and Pivot

regions (**Fig. 5E**), it is expected that Lid and/or Pivot mutations could short-circuit this effect.

The activation mechanism we propose does not specifically involve FtsWI catalytic

residues. However, this model could be expanded to address possible regulation of active site

conformation and dynamics near catalytic residues such as FtsW[D297] [20]. For example, dynamics

in the vicinity of FtsW[D297] revealed potential roles of FtsW[K370] in addition to FtsW[Y379] in regulating

FtsW activity (**Fig. 5D**). In predictions, FtsW[K370] blocks a putative substrate channel [38], suggesting

that FtsW[K370] conformation can regulate substrate or product transport. We also did not address

in detail the hypothesis of a diprotomeric Fts[QLBWI]$_2$ complex or the role of cytoplasmic FtsL-

FtsW interactions [45]. Nevertheless, we observed stable FtsL-FtsW cytoplasmic interaction in our

simulations even though the N-terminal cytoplasmic tail of FtsW was truncated. Additionally, the

location of the FtsN^E binding site and conformational changes in FtsL and FtsB helices observed

in simulations with FtsN^E and FtsI^{R167S} are qualitatively compatible with the diprotomer model that

requires flexibility in this region.

Our simulations for all FtsQLBWI wild-type and mutant complexes, with and without FtsN,

rapidly attained a bent conformation similar to that observed for the *P. aeruginosa* cryo-EM

complex (**Fig. S25B**). Käshammer *et al.*, hypothesized that a transition from the observed bent

conformation to the AF2-predicted straight conformation could be associated with binding of

substrate or proteins such as FtsN. Our results here, when further combined with our previous

single-molecule imaging results for FtsN [18], demonstrate that FtsN^E is associated with FtsQLBWI

on the sPG track, with MD suggesting that FtsN activates the GTase activity of FtsW in the

observed bent conformation. We did not observe stable "straight" conformations in our simulations.

Our model also provides insight into conflicting reports about the role of FtsN in different

organisms. We observed that a shift in FtsI-FtsL interactions, analogous to that observed between

predicted and experimental structures in the *P. aeruginosa* cryo-EM study, only occurred upon

the addition of FtsN^E in the *E. coli* complex (**Fig. S25C**).

There are several limitations on our ability to definitively infer regulatory mechanisms from

our analysis of divisome dynamics. First, despite congruence with the *P. aeruginosa* cryo-EM

structure, we cannot rule out model-dependence or random effects. For this reason, we largely

focused on residues and protein-protein interfaces with well described phenotypes. Determining

the significance of previously undescribed interactions will require reproducing observations in

MD and formulating testable hypotheses. Second, 1 µs of equilibrium MD is insufficient to quantify

binding free energy or to have high confidence in new interfaces formed during simulation. Future

work can address this concern with coarse-grained simulation and enhanced sampling methods.

Lastly, our simulations lack substrates for either FtsW or FtsI, omit termini of some divisome

subunits known to impact cell-wall synthesis [45], and lack other protein components known to

interact with the core divisome complex. Filling in more of this picture will make it possible to link

impacts on divisome structure and dynamics more directly with substrate binding and enzymatic

mechanisms.

In summary**,** our results are not only consistent with past biochemical and genetic studies,

but also shed light on the molecular details of active and inactive conformations of FtsWI. The

approach we developed in this work—structure prediction followed by MD simulation with results

informing experiments to test hypotheses arising from modeled structures—proved powerful in
generating new insights into molecular interactions in the divisome.

DATA AND CODE AVAILABILITY

The plasmids and *E. coli* strains used in this study are available from the corresponding authors
upon request. Code for analyzing single-molecule tracking data is available from the Xiao
Laboratory GitHub repository (XiaoLabJHU) [17]. Full all-atom trajectories will be made available in
the Anton 2 database also upon publication. Files with coordinates for predicted structures and
for structures following 1-µs MD are included as supplementary data.

ACKNOWLEGEMENTS

The authors thank all members of the Hensel, Xiao, and Lau laboratories for helpful discussions
and feedback on the manuscript. ZH received support for this work from FCT - Fundação para a
Ciência e a Tecnologia, I.P., through MOSTMICRO-ITQB R&D Unit (UIDB/04612/2020,
UIDP/04612/2020) and LS4FUTURE Associated Laboratory (LA/P/0087/2020), from a joint
research agreement with the Okinawa Institute of Science and Technology, and from the Google
Cloud Research Credits program with the award GCP20210916. SMC received support from FCT
Fundação para a Ciência e a Tecnologia, I.P., through a PhD fellowship (PD/BD/135480/2018).
Work in the Xiao laboratory was supported by NIH F32GM143895 (B.M.B.), NIH T32GM007445
(J.W.M), and R35GM136436 (J.X.). Work in the Lau laboratory was funded by the Johns Hopkins
Catalyst Award (to A.Y.L.); NIH T32GM135131 (to R.A.Y.). Anton 2 computer time
(MCB130045P) was provided by the Pittsburgh Supercomputing Center (PSC) through NIH grant
R01GM116961 (to A.Y.L.); the Anton 2 machine at PSC was generously made available by D.E.
Shaw Research. We also used resources provided by and Advanced Research Computing at
Hopkins (ARCH) at Johns Hopkins University.

AUTHOR INFORMATION

Authors and Affiliations

Department of Biophysics and Biophysical Chemistry, Johns Hopkins School of Medicine,
Baltimore, MD, 21205, USA

Brooke M. Britton, Remy A. Yovanno, Joshua McCausland, Albert Y. Lau, Jie Xiao

ITQB NOVA, Universidade NOVA de Lisboa, Lisbon, Av. da República, 2780-157 Oeiras,

Portugal.

Sara F. Costa, Zach Hensel

Contributions

B.M.B., J.X., and Z.H. conceived the experiments. R.A.Y. and A.Y.L. designed the simulation

workflow. B.M.B. and J.W.M. constructed plasmids and strains for imaging experiments. B.M.B.

performed single molecule and phenotype imaging, genetic experiments, and all imaging analysis.

R.A.Y. performed molecular dynamic simulations. R.A.Y. and Z.H. wrote analysis code. B.M.B.,

R.A.Y., A.Y.L, J.X., and Z.H. analyzed molecular dynamics data. S.C. and Z.H. analyzed

sequence conservation. B.M.B., R.A.Y., J.X., and Z.H. wrote the original draft. All authors

reviewed and edited the manuscript. B.M.B., J.W.M., J.X., and Z.H. acquired funding.

MATERIALS AND METHODS

Complementation assay

To ensure proper growth of *ΔftsL* and *ΔftsI* strains, arabinose was included during maintenance

and preparation to induce expression of FtsL and FtsI, respectively, from the P$_{BAD}$ promoter. Cells

were grown overnight at 37 °C in LB and 0.2% arabinose from a single colony. The following day,

the saturated culture was reinoculated 1:1000 in fresh LB with 0.2% arabinose. Cells were grown

in log phase at 37 °C to an OD600 of 0.5 for all conditions. Cells were then washed three times

with 0.9% saline. They were then serial diluted in 0.9% saline and plated on LB plates containing

either 0.2% arabinose or 0.4% glucose (to repress expression from the P$_{BAD}$ promoter) and IPTG

as noted. Plates were grown at 37 °C overnight and then imaged.

Plasmid and Strain Construction

Plasmids used in this study (**Supplementary Table S1**) were assembled by Polymerase Chain

Reactions (PCR) amplifying insert and vector DNA fragments followed by In-Fusion cloning

(Takara Biosciences, In-Fusion HD Cloning Kit). Oligonucleotides (Integrated DNA Technologies)

used in PCR amplifications are described in **Supplementary Table S12.** All plasmids were

verified by DNA sequencing. After construction, electroporation was used to transform plasmids

to create strains of interest (**Supplementary Table S2**) under appropriate antibiotic selection.

Depletion strains were maintained under arabinose induction to maintain wild-type phenotype.

Single-molecule tracking sample preparation, imaging, and data analysis

Prior to imaging, cells were grown to log phase at 25 °C in defined minimal M9 medium (0.4%

glucose, 1x MEM amino acids, and 1x MEM vitamins, M9(+) glucose). Cells were incubated for

20 minutes with the addition of 50 nM Janelia Fluor 646 (JF646). Following labeling, cells were

washed 3 times with M9(–) glucose (0.4% glucose and 1x MEM vitamins). Cells were placed

onto a 3% agarose gel pad (M9(–) glucose), sandwiched with a coverslip, and enclosed within an

FSC2 chamber (Bioptechs) for imaging. Cells were imaged after 30 minutes of equilibration on

the microscope.

For experiments with fosfomycin, 200 µg/ml of Fosfomycin was added to the gel pad and to the

cells following the final wash step. Cells were imaged 1 hour after Fosfomycin addition. For

experiments with rich, defined medium, EZDRM (Teknova) was used in place of M9 for the growth

media, wash buffer, and gel pad.

HaloTag-FtsB tracking was performed on a home-built microscope as previously described [17,30].

Briefly, strains were imaged on an Olympus IX-71 microscope body using an UPLANApo

100XOHR Objective (NA1.50/oil) with a 1.6 x field lens engaged. JF646 was imaged with a 674-

nm laser (Coherent) at an excitation power density of ~25 W/cm^2. The exposure time was 500 ms

and the imaging frame rate was 1 Hz.

Single-molecule data analysis was performed as previously described [17,18,30]. Briefly, positions of

single molecules were determined using the ImageJ plug-in, ThunderSTORM [57]. The remaining

data analysis and postprocessing was performed using home-built MATLAB scripts available from

the Xiao Laboratory GitHub repository (XiaoLabJHU) [17]. In short, using a nearest-neighbor

algorithm, single molecules were linked to trajectories. Using the bright-field image as a visual

guide, only trajectories near the midplane of a cell's long axis or visible constriction sites were

chosen to ensure the measurements were made on single molecules at the cell-division site. The

true displacement of tracked molecules around the circumference of a cell is underestimated in

2D single-molecule tracking due to the cylindrical cell envelope. Therefore, the trajectories were

unwrapped to one-dimension. Unwrapped trajectories were then segmented manually to

determine stationary or processive movement. These procedures were described in detail in [17].

The cumulative probability distribution of directional moving velocities was calculated for each
condition and fit to either a single or double log-normal population:

$$CDF = P_1 \frac{1 + erf\left[\frac{lnv - \mu_1}{\sqrt{2\sigma_1}}\right]}{2} + (1 - P_1)\frac{1 + erf\left[\frac{lnv - \mu_1}{\sqrt{2\sigma_2}}\right]}{2},$$

where v is the moving velocity and P_1 is the percentage of the first population. For a single
population, $P_1 = 1$. The parameters μ and σ are the natural logarithmic mean and standard
deviation of the log-normal distribution. The average velocity was calculated using $\exp\left(\mu + \frac{\sigma^2}{2}\right)$.
To estimate errors in parameter estimates, CDF curves were generated for 1,000 bootstrapped
samples and fit, with the standard deviations of parameter fits estimating 1 standard error of
measurement (s.e.m.).

Bright-field sample preparation and imaging

Cells were grown overnight at 37 °C in LB and 0.2% arabinose from a single colony. The following
day, the saturated culture was reinoculated 1:1000 in fresh LB with the appropriate inducer and/or
repressor (0.2% arabinose, 0.4% glucose, and/or ITPG as noted). Cells were grown to log phase
at 37 °C. Cells were then washed three times with 0.9% saline with the appropriate inducer and/or
repressor (0.2% arabinose, 0.4% glucose, and/or ITPG as noted). Cells were placed onto a 3%
agarose gel pad (0.9% saline with the appropriate inducer and/or repressor (0.2% arabinose,
0.4% glucose, and/or ITPG as noted), sandwiched with a coverslip, and enclosed within an FSC2
chamber (Bioptechs) for immediate imaging. Phase imaging was performed on a home-built
microscope as previously described [30] or on the same microscope as that used for single-
molecule tracking experiment with condenser lamp illumination.

Prediction of protein complex structures

We used AlphaFold2 [23] as implemented in ColabFold [32] using the LocalColabFold version and
AlphaFold2 parameters available in September 2021. ColabFold uses the MMseqs2 server [58] to
obtain sequence alignments, greatly reducing local storage and computation requirements.
Predictions were made on a Google Cloud Platform instance with an A100 GPU, taking
approximately 1.5 h including sidechain relaxation. Template structures were not used, and
predictions utilized 48 recycle steps. In preliminary work, we found that AlphaFold2 model 3,
which was trained without templates, typically provided the best performance as measured by
local pLDDT and global pTMscore metrics. Predictions from model 3 for FtsWI, FtsQLBWI, and
FtsQLBWIN were used to generate MD systems. Model 5, which was trained similarly and

produced similar predictions for γ-proteobacterial complexes, was later found to reproduce some

additional reported interactions in other species and was used for *B. subtilis* and *S. pneumonia* in

Fig. S2B. We did not use AlphaFold-multimer [25] for these predictions, as its initial implementation

resulted in structures with steric clashes that could not be directly used for building MD systems.

We note that this has since been addressed and that later versions of AlphaFold-multimer give

similar predictions of divisome complexes to what we report that typically lack clashes and are

appropriate for MD. All structure predictions shown in the manuscript, as well as protein complex

structures following 1 μs of MD, are available as supplementary data.

Conservation of FtsN^E

We constructed sequence logos illustrating FtsN^E conservation following previously reported

methods applied to the *E. coli* FtsN cytoplasmic domain [59]. Sequences annotated as FtsN

(TIGRFAM code TIGR02223) were downloaded from AnnoTree [60] for *Enterobacteriaceae* (513),

Pasteurellaceae (113), and *Vibrionaceae* (227). Sequences that were incorrectly annotated

based on length or other features were manually removed. Multiple sequences alignments were

constructed using MUSCLE [61] and sequence logos for regions corresponding to *E. coli* FtsN$^{73–95}$

were obtained using WebLogo 3 [62]. Only 10 FtsN sequences were identified for

Pseudomonadaceae following this method, so we obtained 304 sequences homologous to *P.*

aeruginosa FtsN$^{68–118}$ in *Pseudomonadaceae* using blastp against the RefSeq Select proteins

database, verifying that this also reproduced conservation observed for *Enterobacteriaceae* when

using *E. coli* FtsN$^{58–108}$ in the same way.

Equilibrium molecular dynamics simulation of FtsQLBWI and FtsQLBWIN complexes
with superfission and dominant negative variants

Simulation systems for molecular dynamics were constructed from AlphaFold models of FtsWI,

FtsQLBWI, and FtsQLBWIN including the following subunits and residues: FtsQ (20–276), FtsL

(1–121), FtsB (1–103), FtsW (46–414), FtsI (19–588), and FtsN (58–108). Mutations of interest

were made using the Mutagenesis Wizard of the PyMOL molecular visualization software.

All systems were embedded in a POPE membrane and solvated (see Supplemental **Table S5** for

system dimensions) with 150 mM NaCl in TIP3P water [63] using the CHARMM-GUI Membrane

Builder. All systems were electrically neutral. N-termini of FtsQ, FtsW, FtsI, and FtsN were capped

with the CHARMM ACE patch, and the C-terminus of FtsN was capped with the CHARMM CT3

patch. Equilibration was performed according to the CHARMM-GUI equilibration protocol

involving a series of gradually relaxing sidechain and backbone restraints. After equilibration, 5

ns pre-production simulations were performed prior to simulation on the special-purpose

supercomputer Anton 2 at the Pittsburgh Supercomputing Center.

Simulations on Anton 2 were performed using the CHARMM36m forcefield in an NPT ensemble

at 310 K, 1 atm, and with a timestep of 2.5 fs. Bond lengths for hydrogen atoms were constrained

using the M-SHAKE algorithm [64]. An r-RESPA integrator [65] was used; long-range electrostatics

were computed every 6 fs. Long-range electrostatics interactions were calculated using the k-

space Gaussian split Ewald method [66]. Trajectories were written at an output frequency of 0.24

ns/frame. Each trajectory was unwrapped using the PBCTools plugin of VMD and aligned to the

backbone atoms of FtsW (segid PROD) to facilitate comparison across complexes. Code for

analyzing simulations was written using the MDAnalysis python package. Hydrogen bonds

between relevant interfaces were computed using the HydrogenBondAnalysis function of

MDAnalysis (v 1.0.0) in which a hydrogen bond is defined by a distance cutoff of 3.0 Å and an

angle cutoff of 150°.

Calculation of structure parameters

Solvent accessible surface area was calculated for all hydrophobic FtsL residues in the region of

FtsL interacting with FtsB and the FtsI head and anchor domains (residues 62–86 inclusive of

FtsL residues L62, L63, A65, L70, V71, L72, A76, L77, I79, W81, L84, I85, and L86) using the

get_area command in PyMOL (version 2.5) with parameters *dot_solvent*, *dot_density*, and

solvent_radius set to 1, 4, and 1.4, respectively. FtsI TPase domain tilt angles were calculated

following methods applied to the *P. aeruginosa* FtsQLBWI structure [41]. Trajectories were aligned

by minimizing RMSD for FtsW C_α atoms, and the coordinates of C_α for FtsI^{S307} were tracked in

this reference frame. The tilt angle is defined as the angle between two vectors connecting C_α

atoms for FtsW^{D297} and FtsI^{S307} in the AF2-predicted structure and in frames of the trajectory. The

P. aeruginosa structure was compared by aligning to *E. coli* FtsW using the default settings for

the *align* command in PyMol (version 2.5) and analyzing C_α coordinates for equivalent residues.

Dynamical network analysis for computing optimal paths

Dynamical network representations of the FtsQBLWI, FtsQLBWI^{R167S}, FtsQL^{R61E}BWI, and

FtsQLBWIN complexes were generated using the dynetan python package [67]. In the network,

each node is defined as the C_α atom of a residue, and pairs of nodes are considered connected

if their heavy atoms are within 4.5 Å of each other for at least 75% of the final 500 ns of the

trajectory. The strength of an edge between two nodes is calculated using a correlation coefficient

computed from a k-nearest-neighbor-based estimator of mutual information [67]. From these

correlations, optimal paths were computed using the Floyd-Warshall algorithm implemented in

dynetan and NetworkX [68]. To illustrate long-range interaction paths, a source node was selected

as the first simulated residue of FtsQ (FtsQ[N20]) and target nodes were selected as the catalytic

residues of FtsW (FtsW[D297]) and FtsI (FtsI[S307]), resulting in two ensembles of optimal paths.

REFERENCES

1 Schneider, T. & Sahl, H. G. An oldie but a goodie - cell wall biosynthesis as antibiotic
target pathway. *Int J Med Microbiol* **300**, 161-169 (2010).
https://doi.org:10.1016/j.ijmm.2009.10.005
2 Silver, L. L. Does the cell wall of bacteria remain a viable source of targets for novel
antibiotics? *Biochem Pharmacol* **71**, 996-1005 (2006).
https://doi.org:10.1016/j.bcp.2005.10.029
3 Paulussen, F. M. *et al.* Covalent Proteomimetic Inhibitor of the Bacterial FtsQB Divisome
Complex. *J Am Chem Soc* **144**, 15303-15313 (2022).
https://doi.org:10.1021/jacs.2c06304
4 Bi, E. F. & Lutkenhaus, J. FtsZ ring structure associated with division in Escherichia coli.
Nature **354**, 161-164 (1991). https://doi.org:10.1038/354161a0
5 Nanninga, N. Cell division and peptidoglycan assembly in Escherichia coli. *Mol Microbiol*
5, 791-795 (1991). https://doi.org:10.1111/j.1365-2958.1991.tb00751.x
6 Levin, P. A. & Janakiraman, A. Localization, Assembly, and Activation of the Escherichia
coli Cell Division Machinery. *EcoSal Plus* **9**, eESP00222021 (2021).
https://doi.org:10.1128/ecosalplus.ESP-0022-2021
7 Taguchi, A. *et al.* FtsW is a peptidoglycan polymerase that is functional only in complex
with its cognate penicillin-binding protein. *Nat Microbiol* **4**, 587-594 (2019).
https://doi.org:10.1038/s41564-018-0345-x
8 Fraipont, C. *et al.* The integral membrane FtsW protein and peptidoglycan synthase
PBP3 form a subcomplex in Escherichia coli. *Microbiology (Reading)* **157**, 251-259
(2011). https://doi.org:10.1099/mic.0.040071-0
9 Marmont, L. S. & Bernhardt, T. G. A conserved subcomplex within the bacterial
cytokinetic ring activates cell wall synthesis by the FtsW-FtsI synthase. *Proc Natl Acad
Sci U S A* **117**, 23879-23885 (2020). https://doi.org:10.1073/pnas.2004598117
10 Park, K. T., Du, S. & Lutkenhaus, J. Essential Role for FtsL in Activation of Septal
Peptidoglycan Synthesis. *mBio* **11** (2020). https://doi.org:10.1128/mBio.03012-20
11 Tsang, M. J. & Bernhardt, T. G. A role for the FtsQLB complex in cytokinetic ring
activation revealed by an ftsL allele that accelerates division. *Mol Microbiol* **95**, 925-944
(2015). https://doi.org:10.1111/mmi.12905
12 Du, S. & Lutkenhaus, J. Assembly and activation of the Escherichia coli divisome. *Mol
Microbiol* **105**, 177-187 (2017). https://doi.org:10.1111/mmi.13696
13 Gerding, M. A. *et al.* Self-enhanced accumulation of FtsN at Division Sites and Roles for
Other Proteins with a SPOR domain (DamX, DedD, and RlpA) in Escherichia coli cell
constriction. *J Bacteriol* **191**, 7383-7401 (2009). https://doi.org:10.1128/JB.00811-09
14 Yahashiri, A., Jorgenson, M. A. & Weiss, D. S. Bacterial SPOR domains are recruited to
septal peptidoglycan by binding to glycan strands that lack stem peptides. *Proc Natl
Acad Sci U S A* **112**, 11347-11352 (2015). https://doi.org:10.1073/pnas.1508536112

15   Liu, B., Persons, L., Lee, L. & de Boer, P. A. Roles for both FtsA and the FtsBLQ
subcomplex in FtsN-stimulated cell constriction in Escherichia coli. *Mol Microbiol* **95**,
945-970 (2015). https://doi.org:10.1111/mmi.12906
16   Pichoff, S., Du, S. & Lutkenhaus, J. Disruption of divisome assembly rescued by FtsN-
FtsA interaction in Escherichia coli. *Proc Natl Acad Sci U S A* **115**, E6855-E6862 (2018).
https://doi.org:10.1073/pnas.1806450115
17   Yang, X. *et al.* A two-track model for the spatiotemporal coordination of bacterial septal
cell wall synthesis revealed by single-molecule imaging of FtsW. *Nat Microbiol* **6**, 584-
593 (2021). https://doi.org:10.1038/s41564-020-00853-0
18   Lyu, Z. *et al.* FtsN maintains active septal cell wall synthesis by forming a processive
complex with the septum-specific peptidoglycan synthases in E. coli. *Nat Commun* **13**,
5751 (2022). https://doi.org:10.1038/s41467-022-33404-8
19   Boes, A., Olatunji, S., Breukink, E. & Terrak, M. Regulation of the Peptidoglycan
Polymerase Activity of PBP1b by Antagonist Actions of the Core Divisome Proteins
FtsBLQ and FtsN. *mBio* **10** (2019). https://doi.org:10.1128/mBio.01912-18
20   Li, Y. *et al.* Genetic analysis of the septal peptidoglycan synthase FtsWI complex
supports a conserved activation mechanism for SEDS-bPBP complexes. *PLoS Genet*
17, e1009366 (2021). https://doi.org:10.1371/journal.pgen.1009366
21   Wissel, M. C. & Weiss, D. S. Genetic analysis of the cell division protein FtsI (PBP3):
amino acid substitutions that impair septal localization of FtsI and recruitment of FtsN. *J
Bacteriol* **186**, 490-502 (2004). https://doi.org:10.1128/jb.186.2.490-502.2004
22   Attaibi, M. & den Blaauwen, T. An Updated Model of the Divisome: Regulation of the
Septal Peptidoglycan Synthesis Machinery by the Divisome. *Int J Mol Sci* **23** (2022).
https://doi.org:10.3390/ijms23073537
23   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021). https://doi.org:10.1038/s41586-021-03819-2
24   Craven, S. J., Condon, S. G. F., Diaz Vazquez, G., Cui, Q. & Senes, A. The coiled-coil
domain of Escherichia coli FtsLB is a structurally detuned element critical for modulating
its activation in bacterial cell division. *J Biol Chem* **298**, 101460 (2022).
https://doi.org:10.1016/j.jbc.2021.101460
25   Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *bioRxiv*,
2021.2010.2004.463034 (2022). https://doi.org:10.1101/2021.10.04.463034
26   Baek, M. *et al.* Accurate prediction of protein structures and interactions using a three-
track neural network. *Science* **373**, 871-876 (2021).
https://doi.org:doi:10.1126/science.abj8754
27   Bryant, P., Pozzati, G. & Elofsson, A. Improved prediction of protein-protein interactions
using AlphaFold2. *Nat Commun* **13**, 1265 (2022). https://doi.org:10.1038/s41467-022-
28865-w
28   Modi, V. & Dunbrack, R. L. Kincore: a web resource for structural classification of protein
kinases and their inhibitors. *Nucleic Acids Res* **50**, D654-D664 (2022).
https://doi.org:10.1093/nar/gkab920
29   Pak, M. A. *et al.* Using AlphaFold to predict the impact of single mutations on protein
stability and function. *bioRxiv*, 2021.2009.2019.460937 (2021).
https://doi.org:10.1101/2021.09.19.460937
30   McCausland, J. W. *et al.* Treadmilling FtsZ polymers drive the directional movement of
sPG-synthesis enzymes via a Brownian ratchet mechanism. *Nat Commun* **12**, 609
(2021). https://doi.org:10.1038/s41467-020-20873-y
31   Buddelmeijer, N. & Beckwith, J. A complex of the Escherichia coli cell division proteins
FtsL, FtsB and FtsQ forms independently of its localization to the septal region. *Mol
Microbiol* **52**, 1315-1327 (2004). https://doi.org:10.1111/j.1365-2958.2004.04044.x

32   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**,
679-682 (2022). https://doi.org:10.1038/s41592-022-01488-1

33   Basu, S. & Wallner, B. DockQ: A Quality Measure for Protein-Protein Docking Models.
PLoS One **11**, e0161879 (2016). https://doi.org:10.1371/journal.pone.0161879

34   Morales Angeles, D., Macia-Valero, A., Bohorquez, L. C. & Scheffers, D. J. The PASTA
domains of Bacillus subtilis PBP2B strengthen the interaction of PBP2B with DivIB.
Microbiology (Reading) **166**, 826-836 (2020). https://doi.org:10.1099/mic.0.000957

35   Bernardo-García, N. *et al.* Allostery, Recognition of Nascent Peptidoglycan, and Cross-
linking of the Cell Wall by the Essential Penicillin-Binding Protein 2x of Streptococcus
pneumoniae. *ACS Chemical Biology* **13**, 694-702 (2018).
https://doi.org:10.1021/acschembio.7b00817

36   Robichon, C., Karimova, G., Beckwith, J. & Ladant, D. Role of leucine zipper motifs in
association of the Escherichia coli cell division proteins FtsL and FtsB. *J Bacteriol* **193**,
4988-4992 (2011). https://doi.org:10.1128/JB.00324-11

37   Sjodt, M. *et al.* Structure of the peptidoglycan polymerase RodA resolved by evolutionary
coupling analysis. *Nature* **556**, 118-121 (2018). https://doi.org:10.1038/nature25985

38   Sjodt, M. *et al.* Structural coordination of polymerization and crosslinking by a SEDS-
bPBP peptidoglycan synthase complex. *Nat Microbiol* **5**, 813-820 (2020).
https://doi.org:10.1038/s41564-020-0687-z

39   Freischem, S. *et al.* Interaction Mode of the Novel Monobactam AIC499 Targeting
Penicillin Binding Protein 3 of Gram-Negative Bacteria. *Biomolecules* **11** (2021).

40   Kureisaite-Ciziene, D. *et al.* Structural Analysis of the Interaction between the Bacterial
Cell Division Proteins FtsQ and FtsB. *mBio* **9** (2018).
https://doi.org:10.1128/mBio.01346-18

41   Kashammer, L. *et al.* Divisome core complex in bacterial cell division revealed by cryo-
EM. *bioRxiv*, 2022.2011.2021.517367 (2022).
https://doi.org:10.1101/2022.11.21.517367

42   Li, Y. *et al.* Identification of the potential active site of the septal peptidoglycan
polymerase FtsW. *PLoS Genet* **18**, e1009993 (2022).
https://doi.org:10.1371/journal.pgen.1009993

43   Park, K. T., Pichoff, S., Du, S. & Lutkenhaus, J. FtsA acts through FtsW to promote cell
wall synthesis during cell division in Escherichia coli. *Proc Natl Acad Sci U S A* **118**
(2021). https://doi.org:10.1073/pnas.2107210118

44   Choi, Y. *et al.* Structural Insights into the FtsQ/FtsB/FtsL Complex, a Key Component of
the Divisome. *Sci Rep* **8**, 18061 (2018). https://doi.org:10.1038/s41598-018-36001-2

45   Craven, S. J., Condon, S. G. F. & Senes, A. A model of the interactions between the
FtsQLB and the FtsWI peptidoglycan synthase complex in bacterial cell division. *bioRxiv*,
2022.2010.2030.514410 (2022). https://doi.org:10.1101/2022.10.30.514410

46   Nagasawa, H. *et al.* Determination of the cleavage site involved in C-terminal processing
of penicillin-binding protein 3 of Escherichia coli. *J Bacteriol* **171**, 5890-5893 (1989).
https://doi.org:10.1128/jb.171.11.5890-5893.1989

47   Hara, H. *et al.* Genetic analyses of processing involving C-terminal cleavage in penicillin-
binding protein 3 of Escherichia coli. *J Bacteriol* **171**, 5882-5889 (1989).
https://doi.org:10.1128/jb.171.11.5882-5889.1989

48   Gonzalez, M. D., Akbay, E. A., Boyd, D. & Beckwith, J. Multiple interaction domains in
FtsL, a protein component of the widely conserved bacterial FtsLBQ cell division
complex. *J Bacteriol* **192**, 2757-2768 (2010). https://doi.org:10.1128/JB.01609-09

49   Goehring, N. W., Petrovska, I., Boyd, D. & Beckwith, J. Mutants, suppressors, and
wrinkled colonies: mutant alleles of the cell division gene ftsQ point to functional
domains in FtsQ and a role for domain 1C of FtsA in divisome assembly. *J Bacteriol* **189**,
633-645 (2007). https://doi.org:10.1128/JB.00991-06

50 Condon, S. G. F. *et al.* The FtsLB subcomplex of the bacterial divisome is a tetramer
with an uninterrupted FtsL helix linking the transmembrane and periplasmic regions. *J*
Biol Chem **293**, 1623-1641 (2018). https://doi.org:10.1074/jbc.RA117.000426
51 Park, K.-T., Du, S. & Lutkenhaus, J. Essential role for FtsL in activation of septal PG
synthesis. *bioRxiv*, 2020.2009.2001.275982 (2020).
https://doi.org:10.1101/2020.09.01.275982
52 Brown, A. M. & Zondlo, N. J. A propensity scale for type II polyproline helices (PPII):
aromatic amino acids in proline-rich sequences strongly disfavor PPII due to proline-
aromatic interactions. *Biochemistry* **51**, 5041-5051 (2012).
https://doi.org:10.1021/bi3002924
53 Williams, K. P. *et al.* Phylogeny of gammaproteobacteria. *J Bacteriol* **192**, 2305-2314
(2010). https://doi.org:10.1128/jb.01480-09
54 Gao, B., Mohan, R. & Gupta, R. S. Phylogenomics and protein signatures elucidating the
evolutionary relationships among the Gammaproteobacteria. *Int J Syst Evol Microbiol*
59, 234-247 (2009). https://doi.org:10.1099/ijs.0.002741-0
55 Melo, M. C. R., Bernardi, R. C., Fuente-Nunez, C. d. l. & Luthey-Schulten, Z.
Generalized correlation-based dynamical network analysis: a new high-performance
approach for identifying allosteric communications in molecular dynamics trajectories.
The Journal of Chemical Physics **153**, 134104 (2020). https://doi.org:10.1063/5.0018980
56 Gómez, M. J., Desviat, L. R., Merchante, R. & Ayala, J. A. in *Bacterial Growth and Lysis:*
Metabolism and Structure of the Bacterial Sacculus   (eds M. A. de Pedro, J. V. Höltje, &
W. Löffelhardt) 309-318 (Springer US, 1993).
57 Ovesný, M., Křížek, P., Borkovec, J., Svindrych, Z. & Hagen, G. M. ThunderSTORM: a
comprehensive ImageJ plug-in for PALM and STORM data analysis and super-
resolution imaging. *Bioinformatics* **30**, 2389-2390 (2014).
https://doi.org:10.1093/bioinformatics/btu202
58 Steinegger, M. & Soding, J. MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028 (2017).
https://doi.org:10.1038/nbt.3988
59 Nierhaus, T. *et al.* Bacterial divisome protein FtsA forms curved antiparallel double
filaments when binding to FtsN. *Nat Microbiol* **7**, 1686-1701 (2022).
https://doi.org:10.1038/s41564-022-01206-9
60 Mendler, K. *et al.* AnnoTree: visualization and exploration of a functionally annotated
microbial tree of life. *Nucleic Acids Res* **47**, 4442-4448 (2019).
https://doi.org:10.1093/nar/gkz246
61 Edgar, R. C. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. *BMC Bioinformatics* **5**, 113 (2004). https://doi.org:10.1186/1471-2105-
5-113
62 Crooks, G. E., Hon, G., Chandonia, J. M. & Brenner, S. E. WebLogo: a sequence logo
generator. *Genome Res* **14**, 1188-1190 (2004). https://doi.org:10.1101/gr.849004
63 Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W. & Klein, M. L.
Comparison of simple potential functions for simulating liquid water. *The Journal of*
Chemical Physics **79**, 926-935 (1983). https://doi.org:10.1063/1.445869
64 Kräutler, V., van Gunsteren, W. F. & Hünenberger, P. H. A fast SHAKE algorithm to
solve distance constraint equations for small molecules in molecular dynamics
simulations. *Journal of Computational Chemistry* **22**, 501-508 (2001).
https://doi.org:https://doi.org/10.1002/1096-987X(20010415)22:5<501::AID-
JCC1021>3.0.CO;2-V
65 Tuckerman, M., Berne, B. J. & Martyna, G. J. Reversible multiple time scale molecular
dynamics. *The Journal of Chemical Physics* **97**, 1990-2001 (1992).
https://doi.org:10.1063/1.463137

66   Shan, Y., Klepeis, J. L., Eastwood, M. P., Dror, R. O. & Shaw, D. E. Gaussian split
Ewald: A fast Ewald mesh method for molecular simulation. *The Journal of Chemical*
Physics **122**, 054101 (2005). https://doi.org:10.1063/1.1839571
67   Melo, M. C. R., Bernardi, R. C., de la Fuente-Nunez, C. & Luthey-Schulten, Z.
Generalized correlation-based dynamical network analysis: a new high-performance
approach for identifying allosteric communications in molecular dynamics trajectories.
The Journal of Chemical Physics **153**, 134104 (2020). https://doi.org:10.1063/5.0018980
68   Hagberg, A. S., Pieter; S Chult, Daniel. Exploring network structure, dynamics, and
function using networkx. (2008).

FIGURES

Figure 1

Figure 1. Characterization and modeling of the *E. coli* FtsQLBWI complex. (A, B) Single-

molecule tracking of Halo-FtsB suggests that FtsQLB remain in complex with FtsWI on both the

fast-moving FtsZ-track and the slow-moving sPG synthesis track. (**A**) Two representative Halo-

FtsB expressing cells with the maximum fluorescence intensity projection images (left),

kymographs of fluorescence line scans at the midcell (middle), and unwrapped one-dimensional

positions of the corresponding Halo-FtsB molecule along the circumference (solid gray line) and

long axis (dotted gray line) of the cell were shown. Measured velocity of each segment and the

corresponding classification (fast-moving, cyan; slow-moving, pink; stationary, gray) are labeled

in the trajectory panels. Scale bar 500 nm. **(B)** Distribution of velocities of single Halo-FtsB

molecules exhibiting directional motion in wild-type *E. coli* cells grown in minimal media in the

absence (top) of fosfomycin was best fit with two moving populations, one slow (red) and one fast

(blue). A dashed line indicates the summed probability. In the presence of fosfomycin that inhibits

cell wall synthesis (bottom), the slow-moving population (red) is drastically reduced. These

dynamics behaviors are similar to those of FtsW or FtsI [17]. **(C)** Modeled structure of *E. coli*

FtsQLBWI within a POPE bilayer (purple) in the last frame of a 1-μs MD simulation. The complex

consists of FtsQ 20–276 (gray), FtsL 1–121 (red), FtsB 1–113 (blue), FtsW 46–414 (orange), and

FtsI 19–588 (magenta). The FtsI TPase, head and anchor domains are labeled in magenta text.

The four interface regions—Pivot, Truss, Hub, and Lid—are highlighted in dashed boxes. **(D)** In

the absence of FtsQLB, FtsI (magenta) collapses to the membrane (purple) at the end of the 1-

μs MD simulation. The position of FtsI at the beginning of the simulation (transparent magenta) is

shown for comparison. **(E)** Zoomed-in view of the Pivot region in FtsQLBWI, in which interactions

between FtsI^{L53} (magenta) and FtsW^{M269} (orange) and between FtsI^{R60} (magenta) and FtsW^{Q266}

(orange) secure the position of the FtsI anchor domain (magenta) with respect to FtsW (orange).

(F) In the absence of FtsQLB, interactions between FtsI^{L53} and FtsW^{M269} are broken, as shown by

the increased Cα-Cα distances between the two residues (light gray) compared to that in the

presence of FtsQLB (WT, dark gray) in the last 500 ns of the MD simulation.

Figure 2

Figure 2. C-terminal extended β-sheet of FtsQLBI in the Truss region is important for cell

division. (A) A detailed view of the Truss region in the final frame of the FtsQLBWI 1-μs MD

simulation illustrates β-sheet interactions between the C-terminal ends of FtsQ (gray), FtsB (blue),

FtsL (red), and FtsI (magenta). **(B)** Cartoon showing FtsL and FtsI β-strand truncation mutants.

Also see **Fig. S12** for details. **(C)** Spot dilution complementation test of FtsL truncation mutants.

E. coli cells depleted of chromosomal wild-type FtsL but contain FtsL expressed from the P$_{BAD}$

promoter (strain MDG279, **Table S2**) complemented in the presence of arabinose (top panel), but

failed in the presence of glucose (No Plasmid, first rows of the bottom panel). The same depletion

strain (MDG279) expressing wild-type FtsL (pCH-FtsL, or pBMB064), FtsL$^{\Delta 6}$ (pCH-FtsL$^{\Delta 6}$, or

pBMB065) and FtsL$^{\Delta 11}$ (pCH-FtsL$^{\Delta 11}$, or pBMB066, **Table S1**) from a *lac* promoter on plasmids

complemented the depletion in the presence of glucose at both no induction and 100 μM IPTG

conditions (middle rows of both panels). pCH-FtsL$^{\Delta 16}$ is unable to complement at both conditions

(bottom rows of the bottom panel). See **Fig. S13** for more induction conditions. **(D)** Images of *E.*

coli cells depleted of wild-type FtsL and expressing an mVenus fusion to FtsL of various

truncations showed that truncations of FtsL of increasing length exhibited increasing cell length

(top) and decreased FtsL midcell localization (bottom) relative to cells expressing mVenus fused

to full-length FtsL. Scale bar 3μm. See Fig. S13 for quantifications. **(E)** Images of *E. coli* cells

depleted of wild-type FtsI and expressing FtsI truncations. A wild-type FtsI fusion and FtsI$^{\Delta 14}$

exhibit near-normal cell lengths even at low induction levels, while FtsI$^{\Delta 11}$ exhibits filamentous

cells at low expression levels. Scale bar 3 μm. See **Fig. S14** for quantifications.

Figure 3

Figure 3. Inhibitory and activating interactions between FtsL and FtsI in the Hub region are
distributed along the CCD (A) and AWI interfaces (B and C). (A) A hydrogen-bonding network
extends from FtsQLB to FtsI in the CCD interface of the Hub region. Shapes indicate residues
involved in hydrogen bonding. Superfission mutant residues (FtsL[N89], FtsL[E88], FtsL[H94], and
FtsB[E56]) are depicted as squares. Dominant negative mutant residues (FtsL[R82] and FtsL[N83])

depicted as tetrahedrons. Spheres indicate other residues involved (FtsI$^{P87,\ S70,\ S85,\ Y168,\ P170,\ R246}$,

FtsL^{E98}, FtsB$^{E69,\ R70}$, and FtsQ$^{R196,\ R213}$). Hydrogen-bond frequencies during the last 500 ns of

FtsQLBWI simulation are indicated by the color bar. **(B)** Hydrophobic packing among FtsL^{I85},

FtsL^{L86}, FtsI^{V86}, FtsI^{V84}, and FtsI^{Y168} is observed in the AWI domain between FtsL (red) and FtsI

(magenta) in the WT FtsQLBWI simulation. **(C)** The same AWI interface exhibits a tighter

hydrophobic packing in the FtsQLBWI^{R167S} complex simulation. Note the relative differences

between FtsI^{V84,Y168} and FtsL^{I85}. **(D)** The distance between the centers of geometry of FtsI^{V84} and

FtsL^{I85} sidechains decreased from $d = 6.4 \pm 0.6$ Å in the WT FtsQLBWI complex to 5.4 ± 0.5 Å in

the FtsQLBWI^{R167S} in the last 500 ns of simulations, consistent with the closer packing between

the two residues in the SF FtsQLBWI^{R167S} complex. **(E)** Top: conformations after 1 µs MD of

FtsL^{G92} (left) and FtsL^{H94} (right) in the WT FtsQLBWI (tan color) and the SF FtsQLBWI^{R167S} (dark

red color) complexes show disruption of the second short α-helix of FtsL in the FtsQLBWI^{R167S}

complex. Bottom: dihedral angles of both FtsL^{G92} (left) and FtsL^{H94} (right) are disrupted in the

FtsQLBWI^{R167S} complex. A shift for FtsL^{H94} dihedral angles was observed from being α-helical

(bottom left quadrant) for WT FtsQLBWI to being β-strand-like in FtsQLBWI^{R167S} (top left quadrant).

Dihedral angles are color-coded from yellow to dark purple in time in the 1 µs MD trajectories. **(F)**

Single-molecule tracking of Halo-FtsB in a strain expressing FtsI^{R167S} shows that Halo-FtsB

molecules only exhibit the slow-moving population. The velocity histogram (gray bars) is best fit

by a single-population (red, $v_{slow} = 8.0 \pm 0.4$ nm/s, $\mu \pm$ s.e.m., $n = 112$ segments).

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024   **Figure 4**

1025

Figure 4. Interactions between FtsB, FtsL, FtsI, and FtsW position the FtsI anchor-loop

near the FtsW active site; FtsI anchor-loop position is correlated with FtsWI activity. (A, B)

Hydrogen bonding in the Lid region. Shapes indicate residues with hydrogen bonding during the

last 500 ns of FtsQLBWI simulation, with a cube specifying SF mutation FtsW[E289G], a tetrahedron

specifying DN mutation FtsL[R61E], and spheres indicate other residues involved (FtsI[R207, R213, Y214,]

[E219, D220], FtsL[R67, E68], FtsB[K23, D35]). Hydrogen bonding frequencies during the last 500 ns of the

simulation are indicated by the color bar. Two views of the Lid region are shown: (**A**) interactions

between the FtsI anchor domain (magenta), FtsL (red), and FtsB (blue), which position the anchor

domain above FtsW ECL4; (**B**) the same region rotated ~180° to show how interactions between

residues in ECL4 of FtsW (orange) with FtsL[R61], FtsI[R213], and FtsI[Y214] position FtsW ECL4 (orange)

below the FtsI anchor-loop (magenta). **(C–E)** Side and top views of a cavity on the periplasmic

face of FtsW (orange) containing the putative catalytic residue FtsW^{D297} (blue) lying below the

FtsW^{E289} residue (yellow; **D** shows SF FtsW^{E289G}) for WT FtsQLBWI (**C**), SF variant

FtsQLBW^{E289G}I (**D**) and DN mutant FtsQL^{R61E}BWI (**E**) complexes following 1 μs MD simulations.

With SF mutation FtsW^{E289G}, the FtsI anchor-loop (magenta) including FtsI^{Y214} (pink and white

spheres) moves away from the cavity, which is expanded as FtsW TM7 (green) tilts into the bilayer.

With DN mutation FtsL^{R61E}, the anchor-loop (magenta) moves and FtsI^{Y214} moves over the cavity,

which is stabilized by an interaction between FtsI^{R216} and FtsW^{E289}.

Figure 5

1045

Figure 5. FtsN^E binding reduces inhibitory interactions and induces conformational

changes observed in SF complexes. (A) Conformer of the FtsQLBWIN complex after 1-μs MD,

with FtsN^E shown in teal and detailed views of FtsN binding in the Hub region. Highly conserved

and essential residues of FtsN, FtsN^W83, FtsN^R84, and FtsN^Y85 are at the AWI interface in the Hub

region, close to residues previously identified in the FtsI[R167S] SF variant complex, including FtsI[V86,]

[I168] and FtsL[L86, N83, A90, E87, L84]. **(B)** Dihedral angle analyses of FtsL[G92] (left) and FtsL[H94] (right) in the

FtsN-bound complex showed similar backbone dihedral angle changes compared to those in the

SF mutant FtsQLBWI[R167S] (**Fig. 3E**). **(C)** The distribution of distances between sidechain centers

of geometry for FtsL[R67] and FtsI[D225] in the FtsQLBWIN simulation (light gray) exhibited a second,

decreased peak compared to that in the FtsQLBWI simulation (dark gray). The distance decrease

in FtsQLBWIN occurred with a conformational change in the last 300 ns of the MD simulation

(trajectory in **Fig. S23**), demonstrating tighter packing between the FtsI anchor domain and FtsL

in the presence of FtsN. **(D)** Interaction between the FtsI anchor-loop residue (magenta) FtsI[Y214]

and FtsW[E289] (orange) is disrupted in FtsQLBWIN as FtsI[D225] in the anchor domain increased its

interaction with FtsL[R67]. This interaction moves the sidechain of FtsI[Y214] (magenta) away from the

putative catalytic residue FtsW[D297] (orange), similar to what was observed in the SF complex

FtsQLBW[E289G]I (**Fig. 4D**). Note that FtsW[K370] forms a salt bridge with FtsW[D297] in FtsQLBWIN, but

not FtsQLBWI. **(E)** Ensembles of optimal paths calculated from FtsQ to FtsW[D297] (teal) showing

the 10 most optimal paths in the final 500 ns of MD for FtsQLBWI (top left), FtsQLBWI[R167S] (top

right), FtsQLBWIN (bottom left), and FtsQL[R61E]BWI (bottom right). Line colors correspond to the

number of paths connecting pairs of residues in the ensemble (colorbar). The DN variant FtsL[R61E]

(bottom right) eliminated all direct paths between FtsL to FtsW, shifting to paths through a small

loop extending from the FtsB helix. Addition of FtsN[E] increased the density of paths through FtsI,

which also includes paths through the Pivot region.