Euphorbia yadgirensis sp. nov., (Euphorbiaceae) from Karnataka state, India

Sarojini Devi N[1] and *Raja Kullayiswamy K[2]

1,2 Dharamavana Nature Ark, Cherlapalli, Hyderabad-500051

sarojini.botany@gmail.com; *raju.botany@gmail.com

Abstract:

Euphorbia yadgirensis is a new species from the scrub forest of Royangole village, Yadgiri

district of Karnataka, India is described and illustrated here. It belongs to the genus

Euphorbia, subg. *Chamaesyce*, sect. *Longistylae* and resembles *Euphorbia kadapensis*.

However, it differs in its decumbent habit (*vs*. erect), sympodial monochasium (*vs*.

dichotomous), laciniate stipules (*vs*. scaly), sutures absent on capsule (*vs*. present).

Keywords:

Euphorbia yadgirensis, *Longistyle, Chamaesyce*, Karnataka, Rayangole, scrub forest

Introduction:

Euphorbia L. includes more than 2000 species (Govaerts et al. 2000) which makes it the most diverse genus of the family Euphorbiaceae Juss. and one of the largest genera within the angiosperms (Horn et al. 2012, Webster 2014). Despite its great variation in morphology, ranging from small annual herbs to large trees, the genus is characterized by a synapomorphy unique among angiosperms. The cyathium (Horn et al. 2012) pseudanthial inflorescence consists of a cup-like involucre with glands along its rim (sometimes with appendages of several types) enclosing a single, central pistillate flower (female floret) surrounded by four or five staminate cymules (male floret) with reduced axes. Both pistillate and staminate flowers are highly reduced to a single pistil and single stamen, respectively (Radcliffe-Smith 2001, Prenner and Rudall 2007, Horn et al. 2012, Webster 2014). This may be further reduced in some species with unisexual cyathia, pistillate or staminate flowers to a pistillode or staminodes (Rizzini 1989). Genus *Euphorbia* plants show all known photosynthetic types C_3, C_4 and CAM (Webster et al. 1975) and a C_2 system that represents an early stage of C_3 to C_4 transition (Sage et al. 2011).

The subgenus *Chamaesyce* has the second largest species diversity containing around 600 taxa worldwide (Yang et al. 2012). It is the most species-rich lineage of *Euphorbia* in India with 36 percent of the species belonging to this subgenus among 8 subgenera that consist of 35 taxa under five sections: *Chamaesyce*, *Elegantes*, *Hypercifoliae, Longistylae* and *Sclerophyllae* (Binojkumar and Balakrishnan 2010) including the recently described *Euphorbia kadapensis* Sarojinidevi and Venkataraju (2014) that fall under the section *Langistylae*. The section *Longistylae* was erected by Binojkumar and Balakrishnan (2010) to accommodate the endemic therophytic Indian taxa with type *E. longistyla* Boissier (1860).

This section was reduced to subsection *Hypericifoliae* by Yang et al. (2012) based on

molecular data.

While doing intensive exploration of the genus *Euphorbia* from India, the present taxon

was collected from the northern part of Karnataka and is here in described as a new species

under genus *Euphorbia* subgen. *Chamaesyce*, sect. *Longistylae*.

Materials and Methods:

Specimens were collected from the scrub forest of Royangole village, Yadgiri district,

Karnataka state, treated with 80% ethanol, and herbarium specimens were prepared following

the standard method (Santapau 1958, Jain and Rao 1977). Flowers and fruits were used for

microscopic observations and imaging and preserved in 70% ethanol for future reference.

Microscopic images were taken with an Olympus stereo microscope SZX16, DP74 camera

attached. Seeds of the species have been provided to the Dharmavana Nature Ark (DNA) for

ex-situ conservation.

Taxonomic treatment:

Euphorbia yadgirensis **SND. & RKS.** *sp. nov*. **(Fig. 1, 2 & 3)**

Euphorbia kadapensis sensu Raja P. et al., JoTT 12(14):17045. 2020.

Type: India (Ind.), Karnataka (KA), Yadgiri district, Shorapur, Royanagole, 16˚16ʹ43.36ʺN,

76˚24ʹ11.39ʺE, elevation 479 m a.s.l., 11 Oct 2022, RKS HJCB-2090 (holotype MH,

iso-BSID, SKU, HDNA).

Other specimens: Ind., KA, Royanagole, Yadgiri district, 16.280°N; 76.393°E, alt 484m,

01.ix.2017, P. Raja 2407 at St. Josephʹs college, Tiruchirappalli, Tamil Nadu.

Diagnosis:

Euphorbia yadgirensis is an annual decumbent herb growing between rocks in scrub forest, stem glabrous, sympodial-monochasium branching; stipules laciniate, limb of gland bi- or tri-fid one fourth of its length, capsules trigonous prominently keeled, stigmas slightly bifid at apex, seeds transversely deeply rugulose. *Euphorbia yadgirensis* resembles *E. kadapensis* Sarojin. & R.R.V. Raju in filiform branches, elliptic leaves, dissected limb, long style, and trigonous capsule, but differs in habit (decumbent versus erect), height of the plant (20 cm versus 30 cm), branching type (sympodial-monochasium versus dichotomous), stipules (laciniate, 2 mm versus scaly, 1 mm), involucre (3 mm versus 4 mm), limb (bi or trifid one by fourth of its length versus bi or trifid one by two of its length), number of staminate florets (9−12 versus 15−18), bract of male floret (laciniate versus simple), capsules (ridges absent along the sutures versus prominently ridged along the sutures), seeds (transversely deeply rugulose versus transversely slightly rugulose).

Etymology:

The species is named after the type locality of the Yadgiri district in the north of Karnataka state. The type locality is dominated by granite rocks and scrub brush.

Description:

Hemicryptophytic decumbent glabrous herb up to 20 cm tall; latex milky. Stem sympodial-monochasium branching. Branches terete, glabrous, filiform towards apices, internodes 2–3.5 cm long, thickened at nodes; stipules interpetiolar, laciniate, 2 mm long. Leaves simple, opposite, equal, sub-sessile; lamina oblong-elliptic, 2(0.5)–2.5 × 1(0.3)–1.5 cm; base oblong (rounded) [apical leaves characters mentioned in parentheses], margin serrulate, apiculate; mid-nerve prominent, lateral nerves obscure, 4–8 pairs; petiole 3 mm long. Cyathia terminal,

up to 10 mm long; peduncles 2.4 mm long; involucre turbinate, 3 mm long, glabrous, green;

involucral lobes 5, laciniate, each 5–8-lobed, appendages of glands 4, 2–3-fid, one by fourth

of its length, 1.5 mm long, glands 4, oblong, green, stipitate, stalks hairy. Staminate florets 9–

12, exserted, 6 mm long, bracteate, bract laciniate, 4 mm long, anther lobes sub-globose,

yellow, and dehiscing transversely. Pistillate flowers laterally pendulous, 10 mm long,

glabrous, pedicel 4 mm long, ovary tricarpellate, 3×5 mm, styles 3, 3.2 mm long, free from

the base, stigmas 3, slightly bifid at apex. Capsules schizocarpic, 2.4 mm long, trigonous,

distinctly keeled, fruiting pedicels recurved. Seeds 3, ecarunculate, brown, 2×1.5 mm, ovoid-

oblong, tetragonal, transversely and obscurely furrowed, truncate at base, transversely

rugulose.

Phenology:

Flowering and fruiting from September to November.

Habitat and distribution:

Euphorbia yadgirensis grows between granite rocks in black gravelly soil, scrub forest near

Rayangole, Yadgiri district, elevation ranges from 470 to 480 m a.s.l. The area is dominated

by the scrub species *Ziziphus mauritiana* Lam., *Maytenus emarginata* (Willd.) Ding Hou,

Searsia mysorensis (G.Don) Moffett, and herbs *Tridax procumbens* L., *Justicia procumbens*

L., *Heteropogon contortus* (L.) P.Beauv. ex Roem. & Schult., *Barleria prionitis* L.,

Andrographis longipedunculata (Sreem.) L. H. Cramer, *Allmania nodiflora* (L.) R.Br. ex

Wight, *Cyperus rotundus* L. growing along with the present species. Thus far, the species has

been collected only from the type locality where there is a threat of grazing and forest fires.

Comparison:

This new species resembles *Euphorbia kadapensis* in its dissected limbs, filiform stem, leaf

shape and long style, but differs in many other morphological characters that are tableted

below along with another closely related species of this section (table 1 and figure 3).

Table1: Morphological comparison of *Euphorbia* sister taxa (the same morphological

characters can be seen in Figure 3 and supporting file)

Characteristics	*Euphorbia yadgirensis*	*E. kadapensis*	*E. longystyla*
Habit	decumbent	erect	erect
Branching	sympodial-monochasium	dichotomous	pseudo-dichotomous
Leaf Margin	serrulate from the base	serrulate from the middle	entire or serrate
Cyathium	terminal, 10 mm long	terminal and axillary 6 – 9 mm long	terminal 8.5 mm long
Limb of Gland	2(3)-fid ¼ of its length	2(3)-fid ½ of its length	Multi-fid from the base
Male floret bract	laciniate	simple	simple
Stigma	slightly bifid	simple	simple
Fruit	2.4 mm long, without ridges along sutures	3 mm long, with ridges along sutures	4 mm long, without ridges along sutures

Artificial key to the section *Longistylae* from India

1. Plants hairy, stem ribbed; seeds wrinkled…………………………………… *E. hispida*

 – Plants glabrous, stem terete; seeds transversely obscurely furrowed or smooth… (2)

2. Branches always filiform towards the apex; leaves elliptic to oblong-elliptic; styles

   more than 1.5 mm long.…………………………..………………………… (3)

 – Branches not filiform towards the apex; leaves oblong or oblong-lanceolate or ovate-

   oblong; style less than 1.5 mm long. …………………………………………….. (6)

3. Branches minutely velutinous at base; limb of glands dissected…………………… (4)

 – Branches not velutinous; limb of glands entire, minutely sinuate…… *E. senguptae*

4. Limb of glands deeply multi-fid from the base up to 12-fid........................ *E. longistyla*

 – Limb of glands slightly dissected ½ to ¼ th of its length, 2–3-fid………………… (5)

5. Plants erect, dichotomously branched; stipules scaly and connate; male floret bract

   simple; limb of gland bifid or trifid from the middle; capsule with prominent ridges

   along sutures; stigmas simple………………………………………… *E. kadapensis*

 – Plants decumbent, sympodial-monochasium branching; stipules laciniate; male floret

   bract laciniate; limb of gland bifid or trifid one fourth of its length; capsule without

   ridges along the sutures; stigmas slightly bifid………………………. **E. yadgirensis**

6. Stem stramineous; leaves too larger on stem than branches; limb of glands 1 mm

   across……………………………………………………………… *E. jodhpurensis*

 – Stem pink or pale pink; all leaves are same as main stem; limbs of glands 2 by 1.5

   mm…………………………………………………………………………… (7)

7. Stem unbranched or rarely dichotomously branched; leaves 3 cm or more long; cyathia

   pubescent…………………………………………………………… *E. katrajensis*

 – Stem branched; leaves less than 3 cm long; cyathia glabrous…………………… (8)

8. Height of the plant 30–60 cm; leaves silvery white below; limb of glands irregularely and shallowly wavy…………………………………………………… *E. erythroclada*

– Height of the plant below 30 cm; leaves pale green below; limb of glands entire or slightly wavy………………………………………………………………… (9)

9. Limb of glands white or yellowish; fruits with 2 or 3 prominent wings on cocci……………………………………………………………… *E. notoptera*

– Limb of glands pink, rosy or pinkish white, never with yellow shade; fruits ridged or keeled, but without wings on cocci………………………………………… (10)

10. Leaves oblong to ovate-oblong, sub-acute to obtuse at apex, 5–11 mm long; capsule prominently broadly and obtusely keeled, distinctly ridged along sutures ………… ………………………………………………………………….. *E. concanensis*

– Leaves oblong-lanceolate, acute to sub-acute at apex, 10–25 mm long; capsule faintly narrowly and acutely keeled, not ridged along the sutures…. *E. deccanensis*

Acknowledgements:

The authors are grateful to the Dharmavana Nature Ark for encouragement and providing the necessary facilities and resources. Thanks to Dr. Kanchian Gandi, of Harvard University for the confirmation of the specific epithet. Also our thanks to Dr. Jahnavi Joshi, Senior Scientist, CCMB for permission for microscopic imaging.

Reference:

Binojkumar, M. S. and Balakrishnan, N. P. 2010. The genus *Euphorbia* L. (Euphorbiaceae) in India. A taxonomic revision. Bishen Sing Mahendra Pal Singh, Dehra Dun.

Boissier, P. E. 1860. Centurria Euphorbiarum. Hermann, Leipzig and Ballierie, Paris.

Govaerts, R. et al. 2000. World checklist and bibliography of Euphorbiaceae (with Pandaceae). – Royel botanical gardens Kew. 2: 417–921.

Horn, J. W. et al. 2012. Phylogenetics and the evolution of major structural characters in the giant genus *Euphorbia* L. (Euphorbiaceae). – Mol. Phylogenetics Evol. 63: 305–326. https://doi.org/10.1016/j.ympev.2011.12.022

Jain, S. K. and Rao, R. R. 1977. A hand book of field and herbarium methods. – Scholarly publication, New Delhi.

Prenner, G. and Rudall, P. J. 2007. Comparative ontogeny of the cyathium in *Euphorbia* (Euphorbiaceae) and its allies: exploring the organ, flower, inflorescence boundary. – Am. J. Bot. 94: 1612–1629. https://doi.org/10.3732/ajb.94.10.1612

Radcliffe-Smith, A. 2001. *Genera Euphorbiacearum*. –Kew Publishing, Kew.

Raja, P. et al. 2020. New distribution record of two endemic plant species, *Euphorbia kadapensis* Sarojin. & R.R.V. Raju (Euphorbiaceae) and *Lepidagathis keralensis* Madhus. & N.P. Singh (Acanthaceae), for Karnataka, India. – JoTT 12: 17045–17048. DOI: https://doi.org/10.11609/jott.4978.12.14.17045-17048

Rizzini, C. T. 1989. Cactiform species of *Euphorbia* from Brazil (Euphorbiaceae). – Rev. Bras. Biil. 49: 979–997.

Sage, T. L. 2011. The occurrence of C_2 photosynthesis in *Euphorbia Chamaesyce*

(Euphorbiaceae). – J. Exp. bot. 62. 3183–3195.

Santapau, H. 1958. Botanical collectioris manual. – Bot.Surv.India, Calcutta.

Sarojinidevi, N. and Venkataraju, R. R. 2014. *Euphorbia kadapensis* (Euphorbiaceae) a new

species from southern India. – Phytotaxa 181:179–183.

Webster, G. L. et al. 1975. Systematics of photosynthetic carbon fixation pathways in

Euphorbia. – Taxon 24: 27–33.

Webster, G. L. 2014. Euphorbiaceae. *In*: Kubitzki, K. (ed.). The Families and Genera of

Vascular Plants. – Springer Berlin, Heidelberg. 11: 51–216. https://doi.org/10.1007/978-

3-642-39417-1_10

Yang, Y. et al. 2012. Molecular phylogenetics and classification of *Euphorbia* subgenus

Chamaesyce (Euphorbiaceae). – Taxon 61: 764–789.

Figure legends:

Figure 1. *Euphorbia yadgirensis sp. nov.* (A) habit, (B) young cyathium, (C) stipules, (D) involucral lobes, limbs and glands, (E) female floret, (F) capsule top view, (G) capsule bottom view, (H) seeds.

Figure 2. *Euphorbia yadgirensis sp. nov.* (A) habit, (B) small branch, (C) cyathium at shoot apex, (D) cyathium, (E) cyathium split opened, (F) portion of involucral cup showing lobes, glands and limbs, (G) mature cyathium, (H) capsule bottom view, (J) capsule top view, (K) ripe capsule, (L) seeds, (M) stipule, (N) male floret.

Figure 3. A-H–*Euphorbia yadgirensis*. (A) cyathium at shoot apex, (B) portion of involucral cup showing lobes, glands and limbs, (C) stipules, (D) male floret (E) capsule bottom view, (F) capsule top view, (G) mature cyathium, (H) seeds. S-Z–*Euphorbia kadapensis*. (S) cyathium at shoot apex, (T) portion of involucral cup showing lobes, glands and limbs, (U) stipules, (V) male floret (W) capsule bottom view, (X) capsule top view, (Y) mature cyathium, (Z) seeds.

Fig: 1

Fig: 2

Fig:3

Euphorbia longistyla Bioss. images for reviewer's understanding purpose

Habit

Young cyathium

Mature cyathium

Capsule

Stipules

Portion of involucral cup showing lobes, glands and limbs