Genome-wide circadian gating of a cold temperature response in bread wheat

Calum A. Graham[1,2], Pirita Paajanen[1], Keith J. Edwards[2], Antony N. Dodd[1,3]

1. Department of Cell and Developmental Biology, John Innes Centre, Norwich Research

Park, Norwich NR4 7RU, UK.

2. School of Biological Sciences, University of Bristol, Bristol Life Sciences Building, 24

Tyndall Avenue, Bristol BS8 1TQ, UK.

3. Corresponding author, antony.dodd@jic.ac.uk

Keywords: circadian regulation, temperature response, wheat, cereals.

Word counts: Intro, Results, Discussion = 4750.

Abstract

Circadian rhythms coordinate the responses of organisms to their daily fluctuating

environments, by establishing a temporal program of gene expression. This schedules

aspects of metabolism, physiology, development and behaviour according to the time of day.

Circadian regulation in plants is extremely pervasive, and is important because it underpins

both productivity and seasonal reproduction. Circadian regulation extends to control of

environmental responses through a regulatory mechanism known as circadian gating,

whereby the circadian clock regulates the response to environmental cues such that the

magnitude of response varies according to the time of day. Here, we show that there is

genome-wide circadian gating of responses to cold temperatures in plants. By using bread

wheat as an experimental model, we establish that circadian gating is crucial to the

programs of gene expression that underlie the environmental responses of a crop of major

socioeconomic importance. Furthermore, we identify that circadian gating of cold

temperature responses are distributed unevenly across the three wheat subgenomes, which

might reflect the geographical origins of the ancestors of modern wheat.

Teaser

There is genome-wide circadian gating of a response to low temperatures in a crop of major

socioeconomic importance.

Introduction

Circadian rhythms are biological cycles with a period of about 24 h that persist in the

absence of environmental cues, and establish a temporal structure within biological systems.

A feature of circadian rhythms is that they determine the time of day (phase) when specific

processes occur (*1, 2*). The process of entrainment aligns the phase of the circadian clock

with the phase of the environment and, in turn, the circadian clock extensively influences the

phasing of gene expression, metabolism and physiology (*3*). One way that this occurs in

plants is through clock controlled gene promoters, which temporally coordinate transcript

accumulation according to the time of day (*1, 4-8*). The regulation of cellular processes by

circadian clocks is important for crop performance, and understanding circadian regulation in

crops forms an important part of the strategic design of climate change resilient crops (*9-12*).

Fluctuating spring temperatures and unpredictable cold periods caused by climate instability

represent a threat to cereal crops worldwide (*13-15*). During multiple stages of development,

wheat is susceptible to low temperatures (*16-18*), including during seedling establishment

(*19, 20*). Low temperatures trigger genome wide responses in wheat that can reduce

photosynthetic capacity, cause photoinhibition, and consequently reduce yield (*18, 21-25*).

Therefore, wheat represents an excellent model to understand the processes that underlie

responses of plants to low temperature conditions, because this has a direct impact upon

crop yield.

One way that the circadian oscillator influences responses to environmental cues, such as

temperature fluctuations, is through circadian gating. Circadian gating is the process

whereby the circadian clock regulates the magnitude of a response to the environment, such

that the response is restricted to certain times of day, or varies according to the time of day

(*26, 27*). In plants, there are widespread examples of the circadian gating of responses to

high and low temperature cues (*9, 28-36*), with temporal gating also occurring in cultivated

rice fields and wild plant populations (*37, 38*). We do not know whether circadian gating of

cold temperature responses occur at a genome-wide scale. This remains an important open

question, because the pervasive nature of circadian gating of temperature responses makes

it a core part of understanding how plants- including crops- respond to their fluctuating

environments.

In hexaploid bread wheat, around 30% of the transcriptome is regulated by the circadian

clock (*39*). Circadian clock gene loci such as *TaPpd* (orthologous to Arabidopsis *PRR3* or

PRR7) determine agriculturally-important photoperiod responses of domesticated wheat (*40-*

43). The complexity of this signal integration is increased in polyploid crops such as wheat

by the presence of multiple gene paralogs or homoeologs that can have differential patterns

of circadian regulation (*39, 44*). In hexaploid wheat, over 50% of genes (approximately

50,000 genes) occur as sets of three gene homoeologs termed triads (*45, 46*), providing an

informative genomic structure to study environmentally-responsive gene expression within

polyploid genomes.

We investigated the involvement of circadian regulation in the genome-wide responses of

hexaploid bread wheat to low temperatures. Given that the cold-responsive CBF pathway is

gated by the circadian clock in Arabidopsis (*28*) and a third of wheat genes are circadian-

regulated (*39*), we reasoned that there might be a pervasive influence of the circadian clock

upon cold temperature responses. Using wheat as a model to investigate this question also

provides insights into a species of vast socioeconomic importance. It also provides insights

into the integration of circadian and environmental cues in polyploids. Taken together, our

data identify a major role for circadian regulation in shaping the cold-responsive

transcriptome in plants.

Results

The time of day affects the number of genes that respond to cold temperature conditions

We wished to discover whether there is circadian gating of responses to temperature cues in

hexaploid bread wheat (*Triticum aestivum).* We used low temperature treatments as an

experimental model, because circadian gating of responses to cold has been detected for

certain genes in Arabidopsis (*28, 47, 48*). Spring wheat varieties are more cold-sensitive

than winter wheat (*49, 50*), and are exposed to low spring temperatures during seedling

establishment. Therefore, we selected for experimentation a cold-responsive hexaploid

spring wheat cultivar (Cadenza) that has no vernalization requirement, some low

temperature tolerance, and good genome sequencing coverage and germplasm resources.

To test for the occurrence of circadian gating, short cold treatments were applied to separate

batches of wheat seedlings under free running (constant) conditions (Fig. 1A). This

comprised identical three-hour cold treatments (4^{o}C), at six sequential timepoints, following

transfer to free running conditions (Fig. 1A). After each treatment, tissue was harvested for

RNA isolation from the second leaf of control temperature and cold-treated seedlings, with

each seedling sampled once only (Fig. 1A). Transcript count values (isoforms) were

condensed to the gene level, so transcript abundance changes described here refer to

changes at the gene level.

The number of differentially expressed genes (DEGs) in response to the cold treatment was

different at each at each timepoint (log fold change > 0.5, false discovery rate < 0.05; Fig.

1B). This indicates that the time of subjective day determines the cold-induced transcriptome

in wheat (Fig. 1B). The greatest number of genes were differentially expressed in response

to cold at zeitgeber time (ZT) 48 (ZT48), with 4350 and 3403 genes significantly up or down-

regulated, respectively (Fig. 1C). Two-thirds fewer genes were differentially expressed in

response to a cold treatment at ZT28 or ZT44 (Fig. 1C), despite a similar total number of

transcripts detected at these timepoints (Table S1). There was a large difference in the

direction of change of DEGs at ZT36 and ZT40, with the number of up-regulated genes

comparable to those at ZT32 and ZT48, but a smaller number of down-regulated genes that

was comparable to the number at ZT28 and ZT44 (Fig. 1C). These data suggest the wheat

transcriptome is more responsive to cold around the middle of the subjective day (ZT32) and

towards the end of the subjective night (ZT48) (Fig. 1C).

There were fewer cold-downregulated than cold-upregulated genes at each timepoint,

whereas the proportion of downregulated genes that were unique to each timepoint was

consistently greater than the proportion upregulated genes that were unique to each

timepoint (Fig. 1D). This suggests that genes downregulated in response to cold are

especially influenced by the time of day. Interestingly, the proportion of cold-responsive

genes that were unique to each timepoint corresponded to the total number DEGs at the

timepoint (Fig. 1C), suggesting that there is a relationship between the total number of cold-

responsive genes and the number of genes that are cold-responsive at specific times of day.

The nature and cold-sensitivity of cold-responsive genes fluctuates across the 24 h cycle

We wished to understand whether the composition of the cold-responsive transcriptome

changes over the 24 h cycle. To achieve this, we compared the identity of the genes that

were cold responsive at each timepoint. Four of the largest six groups of DEGs were cold-

induced at specific timepoints, so more genes responded to cold at specific (unique) times

than were cold responsive at all timepoints (Fig. S1A). Furthermore, the top four sets of

transcripts that were downregulated by cold were unique to single timepoints (Fig. S1B).

Therefore, under free running conditions, the set of genes that responds to cold changes

across the day. Some transcripts were cold responsive at single timepoints, some across

part of the 24 h cycle, and some were cold-responsive regardless of the time (Fig. S1A, B).

Because the set of cold responsive genes changes across the 24 h cycle (Fig. 1D; Fig. S1A,

B), we reasoned that there might be circadian gating of the responses to cold of the wheat

transcriptome. There were a variety of responses to the time course of cold treatments, with

cluster analysis revealing a large proportion of cold-responsive genes to have temporally-

fluctuating response profiles (Fig. 1E). For example, certain genes are upregulated by cold in

the subjective day and down-regulated by cold during the subjective night (Groups 5 and 6;

Fig. 1E), whilst other genes are downregulated by cold during the subjective day, and

upregulated or have little response to cold during the subjective night (Group 3; Fig. 1E).

This indicates that the magnitude of transcriptional responses to cold in wheat can vary

across a 24 hour cycle, suggesting that the circadian clock gates the transcriptional

response to cold of a variety of genes.

The responsiveness of signalling genes can be gated by the circadian clock

To assess how the time of day affects cold-sensitivity of key signalling components, four

genes that are known to be circadian regulated either in Arabidopsis or wheat were studied

as examples. The hexaploid nature of the bread wheat genome means that many transcripts

are present as triads, with gene homoeologs on the A, B and D subgenomes. Therefore, all

three homoeologs were considered to obtain a representative view of how a signalling

component responds to a stimulus.

C-repeat binding factors (CBFs) are well-described cold responsive transcription factors (*51,*

52), with up to 25 CBFs present in wheat (*53*). Certain *CBF* transcripts have a circadian

rhythm in barley and wheat (*39, 54*), and in Arabidopsis the cold-induction of CBFs is gated

by the circadian clock (*28*). Our data indicate that circadian gating of *CBF* cold-

responsiveness is conserved in wheat and can extend across homoeolog triads, such as for

TaCBFIVc-B14, which is a cold-responsive *CBF* ortholog with high homology to *AtCBF2*

(Fig. 2A, B, C) (*24, 53*).

The B-BOX CONTAINING PROTEINs (BBXs) participate in responses to fluctuating

environmental conditions (*55*), such as the circadian modulation of *AtBBX17* transcript

responses to heat (*34, 56*). Transcripts from the *TaBBX17 7ABD* triad, which has highest

homology to *AtBBX17*, share with Arabidopsis rhythmic accumulation under control

temperature conditions (Fig. 2D, E, F). The response to cold varied across the triad, with

cold induced upregulation of *TaBBX17 7A* at the start of the subjective day, compared to

abundance of *7B* and *7D* at the end of the subjective day (Fig. 2D, E, F). This example

suggests that features of circadian gating are not always conserved across triads.

In Arabidopsis, the circadian clock regulates photosynthesis and starch metabolism, with

these thought to provide a selective advantage to plants (*57-59*). A range of related

Arabidopsis transcripts encoding starch degradation enzymes are circadian regulated, but

this is not always conserved in wheat, where certain orthologs of β-amylase (BAM, e.g.

BAM3) have a circadian rhythm, whereas *BAM1* does not (*39*). Under our conditions,

TaBAM1 triad and *TaBAM3* triad were cold-responsive during the subjective day (Fig. 2G-L).

TaBAM1 was not responsive to cold during the subjective night, whereas *TaBAM3* was (Fig.

2G-L). The cold-induced relative abundance of *TaBAM3 1B* and *TaBAM3 1D* was

approximately ten-fold greater than the corresponding homoeologs of *TaBAM1* (Fig. 2H, I, K,

L), supporting the notion of a role for BAM3 in cold conditions.

Circadian gating of transcriptional responses to cold is extensive, and not restricted to

specific times of day

We considered circadian gating to be defined as a 24 h fluctuation in the sensitivity of a

transcript to a stimulus, such that the magnitude to an identical stimulus oscillates with a

24 h cycle. Given that our data identified that different subsets of transcripts are cold-

responsive at different times of day (Fig. 1B, D, E), we reasoned that there might be

widespread circadian gating of the cold-responsive transcriptome. To investigate this, we

obtained a measure of the cold-sensitivity of each transcript, at each timepoint, by

calculating the difference between the mean CPM of the transcript between control and cold

temperature conditions (Fig. 3A). This provided a measure of the response to cold, at each

timepoint, that we termed ΔCPM (Fig. 3A). Using this measure, a 24 h fluctuation in the

sensitivity of any given transcript to the cold treatment will manifest as an oscillation in

ΔCPM (Fig. 3A, lower panel). We tested whether there was a circadian rhythm in this cold-

sensitivity profile (meta2d analysis for rhythmicity) (Wu *et al.*, 2016). Circadian rhythms of

cold sensitivity (ΔCPM) were detected for 4388 transcripts, which we filtered to include only

transcripts that were differentially expressed in response to cold at least once across the

time course. This yielded 1677 transcripts that we consider to undergo circadian gating of

their responses to cold in wheat (Fig. 3B).

We found that this set of transcripts divides into three groups: circadian gated cold-

inductions (where mean[ΔnormalisedCPM] > 0.5), circadian gated cold-reductions

(mean[ΔnormalisedCPM] < -0.5), and cold induced delays in transcript cycling

(mean[ΔnormalisedCPM] between -0.5 and 0.5) (Fig. 3A). In the case of a delay, the

transcript is cold-induced at certain times of day and cold-repressed at other times. Within

these cold response types, we calculated the estimated time of maximum responsiveness

(ETMR) using phase estimates from meta2d (Fig. 3C). This approach allowed us to estimate

ETMR that fell between sampling timepoints, thereby overcoming some limitations of the

sampling frequency. For circadian gated reductions (Fig. 3A), the time of maximum cold-

responsiveness was the trough of the oscillation.

The greatest number of transcripts with a rhythmic response to cold were those whose

cycling was delayed by the cold treatment (Fig. 3C, D). Furthermore, over 500 transcripts

had circadian gating of their induction by cold (Fig. 3C, D). Within these transcripts, the most

common ETMR was between 16 h and 20 h (Fig. 3D), with a large proportion of the

transcripts also phasing between 4 h and 8 h. The largest proportion of cold-repressed

transcripts had greatest cold-sensitivity during the subjective night (Fig. 3D).

This indicates that there is transcriptome-wide circadian gating of the cold-responsive

transcriptome in bread wheat. These responses are not restricted to a particular direction of

change (upregulation *vs.* downregulation), and different transcripts have maximum cold-

responsiveness at different times of day. Our analysis might under-estimate the number of

circadian gated cold-responsive genes, because a longer and higher resolution time course

would provide greater statistical power.

Circadian gating of cold responses have a subtle subgenome bias

The wheat subgenomes arose from hybridisation of three diploid ancestors, resulting in a

circadian oscillator that likely incorporates more components than diploid species. No

specific subgenome is favoured across the circadian regulated transcriptome of wheat (*39*),

although circadian regulated paralogs have differential expression patterns in other crops

(*44*). Furthermore, environmentally-responsive gene triads do not always have expression

that is balanced across the subgenomes (*46*). Therefore, we wished to determine whether

there was subgenome bias in the circadian gating of transcriptomic responses to cold.

The relative proportion of transcripts originating from A, B and D subgenomes remained

relatively constant during the day for circadian gated inductions and those with delayed

cycling (30 - 40%; Fig. 4A). However, during the subjective night, the largest proportion of

circadian gated cold induced transcripts derived from the D subgenome (ETMR > 12 h) (Fig.

4A). This was restricted to cold-induced transcripts, and did not occur for transcripts with

with delayed cycling. The number of genes with circadian-gated repression by cold was

rather small, so the proportional subgenome contribution to cold-repressed transcripts is

noisy (Fig. S2).

Over 60% of circadian-gated cold induced transcripts with a ETMR before 16 h can be

assigned to a triad, with a smaller proportion for transcripts with ETMR over 16 h (Fig. 4B).

Approximately 10 % of transcripts with ETMR < 4 h, 4-8 h, 12-16 h and 16-20 h were

detected as complete triads of three homoeologs, but no complete triads were present in the

ETMR groups 12-16 h and > 20 h (Fig. 4B). Only single transcripts from triads had circadian

gated responses to cold in the 8-12 h ETMR group, suggesting differential behaviour within

triads (Fig. 4A, Fig. 4B). Furthermore, it appears that the time of maximum responsiveness

of some triads is spread across multiple phases, and homeolog-dependent (Fig. 4B). This

suggests there is sometimes circadian gating bias within triads, with only certain

homoeologs responding to the cold stimulus at particular times of day.

The cold-responsiveness of each homoeolog to the overall cold responsiveness of each triad

was assessed (*46*). The relative contribution of each subgenome to the cold-responsiveness

of the triad determined the position of the triad within a ternary plot (example in Fig. 4C). In

general, no subgenome obviously contributed more or less than the other subgenomes (Fig.

4D). Many triads had dynamic responses to cold, in that the contribution from each

subgenome to total transcript level changed after the cold treatment (Fig. S3). The variation

in the contribution of each subgenome to triad cold responsiveness, in each timing group,

suggests that there is no bias towards a particular subgenome despite individual gene

homeologs occupying different EMTR groups.

Genes associated with light responses can be circadian gated to the subjective day

Gene ontology-term analysis identified multiple biological processes associated with

responses to light as over-represented in transcripts that had a maximum cold induction

during the subjective day (ETMR 0-12 h). Processes such as "cellular responses to blue

light", "cellular responses to UV-A" and "anthocyanin containing compound biosynthesis"

were enriched in multiple subjective day ETMR groups (Fig. S3A; adjusted p < 0.001).

Enrichment of terms such as "photoprotection" and "stress-activated MAPK cascade"

suggests the cold stimulus elicited a broad response when delivered in the subjective day, at

a time when light and low temperatures may be expected simultaneously.

There was a clear split between the GO-terms that were enriched in subjective day ETMR

groups and subjective night ETMR groups. "Cold acclimation" is shared across two

subjective night groups (Fig. S4A; adjusted p < 0.001), suggesting that night-time low

temperatures could have a key role in adapting wheat to cold seasonal weather. Also

enriched are terms associated with DNA transcription, which can be an indication of

preparation for a large shift in gene expression (Fig. S4A; adjusted p < 0.001). "Photosystem

II" associated processes were enriched in subjective night ETMR group 16 – 20 h, which

might relate to unexpected exposure to continuous light conditions (Fig. S4A; adjusted p <

0.001). Whilst GO-term analysis interpretation requires caution, it is possible that restriction

of the cold response of certain processes to specific times of day provides an advantage to

wheat by ensuring that energy demanding cold-responsive processes are not induced at

inappropriate times of day (*60*).

Cold treatments delay the oscillation of certain circadian clock components

We noticed that a variety of transcripts had a delay in their cycling in response to the cold

treatment (e.g. Fig. 5A, B). We wished to systematically identify the set of transcripts that

had these delays. We developed an analytical method that used the ΔCPM to consider the

relationship between the transcript levels at adjacent timepoints (Fig. 5C; see methods for

details). Over three-quarters of the 882 transcripts that have a delay and rhythmic ΔCPM

have a delay ratio of < 1, supporting the interpretation that the cycling of these transcripts is

delayed relative to the control (Fig. 5D). A similar proportion of the 3849 transcripts with

rhythmic control and cold CPM profiles had a delay ratio of < 1, suggesting a temporal delay

in the profile of these transcripts also (Fig. 5E). A possible explanation is that the cold

treatment maintained a relatively constant level of these transcripts for the duration of the

treatment, resulting in a pause in the rhythm of the transcript level that is detected as a time-

delay. Only 47% of transcripts with circadian gated induction responses had delay ratios < 1,

suggesting around half of the transcripts with a gated response to cold did not have a cold-

induced delay in cycling.

The GO-term "circadian rhythm" was enriched within the genes that had rhythmic CPM

profiles and a delay ratio < 0.5 in response to cold (adjusted p < 0.001; Fig. S5). This

suggests that the cold treatment might delay the rhythm of certain circadian oscillator

components. All homoeologs of six triads of circadian clock genes (*39*), including morning

phased *TaLHY* and evening phased *TaGI*, had a delay ratio less than 1 (Fig. 5F). Therefore,

the cold treatment delayed the oscillation in the expression of these triads (Fig. 5F).

Homoeologs in the triads *TaELF3*, *TaELF4*-L and *TaTOC1* had a mixture of delay ratios,

indicating that the cold treatment did not equally affect the cycling of homoeologs of all

circadian clock transcripts (Fig. 5F).

Cold treatments did not have a consistent impact upon all circadian oscillator components.

For example, cold had little effect upon expression of the *TaLUX-Lb* and *TaLUX/BOA* triads,

which had lower mean ΔCPM before adjustment in all three homoeologs (Fig. 5F).

Therefore, cold does not appear to affect all circadian clock transcripts equally, as

exemplified by *TaGI* having delayed expression, whereas *TaLUX/BOA* does not (Fig. 5G).

This supports the idea that cold has an unbalanced effect upon circadian clock components,

reflecting a complex relationship between the circadian clock and cold.

Discussion

A dual effect of cold on both the clock and clock-controlled genes

Here, we have identified for the first time extensive genome-wide circadian gating of the response to a cold temperature treatment in bread wheat. This demonstrates that the circadian gating of responses to cold of single genes (*28*) extends across the entire genome, and supports reports of the circadian gating of responses to high temperature stress in Arabidopsis (*32, 61*). Given that the circadian clock gates the responses of certain genes to cold in Arabidopsis, and the considerable phylogenetic distance between wheat and Arabidopsis, we reason that the circadian gating of low temperature responses might be conserved across the angiosperms.

Our results demonstrate the importance of circadian regulation in shaping transcriptomic responses to short-term cold. The diversity of temporal responses to cold suggests considerable complexity in the integration of low temperature and circadian signals (Fig. 1E, Fig. 3). The presence of a circadian rhythm for a transcript under control temperature conditions does not mean that its cold-responsiveness is gated by the circadian clock, because some circadian regulated transcripts are not cold-responsive (Fig. 6A; Fig. S6A). Likewise, some transcripts are cold-responsive, but this is independent from circadian regulation because these transcripts are cold-induced by a similar magnitude, irrespective of time of day (Fig. 6B; Fig. S6B). In one form of circadian gating, the circadian clock restricts the magnitude of response to the stimulus to certain times, and the transcript is completely unresponsive to cold at other times (Fig. 6C, Fig. S6C). This type of circadian gating is bidirectional, because suppression of transcript levels by cold can also be circadian-gated (Fig. 3C). In another form of circadian gating, a circadian modulation of sensitivity occurs across the day, such that a response to cold always occurs, but the response is greater at certain times (e.g. *TaBAM3*; Fig. 3J-L). The phase of maximum cold responsiveness of some transcripts is sometimes aligned with the phase of the underlying circadian rhythm

under control temperature conditions, but in other cases (e.g. *TaBAM3*) the phase was

misaligned (Fig. 3J-L; Fig. 6D; Fig. S6D).

Short cold treatments affect the dynamics of certain wheat circadian clock components, by

delaying their oscillation for approximately the duration of the cold treatment (Fig. 6E). This

delay pervades hundreds of other circadian-regulated cold-responsive transcripts (Fig. 6F). It

is possible that the cold-induced delay in the circadian oscillator causes the delay in the

response of downstream circadian-regulated transcripts. Interestingly, these delays occur for

some circadian clock genes (e.g. *TaPRR73* and *TaGI*) but not others (e.g. *LUX* orthologues)

(Fig. 5F). One potential explanation for this bifurcating response is that the stability of certain

transcripts is insensitive to cold, so the transcript level continues to change under low

temperature conditions. Another explanation could be that different cell types in the leaves of

wheat have differential temperature responses which, combined with a predominance of

different circadian clock components in different cell types (*62*), causes bifurcated dynamics

of distinct circadian clock components in response to cold. Finally, an open question is of

whether the hexaploid wheat genome might support partial decoupling of subgenome-

derived circadian oscillators in single cells, although uneven temperature responses have

also been reported across the Arabidopsis circadian oscillator (*63*). In future, it will be

informative to know whether this delay in cycling in response to cold can act as a zeitgeber

for the wheat circadian clock, in a manner reminiscent of the phase shifts caused in

Arabidopsis by moderate temperature reductions (*64*). It will also be informative to

understand conserved and differing aspects of these responses in winter wheat, given its

different vernalization and photoperiod requirements compared with spring wheat varieties

such as Cadenza. Taken together, these results suggest that fluctuations in environmental

temperature might cause a dynamic adjustment of circadian phase in cultivated wheat (*65*).

Our data suggest two broad processes underlying the dynamics of circadian regulated and

cold-responsive transcripts in spring wheat. In the first case, low temperature delays the

circadian oscillator, maintaining the abundance of circadian-regulated gene targets. In the

second case, the circadian oscillator constrains or amplifies the influence of a cold treatment

on a gene, with that temporal input having a phase that is aligned with or different from the

underlying rhythm of expression of the transcript. This is well demonstrated by the cold

responsiveness of *TaBAM3*, for which the phase of the maximum responsiveness is not

defined by the phase of the transcript under control conditions (Fig. 2J-L). This suggests that

for some genes, separate circadian signalling components might regulate underlying

rhythms and cold responsiveness.

Potential cold induction bias towards D subgenome during subjective night

Genes that had circadian gated inductions by cold, with greatest cold responsiveness during

the subjective day, had a balanced contribution of transcripts from each subgenome. In

contrast, for transcripts with maximum cold induction phased to the subjective night, the

greatest proportion of transcript derived from the D subgenome (Fig. 4A). However, when

comparing the relative contributions from homoeologs that are present as triads, there did

not seem to be a bias towards the D subgenome (Fig. 4E). Therefore, the greater proportion

of transcripts from the D subgenome could arise from genes that are present as diads or

singletons. This might relate to the origins of the D subgenome donor, *Aegilops tauschii*. The

D subgenome hybridizations are thought to involve *A. tauschii* derived from mountainous

regions near the Caspian Sea (*66, 67*), so perhaps the greater cold induction from the D

subgenome during the subjective night relates to adaptation of *A. tauschii* to relatively cold

conditions.

Relationship between the circadian gating of the cold-responsive transcriptome and

circadian clock outputs

Rhythms in transcript abundance do not always lead to rhythms in protein abundance or

activity (*68-71*), although there is circadian gating of the high temperature-responsive

translatome (*56*). In many cases, the abundance of proteins encoded by rhythmic transcripts

remains constant (*71-73*), whilst in some cases (e.g. BAM3 in Arabidopsis) the protein

abundance tracks the transcript abundance (*73*). Circadian regulation of protein turnover

occurs in eukaryotes (*74, 75*), and it is thought that circadian regulation of gene expression

might maintain proteostasis in the presence of rhythmic protein degradation (*76*). This might

explain why rhythms in protein abundance are less prevalent than rhythms of transcript

abundance.

This idea could be relevant to interpreting the physiological significance of circadian gating in

wheat. For approximately half of the genes that have a circadian gated response to cold, the

phase of maximum cold sensitivity was similar to the phase of the rhythm at control

temperatures (delay ratio < 1; Fig. 6C). This likely increases the availability of mRNA at

certain times, perhaps in anticipation of a greater rate of protein turnover (*77*). This idea is

exemplified by the phasing to the subjective day of the cold induction of genes having light

signalling GO terms (Fig. S3). We reason that the regulatory network leads to increased

transcript levels for these components in anticipation of cold- and light-induced oxidative

damage. Conversely, transcripts that are down regulated and circadian gated could encode

proteins that are stabilised in cold or contribute to processes that are not preferred in cold

conditions. Alternatively, down regulated genes may represent repressive regulators like

many components of the circadian clock (*78*), so a reduction in their abundance could lift

repression of cold response mechanisms.

Conclusions

We identified that the circadian clock plays a key role in shaping the cold responsive

transcriptome through the process of circadian gating. This establishes that circadian

regulation is central to the environmental responses of a major cereal crop. Our study

supports the notion that circadian gating is a widespread regulatory mechanism found

across plant life, from peach trees and grape vines (*79, 80*) to Arabidopsis (*34*), and here in

bread wheat. Our findings open opportunities to understand the contributions of circadian

regulation to wheat crop performance, and could underpin biotechnological developments

that optimize the resilience of cereals to an increasingly unpredictable climate.

Materials and methods

Plant materials and growth conditions

Seeds of *Triticum aestivum* cv. Cadenza (donated by James Simmonds, John Innes Centre)

were stratified on damp filter paper for 3 days at 4^{o}C before germination at 22^{o}C in darkness

for two days. Seedlings were cultivated in 24-cell trays for 12 days on a bespoke cereal

cultivation mixture (65% peat, 25% loam, 10% horticultural grit, $3\,$kg m^{-3} dolomitic

limestone, $1.3\,$kg m^{-3} PG Mix fertiliser (Yara), $3\,$kg m^{-3} osmocote extract) under cycles of

12 h light and 12 h darkness, at approximately 200 µmol m^{-2} s^{-1} of white light at 22^{o}C.

Circadian time course sampling

14 day old wheat seedlings at Zadok stage GS1.2 were transferred to constant light for 24 h

before experimentation commenced, to ensure data were free from transitory effects caused

by the final dawn (Fig. S7A). After 25 h of constant conditions (ZT 25), three individuals were

transferred to 4^{o}C with identical light conditions for 3 h, whilst control plants remained at 22

^{o}C. After the 3 h cold treatment, 3 cm of tissue from the second leaf was collected from each

control and treated plant (Fig. S7A), immediately frozen in liquid N$_2$, and stored at -80^{o}C until

processing. Six such treatments were conducted, beginning at ZT 25, ZT 29, ZT 33, ZT 37,

ZT 41 and ZT 45, and ending at ZT 28, ZT 32, ZT 36, ZT 40, ZT 44 and ZT 48, respectively

(Fig. 1A). Each set of samples used a different batch of plants.

RNA sequencing data collection, QC and read mapping

RNA was extracted using Macherey-Nagel NucleoSpin RNA Mini Kit with on-column DNase

treatment, according to manufacturer's instructions. RNA was stored at -80^{o}C in nuclease-

free water. RNA integrity was assessed using an Agilent 2100 Bioanalyzer System at the

John Innes Centre, and again by our sequencing partner Novogene (Cambridge, UK). All

samples had RIN scores > 6 and concentrations > 50 ng RNA µl^{-1}, thereby passing quality

control. Libraries were constructed by Novogene using a mRNA polyA enrichment library

protocol involving fragmentation, cDNA synthesis, end repair, A-tailing, adapter ligation, size

selection, amplification and purification. Libraries were verified on a Bioanalyzer before

pooling for sequencing. After verification, 150bp paired-end sequencing was performed on

the Illumina NovaSeq 6000 platform to generate a minimum of 15 GB data per sample,

generating 675.5 GB total data with an average read depth of 62.5 million reads per sample.

All samples had $\geq$ 98.5 % clean reads suitable for alignment.

RNA sequencing analysis

Transcript quantification was conducted using the RNA-seq pseudoaligner kallisto v0.44.0

(Bray et al., 2016) using 31 bp k-mers and default parameters. The index file was

constructed from the high confidence CDS sequence from the International Wheat Genome

Sequencing Consortium (IWGSC) *T. aestivum* cv. Chinese Spring RefSeqv1.1 (*45*).

Differential gene expression analysis

Kallisto transcript count data were condensed to the gene level using R v2022.07.1 (*81*). The

Bioconductor (*82*) packages edgeR (*83-85*) and Limma (*86*) were used to identify

differentially expressed genes between cold-treated and control seedlings at each timepoint.

Counts per million (CPM) were determined by edgeR and low-expressed transcripts filtered

out of the data (edgeR function filterByExpr, which uses the filtering strategy described in

Chen *et al.* 2016) (Dataset S1, Dataset S2). This filtering strategy retains genes with a CPM

value above *k* in *n* samples, where *k* is determined from the set minimum count value and

sample size, and *n* determined from the smallest group size within the design matrix (*85*).

Around 68.5% of transcripts remained after filtering (Dataset S3). Empirical Bayes

moderation was performed for predictions of gene-wise variability (*87*) and a correction for

multiple testing (Benjamini-Hochberg method) was used to obtain the false discovery rate

(FDR) of differentially expressed genes. Differentially expressed genes were defined as

those with an adjusted p-value < 0.05 and logFC > 0.5 (50% change).

 Data processing and visualisation

CPM and logFC values were assembled using base R and the tidyverse package dplyr (*88,*

89). Graphs were produced using ggplot2 (*90*) and radar charts with fmsb function

radarchart (*91*). Upset plots (Fig. S1) were created with UpSetR (*92*) and GO-terms

presented using code adapted from (*39, 93*). The sensitivity heatmap was produced using

mean normalised CPM values determined by dividing the average CPM at any given

timepoint by the mean CPM for each gene across the entire time course under both

temperature conditions. The heatmap was created with ComplexHeatmap, using the Ward

D.2 method for clustering, with k = 12 (*94, 95*).

Identification of genes having a circadian gated response to cold

We used the difference between the mean control transcript level and mean cold-treated

transcript level (ΔCPM) at each timepoint as a measure of the sensitivity of gene expression

to cold. To test for rhythmicity of cold sensitivity, the ΔCPM profiles were analysed using the

MetaCycle v1.2.0 function meta2d (*96*). The minimum and maximum permitted period

lengths were 20 h and 28 h, respectively (*97*). We considered meta2d to be especially

suitable for this analysis because it integrates period, phase and p-values from three

algorithms, reducing the limitations of individual algorithms, and is appropriate for analysis of

a relatively small number of time-points (*97*). A rhythmicity cut-off of meta2d p-value < 0.05

was used. In general, time-series close to this p-value threshold had approximately 24 h

rhythmicity (Fig. S8). As an additional filter, only genes with rhythmic ΔCPM that were also

differentially expressed at least once in the time-series were considered to be circadian

gated. Only transcripts with average CPM > 1 under control temperature conditions or cold

temperature conditions, across the time-series, were included in the analysis to eliminate

low-expressed genes with large relative variation between replicates due to the low counts.

The difference between mean normalised CPM values (Δmean-normCPM) was used to

assign transcripts to three directional response groups: circadian-gated inductions by cold

(Δmean-normCPM > 0.5), cold induced delay in transcript cycling (-0.5 < Δmean-normCPM

< 0.5) and circadian gated reductions by cold (Δmean-normCPM > -0.5).

The estimated time of maximum cold responsiveness (ETMR) refers to the time of the peak

ΔCPM. This was estimated from the phase measured using meta2d analysis, which provides

the peak phase of the time-series data. For the ETMR for circadian gated cold reductions,

the ETMR was determined using meta2d phase estimates for the inverse ΔCPM values,

because they represent the phase of the trough. This allowed a total of 18 directional ETMR

groups to be defined (Fig. 3C). Extra filters were applied to ETMR groups < 4 h and > 20 h

for inductions and delayed transcripts, to remove 135 trend anomalies that had greater

ΔCPM at the opposite timepoint (+ 12 h) to the meta2d phase estimate.

Gene ontology-term analysis

GO-term enrichment analysis for biological processes was performed using TopGo (*98*) for

each cold-induction ETMR group. The group sizes were sufficiently large for this approach

(n>20). GO-terms were retrieved from the IWGSC RefSeqv1.0 annotation and transferred to

annotation v1.1 using the method described by (*99*). Enrichment analysis was performed on

gene lists through comparison against a gene universe of 75251 genes, which comprised all

genes expressed in our dataset that had high quality annotation and could be associated

with a GO term (nodeSize = 10, algorithm = weight01) (*98*). WeightedFisher values were

adjusted for multiple testing using the Benjamini-Hochberg correction method to generate

adjusted p-values, filtered stringently for adjusted p < 0.001. The same pipeline was used for

genes with CPM profiles that were temporally delayed by cold, but did not have rhythmic

ΔCPM (delay ratio <0.5).

Investigation of the contribution of wheat subgenomes to triad responsiveness

Subgenome occupancy was determined using information within the gene identifier (e.g.

TraesCS**1A**02G123456). Genes were organised into triads with a 1:1:1 correspondence

using the triad numbers in the High Confidence Triad table from (*46*)

(https://github.com/Uauy-Lab/

WheatHomoeologExpression/Data/data/TablesForExploration/HCTriads.csv). Triads were

considered to be part of a circadian gated induction ETMR group if any homoeolog from the

triad was categorised into the group. Individual homeologs of triads were sometimes within

different ETMR groups (Fig. 4D). An average ∆CPM value for the ETMR for each gene was

determined as:

$$\Delta CPM_{ETMR} = \frac{\Delta CPM_{ZTlower} + \Delta CPM_{ZTupper}}{2}$$

519   where ZTlower is the ZT timepoint corresponding to the lower time limit of the ETMR group

520   and ZTupper is the ZT timepoint corresponding to the upper time limit of the ETMR group,

521   e.g for ETMR group 4 – 8 h, ZTlower = ZT 28 and ZTupper = ZT 32. The relative contribution

522   of each homoeolog to the triad responsiveness for the ETMR was determined as:

$$Relative\ Contribution\ [x] = \frac{\Delta CPM_{ETMR}[x]}{\Delta CPM_{ETMR}[x] + \Delta CPM_{ETMR}[y] + \Delta CPM_{ETMR}[z]}$$

where x is the subgenome of the homoeolog being investigated, and y and z are the

remaining two subgenomes. This used a method similar to Ramirez-Gonzalez et al. (*46*).

The relative contributions were determined and visualised in R using ggtern (*100*).

Identifying temporal delays between cold-treated and control CPM profiles

We wished to identify the entire set of transcripts that had a delay in transcript cycling in

response to the cold treatments. To achieve this, transcripts with rhythmicity at control

temperatures and in response to 3 h cold treatments were identified using the ARSER

algorithm in Metacycle (using meta3d) (*96, 101*). Genes previously defined as having a

circadian gated upregulation or downregulation in response to cold were removed from this

group to prevent double-counting of individual genes. Adjusted ∆CPM (Adj.∆CPM) values

were determined for timepoints ZT28-44 for each gene as:

$$Adj. \Delta CPM_{ZTx} = Cold\left[CPM_{ZTx+4}\right] - Control\left[CPM_{ZTx}\right]$$

534 Where ZTx is the is sampling timepoint. A time course mean ΔCPM for each gene was

535 determined as:

$$Mean. \Delta CPM = \frac{\left|\Delta CPM_{ZT28}\right| + \left|\Delta CPM_{ZT32}\right| + \left|\Delta CPM_{ZT36}\right| + \left|\Delta CPM_{ZT40}\right| + \left|\Delta CPM_{ZT44}\right|}{5}$$

536 And a mean for adjusted ΔCPM for each gene was determined as:

$$Mean. Adj. \Delta CPM = \frac{\begin{array}{c}\left|Adj. \Delta CPM_{ZT28}\right| + \left|Adj. \Delta CPM_{ZT32}\right| + \left|Adj. \Delta CPM_{ZT36}\right| + \\ \left|Adj. \Delta CPM_{ZT40}\right| + \left|Adj. \Delta CPM_{ZT44}\right|\end{array}}{5}$$

537 A delay ratio value was determined for each gene in this group using mean ΔCPM and mean

538 adjusted ΔCPM values as:

$$Delay\ ratio = \frac{Mean. Adj. \Delta CPM}{Mean. \Delta CPM}$$

539 Delay ratios less than 1 indicate a reduced mean ΔCPM following adjustment. Gene

540 identifiers for putative wheat circadian clock genes were obtained from (*39*), and delay ratios

541 determined for each.

Acknowledgements

We thank James Simmonds for seeds of *T. aestivum* cv. Cadenza, Paige Panter for experimental design advice, Cristobal Uauy for advice on data interpretation, and Hannah Rees and Rachel Rusholme-Pilcher for advice on data processing.

Funding

BBSRC Institute Strategic Programme GEN BB/P013511/1 (AND, PP).

The Leverhulme Trust RPG-2018-216 (AND).

SWBio DTP grants BB/J014400/1 and BB/M009122/1 (CAG, KJE, AND).

Author contributions

Conceptualization: CAG, KJE, AND.

Investigation: CAG, PP.

Analysis and visualization: CAG, PP.

Writing: CAG, PP, KJE, AND.

Competing interests

Authors declare that they have no competing interests.

Data availability

Raw sequencing data are deposited as European Nucleotide Archive dataset PRJEB56923. Processed data are provided as supplemental information within the manuscript.

References

1.   S. L. Harmer, J. B. Hogenesch, M. Straume, H.-S. Chang, B. Han, T. Zhu, X. Wang,
J. A. Kreps, S. A. Kay, Orchestrated transcription of key pathways in *Arabidopsis* by
the circadian clock. *Science* **290**, 2110-2113 (2000).
2.   J. Aschoff, Temporal orientation: circadian clocks in animals and humans. *Animal
Behaviour* **37**, 881-896 (1989).
3.   A. J. Millar, The intracellular dynamics of circadian clocks reach for the light of
ecology and evolution. *Annual Review of Plant Biology* **67**, 595-618 (2016).
4.   M. F. Covington, J. N. Maloof, M. Straume, S. A. Kay, Global transcriptome analysis
reveals circadian regulation of key pathways in plant growth and development.
Genome Biology **9**, R130 (2008).
5.   P. Y. Hsu, U. K. Devisetty, S. L. Harmer, Accurate timekeeping is controlled by a
cycling activator in *Arabidopsis. eLife* **2**, e00473 (2013).
6.   M. Kamioka, S. Takao, T. Suzuki, K. Taki, T. Higashiyama, T. Kinoshita, N.
Nakamichi, Direct repression of evening genes by CIRCADIAN CLOCK-
ASSOCIATED1 in the Arabidopsis circadian clock. *The Plant Cell* **28**, 696-711
(2016).
7.   N. Nakamichi, T. Kiba, M. Kamioka, T. Suzuki, T. Yamashino, T. Higashiyama, H.
Sakakibara, T. Mizuno, Transcriptional repressor PRR5 directly regulates clock-
output pathways. *Proceedings of the National Academy of Sciences* **109**, 17123-
17128 (2012).
8.   J. M. Gendron, J. L. Pruneda-Paz, C. J. Doherty, A. M. Gross, S. E. Kang, S. A. Kay,
in *Proceedings of the National Academy of Sciences of the United States of America*.
(2012), vol. 109, pp. 3167-3172.
9.   N. Nakamichi, Adaptation to the local environment by modifications of the
photoperiod response in crops. *Plant and Cell Physiology* **56**, 594-604 (2015).
10.   C. R. McClung, Circadian clock components offer targets for crop domestication and
improvement. *Genes* **12**, 374 (2021).
11.   M. W. Oravec, K. Greenham, The adaptive nature of the plant circadian clock in
natural environments. *Plant Physiology* **190**, 968-980 (2022).
12.   G. Steed, D. C. Ramirez, M. A. Hannah, A. A. R. Webb, Chronoculture, harnessing
the circadian clock to improve crop yield and sustainability. *Science* **372**, 479 (2021).
13.   L. Gu, P. J. Hanson, W. M. Post, D. P. Kaiser, B. Yang, R. Nemani, S. G. Pallardy, T.
Meyers, The 2007 eastern US spring freeze: Increased cold damage in a warming
world? *BioScience* **58**, 253-262 (2008).
14.   M. Trnka, R. P. Rötter, M. Ruiz-Ramos, K. C. Kersebaum, J. E. Olesen, Z. Žalud, M.
A. Semenov, Adverse weather conditions for European wheat production will become
more frequent with climate change. *Nature Climate Change* **4**, 637-643 (2014).
15.   K. M. Barlow, B. P. Christy, G. J. O'Leary, P. A. Riffkin, J. G. Nuttall, Simulating the
impact of extreme heat and frost events on wheat crop production: A review. *Field
Crops Research* **171**, 109-119 (2015).
16.   J. M. Whaley, E. J. M. Kirby, J. H. Spink, M. J. Foulkes, D. L. Sparkes, Frost damage
to winter wheat in the UK: the effect of plant population density. *European Journal of
Agronomy* **21**, 105-115 (2004).
17.   P. Thakur, S. Kumar, J. A. Malik, J. D. Berger, H. Nayyar, Cold stress effects on
reproductive development in grain crops: An overview. *Environmental and
Experimental Botany* **67**, 429-443 (2010).
18.   X. Li, H. Pu, F. Liu, Q. Zhou, J. Cai, T. Dai, W. Cao, D. Jiang, Winter wheat
photosynthesis and grain yield responses to spring freeze. *Agronomy Journal* **107**,
1002-1010 (2015).
19.   E. D. Leonardos, L. V. Savitch, N. P. A. Huner, G. Öquist, B. Grodzinski, Daily
photosynthetic and C-export patterns in winter wheat leaves during cold stress and
acclimation. *Physiologia Plantarum* **117**, 521-531 (2003).

20. Y. W. Jame, H. W. Cutforth, Simulating the effects of temperature and seeding depth
on germination and emergence of spring wheat. *Agricultural and Forest Meteorology*
124, 207-218 (2004).
21. V. M. Hurry, N. P. A. Huner, Effect of cold hardening on sensitivity of winter and
spring wheat leaves to short-term photoinhibition and recovery of photosynthesis.
Plant Physiology **100**, 1283-1290 (1992).
22. K. D. Subedi, P. J. Gregory, R. J. Summerfield, M. J. Gooding, Cold temperatures
and boron deficiency caused grain set failure in spring wheat (*Triticum aestivum* L.).
Field Crops Research **57**, 277-288 (1998).
23. Y. V. Venzhik, A. F. Titov, V. V. Talanova, S. A. Frolova, A. V. Talanov, Y. A.
Nazarkina, Influence of lowered temperature on the resistance and functional activity
of the photosynthetic apparatus of wheat plants. *Biology Bulletin* **38**, 132-137 (2011).
24. M. O. Winfield, C. Lu, I. D. Wilson, J. A. Coghill, K. J. Edwards, Plant responses to
cold: transcriptome analysis of wheat. *Plant Biotechnology Journal* **8**, 749-771
(2010).
25. M. P. Fuller, A. M. Fuller, S. Kaniouras, J. Christophers, T. Fredericks, The freezing
characteristics of wheat at ear emergence. *European Journal of Agronomy* **26**, 435-
441 (2007).
26. C. T. Hotta, M. J. Gardner, K. E. Hubbard, S. J. I. N. Baek, N. Dalchau, D. Suhita, A.
N. Dodd, A. A. R. Webb, Modulation of environmental responses of plants by
circadian clocks. *Plant, Cell & Environment* **30**, 333-349 (2007).
27. S. D. Skopik, C. S. Pittendrigh, Circadian systems, II. The oscillation in the individual
Drosophila pupa; its independence of developmental stage. *Proceedings of the*
National Academy of Sciences **58**, 1862-1869 (1967).
28. S. G. Fowler, D. Cook, M. F. Thomashow, Low temperature induction of Arabidopsis
CBF1, 2, and 3 is gated by the circadian clock. *Plant Physiology* **137**, 961-968
(2005).
29. A. N. Dodd, M. K. Jakobsen, A. J. Baker, A. Telzerow, S.-W. Hou, L. Laplaze, L.
Barrot, R. S. Poethig, J. Haseloff, A. A. R. Webb, Time of day modulates low-
temperature Ca signals in Arabidopsis. *The Plant Journal* **48**, 962-973 (2006).
30. Z. Bieniawska, C. Espinoza, A. Schlereth, R. Sulpice, D. K. Hincha, M. A. Hannah,
Disruption of the Arabidopsis circadian clock is responsible for extensive variation in
the cold-responsive transcriptome. *Plant Physiology* **147**, 263-279 (2008).
31. J.-Y. Zhu, E. Oh, T. Wang, Z.-Y. Wang, TOC1–PIF4 interaction mediates the
circadian gating of thermoresponsive growth in Arabidopsis. *Nature Communications*
7, 13692 (2016).
32. E. J. Blair, T. Bonnot, M. Hummel, E. Hay, J. M. Marzolino, I. A. Quijada, D. H.
Nagel, Contribution of time of day and the circadian clock to the heat stress
responsive transcriptome in Arabidopsis. *Scientific Reports* **9**, 4814 (2019).
33. D. O. Grinevich, J. S. Desai, K. P. Stroup, J. Duan, E. Slabaugh, C. J. Doherty, Novel
transcriptional responses to heat revealed by turning up the heat at night. *Plant*
Molecular Biology **101**, 1-19 (2019).
34. T. Bonnot, E. J. Blair, S. J. Cordingley, D. H. Nagel, Circadian coordination of cellular
processes and abiotic stress responses. *Current Opinion in Plant Biology* **64**, 102133
(2021).
35. S. Kidokoro, K. Hayashi, H. Haraguchi, T. Ishikawa, F. Soma, I. Konoura, S. Toda, J.
Mizoi, T. Suzuki, K. Shinozaki, K. Yamaguchi-Shinozaki, Posttranslational regulation
of multiple clock-related transcription factors triggers cold-inducible gene expression
in Arabidopsis. *Proceedings of the National Academy of Sciences* **118**, e2021048118
(2021).
36. B. Li, Z. Gao, X. Liu, D. Sun, W. Tang, Transcriptional profiling reveals a time-of-day-
specific role of REVEILLE 4/8 in regulating the first wave of heat shock–induced
gene expression in Arabidopsis. *The Plant Cell* **31**, 2353-2369 (2019).

37. A. J. Nagano, Y. Sato, M. Mihara, B. A. Antonio, R. Motoyama, H. Itoh, Y. Nagamura,
T. Izawa, Deciphering and prediction of transcriptome dynamics under fluctuating
field conditions. *Cell* **151**, 1358-1369 (2012).
38. D. L. Cano-Ramirez, H. Nishio, T. Muranaka, L. i. L. de Barros Dantas, M. N. Honjo,
J. Sugisaka, H. Kudoh, A. N. Dodd, Circadian and environmental signal transduction
in a natural population of Arabidopsis. *bioRxiv*, (2022).
39. H. Rees, R. Rusholme-Pilcher, P. Bailey, J. Colmer, B. White, C. Reynolds, S. J.
Ward, B. Coombes, C. A. Graham, L. L. de Barros Dantas, A. N. Dodd, A. Hall,
Circadian regulation of the transcriptome in a complex polyploid crop. *PLOS Biology*
20, 1-37 (2022).
40. J. Beales, A. Turner, S. Griffiths, J. W. Snape, D. A. Laurie, A *Pseudo-Response*
Regulator is misexpressed in the photoperiod insensitive *Ppd-D1a* mutant of wheat
(*Triticum aestivum* L.). *Theoretical and Applied Genetics* **115**, 721-733 (2007).
41. A. R. Bentley, R. Horsnell, C. P. Werner, A. S. Turner, G. A. Rose, C. Bedard, P.
Howell, E. P. Wilhelm, I. J. Mackay, R. M. Howells, A. Greenland, D. A. Laurie, N.
Gosman, Short, natural, and extended photoperiod response in BC2F4 lines of bread
wheat with different *Photoperiod-1* (*Ppd-1*) alleles. *Journal of Experimental Botany*
64, 1783-1793 (2013).
42. L. M. Shaw, A. S. Turner, D. A. Laurie, The impact of photoperiod insensitive *Ppd-1a*
mutations on the photoperiod pathway across the three genomes of hexaploid wheat
(*Triticum aestivum*). *The Plant Journal* **71**, 71-84 (2012).
43. A. Turner, J. Beales, S. Faure, R. P. Dunford, D. A. Laurie, The pseudo-response
regulator *Ppd-H1* provides adaptation to photoperiod in Barley. *Science* **310**, 1031-
1034 (2005).
44. K. Greenham, R. C. Sartor, S. Zorich, P. Lou, T. C. Mockler, C. R. McClung,
Expansion of the circadian transcriptome in *Brassica rapa* and genome-wide
diversification of paralog expression patterns. *eLife* **9**, e58993 (2020).
45. IWGSC, Shifting the limits in wheat research and breeding using a fully annotated
reference genome. *Science* **361**, eaar7191 (2018).
46. R. H. Ramírez-González, P. Borrill, D. Lang, S. A. Harrington, J. Brinton, L. Venturini,
M. Davey, J. Jacobs, F. van Ex, A. Pasha, Y. Khedikar, S. J. Robinson, A. T. Cory, T.
Florio, L. Concia, C. Juery, H. Schoonbeek, B. Steuernagel, D. Xiang, C. J. Ridout,
B. Chalhoub, K. F. X. Mayer, M. Benhamed, D. Latrasse, A. Bendahmane, B. B. H.
Wulff, R. Appels, V. Tiwari, R. Datla, F. Choulet, C. J. Pozniak, N. J. Provart, A. G.
Sharpe, E. Paux, M. Spannagl, A. Bräutigam, C. Uauy, The transcriptional landscape
of polyploid wheat. *Science* **361**, 6089 (2018).
47. M. D. Mikkelsen, M. F. Thomashow, A role for circadian evening elements in cold-
regulated gene expression in Arabidopsis. *The Plant Journal* **60**, 328-339 (2009).
48. M. A. Dong, E. M. Farré, M. F. Thomashow, CIRCADIAN CLOCK-ASSOCIATED 1
and LATE ELONGATED HYPOCOTYL regulate expression of the C-REPEAT
BINDING FACTOR (CBF) pathway in *Arabidopsis*. *Proceedings of the National*
Academy of Sciences **108**, 7241-7246 (2011).
49. D. B. Fowler, A. E. Limin, Interactions among factors regulating phenological
development and acclimation rate determine low-temperature tolerance in wheat.
Annals of botany **94**, 717-724 (2004).
50. Q. Li, B. Byrns, M. A. Badawi, A. B. Diallo, J. Danyluk, F. Sarhan, D. Laudencia-
Chingcuanco, J. Zou, D. B. Fowler, Transcriptomic insights into phenological
development and cold tolerance of wheat grown in the field. *Plant Physiology* **176**,
2376-2394 (2017).
51. M. F. Thomashow, Molecular basis of plant cold acclimation: insights gained from
studying the CBF cold response pathway. *Plant Physiology* **154**, 571-577 (2010).
52. M. F. Thomashow, S. J. Gilmour, E. J. Stockinger, K. R. Jaglo-Ottosen, D. G. Zarka,
Role of the Arabidopsis CBF transcriptional activators in cold acclimation.
Physiologia Plantarum **112**, 171-175 (2001).

53. M. Badawi, J. Danyluk, B. Boucho, M. Houde, F. Sarhan, The CBF gene family in
hexaploid wheat and its relationship to the phylogenetic complexity of cereal CBFs.
Molecular Genetics and Genomics **277**, 533-554 (2007).
54. K. Gierczik, A. Novák, M. Ahres, A. Székely, A. Soltész, Á. Boldizsár, Z. Gulyás, B.
Kalapos, I. Monostori, L. Kozma-Bognár, G. Galiba, A. Vágújfalvi, Circadian and light
regulated expression of CBFs and their upstream signalling genes in barley.
International Journal of Molecular Sciences **18**, (2017).
55. D. Xu, COP1 and BBXs-HY5-mediated light signal transduction in plants. *New
Phytologist* **228**, 1748-1753 (2020).
56. T. Bonnot, D. H. Nagel, Time of the day prioritizes the pool of translating mRNAs in
response to heat stress. *The Plant Cell* **33**, 2164-2182 (2021).
57. A. Graf, A. Schlereth, M. Stitt, A. M. Smith, Circadian control of carbohydrate
availability for growth in Arabidopsis plants at night. *Proceedings of the National
Academy of Sciences* **107**, 9458-9463 (2010).
58. S. M. Smith, D. C. Fulton, T. Chia, D. Thorneycroft, A. Chapple, H. Dunstan, C.
Hylton, S. C. Zeeman, A. M. Smith, Diurnal changes in the transcriptome encoding
enzymes of starch metabolism provide evidence for both transcriptional and
posttranscriptional regulation of starch metabolism in *Arabidopsis* leaves. *Plant
Physiology* **136**, 2687-2699 (2004).
59. A. N. Dodd, R. Toth, E. Kevei, A. A. R. Webb, N. Salathia, J. M. Hibberd, F. Nagy, A.
J. Millar, A. Hall, Plant circadian clocks increase photosynthesis, growth, survival,
and competitive advantage. *Science* **309**, 630-633 (2005).
60. K. K. Markham, K. Greenham, Abiotic stress through time. *New Phytologist* **231**, 40-
46 (2021).
61. P. J. Dickinson, M. Kumar, C. Martinho, S. J. Yoo, H. Lan, G. Artavanis, V.
Charoensawan, M. A. Schöttler, R. Bock, K. E. Jaeger, P. A. Wigge, Chloroplast
signaling gates thermotolerance in Arabidopsis. *Cell Reports* **22**, 1657-1665 (2018).
62. M. Endo, H. Shimizu, M. A. Nohales, T. Araki, S. A. Kay, Tissue-specific clocks in
Arabidopsis show asymmetric coupling. *Nature* **515**, 419-422 (2014).
63. T. P. Michael, P. A. Salomé, C. R. McClung, Two *Arabidopsis* circadian oscillators
can be distinguished by differential temperature sensitivity. *Proceedings of the
National Academy of Sciences* **100**, 6878-6883 (2003).
64. P. A. Salomé, C. R. McClung, What makes the Arabidopsis clock tick on time? A
review on entrainment. *Plant, Cell & Environment* **28**, 21-38 (2005).
65. A. A. R. Webb, M. Seki, A. Satake, C. Caldana, Continuous dynamic adjustment of
the plant circadian oscillator. *Nature Communications* **10**, 550 (2019).
66. K. Gaurav, S. Arora, P. Silva, J. Sánchez-Martín, R. Horsnell, L. Gao, G. S. Brar, V.
Widrig, W. John Raupp, N. Singh, S. Wu, S. M. Kale, C. Chinoy, P. Nicholson, J.
Quiroz-Chávez, J. Simmonds, S. Hayta, M. A. Smedley, W. Harwood, S. Pearce, D.
Gilbert, N. Kangara, C. Gardener, M. Forner-Martínez, J. Liu, G. Yu, S. A. Boden, A.
Pascucci, S. Ghosh, A. N. Hafeez, T. O'Hara, J. Waites, J. Cheema, B. Steuernagel,
M. Patpour, A. F. Justesen, S. Liu, J. C. Rudd, R. Avni, A. Sharon, B. Steiner, R. P.
Kirana, H. Buerstmayr, A. A. Mehrabi, F. Y. Nasyrova, N. Chayut, O. Matny, B. J.
Steffenson, N. Sandhu, P. Chhuneja, E. Lagudah, A. F. Elkot, S. Tyrrell, X. Bian, R.
P. Davey, M. Simonsen, L. Schauser, V. K. Tiwari, H. Randy Kutcher, P. Hucl, A. Li,
D.-C. Liu, L. Mao, S. Xu, G. Brown-Guedira, J. Faris, J. Dvorak, M.-C. Luo, K.
Krasileva, T. Lux, S. Artmeier, K. F. X. Mayer, C. Uauy, M. Mascher, A. R. Bentley,
B. Keller, J. Poland, B. B. H. Wulff, Population genomic analysis of *Aegilops tauschii*
identifies targets for bread wheat improvement. *Nature Biotechnology* **40**, 422-431
(2022).
67. Y. Zhou, X. Zhao, Y. Li, J. Xu, A. Bi, L. Kang, D. Xu, H. Chen, Y. Wang, Y.-G. Wang,
S. Liu, C. Jiao, H. Lu, J. Wang, C. Yin, Y. Jiao, F. Lu, Triticum population sequencing
provides insights into wheat adaptation. *Nature Genetics* **52**, 1412-1422 (2020).
68. K. Baerenfaller, C. Massonnet, S. Walsh, S. Baginsky, P. Bühlmann, L. Hennig, M.
Hirsch-Hoffmann, K. A. Howell, S. Kahlau, A. Radziejwoski, D. Russenberger, D.

Rutishauser, I. Small, D. Stekhoven, R. Sulpice, J. Svozil, N. Wuyts, M. Stitt, P.
Hilson, C. Granier, W. Gruissem, Systems-based analysis of Arabidopsis leaf growth
reveals adaptation to water deficit. *Molecular Systems Biology* **8**, 606 (2012).

69. M. K. Choudhary, Y. Nomura, L. Wang, H. Nakagami, D. E. Somers, Quantitative
circadian phosphoproteomic analysis of arabidopsis reveals extensive clock control
of key components in physiological, metabolic, and signaling pathways. *Molecular &*
Cellular Proteomics **14**, 2243-2260 (2015).

70. A. Graf, D. Coman, R. G. Uhrig, S. Walsh, A. Flis, M. Stitt, W. Gruissem, Parallel
analysis of Arabidopsis circadian clock mutants reveals different scales of
transcriptome and proteome regulation. *Open Biology* **7**, 160333 (2017).

71. D. D. Seaton, A. Graf, K. Baerenfaller, M. Stitt, A. J. Millar, W. Gruissem,
Photoperiodic control of the Arabidopsis proteome reveals a translational coincidence
mechanism. *Molecular Systems Biology* **14**, e7962 (2018).

72. R. G. Uhrig, S. Echevarría-Zomeño, P. Schlapfer, J. Grossmann, B. Roschitzki, N.
Koerber, F. Fiorani, W. Gruissem, Diurnal dynamics of the Arabidopsis rosette
proteome and phosphoproteome. *Plant, Cell & Environment* **44**, 821-841 (2021).

73. J. Krahmer, M. Hindle, L. K. Perby, H. K. Mogensen, T. H. Nielsen, K. J. Halliday, G.
van Ooijen, T. Le Bihan, A. J. Millar, The circadian clock gene circuit controls protein
and phosphoprotein rhythms in *Arabidopsis thaliana*. *Molecular & Cellular*
Proteomics **21**, (2022).

74. L. Li, C. J. Nelson, J. Trösch, I. Castleden, S. Huang, A. H. Millar, Protein
degradation rate in *Arabidopsis thaliana* leaf growth and development. *The Plant Cell*
29, 207-228 (2017).

75. S. B. Srikanta, N. Cermakian, To Ub or not to Ub: Regulation of circadian clocks by
ubiquitination and deubiquitination. *Journal of Neurochemistry* **157**, 11-30 (2021).

76. E. Seinkmane, S. Y. Peak-Chew, A. Zeng, N. M. Rzechorzek, A. D. Beale, D. C. S.
Wong, J. S. O'Neill, Circadian regulation of protein turnover and proteome renewal.
bioRxiv, (2022).

77. N. Murata, S. Takahashi, Y. Nishiyama, S. I. Allakhverdiev, Photoinhibition of
photosystem II under environmental stress. *Biochimica et Biophysica Acta* **1767**,
414-421 (2007).

78. K. Fogelmark, C. Troein, Rethinking transcriptional activation in the arabidopsis
circadian clock. *PLOS Computational Biology* **10**, 1-12 (2014).

79. T. S. Artlip, M. E. Wisniewski, C. L. Bassett, J. L. Norelli, CBF gene expression in
peach leaf and bark tissues is gated by a circadian clock. *Tree Physiology* **33**, 866-
877 (2013).

80. M. Rienth, L. Torregrosa, N. Luchaire, R. Chatbanyong, D. Lecourieux, M. T. Kelly,
C. Romieu, Day and night heat stress trigger different transcriptomic responses in
green and ripening grapevine (vitis vinifera) fruit. *BMC Plant Biology* **14**, 108 (2014).

81. R. C. Team, *R: A language and environment for statistical computing* (R Foundation
for Statistical Computing, Vienna, Austria, 2018).

82. W. Huber, V. J. Carey, R. Gentleman, S. Anders, M. Carlson, B. S. Carvalho, H. C.
Bravo, S. Davis, L. Gatto, T. Girke, R. Gottardo, F. Hahne, K. D. Hansen, R. A.
Irizarry, M. Lawrence, M. I. Love, J. MacDonald, V. Obenchain, A. K. Oleś, H. Pagès,
A. Reyes, P. Shannon, G. K. Smyth, D. Tenenbaum, L. Waldron, M. Morgan,
Orchestrating high-throughput genomic analysis with Bioconductor. *Nature Methods*
12, 115-121 (2015).

83. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
139-140 (2010).

84. D. J. McCarthy, Y. Chen, G. K. Smyth, Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research*
40, 4288-4297 (2012).

85. Y. Chen, A. A. T. Lun, G. K. Smyth, From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. *F1000Research* **5**, 1438 (2016).

86. M. E. Ritchie, B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi, G. K. Smyth, limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research* **43**, e47 (2015).

87. G. K. Smyth, Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Statistical Applications in Genetics and Molecular Biology* **3**, Article 3 (2004).

88. H. Wickham, M. Averick, J. Bryan, W. Chang, L. D. A. McGowan, R. François, G. Grolemund, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. L. Pedersen, E. Miller, S. M. Bache, K. Müller, J. Ooms, D. Robinson, D. P. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K. Woo, H. Yutani, Welcome to the tidyverse. *Journal of Open Source Software* **4**, 1686 (2019).

89. H. Wickham, R. François, L. Henry, K. Müller, dplyr: A grammar of data manipulation. (2022).

90. H. Wickham, ggplot2: Elegant graphics for data analysis. (2016).

91. M. Nakazawa, fmsb: functions for medical statistics book with some demographic data. (2022).

92. N. Gehlenborg, UpSetR: a more scalable alternative to Venn and Euler Diagrams for visualizing intersecting sets. (2019).

93. J. J. De Vega, N. Peel, S. J. Purdy, S. Hawkins, L. Donnison, S. Dyer, K. Farrar, Differential expression of starch and sucrose metabolic genes linked to varying biomass yield in Miscanthus hybrids. *Biotechnology for Biofuels* **14**, 98 (2021).

94. Z. Gu, Complex heatmap visualization. *iMeta*, (2022).

95. Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics*, (2016).

96. G. Wu, R. C. Anafi, M. E. Hughes, K. Kornacker, J. B. Hogenesch, MetaCycle: an integrated R package to evaluate periodicity in large scale data. *Bioinformatics* **32**, 3351-3353 (2016).

97. G. Wu, R. Anafi, J. Hogenesch, M. Hughes, K. Kornacker, X. Li, M. Carlucci, MetaCycle: Evaluate periodicity in large scale data. (2019).

98. A. Alexa, J. Rahnenfuhrer, topGO: Enrichment Analysis for Gene Ontology. (2022).

99. P. Borrill, S. A. Harrington, J. Simmonds, C. Uauy, Identification of transcription factors regulating senescence in wheat through gene regulatory network modelling. *Plant Physiology* **180**, 1740-1755 (2019).

100. N. E. Hamilton, M. Ferry, ggtern: Ternary Diagrams Using ggplot2. *Journal of Statistical Software, Code Snippets* **87**, 1-17 (2018).

101. R. Yang, Z. Su, Analyzing circadian expression data by harmonic regression based on autoregressive spectral estimation. *Bioinformatics* **26**, i168-i174 (2010).

 Figure legends

 Figure 1. Different sets of transcripts respond to cold at different times of day in bread

wheat. (A) *Triticum aestivum* cv. Cadenza seedlings were entrained under 12:12 light/dark

cycles before transfer to free-running conditions. Under free running conditions, tissue was

collected after 3 h cold treatments, with six such treatments given over 24 h of free running

conditions. **(B)** Differentially expressed genes after a 3 h cold treatments given at six

timepoints (log fold change > 0.5, FDR < 0.05). Yellow and grey shading indicate subjective

day and night, respectively. **(C)** The number of differentially expressed genes (DEGs) at

each timepoint. **(D)** The proportion of DEGs that only responded to cold at single timepoints.

(E) Summary of the difference between mean normalised control and cold transcript

abundance of DEGs across the time course. Orange and teal colours indicate normalized

change in cold-induced and cold-repressed transcripts, respectively.

Figure 2. Circadian gating of the cold response of selected wheat triads. (A-C)

TaCBFVc-B14; triad no. 15598. **(D-F)** *TaBBX17*; triad no 9634. **(G-I)** *TaBAM1*; triad no

15540. **(J-L)** *TaBAM3*; triad no 15539 (*46*). Yellow/grey shading = subjective day/night. Solid

lines are mean (N = 3 biological replicates; blue/green shading = ± s.e.m). Asterisks indicate

times when the transcript was differentially expressed between the control and cold

temperature conditions (LogFC >0.5, FDR < 0.05). CPM (counts per million) determined by

edgeR (*83*).

Figure 3. The temporal structure of the cold-responsive transcriptome. (A) We defined

circadian gating of the response of genes to cold as a 24 h cycle of sensitivity of the

response of the gene to identical cold treatments. To identify such genes, we calculated the

difference between the mean control and cold-treated CPM (termed ΔCPM). Circadian

gating of a cold response manifests as a 24 h oscillation of ΔCPM, which captures cold

induction, repression, or a delay in transcript cycling. **(B)** The number of significantly cold-

responsive transcripts (LogFC >0.5, FDR <0.05) that have a rhythmic ΔCPM profile across

the time course (meta2d, p <0.05). This represents a set of wheat transcripts with a

circadian gated response to cold. **(C)** Transcripts with circadian gating of their response to

cold, organized according to their estimated time of maximum peak cold responsiveness

(ETMR). Direction of response (induction, reduction, delay) was based on the average

difference between mean normalised control and cold-treated CPM to normalize differences

in ΔCPM profile amplitude across the transcriptome. The direction was formalized as

circadian-gated inductions (Δmean.normCPM > 0.5), cold induced delay (-0.5 <

Δmean.normCPM < 0.5) and circadian gated reductions (Δmean.normCPM > -0.5). ETMR

was derived from meta2d phase estimates. Yellow/grey shading = subjective day/night. Pink

lines indicate mean Δmean.normCPM in each directional ETMR group. **(D)** The number of

genes within each directional cold ETMR group. 1533 genes were included, after filtering to

remove ΔCPM profiles from ETMR groups < 4 h and > 20 h having peak CPM at an

opposing timepoint from the ETMR. Darker colours indicate greater numbers of transcripts.

Figure 4. Subgenome organization of the circadian gating of the cold-responsive

transcriptome. (A) The number and proportion of transcripts from each subgenome that

have a circadian gated response to cold, divided according to the time of maximum cold

responsiveness. Large proportional differences between subgenomes generally occur when

total number of transcripts is small. (**B**) The proportion of transcripts that have a circadian

gated response to cold that belong to an annotated triad (*46*). Not all homeologs within any

given triad are gated, and this is indicated by the bar colour. (**C**) Example ternary plot

showing the relative contribution of each homoeolog within a triad to the combined transcript

accumulation from the triad (*46, 100*). The position of each point on the plot reflects the

proportion of total contribution from the three homeologs within the triad (subgenome A,

yellow axis; subgenome B; green axis, subgenome D, blue axis). A point in the centre of the

plot represents a triad where all three homoeologs are equally cold-responsive. Points

towards corners (shaded areas) represent triads where the majority of the cold response

derives from a specific homoeolog. (**D)** The relative contributions of homoeologs to the cold

responsiveness of each triad, organized according to the time of maximum cold

responsiveness. Each triad contains at least one homoeolog that is within the ETMR group.

Therefore, certain triads are present in only one ETMR (orange points), some in two (dark

pink) and some in three (light).

Figure 5. Cold temperature delays the oscillation of a set of rhythmic transcripts. (A)

Dynamics of a delayed transcript (*TraesCS4A02G091700*), whereby the oscillation in cold

treated plants (blue) is similar to control temperature conditions (green), but delayed in time.

The difference between cold and control treated transcript abundance (ΔCPM) is shown in

dark red. **(B)** Dynamics of a rhythmic transcript (*TraesCS7A02G416300*; meta3d: ARS p

<0.05) that has a delayed oscillation and a more linear ΔCPM profile. **(C)** Artificial data

including a time delay. ΔCPM is the difference between the control and cold CPM at each

timepoint (dark red arrows), and adjusted ΔCPM is the difference between the control

temperature CPM at one timepoint and the cold-treated CPM at the next timepoint (blue

arrows). **(D)** a delay ratio value was calculated from the average adjusted ΔCPM as a

proportion of the original average ΔCPM for transcripts delayed by cold, and **(E)** all

transcripts with rhythmic control and cold-treated CPM profiles (meta3d: ARS p <0.05). In

(D, E) The larger circle represents 100% of these transcripts, the middle circle represents

transcripts with a delay ratio <1, and the smaller circle transcripts with a delay ratio < 0.5.

Delay ratios < 1 suggest a transcript delay. **(F)** Delay ratios of eleven circadian clock gene

triads (Rees *et al.*, 2022) suggest cold-induced delayed cycling for the majority. (**G**)

Transcript dynamics of delayed circadian clock gene *TaGI 3A* (*TraesCS3A02G116300*) and

no-delay *TaLUX/BOA 3A* (*TraesCS3A02G526600*). Solid lines are mean (N = 3). Blue/green

shading= ± s.e.m. Red line represents ΔCPM. Yellow/grey shading = subjective day/night.

Figure 6. Features of the circadian gating of cold temperature responses. The

interaction between the circadian clock and cold produces a diversity of transcript response

profiles across the wheat transcriptome. Diagrams conceptualize how the circadian clock

(red clocks) and cold (blue arrows) can affect underlying control-temperature transcript

levels (dotted black line), producing a diversity of cold-responsive transcript profiles (orange line). **(A)** certain transcripts can be regulated by the circadian clock but transcript levels are unaltered by cold (e.g. *TraesCS5D02G447500* and *TraesCS1D02G372000*). **(B)** transcripts can respond equally to cold, independent of the time of subjective day (e.g. *TraesCS2D02G284400* and *TraesCS3B02G287400*). **(C)** the cold-responsiveness of transcripts can be gated by the circadian clock, resulting in a response to cold at specific times of day (e.g. *TraesCS3D02G325300). **(D)** the circadian clock gates the response to cold of the transcript, with the gate having a different phase from the phase of the underlying circadian rhythm of the transcript (e.g. *TaBAM3*). **(E)** cold affects the circadian oscillator directly (cold blue clock), stabilizing oscillator transcript levels for the duration of cold, thereby stabilising the abundance of clock transcripts. This shifts the phase of the oscillation (e.g. *TaGI* 3A, *TraesCS5B02G271300*). **(F)** the cold-responsiveness of a transcript is restricted to a particular phase, and cold also delays the phase of the circadian clock. In combination, this upregulates and delays the accumulation of the transcript (e.g. *TraesCS1A02G289800* and *TraesCS1B02G095000*).

Figure S1. The number DEGs shared between timepoints. Upset plots visualising the size of the intersections between the groups of **(A)** up-regulated, and **(B)** down-regulated genes at each timepoint. In a similar manner to Venn diagrams, filled circles signify the timepoint of interest and the links between filled circles signify the intersection of interest, the size of which is reported by the height of the bar immediately above. Set size represents the total amount of DEGs detected at each timepoint.

Figure S2. The number and proportion of transcripts from each subgenome that have a circadian gated down regulated response to cold. Small group sizes for each Estimated Time of Maximum Responsiveness group resulted in large variation in the proportion of transcripts derived from each subgenome.

Figure S3. Dynamic contribution of homoeologs to triad cold responsiveness. The

relative contributions of homoeologs to total triad transcript abundance before and after cold

treatment split by the estimated time of maximum responsiveness (ETMR) groups. Ternary

plots as described in Fig. 4C and cold treatment data is that presented in Fig. 4D.

Figure S4. Gene ontology-term analysis of cold-induced transcripts with a circadian

gated response to cold. GO term enrichment within each estimated time of maximum

responsive cold upregulation. Circle size represents the number of transcripts associated

with the GO term, and circle colour indicates the Benjamini-Hochberg adjusted weighted

Fisher p-value.

Figure S5. Gene ontology-term analysis of transcripts with a time delay in cold CPM

profiles relative to control. GO term enrichment for transcripts with rhythmic control and

cold CPM profiles (meta3d p <0.05) and a delay ratio less than 0.5. Circle size represents

the number of transcripts associated with the GO term, and circle colour indicates the

Benjamini-Hochberg adjusted weighted Fisher p-value (adjusted p < 0.001).

Figure S6. **Example transcripts demonstrating the regulatory features described in**

Figure 6. Transcript profiles that exemplify the concepts in Fig. 6. **(A)**

TraesCS5D02G447500. **(B)** *TraesCS3B02G287400*, **(C)** *TraesCS3D02G325300*, **(D)**

TraesCS5A02G468300, **(E)** *TraesCS5B02G271300*, **(F)** *TraesCS5D02G076400*.

Yellow/grey shading = subjective day/night. Solid lines are mean (N = 3 biological

replicates). Blue/green shading = ± s.e.m. Red line represents ∆CPM.

Figure S7. Appearance of seedlings that were sampled, and light spectra used for

cultivation. (**A**) Representative images of bread wheat seedlings after germination on damp

filter paper, followed by 12 days growth on compost under 12 h : 12 h light dark cycles

(22°C) followed by 24 h of constant light. Seedlings were equivalent to Zadok Stage GS1.2

at time of sampling. Red boxes indicate region of second leaf that was sampled. Scale bar

20 mm. (**B)** Spectra of LED lights used for plant cultivation and experimentation, for control

and cold temperature conditions.

Figure S8. Rhythmic transcripts at the upper limit of meta2d statistical threshold are

considered candidate genes with a circadian-gated response to cold. Example

transcripts that have rhythmic ΔCPM (meta2d p <0.05) and p-value close to the cut-off limit,

demonstrating the level of statistical stringency was appropriate for detection of circadian-

gated transcripts. Transcripts are from ETMR grouping **(A)** < 4 h, *TraesCS1A02G210400*,

(B) 4 − 8 h, *TraesCS5D02G343300*, **(C)** 8 − 12 h, *TraesCS5D02G318100*, **(D)** 12 − 16 h,

TraesCS5A02G311100, **(E)** 16 − 20 h, *TraesCS3B02G342936*, **(F)** > 20 h,

TraesCSU02G072000. Yellow/grey shading = subjective day/night. Solid lines are mean (N

= 3 biological replicates). Blue/green shading = ± s.e.m. Red line represents ΔCPM.

Dataset S1. Gene-level counts for entire time-series and transcriptome.

Dataset S2. Counts per million data for entire time-series and transcriptome.

Dataset S3. The number of genes considered from each timepoint/treatment replicate and

the number of genes considered before and after Voom/Limma filtering of low-expressed

genes**.**

1021

1022

(A) Entrainment — Free-running conditions

Separate cold treatments and sampling

3h chilling — X6
Control

14 days
12h light 12h dark
24 h
Constant light
ZT 0   ZT 24   ZT 28   ZT 32   ZT 36
Sample

(B)

Log fold-change

ZT28   ZT32   ZT36   ZT40   ZT44   ZT48

Zeitgeber Time (ZT)

● Up-regulated   ● Down-regulated

(C) Number of DEGs

ZT28
ZT48   ZT32
5000
2500
0
ZT44   ZT36
ZT40

(D) Percentage unique DEGs (%)

ZT28
ZT48   ZT32
100
50
0
ZT44   ZT36
ZT40

● Up-regulated   ● Down-regulated

(E)

Clustering group number: 1 2 3 4 5 6 7 8 9 10 11 12

Relative cold induced change
+4
0
−4

ZT28 ZT32 ZT36 ZT40 ZT44 ZT48

Zeitgeber Time (ZT)

Time in constant light [h]

Control (22°C)   3 h chilling (4°C) treatments

(A)

(B)

(C)

(D)

Estimated time of maximum responsiveness

Number of groups triad represented in: ● 1 ● 2 ● 3

(A) Transcript delay: rhythmic ΔCPM

(B) Transcript delay: rhythmic CPM profiles only

(C) ΔCPM vs Adjusted ΔCPM

↑ ΔCPM
↑ Adj.ΔCPM

(D) Transcript delay: rhythmic ΔCPM

Delay Ratio
Total **882**
< 1 **709** 78.9%
< 0.5 **214** 23.8%
> 1 **190** 21.1%

(E) Transcript delay: rhythmic CPM profiles only

Delay Ratio
Total **3849**
< 1 **3098** 80.5%
< 0.5 **1008** 26.2%
> 1 **751** 19.5%

(F)

TaELF3
TaELF4-L
TaGI
TaLHY
TaLUX-La
TaLUX-Lb
TaLUX/BOA
TaPRR37
TaPRR59
TaPRR73
TaTOC1

Delay Ratio [Mean.Adj.ΔCPM / Mean.ΔCPM]

(G)

TaGI 3A

TaLUX/BOA 3A

Time in Constant Light [h]

Control (22°C)   3 h chilling (4°C) treatments   ΔCPM

(A) Circadian regulated but not cold responsive

(B) Cold-responsive but not circadian regulated

(C) Circadian regulated, and also cold responsive at peak phase

(D) Circadian regulated and cold responsive at different phases

(E) Cold induced pause in circadian rhythm (phase shift)

(F) Cold induced pause in circadian rhythm and cold responsive at shifted peak phase