Computation-guided redesign of promoter specificity of a bacterial RNA polymerase

Xiangyang Liu[1,‡], Anthony T. Meger[2,3,‡], Thomas Gillis[2,‡], and Srivatsan Raman[2,3,4,5,*]

[1]Biophysics Graduate Program, University of Wisconsin-Madison, Madison, Wisconsin 53706, United

States

[2]Department of Biochemistry, University of Wisconsin-Madison, Madison, Wisconsin 53706, United States

[3]Great Lakes Bioenergy Research Center, University of Wisconsin-Madison, Madison, Wisconsin 53706,

United States

[4]Department of Bacteriology, University of Wisconsin-Madison, Madison, Wisconsin 53706, United States

[5]Department of Chemical and Biological Engineering, University of Wisconsin-Madison, Madison,

Wisconsin 53706, United States

[‡]These authors contributed equally to this work

Contact Information

*Corresponding author: Srivatsan Raman (sraman4@wisc.edu)

ABSTRACT

The ability to regulate circuits and pathways is central to cellular control. The existing toolkit is

predominantly comprised of local transcription regulators that are unsuitable for exerting control

at a global genome-wide scale. Bacterial sigma factors are ideal global regulators as they direct

the RNA polymerase to thousands of transcription sites. Here, we redesigned the promoter

specificity of the *E. coli* housekeeping sigma factor, sigma-70, toward five orthogonal promoter

targets not recognized by the native sigma-70. These orthogonal sigma-70s were engineered by

screening a pooled library of computationally-guided designs customized toward each promoter

target. A combination of conserved interactions with the DNA backbone and target-specific

interactions facilitate new promoter recognition. Activity of the top performing redesigned sigma-

70s varied across the promoter targets and ranged from 17% to 77% of native sigma-70 on its

canonical active promoter. These orthogonal sigma factors represent a new suite of regulators

for global transcriptional control.

KEYWORDS: Protein-DNA interactions; Computational design; biological orthogonalization;

specificity; high-throughput screening

INTRODUCTION

Synthetic biology aims to control cellular behavior through synthetic pathways and circuits. These pathways

and circuits depend on orthogonal gene regulation to selectively insulate them from host processes while

sharing central dogma resources and machinery with the host.[1-5] In bacteria, native gene regulation is

governed at the local and global levels.[6-8] At the local level, individual genes and operons are controlled by

transcription factors that are activated by specific metabolites or environmental cues. These local regulators

(i.e., small-molecule inducible transcription factors) have been co-opted and engineered to build a large

toolkit of orthogonal gene regulatory components for synthetic circuits and pathways.[9-12] At a global level,

the RNA polymerase (RNAP) is directed to hundreds to thousands of loci simultaneously by a battery of

sigma factors to maintain homeostasis and respond to environmental changes.[13] In contrast to local

regulators, efforts to engineer RNAP to exert control at a global level remain poorly developed. Although

some genes or operons can be regulated from synthetic promoters with non-native regulation, large

swathes of the bacterial genome, especially genes involved in core metabolic processes, require native

regulation as they are often layered in complex networks that cannot be easily disentangled.[3,14] Orthogonal

regulation in these genes can be introduced by modifying RNAP promoter specificities without disrupting

native regulation.[15] With emerging technologies to synthesize genomes from scratch, genome-scale

orthogonal gene regulation could be programmed into cells.[16,17] Such a system could have different RNAP

promoter specificities encoded in different regions of the genome to compartmentalize them. Thus,

redesigning the RNAP is a powerful but underexplored route to engineer orthogonal genetic regulation.

The sigma subunit of the bacterial RNAP governs the interaction of the RNAP with the promoter.[18-20] The

bacterial RNAP is a multi-subunit molecular machine composed of alpha, beta, omega, and sigma subunits.

The alpha, beta, and omega subunits form the core RNAP enzyme responsible for RNA synthesis using

DNA as a template and ribonucleotides as the substrate.[21,22] To initiate transcription, the core RNAP binds

to a sigma factor which directs the RNAP to specific promoters.[19,20,23,24] Bacteria express several types of

sigma factors that can recognize and bind specific promoters in response to cellular signals and

environmental conditions.[24-26] Of the seven types of sigma factors in *E. coli*, sigma-70 is the dominant as it

is responsible for expressing housekeeping genes, and the other sigma factors are only transiently

expressed during stress.[27-32] Prior strategies to engineer RNAP have used heterologous sigma factors from

other hosts or constructed chimeras of native and non-native sigma factors to engineer new RNAP promoter

specificities.[33-35] However, heterologous sigma factors are often toxic to the host, and chimeric sigma

factors are not orthogonal, limiting the extent to which they may be used in synthetic systems.[36,37]

Additionally, the use of heterologous and chimeric sigma factors is strain-dependent.[33] Despite the

attractiveness of the sigma factor as a target for orthogonal genetic regulation, we do not have well-

developed strategies to engineer them.

In this study, we redesigned the *E. coli* sigma-70 to recognize and transcribe from five orthogonal promoters

that are transcriptionally inactive to wild-type sigma-70, using computation-guided design and pooled cell-

based screens. We used Rosetta to create a customized library of computationally redesigned variants of

the domain of *E. coli* sigma-70 that interacts with each promoter target. Fluorescence-activated cell sorting

followed by deep sequencing of the sigma libraries revealed the sequence determinants of promoter

specificity and how they changed for different targets. We found that recognition of new promoters occurs

through a combination of highly conserved residues that likely make polar interactions with DNA backbone

and target specific adaptations. We identified orthogonal sigma variants for each of the five targets whose

activities ranged from 17-77% of wildtype sigma-70 on a highly active canonical *E. coli* promoter. Our

workflow is generalizable and can be applied to alternative sigma factors in *E. coli* and sigma factors from

other host organisms. The redesigned sigma factor-promoter pairs constitute an important tool for

engineering orthogonal genetic regulation at the genome level.

RESULTS

The partitioned functional domains of sigma-70 enable targeted redesign of promoter specificity

As the "housekeeping" regulator of transcription in *E. coli*, sigma-70 performs three functions. First, sigma-

70 must recognize the consensus -10 (TATAAT) and -35 (TTGACA) promoter elements. The -10 element

is recognized by conserved domains 2 and 3, and the -35 element is recognized by the helix-turn-helix motif

of domain 4 (domain 1 is unstructured) (**Fig. 1a**).[38,39] Second, sigma-70 is responsible for recruiting the

core RNA polymerase complex to the promoter. The interface between RNA polymerase and sigma-70 is

extensive (>8000 Å^2) as it spans the entire length of domains 2-4.[40] The third function of sigma-70 is to

mediate the melting of DNA near the -10 region, which is a critical step for initiating transcription. To operate

as a global regulator of cellular transcription, sigma-70 must perform each of these functions with high

efficiency and fidelity.

To engineer orthogonal sigma variants, we must alter promoter specificity while preserving interactions with

the core RNA polymerase complex and those required for DNA melting. Therefore, we aimed to restrict the

design to key residues of the helix-turn-helix motif of domain 4 to alter promoter specificity through the -35

DNA element. These include residues R584, E585, R586, R588, and Q589 (**Fig. 1a, box**). In the structure

of the *E. coli* transcription initiation complex (PDB ID: 4YLN), the positively charged R586 and R588 form

polar interactions with the negatively charged phosphate backbone to likely promote general DNA affinity.[39]

Specific interactions with the nucleotide bases are formed between R584:Gua-31* and E585:Ade-34* in

the major groove of the -35 element. Although oriented towards the major groove, Q589 does not directly

interact with the -35 element. Instead, Q589 is sterically packed between residues E585, R586 and R588

and, likely plays a role in properly orienting these critical residues within the recognition helix. Given their

structural roles, we hypothesized that redesigning these five key positions could engineer orthogonal

promoter specificity of sigma-70.

Our design workflow begins with identifying -35 element mutants that are not recognized by wildtype sigma-

70. These mutants became recognition targets for the computational redesign of the major groove helix

residues of wildtype sigma-70 (**Fig. 1b**). The top 1000 ranked designs were synthesized as chip-based

oligonucleotide library and evaluated in pooled fashion using a cell-based GFP screen. In this assay, high

GFP fluorescence signifies a specificity switch of sigma-70, as the variant must recognize the orthogonal -

35 element to initiate transcription of GFP. To identify specificity switches from the pooled screen, we

compared the distribution of variants between sorted and presorted states by deep sequencing (**Fig. 1c**).

Clonal screens were then used to validate several successful redesigns for each targeted -35 element.

Fig. 1. Engineering sigma-70 variants with novel promoter specificity. (**a**), Structure of the *E. coli*
transcription initiation complex (PDB ID: 4YLN). sigma-70 (red) bound to the -10 and -35 DNA elements
(gray, cartoon representation) and RNAP core complex (gray, surface representation), composed of α (two
copies), β, β', and ω subunits (left). Canonical -35 element recognized by the helix-turn-helix motif of sigma-
70 domain 4 and -10 element recognized by domains 2 and 3 (middle). Orientation and h-bond contacts of
residues R584, E585, R586, R588, and Q589 of the domain 4 recognition helix in the -35 major groove
(right). (**b**), Workflow for altering sigma-70 promoter specificity *in silico* and characterizing function *in vivo*.
Cartoon schematic of the RNAP holoenzyme with WT sigma-70 bound to the canonical -35 element (left).
Replacement of the canonical -35 with a novel target and computational redesign of the domain 4
recognition helix to engineer sigma-70 variants with orthogonal promoter specificities (middle). *In vivo*
functional characterization using a GFP gene placed downstream of the promoter variants containing novel
-35 targets (right). (**c**), Platform for high-throughput selection and characterization of redesigned sigma-70
variants. FACS to isolate GFP positive (i.e., functional variants) (left), deep sequencing to identify enriched
variants (middle), and clonal testing of redesigned sigma-70s to validate function (right).

Synthetic promoter modifications enhance sigma-70 dependence on the -35 element

As mentioned above, sigma-70 utilizes a two target DNA recognition system, with domain 4 recognizing

the canonical -35 element, and domains 2 and 3 recognizing the canonical -10 element. Due to this two-

target dependence, engineering an orthogonal sigma factor would require the redesign of multiple sigma-

70 DNA binding domains and their respective consensus DNA sequences. However, by artificially

increasing dependence of DNA recognition on the -35 DNA element, redesign is simplified to a one domain-

one target problem. There are several advantages to engineering orthogonality through the -35 element

rather than the -10 element: (1) -35 recognition is mediated by a single domain of sigma-70, (2) interactions

between the helix-turn-helix motif of domain 4 with the -35 major groove are structurally well understood,

and (3) interactions between sigma-70 and the -10 element are much more complex as transcription

initiation is mediated by DNA melting near this region. Thus, we sought to increase sigma-70's dependence

on the -35 element, and then engineer sigma-70 variants that recognize orthogonal -35 targets.

We first placed a gene encoding GFP downstream of a strong constitutive *E. coli* promoter with canonical

-35 'TTGACA' and -10 'TATAAT' sequences (apFab71)[41] on a "reporter plasmid" to use fluorescence as a

measure of transcriptional activity. As expected, a negative control lacking both canonical -10 and -35

promoter elements resulted in very low fluorescence relative to the apFab71 promoter (**Fig. 1a**). We

generated a randomized (NNNNNN) library of the -35 sequence to assess the dependence on the -10 site.

Moderate fluorescence was observed in the population of cells expressing the randomized -35 library,

suggesting that the canonical -10 element alone is partially sufficient to recruit sigma-70. To attenuate the

-10 dependence and increase the -35 dependence, we aimed to identify a promoter variant with high

transcriptional activity with the canonical -35 sequence, but significantly lower activity when paired with a

disrupted -10 sequence. To this end, we characterized a small subset of nucleotide deletion variants (Δ(-

11), Δ(-16,-7), Δ(-8), Δ(-7)) near the -10 region of apFab71. The Δ denotes deletion of a nucleotide at a

specific site within the -10 site. We quantified the -35 dependence of each -10 deletion variant by comparing

the fluorescence ratio of the deletion variant with a clonal canonical -35 and the randomized -35 population.

As desired, all four apFab71 promoter variants retained -35 dependence (i.e., lower fluorescence in the

randomized condition) and displayed approximately a 10-fold reduction in fluorescence relative to the native

apFab71 promoter with a randomized -35 sequence (**Fig. 2a**). The Δ(-7) promoter exhibited the greatest -

35 dependence with a 330-fold difference in fluorescence between the canonical and randomized -35

sequences. Thus, the Δ(-7) promoter was selected as the starting construct for redesigning the promoter

specificity of sigma-70.

Endogenous *E. coli* sigma-70 displays very weak activity on selected -35 targets

We next sought to identify a set of five promoter targets orthogonal to the canonical -35 element recognized

by endogenous *E. coli* sigma-70. From the library of randomized -35 sequences of the apFab71 Δ(-7)

promoter, we used a high-throughput 96-well screening approach to isolate and characterize promoter

variants not recognized by native sigma-70. We isolated 74 unique -35 variants with a 500-fold weaker

sigma-70 transcriptional activity (GFP fluorescence <10^3 RFU) relative to apFab71 (**Supplementary Table**

1). Comparing the sequences of these low activity -35 variants did not reveal a strong pattern suggesting

that there are many unique ways to disrupt sigma-70 promoter recognition (**Fig. 2c**). Weak conservation

was observed at Cyt-31 of the canonical -35 (TTGACA), but other positions deviated from the canonical

sequence which is also reflected by their median Hamming distance of 5 (**Supplementary Table 1**). From

these low activity -35 variants, we selected five diverse target sequences for engineering sigma-70 variants

with novel specificities: TTCATC, GGAACC, CCGCCG, GCTACC, and CCCCTC. These promoter targets

all displayed low transcriptional activity (500- to 900-fold lower fluorescence than apFab71), comparable to

the no promoter control, and vary in their composition and Hamming distances from the canonical -35

sequence (**Fig. 2d**). Due to their diversity, we hypothesized that these five -35 sequences present would

present unique design challenges and demonstrate the breadth of our redesign capabilities.

Fig. 2. Sigma-70 dependence on the -35 element and selection of orthogonal promoter targets. (**a**), Fluorescence distributions of *E. coli* cells expressing native sigma-70 and containing plasmid constructs with a GFP gene placed downstream of no promoter (gray), the strong constitutive apFab71 promoter (green), or the apFab71 promoter with a randomized -35 element (blue). (**b**), Median flow cytometry measured fluorescence of promoter variants containing nucleotide deletions (Δ) near the -10 region, and either a canonical or randomized -35 element. Fold-change between canonical and randomized -35 states was used to evaluate -35 transcriptional dependence. (**c**), Sequence logos (-35 region) of 74 unique clonal isolates with low GFP fluorescence (<10^3 RFU) (**Supplementary Table 1**). Clones were screened in 96-well format from the apFab71 Δ(-7) randomized -35 library. (**d**), Selected -35 DNA targets for engineering orthogonal sigma-70 variants. Nucleotide sequences (left) and median fluorescence (right) using native sigma-70. Error bars denote the standard deviation of replicate flow cytometry experiments (n≥2). RFU, relative fluorescence units. OD, optical density measured at 600nm.

Rosetta calculations reveal the sigma-70 sequence preferences for each promoter target

To engineer sigma-70 variants that recognize each of the five -35 targets (TTCATC, GGAACC, CCGCCG, GCTACC, and CCCCTC), we first performed large-scale combinatorial mutagenesis of residues R584,

E585, R586, R588, and Q589 in the helix-turn-helix (HTH) motif of domain 4 using the Rosetta

macromolecular modeling suite.[42-44] We excluded I587 because it is located along the interface of the HTH

and makes no DNA contacts. The design workflow is as follows. We replace the canonical -35 sequence

(TTGACA) of the sigma-70:DNA co-crystal structure (PDB: 4YLN, **Fig. 1a**) with a target -35 sequence that

we experimentally validated was unable to initiate transcription using native sigma-70 (**Fig. 1b**). Then, we

perform an *in silico* scan to evaluate the stability of the protein-DNA complex for all possible single, double,

triple, and quadruple variants (total ~724,000 sigma-70 variants) of the sigma-70 recognition helix (Fig. 1a,

boxed, middle panel). Calculations for all ~724,000 sigma-70 variants against the five target -35 elements

were performed on a high-throughput computing (HTC) system, requiring approximately 400,000

computing hours (2-3 weeks real-time). Computed protein:DNA interface scores for the ~724,000 sigma

variants in complex with the promoter targets were used to curate designs for experimental testing (**Fig.**

3a, Supplementary Fig. 1a). Lower interface scores indicate a more stable complex. Interestingly,

differences between median interface scores among sigma-70 variants revealed a hierarchy of promoter

preference of the sigma-70 scaffold (TTGACA < TTCATC < GGAACC < CCGCCG < GCATCC < CCCCTC)

with interface scores decreasing with increasing Hamming distance from the consensus -35 sequence,

TTGACA.

This suggests a smooth binding landscape of sigma-70 DNA recognition where incremental mutations from

the canonical -35 sequence decrease binding strength. To assess sigma-70 sequence preferences with

each promoter, we computed position-specific amino acid enrichment scores (E_{score}) for the top 1000

scoring interfaces (**Fig. 3b, Supplementary Fig. 1b**). We observed that substitutions at positions 584 and

589 did not have an impact on the protein-DNA interface scores suggesting that these are indirect contacts

that likely do not drive affinity. Nonetheless, the most enriched residue at both positions was not the wildtype

amino acid and was distinct for each promoter site. This suggests that though these may be indirect

contacts, they could play a role in promoter specificity. In contrast, positions 585, 586 and 588 show subtle,

but distinct preferences for different binding sites. Enrichment heat maps indicate sigma-70 positions 585,

586, and 588 undergo stringent selection across all -35 targets. These positions directly contact DNA in the

crystal structure of native sigma-70 in complex with the canonical (TTGACA) -35 element (**Fig.1a**). As

expected, positively charged and/or large hydrophobic amino acids are generally preferred at these

positions, which can facilitate highly favorable hydrogen bonding, electrostatic and/or van der Waals

interactions with the DNA. For all promoters, arginine and tyrosine were the most enriched amino acids at

positions 586 and 588, respectively. Both residues make non-specific hydrogen bonds with the sugar-

phosphate backbone of DNA (**Fig. 3c, Supplementary Fig. 1c**). Clear differences in amino acid preference

were evident for position 585. For the -35 target TTCATC, arginine was the most enriched amino acid,

which forms two hydrogen bonds with Gua-35* in the Rosetta structural models. Similarly, a single hydrogen

bond is formed by tryptophan at position 585 with Gua-35* or Thy-35* for promoters CCCCTC and

GCATCC, respectively. In contrast, bulky hydrophobic residues phenylalanine and leucine make van der

Waals contacts with the CCGCCG and GGAACC promoters. Because positions 584 and 589 do not directly

interact DNA, substitutions at these positions had less distinct effects on the Rosetta computed interface

scores.

The redesigned sigma-70 variants were ranked by binding energy, and the highest affinity 1000 variants

for each target -35 element were selected for experimental testing. In addition, for targets TTCATC,

GGAACC, and CCGCCG, we selected 1000 variants with binding energies comparable to native sigma-70

and the canonical -35 element (-26.0 REU), as this affinity may be optimal for DNA recognition and promoter

release during transcription initiation. For GCATCC and CCCCTC, the "highest affinity" and "WT-like" sets

were identical because the binding energy distributions of these targets were less favorable (**Fig. 3a**). We

synthesized these variant libraries on an IPTG-inducible plasmid expression system using oligonucleotide

chip synthesis and one-pot cloning.

Fig. 3. Rosetta guided design of sigma-70 variants for each promoter target. (a), Interface scores of
all (gray) or 1000 top scoring (red) sigma-70 variants in complex with each -35 DNA target. Dashed lines
indicate the median interface scores of all single, double, triple, and quadruple combinatorial variants of
sigma-70 positions 584, 585, 586, 588, and 589 modeled with Rosetta. (**b**), Position-specific amino acid
enrichment scores (red gradient) among selected top scoring sigma-70 variants. WT identity (boxed outline)
and most enriched amino acid (*) at each mutable position. (**c**), Cartoon schematic showing H-bonds
formed between each enriched sigma-70 consensus sequence and -35 DNA target in the Rosetta structural
models.

Isolation and identification of sigma-70 variants with novel promoter specificities

To synthesize the computationally curated library of sigma-70 designs for each promoter target, we used

chip-based oligonucleotides and one-pot cloning. The library of sigma-70 designs for each target was

independently cloned into an expression plasmid and placed under IPTG control. Each library was

transformed into *E. coli* harboring their cognate reporter plasmid i.e., GFP gene placed downstream of the

respective target -35 element. Due to the "housekeeping" role of sigma-70, overexpression of a sigma-70

variant can competitively sequester the RNAP core enzyme to inhibit cell viability. This toxicity challenge

has been reported in previous works using heterologous sigmas and domain swaps.[33-35] To identify an

expression level with minimal toxicity, we first varied the IPTG inducer concentration in a clonal growth

assay. Minimal growth deficiency was observed using 5µM IPTG, and thus, this concentration was used

for all subsequent experiments (**Supplementary Fig. 2**).

To isolate and enrich high activity sigma-70 redesigns, we sorted the high fluorescence subpopulation of

each library in the IPTG induced condition. Comparison of the fluorescence distributions of cell populations

in the induced and uninduced states show distinctly higher fluorescence in the induced condition for all five

target promoters (**Fig. 4a, Supplementary Figs. 3,4a**). This suggested that some sigma-70 redesigns may

indeed be able to successfully transcribe from an orthogonal -35 sequence. The presorted and sorted

libraries were deep sequenced to identify sigma-70 redesigns that were enriched after selection.

Approximately 10-15% of all sigma-70 variants within each 1000-member library were enriched (**Fig. 4b,**

Supplementary Fig. 4b). Taken together, these results validate our *in silico* redesign approach and high-

throughput screening platform.

Sequence profiles of redesigned sigma-70 variants are unique to each -35 target

Following deep sequencing, we evaluated the sigma-70 sequence profiles among enriched library variants

of each -35 target. Given that the 5 promoter targets are separated by a median Hamming distance of 4,

we expected the variable positions of sigma to undergo unique sequence selection (i.e., position-specific

amino acid enrichment) to recognize each -35 target (**Supplementary Fig. 5**). While some similarities are

shared across the sequence profiles for the five targets, each consensus sequence is unique (**Fig. 4c,**

Supplementary Fig. 4c). Arginine was highly enriched at position 586, whereas both tyrosine and arginine

were enriched at position 588 for all targets. In the WT sigma-70:DNA co-crystal structure (PDB: 4YLN)

and Rosetta structural models, positions 586 and 588 interact primarily with the sugar-phosphate backbone

of DNA (**Fig. 3c**). Thus, the enrichment of charged and/or polar residues at these positions is consistent

with forming favorable non-specific DNA interactions. These structures also show base-specific interactions

facilitated by position 585, which is where we observe several differences in enrichment across the

promoter targets: arginine is favored by targets TTCATC and CCGCCG, bulky hydrophobic residues (I, F,

and L) by GGAACC, aromatic residues (W, Y, and F) by CCCCTC and a wider range of residue types (W,

R, I, and F) by GCATCC (**Fig. 4c**). Selection was less stringent at positions 584 and 589 of sigma-70 for

all targets, as a broad range of amino acid identities were found among enriched variants. These positions

often do not directly interact with the promoter in the Rosetta models. However, charged residues (R and

H) were more enriched at positions 584 and 589 by targets GGAACC and GCATCC, which may

electrostatically enhance DNA affinity. Additional interactions may be required to facilitate binding because

the interactions by position 585 with the respective -35 targets are less specific. These results demonstrate

that promoter specificity of sigma-70 is dictated by compounding differences in primary sequence that

modulate both base-specific and general DNA affinity interactions.

Fig. 4. Selection and identification of successfully redesigned sigma-70 variants. (**a**), Flow cytometry
fluorescence distributions of uninduced (gray) and IPTG induced (green) sigma-70 variant populations for
each -35 target after FACS-based selection of functional redesigns. Transcriptionally active variants were
enriched using two rounds of sequential GFP positive cell sorting. (**b**), Distributions of log-transformed
enrichment scores of all characterized sigma-70 variants after selection. Deep sequencing was performed
on the presorted and sorted libraries to compute enrichment ratios. (**c**), Sequence logos showing the
weighted amino acid frequencies at each mutable position among functionally enriched sigma-70 variants.
Amino acid identities are colored by chemical properties: polar amino acids (N, Q, S, T) shown in green,
basic (H, K, R) blue, acidic (D, E) red, hydrophobic (A, I, L, M, V) purple, aromatic (F, W, Y) orange, and
other (C, G, P) gray.

Redesigned sigma-70's exhibit varying levels of activity on the promoter targets

To validate the results of our high-throughput screen, we evaluated the activity of clonal isolates from our redesigned sigma-70 libraries. A total of 96 clones from the sorted libraries were randomly sampled, and from this set, we were able to identify redesigned sigma-70s with activity on each of the five promoter targets (**Fig. 5a, Supplementary Table 2**). To assess the performance of each sigma variant, we first measured the fluorescence of cells containing the target promoters upstream of GFP in the presence of endogenous WT sigma-70. Under this condition, the promoter variants yielded low fluorescence (<400 RFU OD^{-1}) for the TTCATC, GGAACC, CCGCCG, GCTACC, and CCCCTC targets (**Fig. 5a**). In comparison, the activity of endogenous WT sigma-70 on the canonical -35 element (TTGACA) resulted in much higher fluorescence (10,200 ± 800 RFU OD^{-1}) (**Supplementary Fig. 6**). Next, we measured the fluorescence of each clone on their respective -35 target sequence. Of the 96 colonies screened, 25-50% exhibited activity resulting in at least a four-fold increase in fluorescence over the endogenous WT sigma-70 baseline on the target promoter (**Supplementary Table 2**). The proportion of 'successful' redesigns in the sorted libraries is consistent with the flow cytometry profiles (**Fig. 4a**). The activities of the top three performing sigma-70 redesigns for each target varied across the targets. The best performing design had 77% of the activity of endogenous WT sigma-70 on the canonical -35 element (**Fig. 5a**). For the target TTCATC, which is closest in Hamming distance to the canonical -35 sequence (**Fig. 2d**), we found the three highest performing redesigned sigma-70s (CRRVY, FIQRY, and FWCRY). However, performance does not scale with Hamming distance, suggesting a more complex relationship exists between a given -35 target and the success of computationally redesigned sigma-70s.

Rosetta structural models reveal potential mechanisms of -35 target recognition

In addition to the low transcriptional activation by WT sigma-70, the five -35 targets were selected due to their sequence deviation (i.e. orthogonality) from the canonical -35 and each other. The utility of engineered sigma factors is dependent on minimizing crosstalk between various sigma factor-target pairs. By targeting diverse -35 sequences with our redesign of sigma-70, we sought to generate novel and specific protein-DNA interactions for each sigma factor-target pair. Evaluating Rosetta structural models generated for the highest performing sigma-70 redesigns, we observe several differences between the hydrogen bonding

patterns of our redesigned sigma-70 variants with their cognate -35 targets (**Fig. 5b).** Some interactions

were largely conserved across target sequences, such as the h-bonds between residues 586 and 588 of

sigma and the sugar-phosphate backbone of nucleotides -35 and -*31, despite the diverse polar and

charged residues employed to maintain these interactions. In contrast to the other -35 targets, the hydrogen

bonding patterns of the top three performing sigma-70 redesigns for targets CCGCCG and CCCCTC are

highly converged. The base-specific interactions between CCGCCG and variants IRRYY, HRRYY, and

CRRYY occur between R585:Gua-33 and Y589:Gua-34*. For CCCCTC, a single base-specific interaction,

W585:Gua-33*, appears sufficient for recognition by sigma-70 variants CWRRP, FWRRP, and MWMRP,

perhaps necessitating the additional non-specific h-bond between R586 and the sugar-phosphate

backbone. However, the number of h-bonds present in these protein-DNA structural models ranges from

one to four, which highlights the importance of other types of molecular interactions (van der Waal's, dipole-

dipole) for promoter recognition. Though computationally derived, these models show how unique sets of

interactions may facilitate -35 target recognition by our successfully redesigned sigma-70 variants.

To understand the relationship between the Rosetta interface scores and the performance of each sigma-

70 redesign, we compared these energies to the enrichment scores from our high-throughput screen (**Fig.**

5c). While the computational screening approach was successful at generating functional variants and

significantly reduced the sequence space search from 3.2 million (20^5) combinations to a manageable

1000-member test set, the highest affinity redesigns (i.e. lowest interface scores) were not among the best

performing. This underscores the necessity of our high-throughput screening platform using FACS and

deep sequencing to isolate functional redesigns. The top three performing sigma-70 variants for each -35

target had Rosetta interface scores comparable to that of WT sigma-70 on the canonical -35 element (-

26.0 REU), suggesting that an 'optimal binding affinity' exists for DNA recognition and transcription initiation

by sigma-70. An optimized affinity may facilitate proper association with the promoter and allow dissociation

of sigma-70 to initiate transcription and elongation.

Fig. 5. Clonal validation of redesigned sigma-70 variants. (**a**), Normalized fluorescence of WT sigma-
70 (gray) and the top three performing variants (red, yellow, or blue) for each promoter target. Clones were
selected from the sorted libraries and assayed in a 96-well fluorescence plate reader. Error bars denote the
standard deviation (**P ≤ 0.01, ***P ≤ 0.001) of replicates (n≥3) and computed fold-changes are relative
to the fluorescence resulting from endogenous WT sigma-70 with each target. Percentages of activity are
relative to the activity of endogenous WT sigma-70 with the promoter containing a canonical -35 element
(**Supplementary Fig. 6**). RFU, relative fluorescence units. OD, optical density at 600nm. (**b**), Cartoon
schematic showing H-bonds formed between each sigma-70 variant and the cognate -35 DNA target in the
Rosetta structural models. (**c**), Rosetta interface scores and enrichment scores from the high-throughput
screen of all tested sigma-70 variants. The top three performing variants are indicated in red (square),
yellow (circle), and blue (triangle).

DISCUSSION

We devised a high throughput *in silico* modeling and *in vivo* screening workflow to redesign promoter specificity of the *E. coli* "housekeeping" regulator of transcription, sigma-70. Using this workflow, we identified multiple sigma-70 variants that activate transcription on 5 target promoters containing diverse -35 elements: TTCATC, GGAACC, CCGCCG, GCTACC, and CCCCTC. These results demonstrate that the promoter specificity of an essential primary transcription regulator can be rationally redesigned. Although the process of transcription initiation (i.e. promoter recognition and DNA melting) by sigma factors is complex, the partitioned functional domains of sigma factors enabled key interactions between the domain 4 recognition helix and the -35 DNA element to be redesigned without perturbing other functions. Our success across all five target sequences demonstrates the extant at which promoter specificity can be altered and the generalizability of our approach. Given that the Hamming distance separation ranges from three to six (of 6 nucleotide positions) between these novel -35 elements and the canonical -35 element, TTGACA, our approach also enables the design of sigma factors with maximal orthogonality from natural sigma factors and thereby, greater potential to minimize crosstalk between synthetic and host expression systems. In contrast, sigma factors generated in previous works using domain swaps of alternative sigma factors or introduced from non-host family bacteria, have been limited in their orthogonality and host compatibility.

One advantage to the engineered sigma-70 variants presented is that they integrate seamlessly into the *E. coli* host, as they utilize the same endogenous transcriptional machinery. While the level of activation varied from 3-fold to 20-fold across our top performing redesigns, we show that this activity was also engineered *de novo*, as the native sigma-70 exhibited little to no activity on each of the orthogonal promoters. Using structural models of these successful redesigns in complex with their cognate promoters, we observe diverse combinations of interactions that facilitate promoter recognition. This diversity extends even among successful redesigns for the same promoter target, demonstrating the utility of computational tools to navigate the energetic landscape and simplify the sequence space to be experimentally tested.

The transcription-level regulatory complexity of a cell is mediated by the concerted efforts of global and

local regulators. While many existing synthetic expression systems are reliant solely on local regulators (i.e.

natural or synthetic transcription factors) to control a sparse set of genes, they lack the complexity and

capabilities of native systems. Only by incorporating global regulatory components, such as engineered

sigma factors, can we expand the utility and complexity of synthetic genetic circuits. However, building or

isolating an extensive repertoire of global regulatory components is challenging because they must be (1)

orthogonal to host systems to minimize crosstalk between the synthetic and native expression networks,

(2) display dynamic control by activating gene expression in response to simple input signals, and (3) be

highly customizable and modular, such that they can be easily integrated into established expression

systems. Here, we developed and validated a generalizable approach to engineer sigma-70 variants that

satisfy all these requirements and can be similarly applied to any host sigma factor to expand the set of

available global regulatory components. Future work could pair these engineered sigma-70 factors with

existing local regulators of transcription in a biosynthetic expression system.

ACKNOWLEDGEMENTS

This work was partly supported by the NIH Director's New Innovator's Award (DP2GM132682-01 to S.R.)

and partly by the Great Lakes Bioenergy Research Center, U. S. Department of Energy, Office of Science,

Office of Biological and Environmental Research (Award Number DE-SC0018409). The content is solely

the responsibility of the authors and does not necessarily represent the official views of the NIH, Department

of Defense, Department of Energy, or other federal agencies.

AUTHOR CONTRIBUTIONS

X.L., A.T.M., T.G., and S.R. conceptualized the project. X.L. devised the cell-based screening system,

cloned the promoter library, and performed sorting and NGS experiments. A.T.M. performed the *in silico*

calculations and designed the sigma variant libraries. T.G. performed clonal screens and identified the top

performing sigma variants for each promoter target. X.L., A.T.M., T.G, and S.R. drafted the manuscript.

A.T.M., X.L., and T.G. created the figures. S.R. supervised the work. All authors edited and approved the

final manuscript.

REFERENCES

Ceroni, F. *et al.* Burden-driven feedback control of gene expression. *Nature Methods* **15**, 387-393 (2018). https://doi.org:10.1038/nmeth.4635

Gyorgy, A. *et al.* Isocost Lines Describe the Cellular Economy of Genetic Circuits. *Biophysical Journal* **109**, 639-646 (2015). https://doi.org:https://doi.org/10.1016/j.bpj.2015.06.034

Müller, I. E. *et al.* Gene networks that compensate for crosstalk with crosstalk. *Nature Communications* **10**, 4028 (2019). https://doi.org:10.1038/s41467-019-12021-y

Slusarczyk, A. L., Lin, A. & Weiss, R. Foundations for the design and implementation of synthetic genetic circuits. *Nature Reviews Genetics* **13**, 406-420 (2012). https://doi.org:10.1038/nrg3227

Costello, A. & Badran, A. H. Synthetic Biological Circuits within an Orthogonal Central Dogma. *Trends in Biotechnology* **39**, 59-71 (2021). https://doi.org:https://doi.org/10.1016/j.tibtech.2020.05.013

Browning, D. F. & Busby, S. J. W. Local and global regulation of transcription initiation in bacteria. *Nature Reviews Microbiology* **14**, 638-650 (2016). https://doi.org:10.1038/nrmicro.2016.103

Martínez-Antonio, A. & Collado-Vides, J. Identifying global regulators in transcriptional regulatory networks in bacteria. *Current Opinion in Microbiology* **6**, 482-489 (2003). https://doi.org:https://doi.org/10.1016/j.mib.2003.09.002

Ishihama, A. & Shimada, T. Hierarchy of transcription factor network in Escherichia coli K-12: H-NS-mediated silencing and Anti-silencing by global regulators. *FEMS Microbiology Reviews* **45**, fuab032 (2021). https://doi.org:10.1093/femsre/fuab032

Taylor, N. D. *et al.* Engineering an allosteric transcription factor to respond to new ligands. *Nature Methods* **13**, 177-183 (2016). https://doi.org:10.1038/nmeth.3696

Rao, C. V. Expanding the synthetic biology toolbox: engineering orthogonal regulators of gene expression. *Current Opinion in Biotechnology* **23**, 689-694 (2012). https://doi.org:https://doi.org/10.1016/j.copbio.2011.12.015

Cameron, D. E., Bashor, C. J. & Collins, J. J. A brief history of synthetic biology. *Nature Reviews Microbiology* **12**, 381-390 (2014). https://doi.org:10.1038/nrmicro3239

Li, J.-W., Zhang, X.-Y., Wu, H. & Bai, Y.-P. Transcription Factor Engineering for High-Throughput Strain Evolution and Organic Acid Bioproduction: A Review. *Frontiers in Bioengineering and Biotechnology* **8** (2020).

Feklístov, A., Sharon, B. D., Darst, S. A. & Gross, C. A. Bacterial Sigma Factors: A Historical, Structural, and Genomic Perspective. *Annual Review of Microbiology* **68**, 357-376 (2014). https://doi.org:10.1146/annurev-micro-092412-155737

Lu, T. K., Khalil, A. S. & Collins, J. J. Next-generation synthetic gene networks. *Nature Biotechnology* **27**, 1139-1150 (2009). https://doi.org:10.1038/nbt.1591

Bervoets, I. *et al.* A sigma factor toolbox for orthogonal gene expression in Escherichia coli. *Nucleic Acids Res* **46**, 2133-2144 (2018). https://doi.org:10.1093/nar/gky010

Gibson, D. G. *et al.* Complete Chemical Synthesis, Assembly, and Cloning of a Mycoplasma genitalium Genome. *Science* **319**, 1215-1220 (2008). https://doi.org:doi:10.1126/science.1151721

Richardson, S. M. *et al.* Design of a synthetic yeast genome. *Science* **355**, 1040-1044 (2017). https://doi.org:doi:10.1126/science.aaf4557

Browning, D. F. & Busby, S. J. W. The regulation of bacterial transcription initiation. *Nature Reviews Microbiology* **2**, 57-65 (2004). https://doi.org:10.1038/nrmicro787

deHaseth Pieter, L., Zupancic Margaret, L. & Record, M. T. RNA Polymerase-Promoter Interactions: the Comings and Goings of RNA Polymerase. *Journal of Bacteriology* **180**, 3019-3025 (1998). https://doi.org:10.1128/JB.180.12.3019-3025.1998

20 Feklistov, A. RNA polymerase: in search of promoters. *Annals of the New York Academy*
of Sciences **1293**, 25-32 (2013). https://doi.org:https://doi.org/10.1111/nyas.12197
21 Burgess, R. R. & Travers, A. A. Escherichia coli RNA polymerase: purification, subunit
structure, and factor requirements. *Fed Proc* **29**, 1164-1169 (1970).
22 A Darst, S. Bacterial RNA polymerase. *Current Opinion in Structural Biology* **11**, 155-162
(2001). https://doi.org:https://doi.org/10.1016/S0959-440X(00)00185-8
23 Helmann, J. D. & Chamberlin, M. J. STRUCTURE AND FUNCTION OF BACTERIAL
SIGMA FACTORS. *Annual Review of Biochemistry* **57**, 839-872 (1988).
https://doi.org:10.1146/annurev.bi.57.070188.004203
24 Paget, M. S. Bacterial Sigma Factors and Anti-Sigma Factors: Structure, Function and
Distribution. *Biomolecules* **5**, 1245-1265 (2015).
25 Kazmierczak Mark, J., Wiedmann, M. & Boor Kathryn, J. Alternative Sigma Factors and
Their Roles in Bacterial Virulence. *Microbiology and Molecular Biology Reviews* **69**, 527-
543 (2005). https://doi.org:10.1128/MMBR.69.4.527-543.2005
26 Gruber, T. M. & Gross, C. A. Multiple Sigma Subunits and the Partitioning of Bacterial
Transcription Space. *Annual Review of Microbiology* **57**, 441-466 (2003).
https://doi.org:10.1146/annurev.micro.57.030502.090913
27 Lonetto, M., Gribskov, M. & Gross, C. A. The sigma 70 family: sequence conservation and
evolutionary relationships. *J Bacteriol* **174**, 3843-3849 (1992).
https://doi.org:10.1128/jb.174.12.3843-3849.1992
28 Marles-Wright, J. & Lewis, R. J. Stress responses of bacteria. *Current Opinion in Structural*
Biology **17**, 755-760 (2007). https://doi.org:https://doi.org/10.1016/j.sbi.2007.08.004
29 Helmann, J. D. The extracytoplasmic function (ECF) sigma factors. *Adv Microb Physiol*
46, 47-110 (2002). https://doi.org:10.1016/s0065-2911(02)46002-x
30 Maeda, H., Fujita, N. & Ishihama, A. Competition among seven Escherichia coli σ
subunits: relative binding affinities to the core RNA polymerase. *Nucleic Acids Research*
28, 3497-3503 (2000). https://doi.org:10.1093/nar/28.18.3497
31 Cowing, D. W. & Gross, C. A. Interaction of Escherichia coli RNA polymerase holoenzyme
containing σ32 with heat shock promoters: DNase I footprinting and methylation
protection. *Journal of Molecular Biology* **210**, 513-520 (1989).
https://doi.org:https://doi.org/10.1016/0022-2836(89)90127-7
32 Koo, B. M., Rhodius, V. A., Nonaka, G., deHaseth, P. L. & Gross, C. A. Reduced capacity
of alternative sigmas to melt promoters ensures stringent promoter recognition. *Genes*
Dev **23**, 2426-2436 (2009). https://doi.org:10.1101/gad.1843709
33 Rhodius, V. A. *et al.* Design of orthogonal genetic switches based on a crosstalk map of
σs, anti-σs, and promoters. *Mol Syst Biol* **9**, 702 (2013).
https://doi.org:10.1038/msb.2013.58
34 Gaida, S. M. *et al.* Expression of heterologous sigma factors enables functional screening
of metagenomic and heterologous genomic libraries. *Nature Communications* **6**, 7045
(2015). https://doi.org:10.1038/ncomms8045
35 Bervoets, I. *et al.* A sigma factor toolbox for orthogonal gene expression in Escherichia
coli. *Nucleic Acids Research* **46**, 2133-2144 (2018). https://doi.org:10.1093/nar/gky010
36 Temme, K., Hill, R., Segall-Shapiro, T. H., Moser, F. & Voigt, C. A. Modular control of
multiple pathways using engineered orthogonal T7 polymerases. *Nucleic Acids Res* **40**,
8773-8781 (2012). https://doi.org:10.1093/nar/gks597
37 Studier, F. W. & Moffatt, B. A. Use of bacteriophage T7 RNA polymerase to direct selective
high-level expression of cloned genes. *J Mol Biol* **189**, 113-130 (1986).
https://doi.org:10.1016/0022-2836(86)90385-2
38 Barne, K. A., Bown, J. A., Busby, S. J. & Minchin, S. D. Region 2.5 of the Escherichia coli
RNA polymerase sigma70 subunit is responsible for the recognition of the 'extended-10'

546		motif at promoters. *Embo j* **16**, 4034-4040 (1997).
547		https://doi.org:10.1093/emboj/16.13.4034
548	39	Zuo, Y. & Steitz, Thomas A. Crystal Structures of the Transcription Initiation Complexes
549		with a Complete Bubble. *Molecular Cell* **58**, 534-540 (2015).
550		https://doi.org:10.1016/j.molcel.2015.03.010
551	40	Paget, M. S. B. & Helmann, J. D. The σ70family of sigma factors. *Genome Biology* **4**, 203
552		(2003). https://doi.org:10.1186/gb-2003-4-1-203
553	41	Mutalik, V. K. *et al.* Precise and reliable gene expression via standard transcription and
554		translation initiation elements. *Nature Methods* **10**, 354-360 (2013).
555		https://doi.org:10.1038/nmeth.2404
556	42	Thyme, S. & Baker, D. in *Homing Endonucleases: Methods and Protocols* (ed David R.
557		Edgell) 265-282 (Humana Press, 2014).
558	43	Ashworth, J. *et al.* Computational redesign of endonuclease DNA binding and cleavage
559		specificity. *Nature* **441**, 656-659 (2006). https://doi.org:10.1038/nature04818
560	44	Joyce, A. P., Zhang, C., Bradley, P. & Havranek, J. J. Structure-based modeling of protein:
561		DNA specificity. *Brief Funct Genomics* **14**, 39-49 (2015).
562		https://doi.org:10.1093/bfgp/elu044
563	45	Campbell, E. A. *et al.* Structural Mechanism for Rifampicin Inhibition of Bacterial RNA
564		Polymerase. *Cell* **104**, 901-912 (2001). https://doi.org:https://doi.org/10.1016/S0092-
565		8674(01)00286-0
566	46	Murakami, K. S., Masuda, S., Campbell, E. A., Muzzin, O. & Darst, S. A. Structural Basis
567		of Transcription Initiation: An RNA Polymerase Holoenzyme-DNA Complex. *Science* **296**,
568		1285-1290 (2002). https://doi.org:10.1126/science.1069595
569	47	Thyme, S. & Baker, D. Redesigning the specificity of protein-DNA interactions with
570		Rosetta. *Methods Mol Biol* **1123**, 265-282 (2014). https://doi.org:10.1007/978-1-62703-
571		968-0_17
572	48	Maguire, E. J., Rocca-Serra, P., Sansone, S.-A. & Chen, M. in *Eurographics Conference*
573		*on Visualization.*
574		
575		

METHODS

Rosetta design

Rosetta design focused on *E. coli* sigma-70 positions R584, E585, R586, R588 and Q589 (PDB: 4YLN) as

previous research has demonstrated that these 5 positions interact with -35 region nucleotides through

hydrogen bonds.[45,46] Rosetta v3.9 was used for all calculations following existing Rosetta design protocol.[47]

5 promoters bearing different -35 sequences (TTCATC, GGAACC, CCGCCG, GCTACC, and CCCCTC)

were chosen after clonal testing, and were desirable due to their low transcription activity in vivo. To

generate starting structures for combinatorial protein mutagenesis, each promoter sequence was first

introduced in the WT sigma-70:DNA complex structure. Nucleotide substitutions were made at the

canonical -35 region (WT=TTGACA).

Input structures were subjected to energy minimization using Rosetta, and lowest energy structures from

50 minimizations were used as the input structure for the combinatorial protein mutagenesis scan at

positions R584, E585, R586, R588, and Q589. Standard resfiles were used to denote mutations at each

position for all variants containing single (95), double (3610), triple (68,590), and quadruple (651,605)

combinatorial mutants. Variants containing 5 mutations (2,476,099) were not tested due to the excessive

burden on computational resources. The structure of each protein variant in complex with all promoters

(including WT) were generated and the Protein-DNA binding affinity was scored. The binding energy was

reported as an average of 10 Rosetta minimized structures. In this minimization procedure, the amino acid

substitutions were first introduced followed by side chain rotomeric state optimization and protein backbone

design. Lastly, the Protein-DNA interface was optimized and scored using the DNA Interface Packer mover.

Out of these designs for each target promoter sequence, 1000 designs were selected based on two criteria.

The first criteria sought to contain variants with the highest binding affinity for target promoter sequence,

i.e. the lowest Rosetta Energy Units (REU). The second criteria sought to contain variants with a binding

energy for target sequence which was closest to the observed binding energy of WT sigma-70 on its native

target sequence.

The reason behind the first criterium is that low REU confers tighter binding to the target promoter

sequence, which could translate into higher transcription level. On the other hand, sigma-70 amino acid

composition and WT -35 sequence are likely the local minima through evolution. Although the pair's REU

is not the lowest among all variants, it might indicate that there is an optimal REU to allow tight enough

binding while permitting the release when RNAP moves downstream. The assumption is that tightest

binding hinders release. For some promoters, there might be overlap between two groups chosen

independently.

Library cloning

A promoter library was ordered via Integrated DNA Technology and amplified by hybridizing with a reverse

primer (hybridization protocol: 95°C for 3min, 55°C for 1min, 72°C for 1min). The product was subsequently

cleaned up using PCR cleanup kit (Omega Biotek) and stored in dH$_2$O. Backbone preparation started with

PCR amplification of plasmid pXL-9. For Rosetta designed sigma variants, 110-base pair (bp) single strand

DNA oligo pool containing Rosetta designed sigma fragments were ordered from Agilent. Oligo design

features unique priming regions at 5' and 3' for individual libraries complemented with BsaI recognition sites

and cutting sites to enable Goldengate cloning. 10ng of the oligo pool was used in the PCR reaction to

amplify individual libraries (PCR protocol: 95°C for 3min, 98°C for 20sec, 55°C for 15sec, 72°C for 8sec,

cycling from step 2 for 20 cycles, 72°C for 30sec). PCR product was subsequently cleaned up using PCR

cleanup kit (Omega Biotek) and stored in dH$_2$O. Backbone preparation started with PCR amplification of

plasmid SC101_LacI_WTsigma containing a WT sigma-70 driven by pLacO promoter. Amplified backbone

bared the corresponding BsaI binding and cutting sites, enabling scarless cloning. Backbone PCR was

cleaned up and subject to sequential digestion of DPN1 and BsaI_HF_V2 (New England Biolab) to remove

the template DNA and expose sticky overhangs. The digested backbone was further incubated with

Antarctic Phosphatase (New England Biolab) to remove 5' and 3' phosphates. The removal of those

phosphates prevents backbones from circularizing and creating false positive transformants. The

Goldengate reaction was done as described on New England Biolab. Briefly, 300ng of the backbone was

combined with 70ng of the library in a 20µL reaction. The mix was incubated at 37°C for 1hr and 65°C for

5min. Cooled reaction mix was dialyzed on 0.02µm filter in dH$_2$O for 1hr at room temperature. The dialyzed

sample was collected and stored at -20°C.

Transformation of promoter and redesigned sigma-70 libraries

Library transformation was conducted using electrocomponent DH10β *E. coli* (New England Biolab). 2µL

of the assembled library reaction mix was transformed with 25µL cells and recovered with 1mL SOC

medium at 37°C for 1hr. Multiple logs of dilutions were plated on appropriate antibiotic plate to measure

transformation efficiency. 4mL of LB and antibiotics were added to ensure proper growth selection. Cultures

were grown overnight before being stored in 25% glycerol at -80°C. For non-library transformations, 10ng

DNA was added directly to 25µL of cells and plated on agar plates containing antibiotics to confer selection.

Co-transformations were conducted by first growing cells containing one plasmid in LB medium with

appropriate antibiotics overnight. 1:50 back dilution was then done into 3mL LB without antibiotics. Cells

were allowed to grow until reaching an OD$_{600nm}$ of 0.6 before being chilled on ice. Chilled cells were spun

down at 5000xg and thoroughly washed with ice cold dH$_2$O twice. The final cell pellet was suspended in

25µL of ice cold dH$_2$O. Electroporation was then repeated similarly to regular plasmid transformation.

Transformed cells were plated onto agar plates with appropriate antibiotics overnight at 37°C before storage

at 4°C.

Induction and fluorescence measurement

Multiple colonies were picked from agar plates and inoculated into 150µL LB medium in a 96-well plate

containing appropriate antibiotics. Cells were allowed to grow shaking (900rpm, multi-well plate shaker,

Southwest Scientific) at 37°C for around 3hr or until OD$_{600nm}$ reached 0.6. Cultures were back inoculated

into two technical replicates of fresh LB medium with appropriate antibiotics at 1:20 dilution factor. After

around 3hr or upon reaching an OD$_{600nm}$ of 0.3, IPTG (at final concentration of 5µM) was added to one of

the two technical replicates. Cells were incubated for 5hr before fluorescence measurement. Fluorescence

measurements were conducted using a multi-well platereader (HTX Biotek). Excitation and emissions

wavelengths of 485nm and 528nm (± 20) were used for all measurements. Measurements were normalized

by dividing fluorescence intensities by the OD$_{600nm}$ before subtracting blank well fluorescence (cells carrying

no GFP). Fold-improvement was calculated by dividing normalized the IPTG induced fluorescence of

sigma-70 variants by the IPTG induced fluorescence of WT sigma-70 containing cells using the same target

promoter sequence. Average fold improvement was calculated from at least 3 biological replicates.

Fluorescence activated cell sorting

Around 25µl of reporter transformed libraries carrying a target promoter upstream of GFP were inoculated

into 3mL LB kan/spec-50 (kanamycin and spectinomycin at 50µg/mL) from glycerol stock and grown

overnight at 37°C in a shaking incubator. Cultures were inoculated into two separate 150µL fresh LB

kan/spec-50 in a 96-well plate at 1:50 dilution. Fresh cultures grew shaking at 37°C for 3hr or until reaching

an OD_{600nm} of 0.3. IPTG was added to a final concentration of 5µM to one technical replicate for induction.

After the addition of IPTG, cultures were grown for 5hr before being chilled on ice. Chilled cultures were

added to ice cold phosphate buffer saline (PBS) at 1:50 ratio and mixed well before being put back onto

the ice. Cell sorting was conducted using a Sony SH800 cell sorter (Sony Biotech) (condition: FCS

threshold=2500, 50% PMT on GFP, ultra-purity, round 8k events/second). Three independent replicates

were sorted. Briefly, cells in PBS were flowed through first to capture the distribution. Under 5µM IPTG

induced condition, top 10% cells were sorted out (at least 50k events). Sorted populations were flowed

through the sorter to capture sorting efficiency (percentage of cells fall back into the sorted gate). Sorted

cells with less than 80% efficiencies were sorted using the same gate again to improve the purity. Sorted

cells were recovered in 1mL LB medium shaking at 37°C. 100µL (1:10 dilution) of cells were plated onto

LB agar plates with appropriate antibiotics. The rest were growing overnight in 3ml LB with appropriate

antibiotics before being miniprepped into plasmid. Promoter library sorting was done by simply isolating 5%

of cells with lowest fluorescence.

Variant identification using next generation sequencing (NGS)

A pair of primers upstream and downstream of the mutated sigma-70 region were used to add partial

Illumina adaptors. Priming sites were at least 20 base pair away from the variable regions to ensure good

quality base calling during NGS. Amplicon size was 120 base pair. Amplicons were sequenced on

Amplicon-EZ Miseq service (Genewiz) with at least 50k reads for each sigma library. Sequenced libraries

were processed with PEAR pair-ended merger (quality threshold set to 35). Merged reads were processed

with custom script to identify variants. Briefly, sigma-70 variable regions were extracted and converted into

amino acids with corresponding frequency. Reads with frequencies of less than 5 were removed. Variant

frequency was normalized against total reads to get percentages. To compute individual variant's

enrichment, a variant's percentage under 10μM IPTG induction was divided by that of the condition without

IPTG. Sequence logos for variants with highest enrichment were generated using ISA-tools.[48]

Clonal identification

To identify functional sigma-70 variants, 100 colonies were sampled from the sorted populations. Highly-

functional (defined by its fold improvement level) colonies were measured using the above protocol for plate

reader measurements, and high performing colonies were subsequently identified using Sanger

sequencing. Unique high-performing variants were then inoculated in 3 mL kan/spec LB media and grown

overnight. Following overnight growth, plasmids from these variants were extracted using the protocol for

plasmid mini-prep. Following plasmid extraction, plasmids from these variants were re-transformed into

DH10β *E. coli* in order to normalize for variations in the *E. coli* genome of sorted cells that may have led to

false positives in GFP transcription during clonal testing. Using three biological replicates, the re-

transformed sigma-70 variants were measured once again using the induction protocol from initial clonal

screens. High-performing sigma-70 variants were measured alongside cells containing WT sigma-70 and

the orthogonal promoter sequences to ensure that fold-improvement scores are consistent across

measurements.

 SUPPLEMENTARY INFORMATION

708 **Supplementary Table 1**. **Normalized fluorescence measurements of clones containing randomized**
709 **-35 sequences.** GFP fluorescence and OD_{600nm} measurements were collected with a 96-well plate reader.
710 Hamming distances for each sequence are relative to the canonical -35 element. The five selected low
711 activity -35 targets are highlighted in green.
712

-35 sequence	Normalized fluorescence (RFU OD^{-1})	Hamming distance from TTGACA
ATGACT	316	2
AAGCCA	426	3
TTATCC	488	3
TTCATC	320	3
TTCTCC	492	3
AGGATG	627	4
ATACCC	340	4
ATATCC	381	4
ATCTCT	420	4
CCGCCC	445	4
CCGCCG	491	4
CGGCCC	474	4
CTCGCC	426	4
CTGCTC	452	4
CTTTCT	436	4
GCGAGT	404	4
GCGCCT	548	4
GCGGCC	413	4
GGAACC	585	4
GTCGCC	439	4
TCTCCC	483	4
TTAGTG	494	4
TTCTTT	514	4
AAACCC	601	5
AAAGTA	497	5
AACGCC	475	5
AAGTGT	341	5
ACGGTT	434	5
AGACCC	376	5
AGAGCC	431	5
AGTTCC	478	5
ATTTTT	479	5
CCCTCC	416	5
CCTTCT	487	5
CGAGCC	413	5
CGCCCG	523	5
CGCGCC	396	5
CGGCTC	527	5
CGTCCT	533	5

CGTGCC	394	5
CGTTCT	489	5
CTCCTT	459	5
GACTCT	422	5
GATCCC	421	5
GATGCT	397	5
GATTCC	436	5
GCATCC	454	5
GCATCT	491	5
GCTTCT	560	5
GGCCCC	310	5
GTCTTT	510	5
GTTTTT	473	5
TACTTT	311	5
TATTTT	376	5
TGCGTT	327	5
TGCTTC	572	5
AATTTT	421	6
ACATTT	402	6
CACTTT	442	6
CATTGC	487	6
CCCCTC	554	6
CCTTTC	521	6
CCTTTT	314	6
CGCTTT	328	6
CGTCTT	509	6
CGTTTC	409	6
CGTTTT	401	6
GAATGT	316	6
GACTTT	354	6
GATCTT	457	6
GATTTT	363	6
GCCTTT	349	6
GCTTTT	400	6
GGCTTT	413	6

713
714

Supplementary Fig. 1. Computation-based selection of redesigned sigma-70 variants for three
promoter targets. (**a**), Interface scores of all (gray) or 1000 most WT-like (Interface score = -26REU, red)
sigma-70 variants in complex with each -35 DNA target. WT-like sets were not created for promoter targets
CCCCTC and GCATCC because they were redundant with the lowest energy sets. All single, double, triple,
and quadruple combinatorial variants of sigma-70 positions 584, 585, 586, 588, and 589 were modeled
using Rosetta. (**b**), Position-specific amino acid enrichment scores (red gradient) among selected wt-like
scoring sigma-70 variants. WT identity (boxed outline) and most enriched amino acid (*) at each mutable
position. (**c**), Cartoon schematic showing H-bonds formed between each enriched sigma-70 consensus
sequence and -35 DNA target in the Rosetta structural models.
725

Supplementary Fig. 2. Sigma-70 expression induced toxicity. Each row corresponds to a different sigma-70 variant. Each column represents a different IPTG concentration. Color coding represents cell density measured by OD_{600}nm after 5 hours of induction. Dark represents high cell density while white represents low cell density.

732
733 **Fig. 3. Selection of successfully redesigned sigma-70 variants.** Flow cytometry fluorescence
734 distributions of uninduced (gray) and IPTG induced (green) sigma-70 variant populations for each -35 target
735 prior to FACS-based selection of functional redesigns. Transcriptionally active variants were enriched by
736 sorting the GFP positive subpopulation of the lowest energy (left) and WT-like (right) libraries of each target.
737

738
Supplementary Fig. 4. Selection and identification of successfully redesigned sigma-70 variants.
(**a**), Flow cytometry fluorescence distributions of uninduced (gray) and IPTG induced (green) sigma-70
variant populations for three -35 targets after FACS-based selection of functional redesigns.
Transcriptionally active variants from the WT-like libraries were enriched using two rounds of sequential
GFP positive cell sorting. (**b**), Distributions of log-transformed enrichment scores of all characterized sigma-
70 variants after selection. Deep sequencing was performed on the presorted and sorted libraries to
compute enrichment ratios. (**c**), Sequence logos showing the weighted amino acid frequencies at each
mutable position among functionally enriched sigma-70 variants. Amino acid identities are colored by
chemical properties: polar amino acids (N, Q, S, T) shown in green, basic (H, K, R) blue, acidic (D, E) red,
hydrophobic (A, I, L, M, V) purple, aromatic (F, W, Y) orange, and other (C, G, P) gray.
749

	CCCCTC	CCGCCG	GCATCC	GGAACC	TTCATC
CCCCTC					
CCGCCG	3				
GCATCC	4	4			
GGAACC	5	5	2		
TTCATC	3	6	5	4	

Supplementary Fig. 5. Hamming distances between each of the five selected -35 targets.

754 **Supplementary Table 2. Clonal screens of sigma variants after FACS-based selection of high**
755 **activity variants.** GFP fluorescence and OD_{600nm} measurements were collected with a 96-well plate reader.
756 96 clones were tested for each -35 target. Normalized GFP fluorescence was compared to WT sigma on
757 each target to compute fold-improvement.
758

-35 target	Total number of clones screened	Number of clones with fold-improvement ≥4
GGAACC	96	50
CCCCTC	96	21
TTCATC	96	51
GCATCC	96	23
CCGCCG	96	35

759
760

762   **Supplementary Fig. 6. Activity of endogenous WT sigma-70 on the canonical and target -35**

763   **elements.** GFP fluorescence and OD_{600nm} measurements (n ≥ 3) were collected with a 96-well plate reader.