AP2XII-2 coordinates transcriptional repression for *Toxoplasma*

***gondii* sexual commitment**

Sandeep Srivastava[1#], Michael J. Holmes[1#], Michael W. White[3],

William J. Sullivan, Jr[1-2*]

[1]Department of Pharmacology & Toxicology, [2]Department of Microbiology &

Immunology, Indiana University School of Medicine, Indianapolis, 46202, Indiana

[3]Department of Global Health, University of South Florida, Tampa, Florida

Running title: AP2XII-2 role in *Toxoplasma* sexual development

[#] Equal contribution

*Corresponding author:

William J. Sullivan, Jr.

Showalter Professor

Indiana University School of Medicine

635 Barnhill Drive

MS A418C

Indianapolis, IN 46202

(317) 274-1573; wjsulliv@iu.edu

ABSTRACT

     Toxoplasma gondii is a widespread protozoan parasite that has significant impact on human and veterinary health. The parasite undergoes a complex life cycle involving multiple hosts and developmental stages. How *Toxoplasma* transitions between life cycle stages is poorly understood, yet central to controlling transmission. Of particular neglect are the factors that contribute to its sexual development, which takes place exclusively in feline intestines. While epigenetic repressors have been shown to play an important role in silencing spurious gene expression of sexually committed parasites, the specific factors that recruit this generalized machinery to the appropriate genes remains largely unexplored. Here, we establish that a member of the AP2 transcription factor family, AP2XII-2, is targeted to genomic loci associated with sexually committed parasites along with the epigenetic regulators of transcriptional silencing, HDAC3 and MORC. Despite widespread association with gene promoters, AP2XII-2 is required for silencing of relatively few genes. Using CUT&Tag methodology, we identify two major genes associated with sexual development downstream of AP2XII-2 control, AP2X-10 and the amino acid hydroxylase AAH1. Our findings show that AP2XII-2 is a key contributor to the gene regulatory pathways modulating *Toxoplasma* sexual development.

IMPORTANCE

Toxoplasma gondii is a parasite that undergoes its sexual stage exclusively in feline intestines, making cats a major source of transmission. A better understanding of the proteins controlling the parasite's life cycle stage transitions is needed for the development of new therapies aimed to treat toxoplasmosis and transmission of the infection. Genes that regulate the sexual stages need to be turned on and off at the appropriate times, activities that are mediated by specific transcription factors that recruit general machinery to silence or activate gene expression. In this study, we identify a transcription factor called AP2XII-2 as being important for repression of a subset of sexual stage genes, including a sexual stage-specific AP2 factor (AP2X-10) and a protein (AAH1) required to construct the infectious oocysts expelled by infected cats.

INTRODUCTION

The ubiquitous parasite *Toxoplasma gondii* (phylum Apicomplexa) infects warm-blooded animals the world over, including approximately one in three people (1). After an initial acute phase of infection by tachyzoites, the parasites redistribute within the solid organs of the infected host and reside as bradyzoites in latent tissue cysts that appear impervious to the immune response and currently available treatments (1). Persistence within the host makes *Toxoplasma* an opportunistic infection in those with compromised immune systems (1). Upon immune suppression, latent bradyzoite stage parasites reconvert into rapidly growing tachyzoites, causing severe tissue destruction that can threaten the life of the infected individual (1).

Given the importance of interconversion between tachyzoites and bradyzoites to disease progression, the molecular mechanisms governing this stage transition have received a great deal of attention (2). The conversion of tachyzoites to bradyzoites requires extensive reprogramming of gene expression at multiple levels, directed in part by chromatin remodeling histone acetyltransferases and deacetylases that are recruited to specific gene loci through their interactions with sequence-specific transcription factors (3, 4). In support of this, members of the AP2 transcription factor family have been found to associate with a cadre of histone modifying complexes including GCN5b/ADA2a and HDAC3/MORC (5, 6). AP2 transcription factors harbor a plant-like DNA-binding domain and have been implicated in contributing to the regulation of gene expression in apicomplexan parasites (7).

In contrast, our understanding of the molecular mechanisms governing the

rewiring of gene expression during *Toxoplasma* sexual development is far less

developed, largely due to a lack of model systems. This is despite evidence showing

that ingestion of the ultimate product of *Toxoplasma* sexual reproduction, sporulated

oocysts, accounts for a substantial portion of worldwide human infections and mass

outbreaks of acute toxoplasmosis (8, 9).

The early stages of *Toxoplasma* sexual development occur in the intestinal lining

of cats. Upon ingestion by the feline, the parasites invade the intestinal enteroepithelial

cells and undergo several rounds of schizogonic replication to produce merozoites (10).

Successive rounds of schizogony have been subcategorized into types A-E of the

enteroepithelial stage. These pre-sexual stages then undergo gametocytogenesis,

ultimately producing motile male microgametes and intracellular female macrogametes.

After fertilization, up to a billion immature oocysts are expelled with the cat's feces (11).

Upon exposure to the air, the oocysts sporulate and become capable of transmitting

infection (10).

It has recently been shown that transcriptional repression mediated by the

HDAC3/MORC complex plays an important role in preventing the aberrant expression

of genes normally restricted to the sexual stages in felids (6). The HDAC3/MORC

complex associates with at least 11 AP2 family transcriptional regulators (6, 12),

suggesting that these factors mediate targeted silencing of felid-specific developmental

gene expression, as proposed for bradyzoite formation (4).

We initially found AP2XII-2 to be in association with AP2IX-4, a transcriptional

repressor that helps coordinate bradyzoite formation (12, 13). We then reported that

AP2XII-2 is required for proper progression of parasites through the S-phase of the

tachyzoite cell cycle and that knockdown of AP2XII-2 increased frequency of bradyzoite

formation *in vitro* (12). We and others have also reported the association between the

HDAC3/MORC complex and AP2XII-2 (6, 12), suggesting that this AP2 family member

controls the transcriptional repression of certain genes. However, the specific genes

regulated by AP2XII-2 remained unexplored.

To better understand the role of AP2XII-2 in *Toxoplasma* biology, we profiled the

genome-wide chromatin-binding sites of AP2XII-2. By adopting Cleavage Under Targets

and Tagmentation (CUT&Tag) for use in *Toxoplasma*, we found find a high degree of

overlap between AP2XII-2 and HDAC3/MORC target genes. Using a transcriptomics

approach, we also found that AP2XII-2 controls the expression of a subset of

HDAC3/MORC-target genes, including some that are restricted to the sexually-

committed developmental stages such as AP2X-10 and the aromatic amino acid

hydroxylase, AAH1. Our results indicate that AP2XII-2 helps coordinate recruitment of

the HDAC3/MORC complex to specific gene loci to repress developmentally-controlled

genes.

RESULTS AND DISCUSSION

AP2XII-2 shows widespread occupancy at gene promoters and high degree of

overlap with the HDAC3/MORC complex

To identify genes that are direct targets of AP2XII-2 in tachyzoites, we applied

the CUT&Tag method (14) to profile the genomic loci bound by the factor. To pulldown

AP2XII-2 for CUT&Tag, we took advantage of a parasite clone we made previously in

ME49 strain that expresses an HA-tagged version of the endogenous protein (12).

Results showed that AP2XII-2-associated regions are widespread across the parasite

genome, displaying a total of 5,527 peaks associated with 3,939 genes (Table S1; Fig.

1A). Consistent with its putative role as a transcriptional regulator, AP2XII-2 is located

near the transcriptional start site of protein coding genes (Fig. 1B).

Previous reports have demonstrated that AP2XII-2 is a member of the multi-

component HDAC3/MORC complex, which silences genes by inducing heterochromatin

formation (6, 12). To corroborate this association, we re-analyzed the existing HDAC3

and MORC ChIPseq data (6) to determine the overlap between HDAC3, MORC, and

AP2XII-2 localization genome-wide (Table S1). Like AP2XII-2, both HDAC3 and MORC

are enriched at the transcriptional start site of protein coding genes (Fig. 1B). Just over

80% of AP2XII-2 enriched regions overlapped with HDAC3 and/or MORC enriched

peaks (Fig. 1C). Additionally, over 85% of genes associated with AP2XII-2 were also

bound by HDAC3 and/or MORC with a clear plurality of all identified genes bound by

the three proteins (Fig. 1D). Given that AP2 domains bind distinct DNA motifs and that

HDAC3 and MORC lack DNA binding domains (15, 16), combined with the previous

observation that they operate within a shared complex (6, 12), the high degree of

coordination between AP2XII-2, HDAC3, and MORC binding sites strongly suggests

that AP2XII-2 is responsible for recruiting the HDAC3/MORC complex to repress

specific genes.

Loss of AP2XII-2 disrupts a small subset of protein coding transcripts in tachyzoites

Given that AP2XII-2 associates with a large number of gene promoters and likely directs transcriptional silencing through the HDAC3/MORC complex, we wanted to assess the consequences of AP2XII-2 depletion on transcriptional regulation. To maximize the robustness of this experiment, we compared the response to AP2XII-2 depletion in both type I (RH strain) and type II (ME49 strain) backgrounds using parasites we made in a previous study (12). These parasites express endogenous AP2XII-2 tagged at its C-terminus with a dual tag comprised of an HA epitope and an auxin-inducible degron (AID). TheAP2XII-2^{AID-HA} protein in both strains was degraded to undetectable levels within 24 hours of adding 500 µM indole-3-acetic acid (IAA) to the culture media (Fig. 2A).

Using these conditions, comparative RNA sequencing (RNAseq) analyses revealed that the loss of AP2XII-2 in tachyzoites led to dysregulation of mRNA levels for a modest number of transcripts in both the RH and ME49 genetic backgrounds (Table S2). In RH parasites lacking AP2XII-2, 66 genes were significantly upregulated and 39 genes were downregulated at least 1.5 fold (Fig. 2B). Similarly, in ME49 parasites, 143 genes were significantly upregulated and 120 genes were downregulated 1.5 fold following depletion of AP2XII-2 (Fig. 2C). The majority of the dysregulated genes in RH strain were also dysregulated in ME49 strain, indicating a high degree of concurrence between strains (Fig. 2D). Surprisingly, there was no correlation between the direction of transcriptional dysregulation and AP2XII-2 gene occupancy as determined by CUT&Tag (Fig. 2E). This observation likely speaks to the complexity of gene regulation

in *Toxoplasma*, which can involve multiple AP2 factors and chromatin remodeling

machinery. Depletion of a single factor like AP2XII-2 may not be sufficient to

significantly disrupt expression of most genes in the network. Since AP2XII-2 was also

found to interact with AP2IX-4 (12), our observations here may point to a model where

AP2 family members cooperatively bind for efficient HDAC3/MORC recruitment to

specific gene loci.

Since AP2XII-2 is a cell cycle regulated factor with expression peaking during the

S/M phases (12, 17), we assessed whether the dysregulated genes were also subject to

cyclic control. Using the cell cycle microarray data available on ToxoDB.org (17, 18), we

determined the average cell cycle profiles for the genes that were up- or downregulated

in response to AP2XII-2 depletion and compared them to the cell cycle pattern exhibited

by AP2XII-2 (Fig. 2F). Downregulated genes tended to be cell cycle regulated with a

valley of expression that spans mid-G1 phase, in line with our previous observation that

AP2XII-2 depletion delays the cell cycle after centrosome duplication (12). The

expanded G1 phase occupancy induced by AP2XII-2 depletion may point to a linkage

between transcriptional repression of AP2XII-2 target genes and the cell cycle. Despite

not displaying cell cycle regulation, the upregulated genes were expressed at lower

robust multichip average (RMA) expression values. The relatively low abundance of

these genes in tachyzoites could indicate that they are normally subjected to

transcriptional repression. Given the association of AP2XII-2 with the HDAC3/MORC

transcriptional repression complex (6, 12), we suspected that these upregulated genes

could be subject to regulation by AP2XII-2 via recruitment of HDAC3/MORC.

Loss of AP2XII-2 increases expression of genes normally restricted to latent and

sexually committed parasites

To identify genes that are most likely to be direct targets of the AP2XII-2-directed

HDAC3/MORC complex, we looked for genes that were upregulated in 1) either ME49

or RH strain parasites upon AP2XII-2 depletion, 2) after 24 hours of MORC depletion

(Fig. S1A), and 3) after 18 hours incubation with the HDAC3 inhibitor FR235222 (Fig.

S1B). These latter two criteria were assessed using publicly available data and were

also generated from a type II parasite strain (Table S3) (6). In addition, we verified that

each gene's promoter was occupied by AP2XII-2, HDAC3, and MORC. Five gene

accession entries met this stringent filtering criteria (Table 1).

Table 1. Genes occupied by AP2XII-2, HDAC3, and MORC that are consistently

upregulated upon their depletion or inhibition.

| Gene ID | Product Description | AP2XII-2 depletion | | MORC[a] | HDAC3[a] |
		ME49[a]	RH[a]		
TGME49_212710	phenylalanine-4-hydroxylase (F4H)	2.18	0.86	4.29	3.74
TGME49_287510	aromatic amino acid hydrolase (AAH1)	0.58	1.40	4.91	3.22
TGME49_318750	deoxyribose-phosphate aldolase (TgDPA)	1.82	1.30	1.39	1.40
TGME49_215340	AP2X-10 (fragment)	1.16	1.01	5.19	3.57
TGME49_215343	hypothetical protein (AP2X-10 fragment)	1.13	1.00	5.27	3.45

[a] $\log_2$FC values

The auxin-induced depletion of MORC as well as inhibition of HDAC3 in

tachyzoites induced upregulation of bradyzoite-specific genes as well as genes

associated with sexual stages (6, 19). Notably, when assessing the degree of

dysregulation for the genes outlined in Table 1, the effect was more pronounced for

MORC and HDAC3 than for AP2XII-2 depletion. This observation is consistent with a

model whereby multiple AP2 family members cooperatively recruit the HDAC3/MORC

complex to genomic loci.

It has been proposed that parasites must pass through the bradyzoite stage

before initiating sexual commitment (6, 20). Our analysis (Table I) identified a single

bradyzoite-specific gene, TgDPA (21), to be reliably upregulated under all three

treatments that were assessed, strongly suggesting that its expression is controlled by

an AP2XII-2-directed HDAC3/MORC complex. The role that TgDPA plays in bradyzoite

biology and their potential commitment to sexual development remain to be elucidated.

Interestingly, the AP2 family member AP2X-10, whose current fragmented gene

annotation incorrectly assigns the gene to both TGME49_215340 and

TGME49_215343, is upregulated when either AP2XII-2, MORC, or HDAC3 is impaired

independent of parasite strain (Table 1). AP2X-10 expression has been noted to be

restricted to the enteroepithelial and oocyst stages; it is the most abundant of all AP2

factors in oocysts, being expressed at the 96[th] percentile (22-24). Seventeen AP2 family

members have been proposed to act downstream of the HDAC3/MORC complex to

regulate various developmental trajectories, with AP2X-10 predicted to guide micro-

and/or macrogamete development (6). Our present study is consistent with AP2XII-2

coordinating the repression of AP2X-10, placing the former above AP2X-10 in a

transcriptional regulatory cascade that contributes to the coordination of *Toxoplasma*

sexual development. Future studies to determine the genes directly controlled by AP2X-

10 and clarify its role in sexual development are warranted.

AAH family gene organization

Two aromatic amino acid hydroxylase enzymes, AAH1 and F4H, are not only

counted amongst the highest upregulated genes upon AP2XII-2, MORC, and HDAC3

inhibition (Table 1), but have also been reported to be important for oocyst maturation

(25). Interestingly, we found that F4H was upregulated in ME49 but not RH strain, AAH1

showed the opposite trend (Table 1). These observations prompted us to further

examine the entire aromatic amino acid hydroxylase family in *Toxoplasma*, which

consists of three annotated members, TGME49_287510 (AAH1), TGME49_212740

(AAH2), and TGME49_212710 (F4H).

Previous reports have indicated that the current ME49 genome assembly

available on ToxoDB.org has incorrectly fused a tandem repeat of the AAH2 gene (25,

26). Before proceeding with our analysis, we had to ensure the fidelity of the AAH2

gene sequence. To that end, we compared the ME49 genome assembly with the RH88

assembly since the former was generated using short read sequencing technology,

which can have difficulty with repetitive regions (Fig. 3A). The RH88 genome was

constructed based on long read sequencing technology, which is better at determining

overall genome architecture (27). This comparison revealed that indeed two members

of the AAH family should appear on chr V of the ME49 genome, spaced apart by

unplaced contigs KE139705 and KE139818 (Fig. 3A).

The three ME49 members of the AAH family share over 99.4% identity at the

nucleotide level in their coding sequences (Fig. 3B), suggesting that the vast majority of

RNAseq reads mapping to the CDS region would have been assigned to all AAH genes

indiscriminately; this spurious read assignment may indicate that a putative strain-

specific upregulation of AAH1 or F4H could occur by chance (Table 1). Given the high

degree of sequence identity between each AAH family member, we examined their

flanking sequences to differentiate them. AAH1 and AAH2 have unique 3'end flanking

sequences (Fig. 3C), however the unplaced contig KE139818 abruptly ends, leaving the

annotated F4H gene (TGME49_212710) without a 3'UTR sequence (Fig. 3D). Analysis

of the unique regions flanking the 5' and 3' ends of the AAH family by sequence

alignment comparison with the RH88 genome architecture allowed us to place AAH1

and AAH2 onto chr V and determine that F4H and AAH2 are likely the same gene (Fig.

3A). However, it should be noted that a previously conducted copy number variant

analysis of the AAH family in *Toxoplasma* indicated that ME49 parasites harbor three

copies of AAH genes whereas type I parasites encode only two (25). We were unable to

map the location of the additional AAH family member with our strategy of comparing

ME49 and RH genome architectures.

Depletion of AP2XII-2 de-represses AAH1 but not AAH2

To distinguish which AAH family member(s) are upregulated upon AP2XII-2

depletion, we endogenously tagged AAH1 and AAH2 with MYC epitopes at their C-

termini in ME49 AP2XII-2^{AID-HA} parasites. The tagging of these genes was confirmed by

verifying successful recombination at each genetic locus by PCR analysis (Fig. S2). In

untreated parasites, AAH1^{MYC} was not detectable by western blotting whereas AAH1^{MYC}

appeared at the expected size of 62 kDa upon IAA-induced loss of AP2XII-2 (Fig. 4A).

Expression of AAH2^{MYC} was not seen under either condition (Fig. 4B).

Neither AAH family member is typically expressed in asexual parasites (26),

suggesting that they are maintained in a transcriptionally repressed state in non-

sexually committed parasites. Our results suggest that AAH genes may be regulated by

different HDAC3/MORC-AP2 complexes, with AP2XII-2 directing the silencing of AAH1

but not AAH2. However, we cannot rule out the possibility that AAH1 and AAH2 have a

parasite strain specific dependence on AP2XII-2. Since parasites lacking AAH1 have a

severe defect in infection of the cat intestine and oocyst production (25), we propose

that AP2XII-2 is a key factor required for completion of *Toxoplasma* sexual

reproduction.

We next examined AAH1 expression by immunofluorescence assay (IFA) and

found AAH1^{MYC} to be localized around the nuclei of tachyzoites only upon AP2XII-2

depletion (Fig. 4C). This staining pattern is reminiscent of the *Toxoplasma*

mitochondrion. To further characterize AAH1^{MYC} localization, we co-stained AP2XII-2$^{AID\text{-}}$

^{HA} knockdown parasites with antibodies to the F$_1$B ATPase mitochondrial marker (28)

and the MYC epitope tag, which confirmed the co-localization of AAH1^{MYC} with the

mitochondrion in tachyzoites (Fig. 4D).

The AAH family of enzymes is responsible for converting phenylalanine to

tyrosine and tyrosine to 3,4-dihydroxy-L-phenylalanine (L-DOPA) (29). The later

compound is a precursor for the formation of di-tyrosine protein crosslinks that are

thought to play a critical structural role in *Toxoplasma* oocyst wall integrity and formation

(25). Given the role that AAH family enzymes play in ensuring oocyst development and

wall integrity, we were surprised to see AAH1 localized to the parasite mitochondrion.

Interestingly, a previous report demonstrated that AAH2 overexpression in bradyzoites

gave rise to a staining pattern consistent with mitochondrial localization (26), suggesting

that the mitochondrion might act as a hub for AAH enzymatic activity. Future studies

using bona fide sexually-committed parasites are needed to verify this possibility.

Conclusion

We have determined that in addition to affecting tachyzoite cell cycle progression

(12), AP2XII-2 also plays a role in coordinating *Toxoplasma* sexual development. This

may indicate that AP2XII-2 plays different roles in different developmental stages or

may suggest that the commitment to sexual development is a cell-cycle dependent

phenomenon as has been demonstrated for the tachyzoite to bradyzoite transition (7,

30). Consistent with this model, either inhibition of HDAC3 or MORC depletion also stall

cell cycle progression and lead to increased bradyzoite and sexual stage gene

expression (6, 19), albeit to a greater extent than does AP2XII-2 depletion. Given the

association of AP2XII-2 with AP2IX-4 (12), and the observation that AP2XII-2 depletion

appears to cause a milder version of the HDAC3 and MORC phenotypes, we propose

that AP2XII-2 is likely one of several AP2 family members that bind cooperatively to

recruit the HDAC3/MORC to specific genomic loci.

On the other hand, the number of reports demonstrating that *Toxoplasma* studies

investigating gene regulation using in vitro culture system lack the fidelity of in vivo

parasite development (31-33). This is particularly relevant when examining the roles of

322 AP2 family members in gene regulation (33). Notably, a previous report indicating that

323 AP2IX-9 is a repressor of bradyzoite formation was later contradicted using a

324 developmentally competent *Toxoplasma* models (15, 22, 33). Given that we have

325 reported that AP2XII-2 impacts both cell cycle regulation (12) and the expression of

326 sexually committed gene expression, we cannot rule out the possibility that mis-

327 expression of AP2XII-2 may have resulted in 'transcriptional confusion' that impacted

328 tachyzoite cell cycle dynamics. Examining the role that AP2XII-2 plays throughout the

329 complete life cycle stages of *Toxoplasma* development is warranted.

330  The intrinsic factors responsible for regulating *Toxoplasma* sexual development

331 have been understudied due to the lack of amenable systems. This is beginning to

332 change with the pivotal discoveries of the upstream extrinsic signals required to initiate

333 sexual development and the requirement of widespread transcriptional repression to

334 coordinate the process (6, 34). Future work will undoubtedly focus on disentangling the

335 gene regulatory networks responsible for initiating and managing *Toxoplasma* sexual

336 development. Here we have begun unravelling one of the networks responsible for this

337 process by placing the AP2XII-2-directed HDAC3/MORC complex above two genes that

338 function in sexual stages, AP2X-10 and AAH1. In addition to utilizing a candidate-based

339 approach as we have relied on for this study, the recent report highlighting inhibitors of

340 AP2-family member activity may prove yet another tool for interrogating the role of AP2

341 factors in *Toxoplasma* developmental transitions (35).

343 **MATERIALS AND METHODS**

Parasite culture

Human fibroblast cells (HFFs, ATCC SCRC-104) were maintained in Dulbecco's modified Eagle medium (DMEM; Corning) supplemented with 10% fetal bovine serum (FBS, R&D systems), 100 µg/ml streptomycin and 100 units/ml penicillin as described elsewhere (36). RH and ME49 parasites were cultured in DMEM supplemented with 1% and 5% heat inactivated FBS, respectively, 100 µg/ml streptomycin and 100 units/ml penicillin. Parasites were maintained in humidified incubator at 37°C and 5% CO_2. Transfected and clonal strains were maintained in selection media supplemented with 1 µM pyrimethamine or a combination of 25 µg mycophenolic acid and 50 µg xanthine as appropriate. Control of AP2XII-2 expression was achieved by adding either 500 µM of indole-3-acetic acid or an equivalent amount of ethanol vehicle for 24 hours to the culture media before analysis.

Generation endogenously tagged AAH1^{MYC} or AAH2^{MYC}

The ME49-TIR1:AP2XII-2^{AID-HA} strain that we generated in a previous study (12) was used as the parental strain for generating dual-tagged AAH1^{MYC} and AAH2^{MYC} parasites. For AAH1, the guide RNA sequence from the pSAG1::CAS9-U6::sgUPRT plasmid (Addgene 54467) (37) was mutagenized to target the stop codon of AAH1. All primers and sequences are listed in supplement (Table S4). A donor repair template with regions of homology to the AAH1 CDS and 3'UTR was made to incorporate the MYC tag and HXGPRT cassette. Parasites were transfected with 25 µg of CAS9 plasmid and 25 µg of donor repair template, were cultured with 25 µg mycophenolic acid and 50 µg xanthine for 7-8 days and single clones were isolated by limiting dilution

in 96 wells. Isolated single clones were confirmed for C-terminal tagging by PCR of their

genomic DNA (Fig. S2)

To generate AAH2^{Myc} parasites, specific primers were used to clone ~2 kb of the

AAH2 gene into the PacI site of the pLIC-3xMYC-HXGPRT plasmid. 50 µg of the final

construct was linearized with NheI restriction endonuclease and transfected into the

ME49-TIR1:AP2XII-2^{AID-HA} strain. Transgenic parasites were selected for 7-8 days

under drug pressure and subjected to limiting dilution in 96 well plates. Single clones

were confirmed for integration of MYC tag at C-terminus (Fig. S2).

Western blotting

Parasites growing in confluent HFF cells were mechanically lysed and filtered to

remove host cell debris followed by centrifugation. Parasite pellets were lysed in

NuPAGE buffer, sonicated, and boiled for 5 minutes. Parasite lysates were centrifuged

briefly to remove debris and separated on 4-12% NuPAGE gels (Invitrogen). Separated

proteins were transferred to PVDF membrane and blocked in 5% Nonfat dry milk-TBST

solution for 1 hour at room temperature. Blocked PVDF membranes were incubated

with anti-HA (1:2000, Roche), anti-MYC (1:3000, Cell Signaling) or anti-SAG1 (1:5000,

ThermoFisher) antibodies in blocking buffer at 4°C overnight with gentle shaking.

Membranes were washed with TBST and incubated with HRP-conjugated anti-rat or

anti-mouse (1:2000; GE healthcare) for 1 hour at room temperature. After washing with

TBST, membranes were developed with SuperSignal West Femto substrate

(ThermoFisher).

Immunofluorescence assay

Parasites growing in HFF monolayers were fixed in 4% paraformaldehyde solution for 15 minutes at room temperature followed by washing with phosphate buffered saline (PBS). Fixed cells were permeabilized and blocked with blocking buffer (3% BSA + 0.2% Triton X-100 in PBS) for 1 h. Cells were incubated with primary antibodies in blocking buffer at 4°C temperature overnight. Cells were washed with PBS and finally incubated with DAPI (1:1000) and Alexa fluor secondary antibodies coupled with desired fluorophore for 1 hour at room temperature. Cells were washed and mounted with Prolong Gold anti-fade reagent (Invitrogen) before analysis with a Leica DMI6000 B inverted microscope. Dilutions for antibodies used for IFA are as follows: Rabbit α-HA (1:2000, Sigma, H6908), Mouse mAb α-MYC (1:1000, Cell signaling, 9B11), Rabbit α-MYC (1:2000, ThermoFisher, PA1-981) and Mouse mAb 5F4 (1:5000, F1B ATPase, a gift from Peter Bradley), Alexa Fluor 488/594 secondary antibodies (1:2000, ThermoFisher) were used as appropriate.

RNA sequencing

Parasites were allowed to invade confluent HFF monolayers for 24 h. Infected HFF monolayers were scraped, and syringe lysed with 23-gauge needle to release intracellular parasites and passed through a 3 μM filter. Filtered parasites were centrifuged at 300 x g for 15 minutes and DMEM media was removed to isolate the parasite pellet. The parasite pellet was resuspended in 1 x PBS and centrifuged at 300 x g for 15 minutes, twice. Total RNA was extracted with TRIzol reagent (Invitrogen) as per manufacturer's protocol and measured by Nanodrop. RNAseq libraries were

414   prepared from polyA-purified mRNA by Azenta using standard Illumina protocols.

415   Libraries were sequenced PE x 150bp.

RNAseq data analysis

RNAseq read data generated from either the RH or the ME49 genetic

background were aligned to the ME49 reference sequence (v52) with hisat2 using

default settings (38). Differential gene expression was conducted with DESeq2 using

default settings (39). Raw sequencing data and processed files generated in this study

are available at GEO GSE217226 and data pertaining to HDAC3 and MORC

impairment (6) were obtained from GEO GSE136123.

CUT&Tag assay

Chromatin occupancy for ME49^{TIR1} parasites expressing AP2XII-2^{AID-HA} was

performed by CUT&Tag profiling (14). Briefly, $2x10^7$ intracellular tachyzoites were

syringe lysed 24 hours post invasion to release fresh parasites. Parasites were

centrifuged at 600 x *g* for 10 minutes at room temperature. Parasite pellets were then

processed as per the manufacturer's protocol (CUT&Tag-IT™ Assay Kit, Anti-Rabbit,

Active motif, Cat no. 53160). Libraries were sequenced by Azenta at PE 2x150 bp.

Chromatin occupancy data analysis

Sequencing data was aligned to the ME49 genome (v52) with bowtie2 (40)

allowing for mate dovetailing. Peaks were called with MACS2 using default settings

(41). Peak annotation was performed with Homer using a 2kb cutoff distance (42). Peak

overlap was conducted with bedtools overlap function (43). AP2XII-2 data was

deposited at GEO GSE217220 and raw data pertaining to ChIPseq of MORC and

HDAC3 (6) were obtained from GEO GSE136060.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. Peter Bradley for the F_1B ATPase antibody, Dr. Vicki

Jeffers for technical advice related to CUT&Tag, and members of the Sullivan lab and

Biology of Intracellular Pathogens group at the IU School of Medicine for helpful

discussions. This research was supported by National Institutes of Health grants

AI124682 (M.W.W.) and AI152583 (W.J.S.). The funders had no role in study design,

data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

1.   Montoya JG, Liesenfeld O. 2004. Toxoplasmosis. Lancet 363:1965-76.

2.   Jeffers V, Tampaki Z, Kim K, Sullivan WJ, Jr. 2018. A latent ability to persist:

differentiation in Toxoplasma gondii. Cell Mol Life Sci 75:2355-2373.

3.   Sullivan WJ, Jr., Hakimi MA. 2006. Histone mediated gene activation in

Toxoplasma gondii. Mol Biochem Parasitol 148:109-16.

4.   Kim K. 2018. The Epigenome, Cell Cycle, and Development in Toxoplasma.

Annu Rev Microbiol 72:479-499.

5.   Harris MT, Jeffers V, Martynowicz J, True JD, Mosley AL, Sullivan WJ, Jr. 2019.

A novel GCN5b lysine acetyltransferase complex associates with distinct

transcription factors in the protozoan parasite Toxoplasma gondii. Mol Biochem

Parasitol 232:111203.

6.   Farhat DC, Swale C, Dard C, Cannella D, Ortet P, Barakat M, Sindikubwabo F,

Belmudes L, De Bock PJ, Couté Y, Bougdour A, Hakimi MA. 2020. A MORC-

driven transcriptional switch controls Toxoplasma developmental trajectories and

sexual commitment. Nat Microbiol 5:570-583.

7.   White MW, Radke JR, Radke JB. 2014. Toxoplasma development - turn the

switch on or off? Cell Microbiol 16:466-72.

8.   Álvarez García G, Davidson R, Jokelainen P, Klevar S, Spano F, Seeber F.

2021. Identification of Oocyst-Driven Toxoplasma gondii Infections in Humans

and Animals through Stage-Specific Serology-Current Status and Future

Perspectives. Microorganisms 9.

9.   Dubey JP. 2021. Outbreaks of clinical toxoplasmosis in humans: five decades of

personal experience, perspectives and lessons learned. Parasit Vectors 14:263.

10.   Attias M, Teixeira DE, Benchimol M, Vommaro RC, Crepaldi PH, De Souza W.

2020. The life-cycle of Toxoplasma gondii reviewed using animations. Parasit

Vectors 13:588.

11.   Zhu S, Shapiro K, VanWormer E. 2022. Dynamics and epidemiology of

Toxoplasma gondii oocyst shedding in domestic and wild felids. Transbound

Emerg Dis 69:2412-2423.

12.   Srivastava S, White MW, Sullivan WJ, Jr. 2020. Toxoplasma gondii AP2XII-2

Contributes to Proper Progression through S-Phase of the Cell Cycle. mSphere

5.

13.   Huang S, Holmes MJ, Radke JB, Hong DP, Liu TK, White MW, Sullivan WJ, Jr.

2017. Toxoplasma gondii AP2IX-4 Regulates Gene Expression during Bradyzoite

Development. mSphere 2.

14.   Kaya-Okur HS, Wu SJ, Codomo CA, Pledger ES, Bryson TD, Henikoff JG,

Ahmad K, Henikoff S. 2019. CUT&Tag for efficient epigenomic profiling of small

samples and single cells. Nat Commun 10:1930.

15.   Radke JB, Lucas O, De Silva EK, Ma Y, Sullivan WJ, Jr., Weiss LM, Llinas M,

White MW. 2013. ApiAP2 transcription factor restricts development of the

Toxoplasma tissue cyst. Proc Natl Acad Sci U S A 110:6871-6.

16.   Campbell TL, De Silva EK, Olszewski KL, Elemento O, Llinás M. 2010.

Identification and genome-wide prediction of DNA binding specificities for the

ApiAP2 family of regulators from the malaria parasite. PLoS Pathog 6:e1001165.

17.   Behnke MS, Wootton JC, Lehmann MM, Radke JB, Lucas O, Nawas J, Sibley

LD, White MW. 2010. Coordinated progression through two subtranscriptomes

underlies the tachyzoite cycle of Toxoplasma gondii. PLoS One 5:e12354.

18.   Harb OS, Roos DS. 2020. ToxoDB: Functional Genomics Resource for

Toxoplasma and Related Organisms. Methods Mol Biol 2071:27-47.

19.   Bougdour A, Maubon D, Baldacci P, Ortet P, Bastien O, Bouillon A, Barale JC,

Pelloux H, Ménard R, Hakimi MA. 2009. Drug inhibition of HDAC3 and epigenetic

control of differentiation in Apicomplexa parasites. J Exp Med 206:953-66.

20.   Dubey JP. 2005. Unexpected oocyst shedding by cats fed Toxoplasma gondii

tachyzoites: in vivo stage conversion and strain variation. Vet Parasitol 133:289-

98.

21.   Ueno A, Dautu G, Munyaka B, Carmen G, Kobayashi Y, Igarashi M. 2009.

Toxoplasma gondii: Identification and characterization of bradyzoite-specific

deoxyribose phosphate aldolase-like gene (TgDPA). Exp Parasitol 121:55-63.

22.   Hong DP, Radke JB, White MW. 2017. Opposing Transcriptional Mechanisms

Regulate Toxoplasma Development. mSphere 2.

23.   Ramakrishnan C, Maier S, Walker RA, Rehrauer H, Joekel DE, Winiger RR,

Basso WU, Grigg ME, Hehl AB, Deplazes P, Smith NC. 2019. An experimental

genetically attenuated live vaccine to prevent transmission of Toxoplasma gondii

by cats. Sci Rep 9:1474.

24.   Fritz HM, Buchholz KR, Chen X, Durbin-Johnson B, Rocke DM, Conrad PA,

Boothroyd JC. 2012. Transcriptomic analysis of toxoplasma development reveals

many novel functions and structures specific to sporozoites and oocysts. PLoS

One 7:e29998.

25.   Wang ZT, Verma SK, Dubey JP, Sibley LD. 2017. The aromatic amino acid

hydroxylase genes AAH1 and AAH2 in Toxoplasma gondii contribute to

transmission in the cat. PLoS Pathog 13:e1006272.

26.   Wang ZT, Harmon S, O'Malley KL, Sibley LD. 2015. Reassessment of the role of

aromatic amino acid hydroxylases and the effect of infection by Toxoplasma

gondii on host dopamine. Infect Immun 83:1039-47.

27. Xia J, Venkat A, Bainbridge RE, Reese ML, Le Roch KG, Ay F, Boyle JP. 2021.

Third-generation sequencing revises the molecular karyotype for Toxoplasma

gondii and identifies emerging copy number variants in sexual recombinants.

Genome Res 31:834-851.

28. Heredero-Bermejo I, Varberg JM, Charvat R, Jacobs K, Garbuz T, Sullivan WJ,

Jr., Arrizabalaga G. 2019. TgDrpC, an atypical dynamin-related protein in

Toxoplasma gondii, is associated with vesicular transport factors and parasite

division. Mol Microbiol 111:46-64.

29. Gaskell EA, Smith JE, Pinney JW, Westhead DR, McConkey GA. 2009. A unique

dual activity amino acid hydroxylase in Toxoplasma gondii. PLoS One 4:e4801.

30. Waldman BS, Schwarz D, Wadsworth MH, 2nd, Saeij JP, Shalek AK, Lourido S.

2020. Identification of a Master Regulator of Differentiation in Toxoplasma. Cell

180:359-372.e16.

31. Xue Y, Theisen TC, Rastogi S, Ferrel A, Quake SR, Boothroyd JC. 2020. A

single-parasite transcriptional atlas of Toxoplasma Gondii reveals novel control of

antigen expression. Elife 9.

32. Goerner AL, Vizcarra EA, Hong DD, Bergersen KV, Alvarez CA, Talavera MA,

Wilson EH, White MW. 2020. An ex vivo model of Toxoplasma

recrudescence. bioRxiv doi:10.1101/2020.05.18.101931:2020.05.18.101931.

33. Hong DD, Brooks K, Goerner AL, Vizcarra EA, Loges LN, Wilson EH, White MW.

2022. Engineering Toxoplasma transgenic tissue cysts. bioRxiv

doi:10.1101/2022.07.21.500998:2022.07.21.500998.

34. Martorelli Di Genova B, Wilson SK, Dubey JP, Knoll LJ. 2019. Intestinal delta-6-
desaturase activity determines host range for Toxoplasma sexual reproduction.
PLoS Biol 17:e3000364.

35. Russell TJ, De Silva EK, Crowley VM, Shaw-Saliba K, Dube N, Josling G, Pasaje
CFA, Kouskoumvekaki I, Panagiotou G, Niles JC, Jacobs-Lorena M, Denise
Okafor C, Gamo FJ, Llinás M. 2022. Inhibitors of ApiAP2 protein DNA binding
exhibit multistage activity against Plasmodium parasites. PLoS Pathog
18:e1010887.

36. Roos DS, Donald RG, Morrissette NS, Moulton AL. 1994. Molecular tools for
genetic dissection of the protozoan parasite Toxoplasma gondii. Methods Cell
Biol 45:27-63.

37. Shen B, Brown KM, Lee TD, Sibley LD. 2014. Efficient gene disruption in diverse
strains of Toxoplasma gondii using CRISPR/CAS9. mBio 5:e01114-14.

38. Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low
memory requirements. Nat Methods 12:357-60.

39. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

40. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat
Methods 9:357-9.

41. Feng J, Liu T, Qin B, Zhang Y, Liu XS. 2012. Identifying ChIP-seq enrichment
using MACS. Nat Protoc 7:1728-40.

42. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C,
Singh H, Glass CK. 2010. Simple combinations of lineage-determining

transcription factors prime cis-regulatory elements required for macrophage and

B cell identities. Mol Cell 38:576-89.

43.   Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing

genomic features. Bioinformatics 26:841-2.

     FIGURE LEGENDS

Figure 1. AP2XII-2 genome occupancy significantly overlaps with HDAC3 and

MORC. A) AP2XII-2 genome occupancy along a segment of chromosome VIIA reveals

its association at the promoters of several genes. Statistically enriched segments are

shown as black bars directly under AP2XII-2 enriched peaks (purple). A similar

distribution and enrichment of HDAC3 and MORC is seen in the orange and teal tracks,

respectively. B) Genome-wide analysis reveals each member of the transcriptional

repression complex associates with the transcriptional start sites (TSS) of genes. C)

Venn diagram indicating that the majority of AP2XII-2 enriched peaks fall within those

identified for HDAC3 and MORC. D) Venn diagram indicating that the majority genes

occupied by AP2XII-2 are also occupied by HDAC3 and MORC.

Figure 2. AP2XII-2 depletion leads to relatively little gene dysregulation. A)

Western blots confirming IAA-dependent downregulation of AP2XII-2^{AID-HA} protein in

RH^{TIR1} or ME49^{TIR1} parasites after 24 hours incubation with 500 µM IAA in culture

media. Blots were probed with anti-SAG1 as a loading control. B-C) MA plot of

differential gene expression analysis of AP2XII-2^{AID-HA}-depleted in RH^{TIR1} (B) or

ME49^{TIR1} (C) parasites 24 hours after IAA addition. Significantly upregulated genes are

colored in blue, downregulated genes in red. Genes of interest for further analysis are

noted. D) Diagrams showing overlap of significantly upregulated and downregulated

genes in ME49 and RH genetic backgrounds. E) Analysis of AP2XII-2 gene occupancy

and directionality of dysregulation that occurs upon AP2XII-2 depletion. F) Meta cell

cycle expression analysis of significantly upregulated (blue) and downregulated (red)

gene sets that respond to AP2XII-2 depletion. The cell cycle profile of AP2XII-2 is

shown in black as a reference.

Figure 3. Disentanglement of the aromatic amino acid hydroxylase family of

genes on chr V. A) Genomic alignment and syntenic analysis of RH88 and ME49

genomes available on ToxoDB.org reveals the correct positioning of unplaced contigs

KE139705 and KE139818 into chr V that was missed due to incorrect fusion of AAH1/2

genes in the ME49 reference strain. Suggested changes to annotation of AAH1 and

AAH2 are based on sequence alignments of 3' flanking regions in both genes across

both parasite strains. B) Multiple pairwise alignments of CDS regions that are present in

each AAH gene fragment reveals a high degree of identity at the nucleotide level which

obfuscates gene calling during differential gene expression analysis. C) Pairwise

alignment of 3' flanking regions reveals a clear way to discriminate both AAH1 and

AAH2 genes for C-terminal tagging at endogenous loci. D) Outline of AAH-family gene

fragments available in ME49 reference strain. Missing fragments on either 5' or 3'

terminus are stated. Conserved CDS regions that were used for multiple sequence

alignment are delimited by dotted line.

Figure 4. Depletion of AP2XII-2 causes de-repression of AAH1 gene expression.

A-B) Western blots demonstrating that AAH1 (A) and not AAH2 (B) is expressed upon

IAA-induced depletion of AP2XII-2. Both genes were endogenously tagged at their C-

termini with MYC epitope tags. Blots were probed with anti-SAG1 as a loading control.

C) AAH1 expression is observed in a concentric pattern around the nucleus upon

AP2XII-2 depletion. D) AAH1 is expressed in the vicinity of the parasite mitochondrial

marker $F_1BATPase$.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Re-analysis of transcriptomic changes that occur upon impairment of

HDAC3/MORC complex. A-B) Differential gene expression analysis for HDAC3

inhibition (A) and IAA-induced MORC depletion (B), each conducted as it was for newly

generated AP2XII-2 depletion data.

Figure S2. Validation for endogenous tagging of AAH1 and AAH2. A) AAH1 was

endogenously tagged with a MYC epitope tag using a CAS9-assisted approach to

facility double homologous recombination as indicated. B) AAH2 was endogenously

tagged with a MYC epitope tag using a single crossover approach as indicated. C-D)

Diagnostic PCRs checking for correct integration (I) or the unmodified endogenous

locus (E) were performed on parental and tagged lines. Specific regions selected for

PCR analysis are highlighted in (A) and (B).

SUPPLEMENTAL TABLE LEGENDS

Table S1. Chromatin-associated regions for AP2XII-2, HDAC3, and MORC as

revealed by CUT&Tag or ChIPseq.

Table S2. Differential gene expression analysis resulting from AP2XII-2 depletion

in RH and ME49 backgrounds.

Table S3. Differential gene expression analysis resulting from HDAC3 inhibition

or MORC depletion.

Table S4. List of primers used in this study.

A

RH88 - long read genome assembly

ME49 Reference Genome - short read assembly missing insert between incorrectly fused AAH2 gene

B

% identity at nucleotide level in aligned CDS

TGME49_287510 (AAH1) ⟩99.6%
99.4% ⟨ TGME49_212710 (F4H)
TGME49_212740 (AAH2) ⟩99.8%

C

% identity at nucleotide level in aligned 3'UTRs

TGME49_287510 (AAH1) ⟩42.2%
TGME49_212740 (AAH2)

D

1kb

TGME49_287510 (AAH1)
5' flank missing
TGME49_212710 (F4H)
3' flank missing
TGME49_212740 (AAH2)
AAH2 full CDS sequence present, UTRs unannotated