Title: The 3-Dimensional Genome Drives the Evolution of Asymmetric Gene Duplicates via Enhancer Capture-Divergence

Authors: UnJin Lee[1,2]*†, Deanna Arsala[1]†, Shengqian Xia[1]†, Cong Li[2], Mujahid Ali[3], Nicolas Svetec[2], Christopher B Langer[2], Débora R. Sobreira[4], Ittai Eres[4], Dylan Sosa[1], Jianhai Chen[1], Li Zhang[5], Patrick Reilly[6], Alexander Guzzetta[7], J.J. Emerson[8], Peter Andolfatto[9], Qi Zhou[3,10], Li Zhao[2] and Manyuan Long[1]*

Affiliations:
[1]Department of Ecology and Evolution, University of Chicago; Chicago, IL, USA
[2]Laboratory of Evolutionary Genetics and Genomics, Rockefeller University; New York, NY, USA
[3]Department of Neuroscience and Developmental Biology, University of Vienna; Vienna, Austria
[4]Department of Human Genetics, University of Chicago; Chicago, IL, USA
[5]Chinese Institute for Brain Research; Beijing, China
[6]Department of Anthropology, Yale University; New Haven, CT, USA
[7]Department of Pathology, University of Chicago; Chicago, IL, USA
[8]Department of Ecology and Evolutionary Biology, University of California, Irvine, CA, USA
[9]Department of Biological Sciences, Columbia University; New York, NY, USA
[10]MOE Laboratory of Biosystems Homeostasis & Protection Life Sciences Institute, Zhejiang University; Hangzhou, Zhejiang, China
*Corresponding authors. Email: ulee@rockefeller.edu, mlong@uchicago.edu
† These authors contributed equally to this work

Abstract: Previous evolutionary models of duplicate gene evolution have overlooked the pivotal role of genome architecture. Here, we show that proximity-based regulatory recruitment of distally duplicated genes (enhancer capture) is an efficient mechanism for modulating tissue-specific production of pre-existing proteins. By leveraging genomic asymmetries in synteny and function that distinguish new genes evolving under enhancer capture-divergence (ECD) from those evolving under previous models, we performed a co-expression analysis on *Drosophila melanogaster* tissue data to show the generality of ECD as a significant evolutionary driver of asymmetric, distally duplicated genes. We use the recently evolved gene *HP6/Umbrea*, which duplicated <15 million years ago (mya), as an example of the ECD process. By assaying genome-wide chromosomal conformations in multiple *Drosophila* species, we show that *HP6/Umbrea* was inserted into a pre-existing, evolutionarily stable 3D genomic structure spanning over 125kb. We then utilize this data to identify a newly discovered enhancer (FLEE1), buried within the coding region of the highly conserved, essential gene *MFS18*, that likely neo-functionalized *HP6/Umbrea*, thereby driving the new duplicate gene copy to fixation. Finally, we demonstrate ancestral transcriptional co-regulation of *HP6/Umbrea*'s future insertion site using single-cell transcriptomics, illustrating how enhancer capture provides a highly evolvable, one-step solution to Ohno's Dilemma. The intuitive molecular mechanism underpinning the ECD model unveils a novel and robust framework to understand the fixation and neofunctionalization of distally duplicated genes.

Main Text:

Newly duplicated genes are at risk of loss in a population through genetic drift or negative selection (*1*) before rare, advantageous neo-functionalizing mutations may occur. The probability of fixation for a slightly deleterious duplicate is more than one to two orders of magnitude lower than a neutral mutant (P_{fix} = 0.085~0.003 x $1/(2N_e)$, Ne ≈ 10^6 as the effective population) for complete and exon duplicates in *Drosophila melanogaster* (*1*, *2*). Consequently, the vast majority of non-fixed duplicate gene copies are likely to be lost in approximately just 2.32 generations or less (**METHODS AND MATERIALS**). Given the low mutation rates in this species (*2*, *3*), it is hardly impossible for a newly fixed duplicate gene to fix or even acquire new mutations, let alone an advantageous one, in such a short time. This problem has also been previously referred to as "Ohno's dilemma" (*4*). Various models have been proposed to resolve this problem including: the duplication, divergence, complementation (DDC)/sub-functionalization model (*5*), the escape from adaptive conflict (EAC) model (*6*), the innovation, amplification, and divergence (IAD) model (*4*, *7*).

The DDC model, also known as subfunctionalization, represents a neutral evolutionary process where symmetric (identical) gene duplicates lose different aspects of their original function due to genetic drift. This random divergence results in the preservation of the duplicated genes, each retaining distinct, yet complementary, functions (Figure 1). Conversely, genes evolving under the EAC model, are under selection for enhanced optimization of specific functions originally held by the parental gene that are partitioned to paralogous copies. The EAC model posits that a single parental gene has intrinsic genetic conflict due to its inability to optimize multiple functions simultaneously, and gene duplication can resolve this evolutionary constraint. While the DDC and EAC models explain how ancestral functions are partitioned among gene duplicates, they fall short in explaining the immediate development of novel expression patterns following gene duplication. These novel expression profiles, often resulting from the gene's new genomic context, can be

instrumental in driving the evolution of new functions—processes not fully captured by the DDC
and EAC models. In contrast, the IAD model describes how shifts in selection pressures can
promote the expression of genes with auxiliary functions by increasing gene copy number (Figure
1). Following the initial increase of auxiliary function through gene amplification, subsequent
relaxation of selection pressure will allow for changes to accumulate on the various copies,
allowing the new copies to diverge and potentially gain a new function (*4*).

While the IAD model provides a reasonable explanation for gene family expansions in
microbial organisms while encountering environmental changes (*7*), the model faces serious
problems when applied to metazoans, as a general and broad increase in gene dosage may be
advantageous in some cell or tissue types but potentially deleterious in others. Similar to the EAC
and DDC models, the IAD model does not directly explain how novel expression patterns arise
immediately following gene duplication, leaving a gap in our understanding of duplicate gene
evolution.

We propose the enhancer capture-divergence (ECD) model, which is an evolutionary model
produced by asymmetric RNA or DNA-based gene duplication processes that allow for distinct
parental and new gene identities and functions (Figure 1). The ECD model first proposes that
selective pressures change for the increased expression of a pre-existing (parental) gene within a
specific tissue or set of tissues. While the evolution of a new enhancer in the parental gene's locus
is plausible, it would require multiple neutral *de novo* substitutions or insertions to generate one
or more necessary transcription factor binding sites that fix within a population and modulates the
expression of the new gene duplicate without disrupting parent gene's expression pattern. Under
the ECD model, duplication of the parental gene into another regulatory environment under the
control of a pre-existing, tissue-specific enhancer is a solution that requires far fewer genomic
changes and can occur in a single step. As the new selection pressures recur, the duplicate copy
that is under new regulatory control will increase in frequency in the population, allowing it to fix.
If the selection pressures change such that the increased tissue-specific expression of the new gene
is no longer advantageous or compensatory mutations appear in the original parent locus, selective
pressures will relax on the new gene copy allowing for divergence. While loss of the new gene
copy by drift or negative selection is one possible fate, if the duplicate gene copy is at high enough
frequency within a population, substitutions may accumulate and result in the gain of new, tissue-
specific protein function.

The previous models addressing Ohno's Dilemma (DDC, EAC, IAD) are symmetric models of
duplication-based evolution which assume that the original parental gene function is randomly
partitioned or entirely retained between identical duplicate copies, making parent and new gene
copies indistinguishable from one another. They offer plausible mechanisms for the retention of
certain types of gene duplicates in various processes of subfunctionalization, conflict resolution
and amplification of ancestral gene functions. The ECD offers a 3D-facilitated efficient route of
neofunctionalization than the development of new enhancers from scratch. It also provides a
coherent and testable framework that describes the evolution of asymmetric or distal gene
duplicates that is currently unexplained by previous models. The asymmetry is a key feature of the
ECD model that distinguishes it from the DDC, EAC and IAC models in different consequences
of functional evolution, and allows for clear identification of genes that evolved under enhancer
capture. (An extended discussion of prior models and genomic symmetry is available in
SUPPLEMENTARY INFORMATION).

91
92

RESULTS

Analysis of Tissue Co-Expression Reveals New Genes Evolve by Enhancer Capture

Central to the IAD model is the observation that gene duplication via unequal crossing over is more likely to occur than a point mutation (*4, 7*). As previously described, one issue with this model is that there is an implicit assumption that during the environmental shift, the increase in fitness gained by over-activity of the auxiliary function must be greater than the decrease in fitness imparted by over-activity of the original gene's function(s). In the case of the enzymatic activity of single-celled organisms where environments are encountered sequentially, it is reasonable to assume that selection might tolerate over-activity of the gene's original function during the transient environment in which the auxiliary function is favored. However, the decrease in fitness for improper expression or activity is larger in multicellular organisms than in single-celled organisms, where a multi-cellular organism's overall phenotype is the cumulative (development) and simultaneous (organ systems) product of many different gene functions.

In the case of multicellular organisms, selection may increase the expression of a gene within a single tissue type (Figure 1). Under the IAD model, a full duplication of the parent gene function and expression pattern drives the duplicate copies to fixation as it provides the most evolvable solution to new conditions. In contrast, under the enhancer capture-divergence model, a copy of the parent gene duplicates into a region of the genome containing an active enhancer(s) that modulates the new gene copy's expression in a tissue-specific manner. Alternatively, the new gene may duplicate into an inactive region of the genome containing unbound transcription factor binding sites, thus activating a previously inert non-coding sequence into a *de novo* enhancer.

Compared to the tissue-specific nature of genes evolving under the ECD model, genes evolving under the IAD model are over-expressed in all tissues, as they are assumed to take on the parent gene expression pattern. We therefore predict that enhancer capture will be more dominant than the IAD model for asymmetrically duplicated genes within multicellular organisms, as it avoids the potentially deleterious effects of increased dosage in multiple tissues resulting from full duplication. However, we stress that the IAD model is likely to drive the evolution of a large number of tandem duplicates as well as a subset of asymmetrically duplicates where the recruitment of pre-existing regulatory elements is unlikely. This increase in fitness caused by the combined output of the new and parental genes thus drives the new gene copy to fixation, providing an alternate resolution to Ohno's Dilemma than the IAD model. Once the tissue-specific selection for the new gene is relaxed, the new gene may then begin to diverge, accumulating substitutions.

Some classes of new genes will continue to evolve under the IAD, DDC, and EAC models. However, the relationships of new genes with their parent genes and neighboring genes differ in expression between those evolving under those previous models and our ECD model, allowing for direct testing of the ECD mechanism as a driver of newly evolved genes. Under the DDC or EAC models, the tissue expression patterns of parental and new genes are complementary, resulting in low co-expression between parental and new gene copies ("parental co-expression"). Since new gene evolution under the DDC and EAC models is assumed to occur in a regulatory-independent context, the tissue expression patterns of the new gene and its neighboring genes should have no relationship, resulting in random co-expression between the new gene and its neighboring gene ("neighboring co-expression"). Under the IAD model, genes and their upstream regulatory sequences are fully duplicated, which predicts a high co-expression between the parent and new gene copies, while the new gene copy and its neighboring genes should have low co-expression.

In the enhancer capture-divergence model, the parent gene is predicted to be more broadly expressed, while the new gene which resides in a distant region of the genome is under the control of one or more tissue-specific enhancers. Here, parental genes are expected to have broad tissue expression patterns, while new genes have expression patterns with high tissue specificity, resulting in low parental co-expression. On the other hand, since the new gene becomes regulated by a locally captured enhancer that is already influencing other genes, neighboring co-expression is high, particularly in gene-dense genomes.

To determine whether the enhancer capture-divergence process is a significant driver of new gene evolution in *D. melanogaster*, we obtained tissue expression data from FlyBase (*8*, *9*)(c.f. Methods and Materials) and calculated co-expression between new/parental and new/neighboring gene pairs (Spearman correlation coefficient) for a random subset of newly evolved genes that 1) underwent a duplication into a different topologically associating domain (TAD) than its parental gene (as defined in (*10*)) and 2) whose essentiality has been validated experimentally ((N=87, Supp. Table S1, Methods and Materials). We focused on experimentally validated genes that were in a different TAD, using distal duplications as a proxy, as their asymmetry allowed us to definitively identify the parent and new gene copies via synteny. To calculate co-expression, we used expression data that contained tissue types extracted from both L3 larvae, pre-pupae, and adult flies, including gut, salivary glands, and imaginal discs from wandering L3 larvae, as well as the head, ovaries, gut, and reproductive organs from adults (c.f. Methods and Materials). For tissues that were represented with multiple experimental runs, data from those tissue types were averaged prior to further analyses to avoid representation bias.

To determine whether a significant number of distally duplicated (non-tandem) genes evolved by enhancer capture, we used two concurrent features of the new genes in our dataset (parent/neighbor tissue co-expression ("PNC") and essentiality) that together determine whether the ECD process is a significant driver of new gene evolution alongside other established models: (Figure 2b). We define "low" and "high" co-expression as being below or above the median co-expression value across new genes in our data set. Under the symmetrical DDC, EAC, and IAD models, the parent and new gene copies are indistinguishable in that all segregable and essential functions of the original gene partition randomly between both parent and duplicate gene copies. In contrast, the ECD model predicts that all original functions, including essential function, are expected to remain with the parental gene while the new copy retains an auxiliary non-essential function. Thus, genes which evolved under enhancer capture are expected to be disproportionately enriched for non-essential functions. Furthermore, as genes under ECD are duplicated into a different regulatory environment from that of their parents, they are expected to appear in the lower right quadrant (quadrant IV) in the PNC plots, with high neighboring co-expression and low parental co-expression (Figure 2b, 2c). In contrast, genes with that have evolved symmetrically via the DDC or EAC models are expected to have a random partitioning of all functions (including essential functions) and should appear in the bottom half of the PNC plots (quadrants III and IV). Specifically, genes evolving under these processes are expected to have low parental co-expression resulting from divergent and complementary expression patterns, while the absence of regulatory context in the DDC and EAC models result in a prediction of random neighboring co-expression, as there is no expected relationship between the new gene and its neighboring genes (Figure 2b, 2c). Similarly, genes that have evolved via the IAD model should also have a random partitioning of essential functions while also appearing in the upper half of the PNC plots (quadrants I and II), with high parental co-expression resulting from full duplication (Figure 2b, 2c).

Alternatively, the enhancer capture-divergence model predicts that most function, including essential gene function, will remain with the parental gene copy, while the tissue-specific expression pattern of the duplicate gene copy serves only to augment the function of the parental gene – a pattern frequently seen in new genes evolving via distal duplication (Figure 2a). Specifically, selection for increased tissue-specific expression of the parental gene predicts the appearance of a distal duplicate of the parental gene copy both with non-essential function and high neighboring co-expression. Meanwhile, the expression pattern and gene function – including all essential function – of the parental copy remains unaltered and is retained. Together, the enhancer capture-divergence model predicts a combination of high neighboring co-expression, low parental co-expression, and non-essentiality. This prediction may be tested by looking for a statistical enrichment of non-essential genes in the lower right quadrant of the PNC plot relative to background (Figure 2b, Supp. Table S1). A distortion in the segregation of essential function is readily identified using the parent/neighbor co-expression plots for distally-duplicated genes in *D. melanogaster*, where the ratio of new essential:new non-essential genes in the lower right quadrant (5:17, 22.7%) was found to be significantly lower than the ratio of remaining new essential:new non-essential genes (29:36, 44.6%) (2.18 fold enrichment, p=0.0294 binomial, p=0.0055, 2D K-S Test based on co-expression data without median thresholding (*11*, *12*)), showing that enhancer capture is a significant driver of new gene evolution alongside previously established processes (Figure 2).

Generality of Enhancer-Capture Divergence

To further address the generality of the enhancer-capture model, we utilized an orthogonally defined, manually curated data set of newly evolved duplicate genes/parental gene pairs in *D. melanogaster* (N=156) which contained information regarding the duplication method of these genes (i.e., tandem, distal, or retro-transposition)(*13*). We also used a separate, publicly available FlyBase data set (*9*) containing 30 classes of developmental tissues produced by the modENCODE consortium (*14*) spanning 0-2 hour embryos to 30-day adults. Using these two data sources, we then calculated the Spearman correlation coefficient for the gene expression each new gene/parent gene pair across all 30 developmental conditions ("developmental co-expression"). Comparison of the developmental co-expression for tandem duplicates vs non-tandem duplicates (i.e., both distal duplicates and retro-transpositions) revealed significantly lower developmental co-expression in non-tandem duplicates than tandem duplicates (p=3.45 x 10^{-10}, Supp. Figure S2). When these comparisons were done with tandem duplicates vs either distal duplicates or retro-transpositions alone, non-tandem duplicates continued to show significantly lower developmental co-expression (distal: p=8.99 x 10^{-9}, retro-transposition: p=5.41 x 10^{-3}, Supp. Figure S2). In support of the generality of the Enhancer Capture-Divergence model, these results demonstrate that regulatory neo-functionalization is a strong driver for non-tandem duplicates, as predicted by the asymmetric ECD model but not the symmetric DDC, EAC, or IAD models. Indeed, similar results have been observed in a wide range of studies, including but not limited to studies of retro-transposons and transposable element domestication as reviewed in (*15*) and (*16*).

HP6/Umbrea as an Illustration of Enhancer Capture-Divergence

Evolution of the *HP6/Umbrea* locus is well-suited for demonstrating the enhancer capture-divergence model, as *HP6/Umbrea* is one of few recently-evolved genes in *D. melanogaster* whose protein evolution has been previously described in the literature (Figure 2b, denoted as (*)) (*17*). *HP1b*, a gene located on the X chromosome, duplicated approximately 12-15 million years

ago (mya) into a gene-poor, intronic region of *dumpy*, located on chromosome 2L (Figure 3). The
new gene, *HP6/Umbrea*, was the result of a full duplication, including its three known domains:
the chromo domain, the chromo-shadow domain, and the hinge domain connecting the two.

Though *HP6/Umbrea* was lost ancestrally to multiple speciation events, suggesting that it was
originally non-essential (*17*), *HP6/Umbrea* continued to evolve in a step-wise manner in the
melanogaster lineage, gaining its semi-lethal phenotype (*18–20*) and rapidly diverging from its
parental gene, *HP1b*. Subsequent to fixation, *HP6/Umbrea* lost its chromo domain approximately
10-12 mya; this was followed by sequence divergence and an accumulation of key substitutions
0-7 mya, resulting in *HP6/Umbrea's* known centromeric protein function in *D. melanogaster* (*17*,
18, *20–22*). The stepwise protein evolution from these results thus eliminates protein neo-
functionalization as the driving force behind the fixation of *HP6/Umbrea*. Sub-functionalization
and/or subsequent optimization of protein function may also be eliminated for similar reasons.

To determine whether enhancer capture is the primary driving force underlying *HP6/Umbrea*'s
fixation, we examined the tissue expression of both *HP6/Umbrea* and *HP1b*. A simple comparison
of *HP6/Umbrea*'s tissue-specific expression pattern to the parental gene *HP1b*'s very broad
expression pattern suggests that *HP1b* is under constitutive regulation (Supp. Figure S1).
Conversely, *HP6/Umbrea* is found only in a subset of tissues which express *HP1b*, suggesting that
the new duplicate is under the control of one or more tissue-specific enhancers. *HP6/Umbrea*'s
expression pattern is not similar to its first neighboring gene, *dumpy*, but its second neighboring
gene, *CR44609*, which expresses in the imaginal discs, larval salivary glands, and male
reproductive organs, which suggests that these genes are likely co-regulated. The non-
complementary nature of the tissue expression patterns of *HP1b* and *HP6/Umbrea* provide further
evidence ruling out sub-functionalization and/or subsequent optimization of regulatory function.

Consistent with the hypothesis that enhancer capture is the driving force being the evolution of
HP6/Umbrea, publicly available modENCODE ChIP-Seq/ChIP-Chip data (*23*) provides positive
evidence that enhancer capture likely drove its early evolution. Using the embryonic S2 cell line
as a negative control where there is little to no *HP6/Umbrea* expression, primed (H3K4me1) and
active (H3K27ac) enhancer marks in whole L3 larvae show strong enhancer activity in an intronic,
gene-poor region of *dumpy*, coinciding with the onset of *HP6/Umbrea* transcription and its co-
regulated, neighboring gene, *CR44609* (Figure 3b). Given the absence of other genes in the region
(Figures 3b, 4a), *HP6/Umbrea* remains the likeliest target of the putative enhancer based on
proximity and expression.

As *HP6/Umbrea* duplicated into a region that appears to be under the control of a pre-existing
enhancer, we tested for further co-regulation in the region by using tissue expression data (c.f. the
section of ***Analysis of Tissue Co-Expression Shows New Genes Evolve by Enhancer Capture***).
We then applied a correlational analysis on this tissue expression data set to determine whether
HP6/Umbrea is co-regulated with other neighboring genes. We took a 500kb region of the genome
centered on the insertion site of *HP6/Umbrea* and calculated the tissue co-expression of each gene
within this region in relation to *HP6/Umbrea*. As enhancers function in a proximity-based manner,
we would expect a distance-dependent effect on the co-expression of neighboring genes across the
genome. To generate a baseline estimate of this distance dependent co-expression distribution, we
sampled 1000 random genic loci within the *D. melanogaster* genome, calculating the degree of
co-regulation expected on proximity alone. Notably, we find that using this distribution, the region
of influence of any given regulatory region of the genome appears to be on the order of 25kb,
suggesting that this is a characteristic distance (1/e reduction) for enhancer interaction in *D.*
melanogaster (Figure 4a). Outside of this region of influence, the likelihood of co-expression

relaxes to the genomic average. Therefore, genes found within this region of influence with high tissue co-expression with neighboring genes are potentially the result of co-regulation with the focal gene. As expected, we find that the neighboring gene, *CR44609*, possesses a similar expression pattern as *HP6/Umbrea*. We also find that a locus of 6 neighboring genes (*CG11929*, *Elba3*, *CG3251*, *Taf12L*, *CG15631*, *CG42523*) located approximately 100kb away from *HP6/Umbrea* also expresses in the same tissues as *HP6/Umbrea*, expressing primarily in the imaginal discs, larval salivary glands, and adult male reproductive organs (Figure 4a).

HP6/Umbrea's Chromosomal Conformations Are Determined by a Larval Enhancer Interaction

While the co-expression of *HP6/Umbrea*'s neighboring gene (*CR44609*) may be explained simply due to its proximity to *HP6/Umbrea*, the co-expression of the 6-gene cluster is not immediately evident as being a result of co-regulation. As the gene cluster of co-expressing genes is distally located along the chromosome beyond *HP6/Umbrea*'s 25kb region of influence, due to the 3-dimensional nature of the eukaryotic genome, these genes may in fact be proximally located near *HP6/Umbrea* in 3D space and thus be co-regulated. Similarly, while active enhancer marks correlating to the onset of expression appear ~50-100kb away from *HP6/Umbrea*, it is not immediately clear that these putative enhancers are driving *HP6/Umbrea* expression, as their distance to *HP6/Umbrea* exceeds the 25kb region of influence. As the 3-dimensional conformations of the genome may still allow these distal genic elements to interact, we tested whether the putative larval enhancer, *HP6/Umbrea*, its neighboring gene (*CR44609*), and the 6-gene cluster are co-regulated by examining high-resolution Hi-C data for *D. melanogaster* (*24*) (Figure 4d).

Like co-expression, the frequency at which two genic elements make physical contact is expected to have a baseline, distance-dependent distribution. We may therefore test for co-regulation by predicting significant physical contact between *HP6/Umbrea*, its putative enhancer, and the cluster of co-expressed neighboring genes using Hi-C data in *D. melanogaster* (Supp. Figure S3). Such an interaction could be detected if contact between these two loci (i.e. *HP6/Umbrea* with enhancer and *HP6/Umbrea* with co-expressing genes) exceeds the baseline distance-dependent distribution of contact frequency. We generated an estimate of this baseline contact frequency distribution using 1000 independent loci that were sampled randomly from the genome, where contact data for the flanking regions were used to generate the baseline distance-dependent contact frequency distribution. We then extracted the contact frequency data for the *HP6/Umbrea* locus alone and compared this to the baseline genome-wide contact frequency distribution (Figure 4d).

We first note that after self-interactions are removed, we find that physical interactions in the genome generally remain highly localized, with most interactions lying near the focal locus as expected. Despite this, we find that *HP6/Umbrea*'s complex contact distribution shows significant contact with two key features: both with the putative larval enhancer as well as the neighboring 6-gene co-expression cluster (Figure 4d). Additionally, when this analysis is repeated for the 6-gene co-expression cluster, we find that this contact is reciprocated, as the 6-gene cluster shows significant contact across the cluster as well as with *HP6/Umbrea* (Figure 4d). Finally, *HP6/Umbrea* has enriched contact with the enhancer region that differentially activates at the onset of *HP6/Umbrea* expression. Combined with the tissue co-expression analysis, these results demonstrate that *HP6/Umbrea* and these 6 genes are likely co-regulated.

The Complex 3D Genome Structure of the HP6/Umbrea Locus Is Conserved Over 25 Million Years, Pre-dating HP6/Umbrea Insertion

While we find evidence that *HP6/Umbrea*, the larval enhancer, and the 6-gene co-expression cluster are co-regulated, it is possible that these interactions evolved after *HP6/ Umbrea*'s insertion. To determine whether these interactions pre-date *HP6/Umbrea*'s insertion, we compared newly generated Hi-C data sets using a second in-group species, *D. yakuba* (Supp. Figure S4), and two out-group species, *D. pseudoobscura* and *D. miranda* (Supp. Figure S5, S6) (Figure 4c, d). While *HP6/Umbrea* inserted 12-15mya, the divergence between *D. melanogaster* and both outgroup species is 25mya (*25*). Within these clades, *D. melanogaster* and *D. yakuba* diverged 6 mya, while *D. pseudoobscura* and *D. miranda* diverged 4 mya. In comparing the Hi-C contact patterns for both *HP6/Umbrea* and its neighboring co-expression cluster, we find that key features of the local chromosomal conformation are conserved in the 3-dimensional structure despite 25 million years of evolution: contact with the larval enhancer, reciprocal contact between *HP6/Umbrea* and its co-expression cluster and contact across the entire co-expression cluster (Figure 4d). These features were found to be conserved in *D. miranda* even in the presence of a large-scale insertion between the future *HP6/Umbrea* locus and the neighboring co-expression cluster. The conservation of this chromosomal structure, despite the subsequent evolution of protein function of *HP6/Umbrea*, suggests that the neo-functionalization event driving the fixation of the original duplication was likely driven by enhancer capture. Specifically, the 3D structure driving enhancer contacts existed prior to *HP6/Umbrea*'s origination, and by duplicating into this region, *HP6/Umbrea* immediately captured this regulatory interaction.

The HP6/Umbrea Locus Structure is Driven by a Tissue-Specific Larval Enhancer

Though the Hi-C data suggested that the captured larval enhancer would be located in the vicinity of chr2L:4500000, the interactions were observed at relatively low genomic resolution. To identify the location of the larval enhancer with higher precision, we utilized 4C-Seq (*26*, *27*) on ~400 dissected L3 larvae and pre-pupae from *D. melanogaster* using a viewpoint of the *HP6/Umbrea* coding sequence, revealing highly enriched contact with a single, distal 394bp locus located approximated 130kb away from the *HP6/Umbrea* locus (Supp. Figure S7). This locus was expanded by approximately 750bp on both 5' and 3' ends and was named the Four-C Larval Enhancer Element (FLEE1)(Supp. File S1). Interestingly, the 2165bp FLEE1 construct was found to be entirely contained within the coding regions of the genes *MFS18* and *Elp3*, which are both highly conserved, essential genes in *D. melanogaster*.

To validate whether FLEE1 contained a functional larval enhancer, we assayed pGreenRabbit reporter plasmids which we site-specifically integrated in *D. melanogaster* (BDSC 79604) (*28*). Compared to control, homozygote transformants that contain empty reporter vectors which drove basal levels of GFP expression, we found that FLEE1 directed GFP expression in the salivary glands of third instar larvae (Figure 5, Movie S1). The result is consistent with prior *in vivo* results of *HP6/Umbrea*'s known expression pattern and key role in polytene chromosome function in larval salivary glands. Anti-body staining of polytene chromosomes showed localization of *HP6/Umbrea* and its parental gene *HP1b*, while tissue-specific RNAi knockdown of *HP6/Umbrea* in the larval salivary glands demonstrated aberrant telomere-telomere attachments (*19*). Consistent with these observations, motif analysis of the FLEE1 locus using FIMO (*29*) and the CIS-BP database (*30*) revealed the presence of a *CrebA* motif, which is a leucine zipper transcription factor associated with regulation of tissue-specific genes in the salivary gland (*9*).

While FLEE1 contains the 3' UTRs of both *MFS18* and *Elp3*, the entirety of FLEE1 is contained within the coding sequences of these two genes, excluding two short intronic regions of 54bp and 66bp within *MFS18* (Supp. Figure S8). Because *MFS18* and *Elp3* are essential genes, we were unable to perform further functional characterization. However, a population genetic analysis of the *MFS18* locus reveals that the coding sequence of *MFS18* is under selective pressure not only to maintain/conserve *MFS18* amino-acid sequence but also to maintain regulatory function as an active larval enhancer. Interestingly, the FLEE1 locus shows strong divergence from *D. yakuba* and *D. simulans* while maintaining low levels of polymorphism within natural populations in *D. melanogaster*, suggesting that the locus is under strong selective pressure (Supp. Figure S9). However, an analysis of the ratio of non-synonymous to synonymous substitution rates from *S. lebanonensis* to *D. melanogaster* for *MFS18* shows that the vast majority of these substitutions are synonymous substitutions (Ka/Ks = 0.033, p=0.0022)(*31*), demonstrating that this locus is under strong purifying selection. Alternatively, the *MFS18* locus fails to show signatures of directional selection, being unable to show significance in the correct direction under the Hudson-Kreitman-Aguadé (HKA)(*32*) and McDonald-Kreitman (MK)(*33*) tests (Table 1). These combined results suggest that the coding sequence of *MFS18* is under selective pressure to maintain/conserve *MFS18* amino-acid sequence while simultaneously maintaining regulatory function as an active larval enhancer, displaying a stereotypically high substitution rate as is common with enhancers under stabilizing selection(*34*). These results stand in sharp contrast to the *HP6/Umbrea* locus, which shows signatures of strong directional selection under both the HKA and MK tests (Table 1).

Co-expression of HP6/Umbrea and Neighboring Genes Pre-dates HP6/Umbrea Insertion

The distal 3D-dimensional contacts (spanning over 250kb from enhancer to co-expression cluster) we identified were found to pre-date *HP6/Umbrea*'s insertion, and these contacts were used to discover a previously uncharacterized enhancer element. However, it is still unclear whether the co-regulation of the co-expressed cluster pre-dates *HP6/Umbrea*'s insertion. The positioning of FLEE1 within a highly conserved, essential gene prevents *in situ* genetic manipulation to validate an ancestral co-regulatory environment. Therefore, to determine whether co-regulation of *HP6/Umbrea*'s insertion site and its neighboring genes pre-dates the insertion of *HP6/Umbrea*, we performed single-cell RNA-Sequencing (scRNA-Seq) using a panel of 3 closely related species: *D. melanogaster* and *D. yakuba*, both containing *HP6/Umbrea*, and *D. ananassae*, which pre-dates *HP6/Umbrea*'s origination. We performed scRNA-seq in the testis tissue because of its high evolutionary importance (*20, 35–37*), the existence of pre-existing high-quality cell type annotations (*38*), and the significantly higher expression levels of *HP6/Umbrea* and its co-expression cluster in this tissue type relative(*38*), and the significantly higher expression levels of HP6/Umbrea and its co-expression cluster compared to imaginal disc or salivary gland tissue (Supp. Figure S1).

After mapping and visualization of the scRNA-Seq data using previous cell type annotations (Figure 6a)(*38*) as well as data from all three species on the same, shared manifold (Figure 6b), it becomes clear that *HP6/Umbrea* is co-regulated on a cellular level with *CG11929*, *Taf12L* and *CG15631*, while overall expression of *Elba3* and *CG3251* are low and restricted mainly to germline stem cells (GSCs)/early spermatagonia. As an internal control, somatic and developmental cell types cluster together as expected. The bulk of the expression is shared across the co-regulated genes, while further cell type-specific expression is also shared within a sub-class of cyst cells, GSCs/early spermatogonia, and early as well as late spermatids. Importantly, the co-

regulation of *CG11929*, *Taf12L*, and *CG15631* pre-dates the insertion of *HP6/Umbrea* as demonstrated by the shared cell type-specific co-expression of these genes in *D. ananassae*. While co-regulation of *CG11929*, *Taf12L*, and *CG15631* and very low expression of *Elba3* and *CG3251* remain evolutionarily conserved, *CG42523* shows significant divergence in its expression pattern within these species. Notably, while its expression pattern shows significant co-regulation with *CG11929*, *Taf12L*, and *CG15631* in *D. ananassae*, it appears that *CG42523* was down-regulated in the *D. yakuba* lineage, while it shows significant functional divergence in its regulation from the co-expression cluster in the *D. melanogaster* lineage.

DISCUSSION

Identification of Distal Larval Enhancer

FLEE1's regulatory activity residing within the primarily exonic regions of the highly-conserved *MFS18* gene constitutes an example of how protein-coding regions of the genome may also have key regulatory functions (*39*, *40*). Such pleiotropy demonstrates how the interpretation of synonymous substitution rates may not necessarily serve as good estimates of neutral evolution rates in commonly used codon table-based tests of molecular evolution. Rather, substitutions typically regarded as synonymous could alternatively be indicative of strong directional or stabilizing selection for the regulatory function of genomic enhancer elements. Furthermore, elucidation of the FLEE1-*HP6/Umbrea* interaction highlights the importance of identifying and characterizing the contributions of structural variations and chromosomal rearrangements in driving phenotypic evolution. Our results demonstrate how stabilizing selection for the conservation of large-scale chromosomal conformations drives the appearance of evolutionary novelty resulting in the development of novel, centromeric function as in the case of *HP6/Umbrea*. While further work will be required to reveal what evolutionary forces underlie the strong conservation of the long-distance interaction of the *HP6/Umbrea* locus prior to the new gene's insertion, our findings demonstrate how complex chromosomal conformations are a key, underappreciated element in the evolution of the eukaryotic genome.

Enhancer Capture Divergence Model

ECD joins various previously proposed models to interpret different evolutionary aspects of gene functionality. Whereas DDC and EAC, for example, explain the duplication-dependent subfunctionalization and resolution of adaptive-caused conflict from ancestral genes with multiple functions respectively, ECD interprets neofunctionalization for creating novel gene functions through duplication. ECD demonstrates how the manner of duplication itself may provide neo-functionalization in an asymmetric, tissue-specific manner. Such neo-functionalization provides a selective advantage in a direct, by the single-step 3D-facilitated acquirement of regulatory elements. The newly acquired elementary functions may maintain the new duplicate for adequate time until the new advantageous mutations occur to solve the Ohno's dilemma.

In addition to partial duplication phenomena such as the generation of gene fusions (*41*) as well as favorable frame-shifts (*42*), our model highlights the under-appreciated evolutionary value of both the act of duplication itself and, more importantly, the genomic context in which these duplications occur. While the role of positional effects in gene regulation and evolution has long been appreciated (*43*, *44*), the advent of new chromosomal conformation capture technologies

allows us to directly connect the conservation of chromosomal domains (*45*, *46*) and the
origination of new genes under a strong conceptual framework.

Under the ECD model, a gene copy duplicates into a pre-existing regulatory context (Figure
7a), gaining a new regulatory interaction. This model thus provides a mechanistic explanation by
which gene interaction networks may rapidly evolve (*36*). Under this model, we have two separate
gene interaction sub-networks for both parental and neighboring genes (Figure 7b). As a new gene
duplicates into a region near the neighboring gene, the new gene acquires the upstream regulatory
function of the neighboring gene as well as the original parental gene's downstream protein
function (Figure 7c) while simultaneously preserving the pre-existing interactions from both
parental and neighboring genes' sub-networks. Since duplication has been observed to occur more
frequently than point mutations (*4*, *7*), enhancer capture provides a faster route to generating
increased tissue-specific expression of a parental gene (Figure 1) than any set of mutations in the
parental gene's regulatory sequence. Duplication in the 3-dimensional looping architecture of the
eukaryotic genome recombines genes and enhancers into new combinations, thus resulting in
regulatory novelty (Figure 7c). As such, this model provides an explanation and mechanism for
the well-described but poorly-understood phenomenon where new gene duplicates often possess
highly tissue-specific expression patterns (Figure 2a)(*37*, *47*, *48*).

One key aspect of the ECD model is the selective advantage imparted by increased tissue-
specific expression. The resolution of genetic conflict, such as sexual antagonism, is becoming
increasingly appreciated as a driver of the evolution of new genes (*49*, *50*). While most new genes
have highly tissue-specific expression patterns, these often favor either the female or male
reproductive organs/germlines in *D. melanogaster* (*37*). A close examination of the expression
pattern of *HP6/Umbrea* demonstrates the same – *HP6/Umbrea* is expressed primarily in the
imaginal discs, larval salivary gland, and the male reproductive organs. As such, it is possible that
the selective advantage imparted by *HP6/Umbrea*'s original duplication may have been a result of
regulatory sexual antagonism and, given that most new genes show expression specific to
reproductive organs, enhancer capture may be a widespread mechanism for the resolution of
sexual antagonism. Furthermore, *HP6/Umbrea*'s repeated ancestral loss suggested it was
originally non-essential follow duplication, but later gained its semi-lethal phenotype in a step-
wise manner. The enhancer capture model may provide a key mechanism by which new essential
genes gain their essential phenotype(*20*).
490

Evidence for the generality of the ECD mechanism
One context in which the mechanism of enhancer capture has been well studied is in the
development of human malignancies. The first discovered example of enhancer capture occurring
in cancer is the t(8;14) translocation in Burkitt's lymphoma, allowing for the oncogene *Myc* to be
expressed under the regulatory control of the immunoglobulin heavy chain gene (*IGH*), which is
expressed in lymphoid cells (*51*, *52*). To date, there have been a variety of other examples of
enhancer capture re-arrangements involved in oncogenesis occurring in diverse tissues (*53–57*).
Although most re-arrangements bring oncogenes into proximity with constitutive regulatory
elements of a given cell type, they may also be brought into proximity with context-specific
regulatory regions. One such translocation in prostate cancer involves the translocation of the
oncogenes *ETV1* or *ERG* within proximity of the promoter region of *TMPRSS2* which contains
several androgen receptor binding sites. In this instance, *ETV1* or *ERG* gains new androgen-
dependent expression, which can be abrogated by androgen deprivation therapy, a common
treatment for prostate cancer(*58*, *59*).

One longstanding question in this literature is why particular re-arrangements are commonly
associated with specific cancers. Are common re-arrangements observed because they are the few
examples that confer a selective advantage in select cell or does the cancer cell type have a
structural predisposition to favor those re-arrangements? Although common rearrangements do
confer a relative fitness advantage to the cancer cells, the evidence has become clear that re-
combining loci are likely to be within physical proximity of one another(*60, 61*). It has been shown
that chromosomes 9 and 22 neighbor each other in hematopoetic cells which may explain the
frequency of the t(9:22) translocation in chronic lymphocytic leukemia which produces the *BCR-*
ABL fusion protein(*62, 63*). Additionally, it has been shown that the *Myc* and *IGH* genes are
brought within close physical proximity during B-cell stimulation (*64*) highlighting the importance
of cell-context specific genomic arrangements in cancer.
The primary difference between enhancer capture in cancer and organismal evolution is the
lack of necessity for cancer cells to preserve an oncogene's previous function via gene duplication
prior to translocation. Additionally, cancer cells typically experience selection at the clonal level
so re-arrangements do not need to confer optimized gene expression within multiple tissue
contexts. However, the cancer literature is clear that enhancer capture is a commonly occurring
one-step mechanism that allows individual cells to gain fitness advantages, and that cell-type
specific 3D genome confirmations selectively favor certain re-arrangements. Given our finding
that the enhancer capture-divergence model is a significant driver of new gene evolution, it is
likely that the inherent 3D configuration of the germline genome imposes a significant and
previously unappreciated constraint on evolutionary novelty.
526
527
528   **METHODS AND MATERIALS**
529
Tissue expression data and analysis
Tissue expression data were retrieved from FlyBase. Pre-computed RPKM data files were
downloaded, with RPKM values for each FlyBase transcript being reported for 29 tissues (*8*). As
many of the tissue-types were repetitive, data from the head, ovary, carcass, and digestive system
were averaged to reduce over-representation bias in further correlational analyses. Gene map data
was also obtained from FlyBase to properly identify neighboring genes (*9*). Parental/new gene pair
information was retrieved from (*20*). Spearman correlation coefficients were calculated using the
tissue expression data between parental and new gene pairs. Due to intronic structures and
variation in gene length, two neighboring genes for each new gene on each side were assessed
using Spearman correlation coefficients and the maximum value of the four neighbors was
recorded. Additionally, correlation coefficients for all genes within 500kb of *HP6/Umbrea* were
reported. To generate a baseline distance-dependent genomic estimate of co-expression, 1000
random genic loci were chosen and co-expression values (Spearman) between the randomly
selected gene and all neighbors within a 500kb range were calculated. This 500kb region was then
divided into 100 non-overlapping windows where means and variances in correlation coefficients
were calculated across all randomly selected loci.
546
547   ***ChIP-Seq data***
548   ChIP-Seq or ChIP-Chip data were obtained for H3K4me1 and H3K27ac for S2 cells as well as
549   whole L3 larvae from modENCODE (*23*). H3K4me1 ChIP-Chip data for S2-DRSC cells were
550   obtained using data ID 304 and 3760. H3K27ac ChIP-Chip data for S2-DRSC cells were obtained

using data ID 296 and 3757. H3K4me1 ChIP-Seq data for whole Oregon-R L3 larvae were obtained using data ID 4986. H3K27ac ChIP-Seq data for whole Oregon-R L3 larvae were obtained using data ID 5084. For all data sets, data was obtained in .gff3 format and visualized using the UCSC Genome Browser.

Hi-C data

We generated Hi-C data of *D. yakuba, D. pseudoobscura,* and *D. miranda*, and used publicly available Hi-C libraries obtained from NCBI for *D. melanogaster*, PRJNA393992. *D. melanogaster* source tissue was S2 cells, *D. yakuba* from adult females (SRR12331759), and *D. pseudoobscura* and *D. miranda* were L3 larvae. Hi-C libraries were preprocessed, mapped, and filtered using HiCUP version 0.8.0 (*65*). Specifically, reads from fastq files were trimmed at ligation junctions, and subsequently each mate of paired-end sequences were independently mapped to the respective genomes using bowtie2 version 2.2.9 (*66*). Reads were mapped to genomes consisting of canonical chromosomes only (i.e., excluding scaffolds and other unplaced sequences). *D. melanogaster* reference genome was dm6 and obtained from FlyBase (*9*). The *D. yakuba* reference genome for the NY73PB line was generated by meta-assembly of two PacBio long read assemblies (FALCON and Canu) using quickmerge, followed by polishing with Quiver, PILON, and a custom FreeBayes homopolymer frameshift polishing step. It can be obtained from NCBI (PRJNA310215). The *D. pseudoobscura* reference genome was obtained directly from Ryan Bracewell (https://www.ryanbracewell.com/data.html) (*67*) and the *D. miranda* reference genome was obtained from NCBI (PRJNA474939), (*68*). HiCUP was used further to remove experimental artifacts based on an *in silico* genome digest as previously described (*65*). HiCUP mapped and filtered .sam files were then converted to formats compatible with HOMER version 4.11 (*69*) and juicer tools version 1.22.01 (*70*). To create matrices, HOMER was used to tile the genome into matrices of fixed-size bins, and assign reads to their correct intersecting bins. HOMER was also used to normalize contact counts in these matrices based on known Hi-C biases, as previously described (*69*). JuicerTools was used to produce .hic files at resolutions of 5kb for *D. melanogaster* and *D. yakuba* and 7.5kb for *D. pseudoobscura* and *D. miranda*, and to create normalized matrices.

Using Hi-C contact matrices, data rows for *HP6/Umbrea* and its neighboring cluster were pulled for a 400kb region centered on *HP6/Umbrea* and self-self interactions were removed. To generate a genome-wide distance-dependent distribution of contact, 1000 random loci were sampled. Contact data for each locus was then normalized with total contact (arb. units) being equal for all loci. The means and variances for each non-overlapping window were calculated and reported and compared to *HP6/Umbrea* and the co-expression clusters' data. To generate genomic coordinates for *HP6/Umbrea* prior to duplication, *D. melanogaster* sequences flanking *HP6/Umbrea*'s insertion site were aligned to the *D. yakuba*, *D. pseudoobscura* and *D. miranda* reference genomes using BLAST. Similarly, the promoter region of CG11929 was aligned to *D. yakuba*, *D. pseudoobscura* and *D. miranda* reference genomes to represent the co-expression cluster.

4C-Seq Data

About 400 *D. melanogaster* L3 larvae and pre-pupae were freshly dissected in 10-minute intervals on ice. A single cell suspension was generated from imaginal disc tissue using collagenase. These suspensions were pooled and formaldehyde-fixed for 10 minutes, followed by glycine quenching. Aliquots of these suspensions were quantified and snap frozen with liquid

nitrogen and stored at -80°C until 10^7 cells were accumulated. All cells were then collected and resuspended in a lysis buffer containing Triton X-100, NP-40, and protease inhibitors followed by homogenization via douncing. Nuclei were then gently lysed using a SDS and Triton-X while shaking (900 RPM) at 37°C for 1 hour each. Restriction enzyme digests were then performed using DpnII. After enzymatic deactivation at 65°C, the resulting solution was diluted in 7 mL of water, and proximity ligation was performed using T4 ligase overnight. This was followed by overnight de-crosslinking using proteinase K. A second restriction enzyme digest was performed with Csp6i followed by a second proximity ligation step performed in 14 mL solution. The resulting circularized library was extracted with ethanol and then purified using a HiPure PCR Cleanup kit. The cleaned library was then amplified using primers specific to HP6/Umbrea with attached Illumina P5/P7 adapters and sequenced on the Illumina HiSeq 2500 platform (PRJNA948431). Results were subsequently aligned to the FlyBase dm6 reference genome, and raw coverage was visualized in R using rtracklayer.

Fly stocks, genetic manipulations and microscopy

All *D. melanogaster* lines were grown on a modified Bloomington cornmeal-molasses formulation. Fly lines for site-specific integration were obtained from Bloomington Drosophila Stock Center. pGreenRabbit reporter plasmids were site-specifically integrated into y[1] w[*] P{y[+t7.7]=nanos-phiC31\int.NLS}X; P{y[+t7.7]=CaryP}attP40 (BDSC 79604). FLEE1 (2L:4444468-4450632) was amplified by PCR and cloned into the pGreenRabbit vector, following traditional cloning methods. We injected an empty pGreenRabbit vector as a negative control and pGreenRabbit with the FLEE1 insert into BDSC 79604 pre-blastoderm embryos. Flies with successful integration were screened for the red eyes phenotype (presence of mini-white). We dissected salivary glands from third instar larvae of homozygous transformants in 1X PBS, fixed in 5% PFA in 1XPBS for 5 minutes, and washed 4 times in 1X PBS for 5 minutes. Fixed salivary glands were stained with DAPI (1:1000) for 10 minutes. All imaging was carried out on an upright laser scanning confocal microscope (Zeiss LSM 710) and similarly processed using ImageJ software.

Population Genetic Analysis

The data analysis. The genomic variants were called from whole genome sequencing of 25 samples of *D. melanogaster* (DRM36, EA87, EA87N, ED10N, EF10N, EF126N, GA01, GA03, GA06, GA07, GH01, GH06, GH12, GH16, GH17, MC23, MC28, RAL900, RG18N, RG4N, UM118, UM37, UM526, ZH16, ZH20), ten samples of *D. simulans* (F11R4, F11R5, F21R2, F21R3, F31R2, F31R3, F31R4, F31R5, F41R1, F41R2), and five samples of D. yakuba (CY02B5, CY08A, CY13A, CY17C, CY22B), with sequencing depths >10(*71*). All these publicly available raw reads were downloaded from NCBI and cleaned with fastp(*72*). The cleaned reads were then mapped to the reference genome of BDGP6.32 with bwa mem v0.7.12(*73*). The variants-calling steps included marking duplicates, recalibrating base quality scores, per-sample calling with HaplotypeCaller, joint-calling with GenotypeGVCFs, and SNPs annotation with snpEff(*71, 74*). Only the biallelic sites with quality score > 30, minimum coverage of 10X, minimum genotype quality of 30, a maximum of 25% missing data were kept.

HKA-like tests (*32, 75*) and MK tests (*33*) were conducted using polarized SNPs by focusing on fixed homologous sites in all outgroup samples (*D. yakuba* and *D. simulans*). The allele frequencies for D. melanogaster and outgroups were estimated with PLINK v1.9(*76*). The expected proportions of diverged and polymorphic sites were calculated using the entirety of

chromosome 2L (547951/ 307551=1.78). The proportions of diverged and polymorphic sites for
genes were compared against the chromosome-wide ones with $\chi 2$ test (d.o.f.=1).

To detect signals of natural selection based on Ka/Ks (also ω) at the loci of *MFS18*, we collected
orthologous sequences of these two genes in 10 Drosophilid species (*D. ananassae, D. erecta, D.*
melanogaster, D. mojavensis, D. pseudoobscura, D. simulans, D. virilis, D. willistoni, D. yakuba,
Scaptodrosophila lebanonensis) from OrthoDB v11(*77*). For *HP6/Umbrea*, Ka/Ks ratio was not
computed due to incomplete ORFs in outgroup species. We used a codon-based alignment
computed with TranslatorX and MAFFT(*78, 79*) for *MFS18* to generate gene trees and conducted
the branch model test implemented by PAML(*80*). To determine the optimal branch model for
substitution rate estimation, we used a dynamic programming method by Zhang et. al. (*31*) to
select the optimal model according to log likelihoods.

The sojourn time of a neutral polymorphic duplicate before loss in a population: The question to
address is how long a newly formed duplicate, if slightly deleterious (as was previously shown for
various polymorphic duplicates (*1*)), can stay in a form of polymorphism in a population before
loss due to genetic drift. The fixation probabilities for various polymorphic duplicates were
calculated using the equation: $u/u_o = S/(1-e^{-S})$ where $u_o = 1/2Ne$ as the fixation probability of a
neutral mutation, $S = 4N_es$ and s the selection coefficient (*81*). The selection component, $\gamma = 2N_es$,
for various polymorphic duplicates in *D. melanogaster* were experimentally measured(*1*). The
average sojourn time before a neutral duplicate mutation disappears from a population was
calculated as $T_0(1/2N) = 2(Ne/N)In(2N)$ where Ne is effective population size and N actual
population size (*82*). The average ratio Ne/N was reported in *D. melanogaster* as 0.027 (*83*) and a
general estimate for metazoans as 0.10(*84*). The $T_0(1/2N) < 1.04 \sim 3.60$ generations (the median
as 2.32 generations) (Ne = 3,300,000 in *D. melanogaster*(*2*)), because all the duplicate variants
are slightly deleterious (*1*) and could disappear even sooner. Furthermore, the point mutation rate,
as reported previously (e.g.(*2, 3*), is in the orders of $10^{-8} \sim 10^{-9}$ per site per generation and the
advantageous ones even much more rare, is unlikely to generate any genetic change that can rescue
the duplicate from extinction in so short a time.

scRNA-Seq
Testes from *D. melanogaster* (*38*), *D. yakuba* (newly generated), and *D. ananassae* (newly
generated) were dissected in drops of cold PBS using forceps on Petri dishes before being
transferred on ice to reduce degradation. We then desheathed testes in lysis buffer (196 µL 1X
TrypLE + 4 µl 100mg/ml collagenase). After spinning down briefly and incubating at room
temperature for 30 minutes with mild vortexing every 10 minutes, the samples were passed
through 35 µm filters before centrifuging for 7 minutes at 163g (1200rpm) at 4°C. We removed
the supernatant, washed the cell pellet with 200 µL cold HBSS, and centrifuged again for 7 minutes
at 163g (1200rpm) at 4°C. We then removed the supernatant before resuspending the cell pellet in
35 µL cold HBSS. We counted cells and checked viability on an automated cell counter using 5
µL of the single cell suspension with 5 µL of trypan blue. Samples were then sent to Rockefeller
Genomics Center for 10X single-cell library preparation and sequencing.
The resulting libraries were processed using cellranger (v7.1.0) and aligned to RefSeq genomes
obtained from NCBI (*mel*: GCF_000001215.4, *yak*: GCF_016746365.2, *ana*: GCF_017639315.1).
Pair-wise alignments were performed for each species' transcriptomes using BLAST and
subsequently imported into SAMAP (v0.3.0) (*85*). Additionally, raw, unprocessed count data from
cellranger was converted into Seurat (v4.3.0) objects in R and then exported as .h5ad AnnData

files. These unprocessed AnnData files were then imported into SAMAP, where expression data
for all three species were mapped to the same manifold and subsequently visualized. Cell type
labels from (*38*) were imported into SAMAP and visualized, allowing for cell type classification.
As overall expression levels of the genes within the visualized cluster varied, resulting in certain
genes' visualizations being saturated and therefore uninterpretable, the scale factor for these genes
was manually adjusted in SAMAP to allow for a consistent interpretation across genes.
695
RNAi and lethality measurements
We used lethality data previously published by our lab (*20, 21*) that was based on RNAi lines
obtained from the Vienna Drosophila Resource Center (VDRC). A quarter of all KK RNAi lines
from VDRC carry an inverted repeat sequence insertion at 30B3. However, a proportion (23–25%)
of KK lines also carry an insertion at 40D3, which is housed within the *tio* locus and produces a
confounding lethal phenotype. To avoid this, we updated the lethality data of new genes reported
in (*20*) by removing the *tio* insertion site in KK lines using a recombination-based approach (*21,
86*) and finally derived lethality data for the new genes. The lethality results for all lines without
insertion in the *tio* locus were reproducible, previously having been analyzed using four replicates,
and again in our analysis in duplicate. Distally duplicated genes had 90% fewer offspring relative
to control flies after Act5c-GAL4 induction were labeled as essential.

References and Notes:

1. J. J. Emerson, M. Cardoso-Moreira, J. O. Borevitz, M. Long, Natural Selection Shapes Genome-Wide Patterns of Copy-Number Polymorphism in Drosophila melanogaster. *Science* **320**, 1629–1631 (2008).

2. M. Kreitman, Nucleotide polymorphism at the alcohol dehydrogenase locus of Drosophila melanogaster. *Nature* **304**, 412–417 (1983).

3. Y. Wang, P. McNeil, R. Abdulazeez, M. Pascual, S. E. Johnston, P. D. Keightley, D. J. Obbard, Variation in mutation, recombination, and transposition rates in Drosophila melanogaster and Drosophila simulans. *Genome Res.* **33**, 587–598 (2023).

4. U. Bergthorsson, D. I. Andersson, J. R. Roth, Ohno's dilemma: Evolution of new genes under continuous selection. *Proc. Natl. Acad. Sci. USA* **104**, 17004–17009 (2007).

5. A. Force, M. F. Lynch, B. Pickett, A. Amores, Y. Yan, J. Postlethwait, Preservation of Duplicate Genes by Complementary, Degenerative Mutations. *Genetics* **151**, 1531–1545 (1999).

6. C. T. Hittinger, S. B. Carroll, Gene duplication and the adaptive evolution of a classic genetic switch. *Nature* **449**, 677–681 (2007).

7. J. Nasvall, L. Sun, J. R. Roth, D. I. Andersson, Real-Time Evolution of New Genes by Innovation, Amplification, and Divergence. *Science* **338**, 384–387 (2012).

8. J. B. Brown, N. Boley, R. Eisman, G. E. May, M. H. Stoiber, M. O. Duff, B. W. Booth, J. Wen, S. Park, A. M. Suzuki, K. H. Wan, C. Yu, D. Zhang, J. W. Carlson, L. Cherbas, B. D. Eads, D. Miller, K. Mockaitis, J. Roberts, C. A. Davis, E. Frise, A. S. Hammonds, S. Olson, S. Shenker, D. Sturgill, A. A. Samsonova, R. Weiszmann, G. Robinson, J. Hernandez, J. Andrews, P. J. Bickel, P. Carninci, P. Cherbas, T. R. Gingeras, R. A. Hoskins, T. C. Kaufman, E. C. Lai, B. Oliver, N. Perrimon, B. R. Graveley, S. E. Celniker, Diversity and dynamics of the Drosophila transcriptome. *Nature* **512**, 393–399 (2014).

9. A. Larkin, S. Marygold, G. Antonazzo, H. Attrill, G. dos Santos, P. Garapati, J. Goodman, L. Gramates, G. Millburn, V. Strelets, C. Tabone, J. Thurmond, T. F. Consortium, FlyBase: updates to the Drosophila melanogaster knowledge base. *Nucleic Acid Research* **49**, D899–D907 (2021).

10. T. Sexton, E. Yaffe, E. Kenigsberg, F. Bantignies, B. Leblanc, M. Hoichman, H. Parrinello, A. Tanay, G. Cavalli, Three-Dimensional Folding and Functional Organization Principles of the Drosophila Genome. *Cell* **148**, 458–472 (2012).

11. G. Fasano, A. Franceschini, A multidimensional version of the Kolmogorov–Smirnov test. *Monthly Notices of the Royal Astronomical Society* **225**, 155–170 (1987).

12. C. Puritz, E. Ness-Cohn, R. Braun, fasano.franceschini.test: An Implementation of a Multidimensional KS Test in R. arXiv arXiv:2106.10539 [Preprint] (2022). https://doi.org/10.48550/arXiv.2106.10539.

13. Q. Zhou, G. Zhang, Y. Zhang, S. Xu, R. Zhao, Z. Zhan, X. Li, Y. Ding, S. Yang, W. Wang, On the origin of new genes in Drosophila. *Genome Res.*, doi: 10.1101/gr.076588.108 (2008).

14. B. R. Graveley, A. N. Brooks, J. W. Carlson, M. O. Duff, J. M. Landolin, L. Yang, C. G. Artieri, M. J. van Baren, N. Boley, B. W. Booth, J. B. Brown, L. Cherbas, C. A. Davis, A. Dobin, R. Li, W. Lin, J. H. Malone, N. R. Mattiuzzo, D. Miller, D. Sturgill, B. B. Tuch, C. Zaleski, D. Zhang, M. Blanchette, S. Dudoit, B. Eads, R. E. Green, A. Hammonds, L. Jiang, P. Kapranov, L. Langton, N. Perrimon, J. E. Sandler, K. H. Wan, A. Willingham, Y. Zhang, Y. Zou, J. Andrews, P. J. Bickel, S. E. Brenner, M. R. Brent, P. Cherbas, T. R. Gingeras, R. A. Hoskins, T. C. Kaufman, B.

Oliver, S. E. Celniker, The developmental transcriptome of Drosophila melanogaster. *Nature* **471**, 473–479
(2011).

15. H. Kaessmann, N. Vinckenbosch, M. Long, RNA-based gene duplication: mechanistic and evolutionary insights.
Nat Rev Genet **10**, 19–31 (2009).

16. D. Jangam, C. Feschotte, E. Betran, Transposable Element Domestication As an Adaptation to Evolutionary
Conflicts. *Trends in Genetics* **33** (2017).

17. B. D. Ross, L. Rosin, A. W. Thomae, M. A. Hiatt, D. Vermaak, A. F. A. de la Cruz, A. Imhof, B. G. Mellone, H. S.
Malik, Stepwise evolution of essential centromere function in a Drosophila neogene. *Science* **240**, 1211–1214
(2013).

18. F. Greil, E. de Wit, H. J. Bussemaker, B. van Steensel, HP1 controls genomic targeting of four novel
heterochromatin proteins in Drosophila. *EMBO Journal* **26** (2007).

19. C. Joppich, S. Scholz, G. Korge, A. Schwendemann, Umbrea, a chromo shadow domain protein in Drosophila
melanogaster heterochromatin, interacts with Hip, HP1 and HOAP. *Chromosome Res* **17**, 19–36 (2009).

20. S. Chen, Y. E. Zhang, M. Long, New genes in Drosophila quickly become essential. *Science* **330**, 1682–1685
(2010).

21. S. Xia, N. W. VanKuren, C. Chen, L. Zhang, C. Kemkemer, Y. Shao, H. Jia, U. Lee, A. S. Advani, A. Gschwent, M.
D. Vibranovski, S. Chen, Y. E. Zhang, M. Long, Genomic analyses of new genes and their phenotypic effects
reveal rapid evolution of essential functions in Drosophila development. *PLoS Genetics* **17**, e1009654 (2021).

22. S. Grill, A. Riley, M. Selvaraj, R. Lehmann, HP6/Umbrea is dispensable for viability and fertility, suggesting
essentiality of newly evolved genes is rare. *Proceedings of the National Academy of Sciences* **120**,
e2309478120 (2023).

23. S. E. Celniker, L. A. L. Dillon, M. B. Gerstein, K. C. Gunsalus, S. Henikoff, G. H. Karpen, M. Kellis, E. C. Lai, J. D.
Lieb, D. M. MacAlpine, G. Micklem, F. Piano, M. Snyder, L. Stein, K. P. White, R. H. Waterston, modENCODE
Consortium, Unlocking the secrets of the genome. *Nature* **459**, 927–930 (2009).

24. Q. Wang, Q. Sun, D. M. Czajkowsky, Z. Shao, Sub-kb Hi-C in D. melanogaster reveals conserved characteristics
of TADs between insect and mammalian cells. *Nature Communications* **9**, 188 (2018).

25. C. A. Russo, N. Takezaki, M. Nei, Molecular phylogeny and divergence times of drosophilid species. *Molecular
Biology and Evolution* **12**, 391–404 (1995).

26. B. Tolhuis, M. Blom, R. M. Kerkhoven, L. Pagie, H. Teunissen, M. Nieuwland, M. Simonis, W. de Laat, M. van
Lohuizen, B. van Steensel, Interactions among Polycomb Domains Are Guided by Chromosome Architecture.
PLOS Genetics **7**, e1001343 (2011).

27. H. J. G. van de Werken, P. J. P. de Vree, E. Splinter, S. J. B. Holwerda, P. Klous, E. de Wit, W. de Laat, 4C
technology: protocols and data analysis. *Methods Enzymol* **513**, 89–112 (2012).

28. B. E. Housden, K. Millen, S. J. Bray, Drosophila Reporter Vectors Compatible with PhiC31 Integrase
Transgenesis Techniques and Their Use to Generate New Notch Reporter Fly Lines. *G3 (Bethesda)* **2**, 79–82
(2012).

29. T. L. Bailey, M. Boden, F. A. Buske, M. Frith, C. E. Grant, L. Clementi, J. Ren, W. W. Li, W. S. Noble, MEME Suite:
tools for motif discovery and searching. *Nucleic Acids Research* **37**, W202–W208 (2009).

30.   M. T. Weirauch, A. Yang, M. Albu, A. G. Cote, A. Montenegro-Montero, P. Drewe, H. S. Najafabadi, S. A.
Lambert, I. Mann, K. Cook, H. Zheng, A. Goity, H. van Bakel, J.-C. Lozano, M. Galli, M. G. Lewsey, E. Huang, T.
Mukherjee, X. Chen, J. S. Reece-Hoyes, S. Govindarajan, G. Shaulsky, A. J. M. Walhout, F.-Y. Bouget, G. Ratsch,
L. F. Larrondo, J. R. Ecker, T. R. Hughes, Determination and Inference of Eukaryotic Transcription Factor
Sequence Specificity. *Cell* **158**, 1431–1443 (2014).

31.   C. Zhang, J. Wang, W. Xie, G. Zhou, M. Long, Q. Zhang, Dynamic programming procedure for searching optimal
models to estimate substitution rates based on the maximum-likelihood method. *Proceedings of the National
Academy of Sciences* **108**, 7860–7865 (2011).

32.   R. R. Hudson, M. Kreitman, M. Aguadé, A Test of Neutral Molecular Evolution Based on Nucleotide Data.
Genetics **116**, 153–159 (1987).

33.   J. H. McDonald, M. Kreitman, Adaptive protein evolution at the Adh locus in Drosophila. *Nature* **351**, 652–654
(1991).

34.   M. Z. Ludwig, C. Bergman, N. H. Patel, M. Kreitman, Evidence for stabilizing selection in a eukaryotic enhancer
element. *Nature* **403**, 564–567 (2000).

35.   M. D. Vibranovski, Y. E. Zhang, C. Kemkemer, H. F. Lopes, T. L. Karr, M. Long, Re-analysis of the larval testis
data on meiotic sex chromosome inactivation revealed evidence for tissue-specific gene expression related to
the drosophila X chromosome. *BMC Biology* **10**, 49 (2012).

36.   W. Zhang, P. Landback, A. R. Gschwend, B. Shen, M. Long, New genes drive the evolution of gene interaction
networks in the human and mouse genomes. *Genome Biology* **202** (2015).

37.   M. Long, N. W. VanKuren, S. Chen, M. D. Vibranovski, New gene evolution: little did we know. *Annual Review
of Genetics* **47**, 307–333 (2013).

38.   E. Witt, S. Benjamin, N. Svetec, L. Zhao, Testis single-cell RNA-seq reveals the dynamics of de novo gene
transcription and germline mutational bias in Drosophila. *eLife* **8**, e47138 (2019).

39.   R. Y. Birnbaum, E. J. Clowney, O. Agamy, M. J. Kim, J. Zhao, T. Yamanaka, Z. Pappalardo, S. L. Clarke, A. M.
Wenger, L. Nguyen, F. Gurrieri, D. B. Everman, C. E. Schwartz, O. S. Birk, G. Bejerano, S. Lomvardas, N. Ahituv,
Coding exons function as tissue-specific enhancers of nearby genes. *Genome Res.* **22**, 1059–1068 (2012).

40.   S. Zhenilo, E. Khrameeva, S. Tsygankova, N. Zhigalova, A. Mazur, E. Prokhortchouk, Individual genome
sequencing identified a novel enhancer element in exon 7 of the CSFR1 gene by shift of expressed allele ratios.
Gene **566**, 223–228 (2015).

41.   W. Wang, J. Zhang, C. Alvarez, A. Llopart, M. Long, The origin of the Jingwei gene and the complex modular
structure of its parental gene, yellow emperor, in Drosophila melanogaster. *Molecular Biology and Evolution*
17, 1294–1301 (2000).

42.   W. Wang, H. Zheng, S. Yang, Y. Haijing, J. Li, H. Jiang, J. Su, L. Yang, J. Zhang, J. McDermott, R. Samudrala, J.
Wang, H. Yang, J. Yun, K. Kristiansen, G. K. S. Wong, J. Wang, Origin and evolution of new exons in rodents.
Genome Research **15**, 1258–1264 (2005).

43.   C. B. Bridges, The Bar "Gene" a Duplication. *Science* **83**, 210–211 (1936).

44.   H. J. Muller, Bar Duplication. *Science* **83**, 528–530 (1936).

45.   N. Harmston, E. Ing-Simmons, G. Tan, M. Perry, M. Markenschlager, B. Lenhard, Topologically associating
domains are ancient features that coincide with Metazoan clusters of extreme noncoding conservation.
Nature Communications **8**, 441 (2017).

46.   J. Krefting, M. A. Andrade-Navarro, J. Ibn-Salen, Evolutionary stability of topologically associating domains is
associated with conserved gene regulation. *BMC Biology*, 87 (2018).

47.   J. Y. Zhang, Q. Zhou, On the Regulatory Evolution of New Genes Throughout Their Life History. *Molecular
Biology and Evolution* **36**, 15–27 (2019).

48.   H. Dai, T. F. Yoshimatsu, M. Long, Retrogene movement within- and between-chromosomes in the evolution
of Drosophila genomes. *Gene* **385**, 96–102 (2006).

49.   L. E. Kursel, H. McConnel, A. F. A. de la Cruz, H. S. Malik, Gametic specialization of centromeric histone paralogs
in Drosophila virilis. *Life Science Alliance* **4**, e202000992 (2021).

50.   N. W. VanKuren, M. Long, Gene Duplicates Resolving Sexual Conflict Rapidly Evolved Essential Gametogenesis
Functions. *Nature Ecology and Evolution* **2**, 705–712 (2018).

51.   L. Zech, U. Haglund, K. Nilsson, G. Klein, Characteristic chromosomal abnormalities in biopsies and lymphoid-
cell lines from patients with burkitt and non-burkitt lymphomas. *International Journal of Cancer* **17**, 47–56
(1976).

52.   V. K. Jain, J. G. Judde, E. E. Max, I. T. Magrath, Variable IgH chain enhancer activity in Burkitt's lymphomas
suggests an additional, direct mechanism of c-myc deregulation. *J Immunol* **150**, 5418–5428 (1993).

53.   B. Gryder, P. C. Scacheri, T. Ried, J. Khan, Chromatin Mechanisms Driving Cancer. *Cold Spring Harb Perspect
Biol* **14**, a040956 (2022).

54.   J. Zheng, Oncogenic chromosomal translocations and human cancer (review). *Oncol Rep* **30**, 2011–2019 (2013).

55.   R. C. Hennessey, K. M. Brown, Cancer regulatory variation. *Curr Opin Genet Dev* **66**, 41–49 (2021).

56.   X. Wang, J. Xu, B. Zhang, Y. Hou, F. Song, H. Lyu, F. Yue, Genome-wide detection of enhancer-hijacking events
from chromatin interaction data in rearranged genomes. *Nat Methods* **18**, 661–668 (2021).

57.   L. E. Montefiori, S. Bendig, Z. Gu, X. Chen, P. Pölönen, X. Ma, A. Murison, A. Zeng, L. Garcia-Prat, K. Dickerson,
I. Iacobucci, S. Abdelhamed, R. Hiltenbrand, P. E. Mead, C. M. Mehr, B. Xu, Z. Cheng, T.-C. Chang, T. Westover,
J. Ma, A. Stengel, S. Kimura, C. Qu, M. B. Valentine, M. Rashkovan, S. Luger, M. R. Litzow, J. M. Rowe, M. L.
den Boer, V. Wang, J. Yin, S. M. Kornblau, S. P. Hunger, M. L. Loh, C.-H. Pui, W. Yang, K. R. Crews, K. G. Roberts,
J. J. Yang, M. V. Relling, W. E. Evans, W. Stock, E. M. Paietta, A. A. Ferrando, J. Zhang, W. Kern, T. Haferlach, G.
Wu, J. E. Dick, J. M. Klco, C. Haferlach, C. G. Mullighan, Enhancer Hijacking Drives Oncogenic BCL11B Expression
in Lineage-Ambiguous Stem Cell Leukemia. *Cancer Discovery* **11**, 2846–2867 (2021).

58.   S. A. Tomlins, D. R. Rhodes, S. Perner, S. M. Dhanasekaran, R. Mehra, X.-W. Sun, S. Varambally, X. Cao, J.
Tchinda, R. Kuefer, C. Lee, J. E. Montie, R. B. Shah, K. J. Pienta, M. A. Rubin, A. M. Chinnaiyan, Recurrent fusion
of TMPRSS2 and ETS transcription factor genes in prostate cancer. *Science* **310**, 644–648 (2005).

59.   M. S. Litwin, H.-J. Tan, The Diagnosis and Treatment of Prostate Cancer: A Review. *JAMA* **317**, 2532–2542
(2017).

60.   K. J. Meaburn, T. Misteli, E. Soutoglou, Spatial genome organization in the formation of chromosomal
translocations. *Semin Cancer Biol* **17**, 80–90 (2007).

61. F. Dubois, N. Sidiropoulos, J. Weischenfeldt, R. Beroukhim, Structural variations in cancer and the 3D genome.
Nat Rev Cancer **22**, 533–546 (2022).

62. H. Neves, C. Ramos, M. G. da Silva, A. Parreira, L. Parreira, The nuclear topography of ABL, BCR, PML, and
RARalpha genes: evidence for gene proximity in specific phases of the cell cycle and stages of hematopoietic
differentiation. *Blood* **93**, 1197–1207 (1999).

63. S. Kozubek, E. Lukásová, A. Marecková, M. Skalníková, M. Kozubek, E. Bártová, V. Kroha, E. Krahulcová, J.
Slotová, The topological organization of chromosomes 9 and 22 in cell nuclei has a determinative role in the
induction of t(9,22) translocations and in the pathogenesis of t(9,22) leukemias. *Chromosoma* **108**, 426–435
(1999).

64. C. S. Osborne, L. Chakalova, J. A. Mitchell, A. Horton, A. L. Wood, D. J. Bolland, A. E. Corcoran, P. Fraser, Myc
dynamically and preferentially relocates to a transcription factory occupied by Igh. *PLoS Biol* **5**, e192 (2007).

65. S. Wingett, P. Ewels, M. Furlan-Magaril, T. Nagano, S. Schoenfelder, P. Fraser, S. Andrews, HiCUP: pipeline for
mapping and processing Hi-C data. *F100Res* **4**, 1310 (2015).

66. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**, 357–359 (2012).

67. R. Bracewell, K. Chatla, M. J. Nalley, B. Bachtrog, Dynamic turnover of centromeres drives karyotype evolution
in Drosophila. *eLife* **Sep 16;8:e49002** (2019).

68. S. Mahajan, K. H.-C. Wie, M. J. Nalley, L. Gibilisco, D. Bachtrog, De novo assembly of a young Drosophila Y
chromosome using single-molecule sequencing and chromatin conformation capture. *PLoS Biology* **16**,
e2006348 (2018).

69. S. Heinz, L. Taxari, M. G. B. Hayes, M. Urbanowski, M. W. Chang, N. Givarkes, A. Rialdi, K. M. White, R. A.
Albrecht, L. Pache, I. Marazzi, A. Garcia-Sastre, M. L. Shaw, C. Benner, Transcription Elongation Can Affect
Genome 3D Structure. *Cell* **174**, 1522–1536 (2018).

70. N. C. Durand, M. S. Shamim, I. Machol, S. S. P. Rao, M. H. Huntley, E. S. Lander, E. L. Aiden, Juicer provides a
one-click system for analyzing loop-resolution Hi-C experiments. *Cell Systems* **3** (2016).

71. J. M. Coughlan, A. J. Dagilis, A. Serrato-Capuchina, H. Elias, D. Peede, K. Isbell, D. M. Castillo, B. S. Cooper, D.
R. Matute, Patterns of Population Structure and Introgression Among Recently Differentiated Drosophila
melanogaster Populations. *Molecular Biology and Evolution* **39**, msac223 (2022).

72. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–
i890 (2018).

73. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**,
1754–1760 (2009).

74. P. Cingolani, A. Platts, L. L. Wang, M. Coon, T. Nguyen, L. Wang, S. J. Land, X. Lu, D. M. Ruden, A program for
annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly* **6**, 80–92 (2012).

75. D. J. Begun, A. K. Holloway, K. Stevens, L. W. Hillier, Y.-P. Poh, M. W. Hahn, P. M. Nista, C. D. Jones, A. D. Kern,
C. N. Dewey, L. Pachter, E. Myers, C. H. Langley, Population Genomics: Whole-Genome Analysis of
Polymorphism and Divergence in Drosophila simulans. *PLOS Biology* **5**, e310 (2007).

76. S. Purcell, B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, D. Bender, J. Maller, P. Sklar, P. I. W. de Bakker,
M. J. Daly, P. C. Sham, PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage
Analyses. *The American Journal of Human Genetics* **81**, 559–575 (2007).

77. D. Kuznetsov, F. Tegenfeldt, M. Manni, M. Seppey, M. Berkeley, E. V. Kriventseva, E. M. Zdobnov, OrthoDB
v11: annotation of orthologs in the widest sampling of organismal diversity. *Nucleic Acids Res* **51**, D445–D451
(2023).

78. F. Abascal, R. Zardoya, M. J. Telford, TranslatorX: multiple alignment of nucleotide sequences guided by amino
acid translations. *Nucleic Acids Res* **38**, W7-13 (2010).

79. K. Katoh, D. M. Standley, MAFFT Multiple Sequence Alignment Software Version 7: Improvements in
Performance and Usability. *Molecular Biology and Evolution* **30**, 772–780 (2013).

80. Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586–1591 (2007).

81. M. Kimura, *The Neutral Theory of Molecular Evolution* (Cambridge University Press, Cambridge, 1983;
https://www.cambridge.org/core/books/neutral-theory-of-molecular-
evolution/0FF60E9F47915B17FFA2620C49400632).

82. D. L. Hartl, A. G. Clark, *Principles of Population Genetics* (Oxford University Press, Oxford, New York, Fourth
Edition, Fourth Edition., 2006).

83. D. A. Briscoe, J. M. Malpica, A. Robertson, G. J. Smith, R. Frankham, R. G. Banks, J. S. F. Barker (1992).
Rapid loss of genetic variation in large captive populations of Drosophila flies: Implications for genetic
management of captive populations. *Conservation Biology* 6, 416-425 (1992).
915
84. R. Frankham, Effective population size/adult population size ratios in wildlife: a review. *Genetics Research* **66**,
95–107 (1995).

85. A. J. Tarashansky, J. M. Musser, M. Khariton, P. Li, D. Arendt, S. R. Quake, B. Wang, Mapping single-cell atlases
throughout Metazoa unravels cell type evolution. *eLife* **10**, e66747 (2021).

86. E. W. Green, G. Fedele, F. Giorgini, C. P. Kyriacou, A Drosophila RNAi collection is subject to dominant
phenotypic effects. *Nat Methods* **11**, 222–223 (2014).

922
923
924

Acknowledgements:
We thank Dr. Marc Halfon and Dr. Sarah Bray for sharing the pGreenRabbit plasmid. We thank
Hong Duan and Connie Zhao at the Genomics Resource Center of Rockefeller University for their
help with the scRNA-seq libraries. This manuscript is dedicated in memory of Frances Lee (F.
LEE).

Funding:
National Institutes of Health grant GM7197-42 (UL)
National Institutes of Health grant 1R01GM116113-01A1 (ML)
National Science Foundation MCB2020667 (ML)
National Institutes of Health postdoctoral fellowship F32GM146423 (DA)
National Institutes of Health grant R01-GM115523 (PA)
National Institutes of Health MIRA R35GM133780 (LZhao)

Author contributions:
Conceptualization: UL, ML
Methodology: UL, ML
Investigation: UL, DA, SX, MA, DRS, IE, DS, JC, PR, NS, CL, CBL, JJE, LZhang
Visualization: UL, DA, PR
Funding acquisition: PA, QZ, LZhao, ML
Project administration: UL, PA, QZ, LZhao, ML
Supervision: PA, QZ, JJE, LZhao, ML
Writing – original draft: UL, DA, AG, ML
Writing – review & editing: UL, DA, PR, AG, NS, LZhao, ML

Competing interests: Authors declare that they have no competing interests.

Data and materials availability:
Publicly available *D. melanogaster* Hi-C libraries form S2 cells: PRJNA393992. Newly
generated *D. yakuba* Hi-C libraries from adult females: SRR12331759. Newly generated *D.
pseudoobscura* and *D. miranda* Hi-C libraries from L3 larvae: PRJNA948678. Newly generated
D. yakuba reference genome: PRJNA310215. Publicly available *D. pseudoobscura* reference
genome: https://www.ryanbracewell.com/data.html. Publicly available *D. miranda* reference
genome: PRJNA474939. Newly generated *D. melanogaster* 4C-Seq data from L3 larve:
PRJNA948431. scRNA-Seq data may: PRJNA995212.

Supplementary Materials
Figs. S1-S9
File S1
Table S1
Movie S1

		HKA				MK			OBSM	
		D	P	p-value		P	D	p-value	Ka/Ks	p-value
MFS18	Obs	31	38	0.018*	N	12	12	0.80	0.033	2.22e-03
	Exp	46	26		S	87	78			
HP6	Obs	50	3	7.5e-4	N	11	0	<<0.001		
	Exp	12	7		S	6	1			

969 **Table 1. Population genetic analysis of enhancer and HP6/Umbrea locus.** The test summaries of
970 HKA, MK, and OBSM for positive selection of *MFS18* and *HP6/Umbrea*. "D, P, N, S, NS, Obs, and
971 Exp" indicate divergence, polymorphism, nonsynonymous sites, synonymous sites, non-significant
972 (p>0.05), observed numbers and proportions, and expected proportions, respectively. *indicates
973 divergence is significantly *lower* than expected, one-sided Fisher's Exact
974
975
976

Figure 1. Comparison of extant models. Presented are illustrations for the (a) Duplication-Divergence-
Complementation (DDC)/Escape-from-Adaptive-Conflict (EAC), the (b) Innovation-Amplification-Divergence
(IAD), and the (c, d) Enhancer Capture-Divergence (ECD) models of duplicate gene evolution, where the gene
regulation of three tissue types are considered. In this scenario, we assume no protein substitutions, so all duplicate
gene copies produce identical proteins. Dotted box represents selection for increased expression, and Δs indicates
the change in selection coefficient. Under the (a) DDC and EAC models, a parental gene duplicates and causes
redundancy in the genome. In the DDC model, redundancy allows for compensation of any single loss-of-function
event, eventually causing the expression pattern of the ancestral gene to be segregated between both new gene
copies in a complementary fashion. Assuming no protein-coding changes, the total output of duplicate gene copies
is identical to the original gene – therefore the DDC model is a neutrally evolving process. In the EAC model,
increased production in a specific tissue cannot occur within a single gene copy due to internal conflict. This
conflict is resolved via the act of duplication, where functions are segregated between duplicate gene copies,
allowing the output of these two genes to increase fitness. Note that the identity of duplicate gene copies may not be
distinguished under the EAC and DDC models (symmetric), resulting in a random segregation of function. Under
the (b) IAD model, an ancestral gene duplicates to increase production of the original protein in a single step. This
increased dosage can potentially cause deleterious effects via misexpression or over-activity in multi-cellular
organisms. Note that the identity of duplicate gene copies also cannot be distinguished in the IAD model
(symmetric), resulting in a redundant segregation of function. Under the (c) ECD model, a parental gene fully
duplicates into a distant region of the genome that is under the control of a pre-existing enhancer. By capturing this
new interaction (d), this duplication increases tissue-specific production of the original protein in a single step.
Notice that the clearly identifiable parental gene copy remains unaltered and thus all original function is retained,
while the duplicate copy acts only to increase protein expression in a single tissue.
1000

c

Model	Parent/New Gene Co-expression	New/Neighboring Gene Co-expression	PNC Plot	Segregation of Essential Function
DDC	low	random	lower half	random
EAC	low	random	lower half	random
IAD	high	random	upper half	random
Enhancer Capture (*)	low	high	lower right	parental gene

Figure 2. Asymmetrically duplicated new genes evolve via enhancer capture. (a) Using new-gene/parent-gene pairs for genes evolving via distal duplication in *D. melanogaster*, the tissue specificity index τ is calculated and plotted above, demonstrating that new genes evolving via distal duplication have higher tissue specificity than parental genes. (b) Shown are parent/neighbor tissue co-expression patterns for new genes in *D. melanogaster* which have duplicated either more than 500kb away or between chromosomes. Tissue co-expression (Spearman correlation coefficient) between new gene/parental gene pairs is plotted on the vertical axis while maximal tissue co-expression between new gene/neighboring genes pairs is plotted on the horizontal axis. Vertical and horizontal lines indicate median co-expression value of all distally duplicated new genes presented here. Genes that evolved via enhancer capture are expected to have low parental co-expression and high neighboring co-expression and should thus be present in the lower right quadrant. (c) While a new gene's essential function is equally likely to be partitioned between either parent or new gene under prior models, new genes evolving via enhancer capture are unlikely to have essential functions, as the expression of the new gene will only augment existing expression of the parental gene, leaving the original essential function intact. Comparing the ratio of new essential to new non-essential genes (29:36, 44.6%) in quadrants I-III to the ratio of new essential to new non-essential genes in quadrant IV showing high neighboring/low parental co-expression (5:17, 22.7%) shows that new genes likely evolve via regulatory capture (2.18 fold enrichment, p=0.0055, 2D K-S test). (* denotes *HP6/Umbrea*)

Figure 3. HP6/Umbrea likely evolved via enhancer capture. (a) *HP6/Umbrea* is a new gene in *D. melanogaster* that arose from a full duplication of *HP1b* into an intronic region of *dumpy*, migrating from chromosome X to 2L. *HP6/Umbrea*'s well characterized, step-wise protein evolution suggests that amino-acid substitutions were unlikely to have driven the duplicate gene copy to fixation. (b) A comparison of ChIP-Seq/ChIP-Chip markers for primed (H3K4me1) and active (H3K27ac) enhancers between embryonic S2 (no/low *HP6/Umbrea* expression) and whole L3 larvae (high *HP6/Umbrea* expression) tracks shows strong activation of a larval enhancer in a 100kb intronic region of *dumpy* that is, aside from *HP6/Umbrea*, devoid of protein coding genes.

Figure 4. *HP6/Umbrea* **co-expression is associated with conserved chromosomal looping that pre-dates its insertion.** (a) Tissue co-expression analysis between *HP6/Umbrea* and neighboring genes reveals the presence of a co-regulated cluster of 6 neighboring genes. Note absence of other genes within *dumpy*'s intronic regions. (b) Two in-group species, *D. melanogaster* and *D. yakuba* (div. ∼ 6mya), contain *HP6/Umbrea*, while two out-group species, *D. pseudoobscura* and *D. miranda* (pse-mir div. ∼ 4mya, pse-mel div. ∼ 25mya), pre-date *HP6/Umbrea*'s insertion (∼ 12-15mya). (c) Cartoon legend illustrating features in (d). Not drawn to scale. (d) Hi-C data tracks for in-group (*D. mel*, *D. yak*) and out-group (*D. pse*, *D. mir*) species are shown for the parental gene *HP1b* (left column) *HP6/Umbrea*'s insertion site (middle column) and the co-regulated 6-gene cluster (right column), with a 95% confidence interval generated from genomic sampling plotted in dotted lines. On the vertical axis is contact in arbitrary units, and on the horizontal axis is genomic coordinates centered on the viewpoint location. Conserved feature (*) shows that *HP6/Umbrea*'s insertion site loops with the active larval enhancers contained in *dumpy*'s intronic gene-desert. Conserved features (†) & (‡) show that *HP6/Umbrea*'s insertion site reciprocally loops with the co-regulated 6-gene cluster. Conserved feature (°) shows that the co-regulated gene cluster loops across the entire 6-gene cluster.

1088
1089 **Figure 5. FLEE1 encodes a larval, salivary gland enhancer.** Green, GFP. White, DAPI (DNA). sg = salivary
1090 gland. fb = fat body. The FLEE1 putative enhancer was tested for enhancer activity in third instar larvae using the
1091 pGreenRabbit reporter vector. (a-c) Basal GFP reporter expression from an empty reporter vector in a third instar
1092 salivary gland and fat body. (d-f) GFP reporter expression directed by FLEE1 in the salivary gland, with minimal
1093 expression in the fat body.
1094

1095
Figure 6. Regulation of *HP6/Umbrea* locus is ancestral. scRNA-Seq data from *D. melanogaster*, *D. yakuba* (both
containing *HP6/Umbrea*) and *D. ananassae* (pre-dating *HP6/Umbrea* origination) are mapped to the same manifold
and visualized. **(a)** Pre-existing data from and their corresponding labels from prior studies were included, allowing
for precise identification of somatic (cyst, hub, and epithelial cells) and developmental (germline stem cell
(GSC)/early spermatogonia, late spermatogonia, early and late spermatocyte, and early and late spermatid) cell type
clusters both within and across species. The developmental trajectory of spermatogenesis is indicated using lines and
arrows. **(b)** Expression data from *HP6/Umbrea* and the co-expression cluster of genes are plotted for all three species.
While the bulk of expression in early spermatids is generally conserved within and across species for *HP6/Umbrea*
(when present), *CG11929*, *Taf12L*, and *CG15631*, finer cell type-specific expression patterns (red arrows) are also
conserved for these genes within each species, indicating conservation of co-regulation pre-dating the insertion of
HP6/Umbrea. Furthermore, low levels of GSC/early spermatogonia-restricted expression of *Elba3* and *CG3251* are
also conserved across species. Interestingly, *CG42523* shares the same cell type-specific expression patterns as
CG11929, *Elba3*, and *CG15631* in *D. ananassae*, but diverges in expression both in *D. yakuba* and *D. melanogaster*.

1109

1110
Figure 7. The 3D organization of the genome allows for rapid rearrangement of genetic networks. Panel (a)
depicts a cartoon illustration of the action of the larval enhancer on the neighboring cluster of co-regulated genes as
well as the future insertion site of *HP6/Umbrea*. Preceding insertion of *HP6/Umbrea*, the larval enhancer was in
contact with both *HP6/Umbrea*'s neighboring gene as well as with the co-regulated 6-gene cluster. (b, c) This looping
structure remains conserved following *HP6/Umbrea*'s insertion, allowing for a rapid recombination of elements
upstream of *HP6/Umbrea*'s neighboring gene (i.e., larval enhancer) with elements downstream of *HP6/Umbrea*'s
parental gene (i.e., *HP1b*'s protein function). A sample gene interaction network, both pre- & post- duplication, is
also depicted above. Note that the parental gene and neighboring gene's original interactions remain intact, preserving
previous function.

SUPPLEMENTARY INFORMATION:

Prior Models and Genomic Symmetries

The first models describing new gene evolution proposed that all new genes likely evolve via duplication-based mechanisms (*1*, *2*), including: the duplication, divergence, complementation (DDC)/sub-functionalization model (*3*), the escape from adaptive conflict (EAC) model (*4*), the innovation, amplification, and divergence (IAD) model. To address how a duplicate, redundant gene copy may rise to fixation, these models all assume multiple functions for any studied gene. In the DDC (sub-functionalization) model, symmetric (identical) copies of a duplicated gene lose function in complementary fashion, resulting in retention of duplicate gene copies with separate but complementary functions. While the DDC model allows each duplicate copy to possess a subset of the parental gene's original functions, the EAC model allows for increased optimization of one or more of the original parental gene's functions that are partitioned to each paralogous copy. The EAC model assumes internal genetic conflict within the parental gene preventing simultaneous optimization of its multiple functions, and duplication thus allows for the resolution of this evolutionary constraint, conferring a selective advantage in both parental and new genes. While the DDC and EAC models can explain how prior gene functions can be partitioned amongst duplicate copies, these models both assume that newly evolved duplicated genes can only retain pre-existing, essential functions from their parental genes and thus fail to describe a mechanism for how truly novel gene function emerges. In contrast, the IAD model proposes that changes in selection pressures may favor the increased expression of a given gene with an auxiliary function. This provides a selective advantage for increased gene dosage through an increase in gene copy number. Following the initial increase of auxiliary function through gene amplification, subsequent relaxation of selection pressure will allow for changes to accumulate on the various copies, allowing the new copies to diverge and potentially gain a new function (*5*). While the IAD model provides a solution for Ohno's dilemma for gene family expansions in microbial organisms while encountering environmental changes (*6*), the model cannot be applied to metazoans due to often conflict effects for same genes in different tissues or cells.

A key factor missing in these previous models is the effect of chromosomal and regulatory context on a gene duplicate's function and spatiotemporal expression. In the DDC, EAC and IAD models, the evolution of new gene duplicates is assumed to occur in a regulatory-independent context and do not describe how the regulatory sequences may shape the evolution of a new gene duplicate. Here, we explain how the regulatory context can promote neofunctionalization of newly duplicated genes through the enhancer-capture divergence (ECD) model. In the ECD model, the duplication of a pre-existing gene into a new regulatory context through a preexisting 3-dimenssional (3D) genome structure results in unique expression pattern from that of its parent gene controlled by a combination of regulatory elements from both the native and new contexts. The single-step evolutionary process of ECD thus allows for rapid neofunctionalization and is dependent on the regulatory architecture of the three-dimensional eukaryotic genome.

Similar to the IAD model, the ECD model first proposes that selective pressures change for the increased expression of a pre-existing (parental) gene within a specific tissue or set of tissues. To achieve this, there are two possible scenarios: 1) the evolution of a new enhancer in the parental gene's locus, either through duplication or substitution, or in the case of the ECD model, 2) the duplication of the parental gene into a distal region of the genome that is already under the control of a pre-existing, tissue-specific enhancer. While the first scenario is possible, this would require multiple neutral *de novo* substitutions or insertions to generate one or more necessary transcription

factor binding sites that fix within a population and modulates the expression of the new gene
duplicate without disrupting parent gene's expression pattern.

In the second scenario under *enhancer capture*, the duplication of the parental gene into another
regulatory environment under the control of a pre-existing, tissue-specific enhancer is a solution
that requires far fewer genomic changes and can occur in a single step. As the new selection
pressures recur, the duplicate copy that is under new regulatory control will increase in frequency
in the population, allowing it to fix. If the selection pressures change such that the increased tissue-
specific expression of the new gene is no longer advantageous or compensatory mutations appear
in the original parent locus, selective pressures will relax on the new gene copy allowing for
divergence. While loss of the new gene copy by drift or negative selection is one possible fate, if
the duplicate gene copy is at high enough frequency within a population, substitutions may
accumulate and result in the gain of new, tissue-specific function.

There are several distinctions between the ECD model and previously classic models of gene
duplication. First, the DDC, EAC and IAD models do not consider the effect of the pre-existing
regulatory and chromosomal environment on a new, distally duplicated gene. Second, compared
to the DDC and EAC models but like IAD, the ECD model is a single-step process in which the
initial duplication event provides a selective advantage. However, unlike IAD, a duplicate gene
copy can immediately integrate into a tissue-specific regulatory network separate from that of its
parent under ECD, providing a fast evolutionary solution to "Ohno's Dilemma." A final and
critical distinction between the ECD model and previous classical models, to address the dilemma,
is that they explain the evolution and retention of different classes of gene duplications. The
previous models are symmetric models of duplication-based evolution which assume that the
original parental gene function is randomly partitioned or entirely retained between identical
duplicate copies, making parent and new gene copies indistinguishable from one another. A similar
genomic symmetry is also seen in tandem duplications, where duplicate copies cannot be
definitively identified as the "parent" or "new" gene copy through synteny. As a result, the DDC,
EAC, and IAD models provide reasonable mechanistic explanations for why a large number of
duplicate gene copies are retained, applying particularly well to tandem and other symmetric gene
duplications. However, these previous models do not consider the role of regulatory and
chromosomal context on newly evolved, asymmetric duplicates and thus cannot explain the
origination of a large number of evolutionarily important genes.

Genes evolving under ECD are asymmetric, as the parental gene remains in its original locus
while the new copy resides in a distal region of the genome under the control of a different, pre-
existing regulatory context. This genomic asymmetry allows for clear distinction between parent
and new gene copies through synteny. A similar asymmetry is also seen in protein and regulatory
function, where the parent gene retains its entire function and spatiotemporal expression pattern,
while the auxiliary tissue-specific function and expression pattern is restricted to the new gene
copy. The asymmetry of both 1) distinguishable gene identity and 2) segregation of expression
and function is a key feature of the ECD model that distinguishes it from the DDC, EAC and IAC
models, and allows for clear identification of genes that evolved under enhancer capture and the
application of genomic tests regarding retention of essential gene function. We utilize these
features of the ECD model to show a statistical enrichment of distally duplicated genes that have
evolved via enhancer capture-divergence within *Drosophila melanogaster*. Under the ECD model,
we predict that newly evolved genes will be enriched for two elements: 1) we predict high degrees
of co-expression with neighboring genes combined with low co-expression with its parent gene
and 2) we predict that genes evolving under ECD should originate as non-essential, as all essential

function should be asymmetrically retained by the parent gene while the auxiliary, non-essential function is retained in the new copy. As the ECD process can occur in a single step in a 3D world of genome, we also predict that the enhancer capture process should be a key mechanism for the evolution of distally duplicated genes alongside the DDC, EAC, and IAD models due to its rapid evolvability. This is supported by the observation that gene duplication occurs more frequently than point mutation (*5*, *6*), where ECD requires fewer genomic alterations than the *de novo* evolution of a new enhancer via substitution.

Under the IAD model, a full duplication of the parent gene function and expression pattern drives the duplicate copies to fixation as it provides the most evolvable solution to new conditions. In contrast, under the enhancer capture-divergence model, a copy of the parent gene duplicates into a region of the genome containing an active enhancer(s) that modulates the new gene copy's expression in a tissue-specific manner. Alternatively, the new gene may duplicate into an inactive region of the genome containing unbound transcription factor binding sites, thus activating a previously inert non-coding sequence into a *de novo* enhancer.

Compared to the tissue-specific nature of genes evolving under the ECD model, genes evolving under the IAD model are over-expressed in all tissues, as they are assumed to take on the parent gene expression pattern. We therefore predict that enhancer capture will be more dominant than the IAD model for asymmetrically duplicated genes within multicellular organisms, as it avoids the potentially deleterious effects of increased dosage in multiple tissues resulting from full duplication. However, we stress that the IAD model is likely to drive the evolution of a large number of tandem duplicates as well as a subset of asymmetrically duplicates where the recruitment of pre-existing regulatory elements is unlikely. This increase in fitness caused by the combined output of the new and parental genes thus drives the new gene copy to fixation, providing an alternate resolution to Ohno's Dilemma than the IAD model. Once the tissue-specific selection for the new gene is relaxed, the new gene may then begin to diverge, accumulating substitutions.

Some classes of new genes will continue to evolve under the IAD, DDC, and EAC models. However, the relationships of new genes with their parent genes and neighboring genes differ in expression between those evolving under those previous models and our ECD model, allowing for direct testing of the ECD mechanism as a driver of newly evolved genes. Under the DDC or EAC models, the tissue expression patterns of parental and new genes are complementary, resulting in low co-expression between parental and new gene copies ("parental co-expression"). Since new gene evolution under the DDC and EAC models is assumed to occur in a regulatory-independent context, the tissue expression patterns of the new gene and its neighboring genes should have no relationship, resulting in random co-expression between the new gene and its neighboring gene ("neighboring co-expression"). Under the IAD model, genes and their upstream regulatory sequences are fully duplicated, which predicts a high co-expression between the parent and new gene copies, while the new gene copy and its neighboring genes should have low co-expression. In the enhancer capture-divergence model, the parent gene is predicted to be more broadly expressed, while the new gene which resides in a distant region of the genome is under the control of one or more tissue-specific enhancers. Here, parental genes are expected to have broad tissue expression patterns, while new genes have expression patterns with high tissue specificity, resulting in low parental co-expression. On the other hand, since the new gene becomes regulated by a locally captured enhancer that is already influencing other genes, neighboring co-expression is high, particularly in gene-dense genomes.

References:

1274
1275
1.   S. Ohno, *Evolution by Gene Duplication* (Springer, New York, 1970).

2.   H. J. Muller, Bar Duplication. *Science*. **83**, 528–530 (1936).

3.   A. Force, M. F. Lynch, B. Pickett, A. Amores, Y. Yan, J. Postlethwait, Preservation of Duplicate Genes by
Complementary, Degenerative Mutations. *Genetics*. **151**, 1531–1545 (1999).

4.   C. T. Hittinger, S. B. Carroll, Gene duplication and the adaptive evolution of a classic genetic switch. *Nature*.
449, 677–681 (2007).

5.   U. Bergthorsson, D. I. Andersson, J. R. Roth, Ohno's dilemma: Evolution of new genes under continuous
selection. *Proc. Natl. Acad. Sci. USA*. **104**, 17004–17009 (2007).

6.   J. Nasvall, L. Sun, J. R. Roth, D. I. Andersson, Real-Time Evolution of New Genes by Innovation,
Amplification, and Divergence. *Science*. **338**, 384–387 (2012).

1286

HP1b

Tissue Expression Level

imaginal disc, larvae L3 wandering
central nervous system, larvae L3
central nervous system, pupae P8
head, virgin 1-day female
head, virgin 4-day female
head, virgin 20-day female
head, mated 1-day female
head, mated 4-day female
head, mated 20-day female
head, mated 1-day male
head, mated 4-day male
head, mated 20-day male
salivary gland, larvae L3 wandering
salivary gland, white prepupae
digestive system, larvae L3 wandering
digestive system, 1-day adult
digestive system, 4-day adult
digestive system, 20-day adult
fat body, larvae L3 wandering
fat body, white prepupae
fat body, pupae P8
carcass, larvae L3 wandering
carcass, 1-day adult
carcass, 4-day adult
carcass, 20-day adult
ovary, virgin 4-day female
ovary, mated 4-day female
testis, mated 4-day male
accessory gland, mated 4-day male

Dumpy

Tissue Expression Level

imaginal disc, larvae L3 wandering
central nervous system, larvae L3
central nervous system, pupae P8
head, virgin 1-day female
head, virgin 4-day female
head, virgin 20-day female
head, mated 1-day female
head, mated 4-day female
head, mated 20-day female
head, mated 1-day male
head, mated 4-day male
head, mated 20-day male
salivary gland, larvae L3 wandering
salivary gland, white prepupae
digestive system, larvae L3 wandering
digestive system, 1-day adult
digestive system, 4-day adult
digestive system, 20-day adult
fat body, larvae L3 wandering
fat body, white prepupae
fat body, pupae P8
carcass, larvae L3 wandering
carcass, 1-day adult
carcass, 4-day adult
carcass, 20-day adult
ovary, virgin 4-day female
ovary, mated 4-day female
testis, mated 4-day male
accessory gland, mated 4-day male

HP6/Umbrea

Tissue Expression Level

imaginal disc, larvae L3 wandering
central nervous system, larvae L3
central nervous system, pupae P8
head, virgin 1-day female
head, virgin 4-day female
head, virgin 20-day female
head, mated 1-day female
head, mated 4-day female
head, mated 20-day female
head, mated 1-day male
head, mated 4-day male
head, mated 20-day male
salivary gland, larvae L3 wandering
salivary gland, white prepupae
digestive system, larvae L3 wandering
digestive system, 1-day adult
digestive system, 4-day adult
digestive system, 20-day adult
fat body, larvae L3 wandering
fat body, white prepupae
fat body, pupae P8
carcass, larvae L3 wandering
carcass, 1-day adult
carcass, 4-day adult
carcass, 20-day adult
ovary, virgin 4-day female
ovary, mated 4-day female
testis, mated 4-day male
accessory gland, mated 4-day male

CR44609

Tissue Expression Level

imaginal disc, larvae L3 wandering
central nervous system, larvae L3
central nervous system, pupae P8
head, virgin 1-day female
head, virgin 4-day female
head, virgin 20-day female
head, mated 1-day female
head, mated 4-day female
head, mated 20-day female
head, mated 1-day male
head, mated 4-day male
head, mated 20-day male
salivary gland, larvae L3 wandering
salivary gland, white prepupae
digestive system, larvae L3 wandering
digestive system, 1-day adult
digestive system, 4-day adult
digestive system, 20-day adult
fat body, larvae L3 wandering
fat body, white prepupae
fat body, pupae P8
carcass, larvae L3 wandering
carcass, 1-day adult
carcass, 4-day adult
carcass, 20-day adult
ovary, virgin 4-day female
ovary, mated 4-day female
testis, mated 4-day male
accessory gland, mated 4-day male

1288

1289 **Figure S1. Expression patterns of *HP6/Umbrea* and other genes.** Unlike the broad expression pattern of parental
1290 gene *HP1b*, the tissue expression pattern of *HP6/Umbrea* is stereotypical of new gene expression patterns, with high
1291 tissue specificity, restricted in this case to primarily the imaginal discs, larval salivary glands, and male reproductive
1292 organs. While *HP6/Umbrea* was inserted into an intronic region of the larger gene *dumpy*, *HP6/Umbrea*'s expression
1293 pattern is shared with *HP6/Umbrea*'s neighboring gene *CR44609*.
1294
1295

Figure S2. Enhancer Capture-Divergence drives regulatory neo-functionalization of new duplicate genes. Parental and new gene co-expression for new duplicate genes arising by tandem, distal duplicates, retro-transposons, and non-tandem (distal + retro-transposons) duplicates were calculated using gene expression data for 30 developmental stages in *D. melanogaster* ("developmental co-expression"). The development co-expression of non-tandem duplicates was significantly lower than the developmental co-expression of tandem duplicates (p=3.45 x 10^{-10}) as well as distal duplicates and retro-transposons alone (distal: p=8.99 x 10^{-9}, retro-transposition: p=5.41 x 10^{-3}). These combined results demonstrate how Enhancer Capture-Divergence is a significant driver of regulatory neo-functionalization in new duplicate genes, which cannot be explained by symmetric models of new duplicate gene evolution.

1307

1308
1309   **Figure S3. Local Hi-C heatmap for *D. melanogaster.*** Shown above is the local chromosomal configuration of
1310   chromosome 2L in the vicinity of *HP6/Umbrea* (chr2L:4570000, center).

1311
1312 **Figure S4. Local Hi-C heatmap for *D. yakuba*.** Shown above is the local chromosomal configuration of
1313 chromosome 2L in the vicinity of *HP6/Umbrea* (chr2L:4680000, center).

1314
1315 **Figure S5. Local Hi-C heatmap for *D. pseudoobscura*.** Shown above is the local chromosomal
1316 configuration of Muller Element B in the vicinity of *HP6/Umbrea*'s future insertion site (Muller
1317 B:29165000, center). Note a large-scale chromosomal inversion event occurred between *D. melanogaster*
1318 and *D. pseudoobscura* (not shown).

1319
1320 **Figure S6. Local Hi-C heatmap for *D. miranda*.** Shown above is the local chromosomal configuration of
1321 chromosome 4 in the vicinity of *HP6/Umbrea*'s future insertion site (chr4:31160000, center).
1322

1323

1324
Figure S7. 4C-Seq in *D. melanogaster*. Shown above are raw read coverage results from 4C-Seq derived from *D.*
melanogaster larval tissue with self-self interactions removed, centered on *HP6/Umbrea*. The strong peak on the left
shows the location of FLEE1.
1328

1329

Figure S8. FLEE1 is located within exonic sequence. The FlyBase gene track and transcript view for FLEE1
shows that it is contained nearly entirely within the coding sequences of *MFS18* and *Elp3* on chromosome 2L.

1330
1331
1332
1333

Figure S9. Polymorphism and divergence in FLEE1/*MFS18* locus. Polymorphism and divergence calculations on chromosome 2L with a 2.5kb window, with the *MFS18* locus highlighted in green, show how the relative number of polymorphisms is low when compared to sequence divergence.

1338   **File S1. Sequence for FLEE1 enhancer.** Sequence for the cloned FLEE1 element in FASTA format.

1339

>FLEE1
ACCAGGGCTTCGGTATGTTGCTGATGGAGGAGGCGGAGCGAATTGCTCGAGAGGAGCACGGCAGCAC
AAAACTGGCGGTCATATCGGGAGTGGGCACCAGAAACTACTATCGCAAAATGGGATACCAACTTGAC
GGACCCTACATGTCAAAGAGCATAGAAGAAATAACTAGGTATAGCGTTAAATGACTGTCTTGGTGGT
ATGTTGGAGGATTAAATATTTGTATTTTATCACGCTAGGAGCAGTAAGATTTCGCTACTTAAAACTACT
CTCTTAAATATATACATTAATATATAGAATGAATCGATTTATTGGCTAAAACTCACAGGGTCCTTTAAA
GTATCAATGATACCACATATTTTTTGGCTTTAACATCTCACAAGAACAACTAAATGATCGTCAATCATA
AACGTGTATACTAAATAATATAAGCAGCATGAACATAAATCGATCCACTCCAATATACCCCACACATA
AATAAATAGGTTAGTTTTTCTGAGGAAAGTGTGCAAGAGAATGTTAAACGATGGCTTCCGCCGAACCA
AAGACTATAAATATGATCCAGCCAACCAAATTGATGCCAGCAGCGGCGCTGAACACCATCGGCCAGC
TTTGTGTGAGCTCCAGAATGTGTCCGGCCAAGTATACTCCGAGAAAGCCAGGAATCGCGCCCACTGTG
TTCATCAGGCCAAAGACGCTGCCCGAATGCAGAGGTGCCAGGTCTTGGGGATTCACTGTTACCGCGTT
GTTGTGGAAGCCCGTGCCGCCAATGATAATGGTCATGCAGATGAGCGCCGTATGGAAGTCCGAGGTG
CGGCTCATCACAAACAGGGCCAGATTCTGAGCGGCAAAGCAGCAACTTTGGATGACCTTGCGCACCGT
CGTCGTGTGCCATTCGCGAGCGAGTAATCTGGTGGTCAAGTACTTGGCGAATAGCGTGCACGGTGGCA
GGGCAAGCCACGGGATCATGTTCACTACCCAACCCTTGGCGTGTGGAAAGCCGTCGTGGAAGTATGTA
GGCAGCCAGGAGAGTAGCACGAAGAAGCAGTTCATCTCGCAGGCGTGAGTCAGCACACAGGCCCAGA
AGGACAGCCTACGAAAGTATCGCAACCAAGGCACGGCTGACGTCTCTGCCGGACTCTTGTTCGCGCAC
AGTCGGGATGGCGTGGCAATATTAATGATTCGGTTTCGCTCGCCGGCCATTGCATAGTAGCGCAGCAC
CAGCGCCCATGCGATGCCCATCAGTCCTATCACCCGGAATACATACGACCAGCCGAAGTAGTCCAGCA
GAAAAGATCCCATAATCCCAGTCAGAAGAGTACCTAGAGCCGATCCCGCTGTGAGCAGCCCAAAGAA
GCTGCTTCTCTCATTGGGGCACAAATTCTGCAAACGATTAAGTTATAGTTTATGTGTAAATTTATAAAA
TTAGCTAAGCACCTGACTGGTTAGACTAATCATGCTAGGAAAGTGCACGCCCTGAAGAGCGCCGTTCA
GGATTCGAATGGCAACAATGAAGGGAATAGCGTAACTCTTGATGGAGCCCGCCGTCCAGATGATAGT
GGGCATTAGGAATGTGATAAGCGACCAGCCGATTGCGGCAAACAGAATGACTCGCTGGCCTCCAAAG
CGGTCGCTGAAGTAGCCGCCCACAACCTGCGTGAGTGTGTAGCCCCAGAAGAAGGAGCTGAGCACAG
TGCCCGAGTCGGTTTTGCTCCACTTTTGGGCGGATGCCACGGCCGGCACAAGAAGTGGCATAGTGGTG
CGGGTGGAGTACAGCATACAGGTGCCCGTAATAAGGGTGATGAACCAGACACGCTTCTCATGCCTGC
AATGATCCGACAAAGGAGTTGTACTTGGGGAGGTTTAGTGAGCTGTATGCTGTGAGACCCACCTGGTC
CAAATGCTCTGCGTGTCCACCAGTTCCCCGCGCAGCAGAGAATATTTTAGCTTCTCGTCCATGGTCACA
AACTGGGTCCGGAACTATTGCCTTTCCTTCACGTCACATATCAACTCCAACTGCTTCGTTGCTTGCCGC
TGTGGCATATTTTACTGCCCTTTGTTTACTTTCATTCACGTTGGCGACTAGACACGCCAAGTATTTGCG
CCTGTTAAAATTATGTTTTTACGTGGCCGTTTTTCCAACAGCCGCTGGACTAGAGCATAG

1373

Table S1. Co-expression and essentiality data for newly evolved distal duplicates. Data for each new gene/parent gene pair with parental coexpression, maximal neighboring coexpression, and essentiality data reported.

new_gene	par_gene	neigh_cor	par_cor	essentiality
CG31875	CG33525	0.544016	0.418772	essential
CG33458	CG30090	0.510544	0.061535	essential
CG33459	CG30090	0.510544	0.170643	essential
CG5372	CG8095	0.533216	0.100566	essential
CG3347	CG17440	0.690522	0.917109	essential
CG31313	CG8066	0.262423	0.262423	essential
CG17802	CG17806	0.754936	0.754936	essential
CG30395	CG15040	0.564165	0.503847	essential
CG8664	CG5107	0.739614	-0.02909	essential
CG4477	CG18223	0.991723	0.87925	essential
CG4907	CG13978	0.473646	0.90807	essential
CG31509	CG31508	0.568805	0.351047	essential
CG33109	CG16826	0.863108	0.855139	essential
CG5609	CG31508	0.568805	0.025656	essential
CG4259	CG3117	0.310205	0.370675	essential
CG6289	CG6663	0.896926	0.896926	essential
CG1736	CG9327	0.476144	0.31898	essential
CG2826	CG13686	0.611623	0.840596	essential
CG10090	CG14666	0.581515	0.72728	essential
CG11597	CG32505	0.294694	0.434793	essential
CG9720	CG31524	1	0.749838	essential
CG9873	CG9091	0.73774	-0.00342	essential
CG15358	CG15818	0.285515	-0.17483	essential
CG11466	CG4486	0.755441	0.854113	essential
CG8358	CG5527	0.195031	0.546997	essential
CG31791	CG31801	0.858564	0.726142	essential
CG32588	CG33252	0.770768	0.809965	essential
CG15527	CG2998	0.546119	0.011347	essential
CG13686	CG2839	0.727607	0.652654	essential
CG17011	CG17799	0.473299	0.954925	essential
CG6687	CG18525	-0.11768	-0.11768	essential
CG17012	CG30031	0.999027	-0.12837	essential
CG30036	CG33145	0.995132	0.12326	essential
CG30037	CG33145	0.995132	0.083991	essential
CG18125	CG15040	-0.12326	-0.30413	non_essential
CG32368	CG18754	0.049468	-0.02339	non_essential
CG16992	CG16761	-0.05143	-0.22744	non_essential
CG12224	CG6392	0.876585	-0.40969	non_essential
CG4580	CG7052	0.23011	0.210476	non_essential
CG17650	CG17945	0.937531	0.744143	non_essential

CG17637	CG7754	0.769495	0.589164	non_essential
CG13463	CG33489	0.63252	0.798583	non_essential
CG4712	CG17734	0.852126	-0.14781	non_essential
CG7815	CG6663	0.795447	0.326526	non_essential
CG33490	CG8622	0.774678	0.288208	non_essential
CG3640	CG33101	0.622135	0.3315	non_essential
CG10799	CG5784	0.639334	-0.58441	non_essential
CG5509	CG11958	0.546399	0.268802	non_essential
CG33920	CG9091	0.512647	0.116871	non_essential
CG8626	CG8095	0.958655	0.224197	non_essential
CG17673	CG33104	0.947211	0.191063	non_essential
CG10232	CG14213	0.122907	0.160009	non_essential
CG18748	CG17799	0.914127	0.711878	non_essential
CG6036	CG5724	0.943401	0.162411	non_essential
CG3217	CG4842	0.753369	-0.22816	non_essential
CG12842	CG18249	0.722673	-0.17571	non_essential
CG31769	CG4486	0.906835	0.57304	non_essential
CG5265	CG15503	0.486472	0.194038	non_essential
CG31918	CG7599	0.726024	-0.07309	non_essential
CG31932	CG6912	0.705203	-0.30682	non_essential
CG13091	CG17843	0.17007	0.685303	non_essential
CG31370	CG17843	0.772457	-0.24621	non_essential
CG7594	CG32107	0.999657	-0.00293	non_essential
CG13656	CG15293	0.64774	0.569462	non_essential
CG6639	CG32284	0.826954	-0.04802	non_essential
CG17268	CG1844	0.544297	0.05546	non_essential
CG33235	CG11598	0.349626	0.556675	non_essential
CG10700	CG10863	0.560563	0.022261	non_essential
CG6208	CG33252	0.250033	0.129214	non_essential
CG14610	CG33145	0.998236	0.043089	non_essential
CG31508	CG30083	0.531385	0.346498	non_essential
CG30450	CG1101	0.585754	-0.32546	non_essential
CG31524	CG33525	1	0.198622	non_essential
CG1840	CG5983	0.447979	0.301494	non_essential
CG13977	CG30362	0.680651	0.590968	non_essential
CG6690	CG1906	0.523537	0.023827	non_essential
CG30473	CG1718	0.760268	0.361607	non_essential
CG8856	CG8036	0.652693	0.330431	non_essential
CG17174	CG3422	0.909234	0.14695	non_essential
CG17176	CG4199	0.923376	0.114519	non_essential
CG12493	CG31508	0.642318	-0.1908	non_essential
CG11833	CG31013	0.057125	0.015086	non_essential
CG7931	CG12359	0.960604	-0.19223	non_essential

CG30494	CG8588	0.818025	0.676018	non_essential
CG15461	CG5302	0.045446	0.361934	non_essential
CG33462	CG1942	0.762024	-0.09881	non_essential
CG15636	CG7041	0.818182	0.087641	non_essential

1377

1378   **Movie S1. Live GFP expression.** Video of live larvae with FLEE1 under the control of an enhancer-reporter

1379   vector. Expression is seen restricted to the larval salivary glands. (MovieS1.mp4)