Antifungal tolerance and resistance emerge at distinct drug concentrations and rely upon different aneuploid chromosomes

Feng Yang[a,b], Eduardo FC Scopel[b], Hao Li[a,c], Liu-liu Sun[a,c], Nora Kawar[b], Yong-bing Cao[d], Yuan-Ying Jiang[a,#], Judith Berman[b,#]

[a]Department of Pharmacy, Shanghai Tenth People's Hospital, School of Medicine, Tongji University, Shanghai, China
[b]Shmunis School of Biomedical and Cancer Research, The George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel
[c]Department of Physiology and Pharmacology, School of Basic Medicine and Clinical Pharmacy, China Pharmaceutical University, Nanjing, China
[d]Department of Vascular Disease, Shanghai TCM-Integrated Hospital, Shanghai University of Traditional Chinese Medicine, Shanghai, China

[#]Address correspondence to jberman@tauex.tau.ac.il (J.B), and jiangyy@tongji.edu.cn (Y-y. J)

Abstract

Antifungal drug tolerance is a response distinct from resistance, in which cells grow slowly above the minimum inhibitory drug concentration (MIC). Here we found that the majority (69.2%) of 133 *Candida albicans* clinical isolates, including standard lab strain SC5314, exhibited *temperature-enhanced* tolerance at 37°C and 39°C, and were not tolerant at 30°C. Other isolates were either *always* tolerant (23.3%) or *never* tolerant (7.5%) at these three temperatures, suggesting that tolerance requires different physiological processes in different isolates. At supra-MIC fluconazole concentrations (8-128 µg/ml), tolerant colonies emerged rapidly at a frequency of ~10^{-3}. In liquid passages over a broader range of fluconazole concentrations (0.25-128 µg/ml), tolerance emerged rapidly (within one passage) at supra-MIC concentrations. By contrast, resistance appeared at sub-MIC concentrations after 5 or more passages. Of 155 adaptors that evolved higher tolerance, all carried one of several recurrent aneuploid chromosomes, often including chromosome R, alone or in combination with other chromosomes. Furthermore, loss of these recurrent aneuploidies was associated with a loss of acquired tolerance, indicating that specific aneuploidies confer fluconazole tolerance. Thus, genetic background and physiology, and the degree of drug stress (above or below the MIC) influence the evolutionary trajectories and dynamics with which antifungal drug resistance or tolerance emerges.

Importance

Antifungal drug tolerance differs from drug resistance: tolerant cells grow slowly in drug, while resistant cells usually grow well, due to mutations in a few known genes. More than half of *Candida albicans* clinical isolates have higher tolerance at body temperature than they do at the lower temperatures used for most lab experiments. This implies that different isolates achieve drug tolerance via several cellular processes. When we evolved different strains at a range of high drug concentrations above inhibitory levels, tolerance emerged rapidly and at high frequency (one in 1000 cells) while resistance only appeared later at very low drug concentrations. An extra copy of all or part of chromosome R was associated with tolerance, while point mutations or different aneuploidies were seen with resistance. Thus, genetic background and physiology, temperature, and drug concentration all influence how drug tolerance or resistance evolves.

Introduction

More than 1.5 million people die from invasive fungal infections every year (1). Increases in the global prevalence of fungal infections has become a major public health concern (2). This is largely because the at-risk population is expanding with the increase in patients with compromised immunity, who are especially vulnerable to fungal infection and the overall increase in lifespan in general. For the vast majority of fungal infections, high morbidity and mortality are caused by species from the genera *Candida*, *Aspergillus* and *Cryptococcus* species (1).

Only three antifungal drug classes are used clinically for monotherapy: the polyenes, azoles and echinocandins (3). Polyenes, such as amphotericin B, have potent and broad-spectrum antifungal activity and resistance is rarely seen. However, polyenes can cause severe toxicity in the kidneys and the liver (4), because they also bind to human cholesterol. Echinocandins are fungicidal against most *Candida* species and are first-line drugs for the treatment of candidemia (5). However, the relatively high cost of echinocandins and the need to administer them intravenously, makes them inappropriate in many settings. Azoles, such as fluconazole (FLC), inhibit 14α-lanosterol demethylase, a key enzyme in ergosterol biosynthesis. Azoles, which are fungistatic, have broad-spectrum antifungal activity, good safety profiles, relatively high bioavailability and are more affordable in many healthcare settings. Therefore, azoles are one of the most widely used antifungal drugs (6, 7).

The clinical prevalence of FLC resistance in *C. albicans,* a prevalent opportunistic human fungal pathogen, is generally less than 1% (8). Despite this, therapeutic failure of FLC against susceptible *C. albicans* isolates is often >30% in systemic infections (9). The fungistatic nature of azoles allows cells to survive and to evolve new traits, such as drug tolerance or resistance.

Since the introduction of first-generation azole drugs in the 1990s, most published research on drug responses did not distinguish between resistance and tolerance.

Antifungal drug tolerance, which is distinct from resistance, has been best characterized

in *C. albicans* cells responding to FLC ((10) and reviewed in (11)). Antifungal drug

resistance is the ability to grow well at drug concentrations above a defined MIC for the

drug. Antifungal drug tolerance is defined as a characteristic of drug-susceptible

genotypes that can grow, albeit slowly, at inhibitory drug concentrations (12).

Furthermore, in a tolerant isolate, not all cells in the isogenic population grow with

similar dynamics. Furthermore, individual clinical isolates exhibit different levels of FLC

tolerance, and the degree of tolerance changes as a function of growth conditions (9,

10). However, the degree to which tolerance in different isolates varies across

environmental conditions remains to be characterized.

Mechanistically, antifungal resistance is usually due to genetic/genomic mutations that

directly affect the drug-target interaction and these mutations affect the response of all

cells in a population. Antifungal drug tolerance depends upon diverse stress pathway

responses, including heat-shock responses, responses to amino acid starvation,

kinases such as protein kinase C, and epigenetic processes (11). Stress pathways

presumably enable the slow growth of some cells, and likely affect drug-target

interactions indirectly. However, the mechanisms that affect tolerance and that

distinguish tolerance and resistance are not well understood. Because the persistence

and mortality of fungal infections is associated with high levels of tolerance (9, 10), we

posit that antifungal tolerance, a property often overlooked in clinical assays, may

explain at least some of the treatment failures caused by clinical isolates that test as

susceptible. We also hypothesize that understanding how genetic and physiological

processes that modulate tolerance should identify new strategies to improve the

outcomes of antifungal drug therapies.

In this study, we first screened a collection of 133 clinical isolates to determine the

prevalence of FLC tolerance under in vitro lab conditions at normal and at febrile body

temperatures. We found three distinct types of temperature responses: 1) *temperature-*

elevated tolerance (TET) at 37 and 39°C, relative to 30°C; 2) *always-tolerant* (AT) and

never-tolerant (NT) isolates at the three temperatures. We then evolved representative

TET and AT isolates for adaptation to FLC at a range of drug concentrations. We found

that aneuploidy appears rapidly and is associated with increased tolerance, especially

at drug concentrations above the MIC. By contrast, resistant isolates emerged at sub-

MIC drug concentrations, only after ~5 days of passaging in sub-MIC drug

concentrations and they proceeded to acquire higher resistance levels with time. Some

of the resistant adaptors also acquired aneuploidies, but different ones from those in

tolerant adaptors. This suggests that tolerance and resistance appear with distinct

evolutionary dynamics and trajectories.

Results

Most clinical isolates tested were fluconazole-tolerant at physiologically relevant temperatures

Previously, we found that temperature and medium composition each affected antifungal tolerance to FLC and ketoconazole (10, 13). Therefore, we asked about how these two physiological factors affected one another. Interestingly, the effect of temperature on FLC tolerance was modulated, sometimes in opposite ways, by medium composition. For example, on RPMI-1640 and casitone plates, SC5314 was tolerant at *both* 30°C and 37°C, yet on chromagar, SD or SDC plates, the same strain was not tolerant at *either* 30°C or 37 °C. Interestingly, on YPD medium (a nutrient-rich medium), SC5314 had *temperature-enhanced* tolerance: it was *non-tolerant* at 30°C and *tolerant* at 37°C (Fig. S1). Thus, FLC tolerance is regulated by the interplay between temperature and medium composition.

To determine the role of temperature in tolerance across a broader set of clinical isolates, we surveyed 133 clinical *C. albicans* isolates for their susceptibility/resistance and tolerance levels in YPD medium under standard lab conditions (30°C) and at normal (37°C) and febrile (39°C) human body temperatures. We used disk diffusion assays analyzed by *diskImageR (14)* to measure the susceptibility/resistance (as the radius of the zone of inhibition (RAD_{20},)) and the tolerance (as the fraction of growth (FoG_{20},) in the zone of inhibition relative to growth outside it) for each strain. For most strains (69.2%), tolerance levels were affected by temperature, with higher tolerance levels at 37°C and 39°C than at 30°C, and little difference between tolerance levels measured at 37°C vs 39°C (Fig. 1A). The susceptibility/resistance levels remained similar at the three temperatures. Thus, tolerance, but not resistance, was affected by both medium composition and temperature.

We classified the isolates into three groups based on the effect of temperature on growth temperature-dependence of the tolerance (FoG_{20} values): temperature-

enhanced tolerant (**TET),** (non-tolerant at 30 °C but highly tolerant at higher

temperatures (n=92, average FoG_{20} values of 0.21 ± 0.03, 0.63 ± 0.17, 0.68 ± 0.15, at

30°C, 37°C and 39°C, respectively)); all-temperature tolerant (**ATT**), which had similar

FoG_{20} values at all three temperatures (n=31, FoG_{20} values of 0.52 ± 0.12, 0.61 ± 0.15,

0.61 ± 0.15, at 30°C, 37°C and 39°C, respectively); and non-tolerant (**NT**), which

exhibited only baseline levels of FoG_{20} at all three temperatures (n=10, average FoG_{20}

values 0.21 ± 0.03, 0.23 ± 0.03, 0.24 ± 0.03, at 30°C, 37°C and 39°C, respectively). At

37°C and 39°C, 92.5% (123 out of 133) of the strains exhibited similar tolerance levels,

while at 30°C, 76.7% (102 out of 133) were not tolerant on the YPD medium (Fig. 1B).

Thus, both body and febrile temperature promoted tolerance on YPD medium in most

isolates.

Growth dynamics of tolerant cells are similar in a broad range of supra-MIC

fluconazole concentrations

To better characterize the similarities and differences in the drug responses of the

different classes of isolates, we used E-Test[®] strips to measure susceptibility as the

MIC after 24 h of growth, and spot dilution assays (13), analyzed at 48h, to measure

tolerance. One representative isolate was used for each of the three temperature-

response classes: SC5314 (the *C. albicans* lab strain) for TET, YJB-T1891, for ATT,

and YJB-T490 for NT strain classes. The three isolates had the same MIC value (1

µg/ml) at 30°C, 37°C and 39°C on YPD (Fig. 2A)). However, spot assays revealed

significant differences in the ability of individual colonies to grow in the presence of FLC

(Fig. 2B). For the TET isolate at 30°C, and for the NT isolates at all three temperatures,

no growth was evident above the MIC, indicating that they were not tolerant under these

conditions. By contrast, the isolates that exhibited tolerance (the ATT isolate at all three

temperatures, and the TET isolate at 37°C and 39°C) grew similarly at all the drug

concentrations tested (up to 128 µg/ml). Thus, for TET and NT tolerant isolates, tolerant

growth does not exhibit much concentration-dependence from 2-128 µg/ml FLC.

We investigated the impact of MIC and tolerance on population growth dynamics in the
presence of FLC by plating 100-200 colony-forming units of each isolate on YPD plates
supplemented with FLC at concentrations from 0.25-128 µg/ml. We monitored growth
dynamics at both 30°C and 37°C for 48 h at 30 min intervals using *ScanLag*, which
reports on the time of appearance and growth rates of individual colonies (10, 15).
205
At 30°C, colonies from TET isolate SC5314 failed to grow at FLC concentrations above
the MIC (1 µg/ml), while at 37°C, growth was detectable. Colonies of the ATT isolate
(YJB-T1891) grew at both 30°C and 37°C in all FLC concentrations, regardless of the
MIC (1 µg/ml) (Fig. 2C) and for the NT isolate (YJB-T490), no colonies grew at either
30°C or 37°C in the supra-MIC FLC concentrations. These results are consistent with
those from the E-test and spot dilution assays above.
212
ScanLag also measures colony growth on agar and reports the time required for a
colony to become detectable on the plate. This time of appearance is considered a
proxy of lag phase length (15). In all three strain types, at both 30°C and 37°C, the
average time of appearance of colonies on drug plates was dose-independent at drug
concentrations below the MIC. Above the MIC, the time of appearance of isolates that
exhibited tolerance in the spot assays was longer than at sub-MIC drug concentrations,
yet was dose-independent: the whole population appeared with a later time of
appearance at all supra-MIC concentrations tested. In contrast to time of appearance,
growth rate was dose-dependent at sub-MIC drug concentrations. Above the MIC, the
growth rate, like the time of appearance, was slower than at sub-MIC concentrations,
yet was drug dose independent. Thus, at supra-MIC drug concentrations, tolerant cells
appeared later, presumably because they have a more prolonged lag phase, and grew
slower than at sub-MIC concentrations. Nonetheless, these growth properties were
similar at concentrations from 2 µg/ml to 128 µg/ml of fluconazole, consistent with the
low degree of concentration-dependence of tolerance.
228
229   **Evolution of higher tolerance from cells in the non-tolerant state**
230

To ask how a non-tolerant isolate adapts to supra-MIC FLC concentrations, we plated

approximately 1x10^6 cells of TET isolate SC5314 and NT isolate YJB-T490 on YPD

plates supplemented with 8 µg/ml to 128 µg/ml FLC. The plates were incubated under

conditions where these isolates were non-tolerant: 30˚C for SC5314 and 30˚C and 37˚C

for YJB-T490. As controls, we plated 1x10^6 cells of SC5314 at 37°C and the ATT strain

(YJB-T1891) at 30˚C and 37˚C on the same range of FLC concentrations. As expected,

all controls produced a lawn of cells. By contrast, the TET strain at 30˚C and the NT

strain at both temperatures gave rise to a few hundred to a few thousand colonies

(adaptors) per plate after 5 days on the drug plates (Fig. S2).

Disk diffusion assays were then performed on 90 randomly selected adaptors from

SC5314 and YJB-T490 parent strains (18 adaptors from each 30°C drug plate). All

these adaptors had a notable elevated tolerance (Fig. 3) when tested at 30°C. When

tested at 37°C, all SC5314-derived adaptors (like the SC5314 ancestor) had high

tolerance (Fig. 3 Top panel), while all YJB-T490-derived adaptors had low tolerance at

37°C (Fig. 3 Middle panel). Thus, adaptors derived from SC5314 at 30˚C acquired

tolerance at 30˚C and maintained their tolerance at 37˚C, and adaptors derived from

YJB-T490 at 30˚C acquired tolerance that was limited to 30˚C. By contrast, many

adaptors derived from YJB-T490 at 37˚C were tolerant at both 30˚C and 37˚C (Fig. 3

Bottom panel). Thus, depending on the temperature used to evolve the adaptors, the

NT isolate adapted to FLC in two distinct manners: at 30°C they acquired a new type of

conditional tolerance that we term _t_emperature-_s_ensitive _t_olerance (**TST)** seen only at

30°C and not at 37°C; when evolved at 37°C, NT isolates acquired tolerance that was

detectable at both 30˚C and 37˚C, a phenotype akin to that of other ATT-like adaptors.

Recurrent aneuploidy enables the acquisition of fluconazole tolerance

To identify genomic mechanisms by which the SC5314 and YJB-T490-derived adaptors

acquired tolerance, we performed deep sequencing of 18 independent adapted isolates

for the TET and NT strains collected at 30˚C on 8 µg/ml, 32 µg/ml, and 128 µg/ml of

FLC. Taken together, 54 adaptors derived at 30˚C from TET and NT isolates were

sequenced.

Aneuploidy was prevalent among the adaptors. Out of the 54 TET-derived adaptors,

one (FY9) that appeared on 8 µg/ml FLC plate grew poorly and was excluded from

sequencing and further analysis. From the others, 50 (94.3%) were aneuploid (Fig. S3)

and three were euploid. Among the 54 NT-derived adaptors at 30˚C, 52 (96.3%) were

aneuploid (Fig. S4) and two were euploid. Four of the five euploid adapters (from TET-

and NT-derived) appeared at 8 µg/ml FLC, the lowest selective drug concentration

tested. Variant calling of the 5 euploid adaptors revealed 11 high confidence _de novo_

SNPs (1 to 4 per adaptor strain; 6 SNPs in the TET-derived adaptors and 8 SNPs in the

NT-derived adaptors) (Table S1). Among them, a G2926C missense mutation of

C1_06340W was detected in 4 of the 5 euploid adaptors; this encodes a protein of

unknown function that has unphased allelic variation and contains ambiguous

sequences (http://www.candidagenome.org/cgi-

bin/locus.pl?locus=C1_06340W&organism=C_albicans_SC5314). Other mutations were

all unique to a single adaptor: four of the remaining SNPs were in genes encoding

proteins with unknown functions. None of the remaining 6 genes encodes any functions

known to directly or indirectly affect drug responses (e.g., ergosterol biosynthesis, efflux

pumps, lipid biosynthesis pathways or known stress response regulators). Mutation in

one of the genes, _PGA63,_ is similar to _S. cerevisiae SEC31,_ which is annotated as

having increased tolerance for aneuploid chromosomes (16), but the role of this gene in

C. albicans will need to be investigated.

Analysis of adaptor karyotypes revealed several recurrent aneuploidies, several of

which were seen in both the TET- and NT-derived isolates. The most prevalent

adaptors (35 TET adaptors; 27 NT adaptors) had aneuploidy involving ChrR, either as

trisomy of the whole chromosome (ChrRx3) (9 from TET, 10 from NT), monosomy of

the left arm distal to the rDNA repeats (SegChrRx1) (6 from TET), or trisomy of ChrR

from rDNA repeats to the right telomere (SegChrRx3) (12 from NT), or a combination of

SegChrRx3 and SegChrRx1 (10 from TET). These ChrR aneuploidies appeared alone

or in combination with trisomy of other chromosome(s) (Chr4 or Chr7 in TET adaptors;

any combination of Chr4 and Chr6 trisomy alone or together with trisomy of Chr7 in NT

adaptors) (Fig. 4). Five TET adaptors and 19 NT adaptors had aneuploidy of

chromosome(s) other than ChrR. The 5 TET adaptors had a single additional

aneuploidy: SegChr1x1, Chr4x3, Chr4x4, SegChr5x1, or Chr6x3, (Fig. S3). Among the

19 NT adaptors, only one had a single aneuploid chromosome (Chr5x1) and this

monosomic strain grew very slowly. The remaining 18 adaptors had at least two

aneuploid chromosomes, mostly Chr4x3 together with Chr6x3 (Fig. S3 and summarized

in Fig. 4).

Taken together, NGS revealed 21 different karyotypes among the 50 TET-derived ATT-

like aneuploid adaptors, and 17 different karyotypes among the 53 NT-derived TST

aneuploid adaptors. Despite differences in genetic background and FLC concentration

used for selection, the same aneuploidies (predominantly ChrRx3 or SegChrRx3) were

recurrently associated with the emergence of FLC tolerance. This occurred in both the

ATT types derived from TET ancestral isolates and the TST types derived from NT

ancestral isolates.

The NT-derived adaptors obtained at 37°C also had diverse karyotypes. Each of the

nine sequenced NT→ATT adaptors had a unique karyotype that was *not* seen in the

NT→TST adaptors obtained at 30°C (Fig. 4): 7 had amplification of Chr6 alone (n=1) or

in combination with aneuploidy (mostly trisomy) of one or more other chromosomes

(n=6). Only two NT→ATT adaptors did not have Chr6 aneuploidy: one adaptor had

SegChr1x1 alone, and one had SegChr2x1 alone.

Taken together, the karyotypes of the adaptors were FLC dose-independent but

temperature-dependent. At 30°C, the TET- and NT-derived karyotypes involved

recurrent ChrR amplification in TET→ATT and NT→TST adaptors selected across the

range of FLC concentrations used. By contrast, from selection at 37°C, NT→ATT

adaptors had distinctly different aneuploidy: predominantly Chr6 trisomy and tetrasomy,

alone or in combination with another aneuploid chromosome. Thus, selection at 30°C

yielded similar karyotypes with different tolerance temperature responses in different

strain backgrounds. Moreover, different selection temperature (30°C vs 37°C) yielded

different karyotypes in the same strain background (NT→TST vs NT→ATT) (Fig. 3 and

Fig. 4).

Loss of aneuploidy is associated with loss of acquired fluconazole tolerance.

Whole chromosome aneuploidies can be gained and lost through mitotic

missegregation. Previously we found that reversible chromosome gain and loss affected

general fitness and drug responses in *C. albicans* (17, 18). Here, we asked if the FLC

tolerance associated with recurrent aneuploidy was also unstable. Adaptors with

different aneuploidies were spread onto YPD (without drug) and grown at 30°C for 36h.

All yielded a mixture of colony sizes, with primarily small and some large colonies (Fig.

5A), although the frequency of appearance of large colonies was different for different

aneuploids. We randomly selected one small and one large colony from each strain and

tested them by disk diffusion assays at 30°C.

To ask if point mutations had any role in the acquisition of tolerance, we performed

variant calling to identify any SNPs of moderate to high effect (missense, nonsense,

frameshift) that passed our quality filters (see Materials and Methods) in the small

(aneuploid) and large colonies of the tolerant adaptors (Table S2). Small and large

colonies from the same parent shared 14 SNPs. We assume that these SNPs were

acquired during selection on FLC and were not drivers of tolerance. Taken together, in

six tolerant adaptors representative of the six different karyotypes, we found variants

that caused missense mutations in 23 genes. None of them were significantly (FDR >

0.1) enriched for gene ontology (GO) terms. Twelve genes were not annotated to any

biological process, and 14 genes had unknown molecular functions. None of the SNPs

found here were in genes encoding known or predicted drug efflux pumps or in known

ergosterol biosynthesis enzymes or regulators.

Importantly, all the small colonies retained the higher tolerance seen in their parent

strain <u>and</u> remained aneuploid like the adaptors from which they were derived. By

contrast, all the large colonies were no longer tolerant (Fig. 5B) <u>and</u> had lost the

aneuploid chromosome(s) to become euploid (Fig. 5C). This strong correlation, between

the presence of aneuploidy and tolerance in all the small and loss of aneuploidy and

tolerance in all the large progeny, supports the idea that the aneuploidies acquired

during passaging on FLC were the major drivers of tolerance in the original adaptors.

Because only some aneuploid chromosomes associated with tolerance, we presume

that these aneuploidies appeared, in some cases recurrently, because they provide a

selective advantage in the presence of drug (17, 19-23).

Evolutionary trajectories differ at sub-MIC versus supra-MIC fluconazole

concentrations

Because the TET and NT isolates acquired tolerance, but not resistance, when exposed

to supra-MIC concentrations of FLC on solid medium, we next asked if this was a

function of the FLC concentration. Therefore, we evolved one TET isolate (SC5314) and

one NT isolate (YJB-T490) by passaging them daily (1:1000 dilutions) in YPD broth

supplemented with FLC ranging from 4X below the MIC (0.25 µg/ml) to 128X above the

MIC (128 µg/ml) for a total of 15 days. As a control, we also passaged the isolates in

YPD without drug.

Every five days we randomly selected 18 or 96 colonies from the TET and NT cultures,

respectively. In total, we measured 18 and 96 TET and NT derivatives per day (1, 5, 10,

and 15 passages) and per drug concentration (0, 0.25, 0.5, 1, 2, 4, 8, 16, 32, 64 and

128 µg/ml FLC) for a total 792 (SC5314 derivatives) and 4224 (YJB-T490 derivatives)

measurements, respectively (Fig 6A and 6B).

For SC5314 (MIC ~ 1 µg/ml FLC), among the 504 adaptors evolved at supra-MIC

concentrations (2-128 µg/ml FLC), most exhibited tolerance (increased FoG_{20}), while

none of the adaptors acquired increased FLC resistance (Fig. 6A). In fact, some

adaptors were *more susceptible* to FLC, as evidenced by a *larger* RAD_{20} than the

parent strain. Similarly, for YJB-T490 (MIC ~ 0.75-1 µg/ml), most of the 2688 adaptors

evolved at supra-MIC concentrations (2-128 µg/ml), exhibited tolerance, and *none*

exhibited resistance (Fig. 6B). Interestingly, both the frequency with which tolerance

arose and the range of tolerance levels achieved were similar at all the supra-MIC

concentrations used for passaging (Fig. 6A). This suggests that once the drug

inactivated its target, additional drug had little if any effect on how often or how much

tolerance emerged.

At the MIC, SC5314 adaptors gained either resistance or tolerance (but not both) after 5

and 10 passages, while only resistant isolates appeared after 15 days of passaging at

the MIC. Thus, it appears that at the MIC, SC5314 adaptors that were tolerant after 5

and 10 passages became resistant after 15 passages. Furthermore, there was stepwise

increase in the level of resistance in adaptors evolved for 5, 10 and 15 days in 1 µg/ml

FLC (Fig. 6A and Fig. S5). For YJB-T490 adaptors, only tolerant isolates were detected

at the MIC. In some assays, the MIC of YJB-T490 is slightly lower than that of SC5314,

and thus 1 µg/ml may be slightly higher than the actual MIC for this strain.

At sub-MIC concentrations (0.25 and 0.5 µg/ml), SC5314 adaptors became either

resistant (decreased RAD_{20}) or tolerant (increased FoG_{20}) (but not both). Among the

adaptors evolved in 0.25 µg/ml and 0.5 µg/ml FLC, resistance emerged in 35 and 30

adaptors, respectively, while tolerance was detected in 1 and 6 adaptors, respectively.

The remaining adaptors appear to have had more transient tolerance that had been lost

during growth in the absence of FLC. Adaptors evolved for 10 and 15 days in 0.5 µg/ml

FLC also exhibited stepwise increases in resistance. YJB-T490 adaptors also acquired

either resistance or tolerance (but not both). Among adaptors evolved in 0.25 µg/ml FLC,

tolerance appeared after ten passages; yet after 15 passages in 0.25 µg/ml FLC for 15

days, ten were tolerant, and 23 were resistant. At 0.5 µg/ml, the proportion of resistant

YJB-T490 adaptors relative to tolerant adaptors increased over time (0 vs 58; 29 vs 18,

and 72 vs 24, for adaptors from passages 5, 10 and 15, respectively (Fig. 6). While this

is consistent with the idea that isolates may have acquired tolerance in early passages

and then become resistant in the later passages, we cannot be sure that the resistant adaptors are progeny of the prior tolerant ones.

The dynamics of the emergence of tolerance and resistance also differed considerably. In both parental strain backgrounds tolerance emerged more rapidly, within a single passage, while the first resistant adaptors appeared after 5 passages and only in SC5314 at 1 µg/ml FLC, while adaptors passaged in 0.25 µg/ml and 0.5 µg/ml only produced resistant and tolerant progeny after 10 days, and resistant adaptors after 15 days. In YJB-T490, *tolerant* adaptors evolved at or below the MIC 1 µg/ml, 0.5 µg/ml and 0.25 µg/ml initially appeared in passages 1, 5, and 10, respectively, while *resistant* adaptors that emerged from 0.5 µg/ml and 0.25 µg/ml FLC initially appeared on days 10 and 15, respectively. Thus, the acquisition of resistance and tolerance occurred at different relative stress levels (sub-MIC vs supra-MIC) and with different temporal dynamics (Fig. 6), and tolerant isolates generally appeared more rapidly than resistant isolates.

Distinct and diverse genetic alterations in fluconazole-resistant versus fluconazole-tolerant adaptors

Individual colonies that arose following the evolution of SC5314 in broth passages had a range of resistance or tolerance levels, suggesting that they may have acquired distinct genetic or genomic changes. To test this hypothesis, we sequenced 73 randomly selected adaptors (34 resistant and 39 tolerant) derived from SC5314. Large-scale genome changes were visualized with Ymap (Fig. 7 and Fig. S6), and variant calling was performed to identify mutations in predicted open reading frames, focusing on missense mutations and frameshift mutations.

Notably, all 39 tolerant adaptors that were sequenced were aneuploid (Fig. 7, lower section): 13 had whole or segmental aneuploidy only for all or segments of ChrR: (ChrRx3 (n=9), SegChrRx3 from 1.89Mb to right telomere (n=3), or SegChrRx1 from 1.89Mb to right telomere (n=1)). An additional 25 adaptors were aneuploid for ChrR in

combination with aneuploidy of another one or two chromosomes (n=22 and n=3,

respectively). Only one tolerant adaptor (Chr6x3, ABB, evolved over 10 passages in 1

µg/ml FLC) did not include copy number changes on ChrR. Thus, ChrR aneuploidy was

the most prevalent aneuploidy associated with tolerance.

Among the 34 resistant adaptors sequenced, the majority (22/34) were euploid and 12

were aneuploid. Among the aneuploid adaptors, two (SY60 and SY61) had Chr3x3, five

(SY20, SY21, SY22, SY51, SY55) had Chr5x3, four (SY50, SY56, SY58, SY59) had

Chr7x3 (AAB), and one adaptor (SY57) had Chr6x3 (AAB) together with Chr7x3 (AAB).

Notably, none of the resistant adaptors affected ChrR copy number and the karyotypes

of the resistant adaptors were distinct from the tolerant adaptors (Fig. 7 and Fig.S5).

Therefore, tolerant and resistant adaptors acquired distinct genomic changes: all

tolerant adaptors were aneuploid, with ChrR aneuploidy as the predominant change

while resistant adaptors were mostly euploid or were trisomic for either Chr3, Chr5, or

Chr7.

Importantly, similar types of karyotypes were detected in SC5314 and YJB-T490

derived tolerant adaptors selected on supra-MIC FLC concentrations on agar plates (Fig.

S3 and Fig. S4). Furthermore, the karyotypes of tolerant adaptors appeared in a dose-

independent and time-independent manner (Table 1 and Fig. 7). For example, ChrRx3

aneuploidies appeared after one day in 2 µg/ml or 128 µg/ml FLC, after 5 days in 1

µg/m FLC, after 10 days in 0.25 µg/ml FLC, and after 15 days in 2 µg/ml FLC. Similarly,

ChrRx3+Chr4x3 appeared after 1 day in 2 µg/ml FLC, 10 days in 1 µg/ml, 2 µg/ml and

128 µg/ml FLC, and after 15 days in 2 µg/ml and 128 µg/ml FLC. And segChrRx3 (from

the left telomere to 1.89 MB) appeared after 10 days in 0.5 µg/ml FLC, and 15 days in 2

µg/ml and 128 µg/ml FLC (Table 1). Thus, the same aneuploidies can arise at different

concentrations and at different times after initial drug exposure.

To ask about point mutations that may influence the degree of tolerance or resistance,

we identified SNPs in the 39 tolerant and 34 resistant adaptors (Table S3). In total we

found from 2 to 5 SNPs per tolerant isolate and 0 to 2 SNPs per resistant isolate. One of

the two ORFs with detectable SNPs in both a tolerant and a resistant adaptor was

C2_08380C, a possible ortholog of *S. cerevisiae DPB11,* that, when mutated, causes

gross chromosomal rearrangements, chromosome loss and an increase in spontaneous

DNA damage (24). While the degree to which these specific SNPs affect the encoded

function is not known, this gene might contribute to increased frequencies of mutations

in the adaptors.

SNPs found only in *tolerant* adaptors were detected in 27 ORFs. However, of these 27

only one could be associated with drug tolerance: C1_04010C. We detected 2 SNPs in

this ORF in 4 different adaptors. This gene encodes a protein with a predicted NADP-

dependent oxidoreductase domain, and its transcript is induced by ketoconazole, yet is

repressed by Upc2p, a transcription factor that up-regulates ergosterol biosynthesis (25).

There was no statistically significant (FDR > 0.1) gene ontology term enrichment of

these 27 genes. Thus, the few SNPs that appeared may be neutral mutations and do

not appear to have any obvious connections to drug resistance or tolerance.

SNPs found only in *resistant* adaptors were detected in 7 genes with no detectable

gene ontology term enrichment (FDR > 0.1), but none were in genes that encode efflux

pumps, or that are known to affect ergosterol biosynthesis, the two major mechanisms

of azole resistance. We suggest that some of these SNPs are probably neutral or

nearly-neutral mutations that arose during the course of passaging and/or false positive

variants that our quality filters were not stringent enough to filter out.

Discussion

Contrary to the concept from bacteria that antimicrobial drugs primarily select for the rapid evolution of drug resistance, here we found that *C. albicans*, a common cause of systemic bloodstream infections, primarily acquired tolerance to the widely used drug FLC. Importantly, this antifungal tolerance appeared rapidly, in some cases within a single day of exposure to drug concentrations above the strain MIC, while resistant isolates appeared after 5 days of exposure to low levels of FLC. Furthermore, the resistant isolates exhibited a stepwise increase in MIC levels (Fig. 6A), and we found no evidence that mutations in classic genes associated with FLC resistance played any role in the resistance acquired under the conditions used.

Here, we found that most clinical *C. albicans* isolates exhibit FLC tolerance at physiologically relevant temperatures, and not at 30°C, the temperature often used in lab studies (Fig. 1). This antifungal tolerance is largely dose-independent at supra-MIC drug concentrations (Fig. 2). Experimental evolution in supra-MIC drug concentrations rapidly selected for the acquisition of tolerance that was associated with a recurrent set of aneuploidies (Figs. 3,4,6,7). Resistant isolates only appeared at or below the MIC, after 5 or more days of passaging (Fig. 6). While most resistant isolates were not aneuploid, those that carried recurrent aneuploidies had extra copies of different chromosomes from those seen in tolerant isolates (Fig. 7). Furthermore, resistance (but not tolerance) appeared to be acquired in a stepwise manner (Fig. 6). Thus, experimental evolution at sub-MIC FLC concentrations selected for isolates that were either only tolerant or only resistant, and a different set of recurrent aneuploidies conferred tolerance vs resistance.

Of the original 133 clinical *C. albicans* isolates tested at 37°C and 39°C, up to 92.5% of the isolates were tolerant, while at 30°C only 23.3% of isolates were tolerant. Thus, it appears that FLC tolerance is prevalent in clinical isolates, at least those collected from Israeli patients. These clinical strains are largely euploid; thus, we suggest that tolerance is primarily due to genomic factors, including inherent genetic backgrounds,

that modulate stress responses in different physiological conditions, such as

temperature and medium composition. In addition, the acquisition of aneuploidy can

increase or modulate tolerance levels. While this study focused on 133 Israeli clinical

isolates, it will be interesting to determine if other clinical isolates exhibit temperature-

enhanced tolerance.

Tolerant isolates were able to grow at supra-MIC FLC concentrations up to 128 µg/ml.

Furthermore, the time of colony appearance, as well as growth rate of colonies on drug

plates were dose-independent at supra-MIC FLC concentrations. Yet, below the MIC,

colony growth rates were dose-dependent with colonies growing more slowly with

increasing drug concentrations. This suggests that *C. albicans* mounts distinct cellular

responses at sub-MIC versus supra-MIC drug concentrations. The concentration-

independent nature of tolerant cell growth rates and the initial time of colony

appearance at supra-MIC, and the concentration-dependent nature of resistance (Fig.

2B and C) at sub-MIC concentrations, highlights probably mechanistic differences

between these processes. We posit that tolerance involves cellular stress mechanisms

that are less sensitive to the intracellular drug concentration and enable growth when

the drug target, in this case lanosterol demethylase, is completely inactivated by drug.

The fact that different aneuploidies or mutations accompany tolerance vs resistance

further supports this idea. Furthermore, isolates with different genetic backgrounds at

the temperatures used and on agar or in broth media, adapted to supra-MIC FLC

primarily by acquiring the same aneuploidies involving ChrR.

All but five tolerant adaptors had at least one aneuploid chromosome. In addition to the

recurrent karyotypes involving ChrR aneuploidies, some tolerant adaptors acquired

other aneuploidies. This suggests that individual cells use a similar strategy,

chromosome instability, to reach non-identical outcomes (e.g., different levels of

acquired tolerance) from different combinations of aneuploid chromosomes that provide

a growth advantage in supra-MIC drug concentrations. Even though we detected a few

SNPs in the aneuploid isolates and we do not know the degree to which they may (or

may not) contribute to tolerance, the fact that loss of the aneuploid chromosome(s) was

accompanied by loss of FLC tolerance suggests that aneuploidy was the primary

mechanism conferring tolerance. The mechanism(s) by which the 5 euploid adaptors

became FLC tolerant are less clear as the few SNPs identified in these strains did not

encode obvious mediators of drug efflux, stress response or alternative strategies to

bypass the deleterious effects of ergosterol depletion due to FLC exposure. While the

SNPs or some other unidentified genetic change can not be ruled out, another

possibility is that non-genetic metabolic shifts contributed to the ability of these isolates

to grow despite the presence of supra-MIC drug concentrations (26).

Strain background and environmental conditions also contribute to the selective

conditions that favor the acquisition of different aneuploidies. Interestingly, in both the

TET and NT backgrounds, amplification of portions of ChrR enabled tolerance at 30°C

but not at 37°C. Yet, for the NT strain, Chr6 trisomy increased tolerance at both 30°C

and 37°C in the presence of the same range of FLC concentrations. Thus, the rapid

emergence of tolerance can occur via multiple routes, and the trajectories of the routes

are affected by the original strain background and the selection conditions. Future work

will require analyzing the trajectories of adaptation of additional TET and NT isolates

and identifying the chromosomal region(s) that specifically affect tolerance at different

temperatures.

Some of the tolerant adaptors also became less FLC resistant (more susceptible, larger

RAD_{20},). This was also seen for isolates passaged in posaconazole, another fungistatic

azole (27). This supports the idea that the acquisition of antifungal resistance and

antifungal tolerance follow different trajectories and that supra-MIC concentrations

preferentially and can rapidly select for azole tolerance along with decreased azole

resistance.

All tolerant strains acquired aneuploidy together and had a similar range of tolerance

levels at all supra-MIC concentrations. By contrast, the resistant mutations that

appeared at concentrations at or below the MIC, continued to become more resistant

over time (Fig. 6 and Fig. S5). Interestingly, this stepwise acquisition of improved fitness

was seen both with aneuploid and euploid resistant adaptors. This highlights the very

different evolutionary trajectories that occur above and below the MIC. At all the supra-

MIC concentrations tested, when the drug target is completely inhibited/saturated by

drug, cells must respond with strategies independent of the drug-target interaction. By

contrast, at sub-MIC concentrations, the drug target is not saturated, all cells are able to

grow to some degree, providing the opportunity for the acquisition of a series of

mutations that incrementally increased resistance. At sub-MIC, tolerance also emerged

occasionally.

Why does aneuploidy appear so rapidly and with such a high frequency ($\sim 10^{-3}$) following

exposure to supra-MIC drug concentrations? We suggest that three forces drive the

rapid appearance of aneuploidy at high frequency: first, random aneuploid

chromosomes maintained in the parent population may provide standing variation;

second, FLC induces chromosome mis-segregation via cell cycle defects that yield

tetraploid intermediates and, subsequently, random aneuploids (28, 29) at high

frequency; and third, strong selection for specific aneuploidies that provide an adaptive

advantage in the presence of the drug allows the aneuploid isolates to compete and

outgrow arrested euploid cells (18). Chromosome mis-segregation events that lead to

whole chromosome aneuploidy are relatively frequent (every 5×10^5 cell divisions in

yeast (30) and on the order of once every 10^4 to 10^5 divisions in mammalian cells (31)).

Since *C. albicans* chromosomes contain between 407-1383 genes per chromosome

(http://www.candidagenome.org/cache/C_albicans_SC5314_genomeSnapshot.html, as

of Nov 24, 2022), it follows that a single aneuploid chromosome should affect the

stoichiometry of many proteins. Since segmental aneuploidies, which are dependent

upon recombination events, also appeared, it is tempting to speculate that either drug

stress increases the likelihood of mitotic recombination as was seen for LOH events

(32), or that recombination events are more frequent in strains carrying aneuploid

chromosomes. Consistent with this idea, haploid *S. cerevisiae* lab strains carrying single

extra chromosomes conferred increased mitotic recombination as well as chromosome

instability (33).

In this study, tolerant adaptors emerged more rapidly (within one passage) than

resistant adaptors. Ultimately, all supra-MIC isolates became tolerant, while only some

of the sub-MIC adapters became resistant. Indeed, even resistant adaptors that had

acquired resistance via aneuploidy appeared much later in the passaging and may have

arisen from tolerant intermediates (e.g., some adaptors evolved in 0.5 µg/ml and 1

µg/ml FLC were tolerant on day 10, but all were resistant on day 15, and some were

aneuploid (Fig. 6)). Thus, while the dynamics of acquiring tolerance is different at sub-

and supra-MIC drug concentrations, a similar set of aneuploidies confers resistance,

and a different set of aneuploidies can confer tolerance.

In *C. albicans*, two major mechanisms contribute to FLC resistance: alteration of the

drug target and increased drug efflux. A combination of these mechanisms causes

stepwise development of FLC resistance ((34); reviewed in (35)) or can arise in a single

segmental aneuploidy (19). Surprisingly, we did not find any of the classic resistance

mutations in resistant adaptors. This could be due to the high fitness cost of mutations

in these genes (34, 36, 37). In the presence of sub-MIC FLC, cells are under weak

stress and mutations with a high fitness cost might be outcompeted. In general, gain-of-

function mutations in *MRR1*, *TAC*, or *UPC2* in FLC-resistant clinical isolates incur a

fitness cost and are outcompeted by the matched susceptible isolates both *in vitro* and

in *vivo* (36). Artificially introducing resistance-enhancing mutations causes a stepwise

increase in resistance that is associated with a gradual reduction in fitness both in *vivo*

and *in vitro* (34). Similarly, experimental evolution with increasing amounts of FLC

yielded resistant mutants with mutations in *TAC1*, *UPC2,* and/or *ERG11* that were less

fit than their parents in macrophages, as well as in the presence of several stresses,

including cell wall stress, cell membrane stress, salt stress, oxidative stress and

temperature stress (37). The longer time required for resistant adaptors to arise, even if

aneuploid, provides indirect support for the idea that resistance is due to the

accumulation of multiple mutational events.

Paradoxically, *C. albicans* strains that we surveyed from the clinic are euploid, not

aneuploid, despite the observation that all adaptors that rapidly acquired increased

tolerance were aneuploid. This implies that multiple changes in chromosome copy

number can confer rapid increases in tolerance levels. Under drug selection, the

aneuploidies should provide a fitness advantage and should be maintained at least until

other, more stable mutations, with improved fitness, out-compete the aneuploids (38).

Future experiments will address whether tolerant adaptors that acquired aneuploidies

eventually evolve to lose the aneuploidy and retain high levels of drug tolerance and/or

resistance.

Cells appear to adopt different adaptive trajectories at supra-MIC versus sub-MIC drug

concentrations. At supra-MIC FLC, extreme stress may favor the immediate

appearance of tolerance, perhaps because aneuploidy can appear within a single cell

division or because aneuploidy may slow growth and thus might slow drug metabolism.

At sub-MIC FLC concentrations, cells experience only mild stress that does not affect

cell survival, which may enable the evolution of resistance in a stepwise manner. The

degree of resistance, measured by the MIC or RAD, can increase over time of exposure

to the drug (Fig. 6 and Fig. S5). Accordingly, more extended passaging at sub-MIC FLC

concentrations yields adaptors with higher MIC levels. We speculate that this may

involve a sequential acquisition of several different types of resistance mutations,

including point mutations that confer stable drug resistance.

In bacteria, both sub- and supra-MIC antibiotic concentrations select for resistant

mutants, but different selection strengths confer different evolutionary trajectories and

drive the acquisition of different mutations (reviewed in (39). When selected in cidal

antibiotics at supra-MIC drug concentrations, resistance mutations in one or a small

number of "classical" resistance genes are selected. But when exposed to low antibiotic

concentrations, bacteria accumulate mutations of several "non-classic" genes that

individually confer weak resistance but together can confer higher resistance levels (40).

Selection in subinhibitory concentrations also increase rates of mutation, recombination,

and horizontal gene transfer, thereby increasing the appearance of genetic changes. In

addition, selection at sub-MIC antibiotic concentrations can select for plasmids carrying

multiple resistance genes that lead to multidrug resistance (MDR) (41).

In summary, FLC tolerance is prevalent in *C. albicans* clinical isolates, especially at 37°C and it exhibits different temperature responses in different strain backgrounds. The acquisition of resistance and tolerance occurs via distinct evolutionary trajectories that are largely a function FLC stress levels: supra-MIC FLC stress drives tolerance, which appears rapidly and enables growth at wide range of supra-MIC drug concentrations; sub-MIC FLC stress selects for either tolerance or stepwise elevated resistance. Different strains have different intrinsic levels of tolerance, but they generally appear to rapidly acquire additional tolerance by becoming trisomic for specific chromosomes. The most commonly acquired aneuploidy in tolerant adaptors is part or all of ChrR, while trisomy of Chr3 or Chr5, are associated with the relatively rapid appearance of increased antifungal drug resistance.

Materials and Methods

Strains and growth conditions

Strains used in this study are listed in Table S4. Stock cultures of all strains were preserved in 35% glycerol and maintained at -80°C. The 133 clinical *C. albicans* were collected from hospitals in Israel and kindly provided by Ronen Ben-Ami. Unless otherwise specified, cells were routinely grown on Yeast extract-Peptone-Dextrose (YPD)-agar media (1% [w/v] yeast extract, 2% [w/v] peptone and 2% [w/v] D-glucose, 2%[w/v] agar) at 30°C. Other media used in this study include casitone agar plates (0.9%[wt/vol] casitone, 0.5% [wt/vol] yeast extract, 1.15% sodium citrate dihydrate [wt/vol], 2% [wt/vol] glucose, 2% [wt/vol] d-glucose, and 2% [wt/vol] agar), SD agar plates (0.67% [wt/vol] yeast nitrogen base without amino acids, 2% [wt/vol] d-glucose, and 2% [wt/vol] agar), and SDC agar plates (0.67% [wt/vol] yeast nitrogen base without amino acids, 2% [wt/vol] d-glucose, 0.2% [wt/vol] complete amino acid mixture, and 2% [wt/vol] agar). Drugs were dissolved in dimethyl sulfoxide (DMSO) and stored at -20°C.

E-test strip assays

E-tests were performed as described previously (20). Strains were streaked from -80°C

freezer onto YPD agar. After incubation at designated temperature (30, 37 or 39°C) for

24h, colonies were chosen randomly and suspended in distilled water. Cell density was

determined using a hemocytometer. Cells were adjusted to 1×10^6 cells/ml, and 100 µl of

the culture was spread on YPD agar. The plate with an FLC E-test strip (BioMerieux,

Marcy l'Etoile, France) at the center was incubated at designated temperature for 24h

and then photographed.

Disk diffusion assays

Disk diffusion assays were performed as previously described (10, 13). The CLSI M44-

A2 guidelines for antifungal disk diffusion susceptibility testing were followed with slight

modifications. Briefly, strains were streaked from glycerol stocks onto YPD agar and

incubated for 36h at designated temperatures. Colonies were suspended in distilled

water and adjusted to 1×10^6 cells/ml. 100 µl of cell suspension were spread onto 15 ml

YPD plates. An empty paper disk (6 mm diameter, and 0.7 mm thickness)

supplemented with 5 µl of 5 mg/ml FLC was placed in the center of each plate. Plates

were then incubated at designated temperature and photographed at 24h and 48h.

Analysis of the disk diffusion assay was done using the *diskImageR* pipeline (14). The

fraction of growth inside the zone of inhibition and radius of inhibition, referred to as

FoG and RAD throughout the manuscript, represent parameters measured at 20% drug

inhibition (FoG_{20} and RAD_{20}, respectively).

Spot assay

Cells were suspended in distilled water and counted using a hemocytometer. Cell

density was adjusted to 1×10^7 cells/ml. 3 µl of 10-fold serial dilutions were spotted on

YPD plates with or without drugs (control). The plates were incubated at the designated

temperature and photographed after 2 days.

ScanLag assay

The *ScanLag* assay was performed as described in (10) with minor modifications.

Approximately 100 cells were spread onto YPD plates with or without FLC. The plates

were placed on the scanners at designated temperature and scanned every 30 min for

48 h. Image analysis was done in MATLAB using the modified "ScanLag" script (10, 15).

Obtaining drug adaptors from plates

Cells were adjusted to 1×10^7 cells/ml as described above. 100 µl of the culture were

spread on YPD plates supplemented with drugs. The plates were incubated at the

designated temperature for 5 days. Adaptors were randomly chosen.

Daily passage in YPD broth supplemented with fluconazole

SC5314 was inoculated into 1ml YPD broth at a final density of approximately 2.5×10^3

cells/ml. The YPD broth was supplemented with DMSO (negative control) or with 2-fold

increase of FLC from 0.25 µg/ml – 128 µg/ml FLC. Every 24h or when the OD600 of

culture was higher than 1.0, 1 µl of each culture was inoculated into 1ml YPD broth

supplemented with the same concentration of FLC. After 1, 5, 10, and 15 days, the

cultures were washed and diluted with distilled water. Approximately 100 cells were

spread on YPD plates. The plates were incubated at 30°C for 36 h and 18 colonies from

each plate were randomly tested with disk diffusion assays using FLC-containing disks.

Colony instability assay

As described previously (18), aneuploid strains were streaked from -80°C freezer to

YPD plates and incubated at 30°C for 36 h. One small colony was randomly chosen and

suspended in distilled water. Cells were diluted with distilled water and approximately

200 cells were spread on a YPD plate and incubated at 30°C for 36 h. One small (S)

colony and one large (L) colony were randomly chosen for further studies.

Next generation sequencing (NGS)

NGS was performed as described in (18).

Variant calling

Paired-end short reads of all sequences obtained from NGS were trimmed with

Trimmomatic (version 0.39; (42)) with default settings and read quality. Trimmed reads

were aligned to both alleles of the *C. albicans* reference genome

(GCF_000182965.3_ASM18296v3) with Burrows-Wheeler Aligner (bwa mem, version

0.7.17; (43)). The alignments were sorted with SAMtools (samtools view; samtools sort;

version 1.15.1; (44)), and duplicates were marked and removed with Picard Tools

(version 2.27.5; available at http://broadinstitute.github.io/picard/). Indels were then

realigned with GATK (version 3.8-1-0; (45)), and the alignments were resorted and

reindexed with SAMtools. Variants (SNPs and indels) were called using FreeBayes (-J -

K --report-genotype-likelihood-max -a -F 0.1; version 0.9.21; (46)). Variants that were

present in both ancestor and evolved strains were removed using BCFtools (isec;

version 1.16-17; (44)), filtered for quality with BCFtools, and annotated with SNPeff

(version 5.1d; (47)). The validity of each high-quality variant was then checked using the

Integrative Genomics Viewer (IGV; version 2.15.1; (48))

GO analysis

Gene Ontology Term Finder tool on Candida Genome Database

(http://www.candidagenome.org/cgi-bin/GO/goTermFinder) was used for functional

enrichment analysis, using default parameters. Only hits with p-value corrected for false

positives FDR < 0.1 were considered as significant.

Data availability

All sequence data are available in the ArrayExpress database at EMBL-EBI

(www.ebi.ac.uk/arrayexpress) under accession numbers E-MTAB-12175, E-MTAB-

12155, E-MTAB-12169 and E-MTAB-12189.

Acknowledgements

We thank Jacob Steenwyk for helpful suggestions. This work was supported by the National Key Research and Development Program of China (2022YFC2303000), the National Natural Science Foundation of China (NO. 82020108032), and the Innovation Program of Shanghai Municipal Education Commission (NO. 202101070007E00094) to Yuan-ying Jiang. National Natural Science Foundation of China (NO. 81872910), and Shanghai Key Basic Research Project (NO. 19JC1414900) to Yong-bing Cao. Judith Berman also received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 951475).

References

1. Bongomin F, Gago S, Oladele RO, Denning DW. 2017. Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. J Fungi (Basel) 3.
2. Enoch DA, Yang H, Aliyu SH, Micallef C. 2017. The Changing Epidemiology of Invasive Fungal Infections. Methods Mol Biol 1508:17-65.
3. Robbins N, Caplan T, Cowen LE. 2017. Molecular Evolution of Antifungal Drug Resistance. Annu Rev Microbiol 71:753-775.
4. Carolus H, Pierson S, Lagrou K, Van Dijck P. 2020. Amphotericin B and Other Polyenes-Discovery, Clinical Use, Mode of Action and Drug Resistance. J Fungi (Basel) 6.
5. Pappas PG, Kauffman CA, Andes DR, Clancy CJ, Marr KA, Ostrosky-Zeichner L, Reboli AC, Schuster MG, Vazquez JA, Walsh TJ, Zaoutis TE, Sobel JD. 2016. Clinical Practice Guideline for the Management of Candidiasis: 2016 Update by the Infectious Diseases Society of America. Clin Infect Dis 62:e1-50.
6. Robbins N, Wright GD, Cowen LE. 2016. Antifungal Drugs: The Current Armamentarium and Development of New Agents. Microbiol Spectr 4.
7. Roemer T, Krysan DJ. 2014. Antifungal drug development: challenges, unmet clinical needs, and new approaches. Cold Spring Harb Perspect Med 4.
8. Pfaller MA, Diekema DJ, Turnidge JD, Castanheira M, Jones RN. 2019. Twenty Years of the SENTRY Antifungal Surveillance Program: Results for Candida Species From 1997-2016. Open Forum Infect Dis 6:S79-S94.
9. Levinson T, Dahan A, Novikov A, Paran Y, Berman J, Ben-Ami R. 2021. Impact of tolerance to fluconazole on treatment response in Candida albicans bloodstream infection. Mycoses 64:78-85.
10. Rosenberg A, Ene IV, Bibi M, Zakin S, Segal ES, Ziv N, Dahan AM, Colombo AL, Bennett RJ, Berman J. 2018. Antifungal tolerance is a subpopulation effect distinct from resistance and is associated with persistent candidemia. Nat Commun 9:2470.
11. Berman J, Krysan DJ. 2020. Drug resistance and tolerance in fungi. Nat Rev Microbiol 18:319-331.

12. Fisher MC, Alastruey-Izquierdo A, Berman J, Bicanic T, Bignell EM, Bowyer P, Bromley M,
Bruggemann R, Garber G, Cornely OA, Gurr SJ, Harrison TS, Kuijper E, Rhodes J,
Sheppard DC, Warris A, White PL, Xu J, Zwaan B, Verweij PE. 2022. Tackling the
emerging threat of antifungal resistance to human health. Nat Rev Microbiol 20:557-571.
13. Xu Y, Lu H, Zhu S, Li WQ, Jiang YY, Berman J, Yang F. 2021. Multifactorial Mechanisms of
Tolerance to Ketoconazole in Candida albicans. Microbiol Spectr 9:e0032121.
14. Gerstein AC, Rosenberg A, Hecht I, Berman J. 2016. diskImageR: quantification of
resistance and tolerance to antimicrobial drugs using disk diffusion assays. Microbiology
(Reading) 162:1059-1068.
15. Levin-Reisman I, Fridman O, Balaban NQ. 2014. ScanLag: high-throughput quantification
of colony growth and lag time. J Vis Exp doi:10.3791/51456.
16. Torres EM, Dephoure N, Panneerselvam A, Tucker CM, Whittaker CA, Gygi SP, Dunham
MJ, Amon A. 2010. Identification of aneuploidy-tolerating mutations. Cell 143:71-83.
17. Yang F, Teoh F, Tan ASM, Cao Y, Pavelka N, Berman J. 2019. Aneuploidy Enables Cross-
Adaptation to Unrelated Drugs. Mol Biol Evol 36:1768-1782.
18. Yang F, Todd RT, Selmecki A, Jiang YY, Cao YB, Berman J. 2021. The fitness costs and
benefits of trisomy of each Candida albicans chromosome. Genetics 218.
19. Selmecki A, Gerami-Nejad M, Paulson C, Forche A, Berman J. 2008. An isochromosome
confers drug resistance in vivo by amplification of two genes, ERG11 and TAC1. Mol
Microbiol 68:624-41.
20. Yang F, Gritsenko V, Lu H, Zhen C, Gao L, Berman J, Jiang YY. 2021. Adaptation to
Fluconazole via Aneuploidy Enables Cross-Adaptation to Amphotericin B and Flucytosine
in Cryptococcus neoformans. Microbiol Spectr 9:e0072321.
21. Yang F, Gritsenko V, Slor Futterman Y, Gao L, Zhen C, Lu H, Jiang YY, Berman J. 2021.
Tunicamycin Potentiates Antifungal Drug Tolerance via Aneuploidy in Candida albicans.
mBio 12:e0227221.
22. Yang F, Kravets A, Bethlendy G, Welle S, Rustchenko E. 2013. Chromosome 5 monosomy
of Candida albicans controls susceptibility to various toxic agents, including major
antifungals. Antimicrob Agents Chemother 57:5026-36.
23. Janbon G, Sherman F, Rustchenko E. 1998. Monosomy of a specific chromosome
determines L-sorbose utilization: a novel regulatory mechanism in Candida albicans.
Proc Natl Acad Sci U S A 95:5150-5.
24. Araki H, Leem SH, Phongdara A, Sugino A. 1995. Dpb11, which interacts with DNA
polymerase II(epsilon) in Saccharomyces cerevisiae, has a dual role in S-phase
progression and at a cell cycle checkpoint. Proc Natl Acad Sci U S A 92:11791-5.
25. Synnott JM, Guida A, Mulhern-Haughey S, Higgins DG, Butler G. 2010. Regulation of the
hypoxic response in Candida albicans. Eukaryot Cell 9:1734-46.
26. Yu JSL, Correia-Melo C, Zorrilla F, Herrera-Dominguez L, Wu MY, Hartl J, Campbell K,
Blasche S, Kreidl M, Egger AS, Messner CB, Demichev V, Freiwald A, Mulleder M, Howell
M, Berman J, Patil KR, Alam MT, Ralser M. 2022. Microbial communities form rich
extracellular metabolomes that foster metabolic interactions and promote drug
tolerance. Nat Microbiol 7:542-555.

27. Kukurudz RJ, Chapel M, Wonitowy Q, Adamu Bukari AR, Sidney B, Sierhuis R, Gerstein
AC. 2022. Acquisition of cross-azole tolerance and aneuploidy in Candida albicans strains
evolved to posaconazole. G3 (Bethesda) 12.
28. Altamirano S, Fang D, Simmons C, Sridhar S, Wu P, Sanyal K, Kozubowski L. 2017.
Fluconazole-Induced Ploidy Change in Cryptococcus neoformans Results from the
Uncoupling of Cell Growth and Nuclear Division. mSphere 2.
29. Harrison BD, Hashemi J, Bibi M, Pulver R, Bavli D, Nahmias Y, Wellington M, Sapiro G,
Berman J. 2014. A tetraploid intermediate precedes aneuploid formation in yeasts
exposed to fluconazole. PLoS Biol 12:e1001815.
30. Hartwell LH, Dutcher SK, Wood JS, Garvik B. 1982. The fidelity of mitotic chromosome
reproduction in S. cerevisiae. Rec Adv Yeast Mol Biol 1:28-38.
31. Rosenstraus MJ, Chasin LA. 1978. Separation of linked markers in Chinese hamster cell
hybrids: mitotic recombination is not involved. Genetics 90:735-60.
32. Forche A, Abbey D, Pisithkul T, Weinzierl MA, Ringstrom T, Bruck D, Petersen K, Berman
J. 2011. Stress alters rates and types of loss of heterozygosity in Candida albicans. mBio
2.
33. Sheltzer JM, Blank HM, Pfau SJ, Tange Y, George BM, Humpton TJ, Brito IL, Hiraoka Y,
Niwa O, Amon A. 2011. Aneuploidy drives genomic instability in yeast. Science
333:1026-30.
34. Sasse C, Dunkel N, Schafer T, Schneider S, Dierolf F, Ohlsen K, Morschhauser J. 2012. The
stepwise acquisition of fluconazole resistance mutations causes a gradual loss of fitness
in Candida albicans. Mol Microbiol 86:539-56.
35. Morschhauser J. 2002. The genetic basis of fluconazole resistance development in
Candida albicans. Biochim Biophys Acta 1587:240-8.
36. Popp C, Hampe IAI, Hertlein T, Ohlsen K, Rogers PD, Morschhauser J. 2017. Competitive
Fitness of Fluconazole-Resistant Clinical Candida albicans Strains. Antimicrob Agents
Chemother 61.
37. Hill JA, O'Meara TR, Cowen LE. 2015. Fitness Trade-Offs Associated with the Evolution of
Resistance to Antifungal Drug Combinations. Cell Rep 10:809-819.
38. Yona AH, Manor YS, Herbst RH, Romano GH, Mitchell A, Kupiec M, Pilpel Y, Dahan O.
2012. Chromosomal duplication is a transient evolutionary solution to stress. Proc Natl
Acad Sci U S A 109:21010-5.
39. Andersson DI, Hughes D. 2014. Microbiological effects of sublethal levels of antibiotics.
Nat Rev Microbiol 12:465-78.
40. Wistrand-Yuen E, Knopp M, Hjort K, Koskiniemi S, Berg OG, Andersson DI. 2018.
Evolution of high-level resistance during low-level antibiotic exposure. Nat Commun
9:1599.
41. Gullberg E, Albrecht LM, Karlsson C, Sandegren L, Andersson DI. 2014. Selection of a
multidrug resistance plasmid by sublethal levels of antibiotics and heavy metals. mBio
5:e01918-14.
42. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30:2114-20.
43. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25:1754-60.

44.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
937   R, Genome Project Data Processing S. 2009. The Sequence Alignment/Map format and
938   SAMtools. Bioinformatics 25:2078-9.
939   45.   Auwera GVd, O'Connor B. 2020. Genomics in the Cloud: Using Docker, GATK, and WDL
940   in Terra. O'Reilly Media.
941   46.   Garrison EP, Marth GT. 2012. Haplotype-based variant detection from short-read
942   sequencing. arXiv: Genomics.
943   47.   Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM.
944   2012. A program for annotating and predicting the effects of single nucleotide
945   polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118;
946   iso-2; iso-3. Fly (Austin) 6:80-92.
947   48.   Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP.
948   2011. Integrative genomics viewer. Nat Biotechnol 29:24-6.
949   49.   Abbey DA, Funt J, Lurie-Weinberger MN, Thompson DA, Regev A, Myers CL, Berman J.
950   2014. YMAP: a pipeline for visualization of copy number variation and loss of
951   heterozygosity in eukaryotic pathogens. Genome Med 6:100.
952
953

Figure legends

Figure 1. Measuring tolerance levels at 30°C, 37°C, and 39°C identifies three classes temperature response.
Disk diffusion assays were performed on 133 different clinical isolates at the indicated temperatures (30°C, 37°C, and 39°C). The isolates were classified as temperature-enhanced tolerant (TET), all-temperature tolerant (ATT), and non-tolerant (NT), based on FoG_{20} values at the three temperatures. ** indicates p-value<0.001 as determined by two-tailed paired t-test (**A**). **B** is the distribution of tolerant isolates at different temperatures (n=133). At 30 °C, 69.2% were TET, 23.3% ATT, 7.5% NT. Thus, the vast majority of strains exhibit tolerance at least at some temperatures.

Figure 2. Temperature dependent growth dynamics of representative TET, ATT and NT strains in different drug concentrations.
Representative isolates (TET, SC5314; ATT, YJB-T1891; and NT, YJB-T490) were assayed on plates with FLC E-Test strips (A) and on 10-fold dilution spot assays (B) at 30°C, 37°C and 39°C on YPD medium at the FLC concentrations indicated. In both cases, plates were photographed after 48h at the indicated temperatures. Growth dynamics of isolates in the presence of FLC were measured using *ScanLag* (10, 15). Approximately 100 colonies of the test strains were plated onto YPD plates supplemented with FLC at the concentrations indicated (C). The growth of colonies was monitored at 30°C and 37°C using *ScanLag*. Colony area was calculated from the area of light pixels in the images, a proxy for colony size, and the change in colony size over time was used to estimate colony growth rate. Note in (B) and (C), for tolerant isolates, growth was similar at all fluconazole concentrations equivalent to and above MIC. The graphs show the distribution of Time of colony Appearance (TOA, top panel) and growth rate (bottom panel) for all colonies on each plate, in 72 hours (top panel). The area of colonies occupied by light pixels and the change in this area over time are proxies for colony size and growth rate.

Figure 3. Selection for adaptors at supra-MIC fluconazole concentrations.

FLC adaptors derived from TET strain SC5314 at 30˚C, and those derived from the NT

strain YJB-T490 at 30˚C and 37˚C, were tested for resistance (RAD_{20}) and tolerance

(Fog_{20}) on disk diffusion assays at 30˚C and 37˚C. The disks contained 25 µg FLC. The

plates were photographed at 24h and 48h for measuring RAD_{20} and FoG_{20}, respectively.

Each plot represents the data from 18 colonies that adapted to fluconazole

concentrations indicated in YPD plates. In the parents, 16 individual colonies were

tested.

Figure 4. Recurrent aneuploidies associated with resistant and tolerant

fluconazole adaptors.

(A) Karyotypes recurrently identified in TET and NT derived fluconazole adaptors

selected at 30˚C. The total number of aneuploid adaptors, and the number of adaptors

bearing each karyotype are also indicated. Chromosomes 1,2,3,5, which were euploid

in these karyotypes, are not shown. (B) Karyotypes of 9 adaptors derived from NT

isolate at 37˚C. None of the karyotypes was identified in (A). The grey colored

karyotypes include adaptors with aneuploidy of different homologs of the same

chromosomes. The detailed karyotypes are shown in Fig. S3 and Fig. S4.

1002

Figure 5. Tolerance acquisition and loss is associated with the gain and loss of

aneuploid chromosomes.

Strains with different aneuploid chromosomes were spread onto YPD plates and grown

for 36 h at 30˚C. Small colonies (S) and large colonies (L), indicated by yellow arrows

and cyan arrows, respectively (A) were tested by standard disc diffusion assays (25 µg

FLC per disk). The plates were incubated at 30˚C for 48 h (B). Both S and L colonies

were sequenced. The karyotypes were visualized by using Ymap (C).

1010

Figure 6. Distinct evolutionary trajectories associated with supra- and sub-MIC

fluconazole concentrations.

Emergence of tolerance (increased FoG_{20}) and resistance (decreased RAD_{20})

measured on disk diffusion assays for 18 isolates from each indicated passage of TET

isolate SC5314 (A) and NT isolate YJB-T490 (B) for 1-15 days of propagation at the

indicated drug concentrations. The MIC (1µg/ml) is highlighted in grey. Note that a

broad range of tolerance emerges at all supra-MIC concentrations starting at passage 1,

while increasing levels of resistance appear with time at or below the MIC only.

1019

Figure 7. Distinct genomic changes in resistant and tolerant adaptors.

DNA sequences of 34 FLC-resistant and 39 FLC-tolerant adaptors evolved from TET

strain SC5314 were analyzed by Ymap (49). The karyotypes, number of adaptors with

this karyotype. and the drug conditions (number of days evolved and FLC concentration)

are indicated to the right of each karyotype diagram. For example, D5-1 indicates that

the adaptors were evolved for 5 days in 1 µg/ml FLC. Colors indicate allele frequencies

in the data as indicated in the key. Black histograms indicate the log1 ratio of DNA copy

number for the strain indicated relative to a diploid control

1028

1029   **Table 1.** Karyotypes of the 34 resistant and 39 tolerant evolved progeny analyzed.

1030

 Table 1. Karyotypes of SC5314 adaptors from passaging experiment

Passage number (day)	Selection pressure FLC (µg/ml)	Adaptor response	Number of sequenced adaptors	Genome features*
1	2	Tolerance	5	ChrRx3 (n=3); ChrRx3+Chr4x3 (n=1); segChrRx1 (from 1.89 Mb to right telomere, n=1)
5	128	Tolerance	2	ChrRx3
	1	Tolerance	1	ChrRx3
		Resistance	4	Euploid
	0.25	Tolerance	1	ChrRx3
		Resistance	5	Euploid
10	0.5	Tolerance	3	SegChrRx3 (from left telomere to 1.89MB, n=2); ChrRx3 +Chr7x3+Chr4x3 (n=1)
		Resistance	3	Chr5x3
	1	Tolerance	3	ChrRx3+Chr4x3 (n=1); ChrRx3+Chr6x3 (n=1); Chr6x3 (n=1)
		Resistance	3	Euploid
	2	Tolerant	6	ChrRx3+Chr4x3 (n=4); ChrRx3+Chr4x3+Chr2x3 (n=1); ChrRx3+Chr4x3+Chr6x3 (n=1)
	128	Tolerance	6	ChrRx3+Chr4x3
15	0.25	Resistance	6	Euploid
	0.5	Resistance	6	Chr5x3 (n=2); Chr7x3 (n=1); Euploid (n=3)
	1	Resistance	6	Chr3x3 (n=2); Chr7x3 (n=3); Chr7x3+Chr6x3 (n=1)
	2	Tolerance	6	SegChrRx3 (from left telomere to 1.89 MB, n=2); ChrRx3 (n=2); ChrRx3+Chr4x3 (n=2)
	128	Tolerance	6	SegChrRx3 (from left telomere to 1.89 MB, n=1); ChrRx3+Chr4x3 (n=5)

1032 [*]See Fig. 7 and Fig. S6 for Ymap views of the karyotypes. Annotation such as "x3"

1033 indicates trisomy, "+" indicates additional aneuploidy in the same adaptor. n, number of

1034 isolates with this karyotype.

1035

1036 **Supplementary materials**
1037

1038 **Figure S1. Physiological factors affect fluconazole tolerance.**
1039 *C. albicans* lab strain SC5314 was tested on disk diffusion assays with drug disks (25
1040 µg FLC) on agar plates containing the indicated media and grown at the indicated
1041 temperatures. The plates were photographed at 48h to highlight tolerance (growth
1042 within the zone of inhibition.
1043

1044 **Figure S2. Growth of tolerant and non-tolerant strains on fluconazole plates.**
1045 Approximately 1x10^6 cells of TET (SC5314), ATT (YJB-T1891), and NT (YJB-T490)
1046 strains were grown on YPD plates supplemented with FLC at the indicated
1047 concentrations and incubated at 30˚C and 37˚C for 5 days. Estimated number of
1048 colonies on plates with 8 µg/ml~128 µg/ml drug plates are also shown.
1049

Figure S3. Karyotypes of all adaptors evolved from SC5314 at 30˚C.
SC5314-derived adaptors from YPD plates supplemented with 8 µg/ml, 32 µg/ml, and
128 µg/ml, were deep-sequenced and karyotype maps were generated by using the
Ymap. Each chromosome is illustrated to scale. The centromere positions are indicated
by an indentation. The major repeat sequence (MRS) positions are indicated by a black
square below the chromosome. Local copy number estimates, scaled to the genome
ploidy, are displayed as black histograms along the length of the chromosome. Allele
frequencies are color coded: homolog "a" is cyan, homolog "b" is magenta and
heterozygous alleles are grey. Karyotypes of all the adaptors sequenced are shown.
The order of the adaptors is the same as the order when they were picked up from the
fluconazole plates.
1061

1062 **Figure S4. Karyotypes of adaptors derived from NT strain YJB-T490 at 30˚C.**
1063 YJB-T490-derived adaptors chosen from YPD plates supplemented with 8 µg/ml, 32
1064 µg/ml, and 128 µg/ml at 30˚C were deep sequenced. Karyotypes of all the 54 adaptors
1065 sequenced are shown. Homozygosis of alleles is indicated in red.
1066

Figure S5. Stepwise increase in MICs for resistant adaptors evolved in 1 µg/ml

fluconazole.

On day 5, day 10, and day 15, the MIC of one resistant adaptor from the 1µg/ml cultures

was tested with FLC Etest strips. As a control, the parent and one adaptor isolated from

passage 1 were also tested. The plates were incubated at 30°C for 48h and then

photographed.

1073

1074　**Figure S6. Karyotypes of progeny evolved in YPD broth supplemented with**

1075　**fluconazole.**

1076　34 FLC-resistant and 39 FLC-tolerant progeny were sequenced. FLC concentration and

1077　timepoint of test are indicated above each section. Strain name and whether it exhibited

1078　resistance (R) or tolerance (T) are indicated to the left of each karyotype.

1079

Table S1: Mutations detected in coding sequences in 5 euploid fluconazole-tolerant

adaptors.

Table S2: Mutations detected in coding sequences in 6 pairs of small and large

colonies.

Table S3: ORFs having mutations identified in the 34 fluconazole-resistant and the 39

fluconazole-tolerant adaptors.

Table S4: Strains used in this study.

1087
1088

A

TET parent selected at 30°C

Chr4 Chr6 Chr7 ChrR   # of adaptors (n=50)

NT parent selected at 30°C

# of adaptors (n=52)   Chr4 Chr6 Chr7 ChrR

B

NT parent selected at 37°C

Chr1   Chr2   Ch4   Chr5   Chr6   Chr7   ChrR   # of adaptors (n=9)

FLC-resistant adaptors (n=34)

	Chr1	Chr2	Chr3	Chr4	Chr5	Chr6	Chr7	ChrR	NO. of adaptors	Condition of evolution (FLC (μg/ml)-Passage)
									22	1-P5; 0.25-P10; 1-P10; 0.25-P15, 0.5-P15, 1-P15
									1	1-P15
									1	1-P15
									5	0.5-P10; 0.5-P15
									4	0.5-P15; 1-P15
									1	1-P15

FLC-tolerant adaptors (n=39)

	Chr1	Chr2	Chr3	Chr4	Chr5	Chr6	Chr7	ChrR		
									4	2-P1; 1-P5; 0.25-P10
									5	2-P1; 128-P1; 2-P15
									1	2-P1
									2	2-P15; 128-P15
									1	2-P15
									2	0.5-P10
									4	1-P10; 2-P15; 128-P15
									5	2-P1; 2-P10; 128-P10; 2-P15; 128-P15
									4	2-P10; 128-P10; 128-P15
									6	128-P10; 128-P15
									1	2-P10
									1	2-P10
									1	0.5-P10
									1	1-P10
									1	1-P10