1
2

Tethering by Uso1 is dispensable: The Uso1 monomeric globular head domain interacts with SNAREs to maintain viability.

Ignacio Bravo-Plaza[1], Víctor G. Tagua[2], Herbert N. Arst, Jr [3], Ana Alonso[1], Mario Pinar[1], Begoña Monterroso[4], Antonio Galindo[5], and Miguel Á. Peñalva[1, $]

Departments of (1) Cellular and Molecular Biology and (4) Structural and Chemical Biology, CSIC Centro de Investigaciones Biológicas, Ramiro de Maeztu 9, 28040 Madrid Spain.

(2) Instituto de Tecnologías Biomédicas, Hospital Universitario Nuestra Señora de Candelaria, Santa Cruz de Tenerife, Spain.

(3) Department of Infectious Diseases, Faculty of Medicine, Flowers Building, Imperial College, Armstrong Road, London, SW7 2AZ, UK

(5) Division of Cell Biology, MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge, CB2 0QH, UK.

($) Corresponding author at the above address

penalva@cib.csis.es
@penalva

phone: +918373112

Summary

Uso1/p115 and RAB1 tether ER-derived vesicles to the Golgi. Uso1/p115 contains a
globular-head-domain (GHD), a coiled-coil (CC) mediating dimerization/tethering and a
C-terminal region (CTR) interacting with golgins. Uso1/p115 is recruited to vesicles by
RAB1. Paradoxically, genetic studies placed Uso1 acting upstream of, or in conjunction
with RAB1 (Sapperstein et al., 1996). We selected two missense mutations in *uso1*
resulting in E6K and G540S substitutions in the GHD permitting growth of otherwise
inviable *rab1*-deficient *Aspergillus nidulans.* Remarkably, the double mutant suppresses
the complete absence of RAB1. Full-length Uso1 and CTRΔ proteins are dimeric and the
GHD lacking the CC/CTR is monomeric irrespective of whether they carry or not
E6K/G540S. Microscopy showed recurrence of Uso1 on puncta (60 sec half-life)
colocalizing with RAB1 and less so with early Golgi markers Sed5 and
GeaA/Gea1/Gea2. Localization of Uso1 but not of Uso1$^{E6K/G540S}$ to puncta is abolished
by compromising RAB1 function, indicating that E6K/G540S creates interactions
bypassing RAB1. By S-tag-coprecipitation we demonstrate that Uso1 is an associate of
the Sed5/Bos1/Bet1/Sec22 SNARE complex zippering vesicles with the Golgi, with
Uso1$^{E6K/G540S}$ showing stronger association. Bos1 and Bet1 bind the Uso1 GHD directly,
but Bet1 is a strong E6K/G540S-independent binder, whereas Bos1 is weaker but
becomes as strong as Bet1 when the GHD carries E6K/G540S. AlphaFold2 predicts that
G540S actually increases binding of GHD to the Bos1 Habc domain. In contrast, E6K
seemingly increases membrane targeting of an N-terminal amphipathic α-helix,
explaining phenotypic additivity. Overexpression of E6K/G540S and wild-type GHD
complemented *uso1Δ*. Thus, a GHD monomer provides the essential Uso1 functions,
demonstrating that long-range tethering activity is dispensable. Therefore, when
enhanced by E6K/G540S, Uso1 binding to Bos1/Bet1 required to regulate SNAREs
bypasses both the contribution of RAB1 to Uso1 recruitment and the reported role of
RAB1 in SNARE complex formation (Lupashin and Waters, 1997), suggesting that the
latter is consequence of the former.

Introduction

Vesicular traffic at the ER/Golgi interface is the cornerstone of the secretory pathway
(Barlowe and Miller, 2013; Weigel et al., 2021). In current models, in which traffic across
the Golgi is driven by cisternal maturation (Day et al., 2013; Pantazopoulou and Glick,
2019), COPII vesicles generated at specialized domains of the ER fuse homotypically
and heterotypically to form and feed the earliest Golgi cisternae (Rexach et al., 1994).
As straightforward as this step might seem, it involves a sophisticated circuitry of
regulation. Actual fusion is in part mediated by compartmental-specific sets of four-
membered SNARE protein complexes (SNARE bundles) (Malsam and Sollner, 2011;
Pelham, 2001; Rizo and Sudhof, 2012). Most SNARES are type II single TMD proteins,
whose N-terminal cytosolic domain contains nearly all the polypeptide, excepting a few
lumenal residues. Like any other transmembrane proteins, SNAREs are synthesized in
the ER. This implies that they have to travel to compartments of the cell as distant as the
plasma membrane in a conformation that precludes them of catalyzing what would be a
calamitous fusion of non-cognate donor and acceptor compartments. Achieving the
strictest specificity is particularly challenging in the ER-to-Golgi stage that, as the first
step in the secretory pathway, represents an obligate point of transit for each and every
transmembrane SNARE. Therefore, the only SNARES acting in this first step are the Qa
Sed5, the Qb Bos1, the Qc Bet1 and the R-SNARE Sec22, which form the bundle
mediating fusion of carriers that coalesce into cisternae (McNew et al., 2000).

Given the central role played by the secretory pathway in the physiology of every
eukaryotic cell, it is unsurprising that this step involves regulatory factors which are
essential for cell survival. One is the SM (Sec1, Munc-18) protein Sly1, which promotes
SNARE bundle formation (Bracher and Weissenhorn, 2002; Peng and Gallwitz, 2002;
Thomas et al., 2019) Another is the TRAPPIII complex, which interacts with the external
coat of COPII carriers and acts as a guanine nucleotide exchange factor (GEF) for RAB1
(Bracher and Weissenhorn, 2002; Cai et al., 2007; Galindo et al., 2021; Joiner et al.,
2021; Lord et al., 2011; Peng and Gallwitz, 2002; Pinar and Peñalva, 2020; Riedel et al.,
2017; Thomas et al., 2018; Thomas et al., 2019). This small GTPase is a key player that
transiently recruits protein effectors from the cytosol to donor and acceptor membranes
(Sogaard et al., 1994) and regulates SNARE assembly through an as yet undefined
mechanism (Lupashin and Waters, 1997; Sapperstein et al., 1996). One RAB1 effector
is a fungal protein denoted Uso1, whose highly conserved metazoan homologue is p115.
These are homodimers with a globular N-terminal head and a long C-terminal coiled-coil
region characteristic of tethering proteins, which bring donor and acceptor membranes
into the distance at which v- and t-SNAREs can engage into the productive *trans*-SNARE

complex that mediates membrane fusion (Cao et al., 1998; Nakajima et al., 1991;

Sapperstein et al., 1996; Sapperstein et al., 1995; Seog et al., 1994; Yamakawa et al.,

1996).

Despite most Golgi tethers are functionally redundant, Uso1 is unique in that it is an

essential protein. *uso1-1*, a *S. cerevisiae* amber mutation truncating most, but not all the

coiled-coil region, is viable, yet further upstream truncation removing the complete

coiled-coil is lethal, which was taken as evidence that tethering is the essential function

of Uso1 (Seog et al., 1994). In addition, as the coiled-coil is predicted to mediate

dimerization, it is broadly accepted that Uso1 is 'just' an essential homodimer that tethers

vesicles to the acceptor membrane. However, genetic evidence stubbornly indicates that

Uso1 plays additional functions related with SNAREs. For example, Sapperstein et al

showed that SNAREs function downstream of Uso1 (Sapperstein et al., 1996). Notably,

the view that Uso1 is a mere RAB1 effector was challenged by the observation that

overexpressing Ypt1 (yeast RAB1) rescues lethality of *uso1Δ*, whereas the reciprocal is

not true, indicating that Uso1 acts upstream of or in conjunction with RAB1 (Sapperstein

et al., 1996).

Our laboratory is interested in deciphering the domains of action of RAB GTPases in the

genetic and cell biological model organism *Aspergillus nidulans* (Pinar and Peñalva,

2021). We have previously gained mechanistic insight into the activation of RAB11 by

TRAPPII by exploiting a forward genetic screen for mutations bypassing, at the restrictive

temperature, the essential role of the key TRAPPII subunit Trs120 (Pinar et al., 2019;

Pinar et al., 2015; Pinar and Peñalva, 2020). In this type of screen, a strain carrying a *ts*

mutation in the gene-of-interest is mutagenized and strains bypassing lethality at the

restrictive temperature are identified and characterized molecularly. A well-

characterized, conditionally lethal *rab1* mutation is available (Pinar et al., 2013), enabling

us to investigate pathways collaborating with RAB1 in anterograde traffic. We isolated

two *uso1* missense mutations causing substitutions in the globular head domain (GHD).

When combined together, these rescued the lethality resulting from *rab1Δ* and promoted

the localization of the protein to early Golgi cisternae by increasing Uso1 binding to the

cytosolic region of the Qa SNARE Bos1 and, potentially, by improving the interaction of

an N-terminal amphipathic α-helix with membranes. Importantly, we show that

endogenous expression of a protein consisting solely of the double mutant GHD, or

overexpression of double mutant or wild-type GHD, rescue the lethality resulting from

uso1Δ, even though the GHD is monomeric. Our results show that one essential role of

RAB1 is recruiting Uso1 to membranes, and that the essential role of Uso1 is not

tethering membranes, but rather regulating the formation of the cognate SNARE bundle,

indicating that Uso1 is a component of the SNARE fusion machinery .

Results

Missense mutations in *uso1* rescue the lethality resulting from *rab1*Δ

rab1^{A136D} (hereby *rab1^{ts}*) mutants do not grow at 37ºC. However, when we plated UV-

mutagenized conidiospores of the *rab1^{ts}* mutant at this temperature, we obtained

colonies showing different degrees of growth, presumably carrying mutations rescuing

the lethality resulting from *rab1^{ts}*. One was chosen for further characterization. By sexual

crosses and parasexual genetics this strain was shown to carry a single suppressor

mutation, denoted *su1rab1^{ts}*, that co-segregated with chromosome VIII. Meiotic mapping

narrowed *su1rab1^{ts}* to the vicinity (2 cM) of *hisC*. 40 kb centromere distal from *hisC* lies

AN0706 (Figure 1A) encoding *Aspergillus nidulans* Uso1, a conserved effector of RAB1.

Sanger sequencing revealed the presence of a G16A transition (denoted E6K) resulting

in Glu6Lys substitution in the *uso1* gene of *su1rab1^{ts}*.

To determine if the remaining suppressor strains were allelic to *su1rab1^{ts}*, we sequenced

uso1 from a further 13 isolates (Figure 1B). Of these, four were *rab1^{ts}* pseudo-revertants

that had acquired a functionally acceptable mutation in the altered codon, and eight

carried *uso1* E6K, suggesting that the screen was close to saturation. However, one

mutation was found to be a different missense allele, *su85rab1ts* (denoted G540S)

resulting in Gly540Ser substitution. Single mutant strains carrying these *uso1* mutations

showed no growth defect, indicating that E6K and G540S were unlikely to result in loss-

of-function, and suggesting instead that mutant strains had acquired features that made

them largely independent of RAB1. These findings were unexpected, because in

Saccharomyces cerevisiae overexpression of Uso1 does not rescue the lethality of *ypt1*Δ

mutants (Ypt1 is the yeast RAB1 homologue)(Sapperstein et al., 1996).

To demonstrate that *uso1^{E6K}* and *uso1^{G540S}* were causative of the suppression, we

reconstructed them by homologous recombination. These reverse-genetic alleles

rescued viability of *rab1^{ts}* at 37ºC to a similar extent as *su1rab1^{ts}* and *su85rab1^{ts}* (Figure

1C). *uso1^{G540S}* was the strongest suppressor, such that *rab1^{ts}* *uso1^{G540S}* double mutants

grew nearly as the wt at 37ºC. Nevertheless, the two alleles showed additivity, and a

triple mutant carrying *uso1^{E6K}*, *uso1^{G540S}* and *rab1^{ts}* grew at 42ºC, unlike either single

mutant (Figure 1C). These data, together with the genetic mapping above, established

that *uso1^{E6K}* and *uso1^{G540S}* are responsible for the suppression phenotype,

RAB1 recruits Uso1/p115 to uncoated COPII vesicles and early Golgi cisternae (Allan et

al., 2000). Therefore, $uso1^{E6K/G540S}$ might, by increasing the affinity of Uso1 for RAB1,

compensate for the reduction in the amount of the GTPase resulting from $rab1^{ts}$.

However, Figure 1D shows that both $uso1^{E6K}$ and $uso1^{G540S}$ rescue the lethality resulting

from the complete ablation of $rab1\Delta$ at 30ºC, with the strongest $uso1^{G540S}$ suppressor

rescuing viability even at 37ºC, and the double mutant rescuing $rab1\Delta$ even at 42ºC

(Figure 1D). In contrast, $uso1^{E6K/G540S}$ did not rescue the lethality resulting from $arf1\Delta$, nor

from $sed5\Delta$ or $sly1\Delta$, the syntaxin and the SM protein which are crucial for the formation

of the ER/Golgi SNARE bundle (Figure 1—figure supplement 1), indicating that Uso1

plays a role acting downstream of RAB1 and upstream of or in conjunction with the

SNARE machinery. This role is essential for survival (Figure 1—figure supplement 1).

E6K affects a previously undetected N-terminal helix, whereas G540S is located

in a loop near the end of the armadillo domain.

1103-residue *A. nidulans* Uso1 is similar in size to 961-residue p115 (bovine) and notably

shorter than *S. cerevisiae* Uso1p (1790 residues) (Yamakawa et al., 1996). Thus far,

atomic structures of Uso1/p115 are limited to the 600-700 residue GHD, which consists

of a highly conserved α-catenin-like armadillo-fold (An et al., 2009; Heo et al., 2020;

Striegl et al., 2009). *In silico* analyses robustly predict that the approximately C-terminal

half of Uso1/p115 consists of a coiled-coil that mediates tethering and dimerization, but

this region has not been characterized beyond low resolution EM studies (Yamakawa et

al., 1996) . Neither crystal structures nor predictions provided information about the N-

terminal extension in which Glu6Lys lies.

Thus, we used AlphaFold2, imposing or not the condition that the protein is a dimer (see

below). Figure 2A shows the monomer and dimer models with the highest confidence

scores (see Figure 2—figure supplement 1). Like their relatives, Uso1 from *A. nidulans*

contains an N-terminal GHD including a previously unnoticed short α-helix in which Glu6

affected by E6K lies. This N-terminal extension is followed by ~34 α-helices arranged

into 12 tandem repetitions of armadillo repeats (ARM1-ARM12; residues 17 through

564), each containing three right-handed α-helices except for the first two repeats.

Altogether, the armadillo repeats resemble the shape of a jai alai basket. Downstream

of the GHD, AlphaFold2 predicts a long extended coiled-coil (CC) between residues 674

through 1082, which would mediate dimerization (see below) (Figure 2A). The CC ends

at a conserved C-terminal region (CTR) (Figure 2A and B), which includes an also C-
terminal segment rich in acidic residues. In Uso1, twelve out of the last seventeen amino
acids are Asp/Glu (Figure 2A and B). The CC and CTR regions will be collectively
denoted the CCD domain.

Even though the GHD contains Glu6 and Gly540 in the N-terminal helix and at the
beginning of armadillo α-helix 29, respectively, intramolecular or intermolecular (in the
context of a homodimer, see below) distances between these residues are long, arguing
against the possibility that they bind a common target as components of the same
interaction surface (Figure 2C). Indeed, the synthetic positive effect of the mutations
would be consistent with their rescuing viability through different mechanisms. The
previously unnoticed short α-helix predicted by AlphaFold between Phe2 and Lys12 is
amphipathic (Figure 2D). Glu6 lies on the polar side of this helix, such that Glu6Lys
increases its overall positive charge (three of the four polar residues are Lys or Arg). As
discussed below, it is tempting to speculate that this helix facilitates membrane
recruitment.

Coiled-coil mediated dimerization of Uso1: the globular head is monomeric
231   It has been proposed that p115 alternates between closed and open
conformations to hide or expose a RAB1 binding site present in the CCD (Beard et al.,
2005). This would be mediated by intramolecular interactions between the globular
domain and the C-terminal acidic region, which would be disrupted by the competitive
binding of golgins GM130 and giantin to the latter. We addressed whether E6K/G540S
promotes a conformational change in Uso1, or, alternatively, a change in the
oligomerization status of the protein, by analytical ultracentrifugation. We designed
seven constructs carrying a C-terminal His tag (Figure 3). Two corresponded to the full-
length protein with or without E6K and G540S substitutions. The second pair included
wild-type and doubly-substituted versions of C-terminally truncated Uso1 lacking the
CTR (Uso1ΔCTR and Uso1$^{E6K/G540S}$ΔCTR). The third corresponded to wild-type and
doubly-substituted versions of the globular domain, denoted Uso1 GHD and Uso1
GHD$^{E6K/G540S}$. The seventh construct corresponded to the CCD domain (i.e. CC + CTR)
and was denoted Uso1 CCD. All seven proteins were expressed in bacteria, purified by
nickel-affinity and size-exclusion chromatography and analyzed by sedimentation
velocity ultracentrifugation. These experiments revealed that all protein preparations
were essentially homogeneous, and thus they were used to determine the corresponding
Svedberg coefficients. In addition, by dynamic light scattering we determined the

translational diffusion coefficients of the constructs. With these values we deduced the
molecular mass of the different proteins using Svedberg's equation.

Wild-type and E6K/G540S full-length Uso1s showed the same sedimentation
coefficients, demonstrating that the mutations do not induce a large conformational
change that would have been reflected in changes in sedimentation velocity due to
differences in frictional forces. Molecular masses deduced from the Svedberg equation
indicated that these full-length proteins are homodimers, in agreement with previous
literature (Figure 3). Ablation of the conserved C-terminal region (CTR) did not result in
any significant change in the sedimentation coefficient (Figure 3, panels 3 and 4 vs. 1
and 2), irrespective of the presence or absence of the substitutions, discarding the model
in which the CTR would interact with the GHD to maintain a hypothetical closed
conformation (Beard et al., 2005). In addition, the molecular masses of the ΔCTR
proteins correspond to a dimer, implying that the acidic region is not involved in
dimerization either.

Notably the GHD, whether wild-type or mutant, behaved as a monomer (Figure 3, panels
5 and 6), which has important implications described below. In contrast, the coiled-coil
domain, with a predicted molecular mass of 52 kDa, behaves as a dimer of *ca.* 100 kDa
(Figure 3, panel 7). The sedimentation coefficient of the CCD is markedly slower than
that of the 70 kDa monomeric GHD, suggesting an elongated shape. These
observations, together with the dimeric nature of the construct lacking the CTR, showed
that dimerization is mediated by the CCD. The absence of 443 residues corresponding
to the CCD plus CTR domains in the GHD construct and the monomeric nature of the
latter compared to full-length Uso1 dimer did not result in a commensurate decrease in
sedimentation coefficient, which changed from 4.8 S to 3.7 S in the wild-type (note that
the change in Mr goes from 246 kDa in full-length Uso1 to only 68 kDa of the
GHD)(Figure 3). These data strongly support AlphaFold2 predictions depicting Uso1 as
a dimer with a globular head and an extended coiled-coil that would retard sedimentation
of the protein very substantially.

As with full-length Uso1, the double substitution did not alter the sedimentation
coefficient of the GHD (Figure 3, panels 5 and 6). To buttress the conclusion that the
GHD is a monomer irrespective of the presence or absence of the mutations, we
performed sedimentation velocity experiments using different protein concentrations
ranging from 0.5 to 5 μM (Figure 3—figure supplement 1, A and B). In all cases the GHD
behaved as a monomer. Sedimentation profiles of Uso1 GHD lacking the His-tag showed

a similar behavior, establishing that the monomeric state of the mutant is not due to the
tag at the C-terminal position hindering dimerization (Figure 3—figure supplement 1,C).
Therefore, sedimentation experiments did not detect any change in tertiary or quaternary
structures between wild-type and mutant GHD, which is important for the interpretation
of genetic data that will be discussed below.

In summary, (i) Uso1 is a dimer; (ii) The C-terminal acidic region is dispensable
for dimerization and does not mediate an equilibrium between closed and open
conformations; (iii) The globular domain of Uso1 is a monomer; (iv) The coiled-coil
domain of Uso1 is a dimer; (v) the double E6K G540S substitution does not promote any
conformational shift in Uso1, nor does it result in a change in the oligomerization state
of the protein.

The punctate pattern of localization of USO1-GFP is dependent on RAB1

The membranous compartments of the Golgi are not generally stacked in fungi,
permitting the resolution of cisternae, which appear as punctate structures in different
steps of maturation, by wide-field fluorescence microscopy (Losev et al., 2006;
Matsuura-Tokita et al., 2006; Pantazopoulou and Peñalva, 2011; Pinar et al., 2013;
Wooding and Pelham, 1998). While Uso1 is predicted to localize to the Golgi, studies of
its localization in fungi are limited (Cruz-García et al., 2014; Sánchez-León et al., 2015).
Therefore, we tagged the *A. nidulans uso1* gene endogenously with GFP

Figure 4A and video 1 depicting a software-shadowed 3D reconstruction of a
Uso1-GFP hypha, as well as consecutive sections of deconvolved z-stacks in Figure 4B
show that Uso1-GFP localizes to puncta polarized towards the tip, often undergoing
short-distance movements (see Figure 4C and Figure 4—figure supplement 1). These
puncta are smaller and more abundant than those reported for other markers of the
Golgi, which suggested that they might represent domains rather than complete
cisternae. Notably, 3D (x, y, t) movies revealed that Uso1 puncta are transient,
recurrently appearing and disappearing with time (Figure 4C). That this recurrence did
not reflect that the puncta go in-and-out of focus was established with 4D (x, y, z, t)
movies, which revealed a similar behavior of Uso1 irrespective of whether 3D or 4D
microscopy was used (video 2) Uso1 foci with confidence, we constructed movies with
middle planes only (i.e. 3D x, y, t series). After careful adjustment of live imaging
conditions, we achieved a 2 fps time resolution with relatively low bleaching for time
series consisting of 400 photograms (video 3). These conditions sufficed to track Uso1
puncta over time using kymographs traced across linear ROIs covering the complete
width of the hyphae (Figure 4C). However, as the abundance of Uso1 puncta made
automated analysis of Uso1 maturation events troublesome, we tracked them manually

with the aid of 3D (x,y,t) representations generated with Imaris software combined with
direct observation of photograms in movies (Figure 4C and Figure 4—figure supplement
1). The boxed event magnified in Figure 4E (see video 4) illustrates a prototypical
example. The right Figure 4E montage shows frames corresponding to this event for
comparison. We analyzed n = 60 events, which gave an estimation of the average half-
life of Uso1 residing in puncta of 60 sec +/- 25.26 S.D. (Figure 4D).

Nakano and co-workers have proposed that the transfer of lipids and proteins between
ER exit sites (ERES) and the early Golgi occurs through a kiss-and-run mechanism
(Kurokawa et al., 2014). Because Uso1-GFP punctate structures resemble, in size and
abundance, ER exit sites labelled with COPII components, we studied Uso1-GFP cells
co-expressing Sec13 endogenously labeled with mCherry (Bravo-Plaza et al., 2019).
The maximal intensity projection (MIP) shown on Figure 5A, and video 5 show that the
two markers are closely associated, but only in a few instances they showed
colocalization. These examples did not represent simple overlap, as they were found to
colocalize in the Z dimension using orthogonal views or montages (Figure 5B and C).
These observations have not been pursued further with time-resolved sequences, but at
the very least we can conclude that the reporters are closely associated in space. In view
of this, we determined that Uso1 structures originate downstream of COPII-mediated ER
exit. Therefore, we investigated, using *sarA*6, a temperature-sensitive allele of the gene
encoding *A. nidulans* SAR1 (Hernández-González et al., 2014), whether the punctate
Uso1 structures are dependent on this master GTPase regulating COPII biogenesis.
Figure 5D shows that this is indeed the case; the number of Uso1-GFP puncta was
significantly reduced relative to the wt when cells were shifted from 28°C to 37°C,
indicating that Uso1 populates a membrane compartment with Golgi identity.

To determine the 'sub-Golgi' localization of Uso1 puncta, we filmed Uso1-GFP along with
different Golgi markers (Figure 6). Uso1-GFP showed no overlap (Pearson's coefficient
0.17 ± 0.06 S.D., n = 16 cells) with cisternae labeled with mCherry-Sec7, the late Golgi
ARF1 GEF that is a prototypic marker of the TGN (Arst et al., 2014; Day et al., 2018;
Galindo et al., 2016; Halaby and Fromme, 2018; Losev et al., 2006; McDonald and
Fromme, 2014; Pantazopoulou, 2016; Pantazopoulou and Glick, 2019; Richardson et
al., 2016; Richardson et al., 2012) (Figure 6; video 6). In contrast, visual observation of
cells expressing mCh-Sed5 and Uso1-GFP revealed substantial, yet incomplete, overlap
of the reporters (Figure 6), reflected in a Pearson's coefficient of 0.44 ± 0.04 S.D., n =
15 cells (Figure 6). The Qa syntaxin Sed5 drives fusion of COPII vesicles with early Golgi
cisternae, with Qb, Qc and R-SNAREs Bet1, Bos1 and Sec22, and mediates intra-Golgi

trafficking, with Qb, Qc and R-SNARES Sft1, Gos1 and Ykt6), respectively (Banfield et
al., 1995; McNew et al., 2000; Parlati et al., 2002; Pelham, 1999; Wooding and Pelham,
1998). These data suggest that Uso1 localizes to a subset of early Golgi
cisternae/membranes containing Sed5. GeaA^{Gea1,2} is the only *A. nidulans* homologue of
the *S. cerevisiae* early Golgi ARF1 GEFs Gea1 and Gea2 dwelling at the early Golgi
(Arst et al., 2014; Gustafson and Fromme, 2017; Muccini et al., 2022; Pantazopoulou,
2016; Park et al., 2005; Wright et al., 2014). Overlapping of Uso1 with GeaA^{Gea1,2} was
more conspicuous than with Sed5 (Figure 6), which was reflected in an increased
Pearson's coefficient to 0.52 ± 0.06 S.D., *n* = 16 cells. Of note, mammalian GeaA (GBF1)
and Uso1 (p115) interact (Garcia-Mata and Sztul, 2003). Taken together, these data
indicate that Uso1 localizes to Golgi cisternae in early stages of maturation.

In both mammalian cells and in yeasts Uso1/p115 has been shown to be recruited to
early Golgi membranes by RAB1, which is activated by the TRAPPIII GEF on COPII
vesicles after they bud from the ER (Allan et al., 2000; Cai et al., 2007; Lord et al., 2011;
Yuan et al., 2017). Therefore, we imaged Uso1 and RAB1, which revealed that indeed
Uso1 colocalized with RAB1 (Pearson's 0.61 ± 0.07 S.D., *n* = 20 cells) (Figure 6).
Altogether, the above microscopy data strongly suggest that Uso1 is transiently recruited
to early Golgi membrane domains enriched in RAB1, agreeing with the accepted view
that RAB1 acts by recruiting Uso1.

Uso1 delocalization after RAB1 impairment rescued by E6K G540S.

We next tested if the subcellular localization of Uso1 is dependent on RAB1, and
if this dependency can be bypassed by E6K/G540S. To this end we first showed that in
a *RAB1*⁺ background endogenously tagged wild-type and E6K/G540S Uso1-GFP have
the same punctate localization pattern (Figure 7A), and that both supported vigorous wt
growth (Figure 7B lanes 1, 3, 5 and 7), indicating that the tagged proteins are functional.
Next, we introduced in these strains *rab1^{ts}* (Jedd et al., 1995; Pinar et al., 2013) by
crossing. This allele, which completely prevents growth at 37ºC, is a hypomorph at
permissive (25-30ºC) temperatures, which permitted testing RAB1 dependence under
standard microscopy conditions (28ºC). Figure 7A shows that wt Uso1-GFP was largely
delocalized to the cytosol by *rab1^{ts}*. This establishes that Uso1 localization to membranes
is subordinated to RAB1. Notably, the wt punctate pattern of Uso1 localization was
restored by the Uso1 E6K/G540S mutant substitution, correlating with correction of the
synthetic growth defect (Figure 7B, compare lanes 2, 4, 6 and 8). All these data indicated
that the localization of Uso1 is compromised when RAB1 function is impaired, and that

the E6K/G540S substitutions augment Uso1 affinity for a membrane anchor(s)

independent of RAB1. They additionally suggest that the principal physiological role of

RAB1 is ensuring the proper localization of Uso1.

Delocalization of Uso1 in the *rab1^ts* background was not solely dependent on RAB1.

Wild-type *uso1-GFP* displayed a synthetic negative interaction with *rab1^ts* (Figure 7B,

compare lanes 2 and 4 at 30ºC), suggesting that the presence of GFP in the C-terminus

interferes with a RAB1-independent mechanism that facilitates its recruitment to

membranes (see below)

Genetic evidence that a network involving the CTR and the Grh1/Bug1 golgin

contributes to the recruitment of Uso1 to membranes.

To follow up the above observation, we focused on golgins. In mammalian cells, the C-

terminal region of p115 interacts with GM130, a golgin which is recruited to the early

Golgi by GRASP65 (Beard et al., 2005). The equivalent proteins in budding yeast are

denoted Bug1 and Grh1 (Behnia et al., 2007)(Figure 8A).

GRASP65 contains two C-terminal PDZ [post synaptic density protein (PSD95),

Drosophila disc large tumor suppressor (Dlg1), and zonula occludens-1 protein (zo-1)]

domains, which bind an also C-terminal peptide in GM130 (Hu et al., 2015).. Similar to

its metazoan counterparts, *Aspergillus* Grh1 contains an N-terminal α-helix and two PDZ

domains, in this case followed by ~130 disordered residues (Figure 8B, Figure 8—figure

supplement 1A and C). We modelled the Grh1-Bug1 interaction using AlphaFold2.

Residues 666-675 of the C-terminal peptide of Bug1 bind to a hydrophobic cleft located

between PDZ1 and PDZ2. Bug1 Leu668 and 670 coordinate their side chains with

residues from both PDZ domains (e.g. Phe45 and Trp44 in PDZ1 and Trp171 and Val179

in PDZ2). A second interaction involves Bug1 C-terminal residues 683-690 fitting within

a second groove in PDZ1, such that the four C-terminal residues form a β-strand

extending the β-sheet of N-terminal PDZ1 domain (Figure 8B, Figure 8—figure

supplement 1B). Further genetic evidence that a network of interactions similar to that

acting in yeast and mammalian cells operates in the *A. nidulans* ER-Golgi interface was

obtained by constructing strains with combinations of gene-replaced alleles. These

consisted of *uso1ΔCTR* encoding Uso1 lacking the C-terminal region (residues 1-1041)

and containing or not E6K G540S, *rab1^ts*, and deletion alleles of the *Aspergillus BUG1*

(AN7680) and *GRH1* (AN11248) genes (Figure 8A). Combining *rab1^ts* with *uso1ΔCTR*

resulted in a synthetic negative interaction at 30ºC akin to that seen with *uso1-GFP*

(lanes 2 and 4 in Figure 8C). That the E6K/G540S double substitution rescued this

negative interaction strongly indicates that the CTR cooperates with RAB1 in the

recruitment of Uso1 to membranes (Figure 8C, lanes 2, 4, 6 and 8). We note that the

control wild-type strain used in these experiments contains a construct completely

analogous to the mutant allele, ruling out that the genetic manipulation (for example the

introduction, linked to the *uso1* locus, of a selection marker, potentially chromatin-

disruptive) is causative of the observed phenotype. Another trivial explanation that we

ruled out by Western-blot analysis was that the deletion off the 62 C-terminal residues in

uso1ΔCTR resulted in increased degradation, which it did not (Figure 8D).

If interactions involving the CTR of p115 were conserved in fungi, *BUG1* (GM130

equivalent) and *GRH1* (GRASP65) should also show a synthetic negative interaction

with *rab1ts*. Figure 8E (lanes 1,3 and 5) shows that neither *bug1Δ* nor *grh1Δ* affects

growth. However, both deletion alleles showed a strong synthetic negative interaction

with *rab1ts* (Figure 8E, lanes 1,3 and 5). Remarkably, the synthetic negative phenotype

was rescued by the presence of E6K/G540S substitutions in Uso1, further suggesting

that they promote Uso1 recruitment to its locale of action, compensating for the loss of

the Bug1-Uso1 CTR interaction. We conclude that interaction involving the CTR of Uso1

and the BUG1/GRH1 complex cooperates with RAB1-mediated mechanisms to recruit

Uso1 to membranes.

The globular head domain (GHD) of Uso1 carrying the double E6K/G540S

substitution supports cell viability

Uso1 has traditionally been considered the archetype of a coiled-coil tether

recruiting ER-derived vesicles to the *cis*-Golgi. In *S. cerevisiae*, *uso1-1* (Sapperstein et

al., 1996), is a C-terminally truncating, conditionally-lethal *ts* allele whose encoded

protein still retains 20% of the coiled-coil region. In contrast, *uso1-12* and *uso1-13*

removing the complete coiled-coil region are lethal (Seog et al., 1994). Thus, we asked

whether the double E6K/540S substitution bypasses the requirement of the coiled-coil

region. To this end, we constructed, by gene replacement, a *uso1GHD* allele expressing

a protein truncated immediately after the GHD, lacking residues 660-1103. By

heterokaryon rescue, we demonstrated that this allele is lethal (Figure 9A).

Unexpectedly, given that the *rab1ts* suppressor substitutions lie outside the coiled-coil

region, the equivalent allele containing E6K/G540S sufficed for the fungus to survive at

30ºC. This result has two key implications: that the coiled-coil of Uso1 is not essential for

survival and as the GHD of Uso1 is a monomer in solution, it follows that dimerization of

Uso1 is not essential either (see also below).

In view of these unexpected results we wondered whether the structure of the GHD

synthesized in bacteria differed from the physiological form in *Aspergillus*, such that the

GHD were a dimer *in vivo.* To address this possibility, we ran an *Aspergillus* cell-free

extract expressing HA3-tagged Uso1 GHD through a Sepharose column. As control, we

ran in parallel a sample of bacterially-expressed His-tagged GHD used in sedimentation

velocity experiments. Western blot analysis of the fractions (Figure 9B) demonstrated

that both proteins eluted at the same position, corresponding to that expected for a

globular protein with the size of the GHD. As controls we ran in the same column a similar

pair of proteins corresponding to full length Uso1. The bacterially-expressed and the

native Uso1 proteins also co-eluted (Figure 9B), but this time at a position corresponding

to a highly elongated dimer, consistent with sedimentation velocity experiments. Thus,

in vivo, the GHD, expressed from its own promoter, is a monomer, and it is sufficient to

sustain viability if it carries the double mutant substitution that bypasses the requirement

for RAB1. These data strongly argue against tethering being *the* essential physiological

role that Uso1 plays.

We next investigated why the E6K/G540S GHD suffices for viability only at 30°C. Often

thermo-sensitivity results from protein instability, which is enhanced at high

temperatures. Western blot analysis of the allele-replaced strain expressing E6K G540S

GHD as the only source of Uso1 revealed that levels of the truncated Uso1 mutant were

minuscule relative to the wt or to the equivalent Δ*CTR* allele (Figure 9C). We reasoned

that increasing expression would result in E6K/G540S GHD supporting growth over a

wider range of temperatures. Thus, we drove its expression with the promoter of the

inulinase *inuA* gene, which is inducible by the presence of sucrose in the medium and

almost completely shut off on glucose (Hernández-González et al., 2018; Peñalva et al.,

2020). Initially we tested wild-type and E6K/G540S GHD in a *uso1+* background. This

had no phenotypic consequences despite the fact that western blots confirmed that that

the truncated proteins were being overexpressed (Figure 9D; Figure 9E, lanes 1, 3, 5, 7

and 10;). Then we proceeded to delete the resident *USO1* gene in the wild-type and

mutant GHD overexpressing strains. As expected, neither of the resulting pair of strains

was able to grow on medium with glucose as the only carbon source (Figure 9E, lanes

2 and 6). Notably, the strain expressing E6K/G540S GHD as sole Uso1 source grew

essentially as the wild-type and 30ºC and, although debilitated, was viable at 37ºC,

showing a substantial improvement of the growth capacity displayed by the gene-

replaced mutant (Figure 9, lanes 7,8 and 9). Thus, if expressed at sufficiently high levels,

E6K/G540S GHD maintains viability at the optimal growth temperature.

Unexpectedly, the wild-type GHD also rescued the viability of the *uso1Δ* mutant when

this was cultured with sucrose as carbon source at 30ºC and 37ºC. In fact, at 30ºC, the

uso1Δ inuAp::GHD strain grew like the wt (Figure 9E, lanes 3 and 4)(note that these

experiments were carried out in a *RAB1+* background), suggesting that increased

binding to a Golgi receptor facilitated by mass action compensated for the loss of the

coiled-coil region and associated dimerization. The E6K/G540S GHD would have gained

affinity for this receptor, explaining why the doubly substituted GHD suppressed mis-

localization of Uso1-GFP when RAB1 is compromised, even when its steady-state levels

were very low. Forced expression, combined with a potentially increased binding affinity

of E6K/G540S GHD to such a hypothetical receptor might be toxic.

Uso1 is an associate of the early Golgi SNARE machinery, with the double

substitution E6K/G540S increasing this association

What is the nature of this hypothetical receptor? To address this question, we screened

for interactors of Uso1 among proteins acting at the same functional level (consumption

of COPII vesicles by the early Golgi) using a modified version of the S-tag co-

precipitation approach that we used to characterize of TRAPP complexes (Pinar et al.,

2019)(Figure 10A). We constructed strains expressing wild-type or mutant Uso1, tagged

endogenously with the S-tag and, as negative unrelated control, BapH (an effector of

RAB11 acting in late steps of the secretory pathway (Pinar and Peñalva, 2017). Then,

derivatives of these three strains co-expressing each of the candidate Uso1 GHD

targets, tagged with HA3 (also endogenously), were constructed. The resulting panel

(Figure 10B) was screened for HA-tagged proteins co-precipitating more efficiently with

the E6K/G540S version of Uso1 than with the wild-type, and satisfying the criterium of

not co-purifying with BapH. To this end cell-free extracts of these strains were incubated

with S-agarose beads that were recovered by centrifugation. Proteins associating with

the S-baits were revealed by anti-HA western blotting.

That not every protein specifically co-purified with Uso1 baits was demonstrated by the

results obtained with the COG component COG2, which did not associate with any of

the three baits (Figure 10C, 10). In contrast, β-COP was a promiscuous non-specific

interactor pulled down by all three baits (Figure 10C, 11). Notably, the screen identified

the Golgi syntaxin Sed5 within the specific Uso1 associates (Figure 10C, 1), an

association reported previously by others for both p115 and fungal Uso1 (Allan et al.,

2000; Sapperstein et al., 1996). That the PM SNARE Sso1 did not interact at all with any

of the S-baits demonstrated that Uso1 does not bind promiscuously to syntaxins (Figure

10C, 5). Importantly, Sed5 was brought down more efficiently by E6K/G540S Uso1, and

not at all by BapH, even though levels of this unrelated bait, as assessed by silver

staining of pull-downs, were markedly higher than those of either Uso1 version.

Therefore, Sed5 (or its associates) might represent a potential anchor bound by

E6K/G540S Uso1 with increased affinity to compensate for the lack of RAB1-mediated

recruitment.

This analysis was extended to other members of the SNARE bundle forming in the

ER/Golgi interface with the Qa Sed5: the Qb Bos1, the Qc Bet1 and the R-SNARE Sec22

(Parlati et al., 2002; Pelham, 1999; Tsui et al., 2001). Sec22 was slightly enriched in the

E6K/G540S pull-down relative to wild-type Uso1 (Figure 10C, 4). The results with Bos1

and Bet1 were most noteworthy (Figure 10C2 and 3). Both were markedly increased in

the E6K/G540S pull-downs, with Bet1 increased most. The AAA ATPase Sec18

disassembling *cis*-SNARE complexes also bound Uso1 and was slightly enriched in the

sample of E6K/G540S associates, as was the Uso1-interacting Golgin Bug1, but not its

membrane anchor Grh1 (Figure 10, C6-8). We conclude that Uso1 is a component of

the SNARE machinery and that this association is augmented by E6K/G540S.

S-tag-coprecipitations in Figure 10 clearly singled out Bet1 (Qc) and Bos1(Qb) as the

preys that were most strongly enriched with the mutant E6K/G540S bait relative to wild-

type, and therefore with the highest probability of being direct interactors bound with

greater affinity by the doubly substituted Uso1 mutant.

During these analyses we also addressed the physiological role of golgins, which were

expected to be dispensable for growth, given their functional redundancy (Gillingham,

2018; Gillingham and Munro, 2016; Muschalik and Munro, 2018). We examined the role

of three golgins acting at the early Golgi: The Grh1 (AN11248)/Bug1(AN7680) complex,

discussed above, Coy1 (the product of AN0762) and Rud3 (AN10186). Consistent with

their roles, AlphaFold2 predicts that they form long coiled-coils carrying C-terminal

membrane anchors, characteristics of golgins. Coy1 contains a C-terminal TMD anchor

and an adjacent CLASP domain that, according to AlphaFold2, consists of α-helices

(Figure 10—figure supplement 1). Rud3 is a dimer consisting of a long coiled-coil with a

ARF1-binding GRIP domain composed of four short helices, near its C-terminus (Figure

10—figure supplemental 1). Ablation of *grh1*, *bug1*, *coy1* or *rud3* did not prevent growth

of the corresponding mutants, with only *coy1*Δ displaying a minor growth phenotype

(Figure 10—figure supplement 2). Next, we deleted the corresponding genes in a *rab1*Δ

uso1^{E6K G540S} and tested whether any of the golgins was required for viability rescue. Grh1

and Bug1 were partially required, as their absence precluded rescue at 42° (Figure 10—

figure supplement 2). In contrast, RUD3 was not required. With regard to COY1, the fact

that combining *coy1∆* and *rab1∆ uso1*^{E6K G540S} results in lethality at 30ºC, 37ºC and 42ºC

precluded conclusions on the role of COY1 in suppression. However, we can conclude

that proper assembly of *A. nidulans* Golgi cisternae is supported by a redundant set of

tethers (Behnia et al., 2007), of which in Coy1 appears to be the least redundant of those

tested. Coy1 has been implicated in retrograde traffic within the Golgi itself (Anderson et

al., 2017)

Golgi SNAREs bind directly to the Uso1 GHD; effects of Uso1 E6K/G540S

Work by others implicated Uso1 in the assembly of the early Golgi SNARE bundle

(Sapperstein et al., 1996). Thus, prompted by co-association experiments, we predicted

that Uso1 would bind directly Bet1, Bos1 and perhaps other SNAREs implicated in the

biogenesis of the early Golgi. We anticipated that binding to Bet1 and Bos1 would be

insufficient to recruit Uso1 to membranes in the absence of RAB1, but that once

reinforced by E6K/G540S, Uso1 would not require RAB1 for its recruitment. To test this

possibility, we searched for direct and E6K/G540S-enhanced interactions between the

GHD and SNAREs with pull down assays carried out with purified SNARE-GST fusion

proteins as baits and Uso1-His6 constructs as preys.

Full-length Uso1 bound, weakly, to the Sec22 R-SNARE and to the Qb Bos1, and very

efficiently [with *circa* 70% of the prey being pulled down (Figure 11A, B)] to the Qc Bet1.

In contrast, Uso1 did not bind the Qa syntaxins tested, Sed5 and Sso1 (Figure 11A). The

absence of interaction between Sed5 and Uso1, be it the wild-type or the E6K/G540S

mutant version, strongly indicated that the association detected with S-tag pull-downs

between Uso1 and Sed5 is bridged by other protein(s). This absence of binding cannot

be attributed to Sed5-GST being incompetent for binding because Sed5-GST was

competent in pulling-down highly efficiently its cognate SM protein Sly1, an interaction

that did not occur with Sso1-GST (Figure 11C). Notably, the presence of the E6K/G540S

double substitution in Uso1 (indicated with ** for simplicity on Fig 11) increased five times

the amount of protein retained by the Qb Bos1 bait (Figure 11A and B), whereas

interaction with Bet1 did not change (Figure 11A and B) . The double substitution in Uso1

did not promote interaction with Sed5 either. Thus, under normal circumstances Uso1 is

able to bind directly to three of the four SNAREs in the ER/Golgi interface, with binding

to Bet1 being the strongest. The double E6K/G540S substitution increases binding to

Bos1 very markedly and specifically, bringing it to up to the levels of Bet1 without
affecting, for example, binding to Bet1 or Sec22. Consistently, the GHD is sufficient to
mediate interaction with Bet1 and Bos1, as well as, if E6K G540S-substituted, the
increased binding of Bos1 to Uso1 (Figure 11D and E). The GHD did not interact with
Sec22, suggesting either that this R-SNARE is recruited by other parts of the protein or
that binding is dimerization dependent.

Besides the Sed5/Bos1/Bet1/Sec22 combination ( ), across cisternal maturation Sed5
forms SNARE bundles in Golgi compartments located downstream of Uso1 domains
(Pelham, 1999). In fungi, membrane fusion in the medial Golgi involves the Sed5
partners Gos1 (Qb) and Sft1 (Qc) substituting for Bos1 and Bet1, respectively, but
neither Gos1 nor Sft1 bound wild-type or E6K/G540 Uso1 GHD (Figure 11F),
demonstrating that interaction of Uso1 with Bos1 and Bet1 is highly specific. Therefore,
it seems fair to conclude that increased binding for a SNARE receptor underlies the
mechanism by which mutant Uso1 bypasses the need for RAB1 in the ER/Golgi
interface.

In summary, Uso1 is an essential protein acting in the ER/Golgi interface, and we report
here several important findings. We show that (i) the Golgi GTPase RAB1, which is
essential for viability, becomes dispensable if there is an alternative method to recruit
Uso1 to Golgi membranes; (ii) the coiled-coil region of Uso1 is dispensable to sustain
viability, implying that the tethering role of the protein is not essential either, consistent
with the redundant roles of other Golgin tethers; (iii) the Uso1 GHD is essential for
viability; (iv) the Uso1 GHD monomer, if present at suitably high levels, is sufficient to
maintain viability; (v) that the Uso1 GHD is a direct and specific binder of the Qb SNARE
Bos1, and a strong binder of the Qc Bet1; (vi) the mutation bypassing the need for RAB1
markedly increases the affinity of the GHD for Bos1, indicating that $rab1\Delta$ viability rescue
by E6K/G540S Uso1 occurs because SNARE anchoring provides an alternative mode
of recruitment to Golgi membranes to that provided physiologically by RAB1.

Mechanistic insights guided by AlphaFold2 predictions

To gain further insight into the mechanisms by which the double E6K/G540S bypasses
RAB1 we exploited AlphaFold2 to model the Uso1^{GHD} domain alone, or together with
each of the individual SNAREs, with both Bos1 and Bet1 simultaneously, and with RAB1.

Consistent with biochemical data, in the predicted model Bet1 and Bos1 interact with a

medial and a C-terminal region, respectively, of Uso1^{GHD}. (Figure 12; Figure 12—

supplement 1 and —supplement 2). Bet1 docks against a region of the GHD that is not

affected by the mutations. In contrast, Gly540 and its environs dock against the N-

terminal, triple α-helical Habc domain of Bos1. In all likelihood Gly540Ser mediates the

increased binding of mutant Uso1^{GHD} to Bos1. AlphaFold2 predicts that Uso1^{GHD}

interacts with Bos1 through a surface composed by the N-terminal part of the first Bos1

Habc α-helix and the loop between α-helices 2 and 3 (Figure 12A and B). The binding

surface in Uso1^{GHD} involves the second α-helices of the ARM10 and ARM11 repeats,

(α-helices 26 and 29), and the loop connecting the first two α-helices of ARM10 (Figure

12B). In AlphaFold2 models, Uso1 Gly504 (wild-type) is located at the beginning of

Uso1^{GHD} α-helix 29, at the heart of the interaction surface, contributing to the Uso1-Bos1

interaction by coordinating the amide group of the Uso1 backbone with the Bos1 Glu58

carboxylate to create a hydrogen bond (Figure 12C). According to the most confident

prediction, Gly504Ser results in the hydroxymethyl side chain protruding into a small

pocket rimmed by the side chains of Bos1 Leu59 and Ile60, such that the Uso1 Ser504

hydroxyl group hydrogen-bonds the amide group of Bos1 Glu58 (Figure 12C). Besides

creating a new hydrogen bond, Gly540Ser would increase the surface of interaction

slightly, from ~825.2 Å^2 to ~831.1 Å^2, implying that these alterations, together with a

minor shift in the environs of Gly540 detected by models (which might facilitate additional

interactions), would strengthen the binding of the Uso1^{GHD} to the Bos1 surface.

The mechanism by which Glu6Lys contributes to increase the recruitment of Uso1 to

Golgi membranes appears to be different. This glutamate is located in a region with low

pLDDT score which shows different conformations depending on the model (Figure 2D,

boxed), suggesting that this region is difficult to predict due to flexibility. However,

models concur in the prediction of an amphipatic N-terminal α-helix containing Glu6,

whose substitution by Lys (as in E6K) reinforces the positive charge of this α-helix, which

would be inserted into the vesicle membrane, potentially contributing to Uso1 recruitment

to COPII vesicles

In the case of the strong Uso1^{GHD}-Bet1 interaction (Figure 12—figure supplement 1), the

N-terminal region of Bet1 consisting of *circa* 80 amino acids is disordered, and the

pLDDT score of the different models is understandably low. However, all structural

models depicting Bet1 interacting with Uso1^{GHD} (e.g. in the context of the whole SNARE

complex, Bet1 alone or the isolated Bet1 N-terminal) consistently show a region where

the pLDDT is higher. This region forms a kink in this N-terminal part of the Bet1 that

protrudes into the surface created by the α-helices 13 and 16 of ARM4 and ARM5 (Figure

12—figure supplement 1). Surface representations show that this section of the Bet1

polypeptide covers $\sim$ 1300 $Å^2$ of the GHD, docking against the same side of the

boomerang-shaped solenoid as Bos1, which is consistent with the orientation that these

SNAREs should take during the formation of the SNARE pin (Figure 12—figure

supplement 2)

We next modelled RAB1 binding to the GHD in the absence and presence of Uso1

binders Bos1 and Bet1. AlphaFold2 predicts that the GHD interacts with RAB1 through

a binding surface formed by the ARM3-5. repeats. On the other hand, the interactive

region of RAB1 conforms to canons, as Uso1 α-helices 8 and 9 (ARM3) interact with the

Switch II region while α-helices 11 (ARM-4) and 14 (ARM-5) interact with the Switch I)

(Figure 12—figure supplement 2, A and B). Importantly, the model indicates that within

the Uso1-GHD jai alai basket, RAB1 binds to the opposite (convex) side of the Bet1

interacting area at the concave side, and away from the C-terminal helix where the Bos1

Habc domain predictably binds, which would allow Uso1 to bind these three interactors

simultaneously (Figure 12—figure supplement 2C). In addition, the predicted models

supported two highly suggestive but as yet speculative implications. One is that the

position of the RAB1 hypervariable domain, which anchors the GTPase to the membrane

through is prenylated C-terminus, is compatible whith the hypothetical membrane

binding of the N-terminal Uso1 amphipatic helix (Figure 12—figure supplement 2D); The

second is that Uso1-RAB1 would be in an orientation that facilitates the docking of RAB1-

loaded ER-derived vesicles with an acceptor membrane where SNARE zippering occurs

(see discussion); this orientation implies that the donor membrane (the position of the N-

terminal Uso1 helix) and the acceptor membrane (the TMDs of the SNAREs) is

compatible with the Uso1 CTR contributing to tethering through interaction with

Bug1/Grh1.

714   **Discussion**

716   RAB1 regulates transport at the ER/Golgi interface. Using an unbiased forward genetic

717   screen to identify subordinated genes accounting for its essential role, we isolated two

718   extragenic mutations resulting in single-residue substitutions in the RAB1 effector Uso1,

usually regarded as a tether. The single-residue substitutions lie at opposite ends of the jai alai basket-shaped GHD; both are individually able to rescue viability of *rab1Δ* strains at 30ºC and, when combined, even at 42ºC. Subcellular localization experiments hinted at the mechanism by which the double mutation rescues *rab1Δ* lethality. Uso1 plays its physiological role on an early Golgi compartment, where it largely colocalizes with RAB1, with relocation to the cytosol in a RAB1-deficient background. Under normal circumstances, the Uso1 CTR acts in concert with RAB1 to recruit the protein to the Golgi. Genetic evidence (Figure 8) showed that this contribution requires the Golgi-localized tether composed of the membrane anchor Grh1 and its associated golgin BUG1 (Behnia et al., 2007), homologues of human GRASP65 and GM130, respectively. In the absence of RAB1, engagement of the CTR with BUG1 is insufficient to stabilize wild-type Uso1 on membranes. However, the double E6K/G540S substitution relocalizes Uso1 to Golgi structures, suggesting that mutant Uso1 GHD had gained affinity for another element, thereby compensating for the loss of RAB1.

By S-tag co-precipitation experiments we identified proteins associating with wild-type and E6K/G540S Uso1 baits. Prominent among these were the four SNAREs, Sed5, Bos1, Bet1 and Sec22, mediating fusion events at the ER/ Golgi interface, and the SNARE regulator Sec18, indicating that Uso1 is a component of the SNARE fusion machinery. This was suggested by previous studies with *S. cerevisiae* showing that overexpression of Bet1, Bos1 and Sec22 suppresses *uso1-1* and that of Sec22 and Bet1 weakly suppresses *uso1Δ* (Sapperstein et al., 1996). Moreover, in mammalian cells crosslinking studies with p115 identified Sed5, membrin (Bos1) and mBet1 as its weak interactors (Allan et al., 2000), suggesting that contributing to the ER/Golgi SNARE machinery is a conserved feature of Uso1/p115 family members. Of the above four *A. nidulans* SNAREs, Bet1 and Bos1, which co-precipitated weakly with the wild-type, were dramatically enriched with the mutant. Pull-down assays showed that full-length *Aspergillus* Uso1 interacts weakly with Sec22 and strongly with Bet1, irrespective of whether the bait was wild-type or E6K/G540S, indicating that they are direct physiological interactors. In sharp contrast, the interaction of E6K/G540S Uso1 with the Qb Bos1 was markedly augmented, strongly suggesting that this increase contributes substantially to bypass RAB1. Both the high levels of binding to Bet1, and the marked increase in binding to Bos1 resulting from E6K/G540S were tracked down to the GHD, which is sufficient to bind these SNAREs.

AlphaFold2 models predicting the regions of interaction of the GHD with Bos1 and Bet1

showed that, of the two Uso1 residue substitutions, only Gly540Ser maps to the region

mediating the interaction with Bos1, whereas neither affected the interacting region with

Bet1, agreeing with GST pull-downs. Interaction with this SNARE is predicted to involve

a large surface, consistent with the strong "constitutive" binding of Bet1 with the GHD.

AlphaFold2 also predicted that RAB1 binds to the convex face of the GHD solenoid, in a

position that would permit the simultaneous binding of Qb and Qc SNARES and the

GTPase. In addition, AlphaFold2 also detected a previously unnoticed amphipathic α-

helix in the N-terminal region of Uso1. The Glu6Lys substitution falls within this helix,

making its global positive charge even greater, strongly suggesting that this substitution

increases Uso1 GHD binding to membranes. That the two residue substitutions act by

different mechanisms is coherent with their showing additivity to suppress RAB1 deficit.

The Uso1/p115 family has been implicated in the regulation of SNARE complexes (Allan

et al., 2000). (Shorter et al., 2002) first reported that p115 binds to SNAREs and

proposed that they would 'catalyze' the formation of the ER/Golgi SNARE bundle. The

direct interaction of p115 with unassembled mBet1 and Sec22 agrees with this role

(Wang et al., 2014). However, there are significant gaps in this model: Uso1

overexpression suppresses a partial deficit of *ypt1/RAB1*, but not *ypt1Δ, whereas* Ypt1

overexpression suppresses *uso1Δ*, indicating that Uso1 acts and in concert or upstream

of RAB1. [Of note, overexpression of Uso1 also suppresses *bet3-1*, a *ts* allele

inactivating the TRAPPIII complex, which is the RAB1 GEF (Galindo et al., 2021; Jiang

et al., 1998; Pinar et al., 2019; Riedel et al., 2017; Thomas et al., 2018)]. The grid of

reported genetic interactions strongly indicates that RAB1 regulates SNAREs (Brandon

et al., 2006; Lupashin and Waters, 1997; Sapperstein et al., 1996). Notably, our viability

rescue experiments show that E6K/G540S bypasses this non-canonical, yet essential

role of RAB1. As the double mutant substitution increases the recruitment of USO1 to

the SNARE machinery and as E6K/G540S suppresses *rab1Δ*, the most parsimonic

interpretation of the data is that the role of RAB1is cooperating to recruit Uso1 to the

SNARE complexes, a cooperation that is no longer needed when recruitment is ensured

by other means. That p115 regulates SNAREs' assembly was proposed by (Shorter et

al., 2002). However, a fundamental difference with our conclusions is that they attributed

this role to the CCD, whereas we identify here the GHD as the positively-acting player.

We also note that our experiments provide a mechanistic interpretation as to why

overexpression of Bet1, Bos1, Sec22 and Ypt1/RAB1 rescue the viability of *uso1-1*

removing a substantial portion of the yeast Uso1 CCD (Sapperstein et al., 1996; Seog

et al., 1994). A similar *Aspergillus* allele results in marked protein instability (Figure 9).

Thus, in all likelihood, suppression by overexpression of known direct interactors

involves stabilization of the *uso1-1* product.

In summary, our work firmly establishes that the essential role of Uso1 resides not in its

CCD domain tethering donor and acceptor membranes, but in the GHD. When

expressed at sufficient levels this domain is capable of fully complementing *uso1Δ*, which

is definitive evidence that neither the tethering function of Uso1 nor dimerization (the

GHD is monomeric) is required for the protein to play its essential role. As the GHD binds

two of the four SNAREs of the bundle, the simplest interpretation is that the essential

role of Uso1 is regulating SNAREs.

Ideas and Speculation

While the molecular details of this regulation will be addressed in future, it is tempting to

speculate that the GHD contributes, with Sly1, to orientate SNAREs to form a productive

bundle, acting as chaperones, similarly to the HOPS SM component Vps33, which

appears to align SNAREs in a pre-zippering stage, facilitating their assembly (Baker and

Hughson, 2016; Baker et al., 2015; Ren et al., 2009; Yu and Hughson, 2010; Zhang and

Hughson, 2021; Zhang and Yang, 2020). A second speculative interpretation is that

tethering occurs in two steps, with golgins acting at long distances, approximating

vesicles to the vicinity of SNAREs. Then SNAREs would engage RAB1–Uso1 to serve

as short-range tether preceding the zippering up of membranes. The > 700 Å-long Uso1

CCD would cooperate with Bug1 in the first step, whereas the Uso1 GHD would

cooperate with RAB1 and Bet1/Bos1 in the second, exploiting the fact that GHD binders

use surfaces located at opposite sides of the α-solenoid (Figure 12—figure supplement

2). Such arrangement implies that RAB1 C-terminal isoprenoids inserting into the donor

vesicle membrane would be *circa* 220 Å apart from the C-termini of the SNAREs inserted

in the acceptor membrane. Two-step tethering might impose one additional level of

specificity, preventing unproductive fusion events mediated by other SNARES circulating

through the ER/ interface and directing, by way of Uso1 interactions, incoming vesicles

to fusion-competent areas enriched in target SNAREs (Bentley et al., 2006).

Materials and Methods

Aspergillus techniques

Standard *A. nidulans* media were used for growth tests, strain maintenance and conidiospore harvesting (Cove, 1966). GFP-, HA3- and S-tagged alleles were introduced by homologous recombination-mediated gene replacement, using transformation (Tilburn et al., 1983) of recipient *nkuAΔ* strains deficient in the non-homologous end joining pathway (Nayak et al., 2005). Complete strain genotypes are listed in supplemental table I. These alleles were usually mobilized into the different genetic backgrounds by meiotic recombination (Todd et al., 2007).

Null mutant strains were constructed by transformation-mediated gene replacement, using as donor DNA cassettes made by fusion PCR (primers detailed in supplemental table II) carrying appropriate selectable markers (Szewczyk et al., 2006). Integration events were confirmed by PCR with external primers. When allele combinations were expected to be synthetically lethal or severely debilitating, the corresponding strains were constructed by sequential transformation, but the second such manipulation was always carried out using $pyrG^{Af}$ as selective marker, which favors the formation of heterokaryons in which untransformed nuclei supported growth (Osmani et al., 1988). Conidiospores, in which single nuclei had segregated (i.e., homokaryotic nuclei), were scrapped and streaked onto plates carrying doubly-selective medium. Absence of growth or appearance of microcolonies, combined with a positive PCR diagnostic of heterokaryosis of the primary transformants, was taken as indication of lethality. Whenever possible, colony PCR of microcolonies was always used to genotype the desired genetic intervention

The following proteins were C- or N-terminally tagged endogenously, using cassettes constructed by fusion PCR (Nayak et al., 2005; Szewczyk et al., 2006): Uso1-GFP and Uso1$^{E6K/G540S}$-GFP, Uso1-HA3 and Uso1$^{E6K/G540S}$-HA3; Uso1-S and Uso1$^{E6K/G540S}$-S; BapH-S (Pinar and Peñalva, 2017), Sec13-mCherry (Bravo-Plaza et al., 2019; Hernández-González et al., 2019), Gea1-mCherry and Sec7-mCherry (Arst et al., 2014), mCherry-Sed5 (Pantazopoulou and Peñalva, 2011), mCherry-RAB1 (Pinar et al., 2013), HA3-Sed5, HA3-Bet1, HA3-Bos1, HA3-Sec22, Sec18-HA3, Grh1-HA3, Bug1-HA3, Coy1-HA3, COG2-HA3, and β-COP-HA3.

860 **Antibodies for western blotting**

Antibody	dilution	Origin	Reference
Primary			
α-HA	1:1000	rat	3F10 clone, Roche
α-His6x tag	1:10 000	mouse	#631212, Clontech
α-tubulin	1:5000	mouse	DM1A clone, Sigma
α-Uso1	1:1000	rabbit	Polyclonal antiserum, Davids Biotechnologie
Secondary (HRP-conjugated)			
α-rat IgG	1:4000	goat	#3010-05, Southern Biotech
α-mouse IgG	1:5000	goat	#A9044, Sigma
α-rabbit IgG	1:2000	donkey	#NA934, Amersham

Antiserum against Uso1 was raised in rabbits by Davids Biotechnology. Animals were
immunized with the Uso1 GHD (residues 1-659), tagged with His6x. Recombinant
expression in *E. coli* and Ni^{2+} affinity purification is described below. Target antibodies
were purified from raw antiserum by affinity chromatography through Hi-Trap NHS
columns (#17-0716-01, Cytiva) charged with Uso1 antigen following the manufacturer's
instructions. Affinity-bound antibodies were eluted with 100 mM glycine (pH 3.0), then
neutralized with 2 M Tris to a pH of 7.5 and stored at -20ºC.

inuA* promoter-driven expression of Uso1 GHD in *Aspergillus
The Uso1 GHD was expressed in a sucrose-inducible manner from an *in locus*
replacement of the *inuA* gene (AN11778) ORF encoding inulinase by the GHD coding
sequence, such that its expression was driven by the *inuA* promoter, which is induced
on sucrose and non-induced on glucose (Hernández-González et al., 2018). The gene
replacement cassette was assembled through fusion PCR of 4 different elements, listed
here from 5' to 3': (1) inuA promoter, (2) cDNA sequence encoding Uso1 wt or mutant
E6K/G540S GHD (residues 1 – 659), (3) *Aspergillus fumigatus riboB* gene as selection
marker and (4) *inuA* gene 3'-flanking region. A *pyrG*89, *nkuA*Δ::bar, *riboB*2 *A. nidulans*
strain was transformed with this cassette, replacing the inuA gene. The resulting strain
was subsequently transformed with a *uso1*Δ deletion cassette (*A. fumigatus pyrG* as
selection marker) to ablate endogenous of Uso1, such that the GHD was the only moiety
of Uso1 present.

Plasmids for protein expression in *E. coli*
882
(I) His6x-tagging constructs

pET21b-Uso1-His6x and pET21b-Uso1(E6K/G540S)-His6x: cDNA encoding full length
Uso1 (residues 1 -1103) was cloned into a pET21b *Nde*I/*Not*I linearized vector.

pET21b-Uso1ΔCTR-His6x and pET21b-Uso1(E6K/G540S)ΔCTR-His6x: lacking the C-
Terminal Region of the Coiled Coil Domain (residues 1 – 1040)

pET21b-Uso1GHD-His6x and pET21b-Uso1(E6K/G540S)GHD-His6x: cDNA encoding
the Globular Head Domain of Uso1 (residues 1 - 659) was cloned as a *Nde*I/*Xho*I insert
into a pET21b *Nde*I/*Xho*I linearized vector.

pET21b-Uso1 CCD-His6x: cDNA encoding Uso1 Coiled Coil Domain (residues 660-
1103) was cloned as a *Nde*I/*Not*I insert into a pET21b *Nde*I/*Not*I linearized vector.

(II) TNT® expression plasmids

pSP64-Sly1-HA: this plasmid carries cDNA encoding full length Sly1 (AN2518) C-
terminally tagged with a HA3x epitope, cloned as an *Nsi*I/*Sac*I insert into *Pst*I/*Sac*I
pSP64(PolyA) vector

(III) GST-tagging constructs

pET21b-Sed5-GST: cDNA encoding Sed5/AN9526 cytoplasmic domain (residues 1–
322) C-terminally tagged with GST, was cloned as a *Nde*I/*Sal*I insert into a pET21b
*Nde*I/*Xho*I linearized vector.

pET21b-Bos1-GST: cDNA encoding Bos1/AN11900 cytoplasmic domain (residues 1–
219) C-terminally tagged with GST, was cloned as a *Nde*I/*Sal*I insert into a pET21b
*Nde*I/*Xho*I linearized vector.

pET21b-Bet1-GST: cDNA encoding Bos1/AN5127 cytoplasmic domain (residues 1–71)
C-terminally tagged with GST, was cloned as a *Nde*I/*Sal*I insert into a pET21b *Nde*I/*Xho*I
linearized vector.

pET21b-Sec22-GST: cDNA encoding Sec22/ ASPND00903 cytoplasmic domain
(residues 1–198) C-terminally tagged with GST, was cloned as a *Nhe*I/*Sal*I insert into a
pET21b *Nhe*I/*Xho*I linearized vector.

pET21b-Sso1-GST: cDNA encoding Sso1/AN3416 cytoplasmic domain (residues 1–
271) C-terminally tagged with GST, was cloned as a *Nde*I/*Sac*I insert into a pET21b
*Nde*I/*Sac*I linearized vector.

pET21b-Gos1-GST: cDNA encoding Gos1/AN1229 cytoplasmic domain (residues 1–
208) C-terminally tagged with GST, was cloned as a *Nde*I/*Sal*I insert into a pET21b
*Nde*I/*Xho*I linearized vector.

<u>pET21b-Sft1-GST:</u> cDNA encoding Sft1/AN10508 cytoplasmic domain (residues 1–73)
C-terminally tagged with GST, was cloned as a *Nde*I/*Sal*I insert into a pET21b *Nde*I/*Xho*I
linearized vector.

Co-precipitation experiments with total cell extracts

Preparation of *Aspergillus* total cell extracts was done as described, with minor
modifications (Pinar et al., 2019) (Pinar et al., 2019). 70 mg of lyophilized mycelium were
ground with a ceramic bead in a Fast Prep (settings: 20 sec, power 4). The resulting fine
powder was resuspended in 1.5 ml of extraction buffer [25 mM HEPES-KOH (pH 7.5),
200 mM KCl, 4 mM EDTA, 1% (v/v) IGEPAL CA-630 (NP-40 substitute, #I8896, Sigma),
1 mM DTT, 2 µM MG-132 proteasome inhibitor (#S2619, SelleckChem) and cOmplete®
ULTRA EDTA-free inhibitor cocktail (#5892953001, Roche). Approximately 0.1 ml of 0.6
mm glass beads were added and thoroughly mixed. This suspension was homogenized
with a 10 sec pulse at the Fast Prep (power 6) followed by a 10 min incubation at 4ºC.
This homogenization step was repeated two times before clarifying the extract by
centrifugation at 4ºC and 15,000 × *g* in a microcentrifuge. Total protein concentration of
the extracts was determined by Bradford. Bovine Serum-Albumin BSA was then added
as a blocking agent to the cell extract (final concentration 1% (w/v)). Binding reactions
were carried out in 0.8 ml Pierce centrifuge columns (#89869, ThermoFisher): 9 mg of
protein were mixed with 20 µL of S-protein Agarose beads (#69704, Novagen), that had
been previously washed in extraction buffer with 1% (w/v) BSA. This buffer was also
added to complete final reaction volume of 0.6 ml. The mix was incubated for 3 h at 4°C
in a rotating wheel. Columns were then opened at the bottom and gently centrifuged to
remove the supernatant and collect the protein-bound beads. These were resuspended
in extraction buffer without inhibitors and incubated in rotation for 10 min at 4ºC, followed
by two more washing steps in extraction buffer without detergent and inhibitors. To elute
proteins bound to the beads, 30 µl of Laemmli loading buffer [62.5 mM Tris-HCl (pH 6.0),
6 M urea, 2% (w/v) SDS and 5% (v/v) β-mercaptoethanol] were added and the columns
incubated at 90ºC for 2 min. The columns were centrifuged to collect the eluate, of which
a 40% of the final volume were resolved in a SDS-polyacrylamide gel and then
transferred to nitrocellulose for α-HA (#3F10, Roche) western blotting.

Purification of Uso1 constructs tagged with His6x

Full-length His6-tagged Uso1, Uso1∆CTR, Uso1 GTD and Uso1 CCD constructs, wild
type and mutant versions, were expressed in *E. coli* BL21(DE3) cells harboring pET21b-
His6 derivatives and pRIL. Bacteria were cultured at 37ºC in LB medium containing
ampicillin and chloramphenicol until reaching an OD$_{600nm}$ of 0.6. Then, IPTG was added
to a final concentration of 0.1 mM. Cultures were shifted to 15ºC and incubated for 20 h.

Bacterial cells were collected by centrifugation and pellets stored at -80ºC. For
purification, frozen pellets were thawed in ice and resuspended in ice-cold bacterial cell
lysis buffer [20 mM sodium phosphate buffer, pH 7.4, 500 mM KCl, 30 mM imidazole,
5% (v/v) glycerol, 1 mM β-mercaptoethanol, 1 mM MgCl$_2$, 0.2 mg/ml lysozyme and 1
µg/ml of DNAse I and cOmplete® protease inhibitor cocktail (#11873580001, Sigma)].
This cell suspension was mechanically lysed in a French press (1500 kg/cm^2) and the
resulting lysate was centrifuged at 10,000 × g and 4ºC for 20 min to remove the cell
debris. The supernatant was then transferred to polycarbonate tubes and centrifuged at
100,000 × g and 4ºC for 1 h in a XL-90 ultracentrifuge (Beckman Coulter) . 50 ml of
cleared lysate were incubated with 400 µL of Ni-Sepharose High Performance beads
(#17526801, Cytiva) for 2 h at 4ºC. After this step, His-tagged protein-bound beads were
pelleted at low-speed centrifugation and washed three times in lysis buffer [20 mM
sodium phosphate buffer, pH 7.4, 500 mM KCl, 5% (v/v) glycerol, 1 mM β-
mercaptoethanol] with increasing concentrations of imidazole. Finally, Ni2+bound His6
proteins were eluted 0.5 M imidazole buffer. 5 ml of eluted protein were loaded onto a
HiLoad 16/600 Superdex 200 column (Cytiva) and run at 1 ml/min flow rate on an AKTA
HPLC system, using phosphate buffered saline PBS containing 5% (v/v) glycerol and 1
mM β-mercaptoethanol. Fractions containing protein were pooled, analyzed for purity by
SDS-PAGE followed by Coomassie staining, and finally quantified on a UV-Vis
spectrophotometer before being stored at -80ºC.

Purification of SNARE constructs tagged with GST
cDNAs encoding the cytosolic domains of SNAREs fused to a C-terminal GST were
cloned into pET21b. Bacterial cultures and protein expression conditions were as
described above for Uso1-His6 constructs. Frozen pellets were thawed in ice and
resuspended in chilled bacterial cell lysis buffer [25 mM Tris-HCl (pH 7.4), 300 mM KCl,
5 mM MgCl$_2$, 1 mM DTT, 0.5 mg/ml lysozyme, 1 µg/ml of DNAse I and cOmplete®
protease inhibitor cocktail (#11873580001, Sigma)]. This cell suspension was incubated
for 30 min in ice before being mechanically lysed in a French press (1500 kg/cm^2). The
lysate was incubated for a further 30 min on ice and centrifuged at 20,000 × g and 4ºC
for 30 min. After adding 10 mM EDTA to the clarified supernatant to stop DNAse I activity,
it was transferred to a 50 ml tube, mixed with 500 µL of glutathione Sepharose beads 4B
(#17075601, Cytiva) and rotated for 2 h at 4ºC. After incubation, SNARE-GST-bound
beads were pelleted by gentle centrifugation and washed three times for 10 min at 4ºC
in 25 mM Tris-HCl (pH 7.4), 500 mM KCl, 5 mM EDTA, 1 mM DTT and subsequently
transferred to a 0.8 ml Pierce column. Beads were washed 6 times (10 min at RT) in 200
µL of elution buffer [50 mM Tris-HCl (pH 8.0), 200 mM KCl, 10 mM glutathione and 1

mM DTT]. These fractions were collected and pooled (~1 ml), then buffer-exchanged to

storage buffer (PBS, 5% (v/v) glycerol and 0.1 mM DTT) in a PD MidiTrap G-25 column.

Protein concentration and purity was assessed by spectrophotometry and SDS-PAGE

followed by Coomassie staining. Protein stocks were kept frozen at -80ºC.

SNARE-GST pull-downs with purified Uso1-His6 constructs (Uso1, Uso1 GHD)

Binding reactions were performed in 0.8 ml Pierce centrifuge columns. 75 µg of purified

SNARE-GST were mixed with 15 µL of glutathione Sepharose 4B beads and storage

buffer to a final volume of 0.3 ml. Columns were rotated at 4ºC for 2 h before the

supernatant was removed after low speed centrifugation. Subsequently, His6 preys were

added to a final concentration of 0.2 µM in a total volume of 0.4 ml of pull-down binding

buffer [25 mM HEPES-KOH (pH 7.5), 150 mM NaCl, 10% (v/v) glycerol, 0.1% (v/v) Triton

X-100 and 0.1 mM DTT]. Columns were rotated overnight at 4ºC. Beads were collected

by gentle centrifugation and washed three times for 10 min with ice-cold binding buffer,

before eluting bound proteins with 30 µL of Laemmli loading buffer pre-heated at 90ºC.

0.5 % of the samples (eluted material or flow-through) were run onto 8% SDS-

polyacrylamide gels that were transferred to nitrocellulose membranes which were

reacted with α-His tag antibody (#631212, Clontech). Quantitation of band intensities

was done with ImageLab software (BioRad). In parallel, 4% of the elution sample volume

was loaded onto 10% SDS-polyacrylamide gel and stained with coomassie dye

(BlueSafe, NZY) to confirm recovery of SNARE-GST baits.

1008

Pull-down of TNT®-expressed Sly1-HA3

Sly1-HA3 was synthesized with the TNT® SP6 Quick Coupled Transcription/Translation

system (#L2080, Promega), according to the instructions of the manufacturer. The

reaction was primed with 1 µg of pSP64::Sly1-HA3 cDNA. 10 µL of the resulting mix

were combined with 15 µL of glutathione-Sepharose beads, previously loaded with

SNARE-GST baits as described above, in 0.4 ml of pull-down binding buffer, using 0.8

ml Pierce columns that were rotated overnight at 4ºC before beads and flow-through

were recovered after gentle centrifugation. Beads were washed three times for 10 min

at 4ºC in pull-down binding buffer before eluting bound material with 30 µL of Laemmli

loading buffer for 2 min at 90ºC. 20% of the elution sample volume was analyzed by

western blotting with α-HA tag antibody (#3F10, Roche) for Sly1-HA immunodetection.

Size exclusion chromatography of HA-tagged cell extracts

Gel filtration experiments were performed as described (Bravo-Plaza et al., 2019).

Briefly, 200 µL of cell extract were loaded onto a Superose 6 10/300 column (Pharmacia)

equilibrated with running buffer [25 mM Tris-HCl (pH 7.5), 600 mM KCl, 4 mM EDTA, 1

mM DTT]. Fractions of 0.5 ml were collected, from which 80 µL were mixed with 40 µL

of Laemmli loading buffer and denatured at 90ºC. 25 µL of these samples were resolved

by SDS-PAGE and analyzed by western blotting with α-HA3 tag antibody (#3F10,

Roche) for western blotting. Sizing standards were myoglobin (17 kDa), BSA (67 kDa),

aldolase (158 kDa) ferritin (449 kDa), thyroglobulin (669 kDa) and dextran blue (Vo).

Analytical ultracentrifugation: sedimentation velocity

1029

Sedimentation velocity assays and subsequent raw data analysis were performed in the

Molecular Interactions Facility of the Centro de Investigaciones Biológicas Margarita

Salas. Samples (320 µL) in PBS containing 5% (v/v) glycerol and 1 mM β-

mercaptoethanol were loaded into analytical ultracentrifugation cells, which were run at

20°C and 48,000 rpm in a XL-I analytical ultracentrifuge (Beckman-Coulter Inc.)

equipped with UV-VIS absorbance and Raleigh interference detection systems, using an

An-50Ti rotor, and 12 mm Epon-charcoal standard double-sector centerpieces.

Sedimentation profiles were recorded at 230 nm. Differential sedimentation coefficient

distributions were calculated by least-squares boundary modelling of sedimentation

velocity data using the continuous distribution $c(s)$ Lamm equation model as

implemented by SEDFIT(Schuck, 2000). Experimental Svedberg coefficient values were

corrected to standard conditions ($s_{20,w}$: water, 20°C, and infinite dilution) using

SEDNTERP software

Dynamic Light Scattering, DLS

1043

DLS experiments were carried out in a Protein Solutions DynaPro MS/X instrument at

20C using a 90° light scattering cuvette. DLS autocorrelation functions, average of at

least 18 replicates, were collected with Dynamics V6 software. Analysis evidenced in

most cases the presence of two diffusing species, one with faster diffusion corresponding

to a discrete major species and a second with substantially slower diffusion,

corresponding to higher order species, with a minor contribution to the whole population

within the sample. Exceptions were the globular domain GHD that appeared as a single

species, and the coiled-coil construct with a larger contribution of the higher order

species. Dynamics software was also employed to export the data as text files for parallel

analysis using user-written scripts and functions in MATLAB (Version 7.10, MathWorks,

Natick, MA). A double exponential decay model was fit to the data *via* nonlinear least

squares, using as starting values the translational diffusion coefficients and relative

amounts of the two different species, using as starting values those from the

regularization analysis and their masses (those of the discrete species as estimated by

the Svedberg equation). These values were compatible with the experimental data,

rendering a similar best-fit value of the diffusion coefficient for the major species and
allowing to assess its probability distribution.

Estimate of molar mass from hydrodynamic measurements.
The apparent molar masses of Uso1 and its mutants were calculated via the Svedberg
equation, using the *s*- and *D*-values of the major species independently measured by
sedimentation velocity and DLS, respectively.

Fluorescence Microscopy
A. nidulans hyphae were cultured in Watch Minimal Medium WMM (Peñalva, 2005).
Image acquisition equipment, microscopy culture chambers and software have been
detailed (Pinar et al., 2022; Pinar and Peñalva, 2020). Simultaneous visualization of
green and red emission channels was achieved with a Gemini Hamamatsu beam splitter
coupled to a Leica DMi8 inverted microscope. Z-Stacks were deconvolved using
Huygens Professional software (version 20.04.0p5 64 bits, SVI). Images were contrasted
with Metamorph (Molecular Devices). Statistical analysis was performed with GraphPad
Prism 8.02 (GraphPad). Uso1-GFP time of residence in cisternae was estimated from
3D movies consisting of middle planes with 400 photograms at 2 fps time resolution.
Each Uso1 puncta considered in the analysis was tracked manually with 3D (x,y,t)
representations generated with Imaris software (Oxford Instruments) combined with
direct observation of photograms in movies and kymograph representations traced
across >25 px-wide linear ROI covering the full width of the hyphae.

AlphaFold predictions
AlphaFold2 (Jumper et al., 2021) predictions were run using versions of the program
installed locally and on ColabFold (Mirdita et al., 2022) with the
AlphaFold2_advanced.ipynb notebook and the MMseqs2 MSA option. In all cases, the
five solutions predicted by AlphaFold2 by default were internally congruent, and we
always chose the one ranked first by the software. Uso1 GHD (1-674), Bos1, Bet1 and
RAB1 were initially submitted as hetero-oligomers. Subsequently, predictions were
submitted as 1:1 complexes of Uso1 GHD with Bos1, Bet1 and RAB1, as described in
the table below. The solutions were also very similar when comparing the different
combinations displayed in the table were fed to the software, strongly supporting the
validity of the results.
1090

AlphaFold2 runs, subunits included	
Uso1 GHD	Bet1

Uso1 GHD	Bet1	Bos1	
Uso1 GHD	Bet1	Bos1	Rab1
Uso1 GHD	Bet1 (1-49)		
Uso1 GHD (N-terminal)	Bet1 (1-49)		
Uso1 GHD	Bet1 (1-49)	Bos1 Habc (1-126)	
Uso1 GHD E6K/G540S	Bet1 (1-49)	Bos1 Habc (1-126)	
Uso1 GHD		Bos1	
Uso1 GHD		Bos1 Habc (1-126)	
Uso1 GHD dimer		Bos1 Habc (1-126)	
Uso1 GHD E6K/G540S		Bos1	
Uso1 GHD E6K/G540S		Bos1 Habc (1-126)	
Uso1			
Uso1 dimer			
Uso1 dimer E6K/G540S			
Bet1	Bos1	Sec22	Sed5
Rud3 dimer			
Coy1 dimer			
Grh1	Bug1		

1091

Acknowledgements

We thank Juan R. Luque (Molecular Interactions Facility, Centro de Investigaciones Biológicas) for his help with the analytical ultracentrifuge experiments, Manuel Sánchez-Berges for HA3-tagged SNARE strains, Sara Abib and Elena Reoyo for skillful technical assistance. Thanks are due to Spain's Ministerio de Ciencia e Innovación for grants RTI2018-093344-B100 (MAP) and predoctoral contract BES-2016-077440 (IB.-P/MAP), and to the Comunidad de Madrid for grant S2017/BMD-3691 (MAP). Grants were co-funded by European Regional Development and European Social Funds. The authors declare that they do not have any competing financial interests.

1101

Additional information

Competing interests

The authors declare that there are no competing interests

1105

Availability statement

All DNA molecules used here may reconstructed by standard techniques using primers listed in supplemental Table II. All strains listed under supplemental Table I are available for academic purposes upon reasonable request to the corresponding author. They are deposited and maintained by the corresponding laboratory in the so-denoted Madrid (MAD) collection.

Funding

Funding Agency	Grant reference number	Principal Investigator
Agencia Estatal de Investigación, Ministerio de Ciencia e Innovación, Spain	RTI2018-093344-B100	Miguel A. Peñalva
Agencia Estatal de Investigación, Ministerio de Ciencia e Innovación, Spain	predoctoral contract BES-2016-077440	Ignacio Bravo-Plaza
Agencia Estatal de Investigación, Ministerio de Ciencia e Innovación, Spain	PID2021-124278OB-I00	Miguel A. Peñalva & Eduardo A. Espeso, co-IPs
Comunidad de Madrid	S2017/BMD-3691	Miguel A. Peñalva
Grants co-funded by European Regional Development and European Social Funds		

Author contributions

Ignacio Bravo Plaza, Conceptualization, Data curation, Formal analysis, Validation, Investigation, Visualization, Writing — review and editing; Víctor G. Tagua, Data curation, Formal analysis, Validation, Investigation; Herbert N. Arst, Jr., Conceptualization, Data curation, Formal analysis, Validation, Investigation, Supervision, Writing — review and editing; Ana M. Alonso, Data curation, Formal analysis, Validation, Investigation; Mario Pinar, Conceptualization, Data curation, Formal analysis, Validation, Investigation, Supervision, Writing — review and editing ; Begoña Monterroso, Conceptualization, Data curation, Formal analysis, Validation, Supervision, Writing — review and editing; Antonio Galindo, Conceptualization, Data curation, Formal analysis, Validation, Investigation, Visualization, Writing — review and editing and Miguel Á. Peñalva, Conceptualization, Data curation, Formal analysis, Validation, Investigation, Supervision, Funding acquisition, Visualization, Writing – original draft, Writing — review and editing.

1131

Authors ORCIDs

Ignacio Bravo-Plaza (0000-0003-0934-9084) - ORCID

Victor García-Tagua (0000-0003-1494-6895) - ORCID

Ana M. Alonso (0000-0002-1228-7331) - ORCID

Mario Pinar (0000-0002-2415-8721) - ORCID

Begoña Monterroso (0000-0003-2538-084X)- ORCID

Antonio Galindo (0000-0002-1108-652X) - ORCID

Miguel A. Peñalva (0000-0002-3102-2806)- ORCID

1140

1141

1142

1143

1144

Legends to figures

Figure 1. Characterization of mutations bypassing the essential role of RAB1.

(A). Genetic map in the region surrounding *uso1* with genetic markers used as landmarks

for mapping.

(B). Molecular identification of the nucleotide changes in *suArab1ts* strains

(C) and (D): growth tests showing *rab1ts*- and *rab1Δ*-rescuing phenotypes, respectively,

of individual mutations, and synthetic positive interaction between E6K and G540S.

Strains produce either green or white conidiospores (conidiospore colors are used as

genetic markers). In (C), strains were point-inoculated. In (D) conidiospores were spread

on agar plates to give individual colonies.

Figure 1—figure supplement 1: E6K/G540S do not rescue lethality resulting from

***arf1Δ, sly1Δ* or *sed5Δ*.**

Top, *uso1* is an essential gene. Singly-nucleated conidiospores derived from a

heterokaryotic strain in which one class of nuclei carries a deficient *pyrG* uracil

biosynthetic gene whereas the second class contains a *uso1Δ* allele tagged with

functional *pyrG* were unable to grow on medium lacking pyrimidines at any of the

tested temperatures. Bottom: Similar experiments showing that unlike *rab1Δ*

strains, strains carrying lethal *arf1Δ, sly1Δ* and *sed5Δ* alleles cannot be rescued by

uso1$^{E6K/G540S}$. Top panel, strains with green conidiospores; bottom, strains with

white conidiospores.

Figure 2: Localization of the amino acid substitutions within the Uso1

AlphaFold2 structure

(A). AlphaFold2 cartoon representations of A. nidulans Uso1 in monomeric and dimeric

forms. Confidence estimations for this and other models are detailed under extended

data S2. Red, N-terminal tail; marine blue, globular head domain; gray, coiled-coil; green,

limit of the CTR. The rest of the CTR is shown as surface representation.

(B). Amino acid alignment of fungal sequences with mammalian p115 showing strong

conservation within the CTR: ANIDU, *Aspergillus nidulans;* PRUBE, *Penicillium rubens*;

TREES, *Thrichoderma ressei*; SSCLE, *Sclerotinia scleriotorum*; MORYZ, *Magnaporthe*

*oryza*e; CIMM, *Coccidioides immitis;* BOVIN, *Bos taurus*.

(C). Position of the Gly6Lys and Gly540Ser substitutions. Only the GHD of dimeric full-

length Uso1 are shown. The two different chains are colored in green and yellow,

respectively. Distances between mutated residues are displayed in armstrongs.

(D). The N-terminal amphipathic α-helix affected by the Glu6Lys substitution.

Figure 2—figure supplement 1. AlphaFold2 predictions of Uso1

Ribbon representation of AlphaFold 2-predicted structures of full-length Uso1 (A) and Uso1 GHD (B), color-coded by pLDDT values. Graphs at the bottom are the corresponding plots of predicted aligned error of the residues (PAE).

Figure 3: Determining molecular masses and oligomerization status of the different Uso1 constructs by velocity sedimentation analysis

The different panels display the sedimentation profiles of the protein being analyzed, with % of the main species, scheme of the different constructs and their limits and pictures of Coomassie stained-gels showing the purity of the protein preparations. The table below depicts biophysical parameters of the constructs used to obtain relative molecular masses. s_{exp} is the experimentally determined Svedberg coefficient; D_{exp}, translational diffusion coefficient of the main species; Mr, molecular mass deduced from Svedberg equation; M1 predicted molecular mass of the monomer; n = (Mr /M1).

Figure 3—figure supplement 1: GHD is a monomer across a range of concentrations

(A) and (B). Sedimentation velocity experiments with wild-type and E6K/G540S mutant GHD, respectively, showing that they behave as monomers at concentrations up to 5 µM.(C). Sedimentation velocity profile of E6K/G540S mutant GHD lacking the His-tag, showing that the presence of the latter does not interfere with oligomerization, and a picture of a Coomassie stained gel showing the purity of the protein preparation on the right.

Figure 4: Subcellular localization of Uso1

(A). Uso1-GFP localizing to punctate cytoplasmic structures, 3D shaded by software.

(B). Sections of a deconvolved Z-stack and its corresponding MIP. Uso1-GFP in inverted greyscale for clarity

(C). Kymograph showing the transient recruitment of Uso1 to punctate cytoplasmic structures.

(D). Average time of residence of Uso1 in these structures. Error bars, 95% CI.

(E). Example of one such structures visualized with a kymograph and with the corresponding movie frames (Movie 4).

Figure 4—figure supplement 1: Methodology for tracking the half-life of Uso1-GFP on punctate structures.

(1) 3D movies were acquired by streaming pictures to the computer RAM at 2 fps.

Appropriate reduction of excitation light intensity permitted acquisition of 400

frames without apparent phototoxicity.

(2) The behavior of punctate structures over time was represented in kymographs,

in which vertical lines represent the time of residence of Uso1 on membranes.

(3). As vertical lines frequently overlapped, jeopardizing the quality of this analysis,

we imported the time series into a 3D viewer as if they were (x, y, z) series. Rotation

across the different axes facilitated unambiguous tracking of the trajectories across

time.

(4) The length of the trajectories was measured and converted to time units.

Bottom graphs display examples of time trajectories.

1224

Figure 5: Uso1 puncta do not colocalize with ERESs

(A). Low extent co-localization of Sec13 ERES and Uso1 structures. Z-stacks for the two

channels were acquired simultaneously, deconvolved and represented as MIPs. Two

rare examples of colocalization are arrowed.

(B). Photograms of a dual channel Z-stack with a Sec13-labeled nuclear envelope

focused in the middle plane, illustrating that while some puncta show colocalization, the

red Sec13 signal and the green Uso1 signal do not usually overlap.

(C). A MIP of the same z-stack showing orthogonal views with some overlapping puncta

(arrows).

(D). A ts mutation in the *sarA* gene encoding the SarA^{Sar1} GTPase governing ER exit

markedly reduces the number of Uso1-GFP puncta upon shifting cells to restrictive

conditions. Box-and-whisker plots: Statistical comparison was made using one-way

ANOVA with Dunn's test for multiple comparisons. Whiskers are in Tukey's style: Only

significant differences were indicated, using asterisks.

1239

1240 **Figure 6: Uso1 localizes to RAB1-containing Golgi cisternae**

1241 (A). Tip cells showing Uso1 colocalization with the indicated subcellular markers. Images

1242 are MIPs of deconvolved Z-stacks.

1243 (B). Magnified images of the color-coded shaded regions of the cells shown in A.

1244 (C). Pearson's coefficients of the different combinations.

1245

Figure 7: Uso1 localization to punctate structures is dependent on RAB1

(A). Complete de-localization of Uso1-GFP to the cytosol by rab1ts and relocalization by

E6K/G540S.

(B). Uso1-GFP and *rab1ts* show a synthetic negative interaction that is rescued by the

E6K/G540S double substitution. Strains in lanes 7 and 8 carry the *wA2* mutation resulting

in white conidiospores.

1252

Figure 8. Genetic evidence showing that the CTR region of Uso1 contributes to

its recruitment to membranes.

(A). Top, scheme of the predicted interactions. Bottom, engineering a gene-replaced

allele lacking the CTR domain by homologous recombination.

(B). The Bug1 C-terminal residues fit into the groove formed between the two Grh1 PDZ

domains and into the pocket of the N-terminal PDZ domain (PDZ1).

(C). A gene-replaced *uso1ΔCTR* allele encoding a protein truncated for the CTR domain

shows a synthetic negative interaction with *rab1ts*.

(D). Western blot analysis. Removal of the CTR does not result in Uso1 instability.

(E). *bug1Δ* and *grh1Δ* show a synthetic negative interaction with *rab1ts* that is rescued

by the double E6K/G540S substitution in Uso1

Figure 8—figure supplement 1: AlphaFold2 modelling of Grh1-Bug1.

(A). Cartoon, with alpha-helices shown as cylinders, of the nearly N-terminal PDZ

domains of Grh1

(B). AlphaFold 2 prediction of a 1:1 Grh1-Bug1 complex, trimmed of disordered

regions

(C). complete AlphaFold2 model of Grh1-Bug1 with color-coded model

confidences values.

Figure 9: The GHD of Uso1 is sufficient to support cell viability

(A). Gene-replaced *uso1^{GHD}* allele carrying the double E6K/G540S substitution is

sufficient to rescue viability at 30°C, but not at higher temperatures. (B). The Uso1 GHD

is a monomer in vivo. Fractions collected from Superose columns loaded with the

indicated protein extracts and reference His-tagged proteins were collected and

analyzed by western blotting with α-HA and α-His antibodies. (C) Truncating Uso1 after

the GHD results in markedly reduced protein levels, as determined by α-Uso1 GHD

western blotting. The band (yellow asterisks) moving slower than Uso1 (magenta

asterisk) and at nearly the same position of Uso1 ΔCTR (blue asterisk) represents cross-

reacting contaminants unrelated to Uso1. The right panel shows a longer exposure for

the indicated region, to reveal the faint GHD band (green dot). (D). Overexpression of Uso1, wild-type and E6K/G540S mutant, under the control of the *inuA* promoter, which is turned off on glucose and induced on sucrose. Western blots reacted with α-Uso1 GHD antiserum. (E). Overexpressed GHD, be it E6K/G540S or wild-type, as the only source of Uso1 supports viability.

Figure 10: Screening the preferential association of proteins acting in the ER/Golgi interface with E6K/G540S Uso1.

(A). S-tagged baits (Uso1, wt and E6K/G540S, and the unrelated protein BapH), expressed after gene replacement, were captured with their associated polypeptides on S-protein agarose beads. Candidate associates, also expressed after gene replacement, were tagged with HA3. (B). Schematic depiction of the proteins listed in these experiments showing their sites of action.(C). Anti-HA3 western blot analysis of the indicated S-bait and HA3-prey combinations. Equal loading of Uso1 proteins was confirmed by silver staining of precipitates. Note that BapH, chosen as negative control, is expressed at much higher levels than Uso1 proteins. Each panel is a representative experiment of three experimental replicates.

Figure 10—figure supplement 1.

> (A) AlphaFold2 model, with PAE plot, of the GRIP domain of A. nidulans RUD3, predicted to be a dimer.

> (B) AlphaFold2 model of Coy1, calculated as a monomer. TMH is the nearly C-terminal transmembrane helix that contributes to its recruitment to membranes

Figure 10—figure supplement 2: Growth phenotypes of null mutants of genes encoding golgins.

> (A). Ablation of individual golgins Bug1/Grh1 and Rud3 does not result in detectable growth defects. *coy1*Δ strains have a subtle growth phenotype.

> (B). Negative effects of *grh1*Δ, *bug1*Δ, coy1Δ and *rud3*Δ on the ability of *uso1^{E6K/G540S}* to rescue *rab1*Δ. Note that *coy1*Δ and *rab1*Δ *uso1^{E6K/G540S}* are synthetically lethal. For convenience, this set of strains carried a mutation resulting in white conidiospores, as opposed to the wild-type green color.

Figure 11: The Uso1 GHD interacts directly with Bos1 and Bet1 SNAREs acting in the ER/Golgi interface

(A). Purified fusion proteins in which the cytosolic domains of the indicated SNAREs have been fused to GST were used in pulldown experiments with His-tagged, purified

wild-type and E6K/G540S Uso1. The plasma membrane Qa syntaxin Sso1 was used as

negative control. Pulled-down material was analyzed by anti-His western blotting. (B).

Quantitation of the above experiment; significance was determined by unpaired *t*-student

tests. Error bars represent S.E.M. (C). As in A, but using in vitro synthesized, HA3-tagged

Sly1 as prey. Samples were analyzed by anti-HA western blotting. (D). As in A, but using

wild-type and mutant GHD as preys, rather than full-length Uso1. (E). Quantitation of the

experiment in D. (F). GST pull-down experiment comparing the ability of the GHD to

interact with the early Golgi Qb and Qc SNAREs (Bos1 and Bet1), with that of their

medial Golgi counterparts (Qb Gos1 and Qc Sft1).

1325

Figure 11—figure supplement 1. AlphaFold2 prediction of the ER/Golgi SNARE

bundle.

(A) Sec5/Bos1/Bet1/Sec22 predicted SNARE bundle. (B) Quality control (pLDDT,

color coded, and PAE) of the model.

Figure 12. AlphaFold2 models provide insight into the additive mode of

suppression shown by E6K and G540S.

(A). Model of full length Uso1 bound to the ER/Golgi SNAREs Bos1 and Bet1.

(B). Top, ribbon representation of the Bos1 N-terminal Habc domain and Uso1GHD.

Bottom, Inset combining surface and ribbon depiction.

(C). Increased binding of Bos1 to G540S Uso1 appears to involve insertion of Ser540

into a pocket located in the Habc domain of the Qb SNARE. Partial view of the Bos1-

Uso1 GHD surface of interaction in the wild type (left) and mutant (right) models. G540

and S540 are annotated.

(D). The N-terminal amphipathic α-helix of Uso1 comprising the E6K substitution lies

within a flexible stretch of the protein that might facilitate its insertion into membranes.

Alignment of six independent predictions, with Glu6 highlighted in red. The Uso1 GHD

was modeled alone, in a complex with SNARE proteins or with Ypt1. The N-terminal a-

helix (boxed) adopts different positions, suggesting high flexibility.

Figure 12—figure supplement 1. AlphaFold2 prediction of the Bet1-GHD

interaction.

The putative binding surface of Bet1 and Uso1 as determined by AlphaFold2. Top

images, cartoon of Bet1-GHD interactions, colored by pLDDT score. Alignment of

four independent predictions involving the Bet1 N-terminal region and Uso1 GHD.

A single model for Uso1 GHD is shown on the top representation for simplicity. In

spite of the disordered nature of the N-terminal Bet1 region, the Bet1-Uso1 binding

interface is consistent among models. Bottom, surface representation of the N-terminal Bet1 region (orange) in complex with the GHD. Also indicated is the Habc domain of Bos1 (yellow) bound to the GHD.

Figure 12—figure supplement 2. AlphaFold 2 prediction of the RAB1 binding site on the Bet1/Bos1/Uso1 GHD complex.

(A) and (B): cartoon representation of the GHD-RAB1 complex. The model is depicted as pipes and planks

(C): Orthogonal views of the Uso1 GHD-RAB1-Bet1-Bos1Habc structural model. The Uso1 GHD is shown as surface to emphasize the distant binding sites of the Bos1 Habc domain, RAB1 and the Bet1 N-terminal region

(D): Ribbon representation of the model shown in C but including the full-length SNARE subunits, i.e. the GHD domain of Uso1, RAB1 and the SNARES Bet1 and Bos1. Proteins are in the correct orientation to connect membranes separated by 23 nm, counting from the SNARE TMDs to the prenylated RAB1 residues.

Figure 12—figure supplement 3. Quality control assessment of AlphaFold2 predictions for the indicated complexes.

Supplemental Table I

List and complete genotypes of *A. nidulans* strains used in this work

Supplemental Table II

Primers used for PCR-based genetic manipulations

Figure 8 source data

Raw images for western blots in panel D and uncropped pictures with used exposures and regions indicated.

Figure 9 source data

For panels B, C, D; raw images for western blots and uncropped pictures with used exposures and regions indicated.

Figure 10 source data

Raw images for western blots and silver-stained gels and uncropped pictures with used exposures and regions indicated.

1383 **Figure 11 source data**

1384 Raw images for western blots and Coomassie-stained gels and uncropped pictures

1385 with used exposures and regions indicated.

1386

Rich file media

Video 1: Shaded 3D reconstruction of a hypha expressing Uso1-GFP

Video 2: 4D acquisition showing the dynamics of Uso1-GFP.

4D (x, y, z, t) in which Z-stacks were acquired at a rate of 1 frame every 2.6 sec

Video 3: Dynamics of Uso1-GFP at 2 fps

3D acquisition (200 frames) showing the dynamics of Uso1-GFP. Time resolution, 2 fps

Video 4: Single Uso1-GFP cisterna tracked over time

Example of Uso1-GFP cisterna. The video contains 96 photograms acquired at 2fps

Video 5: ·3D reconstruction of a hypha expressing fluorescently labeled Uso1-

GFP and Sec13-mCh

There is little colocalization between Uso1-GFP and Sec13 ERES

Video 6: ·4D video (1 fpm) of a hypha expressing fluorescently labeled Uso1-GFP

and Sec7-mCh

Uso1 does not colocalize at all with the TGN marker Sec7

1401

1402

1403   **References**

1404
1405
Allan, B.B., B.D. Moyer, and W.E. Balch. 2000. Rab1 recruitment of p115 into a cis-
SNARE complex: programming budding COPII vesicles for fusion. *Science*.
289:444-448. 10.1126/science.289.5478.444

An, Y., C.Y. Chen, B. Moyer, P. Rotkiewicz, M.A. Elsliger, A. Godzik, I.A. Wilson, and
W.E. Balch. 2009. Structural and functional analysis of the globular head
domain of p115 provides insight into membrane tethering. *J Mol Biol*. 391:26-
41. PM:19414022

Anderson, N.S., I. Mukherjee, C.M. Bentivoglio, and C. Barlowe. 2017. The Golgin
protein Coy1 functions in intra-Golgi retrograde transport and interacts with the
COG complex and Golgi SNAREs. *Mol Biol Cell*. 10.1091/mbc.E17-03-0137

Arst, H.N., Jr., M. Hernández-González, M.A. Peñalva, and A. Pantazopoulou. 2014.
GBF/Gea mutant with a single substitution sustains fungal growth in the
absence of BIG/Sec7. *FEBS Lett*. 588:4799-4786.
10.1016/j.febslet.2014.11.014

Baker, R.W., and F.M. Hughson. 2016. Chaperoning SNARE assembly and
disassembly. *Nat Rev Mol Cell Biol*. 17:465-479. 10.1038/nrm.2016.65

Baker, R.W., P.D. Jeffrey, M. Zick, B.P. Phillips, W.T. Wickner, and F.M. Hughson.
2015. A direct role for the Sec1/Munc18-family protein Vps33 as a template for
SNARE assembly. *Science*. 349:1111-1114. 10.1126/science.aac7906

Barlowe, C.K., and E.A. Miller. 2013. Secretory protein biogenesis and traffic in the
early secretory pathway. *Genetics*. 193:383-410.
http://www.ncbi.nlm.nih.gov/pubmed/23396477

Beard, M., A. Satoh, J. Shorter, and G. Warren. 2005. A cryptic Rab1-binding site in
the p115 tethering protein. *J Biol Chem*. 280:25840-25848.
10.1074/jbc.M503925200

Behnia, R., F.A. Barr, J.J. Flanagan, C. Barlowe, and S. Munro. 2007. The yeast
orthologue of GRASP65 forms a complex with a coiled-coil protein that
contributes to ER to Golgi traffic. *J Cell Biol*. 176:255-261. PM:17261844

Bentley, M., Y. Liang, K. Mullen, D. Xu, E. Sztul, and J.C. Hay. 2006. SNARE Status
Regulates Tether Recruitment and Function in Homotypic COPII Vesicle
Fusion*. *J Biol Chem*. 281:38825-38833.
https://doi.org/10.1074/jbc.M606044200

Bracher, A., and W. Weissenhorn. 2002. Structural basis for the Golgi membrane
recruitment of Sly1p by Sed5p. *EMBO J*. 21:6114-6124. PM:12426383

Brandon, E., T. Szul, C. Alvarez, R. Grabski, R. Benjamin, R. Kawai, and E. Sztul.
2006. On and off membrane dynamics of the endoplasmic reticulum-golgi
tethering factor p115 in vivo. *Mol Biol Cell*. 17:2996-3008. 10.1091/mbc.E05-09-
0862

Bravo-Plaza, I., M. Hernandez-Gonzalez, M. Pinar, J.F. Diaz, and M.A. Penalva. 2019.
Identification of the guanine nucleotide exchange factor for SAR1 in the
filamentous fungal model Aspergillus nidulans. *Biochim Biophys Acta Mol Cell*
Res. 1866:118551. 10.1016/j.bbamcr.2019.118551

Cai, H., S. Yu, S. Menon, Y. Cai, D. Lazarova, C. Fu, K. Reinisch, J.C. Hay, and S. Ferro-Novick. 2007. TRAPPI tethers COPII vesicles by binding the coat subunit Sec23. *Nature*. 445:941-944. PM:17287728

Cao, X., N. Ballew, and C. Barlowe. 1998. Initial docking of ER-derived vesicles requires Uso1p and Ypt1p but is independent of SNARE proteins. *EMBO J*. 17:2156-2165. 10.1093/emboj/17.8.2156

Cove, D.J. 1966. The induction and repression of nitrate reductase in the fungus *Aspergillus nidulans. Biochim Biophys Acta*. 113:51-56. 10.1016/S0926-6593(66)80120-0

Day, K.J., L.A. Staehelin, and B.S. Glick. 2013. A three-stage model of Golgi structure and function. *Histochem Cell Biol*. 140:239-249. 10.1007/s00418-013-1128-3

Galindo, A., V.J. Planelles-Herrero, G. Degliesposti, and S. Munro. 2021. Cryo-EM structure of metazoan TRAPPIII, the multi-subunit complex that activates the GTPase Rab1. *Embo J*. 40:e107608. https://doi.org/10.15252/embj.2020107608

Gillingham, A.K. 2018. At the ends of their tethers! How coiled-coil proteins capture vesicles at the Golgi. *Biochem Soc Trans*. 46:43-50. 10.1042/BST20170188

Gillingham, A.K., and S. Munro. 2016. Finding the Golgi: Golgin Coiled-Coil Proteins Show the Way. *Trends Cell Biol*. 26:339-408. http://dx.doi.org/10.1016/j.tcb.2016.02.005

Heo, Y., H.-J. Yoon, H. Ko, S. Jang, and H.H. Lee. 2020. Crystal structures of Uso1 membrane tether reveal an alternative conformation in the globular head domain. *Scientific Reports*. 10:9544. 10.1038/s41598-020-66480-1

Hernández-González, M., I. Bravo-Plaza, V. de Los Ríos, M. Pinar, A. Pantazopoulou, and M.A. Peñalva. 2019. COPI localizes to the early Golgi in *Aspergillus nidulans. Fungal Genet Biol*. 123:78-86. 10.1016/j.fgb.2018.12.003

Hernández-González, M., A. Pantazopoulou, D. Spanoudakis, C.L.C. Seegers, and M.A. Peñalva. 2018. Genetic dissection of the secretory route followed by a fungal extracellular glycosyl hydrolase. *Mol Microbiol*. 109:781-800. doi:10.1111/mmi.14073

Hu, F., X. Shi, B. Li, X. Huang, X. Morelli, and N. Shi. 2015. Structural basis for the interaction between the Golgi reassembly-stacking protein GRASP65 and the Golgi matrix protein GM130. *J Biol Chem*. 290:26373-26382. 10.1074/jbc.M115.657940

Jiang, Y., A. Scarpa, L. Zhang, S. Stone, E. Feliciano, and S. Ferro-Novick. 1998. A high copy suppressor screen reveals genetic interactions between BET3 and a new gene. Evidence for a novel complex in ER-to-Golgi transport. *Genetics*. 149:833-841. 10.1093/genetics/149.2.833

Joiner, A.M., B.P. Phillips, K. Yugandhar, E.J. Sanford, M.B. Smolka, H. Yu, E.A. Miller, and J.C. Fromme. 2021. Structural basis of TRAPPIII-mediated Rab1 activation. *Embo J*. 40:e107607. https://doi.org/10.15252/embj.2020107607

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Zidek, A. Potapenko, A. Bridgland, C. Meyer, S.A.A. Kohl, A.J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A.W. Senior, K. Kavukcuoglu, P. Kohli, and D. Hassabis. 2021. Highly accurate protein structure prediction with AlphaFold. *Nature*. 596:583-589. 10.1038/s41586-021-03819-2

Lord, C., D. Bhandari, S. Menon, M. Ghassemian, D. Nycz, J. Hay, P. Ghosh, and S. Ferro-Novick. 2011. Sequential interactions with Sec23 control the direction of vesicle traffic. *Nature*. 473:181-186. 10.1038/nature09969

Lupashin, V.V., and M.G. Waters. 1997. t-SNARE activation through transient interaction with a Rab-like guanosine triphosphatase. *Science*. 276:1255-1258. http://www.ncbi.nlm.nih.gov/pubmed/9157884

Malsam, J., and T.H. Sollner. 2011. Organization of SNAREs within the Golgi stack. *Cold Spring Harb Perspect Biol*. 3:a005249. 10.1101/cshperspect.a005249

McNew, J.A., F. Parlati, R. Fukuda, R.J. Johnston, K. Paz, F. Paumet, T.H. Sollner, and J.E. Rothman. 2000. Compartmental specificity of cellular membrane fusion encoded in SNARE proteins. *Nature*. 407:153-159. 10.1038/35025000.

Mirdita, M., K. Schütze, Y. Moriwaki, L. Heo, S. Ovchinnikov, and M. Steinegger. 2022. ColabFold: making protein folding accessible to all. *Nature Methods*. 19:679-682. 10.1038/s41592-022-01488-1

Muschalik, N., and S. Munro. 2018. Golgins. *Curr Biol*. 28:R374-R376. https://doi.org/10.1016/j.cub.2018.01.006

Nakajima, H., A. Hirata, Y. Ogawa, T. Yonehara, K. Yoda, and M. Yamasaki. 1991. A cytoskeleton-related gene, uso1, is required for intracellular protein transport in Saccharomyces cerevisiae. *J Cell Biol*. 113:245-260. 10.1083/jcb.113.2.245

Nayak, T., E. Szewczyk, C.E. Oakley, A. Osmani, L. Ukil, S.L. Murray, M.J. Hynes, S.A. Osmani, and B.R. Oakley. 2005. A versatile and efficient gene targeting system for *Aspergillus nidulans. Genetics*. 172:1557-1566. 10.1534/genetics.105.052563

Osmani, S.A., D.B. Engle, J.H. Doonan, and N.R. Morris. 1988. Spindle formation and chromatin condensation in cells blocked in interphase by mutation of a negative cell cycle control gene. *Cell*. 52:241-251. 10.1016/0092-8674(88)90513-2.

Pantazopoulou, A., and B.S. Glick. 2019. A Kinetic View of Membrane Traffic Pathways Can Transcend the Classical View of Golgi Compartments. *Frontiers in Cell and Developmental Biology*. 7:153. https://www.frontiersin.org/article/10.3389/fcell.2019.00153

Pantazopoulou, A., and M.A. Peñalva. 2011. Characterization of *Aspergillus nidulans* RabC[Rab6]. *Traffic*. 12:386-406. 10.1111/j.1600-0854.2011.01164.x

Parlati, F., O. Varlamov, K. Paz, J.A. McNew, D. Hurtado, T.H. Sollner, and J.E. Rothman. 2002. Distinct SNARE complexes mediating membrane fusion in Golgi transport based on combinatorial specificity. *Proc Natl Acad Sci USA*. 99:5424-5429. 10.1073/pnas.082100899

Pelham, H.R. 1999. SNAREs and the secretory pathway-lessons from yeast. *Experimental Cell Research*. 247:1-8. https://doi.org/10.1006/excr.1998.4356

Pelham, H.R. 2001. SNAREs and the specificity of membrane fusion. *Trends Cell Biol*. 11:99-101.

Peng, R., and D. Gallwitz. 2002. Sly1 protein bound to Golgi syntaxin Sed5p allows assembly and contributes to specificity of SNARE fusion complexes. *J Cell Biol*. 157:645-655. 10.1083/jcb.200202006

Peñalva, M.A. 2005. Tracing the endocytic pathway of *Aspergillus nidulans* with FM4-64. *Fungal Genet Biol*. 42:963-975. 10.1016/j.fgb.2005.09.004

Peñalva, M.A., E. Moscoso-Romero, and M. Hernández-González. 2020. Tracking exocytosis of a GPI-anchored protein in *Aspergillus nidulans*. *Traffic*. 21:675-688. https://doi.org/10.1111/tra.12761

Pinar, M., A. Alonso, V. de los Ríos, I. Bravo-Plaza, Á. de la Gándara, A. Galindo, E. Arias-Palomo, and M.Á. Peñalva. 2022. The type V myosin-containing complex HUM is a RAB11 effector powering movement of secretory vesicles. *iScience*. 25. 10.1016/j.isci.2022.104514

Pinar, M., E. Arias-Palomo, V. de los Ríos, H.N. Arst, Jr., and M.A. Peñalva. 2019. Characterization of *Aspergillus nidulans* TRAPPs uncovers unprecedented similarities with metazoans and reveals the modular assembly of TRAPPII. *PLoS Genetics*. 15:e1008557. https://doi.org/10.1371/journal.pgen.1008557

Pinar, M., H.N. Arst, Jr., A. Pantazopoulou, V.G. Tagua, V. de los Ríos, J. Rodríguez-Salarichs, J.F. Díaz, and M.A. Peñalva. 2015. TRAPPII regulates exocytic Golgi exit by mediating nucleotide exchange on the Ypt31 orthologue RabE/RAB11. *Proc Natl Acad Sci USA*. 112:4346-4351. 10.1073/pnas.1419168112

Pinar, M., A. Pantazopoulou, H.N. Arst, Jr., and M.A. Peñalva. 2013. Acute inactivation of the *Aspergillus nidulans* Golgi membrane fusion machinery: correlation of apical extension arrest and tip swelling with cisternal disorganization. *Mol Microbiol*. 89:228-248. 10.1111/mmi.12280

Pinar, M., and M.A. Peñalva. 2017. *Aspergillus nidulans* BapH is a RAB11 effector that connects membranes in the Spitzenkörper with basal autophagy. *Mol Microbiol*. 106:452-468. 10.1111/mmi.13777

Pinar, M., and M.A. Peñalva. 2020. *En bloc* TGN recruitment of *Aspergillus* TRAPPII reveals TRAPP maturation as unlikely to drive RAB1-to-RAB11 transition. *J Cell Sci*. 133:jcs241141. 10.1242/jcs.241141

Pinar, M., and M.A. Peñalva. 2021. The fungal RABOME: RAB GTPases acting in the endocytic and exocytic pathways of Aspergillus nidulans (with excursions to other filamentous fungi). *Mol Microbiol*. 116:53-70. https://doi.org/10.1111/mmi.14716

Ren, Y., C.K. Yip, A. Tripathi, D. Huie, P.D. Jeffrey, T. Walz, and F.M. Hughson. 2009. A structure-based mechanism for vesicle capture by the multisubunit tethering complex Dsl1. *Cell*. 139:1119-1129. 10.1016/j.cell.2009.11.002

Rexach, M.F., M. Latterich, and R.W. Schekman. 1994. Characteristics of endoplasmic reticulum-derived transport vesicles. *J Cell Biol*. 126:1133-1148. 10.1083/jcb.126.5.1133

Riedel, F., A. Galindo, N. Muschalik, and S. Munro. 2017. The two TRAPP complexes of metazoans have distinct roles and act on different Rab GTPases. *J Cell Biol*. 217:601-617. 10.1083/jcb.201705068

Rizo, J., and T.C. Sudhof. 2012. The membrane fusion enigma: SNAREs, Sec1/Munc18 proteins, and their accomplices--guilty as charged? *Annu Rev Cell Dev Biol*. 28:279-308. 10.1146/annurev-cellbio-101011-155818

Sapperstein, S.K., V.V. Lupashin, H.D. Schmitt, and M.G. Waters. 1996. Assembly of the ER to Golgi SNARE complex requires Uso1p. *J Cell Biol*. 132:755-767. http://www.ncbi.nlm.nih.gov/pubmed/8603910

Sapperstein, S.K., D.M. Walter, A.R. Grosvenor, J.E. Heuser, and M.G. Waters. 1995. p115 is a general vesicular transport factor related to the yeast endoplasmic reticulum to Golgi transport factor Uso1p. *Proc Natl Acad Sci USA*. 92:522-526. 10.1073/pnas.92.2.522

Schuck, P. 2000. Size-distribution analysis of macromolecules by sedimentation velocity ultracentrifugation and lamm equation modeling. *Biophys J*. 78:1606-1619. 10.1016/S0006-3495(00)76713-0

Seog, D.H., M. Kito, K. Yoda, and M. Yamasaki. 1994. Uso1 protein contains a coiled-coil rod region essential for protein transport from the ER to the Golgi apparatus in Saccharomyces cerevisiae. *J Biochem*. 116:1341-1345. 10.1093/oxfordjournals.jbchem.a124685

Shorter, J., M.B. Beard, J. Seemann, A.B. Dirac-Svejstrup, and G. Warren. 2002. Sequential tethering of Golgins and catalysis of SNAREpin assembly by the vesicle-tethering protein p115. *J Cell Biol*. 157:45-62. 10.1083/jcb.200112127

Sogaard, M., K. Tani, R.R. Ye, S. Geromanos, P. Tempst, T. Kirchhausen, J.E. Rothman, and T. Sollner. 1994. A rab protein is required for the assembly of SNARE complexes in the docking of transport vesicles. *Cell*. 78:937-948. 10.1016/0092-8674(94)90270-4

Striegl, H., Y. Roske, D. Kummel, and U. Heinemann. 2009. Unusual armadillo fold in the human general vesicular transport factor p115. *PLoS ONE*. 4:e4656. 10.1371/journal.pone.0004656

Szewczyk, E., T. Nayak, C.E. Oakley, H. Edgerton, Y. Xiong, N. Taheri-Talesh, S.A. Osmani, and B.R. Oakley. 2006. Fusion PCR and gene targeting in *Aspergillus nidulans*. *Nat Protoc*. 1:3111-3120. 10.1038/nprot.2006.405

Thomas, L.L., A.M.N. Joiner, and J.C. Fromme. 2018. The TRAPPIII complex activates the GTPase Ypt1 (Rab1) in the secretory pathway. *J Cell Biol*. 217:283-298. 10.1083/jcb.201705214

Thomas, L.L., S.A. van der Vegt, and J.C. Fromme. 2019. A Steric Gating Mechanism Dictates the Substrate Specificity of a Rab-GEF. *Dev Cell*. 48:100-114.e109. https://doi.org/10.1016/j.devcel.2018.11.013

Tilburn, J., C. Scazzocchio, G.G. Taylor, J.H. Zabicky-Zissman, R.A. Lockington, and R.W. Davies. 1983. Transformation by integration in *Aspergillus nidulans*. *Gene*. 26:205-211. 10.1016/0378-1119(83)90191-9.

Todd, R.B., M.A. Davis, and M.J. Hynes. 2007. Genetic manipulation of Aspergillus nidulans : meiotic progeny for genetic analysis and strain construction. *Nat Protoc*. 2:811-821. 10.1038/nprot.2007.113.

Tsui, M.M., W.C. Tai, and D.K. Banfield. 2001. Selective formation of Sed5p-containing SNARE complexes is mediated by combinatorial binding interactions. *Mol Biol Cell*. 12:521-538. http://www.ncbi.nlm.nih.gov/pubmed/11251068

Wang, T., R. Grabski, E. Sztul, and J.C. Hay. 2014. p115-SNARE Interactions: A Dynamic Cycle of p115 Binding Monomeric SNARE Motifs and Releasing Assembled Bundles. *Traffic*. 10.1111/tra.12242

Weigel, A.V., C.L. Chang, G. Shtengel, C.S. Xu, D.P. Hoffman, M. Freeman, N. Iyer, J. Aaron, S. Khuon, J. Bogovic, W. Qiu, H.F. Hess, and J. Lippincott-Schwartz. 2021. ER-to-Golgi protein delivery through an interwoven, tubular network extending from ER. *Cell*. 184:2412-2429 e2416. 10.1016/j.cell.2021.03.035

Yamakawa, H., D.H. Seog, K. Yoda, M. Yamasaki, and T. Wakabayashi. 1996. Uso1 protein is a dimer with two globular heads and a long coiled-coil tail. *Journal of Structural Biology*. 116:356-365. 10.1006/jsbi.1996.0053

Yu, I.M., and F.M. Hughson. 2010. Tethering factors as organizers of intracellular vesicular traffic. *Annu Rev Cell Dev Biol*. 26:137-156. 10.1146/annurev.cellbio.042308.113327

1638   Zhang, Y., and F.M. Hughson. 2021. Chaperoning SNARE Folding and Assembly.
1639   *Annual Review of Biochemistry*. 90:581-603. 10.1146/annurev-biochem-
1640   081820-103615

1641   Zhang, Y., and J. Yang. 2020. Securing SNAREs for assembly. *J Biol Chem*.
1642   295:10136-10137. 10.1074/jbc.H120.014815

1643

Figure 1. Characterization of mutations bypassing the essential role of RAB1

(A). Genetic map in the region surrounding *uso1* with genetic markers used as landmarks for mapping.B). Molecular identification of the nucleotide changes in *suArab1^ts* strains

(C) and (D): growth tests showing *rab1^ts*- and *rab1Δ*-rescuing phenotypes, respectively, of individual mutations, and synthetic positive interaction between E6K and G540S. Strains produce either green or white conidiospores (conidiospore colors are used as genetic markers). In (C), strains were point-inoculated. In (D) conidiospores were spread on agar plates to give individual colonies.

Figure 1—figure supplement 1: E6K/G540S do not rescue lethality resulting from *arf1∆*, *sly1∆* or *sed5∆*.

Top, *uso1* is an essential gene. Singly-nucleated conidiospores derived from a heterokaryotic strain in which one class of nuclei carries a deficient *pyrG* uracil biosynthetic gene whereas the second class contains a *uso1∆* allele tagged with functional *pyrG* were unable to grow on medium lacking pyrimidines at any of the tested temperatures. Bottom: Similar experiments showing that unlike *rab1∆* strains, strains carrying lethal *arf1∆ ,sly1∆* and *sed5∆* alleles cannot be rescued by *uso1^{E6K/G540S}*. Top panel, strains with green conidiospores; bottom, strains with white conidiospores.

Figure 2: Localization of the amino acid substitutions within the Uso1 AlphaFold2 structure

(A). AlphaFold2 cartoon representations of *A. nidulans* Uso1 in monomeric and dimeric forms. Red, N-terminal tail; marine blue, globular head domain; gray, coiled-coil; green, limit of the CTR. The rest of the CTR is shown as surface representation.

(B). Amino acid alignment of fungal sequences with mammalian p115 showing strong conservation within the CTR: ANIDU, *Aspergillus nidulans*; PRUBE, *Penicillium rubens*; TREES, *Thrichoderma ressei;* SSCLE, *Sclerotinia scleriotorum*; MORYZ, *Magnaporthe oryzae*; CIMM, *Coccidioides immitis*; BOVIN, *Bos taurus*.

©. Position of the Gly6Lys and Gly540Ser substitutions. Only the GHD of dimeric full-length Uso1 are shown. The two different chains are colored in green and yellow, respectively. Distances between mutated residues are displayed in armstrongs.

(D). The N-terminal amphipathic α–helix affected by the Glu6Lys substitution.

Figure 2— figure supplement 1. AlphaFold2 predictions of Uso1
Ribbon representation of AlphaFold 2-predicted structures of full-length Uso1 (A) and
Uso1 GHD (B), color-coded by pLDDT values. Graphs at the bottom are the
corresponding plots of predicted aligned error of the residues (PAE).

Figure 3: Determining molecular masses and oligomerization status of the different Uso1 constructs by velocity sedimentation analysis

The different panels display the sedimentation profiles of the protein being analyzed, with % of the main species, scheme of the different constructs and their limits and pictures of Coomassie stained-gels showing the purity of the protein preparations. The table below depicts biophysical parameters of the constructs used to obtain relative molecular masses. s_{exp} is the experimentally determined Svedberg coefficient; D_{exp}, translational diffusion coefficient of the main species; Mr, molecular mass deduced from Svedberg equation; M_1 predicted molecular mass of the monomer; $n = (M_r/M_1)$

Figure 3—figure supplement 1: GHD is a monomer across a range of concentrations
(A) and (B). Sedimentation velocity experiments with wild-type and E6K/G540S mutant GHD, respectively, showing that they behave as monomers at concentrations up to 5 μM.(C). Sedimentation velocity profile of E6K/G540S mutant GHD lacking the His-tag, showing that the presence of the latter does not interfere with oligomerization, and a picture of a Coomassie stained gel showing the purity of the protein preparation on the right.

Figure 4: Subcellular localization of Uso1

(A). Uso1-GFP localizing to punctate cytoplasmic structures, 3D shaded by software.

(B). Sections of a deconvolved Z-stack and its corresponding MIP. Uso1-GFP in inverted greyscale for clarity

(C). Kymograph showing the transient recruitment of Uso1 to punctate cytoplasmic structures.

(D). Average time of residence of Uso1 in these structures. Error bars, 95% CI.

(E). Example of one such structures visualized with a kymograph and with the corresponding movie frames (Movie 4).

① With wide-field optics and a 1.4 NA, 63x objetive stream frames to the computer RAM at a time resolution of 2 fps.

③ Convert time series into x,y,z series and visualize with 3D software to resove overlaps

② Draw kymograph x,y,t a number of Uso1 traces cannot be resolved because they overlap

④ measure lifetime of individual cisternae as the length of vertical lines & plot

Figure 4—figure supplement 1: Methodology for tracking the half-life of Uso1-GFP on punctate structures.

1) 3D movies were acquired by streaming pictures to the computer RAM at 2 fps. Appropriate reduction of excitation light intensity permitted acquisition of 400 frames without apparent phototoxicity.

2) The behavior of punctate structures over time was represented in kymographs, in which vertical lines represent the time of residence of Uso1 on membranes.

3). As vertical lines frequently overlapped, jeopardizing the quality of this analysis, we imported the time series into a 3D viewer as if they were (x, y, z) series. Rotation across the different axes facilitated unambiguous tracking of the trajectories across time.

4) The length of the trajectories was measured and converted to time units. Bottom graphs display examples of time trajectories.

Figure 5: Uso1 puncta do not colocalize with ERESs

(A). Low extent co-localization of Sec13 ERES and Uso1 structures. Z-stacks for the two channels were acquired simultaneously, deconvolved and represented as MIPs. Two rare examples of colocalization are arrowed.

(B). Photograms of a dual channel Z-stack with a Sec13-labeled nuclear envelope focused in the middle plane, illustrating that while some puncta show colocalization, the red Sec13 signal and the green Uso1 signal do not usually overlap.

(C). A MIP of the same z-stack showing orthogonal views with some overlapping puncta (arrows).

(D). A *ts* mutation in the *sarA* gene encoding the SarA^{Sar1} GTPase governing ER exit markedly reduces the number of Uso1-GFP puncta upon shifting cells to restrictive conditions. Box-and-whisker plots: Statistical comparison was made using one-way ANOVA with Dunn's test for multiple comparisons. Whiskers are in Tukey's style: Only significant differences were indicated, using asterisks.

Figure 6: Uso1 localizes to RAB1-containing Golgi cisternae

(A). Tip cells showing Uso1 colocalization with the indicated subcellular markers. Images are MIPs of deconvolved Z-stacks.

(B). Magnified images of the color-coded shaded regions of the cells shown in A.

©. Pearson's coefficients of the different combinations

Figure 7: Uso1 localization to punctate structures is dependent on RAB1

(A). Complete de-localization of Uso1-GFP to the cytosol by *rab1ts* and relocalization by E6K/G540S.

(B). Uso1-GFP and *rab1ts* show a synthetic negative interaction that is rescued by the E6K/G540S double substitution. Strains in lanes 7 and 8 carry the *wA2* mutation resulting in white conidiospores.

Figure 8. Genetic evidence showing that the CTR region of Uso1 contributes to its recruitment to membranes.

(A). Top, scheme of the predicted interactions. Bottom, engineering a gene-replaced allele lacking the CTR domain by homologous recombination.

(B). The Bug1 C-terminal residues fit into the groove formed between the two Grh1 PDZ domains and into the pocket of the N-terminal PDZ domain (PDZ1).

(C). A gene-replaced *uso1ΔCTR* allele encoding a protein truncated for the CTR domain shows a synthetic negative interaction with *rab1ts.*

(D). Western blot analysis. Removal of the CTR does not result in Uso1 instability.

(E). *bug1Δ* and *grh1Δ* show a synthetic negative interaction with *rab1ts* that is rescued by the double E6K/G540S substitution in Uso1

Figure 8 —figure supplement 1: AlphaFold2 modelling of Grh1-Bug1.
(A). Cartoon, with alpha-helices shown as cylinders, of the nearly N-terminal PDZ domains of Grh1
(B). AlphaFold 2 prediction of a 1:1 Grh1-Bug1 complex, trimmed of disordered regions
©. complete AlphaFold2 model of Grh1-Bug1 with color-coded model confidence values.

Figure 9: The GHD of Uso1 is sufficient to support cell viability
(A). Gene-replaced *uso1*^GHD^ allele carrying the double E6K/G540S substitution is sufficient to rescue viability at 30°C, but not at higher temperatures. (B). The Uso1 GHD is a monomer in vivo. Fractions collected from Superose columns loaded with the indicated protein extracts and reference His-tagged proteins were collected and analyzed by western blotting with α−HA and α−His antibodies. (C) Truncating Uso1 after the GHD results in markedly reduced protein levels, as determined by α-Uso1 GHD western blotting. The band (yellow asterisks) moving slower than Uso1 (magenta asterisk) and at nearly the same position of Uso1 ΔCTR (blue asterisk) represents cross-reacting contaminants unrelated to Uso1. The right panel shows a longer exposure for the indicated region, to reveal the faint GHD band (green dot). (D). Overexpression of Uso1, wild-type and E6K/G540S mutant, under the control of the *inuA* promoter, which is turned off on glucose and induced on sucrose. Western blots reacted with α−Uso1 GHD antiserum. (E). Overexpressed GHD, be it E6K/G540S or wild-type, as the only source of Uso1 supports viability.

Figure 10: Screening the preferential association of proteins acting in the ER/Golgi interface with E6K/G540S Uso1.

(A). S-tagged baits (Uso1, wt and E6K/G540S, and the unrelated protein BapH), expressed after gene replacement, were captured with their associated polypeptides on S-protein agarose beads. Candidate associates, also expressed after gene replacement, were tagged with HA3. (B). Schematic depiction of the proteins listed in these experiments showing their sites of action.(C). Anti-HA3 western blot analysis of the indicated S-bait and HA3-prey combinations. Equal loading of Uso1 proteins was confirmed by silver staining of precipitates. Note that BapH, chosen as negative control, is expressed at much higher levels than Uso1 proteins. Each panel is a representative experiment of three experimental replicates.

Figure 10—figure supplement 1.
/A) AlphaFold2 model, with PAE plot, of the GRIP domain of *A. nidulans* RUD3, predicted to be a dimer.
(B) AlphaFold2 model of Coy1, calculated as a monomer. TMH is the nearly C-terminal transmembrane helix that contributes to its recruitment to membranes

Figure 10—figure supplement 2: Growth phenotypes of null mutants of genes encoding golgins.

(A). Ablation of individual golgins Bug1/Grh1 and Rud3 does not result in detectable growth defects. *coy1Δ* strains have a subtle growth phenotype.

(B). Negative effects of *grh1Δ, bug1Δ, coy1Δ* and *rud3Δ* on the ability of *uso1*$^{E6K/G540S}$ to rescue *rab1Δ*. Note that *coy1Δ* and *rab1Δ uso1*$^{E6K/G540S}$ are synthetically lethal. For convenience, this set of strains carried a *wA1* mutation resulting in white conidiospores, as opposed to the wild-type green color.

Figure 11: The Uso1 GHD interacts directly with Bos1 and Bet1 SNAREs acting in the ER/Golgi interface

(A). Purified fusion proteins in which the cytosolic domains of the indicated SNAREs have been fused to GST were used in pulldown experiments with His-tagged, purified wild-type and E6K/G540S Uso1. The plasma membrane Qa syntaxin Sso1 was used as negative control. Pulled-down material was analyzed by anti-His western blotting. (B). Quantitation of the above experiment; significance was determined by unpaired *t*-student tests. Error bars represent S.E.M. (C). As in A, but using in vitro synthesized, HA3-tagged Sly1 as prey. Samples were analyzed by anti-HA western blotting. (D). As in A, but using wild-type and mutant GHD as preys, rather than full-length Uso1. (E). Quantitation of the experiment in D. **(F)**. GST pull-down experiment comparing the ability of the GHD to interact with the early Golgi Qb and Qc SNAREs (Bos1 and Bet1), with that of their medial Golgi counterparts (Qb Gos1 and Qc Sft1).

Figure 11—figure supplement 1. AlphaFold2 prediction of the ER/Golgi SNARE bundle.
(A) Sec5/Bos1/Bet1/Sec22 predicted SNARE bundle.
(B) Quality control (pLDDT, color coded, and PAE) of the model.

Figure 12. AlphaFold2 models provide insight into the additive mode of suppression shown by E6K and G540S.

(A). Model of full length Uso1 bound to the ER/Golgi SNAREs Bos1 and Bet1.

(B). Top, ribbon representation of the Bos1 N-terminal Habc domain and Uso1GHD. Bottom, Inset combining surface and ribbon depiction.

(C). Increased binding of Bos1 to G540S Uso1 appears to involve insertion of Ser540 into a pocket located in the Habc domain of the Qb SNARE. Partial view of the Bos1-Uso1 GHD surface of interaction in the wild type (left) and mutant (right) models. G540 and S540 are annotated.

(D). The N-terminal amphipathic α-helix of Uso1 comprising the E6K substitution lies within a flexible stretch of the protein that might facilitate its insertion into membranes. Alignment of six independent predictions, with Glu6 highlighted in red. The Uso1 GHD was modeled alone, in a complex with SNARE proteins or with Ypt1. The N-terminal α-helix (boxed) adopts different positions, suggesting high flexibility.

Figure 12—figure supplement 1. AlphaFold2 prediction of the Bet1-GHD interaction.
The putative binding surface of Bet1 and Uso1 as determined by AlphaFold2. Top images, cartoon of Bet1-GHD interactions, colored by pLDDT score. Alignment of four independent predictions involving the Bet1 N-terminal region and Uso1 GHD. A single model for Uso1 GHD is shown on the top representation for simplicity. In spite of the disordered nature of the N-terminal Bet1 region, the Bet1-Uso1 binding interface is consistent among models. Bottom, surface representation of the N-terminal Bet1 region (orange) in complex with the GHD. Also indicated is the Habc domain of Bos1 (yellow) bound to the GHD.

Figure 12—figure supplement 2. AlphaFold 2 prediction of the RAB1 binding site on the Bet1/Bos1/Uso1 GHD complex.

(A) and (B): cartoon representation of the GHD-RAB1 complex. The model is depicted as pipes and planks

(C): Orthogonal views of the Uso1 GHD-RAB1-Bet1-Bos1Habc structural model. The Uso1 GHD is shown as surface to emphasize the distant binding sites of the Bos1 Habc domain, RAB1 and the Bet1 N-terminal region

(D): Ribbon representation of the model shown in C but including the full-length SNARE subunits, i.e. the GHD domain of Uso1, RAB1 and the SNARES Bet1 and Bos1. Proteins are in the correct orientation to connect membranes separated by 23 nm, counting from the SNARE TMDs to the prenylated RAB1 residues.

Figure 12—figure supplement 3. Quality control assessment of AlphaFold2 predictions for the indicated complexes.