Cell surface auxin signalling directly targets PIN-mediated auxin fluxes for adaptive

plant development

Lesia Rodriguez[1], Lukáš Fiedler[1], Minxia Zou[1], Caterina Giannini[1], Aline Monzer[1], Zuzana Gelová[1],

Inge Verstraeten[1], Jakub Hajný[1,2], Shutang Tan[1], Lukas Hoermayer[1], Lanxin Li[1], Maria Mar Marques-

Bueno[3,5], Gergely Molnár[1], Tongda Xu[4], Ivan Kulich[1], Yvon Jaillais[3] and Jiří Friml[1*]

[1]Institute of Science and Technology Austria (ISTA), Am Campus 1, 3400, Klosterneuburg, Austria

[2]Laboratory of Growth Regulators, The Czech Academy of Sciences, Institute of Experimental Botany

& Palacký University, Šlechtitelů 27, CZ-78371 Olomouc, Czech Republic

[3]Laboratoire Reproduction et Développement des Plantes Université de Lyon, ENS de Lyon, UCB Lyon

1, CNRS, INRA, F-69342 Lyon, France

[4]Plant Synthetic Biology Center, Haixia Institute of Science and Technology, and College of Life

Sciences, Fujian Agriculture and Forestry University, Fuzhou, Fujian, P.R.China

[5]Present Address: Center for Research in Agricultural Genomics (CRAG) CSIC-IRTA-UAB-UB,

Barcelona 01893, Spain

* Corresponding author: jiri.friml@ista.ac.at

24
25

SUMMARY

The plant hormone auxin and its directional transport across tissues mediate much of the remarkably adaptive and plastic development of higher plants. Positive feedback between auxin signalling and transport is a key prerequisite for self-organizing development including flexible vascular tissue formation and directional growth responses, such as gravitropism. Here we identified a mechanistic link between cell surface ABP1-based auxin perception, the associated TMK1 kinase and PIN auxin transporters. *abp1* and *tmk1* mutants are defective in auxin-triggered phosphorylation of PIN proteins and in PIN-mediated directional auxin transport. Auxin, via ABP1, induces activation and stabilization of TMK1, thus promoting direct interaction with and phosphorylation of PIN2. Following gravistimulation, the auxin-activated TMK1 acts along the lower root side to reinforce asymmetry in PIN2-mediated auxin fluxes for gravitropic root bending. This ABP1-TMK1-dependent positive feedback on PIN-mediated directional auxin transport is fundamental for robust root gravitropism and presumably also for other self-organizing developmental processes.

INTRODUCTION

Being sedentary, plants utilize directional growth to spread and position themselves in various environments. Such plant 'movements' according to gravity, light, water availability and other factors are a crucial part of their survival strategy[1]. Other spectacular examples of the superior adaptability of plant development are self-organizing processes such as the formation of an embryonal axis, the regular arrangement of leaves and flowers, the formation of intricate leaf vein patterns, and flexible regeneration of vasculature around a wound. The phytohormone auxin serves as the major endogenous signal behind all this impressive adaptive development[2]. It translates environmental stimuli such as light and gravity into growth responses and aids self-organizing development on account of a feedback regulation between auxin signalling and polar auxin flow[3,4,5].

Directional cell-to-cell auxin transport is a unique, plant-specific mechanism[6] that relies on several plasma membrane-localized transporters[7,8]. Chief among these are the AUX1/LAX auxin importers[9] and the PIN auxin exporters, with the latter determining auxin flow direction by their polarized subcellular localization[10,11]. Inside the cell, auxin triggers a well-studied, predominantly nuclear signalling pathway dependent on TIR1/AFB auxin receptors leading to developmental reprogramming[12,13] and growth regulation[14]. The less well characterized, cell surface auxin signalling involves the AUXIN BINDING PROTEIN 1 (ABP1) receptor and the leucine-rich repeat receptor-like kinase (LRR RLK) TRANSMEMBRANE RECEPTOR-LIKE KINASEs (TMKs)[15,16]. Both these signalling mechanisms are required for auxin feedback on PIN-mediated auxin transport as manifested during the so-called 'auxin canalization', the formation of polarized auxin-transporting channels from a local auxin source[15,17]. Auxin canalization is a process relying on a positive feedback between auxin signalling and auxin transport[3]. This is consistent with theoretical models of auxin feedback-based PIN polarization encompassing both intracellular and cell surface auxin signalling[18]. It has been speculated that auxin impacts PIN activity and polarity by affecting endocytic trafficking[19] involving clathrin-mediated PIN endocytosis[20] and ARF-GEF GNOM-dependent PIN endosomal recycling[21,22] as well as post-translational modifications, chiefly reversible phosphorylation[23,24,25,26]. Nonetheless, the mechanistic link between auxin signalling and PIN-mediated transport has remained elusive.

Fast root gravitropism, mediated by directional auxin flow from the site of gravity perception at the root cap towards the elongation zone where the growth response takes place, is among the latest evolutionary innovations in the quest of plants to grow upright[27]. Besides being a fascinating example of adaptive plant development[28], it is also a perfect model for studying the coordination of auxin signalling and transport. After gravity sensing by sedimenting statoliths in root cap cells[29], the auxin flux is redirected towards the lower side of the root by PIN polarization to bottom cell sides[30,31]. This initial auxin asymmetry is propagated from the root tip to the elongation zone by AUX1- and PIN2-mediated transport[32,33,27], and translated into growth redirection through inhibition of cell elongation at the lower root side[34]. During this response, PIN2 distribution also becomes asymmetric with PIN2 being more

abundant at the lower side of the root as compared to the upper side[35,36]. Such asymmetric PIN2
distribution thus not only propagates, but also reinforces the initial auxin gradient at the root tip, crucially
contributing to the robustness of the root bending response as well as to the fine-tuning of gravitropism
by other hormonal cues[37]. Nonetheless, the underlying mechanism of this auxin flow reinforcement
remains unclear.

Here we uncovered a role of the ABP1-TMK1 cell surface auxin signalling in direct phosphorylation
and regulation of PIN auxin transporters. In the context of gravitropic root bending, TMK1 interacts in
an auxin-dependent manner with PIN2 and phosphorylates it. This provides a direct mechanism for
feedback regulation of auxin transport, applicable not only to gravitropism but also to other
developmental processes.

RESULTS

ABP1-TMK1 auxin perception mediates auxin-triggered PIN phosphorylation

ABP1-TMK-mediated cell surface auxin signalling was recently implicated in self-organizing aspects
of plant development such as vasculature regeneration and *de novo* establishment of auxin-transporting
channels[15]. These phenomena are encompassed by the broader term 'auxin canalization' and are
postulated to rely on a direct feedback between auxin signalling and auxin transport[4]. The molecular
insights into canalization remained largely enigmatic until the recent identification of CAMEL and
CANAR, two LRR RLKs, which target PIN auxin efflux carriers (PINs) to polarize cells for auxin
transport across tissues[38].

Motivated by this possible link between phospho-signalling and canalization, we decided to investigate
whether auxin perception by ABP1-TMKs might itself feed into PIN-mediated polar auxin transport.
We took advantage of a published rapid phospho-proteomic dataset recorded in roots of the *tmk1-1*
mutant or WT exactly after 2 minutes of auxin treatment[15,39,40] and examined it for evidence of TMK1-
regulated PIN phosphorylation. We identified 12 phospho-sites pertaining to four canonical PINs as
putative targets of TMK1 (Figure 1A). Notably, all twelve sites mapped to hydrophilic PIN loops which
is the expected location for post-translational modifications regulating PIN polarity and auxin efflux
activity[41].

We found two PIN1 phospho-sites (S271, S337), five PIN2 phospho-sites (S179, S183, T233, S393,
S439), and two PIN3 phospho-sites (S366, S389) significantly hypo-phosphorylated in the *tmk1-1*
mutant compared to WT (Figure 1A). The two PIN1 phospho-sites were previously characterized, PIN1
S271 as a target of the D6PK protein kinase[26] and PIN1 S337 as a target of the MKK7-MPK6 module[42].

There were also three hyper-phosphorylated PIN phospho-sites in the *tmk1-1* mutant, one belonging to

PIN3 (S294) and two pertaining to PIN7 (S297, S431) (Figure 1A). We next examined the role of ABP1,

the auxin receptor upstream of TMK1[15]. We found that eight out of the twelve TMK1-dependent PIN

phospho-sites were also significantly hypo-phosphorylated in the *abp1-TD1* mutant compared to WT

(Figure 1B).

These observations suggest that auxin very rapidly changes the phospho-status of PIN proteins by a

mechanism depending on ABP1-TMK1 signalling.

ABP1-TMK1 is required for polar auxin transport

We next addressed whether the ABP1-TMK1-mediated PIN phosphorylation has any functional

relevance for PIN-mediated polar auxin transport. To this end, we tested basipetal/rootward auxin

transport in hypocotyls and basipetal/shootward transport in roots of WT, *abp1*, and *tmk1* mutants.

Application of labelled ^{3}H-IAA on decapitated etiolated hypocotyls allows monitoring of the NPA-

sensitive[43], mainly PIN1-dependent downward auxin transport[38]. Indeed, *tmk1-1* mutant hypocotyls as

well as those of both *abp1* mutant alleles (*abp1-C1* and *abp1-TD1*) showed a decreased rate of ^{3}H-IAA

transport (Figure 1C and 1D). Similarly, application of ^{3}H-IAA on the root tip and monitoring of

predominantly PIN2-mediated upward auxin transport revealed that the *tmk1-1* and *abp1* mutant roots

show a decreased ^{3}H-IAA transport relative to WT (Figure 1E). Importantly, these defects are

complemented in the *TMK1::TMK1-FLAG/tmk1-1* (*comp-tmk1-1*) and *ABP1::GFP-ABP1/abp1-C1*

(*comp-C1*) lines (Figure 1F).

Collectively, this shows that deregulated PIN phosphorylation in *abp1* and *tmk1* mutants correlates with

impaired PIN-mediated polar auxin transport in shoots and roots.

ABP1 mediates auxin-induced activation of TMK1 in roots

The action of ABP1 as an auxin receptor, auxin-induced ABP1-TMK1 interaction, and similar

developmental and cellular phenotypes of the corresponding mutants[15,16], suggest that the ABP1 receptor

mediates activation of the TMK1 pathway by auxin. This has been also suggested by previous studies,

where auxin stimulates TMK1-dependent ROP2 and ROP6 signalling[16] and TMK1 nanoclustering[44] in

pavement cells and MEMBRANE ASSOCIATED KINASE REGULATOR2 (MAKR2) membrane

dissociation during root gravitropism[45].

TMKs form a redundant family of 4 LRR-RLKs, for which single mutants have only subtle phenotypes,

while higher-order mutants show strong defects in many developmental processes[46]. To focus on TMKs

in roots we used available global transcriptomic data (Figure S1A and S1B) and generated *TMK1::GUS*,

TMK2::GUS, *TMK3::GUS* and *TMK4::GUS* lines reporting the corresponding promoter activities. This revealed that TMK1 is the dominant family member highly expressed in roots, followed by TMK3 and TMK4 with lower expression levels (Figure S1C).

As an indirect proxy for TMK1 activation in roots we examined the membrane association of ROP6, which has been shown to act downstream of TMK1[16]. Immunoblotting microsomal root protein extracts from auxin-treated (IAA, 5 nM, 0–120 minutes) plants with an anti-ROP6 antibody revealed auxin-induced membrane association of ROP6. This effect was abolished in *tmk1-1* mutant roots (Figure 2A and S2E), confirming that ROP6 activation by auxin is TMK1-dependent.

To further confirm these observations and test ABP1 involvement in this process, we detected TMK1-mediated activation of ROP signalling using an orthogonal method. As usual for small GTPases, only GTP-bound (active) but not GDP-bound (inactive) ROP proteins engage in protein-protein interactions with their effectors. We purified the Cdc42/Rac interactive binding motif (CRIB) domain of the RIC1 ROP effector from bacteria and used it to pull down active ROPs from root protein extracts. Immunoblotting with an anti-ROP2 antibody revealed auxin-induced (IAA, 100 nM, 10 minutes) ROP2 activation in WT but not in *tmk1-1* roots and importantly neither in *abp1-C1* or *abp1-TD1* roots (Figure 2B). These results demonstrate that in roots auxin triggers ROP2 and ROP6 signalling downstream of ABP1-TMK1 signalling.

Next, we analysed TMK1 phosphorylation in response to auxin. The TMK1 kinase domain is known to be capable of auto-phosphorylation[47], and numerous studies have shown that changes in phosphorylation of the cytoplasmic domain of LRR-RLKs are indicative of LRR-RLK activation in plants[48,49]. Therefore, we examined TMK1 phosphorylation by performing a Phos-tag Biotin Probe binding analysis on TMK1-FLAG immunoprecipitated from auxin-treated (IAA, 10 nM, 1 hour) *TMK1::TMK1-FLAG*/*tmk1-1* roots. With this approach, we revealed that auxin induced phosphorylation and therefore activation of TMK1 in roots (Figure 2C and S2C). Unexpectedly, we observed that auxin treatment increased not only the [phospho-TMK1/total TMK] ratio (Figure S2C) but also the total TMK1 abundance (Figure 2C). Immunoblotting microsomal root protein extracts from WT (anti-TMK1 antibody), *TMK1::TMK1-FLAG* (anti-FLAG antibody), and *UBQ10::TMK1-3xHA* (anti-HA antibody) confirmed auxin-mediated (anti-TMK1: IAA, 5 nM, 0–120 minutes; anti-FLAG/HA: IAA, 5 or 10 or 100 nM, 1 hour) increase in TMK1 abundance (Figure 2A, S2A, S2B and S2E). In contrast, *TMK1* mRNA levels as determined by RT-qPCR did not increase but rather decreased with IAA treatment (IAA, 10 or 100 nM, 0–120 minutes: Figure S2D). This suggests that auxin upregulates TMK1 abundance on the protein level rather than transcriptionally. We also monitored TMK1-GFP signal after auxin treatment (IAA, 5 or 10 nM, 3.5-hour time course) of *TMK1::TMK1-GFP*/WT roots. In agreement with biochemical data, concentrations as low as 5 nM of IAA led to a strong increase in the TMK1-GFP signal (Figure 2D and 2E).

185 In totality, our experiments in roots suggest that auxin, via ABP1, promotes TMK1 phosphorylation and
186 stabilization at the plasma membrane, underlying TMK1 activation and leading to downstream
187 signalling. This provides a possible mechanism for how auxin could act through the ABP1-TMK1
188 pathway to trigger phosphorylation of PINs and other targets.

TMK1 interacts with and phosphorylates PIN2

Given that most TMK1-regulated PIN phospho-sites were identified in PIN2 (Figure 1A) and its
prominent role in root gravitropism, we further focused on PIN2 as a representative of PINs.

To address whether TMK1 regulates PIN2 directly, we performed a co-immunoprecipitation analysis
(Co-IP) between endogenous PIN2 and TMK1. We immunoprecipitated TMK1-FLAG from
TMK1::TMK1-FLAG/tmk1-1 roots and then used an anti-PIN2 antibody for detection of native PIN2.
PIN2 did not co-immunoprecipitate with TMK1-FLAG in untreated samples. However, in auxin-treated
roots (IAA, 5 or 20 nM, 15 or 30 minutes) we could co-immunoprecipitate PIN2 with TMK1-FLAG
and the Co-IP efficiency increased with the duration of IAA treatment (Figure 3A and S3A), suggesting
an auxin-dependent interaction of TMK1 with PIN2.

To verify the biochemical evidence for TMK1-PIN2 interaction, we employed the quantitative FRET-
FLIM technique. We co-expressed *35S::TMK1-GFP* and *35S::PIN2-mCherry* in *Arabidopsis* root
protoplasts and measured the fluorescence lifetime of the GFP signal. The lifetime of TMK1-GFP was
reduced after co-expression with PIN2-mCherry, which supported the interaction between TMK1 and
PIN2 (Figure 3B). The negative controls, an unrelated receptor-like kinase FLS2-GFP and TMK1$^{\Delta KD}$-
GFP with a truncated kinase domain, showed a less pronounced decrease in fluorescence lifetime,
confirming the specificity of the TMK1-PIN2 interaction (Figure 3B and S3B).

Given the direct interaction of TMK1 with PIN2, it seemed likely that at least some of the TMK1-
regulated PIN2 phospho-sites (Figure 1A) could be ascribed to direct phosphorylation by TMK1. To
test this, we performed an *in vitro* phosphorylation assay with ^{32}P-ATP as a phosphate donor by
incubating a purified N-terminal His-tagged hydrophilic loop (HL) of PIN2 with TMK1-3xHA
immunoprecipitated from a root protein extract. These *in vitro* assays showed that TMK1-3xHA
mediated the phosphorylation of His-PIN2 HL. Notably, only intact TMK1-HA but not the kinase-dead
version TMK1^{K616R}-HA was able to phosphorylate His-PIN2-HL (Figure 3C). We did not observe any
auxin effect in these kinase assays (Figure S3C), but this could be explained if TMK1-3HA activity was
already saturated by endogenous auxin during immunoprecipitation from TMK1-3xHA roots.

In conclusion, our data demonstrate that the receptor-like kinase TMK1 interacts, in an auxin-dependent
manner, with the PIN2 auxin efflux carrier and phosphorylates its hydrophilic loop.

TMK1 modulates PIN2 levels during the root response to exogenous auxin

The compromised PIN2-mediated shootward auxin transport in *abp1* and *tmk1* mutant roots (Figure 1) suggested an involvement of ABP1-TMK1 signalling in PIN2 functionality. Therefore, we tested whether TMK1-mediated PIN2 phosphorylation influences its polarity and abundance, both of which are known to be regulated by phosphorylation[41]. We did not observe any obvious changes in PIN2 polarity as assessed by anti-PIN2 immunolocalization experiments in *tmk1* mutants (data not shown), thus we examined the effects of auxin on PIN2 levels in WT and *tmk1-1* by immunoblotting microsomal root samples with an anti-PIN2 antibody. In WT, 5 nM IAA stabilized PIN2 over the course of a 1-hour treatment, but after 2 hours PIN2 decreased below starting levels (Figure 4A and S4A). With 10 nM IAA, however, PIN2 was destabilized already after 1 hour (Figure S4B). Such complex PIN2 dynamics in response to short versus long auxin treatments were described before for the synthetic auxin NAA[35,50]. Investigation of PIN2 transcription by RT-qPCR confirmed that regulation of PIN2 stability by auxin is independent of transcription (Figure S4C). Thus, auxin initially stabilizes and after a prolonged exposure destabilizes PIN2, consistent with previous reports[19,35].

Notably, the *tmk1-1* mutant exhibited premature PIN2 stabilization by IAA and showed resistance to long-term PIN2 downregulation (Figure 4A, S4A and S4B). We further assessed PIN2 abundance in auxin-treated WT and *tmk1-1* roots by live imaging. IAA treatment of PIN2-GFP roots caused a decrease in the GFP signal in a concentration-dependent manner. This auxin effect was not observed or even reverted in the *tmk1-1* mutant (Figure 4b and 4C), overall indicating that TMK1 mediates PIN2 destabilization, which is triggered by exogenous auxin.

TMK1 kinase activity is required for the root gravitropic response

To test the relevance of our observations on PIN2 regulation by auxin and by TMK1 in a physiological context, we focused on gravitropism, a classic developmental process during which membrane PIN2 levels undergo dynamic spatiotemporal changes[35,36], and which has been proposed to be regulated by TMK1 and its inhibitory protein MAKR2[45]. Analysis of root growth in verified *tmk* single mutants (Figure S5A) did not show obvious defects in growth rate (Figure S5C), but time-lapse imaging of root bending following gravistimulation revealed slower gravitropic response kinetics compared to WT plants. This defect was most pronounced in the *tmk1-1* mutant and complemented in the *TMK1::TMK1-FLAG/tmk1-1* line (Figure 5A and 5B; Figure S5B and S5C; Table S1 and S2). *tmk4-1* roots also bent less efficiently; *tmk3-1* had a very weak defect, whereas *tmk2-1* showed normal gravitropic bending (Figure S5B and S5C; Table S2), which is consistent with the very low expression of *TMK2* in the root (Figure S1A, S1B and S1C). Notably, the kinase activity of TMK1 is required for its role in root

gravitropism since the expression of TMK1 without a kinase domain ($35S::TMK1^{\Delta KD}$-GFP) or TMK1 carrying a mutation in the ATP-binding site ($UBQ10::TMK1^{K616R}$-$mCherry$) also conveyed defects in root gravitropism, even when introduced into WT plants (Figure 5A and 5B; Figure S5D and S5E; Table S2). Of note, the two kinase-dead lines showed fluorescent signal distribution comparable to that of *TMK1::TMK1-GFP* (Figure S1D and S5F).

Next, we tested whether TMK1 is involved in gravity-induced auxin re-distribution by introducing the *DR5rev::GFP* auxin response reporter into the *tmk1-1* loss-of-function mutant. Monitoring the DR5 signal following gravistimulation showed significantly less auxin response asymmetry in the *tmk1-1* mutant as compared to WT control (Figure 5C and 5D).

These results show that, during root gravitropism, TMK1 is involved in both the asymmetric distribution of auxin and in root bending.

TMK1 mediates PIN2 asymmetry during root gravitropic response

Having confirmed that TMK1 signalling is required for the root gravitropic response and given that TMK1 mediates PIN2 phosphorylation (Figure 1A and 3C) and stability (Figure 4), we examined the dynamic regulation of PIN2 abundance in response to gravity by immunoblotting microsomal root protein extracts. After 1 hour of gravistimulation, WT roots showed modest PIN2 stabilization; this was followed by a decrease below starting levels at the 2-hours timepoint without further changes for the rest of the time course (Figure 6A and S6C). The *tmk1-1* mutant PIN2 levels, however, increased dramatically after 1 hour of gravistimulation and then slowly recovered back to the starting value after 4 hours of gravistimulation. This biochemical evidence reveals dynamic regulation of total PIN2 abundance during gravitropism, which is mediated by TMK1.

To gain more spatial insights into the behaviour of PIN2 during the gravity response, we performed live imaging and followed PIN2-GFP asymmetry in WT and *tmk1-1* roots after gravistimulation. While WT roots established asymmetric PIN2 distribution as expected, this response was abrogated in the *tmk1-1* mutant (Figure 6C and 6D). Importantly, the *tmk1-1* mutant showed defects in both PIN2 stabilization at the lower side of the root (from 2 hours onwards) and in the destabilization of PIN2 at the upper side of the root (the 3 hours timepoint) (Figure 6D).

Thus, TMK1 orchestrates a complex, dynamic regulation of PIN2 abundance during gravitropism which provides a likely explanation for the gravitropic phenotype of the *tmk1* mutant.

TMK1 activity and asymmetric abundance during root gravitropism

The gravitropic phenotype of *tmk1-1* roots suggested to us that TMK1 might become activated during

the root gravitropic response. Immunoblotting WT root microsomal fractions from gravistimulated

seedlings with an anti-ROP6 antibody confirmed that gravistimulation activated ROP6 signalling. The

absence of these dynamics in the *tmk1-1* mutant showed that gravity-induced ROP6 activation was

TMK1-dependent (Figure 6A and S6C). We subsequently conducted a Phos-tag Biotin Probe binding

analysis on TMK1-FLAG immunoprecipitated from gravistimulated *TMK1::TMK1-FLAG/tmk1-1*

roots. This revealed a significant increase in TMK1 phosphorylation already after 1 hour of

gravistimulation (Figure 6B and S6D) demonstrating that gravistimulation activates TMK1. Similar to

our experiments with a time course of exogenous auxin (Figure 2A and 2C), we noted an increase in

TMK1 abundance over the course of gravistimulation (Figure 6B). Increasing TMK1 levels were also

apparent from Western blots performed on microsomal root extracts from gravistimulated seedlings

(Figure 6A; Figure S6A, S6B and S6C). Collectively, these data show that gravistimulation activates

TMK1 signalling in roots by impacting both the TMK1 phosphorylation status and stability.

To gain additional spatial information on this regulation, we performed live imaging on the

TMK1::TMK1-GFP line. Strikingly, we observed that after 1 hour of gravistimulation TMK1-GFP

developed an asymmetric pattern with higher signal intensity along the lower root side, which was

getting more pronounced over time (Figure 6E and 6F). This result is consistent with auxin accumulating

at the lower root side stabilizing TMK1 as seen for exogenous auxin application (Figure 2D and 2E).

This implies the existence of additional signals/regulations at the lower side of the root during the

graviresponse or some compensatory mechanism in the *tmk1-1* mutant, which causes bulk stabilization

of PIN2 and overshadows the fact that, in WT roots, the function of TMK1 is to stabilize PIN2 at the

lower root side shortly after gravistimulation. It is tempting to propose a model in which TMK1

stabilization at the lower root side reinforces PIN2 asymmetry to confer robustness on root gravitropism

in land plants, however, it remains a challenge for future studies to fully consolidate all the above-

mentioned intriguing observations.

Nonetheless, our data reveal crucial importance of TMK1 for dynamic PIN2-mediated fluxes and auxin

asymmetry in driving root gravitropism. Notably, auxin-triggered TMK1-PIN2 interaction and

phosphorylation provides a plausible mechanism for how this regulation occurs.

DISCUSSION

ABP1 relays auxin signal for TMK1 activation and stabilization

Cell surface auxin signalling, unlike the TIR1/AFB transcriptional pathway based predominantly in the

nucleus[51], has been a matter of considerable controversy[52]. Only recently was it confirmed in

Arabidopsis that ABP1 acts as an apoplastic auxin-binding receptor[15], which associates to the

extracellular domain of TMK1[16]. This provides a plausible mechanism for how extracellular auxin can

induce via ABP1 the phosphorylation of proteins in the cell interior, either directly through the cytosolic

kinase domain of TMK1 or through downstream kinase cascades. Indeed, *abp1* and *tmk1* mutants show

aberrant and highly inter-correlated phospho-proteomes[15,39]. Overall, the auxin-induced ABP1-TMK1

association[16], failure of *abp1* mutants to maintain steady-state TMK phosphorylation[15], and the lack of

ROP signalling downstream of TMK1 in response to auxin (Figure 2B and [16]) strongly support the notion

that ABP1 acts as a receptor communicating auxin for TMK-mediated phosphorylation.

It comes as a surprise that auxin not only induces TMK1 phosphorylation but also its persistent

stabilization. This is to be contrasted with the usual textbook signalling paradigm where prolonged

stimulation desensitizes the pathway in question. For instance, mammalian receptor tyrosine kinases

such as the EPIDERMAL GROWTH FACTOR RECEPTOR (EGFR) are endocytosed and almost

completely degraded within 1-2 hours of ligand application[53,54]. This peculiar TMK1 behaviour may

constitute a part of the positive feedback linking auxin to its transport that has been proposed as a central

mechanism behind canalization-mediated self-organized plant development[3]. It is conceivable that

canalization would rely on a positive feedback loop which persists until the process is completed rather

than on rapid desensitization. Furthermore, extracellular auxin perception has been predicted as a key

signalling input parameter for auxin canalization[18]. Consistent with these notions, auxin canalization

and canalization-mediated vascular tissue regeneration are severely defective in *abp1* and *tmk* mutants[15].

Despite these strong, specific defects, overall growth and development is not strongly affected following

single-gene lesions within the ABP1-TMK pathway, which can be explained by the genetic redundancy

between TMKs[46] and between ABP1 and its homologues[55]. Thus, ABP1-TMK cell surface signalling is

a likely input component of the positive feedback central to the auxin canalization mechanism.

Auxin induced TMK1-PIN2 interaction and phosphorylation for PIN2 stability

The biological roles of the global, ultrafast phosphorylation downstream of ABP1-TMK1 remain

unclear but prominent among the phosphorylated targets are multiple PIN auxin transporters (Figure

1A, 1B and [39]). The activity, stability and polar localization of PINs are known to be regulated heavily

by complex phosphorylation on multiple sites by different types of kinases in response to various

signals[41]. Accordingly, PIN-mediated auxin transport in shoots and roots of *abp1* and *tmk1* mutants is

compromised (Figure 1C-F) implying that PIN regulation by ABP1-TMK1 is functionally relevant.

It appears clear that ABP1 transmits the auxin signal to TMKs for their phosphorylation and activation,

thus enabling the phosphorylation of downstream targets such as PINs. What part of the phosphorylation

response emanating from ABP1-TMK1 is attributable to direct TMK1 action on substrate proteins,

however, remains elusive. At least in the case of PINs, as exemplified by PIN1[56] and PIN2 (Figure 3),

TMK1 engages in a direct interaction and in an auxin-dependent manner phosphorylates multiple

residues within their hydrophilic loops. While for PIN1 the phosphorylation seems relevant to its polar

localization in tissues[56], PIN2 undergoes dynamic stabilization and destabilization in response to the

TMK1-transduced auxin phospho-signalling. The former scenario is likely relevant to classical auxin

canalization, the latter for fine-tuning of auxin fluxes during root gravitropism. Thus, distinct PINs

phosphorylated by TMK1 at different residues can be regulated by auxin in ways depending on the

developmental context. Regardless of specifics of the regulation, the auxin input via ABP1-TMK1 and

its direct modulation of PIN activity represent a core mechanism for the positive feedback between auxin

signalling and directional transport.

TMK1-PIN2 module during root gravitropism

PIN2 evolved as a specific component of efficient root gravitropism of seed plants, which contributed

to their successful colonization of land habitats[27]. The key evolutionary innovation has been the

directional, PIN2-mediated auxin flow that carries the signal from the site of gravity perception at the

root tip to the site of growth response in the elongation zone[28]. Notably, PIN2 abundance also becomes

asymmetric with more PIN2 at the lower root side[35,36], which reinforces the initial auxin gradient and

contributes to the robustness of the root bending response[37]. This provides an excellent model to study

the interplay of auxin signalling and transport. TMK1 is required for the full extent of asymmetric auxin

distribution and for root bending during the gravitropic response. This is elegantly explained by the fact

that *tmk1* roots cannot establish asymmetric PIN2 distribution during the graviresponse (Figure 6C, 6D

and [57]). TMK1 protein abundance increases dynamically during root gravitropism along the lower side

of the root, mirroring the asymmetric auxin distribution and PIN2 accumulation.

The above observations lead to a model in which the initial auxin flow asymmetry following

gravistimulation activates and stabilizes TMK1 at the lower root side and promotes its interaction with

PIN2, leading to PIN2 phosphorylation. This positive feedback loop between auxin and PIN2-mediated

transport reinforces the auxin flow at the lower root side and contributes to a robust gravitropic response.

However, TMK1-mediated persistent PIN2 destabilization in response to exogenous auxin treatment

(Figure 4) is not consistent with PIN2 stabilization at the place of elevated auxin along the lower side of

the gravistimulated root, indicating a possible hidden complexity and variables. Regardless of these

additional regulations, the asymmetric action of TMK1 following gravistimulation and the TMK1-PIN2

feedback loop clearly are required for PIN2 asymmetry during gravitropism, and consequently for a

robust bending response.

Conclusions

Here we identified a mechanism for the positive feedback between auxin and its transport, which is a

key prerequisite for canalization-guided plant development. It involves auxin-triggered phosphorylation

of multiple PIN auxin transporters. Exemplified by PIN2, this regulation involves a direct interaction

with the TMK1 receptor-like kinase. The input for TMK1 activation comes from auxin binding to the

ABP1 receptor at the cell surface, resulting in TMK1 phosphorylation, long-term TMK1 stabilization at the plasma membrane, and activation of downstream processes. This positive feedback regulation is required in the case of PIN2 for robust asymmetric auxin fluxes during root gravitropism and in the case of PIN1[56] most likely for auxin canalization-mediated flexible vascular tissue formation and regeneration. Thus, TMK-based auxin feedback regulation may represent a more general mechanism acting in different developmental contexts with different PIN transporters, thereby mediating specialized aspects of adaptive and self-organized development in plants. It is remarkable that the herein identified auxin feedback on its transport via TMK-based cell surface signalling likely evolved from an ancestral and ancient phosphorylation response to the small molecule auxin[39,40], later diversifying to guide the complex and flexible auxin-regulated development of morphologically complex land plants.

ACKNOWLEDGEMENTS

We thank W. Gray for providing material, N. Gnyliukh and E. Cervenova for help with manuscript preparation. To Dolf Weijers and Mark Roosjen, we thank for encouraging discussions and support with phospho-proteomic analyses. We thank Bioimaging and Life Science facilities at IST Austria for an excellent service and assistance. The research leading to these results has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme grant agreement No 742985 and Austrian Science Fund (FWF): I3630-775 B25 to J.F; National Natural Science Foundation of China (Grant 32130010, 31422008), start-up funds from FAFU to T.X.. Y.J. was funded by ERC no. 3363360-APPL under FP/2007-2013. L.R. was supported by FP7-PEOPLE-2011-COFUND ISTFELLOW program (IC1023FELL01) and the European Molecular Biology Organization (EMBO) long-term postdoctoral fellowship (ALTF 985-2016).

AUTHOR CONTRIBUTIONS

Conceptualization: J.F., L.R.; Methodology: L.R., I.K.; Investigation: L.R., L.F., M.Z., C.G., A.M., Z.G., I.V., J.H., M.R., S.T., G.M., L.H., L.L.; Resources: L.R., M.M.M.-B, Y.J., T.X.; Writing: J.F., L.F., L.R.; Review & Editing: all authors; Visualization: L.R., L.F.; Supervision: L.R., J.F.; Funding Acquisition: J.F., D.W.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE TITLES AND LEGENDS

Figure 1.

Figure 1. ABP1-TMK1 auxin perception targets PIN phosphorylation and polar auxin transport.

(A) Overview of PIN phospho-sites significantly regulated (FDR < 0.05) in *tmk1-1* mutant roots
compared to WT after treatment with auxin (100 nM, 2 min). 4 independent biological replicates, mean
$\pm$ SD, permutation-based t-tests with FDR-controlled p-values. Highlighted in bold are sites which are
co-regulated in the *abp1-TD1* mutant. Data adapted from [15]. *p < 0.05, **p < 0.01, ***p < 0.001, ****p
< 0.0001.

(B) Overview of PIN phospho-sites significantly regulated (FDR < 0.05) in *abp1-TD1* mutant roots
compared to WT after treatment with auxin (100 nM, 2 min). 4 independent biological replicates, mean
$\pm$ SD, permutation-based t-tests with FDR-controlled p-values. Highlighted in bold are sites which are
co-regulated in the *tmk1-1* mutant. Data adapted from [15]. *p < 0.05, **p < 0.01, ***p < 0.001, ****p <
0.0001.

(C) Basipetal/rootward transport of radioactive ^{3}H-IAA in decapitated hypocotyls is defective in the
tmk1-1 mutant (genotype/treatment fixed effect: χ^2 = 60.64, df = 2, p = 6.78e-14). 10 μM NPA, an
inhibitor of auxin transport, was used as a negative control. Pooled data from two independent
experiments with similar results (repetition random effect: ΔAIC = 2.0, df = 1, p = 0.9999), mean $\pm$ SD.
Asterisks show Holm-corrected unpaired two-sided t-tests. *p < 0.05, **p < 0.01, ***p < 0.001, ****p
< 0.0001.

D) Basipetal/rootward transport of radioactive ^{3}H-IAA in decapitated hypocotyls is defective in *abp1*
mutants (one-way ANOVA: $F_{3,8}$ = 60.13, p = 7.86e-06). 10 μM NPA, an inhibitor of auxin transport,
was used as a negative control. Single experiment, n = 3 for each condition, mean $\pm$ SD. Asterisks show
Holm-corrected unpaired two-sided t-tests. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

(E) Basipetal/shootward transport of radioactive ^{3}H-IAA in roots is defective in *tmk1-1* and *abp1*
mutants (genotype/treatment fixed effect: χ^2 = 32.30, df = 4, p = 1.66e-06). Pooled data from two
independent experiments with similar results (repetition random effect: ΔAIC = 1.1, df = 1, p = 0.0804),
mean $\pm$ SD. Asterisks show Holm-corrected unpaired two-sided t-tests. *p < 0.05, **p < 0.01, ***p <
0.001, ****p < 0.0001.

(F) Complementation of defects in root basipetal/shootward transport of radioactive ^{3}H-IAA for the *TMK1::TMK1-FLAG/tmk1-1* (*comp-tmk1-1*) and *ABP1::GFP-ABP1/abp1-C1* (*comp-abp1-C1*) lines. Single experiment, n = 10 for each condition, mean ± SD.

Figure 2.

Figure 2. Auxin-induced activation of cell surface ABP1-TMK1 signalling in roots.

(A) Western Blot: time course of endogenous TMK1 and ROP6 protein levels in the microsomal fraction of 4-day-old WT and *tmk1-1* roots treated with 5 nM IAA (see Figure S2E for quantification). The experiment was repeated 3 times and representative data are shown. The loading control is reused in Figure 4A which depicts more data from the same experiment.

(B) Auxin-mediated activation of ROP signalling is defective in *tmk1-1* and *abp1* mutant backgrounds. Pull-down of active ROP GTPases from root protein extracts using a heterologously expressed GST-tagged CRIB domain from the RIC1 ROP effector. Both input and pull-down gels were blotted with an anti-ROP2 antibody.

(C) Comparison of TMK1-FLAG phosphorylation in untreated roots and roots treated with 10 nM IAA for 1h. Root *TMK1::TMK1-FLAG/tmk1-1* extracts treated with 10 nM IAA or DMSO control for 1 hour were used for immunoprecipitation with anti-Flag antibody. Phos-tag Biotin Probe was used to analyse TMK1 phosphorylation levels and anti-Flag antibody for total amount of immunoprecipitated TMK1-FLAG (see figure S2C for quantification).

(D) Representative intensity-coloured images of *TMK1::TMK1-GFP* expression in 4-day-old seedlings after a 3-hour treatment with 5 or 10 nM IAA. Scale bars represent 20 μm.

(E) Time-course quantification of *TMK1::TMK1-GFP* signal in root tips treated with IAA (5 or 10 nM) or DMSO control, normalized to the initial fluorescence value. Mean ± SD (n = 20; experiment repeated 3 times with representative data depicted).

Figure 3.

Figure 3. Auxin-mediated TMK1 interaction with and phosphorylation of PIN2.

(A) Auxin promotes the interaction between TMK1 and PIN2 in a co-immunoprecipitation (Co-IP) assay. Microsomal protein fraction from 20 nM IAA- or DMSO-treated roots expressing *TMK1::TMK1-FLAG/tmk1-1* was immunoprecipitated with an anti-FLAG antibody, and endogenous PIN2 was detected by blotting with an anti-PIN2 antibody. See Figure S3A for Co-IP with 5 nM IAA where input PIN2 abundance in the protein microsomal fraction is increased.

(B) FRET-FLIM analysis on transiently expressed *35S::TMK1-GFP, 35S::TMK1^{ΔKD}-GFP, 35S::FLS2-GFP* and *35S::PIN2-mCherry* in root protoplasts. Fluorescence lifetime values are displayed as a heat map (for additional images see Figure S3B). Two-way ANOVA with post hoc tests ($*p < 0.05$; $****p < 0.0001$).

(C) *In vitro* kinase assay showing that TMK1-3xHA directly phosphorylates the 6His-PIN2 hydrophilic loop (HL). Top panel, ^{32}P autoradiography; middle panel, CBB staining; bottom panel, immunoblot with an anti-HA antibody.

Figure 4.

Figure 4. TMK1 dynamically regulates root PIN2 levels in response to exogenous auxin.

(A) Western Blot: time course of endogenous PIN2 protein levels in the microsomal fraction of 4-day-old WT and *tmk1-1* roots treated with 5 nM IAA (see Figure S4A for quantification). The experiment was repeated 3 times and representative data are shown. The loading control (reused) and the anti-PIN2 immunoblot come from the same experiment as that shown in Figure 2A.

(B) Representative intensity coloured images of *PIN2::PIN2-GFP* after a 3-hour treatment with 10 nM IAA. Scale bars, 20 μm.

(C) Quantification of *PIN2::PIN2-GFP* in root tips of WT or *tmk1-1* seedlings treated with IAA (5, 10 nM or DMSO control), normalized to the initial fluorescence value. Mean ± SD (n = 20; experiment repeated 3 times with representative data depicted).

Figure 5.

Figure 5. TMK1 kinase activity is required for the gravitropic response.

(A) Representative images of root bending 3 hours after a 90° rotation for 4-day-old WT, mutant *tmk1-1*, kinase-dead *UBQ10::TMK1^{K616R}-mCherry,* and genetically complemented *TMK1::TMK1-FLAG/tmk1-1* seedlings.

(B) Time course of the gravitropic response for lines shown in (A) measured every 30 minutes, mean ± SD (n = 10; experiment repeated 3 times, a representative run is depicted). See Figure S5E for complete analysis and root growth rate data, and Table S2 for statistical analysis.

(C) Representative confocal images of asymmetric auxin response (visualized by *DR5rev::GFP*) at the lower side of the root before and after a 3-hour gravistimulation in WT and *tmk1-1* 4-day-old roots.

(D) Quantification of *DR5rev::GFP* asymmetry as a ratio of fluorescence intensity at the lower side to the upper side of gravistimulated roots at the indicated time points, normalized to the initial fluorescence value. Mean ± SEM of at least 3 independent experiments, n = 10. Two-way ANOVA with Dunnett's multiple comparison tests (*$p < 0.05$, **$p < 0.01$). 'g' represents the gravity vector.

Figure 6.

Figure 6. TMK1 mediates PIN2 asymmetry during the root gravitropic response.

(A) Western Blot: time course of endogenous TMK1, ROP6 and PIN2 protein levels in the microsomal fraction of 4-day-old WT and *tmk1-1* roots after 0, 1, 2, 3 and 4 hours of gravistimulation. The experiment was repeated 3 times and representative data are shown. For quantification of protein levels see Figure S6C.

(B) TMK1-FLAG phosphorylation after gravistimulation. *TMK1::TMK1-FLAG/tmk1-1* was immunoprecipitated with an anti-FLAG antibody and detected using a Phos-tag Biotin Probe (see Figure S6D for quantification).

(C) Representative images of *PIN2::PIN2-GFP* in 4-day-old WT and *tmk1-1* seedlings after 4 hours of gravistimulation within a time course of 0, 2, 3, 4 and 5 hours of gravistimulation. 'g' represents the gravity vector. Scale bars, 20 μm.

(D) Quantification of PIN2-GFP ratio between the lower and upper sides of gravistimulated roots, normalized to the initial fluorescence value at the indicated time point. Mean ± SD (n = 8; experiment repeated 3 times with representative data depicted). Significant differences to timepoint 0 are shown by *p*-values, two-way ANOVA with Dunnett's post hoc tests (**** *$p < 0.0001$*).

(E) Intensity-coloured map of 4-day-old *TMK1::TMK1-GFP* root tips before and after 3 hours of a 90° gravitropic stimulus. White arrows indicate asymmetric TMK1-GFP distribution. Scale bars, 20 μm.

(F) Quantification of TMK1-GFP in the epidermis and cortex of the upper and lower root sides normalized to the initial fluorescence value, mean ± SD (n = 10). Experiment repeated 3 times with representative data depicted. Significant differences to timepoint 0 are shown by *p*-values, two-way ANOVA with Dunnett's post hoc tests (**** *$p < 0.0001$*).

Figure S1.

Figure S1. Expression of the TMK leucine-rich repeat receptor-like (LRR-RLK) kinase family in roots.

(A) Relative transcription levels of *TMK1*, *TMK2*, *TMK3* and *TMK4* in *Arabidopsis* roots from 803 samples assayed with the Affymetrix Arabidopsis ATH1 Genome Array. Data was obtained from the Genevestigator database (https://genevestigator.com). Mean ± SD.

(B) Relative expression levels of *TMK1, TMK2, TMK3* and *TMK4* in different root cell types obtained from high-resolution spatiotemporal microarray analysis of 5–6-day old roots reported in [58] (ePlant http://bar.utoronto.ca/eplant).

(C) Root tip expression pattern of *TMK1*, *TMK2*, *TMK3*, and *TMK4* promoters fused to the GUS transcriptional reporter. Scale bars, 100 μm.

(D) Representative intensity-colored image of *TMK1::TMK1-GFP* expression in 4-day-old root tips.

Figure S2.

Figure S2. Auxin promotes TMK1 stability, phosphorylation state, and downstream signalling.

(A) Western blotting of total root protein extracts of 4-day-old *TMK1::TMK1-FLAG/tmk1-1* and *UBQ10::TMK1-3xHA* transgenic plants after 1-hour treatment with different concentrations of IAA (5, 10 and 100 nM IAA) or DMSO control. Ponceau staining was used as a loading control.

(B) Corresponding quantification of TMK1 protein levels normalized to Ponceau, and subsequently to DMSO condition, mean ±SD from 3 independent experiments. One-way ANOVA with Dunnett's multiple comparisons relative to DMSO (****$p<0.0001$).

(C) Quantification of TMK1-FLAG phosphorylated upon auxin treatment (10 nM, 1 hour) (see Figure 2C). Phosphorylation levels detected with a Phos-tag Biotin Probe were normalized to signal obtained from anti-FLAG detection, and subsequently to the initial value from time point 0. Mean ± SD of 3 independent biological replicates. *p*-value for comparison to DMSO was calculated with a two-tailed unpaired t-test (**$p < 0.01$).

(D) RT-qPCR time course of relative *TMK1* expression after 10 or 100 nM IAA treatment. Normalization was done to *PP2A* as a reference gene. The expression level at the start of the experiment (T0) was set to a relative value of 1 and statistical comparisons were made to this expression level (*$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$).

(E) Time course of TMK1 and ROP6 protein levels in the membrane fraction of 4-day-old WT or *tmk1-1* roots upon auxin treatment (5 nM). Plasma membrane AHA2 H$^+$ ATPase (immunoblot shown in Figures 2A and 4A) was used as a loading control. We then normalized to initial protein levels at time point 0. Values are mean ± SD from 3 independent experiments. Two-way ANOVA with Dunnett's multiple comparisons (****$p < 0.0001$).

Figure S3.

Figure S3. TMK1 interacts with PIN2 in an auxin-dependent manner and phosphorylates it.

(A) Co-IP assay showing the interaction of TMK1 with PIN2 upon low auxin treatment (5 nM IAA).

Microsomal protein fraction from WT was used as a control for unspecific binding of endogenous

PIN2 (shown in Figure 3A, same experiment).

(B) FRET-FLIM analysis on transiently expressed *35S::TMK1^{ΔKD}-GFP, 35S::TMK1^{ΔKD}-*

GFP/35S::PIN2-mCherry, 35S::FLS2-GFP and *35S::FLS2-GFP/35S::PIN2-mCherry* in root

protoplasts (the same experiment is shown in Figure 3B). GFP fluorescence lifetime was calculated as

described in the Methods section. The heat map depicts fluorescence lifetime values (see Figure 3B for

lifetime analysis).

(C) *In vitro* kinase assay showing phosphorylation of the 6His-PIN2 HL substrate by TMK1-3xHA in

the presence or absence of auxin. Top panel, ^{32}P-autoradiography; middle panel, CBB staining; bottom

panel, immunoblot with an anti-HA antibody.

Figure S4.

Figure S4. TMK1 regulates PIN2 stability on the post-transcriptional level.

(A) Time course of PIN2 protein levels in the membrane fraction of 4-day-old WT or *tmk1-1* roots upon

auxin treatment (5 nM). Plasma membrane AHA2 H$^+$-ATPase (immunoblot shown in Figures 2A and

4A) was used as a loading control. We then normalized to initial protein levels at time point 0. Values

are mean ± SD from 3 independent experiments. Two-way ANOVA with Dunnett's multiple

comparisons (****$p < 0.0001$).

(B) Left panel: immunoblot of endogenous TMK1 and PIN2 in the microsomal fraction in response to

1-hour treatment with 10 nM IAA in WT versus *tmk1-1* roots. Right panel: quantification of protein

levels detected with anti-TMK1 and anti-PIN2 antibodies normalized to the loading control, and then to

DMSO. Values are mean ± SD from 3 independent experiments. Two-way ANOVA with Dunnett's

multiple comparisons (****$p<0.0001$).

(C) Comparison of PIN2 expression in WT versus *tmk1-1* upon auxin treatment. Relative expression of

PIN2 upon 10 and 100 nM IAA treatment at different time points (0, 5, 60, 120 and 180 minutes).

Normalization was done to *PP2A* as a reference gene. The expression level at the start of the experiment

was set to the relative value of 1.

Figure S5.

Figure S5. *TMK* genes are required for robust gravitropic root bending.

(A) Expression levels of *TMK1, TMK2, TMK3* and *TMK4* determined by RT-qPCR in root samples of 4-day-old WT and *tmk1-1, tmk2-1, tmk3-1* and *tmk4-1* seedlings. RNA was extracted in 4 independent biological repeats and the RT-qPCR analysis was performed in 3 technical repetitions. Expression levels were normalized against the expression level of PP2A as a housekeeping gene. Asterisks indicate significantly lower expression levels in the respective mutants, two-tailed unpaired t-tests (****$p < 0.0001$ and **$p < 0.01$).

(B) Root gravitropism in single mutants of TMK family genes. Root curvature angles were measured every 30 min after reorientation by a 90° rotation. Representative images of 4-day-old WT, *tmk1-1, tmk2-1, tmk3-1,* and *tmk4-1* single mutant roots after 3 hours of reorientation.

(C) Time course of root tip angle changes and root growth rates, mean ± SEM from 3 independent experiments (n= 30-40). See Table S1 for statistical analysis.

(D) Representative images of root bending 3 hours after a 90° rotation of 4-day-old WT and the overexpression line of TMK1 without a cytoplasmic kinase domain (*35S::TMK1^{ΔKD}-GFP*).

(E) Time course of the gravitropic response including *35S::TMK1^{ΔKD}-GFP* lines and root growth rate from the same experiment at the indicated time points, mean ± SD (n = 10; experiment repeated 3 times with representative data depicted).

(F) Representative images of primary roots expressing the *UBQ10::TMK1^{K616R}-mCherry* and *35S::TMK1^{ΔKD}-GFP* constructs.

Figure S6.

Figure S6. Gravistimulation promotes TMK1 stability, phosphorylation state, and downstream signalling

(A) Western blotting of total root protein extracts of 4-day-old *TMK1::TMK1-GFP, TMK1::TMK1-FLAG/tmk1-1* and *UBQ10::TMK1-3xHA* transgenic plants after 0, 1, 2, 3h of gravistimulation. Ponceau staining was used as a loading control.

(B) Corresponding quantification of the relative TMK1 protein levels normalized first to Ponceau, and then to initial protein levels at time point 0, values are mean ± SD from 3 independent experiments. One-way ANOVA with Dunnett's multiple comparisons to timepoint 0 (****$p < 0.0001$).

(C) Comparison of TMK1, PIN2, and ROP6 endogenous levels in the membrane fraction during gravistimulation of WT and *tmk1-1* 4-day-old roots. Normalization was to plasma membrane AHA2 ATPase (immunoblot analysis shown in Figure 6A), and then to initial protein levels at timepoint 0. Values are mean ± SD from 3 independent experiments. Two-way ANOVA with Dunnett's multiple comparisons to timepoint 0 (****$p < 0.0001$).

(D) Quantification of phosphorylated TMK1 levels during gravistimulation (see Figure 6B). TMK1 protein levels detected with biotinylated Phos-tag Probe were normalized to the signal obtained from anti-FLAG detection, and then to the initial value from timepoint 0. Values are mean ± SD from 3 experiments with 3 independent biological replicates. Two-way ANOVA with Dunnett's multiple comparisons to timepoint 0 (****$p < 0.0001$).

653   **SUPPLEMENTARY TABLES**

655   **Table S1. Statistical analysis of the gravitropic kinetics presented in Figure S5B and S5C.** Asterisks
656   are from Dunnett's multiple comparison tests versus WT. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p <$
657   0.0001.

Time (h)	tmk1-1	tmk2-1	tmk3-1	tmk4-1
0.5	ns	ns	ns	ns
1	**	ns	ns	ns
1.5	*	ns	ns	ns
2	*	ns	ns	ns
2.5	*	ns	ns	ns
3	*	ns	ns	ns
3.5	**	ns	ns	ns
4	**	ns	ns	**
4.5	***	ns	ns	*
5	****	ns	ns	*
5.5	****	ns	ns	**
6	****	ns	ns	**
6.5	****	ns	ns	*
7	***	ns	*	*

669 **Table S2. Statistical analysis of the gravitropic kinetics presented in Figure 5A, 5B, S5D and**
670 **S5E.** Asterisks are from Dunnett's multiple comparison tests versus WT. *p < 0.05, **p < 0.01, ***p
671 < 0.001, ****p<0.0001.

Time (h)	*tmk1-1*	*TMK1::TMK1-FLAG/ tmk1-1*	*35S::TMK1ΔKD -GFP WT (L9)*	*35S::TMK1ΔKD-GFP WT (L8)*	*UBQ10::TMK1K616R- mCherry WT (L11)*	*UBQ10::TMK1K616R- mCherry WT (L14)*
0.5	****	ns	*	**	*	ns
1	****	ns	**	**	****	**
1.5	***	ns	*	**	****	**
2	***	ns	*	***	****	**
2.5	****	ns	*	***	****	**
3	****	ns	*	***	****	*
3.5	****	ns	*	*	***	*
4	****	ns	ns	*	***	ns
4.5	****	ns	ns	*	**	ns
5	****	ns	ns	ns	*	*
5.5	****	ns	ns	ns	*	*
6	****	ns	ns	ns	*	**
6.5	****	ns	ns	ns	*	ns
7	***	ns	ns	ns	*	ns

681 **Table S3. List of primer sequences used in this study.**

Primer name	Use	Sequence 5'→ 3'
PP2A FW	RT-qPCR	TAACGTGGCCAAAATGATGC
PP2A RE	RT-qPCR	GTTCTCCACAACCGCTTGGT
EF1a FW	RT-qPCR	TGAGCACGCTCTTCTTGCTTTCA
EF1a RE	RT-qPCR	GGTGGTGGCATCCATCTTGTTACA
TMK1 FW	RT-qPCR	TGGCACCCGAATACGCAGTAAC
TMK1 RE	RT-qPCR	GCTTTCTTCTGGTTGCGATTCGTC
TMK2 FW	RT-qPCR	TTTGGTTTAGTCCGGCTTGC
TMK2 RE	RT-qPCR	CGACCCGTTAGTAGCTCCAT
TMK3 FW	RT-qPCR	ACTCGAGTAGCTGGGACTTTCG
TMK3 RE	RT-qPCR	CACTCTTCCCGTCACTGCGTATTC
TMK4 FW	RT-qPCR	CAGAATACGCCGCTACTGGAAG
TMK4 RE	RT-qPCR	TCCATGAGAACCACACCAAATGC
PIN1 FWa	RT-qPCR	TTACTCCACCGCTACGAAC
PIN1 REa	RT-qPCR	ACTAAACTCTTCCCTCTCCAC
PIN2 FWa	RT-qPCR	TCACGACAACCTCGCTACTAAAGC
PIN2 REa	RT-qPCR	TGCCCATGTAAGGTGACTTTCCC
TMK1-FL-B1-F	cloning	GGGGACAAGTTTGTACAAAAAAGCAGGCTCAATGAAGAAAAGAAGAACCTTTC
TMK1-FL-B2-R	cloning	GGGGACCACTTTGTACAAGAAAGCTGGGTCTCGTCCATCTACTGAAGTGAAT
TMK1-ΔKD-B2-R	cloning	GGGGACCACTTTGTACAAGAAAGCTGGGTCGTTGTTAGTCACAGAACGAAG
TMK1KD-K616F	cloning	ATTGCGGTTAGGAGAATGGAGAATGGAGTTATTGCTGG
TMK1KD-K616R	cloning	CCATTCTCCATTCTCCTAACCGCAATCTTCGTTCCATC
pTMK1-B4-F	cloning	GGGGACAACTTTGTATAGAAAAGTTGCATCGTTTGGAACAGATCTTTCAGC
pTMK1-B1-R	cloning	GGGGACTGCTTTTTTGTACAAACTTGCAGCTTGAAGAAACAGAGGATTGAAG

pTMK2-B4-F	cloning	GGGGACAACTTTGTATAGAAAAGTTGCAGTAGAGGAATGTCGTTGATACCAC
pTMK2-B1-R	cloning	GGGGACTGCTTTTTTGTACAAACTTGCTTTCAACTGAGAAACAAACGCAGGG
pTMK3-B4-F	cloning	GGGGACAACTTTGTATAGAAAAGTTGCAGCCAATCAACAACCCTTTAATCG
pTMK3-B1-R	cloning	GGGGACTGCTTTTTTGTACAAACTTGCTTGTTGGAGAAAGAAACAGAGTGGA
pTMK4-B4-F	cloning	GGGGACAACTTTGTATAGAAAAGTTGCAACTCACAACTCCGTCGATCA
pTMK4-B1-R	cloning	GGGGACTGCTTTTTTGTACAAACTTGCATTAGAGAACACGCGCGGC
tmk1-1-F	genotyping	CCAGTTCCTGCGTCTTTGTT
tmk1-1-R	genotyping	TTAACCGCAATCTTCGTTCC
tmk2-1-R	genotyping	TTTCTCATTCCATGTTCCAGC
tmk2-1-F	genotyping	ACAAGGGAGAGTTGCATGATG
tmk3-1-R	genotyping	TATCAACACCAACGGATTTGC
tmk3-1-F	genotyping	TTGCCAGCTGTATTCACACAC
tmk4-1-F	genotyping	GTGGAATATCTCCATAGCTTGGCT
tmk4-1-R	genotyping	CCACAATTAAAACACAGCCTGTAA

 METHODS

Plant materials and growth conditions.

All mutants and transgenic lines are in the *Columbia-0* (*Col-0*) ecotype background with the exception

of *abp1-TD1*, which is in *Columbia-4* (*Col-4*). The T-DNA insertion lines *eir1-4* (SALK_091142) [*pin2*

mutant used in Figure 1], *tmk1-1* (SALK_016360), *tmk2-1* (SAIL_1242_H07), *tmk3-1* (SALK_129759)

and *tmk4-1* (GABI_348E01) were previously reported[35,59], as was *abp1-C1* and *abp1-TD1*[60], and *comp-*

C1 = (ABP1::GFP-ABP1/abp1-C1)[61]. The primers for genotyping are described in Supplementary

Table 3. *PIN2::PIN2-GFP*, *DR5rev::GFP* and *TMK1::TMK1-FLAG* in *tmk1-1* (*comp-tmk1-1*) were

described[59,62,63]. *DEX::TMK1^{K616R}-HA* line (unpublished) was kindly provided by William Gray. We

generated transgenic lines *PIN2::PIN2-GFP* in *tmk1-1* and *DR5rev::GFP* in *tmk1-1* by crossing the

respective markers and the mutant line.

To generate *TMK1::GUS, TMK2::GUS, TMK3::GUS, and TMK4::GUS* transgenic lines, genomic

fragments covering 2000 bp upstream from the start codon of *TMK1, TMK2, TMK3* and *TMK4* were

amplified from *Col-0* genomic DNA as a promotor region. We used these primers for amplification:

pTMK1-B4-F/pTMK1-B1-R, pTMK2-B4-F/pTMK2-B1-R, pTMK3-B4-F/pTMK3-B1-R, pTMK4-B4-

F/pTMK4-B1-R (see sequences in Supplemental Table 3). The promoter fragments were inserted into

pDONR P4-P1r vector by BP recombination reaction, resulting in *pDONR P4-P1R pTMK1, pDONR*

P4-P1R pTMK2, pDONR P4-P1R pTMK3, and pDONR P4-P1R pTMK4 plasmids. Those products were

subsequently cloned into pB7m24GW.3 vector together with pEN-L1-GUS-L2 (Invitrogen) by

MultiSite Gateway LR recombination reaction. To generate *TMK1::TMK1-GFP, 35S::TMK1$^{\Delta KD}$-GFP*

expressing lines, the *TMK1* full length/ *TMK1$^{\Delta KD}$* (amino acid 1-587) genomic DNA without stop codon

were amplified from DNA of *Col-0* seedlings by PCR using *TMK1-FL-B1-F* and *TMK1-FL-B2-*

R/TMK1-ΔKD-B2-R primers (see sequences Supplemental Table 3), respectively. The corresponding

TMK1 DNA sequences were inserted into pDONR221 vector by BP recombination reaction. Next, for

TMK1::TMK1-GFP construct, *pDONR P4-P1R pTMK1, pDONR221 gTMK1* and *pDONR P2R-P3*

GFP; for *35S::TMK1$^{\Delta KD}$-GFP, pDONR P4-P1R p35S, pDONR221 gTMK1$^{\Delta KD}$* and *pDONR P2R-P3*

GFP were cloned into pB7m34GW vector by MultiSite Gateway LR recombination reaction. To

generate *UBQ10::TMK1^{K616R}-2xmCherry, TMK1^{K616R}noSTOP/pDONR221* was obtained by site directed

mutagenesis, amplifying *pDONR221 TMK1* with the *TMK1KD-K616F/TMK1KD-K616R* primer pair.

pDONR221 TMK1^{K616R} was then recombined by LR reaction with *pDONR P4-P1R pUBQ10*[64], *pDONR*

P2R-P3 2xmCHERRY-4xMyc[65] and pH7m34GW[66] to introduce *UBQ10::TMK1^{K616R}-2xmCherry* in

pH7m34GW. All constructs were transformed into the *Agrobacterium tumefaciens* strain pGV3101 by

electroporation and further into *Col-0* plants by floral dip method. After being selected for single-copy

insertions, two individual homozygous T3 transgenic lines were used for further studies.

*Arabidopsi*s *thaliana* seeds were surface-sterilized with 70% (v/v) ethanol for 20 min, followed by

commercial bleach (2.5% [v/v] sodium hypochlorite) containing 0.05% (v/v) Triton X-100 for 10 min,

and finally washed four times with sterile distilled water. The stratification of the seeds was conducted

in the dark at 4°C for 3 days. *Arabidopsis thaliana* seedlings were grown at 22°C on ½ MS plates with

1% agar and 1% sucrose, or in soil with 16h light/8h dark cycles, photoperiod at 80 to 100 mE m^{-2} sec^{-}

^{1}.

Measurement of basipetal auxin transport in roots

Root basipetal auxin transport measurement was executed as previously described[67]. Briefly, 5-day-old

Arabidopsis seedlings were neatly placed on ½ MS plates and vertically grown for around 12 hours. 100

nM ^{3}H -IAA droplets were prepared by adding 2 µl of 25 Ci mmol^{-1} ^{3}H -IAA to 1 ml ½ MS with 1.25

% agar and quickly added dropwise onto a humid petri dish lid. Then the auxin-containing droplets were

carefully placed on root tips with about 0.5 mm overlap. Transport was allowed for 5 hours in darkness

at room temperature. 2 mm of root tissue proximal to the site of auxin application was discarded, and

an 8-mm root segment was cut from the rest of the root tip and harvested to 3 ml of scintillation fluid.

Radioactivity was counted with a scintillation counter (HIDEX 300 SL) for 100 seconds. 10 seedlings

were used for each treatment with 3 repetitions.

Measurement of basipetal auxin transport in shoots

Shoot basipetal auxin transport was carried out as previously described[68]. Briefly, surface-sterilized

Arabidopsis thaliana seeds were stratified in the dark at 4°C for 72 h, placed in the light for 6 hours to

induce germination and transferred to the darkness at 22°C for 5 days. Then, 5-day-old etiolated

hypocotyls were decapitated to exclude the effect of auxin biosynthesis in cotyledons and ^{3}H -IAA

droplets (60 nM: 12 µL ^{3}H-IAA in 10 mL of ½ MS medium with 1.25% agar) were carefully applied to

the apical part of the hypocotyls. After 6 hours incubation, treated hypocotyls were collected,

homogenized in liquid nitrogen and incubated overnight in Opti-Fluor scintillation cocktail (Perkin

Elmer). Finally, the radioactivity was measured with a scintillation counter (Hidex 300SL) for

300 seconds. 10 seedlings were used for each treatment with 3 repetitions. An additional background

control for any diffusion though the agar was performed by inserting the droplet above the decapitated

hypocotyl and treatment with 10 µM NPA as a negative control.

Mutant evaluation and expression analysis by RT-qPCR.

For RT-qPCR analyses, total RNA was prepared from max. 100mg of roots of 4-day-old seedlings

according to the RNeasy Plant Mini Kit (Qiagen). cDNA was synthesized from 1µg of total mRNA

using the QuantiNova Reverse Transcription Kit (Qiagen). For all mutant expression analyses, four

biological replicates were considered, and the samples were pipetted in three technical replicates into a

384 well plate using an automated JANUS Workstation (PerkinElmer). According to the manufacturer's

instructions, 5 µL reaction volume contained 2.5 µL Luna® Universal qPCR mastermix (NEB). RT-

qPCR analyses were performed using the Real-time PCR Roche Lightcycler 480 and the expression of

PP2AA3 (At1G13320) was used as in[69]. For each of the evaluated genes, three different primer pairs

were evaluated, and representative graphs are enclosed in the manuscript. The primers used for the

presented analyses are listed in Supplemental Table 3.

Förster Resonance Energy Transfer with Fluorescence Lifetime Imaging Microscopy (FRET-

FLIM).

FRET-FLIM experiments were performed in protoplast systems isolated from *Arabidopsis* root cell

suspension as described previously[70]. 10 µg of plasmid DNA (*TMK1-GFP, TMK1$^{\Delta KD}$-GFP, PIN2-*

mCherry, FLS2-GFP) were used for protoplast transfection and then incubated in a sterile 24-well

microtiter plate overnight in the dark at room temperature. FRET-FLIM experiments were performed

using a TriM Scope II inverted 2-photon microscope equipped with a FLIM X16 TCSPC Detector for

time correlated single photon counting (LaVision BioTec). Fluorescence lifetime image stacks (150

slices, with 0,082 ns time interval) were acquired. Image analyses were done in Fiji by performing a

threshold mask from the sum projection of each stack and by averaging all the pixels at each time point

of the stack. In order to yield the exponential decay with offset, the intensity at time point 0 was

normalized.

Histochemical GUS staining.

4-day-old seedlings of *TMK1::GUS, TMK3::GUS and TMK4::GUS* were stained in 0.1 M sodium

phosphate buffer (pH 7.0) containing 0.1% X-GlcA sodium salt, 1 mM $K_3[Fe(CN)_6]$, 1 mM $K_4[Fe(CN)_6]$

and 0.05% Triton X-100 for 20 min (TMK1::GUS), 2 h (TMK4::GUS) or 6 h (TMK3::GUS) at 37°C.

Further, samples were incubated overnight in 80% (v/v) ethanol at room temperature. Tissue clearing

was conducted as previously described[71]. DIC microscopy for analysis of GUS staining assay was

performed using an Olympus BX53 microscope equipped with 10x and 20x air objectives and a DP26

CCD camera**.**

Root Gravitropic Assay.

For measurements of root gravitropic curvature kinetics and growth rate, 4-day-old seedlings were

rotated 90° and roots were imaged every 30 min on a vertically placed flatbed scanner (Epson Perfection

V370 Photo). Image time series were stabilized using the StackReg Fiji plugin. Root curvature angle

(including root tip angle and rate of change) along with growth rate were analyzed using the Manual

Tracking Fiji plugin and Microsoft Excel.

Imaging with confocal laser scanning microscopy (CLSM).

Confocal laser scanning microscopy for analysis of fluorescent intensity was performed on Zeiss LSM800 microscope, positioned vertically[72], assisted with Zeiss Zen 2011 software. Pictures were taken by 20x objectives. Fluorescence signals for GFP (excitation 488 nm, emission 507 nm) and mCherry (excitation 561 nm; emission 580-650 nm)-tagged proteins were detected.

For the analysis of *PIN2::PIN2-GFP, DR5rev::GFP* and *TMK1::TMK1-GFP* asymmetric distribution during gravistimulation, 4-day-old seedlings were placed in a 1-well chambered coverglass (VWR, Kammerdeckgla¨ ser, Lab-Tek, Nunc - eine kammer, catalog number 734-2056) with a block of solid 0.5 MS medium (as described in [72]). In order to adjust seedlings to the new gravity vector, the chamber with samples was transferred to the darkness 1 h prior CLSM analysis, at the vertical position. Ten Z-sections spaced 4.5 mm for *PIN2* line and 1 mm for *DR5rev* and *TMK1* promoter analysis were collected in the median root section before and after 90° rotation at the indicated time points. Pictures were subsequently combined into the *sum* slices intensity projection. Marking epidermal and cortical regions together, the mean grey value of the signal intensity was collected and used for quantification (as described previously in [36]).

Protein extraction and western blot analysis.

Roots from 4-day-old seedlings expressing *TMK1::TMK1-GFP*, *TMK1::TMK1-FLAG* or *UBQ10::TMK1-3xHA* after gravistimulation or auxin treatments were frozen and grounded to powder in a liquid nitrogen. To analyze the *TMK1*-tagged expression level, proteins were isolated in a mixture of: 50 mM Tris–HCl pH 7.5, 150 mM NaCl, 5 mM dithiothiothreitol (DTT), 1 mM $Na_2MoO_4•2H_2O$, 1 mM NaF, 1% TritonX100, Complete EDTA-free Protease Inhibitor cocktail (Roche), PhosSTOP phosphatase inhibitor cocktail (Roche) and clarified by 15 min centrifugation step at 12000 g, 4°C. The analysis was carried out by SDS-PAGE and western blots using Anti-GFP HRP-conjugated (MACS Miltenyi Biotec), anti-FLAG®M2-Peroxidase (HRP) (Sigma), Anti-HA HRP-conjugated, High Affinity (3F10) (Roche) and SuperSignal West Femto Maximum Sensitivity Substrate Detection System (Thermo Scientific). Equal loading was ensured by Bradford protein quantification before loading, and by Ponceau staining of the membrane after protein transfer. Image capture was done using the image analyzer Amersham 600RGB 604 (GE Healthcare), and quantification of the protein signal was done using Fiji software.

Protein microsomal fraction extraction.

4-day-old seedling roots were harvested, grounded to powder in a liquid nitrogen, and vortexed vigorously in extraction buffer 1 (50 mM Tris–HCl pH 7.5, 150 mM NaCl, Complete EDTA-free Protease Inhibitor cocktail (Roche), PhosSTOP phosphatase inhibitor cocktail (Roche)) in 1/10 (w/v) ratio. The resulting homogenate was centrifuged at 20000 g for 30 min at 4°C. The pellet was

resuspended in extraction buffer 2 (50 mM Tris–HCl pH 7.5, 150 mM NaCl, 0.5% Nonidet P-40, 1%
TritonX100, Complete EDTA-free Protease Inhibitor cocktail (Roche), PhosSTOP phosphatase
inhibitor cocktail (Roche)), and centrifuged at 12000 g for 20 min at 4°C. The supernatant was used for
total protein concentration quantification by Bradford assay. Samples were separated by SDS-PAGE (4-
15% Mini-PROTEAN®TGX™ Precast Protein Gels (Bio-RAD)), transferred to a PVDF
(polyvinylidene difluoride) membrane and analyzed by immunoblot using the following primary
antibodies: affinity-purified TMK1 (1:1000, [59]), AHA2 (1:1000, [73]) and PIN2 (1:1000, [35]) antibodies and
anti-ROP6 (C) (1:1000, Abiocode), using anti-Rabbit HRP-conjugated (1:5000, GE Healthcare) as a
secondary antibody. Detection was performed using the SuperSignal West Femto Maximum Sensitivity
Substrate detection kit (ThermoFisher Scientific). Image capture was done using the image analyzer
Amersham 600RGB (GE Healthcare), and quantification of the protein signal was done using Fiji
software.

Western blot analysis of phosphorylated proteins.
To analyze the phosphorylation status of TMK1 *in planta*, after corresponding treatment, 4-day-old
tmk1-1 roots expressing *TMK1::TMK1-FLAG* were grounded to powder in liquid nitrogen and
homogenized in ice-cold sucrose buffer (20 mM Tris–HCl pH 8, 0.33 M sucrose, Complete EDTA-free
Protease Inhibitor cocktail (Roche), PhosSTOP phosphatase inhibitor cocktail (Roche)), followed by a
centrifugation step at 5000 g for 10 min at 4°C. To obtain the membrane protein fraction, the supernatant
was centrifuged at 20000 g for 30 min at 4°C and the resulting pellet was solubilized with lysis buffer
(20 mM Tris–HCl pH 8, 150 mM NaCl, 1% TritonX100, Complete EDTA-free Protease Inhibitor
cocktail (Roche), PhosSTOP phosphatase inhibitor cocktail (Roche)) and centrifuged at 20000 g for 10
min at 4°C. The corresponding supernatant was used for immunoprecipitation assay with anti-FLAG
micro beads according to the manufacturer's instructions (μMACS Epitope Tag Protein Isolation Kit
(MACS Miltenyi Biotec)). Samples were separated by SDS-PAGE (4-15% Mini-PROTEAN®TGX™
Precast Protein Gels (Bio-RAD)) and transferred to a PVDF membrane followed by probe with Phos-
tag BTL-111 according to the manufacturer's instructions (www.wako-chem.co.jp) to detect protein
phosphorylated levels and anti-FLAG®M2-Peroxidase (HRP) (Sigma) antibody to detect the total
amount of immunoprecipitated TMK1-FLAG.

Co-Immunoprecipitation assays.
4-day-old *tmk1-1* seedlings expressing *TMK1::TMK1-FLAG* were treated with 5, 20 nM IAA or DMSO
control for 30 minutes. Roots were harvested, grounded to powder in liquid nitrogen, and subjected to
protein microsomal fraction extraction. Solubilized proteins from microsomal fraction were
immunoprecipitated using super-paramagnetic micro MACS beads coupled to monoclonal anti-FLAG
antibody according to the manufacturer's instructions (Miltenyi Biotec). WT *Col-0* extract was used as
a control of the unspecific binding of endogen PIN2. Proteins immunoprecipitated with anti-FLAG

antibody were separated by SDS-PAGE (4-15% Mini-PROTEAN®TGX™ Precast Protein Gels (Bio-RAD)), transferred to a PVDF membrane and analyzed by immunoblot with anti-FLAG®M2-Peroxidase (HRP) (Sigma) antibody to detect TMK1-FLAG and with anti-PIN2 antibody to detect co-immunoprecipitated endogenous PIN2.

Active ROP assay of non-overexpressing plants

5-day-old seedling roots were excised and incubated in ½ MS liquid medium. The roots were treated with 100 nM IAA for 10 minutes and frozen in liquid nitrogen. Total protein was extracted from the treated root tissues using Extraction buffer (25mM HEPES pH 7.4, 100mM KCl, 10mM MgCl2, 1 mM PMSF, 5 mM Na3VO4, 5 mM NaF, 1 mM TCEP, cOmplete Protease Inhibitor Cocktail, Roche) in 2 steps. For every 100 mg of grounded tissue, 400 μl of extraction buffer were added, then incubated with another 400 μl of extraction buffer containing 5% Triton X-100 for one hour. Active ROP proteins were pulled down using His-MBP-CRIB (originally from RIC1 effector, obtained from expressing BL21 *E.coli* total cell extract) conjugated to HisPur™ Ni-NTA Magnetic Beads (Thermo Fisher) by incubation of 200 μl total protein extract with 25 μl (initial volume) of beads for 2 hours at 4°C. The beads were washed by Washing buffer (25 mM HEPES pH 7.4, 300 mM KCl, 10 mM MgCl$_2$, 12.5 mM imidazole) for 3 times and boiled at 95°C with 30 μl SDS loading dye (BioRad). After removing the beads, the protein samples were separated by 12% SDS-PAGE and analyzed by immunoblot with anti-ROP2 specific antibody (1:10,000, Abiocode). Input samples were isolated from total protein extracts, before addition of Triton X-100 to get an estimate of the present total ROP content.

Quantitative immunoblot analysis.

To determine the relative amount of protein in time course experiments, grey values were calculated by ImageJ Gel Analysis tool, where for each band of protein of interest (PI) and normalizing control (NC), membrane background was removed individually. To get relative NC value, all the NC values were divided by the highest one. Subsequently, all the PI values were divided by their respective relative NC values. Final obtained values of PI were normalized dividing by the time point 0 or DMSO control treatment value. Values from three independent experiments were used to determine average and ±SD. For example, for quantitative immunoblot analysis of phosphorylated proteins, it was first determined grey values of TMK1 obtained with Phos-tag BTL-111 probe and anti-Flag antibody at different time points/treatments, tacking into account each membrane background. TMK1-Flag values were used as a protein amount normalizing control (NC). To get NC values of each experiment point, TMK1-Flag values were divided by the highest one. Subsequently, TMK1 Phos-tag values were divided by their respective relative NC values. To compare differences between time points/ treatments, as a final step,

the normalized at protein level TMK1 Phos-tag values were divided by the corresponding time point 0
or DMSO control treatment values.

Recombinant protein expression and purification from *E. coli.*

The 6xHis-PIN2 HL recombinant protein was expressed using the 6xHis-PIN2 HL vector [74] in *E. coli*
BL21 (DE3) strain upon induction by 0.5 mM IPTG (Isopropyl β-D-1-Thiogalactopyranoside) at 16°C
for 12 h. Cells were harvested by centrifugation at 4000 g for 15 min and washed with water. They were
then lysed by sonication in lysis buffer (50 mM Tris-HCl pH7.5; 150 mM NaCl; 1 mM EDTA; 1 mM
DTT and 1% TritonX100). The lysed solution was centrifuged at 12000 g for 15 min. The supernatant
was then purified using Ni-NTA His affinity agarose (Thermo Scientific) according to the
manufacturer's instructions. 6xHis- PIN2 HL protein was eluted from the beads in elution buffer (50
mM Tris-HCl pH7.4; 150 mM NaCl and 250 mM imidazole). Eluted protein samples were checked by
SDS-PAGE and Coomassie brilliant blue staining (Bio-Safe™ Coomassie Stain, Bio-Rad). The protein
concentration was determined with the Bradford method (Quick Start™ Bradford Reagent, Bio-Rad).

TMK1-HA immunoprecipitation.

To immunoprecipitate HA-tagged TMK1 protein, roots of 7-day-old *pUBQ10::TMK1-3xHA* and
DEX::TMK1^{K616R}-HA (previously induced for 48 h with 30 µM dexamethasone) seedlings were frozen
and grounded in a liquid nitrogen, and, subsequently, homogenized in protein extraction buffer (PEB:
50 mM Tris-HCl pH7.5, 200 mM NaCl, 1% TritonX100, 10 mM MgCl$_2$, 1 mM MnCl$_2$, Complete
(Roche) protease cocktail and PhosSTOP phosphatase inhibitor cocktail (Roche)), followed by the 20
min centrifugation step at 14000 g, 4°C. In a fresh Eppendorf tube, 50 µL anti-HA agarose beads were
added (Anti-HA Affinity Matrix, SIGMA; pre-washed with 200 µL PEB) to the supernatant. After 4 h
of rotating at 4°C, the samples were spun down again at 2000 g, 4°C, and the supernatant was discarded.
The Agarose beads were washed twice with 200 µL PEB, and finally re-suspended into 50 µL PEB to
allow further biochemical reactions.

***In vitro* kinase assay.**

The *in vitro* kinase assay with [γ-^{32}P]-ATP was conducted as reported[74] with minor modifications.
Immunoprecipated TMK1-HA and TMK1^{K616R}-HA (5 µL) from 7-day-old *Arabidopsis* seedlings,
together with the recombinant 6xHis-PIN2 HL (10 µL) from *E.coli*, were added to the kinase reaction
buffer (50 mM Tris-HCl pH 7.5, 10 mM MgCl$_2$, 5 mM NaCl, 2.5 mM cold ATP (adenosine 5'-
triphosphate), and 1 mM DTT) in the presence of 5 µCi [γ-^{32}P]-ATP (NEG502A001MC; Perkin-Elmer).
Reactions were incubated at 25°C for 90 min and afterwards terminated by adding 10 µL 5×SDS loading
buffer. 20 µL reaction samples were then separated with 10% SDS-PAGE gel, developed with a
phosphor-plate overnight. Eventually, the phosphor-plate was imaged with a Fujifilm FLA 3000 plus
DAGE system.

Statistical analysis.

Data visualization was conducted using the GraphPad Prism 8 Software or in R version 4.0.4. For statistical comparisons we used standard linear models (t-test, one- or two-way ANOVA with Dunnett's multiple comparisons) after first checking their assumptions. Some charts in Figure 1 represent pooled data from multiple experiments. In such cases, we fitted mixed linear models with *restricted maximum likelihood* using the *lme()* function implemeted as part of the *nlme* R package. We tested the influence of genotype/treatment as a fixed effect (type II analysis of deviance) and included experimental repetition as a random effect. The random effect was tested by comparing a model where it was included to a model with just the fixed effect using the *gls()* function. The result was reported as the amount by which the Akaike Information Criterion (AIC) of the model increased when excluding the random effect, along with the associated p-value.

Accession Numbers

Sequence data from this article can be found in the *Arabidopsis* Genome Initiative databases under the following accession numbers: At4g02980 for *ABP1*, At1g66150 for *TMK1*, At1g24650 for *TMK2*, At2g01820 for *TMK3*, At3g23750 for *TMK4*, At5g57090 for *PIN2*, At1g20090 for *ROP2*, At4g35020 for *ROP6*, At4g30190 for *AHA2*, At5g46330 for *FLS2*.

REFERENCES

1. Darwin, C., and Darwin, F. (1897). The Power of Movement in Plants. Appleton.

2. Friml, J. (2021). Fourteen Stations of Auxin. Cold Spring Harb. Perspect. Biol. *14*, a039859. 10.1101/cshperspect.a039859.

3. Berleth, T., and Sachs, T. (2001). Plant morphogenesis: Long-distance coordination and local patterning. Curr. Opin. Plant Biol. *4*, 57–62. 10.1016/S1369-5266(00)00136-9.

4. Hajný, J., Tan, S., and Friml, J. (2022). Auxin canalization: From speculative models toward molecular players. Curr. Opin. Plant Biol. *65*, 102174. 10.1016/J.PBI.2022.102174.

5. Zhang, J., Mazur, E., Balla, J., Gallei, M., Kalousek, P., Medveďová, Z., Li, Y., Wang, Y., Prát, T., Vasileva, M., et al. (2020). Strigolactones inhibit auxin feedback on PIN-dependent auxin transport canalization. Nat. Commun. *11*, 1–10. 10.1038/s41467-020-17252-y.

6. Adamowski, M., and Friml, J. (2015). PIN-dependent auxin transport: Action, regulation, and evolution. Plant Cell *27*, 20–32. 10.1105/tpc.114.134874.

7. Luschnig, C., and Vert, G. (2014). The dynamics of plant plasma membrane proteins: PINs and beyond. Dev. *141*, 2924–2938. 10.1242/dev.103424.

8. Park, J., Lee, Y., Martinoia, E., and Geisler, M. (2017). Plant hormone transporters: what we know and what we would like to know. BMC Biol. *15*, 1–15. 10.1186/S12915-017-0443-X.

9. Péret, B., Swarup, K., Ferguson, A., Seth, M., Yang, Y., Dhondt, S., James, N., Casimiro, I., Perry, P., Syed, A., et al. (2012). AUX/LAX Genes Encode a Family of Auxin Influx Transporters That Perform Distinct Functions during Arabidopsis Development. Plant Cell *24*, 2874–2885. 10.1105/tpc.112.097766.

10. Zhang, J., Nodzyński, T., Pěnčík, A., Rolčík, J., and Friml, J. (2010). PIN phosphorylation is sufficient to mediate PIN polarity and direct auxin transport. Proc. Natl. Acad. Sci. U. S. A. *107*, 918–922. 10.1073/pnas.0909460107.

11. Wisniewska, J., Xu, J., Seifartová, D., Brewer, P.B., Růžička, K., Blilou, L., Rouquié, D., Benková, E., Scheres, B., and Friml, J. (2006). Polar PIN localization directs auxin flow in plants. Science *312*, 883. 10.1126/science.1121356.

12. Prigge, M.J., Platre, M., Kadakia, N., Zhang, Y., Greenham, K., Szutu, W., Pandey, B.K., Bhosale, R.A., Bennett, M.J., Busch, W., et al. (2020). Genetic analysis of the Arabidopsis

TIR1/ AFB auxin receptors reveals both overlapping and specialized functions. Elife *9*,
e54740.

13.   Qi, L., Kwiatkowski, M., Chen, H., Hoermayer, L., Sinclair, S., Zou, M., Del Genio, C.I.,
Kubeš, M.F., Napier, R., Jaworski, K., et al. (2022). Adenylate cyclase activity of TIR1/AFB
auxin receptors in plants. Nature *611*, 133–138. https://doi.org/10.1038/s41586-022-05369-7.

14.   Li, L., Verstraeten, I., Roosjen, M., Takahashi, K., Rodriguez, L., Merrin, J., Chen, J., Shabala,
L., Smet, W., Ren, H., et al. (2021). Cell surface and intracellular auxin signalling for H+
fluxes in root growth. Nature *599*, 273–277. 10.1038/s41586-021-04037-6.

15.   Friml, J., Gallei, M., Gelová, Z., Johnson, A., Mazur, E., Monzer, A., Rodriquez, L., Roosjen,
M., Verstraeten, I., Živanovič, B.D., et al. (2022). ABP1–TMK auxin perception for global
phosphorylation and auxin canalization. Nature *609*, 575–581. 10.1038/s41586-022-05187-x.

16.   Xu, T., Dai, N., Chen, J., Nagawa, S., Cao, M., Li, H., Zhou, Z., Chen, X., De Rycke, R.,
Rakusová, H., et al. (2014). Cell surface ABP1-TMK auxin-sensing complex activates ROP
GTPase signaling. Science *343*, 1025–1028. 10.1126/science.1245125.

17.   Mazur, E., Kulik, I., Hajný, J., and Friml, J. (2020). Auxin canalization and vascular tissue
formation by TIR1/AFB-mediated auxin signaling in Arabidopsis. New Phytol. *226*, 1375–
1383. 10.1111/NPH.16446.

18.   Wabnik, K., Kleine-Vehn, J., Balla, J., Sauer, M., Naramoto, S., Reinö Hl, V., Merks, R.M.,
Govaerts, W., and Friml, J. (2010). Emergence of tissue polarization from synergy of
intracellular and extracellular auxin signaling. Mol. Syst. Biol. *6*, 447. 10.1038/msb.2010.103.

19.   Narasimhan, M., Gallei, M., Tan, S., Johnson, A., Verstraeten, I., Li, L., Rodriguez, L., Han,
H., Himschoot, E., Wang, R., et al. (2021). Systematic analysis of specific and nonspecific
auxin effects on endocytosis and trafficking. Plant Physiol. *186*, 1122–1142.
10.1093/PLPHYS/KIAB134.

20.   Narasimhan, M., Johnson, A., Prizak, R., Kaufmann, W.A., Tan, S., Casillas-Pérez, B., and
Friml, J. (2020). Evolutionarily unique mechanistic framework of clathrin-mediated
endocytosis in plants. Elife *9*, 1–30. 10.7554/eLife.52067.

21.   Naramoto, S., Kleine-Vehn, J., Robert, S., Fujimoto, M., Dainobu, T., Paciorek, T., Ueda, T.,
Nakano, A., Van Montagu, M.C.E., Fukuda, H., et al. (2010). ADP-ribosylation factor
machinery mediates endocytosis in plant cells. Proc. Natl. Acad. Sci. *107*, 21890–21895.
10.1073/pnas.1016260107.

22.   Naramoto, S., Otegui, M.S., Kutsuna, N., de Rycke, R., Dainobu, T., Karampelias, M.,
Fujimoto, M., Feraru, E., Miki, D., Fukuda, H., et al. (2014). Insights into the Localization and

Function of the Membrane Trafficking Regulator GNOM ARF-GEF at the Golgi Apparatus in
Arabidopsis. Plant Cell *26*, 3062–3076. 10.1105/TPC.114.125880.

23. Glanc, M., Van Gelderen, K., Hoermayer, L., Tan, S., Naramoto, S., Zhang, X., Domjan, D.,
Včelařová, L., Hauschild, R., Johnson, A., et al. (2021). AGC kinases and MAB4/MEL
proteins maintain PIN polarity by limiting lateral diffusion in plant cells. Curr. Biol. *31*, 1918–
1930. 10.1016/J.CUB.2021.02.028.

24. Tan, S., Zhang, X., Kong, W., Yang, X.L., Molnár, G., Vondráková, Z., Filepová, R., Petrášek,
J., Friml, J., and Xue, H.W. (2020). The lipid code-dependent phosphoswitch PDK1–D6PK
activates PIN-mediated auxin efflux in Arabidopsis. Nat. Plants *6*, 556–569. 10.1038/s41477-
020-0648-9.

25. Xiao, Y., and Offringa, R. PDK1 regulates auxin transport and Arabidopsis vascular
development through AGC1 kinase PAX. *6*, 544–555. 10.1038/s41477-020-0650-2.

26. Zourelidou, M., Absmanner, B., Weller, B., Barbosa, I.C.R., Willige, B.C., Fastner, A., Streit,
V., Port, S.A., Colcombet, J., van Bentem, S. de la F., et al. (2014). Auxin efflux by PIN-
FORMED proteins is activated by two different protein kinases, D6 PROTEIN KINASE and
PINOID. Elife *2014*. 10.7554/ELIFE.02860.

27. Zhang, Y., Xiao, G., Wang, X., Zhang, X., and Friml, J. (2019). Evolution of fast root
gravitropism in seed plants. Nat. Commun. *10*, 1–10. 10.1038/s41467-019-11471-8.

28. Sato, E.M., Hijazi, H., Bennett, M.J., Vissenberg, K., and Swarup, R. (2015). New insights into
root gravitropic signalling. J. Exp. Bot. *66*, 2155–2165. 10.1093/JXB/ERU515.

29. Morita, M.T. (2010). Directional Gravity Sensing in Gravitropism. Annu. Rev. Plant Biol. *61*,
705–720. 10.1146/annurev.arplant.043008.092042.

30. Friml, J., Wisniewska, J., Benkova, E., Mendgen, K., and Palme, K. (2002). Lateral relocation
of auxin efflux regulator PIN3 mediates tropism in Arabidopsis. Nature *415*, 806–809.
10.1038/415806a.

31. Kleine-Vehn, J., Ding, Z., Jones, A.R., Tasaka, M., Morita, M.T., and Friml, J. (2010).
Gravity-induced PIN transcytosis for polarization of auxin fluxes in gravity-sensing root cells.
Proc. Natl. Acad. Sci. U. S. A. *107*, 22344–22349. 10.1073/pnas.1013145107.

32. Luschnig, C., Gaxiola, R.A., Grisafi, P., and Fink, G.R. (1998). EIR1, a root-specific protein
involved in auxin transport, is required for gravitropism in Arabidopsis thaliana. Genes Dev.
12, 2175–2187. 10.1101/gad.12.14.2175.

33. Swarup, R., Friml, J., Marchant, A., Ljung, K., Sandberg, G., Palme, K., and Bennett, M.

(2001). Localization of the auxin permease AUX1 suggests two functionally distinct hormone

transport pathways operate in the Arabidopsis root apex. Genes Dev. *15*, 2648–2653.

10.1101/gad.210501.

34.   Fendrych, M., Akhmanova, M., Merrin, J., Glanc, M., Hagihara, S., Takahashi, K., Uchida, N.,

Torii, K.U., and Friml, J. (2018). Rapid and reversible root growth inhibition by TIR1 auxin

signalling. Nat. Plants *4*, 453–459. 10.1038/s41477-018-0190-1.

35.   Abas, L., Benjamins, R., Malenica, N., Paciorek, T.T., Wiřniewska, J., Moulinier-Anzola, J.C.,

Sieberer, T., Friml, J., and Luschnig, C. (2006). Intracellular trafficking and proteolysis of the

Arabidopsis auxin-efflux facilitator PIN2 are involved in root gravitropism. Nat. Cell Biol. *8*,

249–256. 10.1038/ncb1369.

36.   Baster, P., Robert, S., Kleine-Vehn, J., Vanneste, S., Kania, U., Grunewald, W., De Rybel, B.,

Beeckman, T., and Friml, J. (2013). SCFTIR1/AFB-auxin signalling regulates PIN vacuolar

trafficking and auxin fluxes during root gravitropism. EMBO J. *32*, 260–274.

10.1038/emboj.2012.310.

37.   Retzer, K., Akhmanova, M., Konstantinova, N., Malínská, K., Leitner, J., Petrášek, J., and

Luschnig, C. (2019). Brassinosteroid signaling delimits root gravitropism via sorting of the

Arabidopsis PIN2 auxin transporter. Nat. Commun. *10*, 1–15. 10.1038/s41467-019-13543-1.

38.   Hajný, J., Prát, T., Rydza, N., Rodriguez, L., Tan, S., Verstraeten, I., Domjan, D., Mazur, E.,

Smakowska-Luzan, E., Smet, W., et al. (2020). Receptor kinase module targets PIN-dependent

auxin transport during canalization. Science *370*, 550–557. 10.1126/science.aba3178.

39.   Roosjen, M., Kuhn, A., Mutte, S.K., Boeren, S., Krupar, P., Koehorst, J., Fendrych, M., Friml,

J., and Weijers, D. (2022). An ultra-fast, proteome-wide response to the plant hormone auxin.

bioRxiv, 2022.11.25.517949. https://doi.org/10.1101/2022.11.25.517949.

40.   Kuhn, A., Roosjen, M., Sumanth, M., Dubey, S.M., Carasco, P.C., Monzer, A., Kohchi, T.,

Nishihama, R., Fendrych, M., Friml, J., et al. (2022). A RAF-like kinase mediates a deeply

conserved, ultra-rapid auxin response. bioRxiv, 2022.11.25.517951.

https://doi.org/10.1101/2022.11.25.517951.

41.   Tan, S., Luschnig, C., and Friml, J. (2021). Pho-view of Auxin: Reversible Protein

Phosphorylation in Auxin Biosynthesis, Transport and Signaling. Mol. Plant *14*, 151–165.

10.1016/J.MOLP.2020.11.004.

42.   Jia, W., Li, B., Li, S., Liang, Y., Wu, X., Ma, M., Wang, J., Gao, J., Cai, Y., Zhang, Y., et al.

(2016). Mitogen-Activated Protein Kinase Cascade MKK7-MPK6 Plays Important Roles in

Plant Development and Regulates Shoot Branching by Phosphorylating PIN1 in Arabidopsis.

PLOS Biol. *14*, e1002550. 10.1371/journal.pbio.1002550.

43.   Abas, L., Kolb, M., Stadlmann, J., Janacek, D.P., Lukic, K., Schwechheimer, C., Sazanov,
L.A., Mach, L., Friml, J., and Hammes, U.Z. (2020). Naphthylphthalamic acid associates with
and inhibits PIN auxin transporters. Proc. Natl. Acad. Sci. U. S. A. *118*, 1–8.
10.1073/pnas.2020857118.

44.   Pan, X., Fang, L., Liu, J., Senay-Aras, B., Lin, W., Zheng, S., Zhang, T., Guo, J., Manor, U.,
Van Norman, J., et al. (2020). Auxin-induced signaling protein nanoclustering contributes to
cell polarity formation. Nat. Commun. *11*, 1–14. 10.1038/s41467-020-17602-w.

45.   Marquès-Bueno, M.M., Armengot, L., Noack, L.C., Bareille, J., Rodriguez, L., Platre, M.P.,
Bayle, V., Liu, M., Opdenacker, D., Vanneste, S., et al. (2021). Auxin-Regulated Reversible
Inhibition of TMK1 Signaling by MAKR2 Modulates the Dynamics of Root Gravitropism.
Curr. Biol. *31*, 228–237. 10.1016/J.CUB.2020.10.011.

46.   Dai, N., Wang, W., Patterson, S.E., and Bleecker, A.B. (2013). The TMK Subfamily of
Receptor-Like Kinases in Arabidopsis Display an Essential Role in Growth and a Reduced
Sensitivity to Auxin. PLoS One *8*, 60990. 10.1371/journal.pone.0060990.

47.   Chang, C., Eric Schaller, G., Patterson, S.E., Kwok, S.F., and Meyerowitz, E.M. (1992). The
TMKl Gene from Arabidopsis Codes for a Protein with Structural and Biochemical
Characteristics of a Receptor Protein Kinase. Plant Cell *4*, 1263–1271. 10.1105/tpc.4.10.1263.

48.   Oh, M.H., Wang, X., Kota, U., Goshe, M.B., Clouse, S.D., and Huber, S.C. (2009). Tyrosine
phosphorylation of the BRI1 receptor kinase emerges as a component of brassinosteroid
signaling in Arabidopsis. Proc. Natl. Acad. Sci. U. S. A. *106*, 658–663.
10.1073/PNAS.0810249106.

49.   Wang, X., Goshe, M.B., Soderblom, E.J., Phinney, B.S., Kuchar, J.A., Li, J., Asami, T.,
Yoshida, S., Huber, S.C., and Clouse, S.D. (2005). Identification and Functional Analysis of in
Vivo Phosphorylation Sites of the Arabidopsis BRASSINOSTEROID-INSENSITIVE1
Receptor Kinase. Plant Cell *17*, 1685–1703. 10.1105/TPC.105.031393.

50.   Paciorek, T., Zažímalová, E., Ruthardt, N., Petrášek, J., Stierhof, Y.D., Kleine-Vehn, J.,
Morris, D.A., Emans, N., Jürgens, G., Geldner, N., et al. (2005). Auxin inhibits endocytosis
and promotes its own efflux from cells. Nature *435*, 1251–1256. 10.1038/nature03633.

51.   Morffy, N., and Strader, L.C. (2022). Structural Aspects of Auxin Signaling. Cold Spring Harb.
Perspect. Biol. *14*, a039883. 10.1101/cshperspect.a039883.

52.   Napier, R. (2021). The Story of Auxin-Binding Protein 1 (ABP1). Cold Spring Harb. Perspect.
Biol. *13*, a039909. 10.1101/cshperspect.a039909.

53. Er, E.E., Mendoza, M.C., Mackey, A.M., Rameh, L.E., and Blenis, J. (2013). AKT Facilitates
EGFR Trafficking and Degradation by Phosphorylating and Activating PIKfyve. Sci. Signal. *6*,
1–11. 10.1126/scisignal.2004015.

54. Goh, L.K., and Sorkin, A. (2013). Endocytosis of Receptor Tyrosine Kinases. Cold Spring
Harb. Perspect. Biol., a017459. 10.1101/cshperspect.a017459.

55. Yu, Y., Tang, W., Lin, W., Zhou, X., Li, W., Li, Y., Chen, R., Cao, W., Perez-Henriquez, P.,
Huang, R., et al. (2022). ABLs and TMKs are co-receptors for extracellular auxin. bioRxiv,
2022.11.28.518138. https://doi.org/10.1101/2022.11.28.518138.

56. Wang, J., Chang, M., Huang, R., Jiri, F., Yu, Y., Wen, M., Yang, Z., and Xu, T. (2022). Self-
regulation of PIN1-driven auxin transport by cell surface-based auxin signaling in Arabidopsis.
co-submitted to Cell.

57. Wang, Y., Yan, X., Xu, M., Qi, W., Shi, C., Li, X., Ma, J., Tian, D., Shou, J., Wu, H., et al.
(2022). TMK1-mediated auxin signal regulates membrane-associated clathrin in Arabidopsis
roots. J. Integr. Plant Biol. 10.1111/jipb.13366.

58. Brady, S.M., Orlando, D.A., Lee, J.Y., Wang, J.Y., Koch, J., Dinneny, J.R., Mace, D., Ohler,
U., and Benfey, P.N. (2007). A high-resolution root spatiotemporal map reveals dominant
expression patterns. Science *318*, 801–806. 10.1126/SCIENCE.1146265.

59. Cao, M., Chen, R., Li, P., Yu, Y., Zheng, R., Ge, D., Zheng, W., Wang, X., Gu, Y., Gelová, Z.,
et al. (2019). TMK1-mediated auxin signalling regulates differential growth of the apical hook.
Nature *568*, 240–243. 10.1038/s41586-019-1069-7.

60. Gao, Y., Zhang, Y., Zhang, D., Dai, X., Estelle, M., and Zhao, Y. (2015). Auxin binding
protein 1 (ABP1) is not required for either auxin signaling or Arabidopsis development. Proc.
Natl. Acad. Sci. U. S. A. *112*, 2275–2280. 10.1073/PNAS.1500365112.

61. Gelová, Z., Gallei, M., Pernisová, M., Brunoud, G., Zhang, X., Glanc, M., Li, L., Michalko, J.,
Pavlovičová, Z., Verstraeten, I., et al. (2021). Developmental roles of Auxin Binding Protein 1
in Arabidopsis thaliana. Plant Sci. *303*, 110750. 10.1016/j.plantsci.2020.110750.

62. Friml, J.J., Vieten, A., Sauer, M., Weijers, D., Schwarz, H., Hamann, T., Offringa, R., Jurgens,
G., Cao, M., Chen, R., et al. (2003). Efflux-dependent auxin gradients establish the apical-basal
axis of Arabidopsis. Nature *426*, 147–153. 10.1038/nature02085.

63. Xu, J., and Scheres, B. (2005). Dissection of arabidopsis ADP-ribosylation factor 1 function in
epidermal cell polarity. Plant Cell *17*, 525–536. 10.1105/tpc.104.028449.

64. Jaillais, Y., Hothorn, M., Belkhadir, Y., Dabi, T., Nimchuk, Z.L., Meyerowitz, E.M., and

Chory, J. (2011). Tyrosine phosphorylation controls brassinosteroid receptor activation by

triggering membrane release of its kinase inhibitor. Genes Dev. *25*, 232–237.

10.1101/gad.2001911.

65.   Simon, M.L.A., Platre, M.P., Assil, S., van Wijk, R., Chen, W.Y., Chory, J., Dreux, M.,

Munnik, T., and Jaillais, Y. (2014). A multi-colour/multi-affinity marker set to visualize

phosphoinositide dynamics in Arabidopsis. Plant J. *77*, 322–337. 10.1111/tpj.12358.

66.   Karimi, M., Depicker, A., and Hilson, P. (2007). Recombinational cloning with plant gateway

vectors. Plant Physiol. *145*, 1144–1154. 10.1104/pp.107.106989.

67.   Lewis, D.R., and Muday, G.K. (2009). Measurement of auxin transport in Arabidopsis thaliana.

Nat. Protoc. *4*, 437–451. 10.1038/nprot.2009.1.

68.   Grones, P., Abas, M., Hajný, J., Jones, A., Waidmann, S., Kleine-Vehn, J., and Friml, J.

(2018). PID/WAG-mediated phosphorylation of the Arabidopsis PIN3 auxin transporter

mediates polarity switches during gravitropism. Sci. Rep. *8*, 10279. 10.1038/s41598-018-

28188-1.

69.   Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K., and Scheible, W.R. (2005). Genome-

wide identification and testing of superior reference genes for transcript normalization in

arabidopsis. Plant Physiol. *139*, 5–17. 10.1104/pp.105.063743.

70.   Grones, P., Chen, X., Simon, S., Kaufmann, W.A., De Rycke, R., Nodzyński, T., Zažímalová,

E., and Friml, J. (2015). Auxin-binding pocket of ABP1 is crucial for its gain-of-function

cellular and developmental roles. J. Exp. Bot. *66*, 5055–5065. 10.1093/jxb/erv177.

71.   Malamy, J.E., and Benfey, P.N. (1997). Organization and cell differentiation in lateral roots of

Arabidopsis thaliana. Development *124*, 33–44. 10.1242/dev.124.1.33.

72.   von Wangenheim, D., Hauschild, R., Fendrych, M., Barone, V., Benková, E., and Friml, J.

(2017). Live tracking of moving samples in confocal microscopy for vertically grown roots.

Elife *6*. 10.7554/eLife.26792.

73.   Hayashi, Y., Nakamura, S., Takemiya, A., Takahashi, Y., Shimazaki, K., and Kinoshita, T.

(2010). Biochemical Characterization of In Vitro Phosphorylation and Dephosphorylation of

the Plasma Membrane H+-ATPase. Plant Cell Physiol. *51*, 1186–1196. 10.1093/pcp/pcq078.

74.   Tan, S., Abas, M., Verstraeten, I., Glanc, M., Molnár, G., Hajný, J., Lasák, P., Petřík, I.,

Russinova, E., Petrášek, J., et al. (2020). Salicylic Acid Targets Protein Phosphatase 2A to

Attenuate Growth in Plants. Curr. Biol. *30*, 381–395. 10.1016/j.cub.2019.11.058.

1201

1202

1203

1204

Figure 1.

Figure 1. ABP1-TMK1 auxin perception targets PIN phosphorylation and polar auxin transport.

(A) Overview of PIN phospho-sites significantly regulated (FDR < 0.05) in *tmk1-1* mutant roots compared to WT after treatment with auxin (100 nM, 2 min). 4 independent biological replicates, mean ± SD, permutation-based t-tests with FDR-controlled p-values. Highlighted in bold are sites which are co-regulated in the *abp1-TD1* mutant. Data adapted from [15]. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

(B) Overview of PIN phospho-sites significantly regulated (FDR < 0.05) in *abp1-TD1* mutant roots compared to WT after treatment with auxin (100 nM, 2 min). 4 independent biological replicates, mean ± SD, permutation-based t-tests with FDR-controlled p-values. Highlighted in bold are sites which are co-regulated in the *tmk1-1* mutant. Data adapted from [15]. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

(C) Basipetal/rootward transport of radioactive ^{3}H-IAA in decapitated hypocotyls is defective in the *tmk1-1* mutant (genotype/treatment fixed effect: χ^2 = 60.64, df = 2, p = 6.78e-14). 10 μM NPA, an inhibitor of auxin transport, was used as a negative control. Pooled data from two independent experiments with similar results (repetition random effect: ΔAIC = 2.0, df = 1, p = 0.9999), mean ± SD. Asterisks show Holm-corrected unpaired two-sided t-tests. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

D) Basipetal/rootward transport of radioactive ^{3}H-IAA in decapitated hypocotyls is defective in *abp1* mutants (one-way ANOVA: $F_{3,8}$ = 60.13, p = 7.86e-06). 10 μM NPA, an inhibitor of auxin transport, was used as a negative control. Single experiment, n = 3 for each condition, mean ± SD. Asterisks show Holm-corrected unpaired two-sided t-tests. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

(E) Basipetal/shootward transport of radioactive ^{3}H-IAA in roots is defective in *tmk1-1* and *abp1* mutants (genotype/treatment fixed effect: χ^2 = 32.30, df = 4, p = 1.66e-06). Pooled data from two independent experiments with similar results (repetition random effect: ΔAIC = 1.1, df = 1, p = 0.0804), mean ± SD. Asterisks show Holm-corrected unpaired two-sided t-tests. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

(F) Complementation of defects in root basipetal/shootward transport of radioactive ^{3}H-IAA for the *TMK1::TMK1-FLAG/tmk1-1* (*comp-tmk1-1*) and *ABP1::GFP-ABP1/abp1-C1* (*comp-abp1-C1*) lines. Single experiment, n = 10 for each condition, mean ± SD.

A

	WT					tmk1-1					Time (min),
	0	15	30	60	120	0	15	30	60	120	5 nM IAA

α-TMK1 — -130 kDa

α-ROP6 — -25 kDa

α-AHA2 — -100 kDa

B

	WT		tmk1-1		abp1-C1		abp1-TD1		100 nM IAA
	−	+	−	+	−	+	−	+	10 min

Pull-down:
α-ROP2   -25 kDa
                                         *empty well

	WT		tmk1-1		abp1-C1		abp1-TD1		100 nM IAA
	−	+	−	+	−	+	−	+	10 min

Input:
α-ROP2   -25 kDa

C

TMK1::TMK1-FLAG/
tmk1-1

DMSO   10   nM IAA, 1 h

Phos-tag
Biotin   -130 kDa
                          ◄ TMK1-P

Total
α-Flag   -130 kDa
                          ◄ TMK1

E

D

TMK1::TMK1-GFP

DMSO   5 nM IAA   10 nM IAA

3 h   3 h   3 h

0 ▬▬▬ 256

Figure 2.

Figure 2. Auxin-induced activation of cell surface ABP1-TMK1 signalling in roots.

(A) Western Blot: time course of endogenous TMK1 and ROP6 protein levels in the microsomal fraction of 4-day-old WT and *tmk1-1* roots treated with 5 nM IAA (see Figure S2E for quantification). The experiment was repeated 3 times and representative data are shown. The loading control is reused in Figure 4A which depicts more data from the same experiment.

(B) Auxin-mediated activation of ROP signalling is defective in *tmk1-1* and *abp1* mutant backgrounds. Pull-down of active ROP GTPases from root protein extracts using a heterologously expressed GST-tagged CRIB domain from the RIC1 ROP effector. Both input and pull-down gels were blotted with an anti-ROP2 antibody.

(C) Comparison of TMK1-FLAG phosphorylation in untreated roots and roots treated with 10 nM IAA for 1h. Root *TMK1::TMK1-FLAG/tmk1-1* extracts treated with 10 nM IAA or DMSO control for 1 hour were used for immunoprecipitation with anti-Flag antibody. Phos-tag Biotin Probe was used to analyse TMK1 phosphorylation levels and anti-Flag antibody for total amount of immunoprecipitated TMK1-FLAG (see figure S2C for quantification).

(D) Representative intensity-coloured images of *TMK1::TMK1-GFP* expression in 4-day-old seedlings after a 3-hour treatment with 5 or 10 nM IAA. Scale bars represent 20 μm.

(E) Time-course quantification of *TMK1::TMK1-GFP* signal in root tips treated with IAA (5 or 10 nM) or DMSO control, normalized to the initial fluorescence value. Mean ± SD (n = 20; experiment repeated 3 times with representative data depicted).

Figure 3.

Figure 3. Auxin-mediated TMK1 interaction with and phosphorylation of PIN2.

(A) Auxin promotes the interaction between TMK1 and PIN2 in a co-immunoprecipitation (Co-IP) assay. Microsomal protein fraction from 20 nM IAA- or DMSO-treated roots expressing *TMK1::TMK1-FLAG/tmk1-1* was immunoprecipitated with an anti-FLAG antibody, and endogenous PIN2 was detected by blotting with an anti-PIN2 antibody. See Figure S3A for Co-IP with 5 nM IAA where input PIN2 abundance in the protein microsomal fraction is increased.

(B) FRET-FLIM analysis on transiently expressed *35S::TMK1-GFP, 35S::TMK1^{ΔKD}-GFP, 35S::FLS2-GFP* and *35S::PIN2-mCherry* in root protoplasts. Fluorescence lifetime values are displayed as a heat map (for additional images see Figure S3B). Two-way ANOVA with post hoc tests (*$p < 0.05$; ****$p < 0.0001$).

(C) *In vitro* kinase assay showing that TMK1-3xHA directly phosphorylates the 6His-PIN2 hydrophilic loop (HL). Top panel, ^{32}P autoradiography; middle panel, CBB staining; bottom panel, immunoblot with an anti-HA antibody.

A

WT   tmk1-1

Time (min),
5 nM IAA

0 15 30 60 120   0 15 30 60 120

α-PIN2   -70 kDa

α-AHA2   -100 kDa

B

PIN2::PIN2-GFP

WT   tmk1-1

DMSO   10 nM IAA   DMSO   10 nM IAA

3 h   3 h   3 h   3 h

0   256

C

Ratio fluorescence intensity T/T0

WT DMSO   tmk1-1 DMSO
WT 5 nM IAA   tmk1-1 5 nM IAA
WT 10 nM IAA   tmk1-110 nM IAA

Time (h)

Figure 4.

Figure 4. TMK1 dynamically regulates root PIN2 levels in response to exogenous auxin.

(A) Western Blot: time course of endogenous PIN2 protein levels in the microsomal fraction of 4-day-old WT and *tmk1-1* roots treated with 5 nM IAA (see Figure S4A for quantification). The experiment was repeated 3 times and representative data are shown. The loading control (reused) and the anti-PIN2 immunoblot come from the same experiment as that shown in Figure 2A.

(B) Representative intensity coloured images of *PIN2::PIN2-GFP* after a 3-hour treatment with 10 nM IAA. Scale bars, 20 μm.

(C) Quantification of *PIN2::PIN2-GFP* in root tips of WT or *tmk1-1* seedlings treated with IAA (5, 10 nM or DMSO control), normalized to the initial fluorescence value. Mean ± SD (n = 20; experiment repeated 3 times with representative data depicted).

Figure 5.

Figure 5. TMK1 kinase activity is required for the gravitropic response.

(A) Representative images of root bending 3 hours after a 90° rotation for 4-day-old WT, mutant *tmk1-1*, kinase-dead *UBQ10::TMK1^{K616R}-mCherry,* and genetically complemented *TMK1::TMK1-FLAG/tmk1-1* seedlings.

(B) Time course of the gravitropic response for lines shown in (A) measured every 30 minutes, mean ± SD (n = 10; experiment repeated 3 times, a representative run is depicted). See Figure S5E for complete analysis and root growth rate data, and Table S2 for statistical analysis.

(C) Representative confocal images of asymmetric auxin response (visualized by *DR5rev::GFP*) at the lower side of the root before and after a 3-hour gravistimulation in WT and *tmk1-1* 4-day-old roots.

(D) Quantification of *DR5rev::GFP* asymmetry as a ratio of fluorescence intensity at the lower side to the upper side of gravistimulated roots at the indicated time points, normalized to the initial fluorescence value. Mean ± SEM of at least 3 independent experiments, n = 10. Two-way ANOVA with Dunnett's multiple comparison tests (*$p < 0.05$, **$p < 0.01$). 'g' represents the gravity vector.

A

	WT						tmk1-1					
α-TMK1												-130 kDa
α-ROP6												-25 kDa
α-PIN2												-70 kDa
α-AHA2												-100 kDa

Time (h): 0 1 2 3 4   0 1 2 3 4

B

TMK1::TMK1-FLAG/
tmk1-1

Time (h): 0 1 2 3

Phos-tag Biotin — -130 kDa ◄TMK1-P

Total α-Flag — -130 kDa ◄TMK1

C

WT

tmk1-1

0 — 256

D

PIN2::PIN2-GFP WT
■ Lower   ○ Upper

PIN2::PIN2-GFP *tmk1-1*
■ Lower   ○ Upper

Ratio fluorescence intensity T/T0

Time after gravistimulation (h)

E

0 h   3 h

— *TMK1::TMK1-GFP*

0 — 256

F

TMK1::TMK1-GFP
■ Lower   ○ Upper

Ratio fluorescence intensity T/T0

Time after gravistimulation (h)

Figure 6.

Figure 6. TMK1 mediates PIN2 asymmetry during the root gravitropic response.

(A) Western Blot: time course of endogenous TMK1, ROP6 and PIN2 protein levels in the microsomal fraction of 4-day-old WT and *tmk1-1* roots after 0, 1, 2, 3 and 4 hours of gravistimulation. The experiment was repeated 3 times and representative data are shown. For quantification of protein levels see Figure S6C.

(B) TMK1-FLAG phosphorylation after gravistimulation. *TMK1::TMK1-FLAG/tmk1-1* was immunoprecipitated with an anti-FLAG antibody and detected using a Phos-tag Biotin Probe (see Figure S6D for quantification).

(C) Representative images of *PIN2::PIN2-GFP* in 4-day-old WT and *tmk1-1* seedlings after 4 hours of gravistimulation within a time course of 0, 2, 3, 4 and 5 hours of gravistimulation. 'g' represents the gravity vector. Scale bars, 20 μm.

(D) Quantification of PIN2-GFP ratio between the lower and upper sides of gravistimulated roots, normalized to the initial fluorescence value at the indicated time point. Mean ± SD (n = 8; experiment repeated 3 times with representative data depicted). Significant differences to timepoint 0 are shown by *p*-values, two-way ANOVA with Dunnett's post hoc tests (**** *p < 0.0001*).

(E) Intensity-coloured map of 4-day-old *TMK1::TMK1-GFP* root tips before and after 3 hours of a 90° gravitropic stimulus. White arrows indicate asymmetric TMK1-GFP distribution. Scale bars, 20 μm.

(F) Quantification of TMK1-GFP in the epidermis and cortex of the upper and lower root sides normalized to the initial fluorescence value, mean ± SD (n = 10). Experiment repeated 3 times with representative data depicted. Significant differences to timepoint 0 are shown by *p*-values, two-way ANOVA with Dunnett's post hoc tests (**** *p < 0.0001*).

Figure S1.

Figure S1. Expression of the TMK leucine-rich repeat receptor-like (LRR-RLK) kinase family in roots.

(A) Relative transcription levels of *TMK1*, *TMK2*, *TMK3* and *TMK4* in *Arabidopsis* roots from 803 samples assayed with the Affymetrix Arabidopsis ATH1 Genome Array. Data was obtained from the Genevestigator database (https://genevestigator.com). Mean ± SD.

(B) Relative expression levels of *TMK1, TMK2, TMK3* and *TMK4* in different root cell types obtained from high-resolution spatiotemporal microarray analysis of 5–6-day old roots reported in [58] (ePlant http://bar.utoronto.ca/eplant).

(C) Root tip expression pattern of *TMK1*, *TMK2*, *TMK3*, and *TMK4* promoters fused to the GUS transcriptional reporter. Scale bars, 100 μm.

(D) Representative intensity-colored image of *TMK1::TMK1-GFP* expression in 4-day-old root tips.

Figure S2.

Figure S2. Auxin promotes TMK1 stability, phosphorylation state, and downstream signalling.

(A) Western blotting of total root protein extracts of 4-day-old *TMK1::TMK1-FLAG/tmk1-1* and *UBQ10::TMK1-3xHA* transgenic plants after 1-hour treatment with different concentrations of IAA (5, 10 and 100 nM IAA) or DMSO control. Ponceau staining was used as a loading control.

(B) Corresponding quantification of TMK1 protein levels normalized to Ponceau, and subsequently to DMSO condition, mean ±SD from 3 independent experiments. One-way ANOVA with Dunnett's multiple comparisons relative to DMSO (****$p<0.0001$).

(C) Quantification of TMK1-FLAG phosphorylated upon auxin treatment (10 nM, 1 hour) (see Figure 2C). Phosphorylation levels detected with a Phos-tag Biotin Probe were normalized to signal obtained from anti-FLAG detection, and subsequently to the initial value from time point 0. Mean ± SD of 3 independent biological replicates. *p*-value for comparison to DMSO was calculated with a two-tailed unpaired t-test (**$p < 0.01$).

(D) RT-qPCR time course of relative *TMK1* expression after 10 or 100 nM IAA treatment. Normalization was done to *PP2A* as a reference gene. The expression level at the start of the experiment (T0) was set to a relative value of 1 and statistical comparisons were made to this expression level (*$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$).

(E) Time course of TMK1 and ROP6 protein levels in the membrane fraction of 4-day-old WT or *tmk1-1* roots upon auxin treatment (5 nM). Plasma membrane AHA2 H$^+$ ATPase (immunoblot shown in Figures 2A and 4A) was used as a loading control. We then normalized to initial protein levels at time point 0. Values are mean ± SD from 3 independent experiments. Two-way ANOVA with Dunnett's multiple comparisons (****$p < 0.0001$).

Figure S3.

Figure S3. TMK1 interacts with PIN2 in an auxin-dependent manner and phosphorylates it.

(A) Co-IP assay showing the interaction of TMK1 with PIN2 upon low auxin treatment (5 nM IAA). Microsomal protein fraction from WT was used as a control for unspecific binding of endogenous PIN2 (shown in Figure 3A, same experiment).

(B) FRET-FLIM analysis on transiently expressed *35S::TMK1$^{\Delta KD}$-GFP, 35S::TMK1$^{\Delta KD}$-GFP/35S::PIN2-mCherry, 35S::FLS2-GFP* and *35S::FLS2-GFP/35S::PIN2-mCherry* in root protoplasts (the same experiment is shown in Figure 3B). GFP fluorescence lifetime was calculated as described in the Methods section. The heat map depicts fluorescence lifetime values (see Figure 3B for lifetime analysis).

(C) *In vitro* kinase assay showing phosphorylation of the 6His-PIN2 HL substrate by TMK1-3xHA in the presence or absence of auxin. Top panel, ^{32}P-autoradiography; middle panel, CBB staining; bottom panel, immunoblot with an anti-HA antibody.

Figure S4.

Figure S4. TMK1 regulates PIN2 stability on the post-transcriptional level.

(A) Time course of PIN2 protein levels in the membrane fraction of 4-day-old WT or *tmk1-1* roots upon auxin treatment (5 nM). Plasma membrane AHA2 H$^+$-ATPase (immunoblot shown in Figures 2A and 4A) was used as a loading control. We then normalized to initial protein levels at time point 0. Values are mean ± SD from 3 independent experiments. Two-way ANOVA with Dunnett's multiple comparisons (****$p < 0.0001$).

(B) Left panel: immunoblot of endogenous TMK1 and PIN2 in the microsomal fraction in response to 1-hour treatment with 10 nM IAA in WT versus *tmk1-1* roots. Right panel: quantification of protein levels detected with anti-TMK1 and anti-PIN2 antibodies normalized to the loading control, and then to DMSO. Values are mean ± SD from 3 independent experiments. Two-way ANOVA with Dunnett's multiple comparisons (****$p<0.0001$).

(C) Comparison of PIN2 expression in WT versus *tmk1-1* upon auxin treatment. Relative expression of PIN2 upon 10 and 100 nM IAA treatment at different time points (0, 5, 60, 120 and 180 minutes). Normalization was done to *PP2A* as a reference gene. The expression level at the start of the experiment was set to the relative value of 1.

Figure S5.

Figure S5. *TMK* genes are required for robust gravitropic root bending.

(A) Expression levels of *TMK1, TMK2, TMK3* and *TMK4* determined by RT-qPCR in root samples of 4-day-old WT and *tmk1-1, tmk2-1, tmk3-1* and *tmk4-1* seedlings. RNA was extracted in 4 independent biological repeats and the RT-qPCR analysis was performed in 3 technical repetitions. Expression levels were normalized against the expression level of PP2A as a housekeeping gene. Asterisks indicate significantly lower expression levels in the respective mutants, two-tailed unpaired t-tests (****$p <$ 0.0001 and **$p < 0.01$).

(B) Root gravitropism in single mutants of TMK family genes. Root curvature angles were measured every 30 min after reorientation by a 90° rotation. Representative images of 4-day-old WT, *tmk1-1, tmk2-1, tmk3-1,* and *tmk4-1* single mutant roots after 3 hours of reorientation.

(C) Time course of root tip angle changes and root growth rates, mean ± SEM from 3 independent experiments (n= 30-40). See Table S1 for statistical analysis.

(D) Representative images of root bending 3 hours after a 90° rotation of 4-day-old WT and the overexpression line of TMK1 without a cytoplasmic kinase domain (*35S::TMK1^{ΔKD}-GFP*).

(E) Time course of the gravitropic response including *35S::TMK1^{ΔKD}-GFP* lines and root growth rate from the same experiment at the indicated time points, mean ± SD (n = 10; experiment repeated 3 times with representative data depicted).

(F) Representative images of primary roots expressing the *UBQ10::TMK1^{K616R}-mCherry* and *35S::TMK1^{ΔKD}-GFP* constructs.

Figure S6.

Figure S6. Gravistimulation promotes TMK1 stability, phosphorylation state, and downstream signalling

(A) Western blotting of total root protein extracts of 4-day-old *TMK1::TMK1-GFP, TMK1::TMK1-FLAG/tmk1-1* and *UBQ10::TMK1-3xHA* transgenic plants after 0, 1, 2, 3h of gravistimulation. Ponceau staining was used as a loading control.

(B) Corresponding quantification of the relative TMK1 protein levels normalized first to Ponceau, and then to initial protein levels at time point 0, values are mean ± SD from 3 independent experiments. One-way ANOVA with Dunnett's multiple comparisons to timepoint 0 (****$p < 0.0001$).

(C) Comparison of TMK1, PIN2, and ROP6 endogenous levels in the membrane fraction during gravistimulation of WT and *tmk1-1* 4-day-old roots. Normalization was to plasma membrane AHA2 ATPase (immunoblot analysis shown in Figure 6A), and then to initial protein levels at timepoint 0. Values are mean ± SD from 3 independent experiments. Two-way ANOVA with Dunnett's multiple comparisons to timepoint 0 (****$p < 0.0001$).

(D) Quantification of phosphorylated TMK1 levels during gravistimulation (see Figure 6B). TMK1 protein levels detected with biotinylated Phos-tag Probe were normalized to the signal obtained from anti-FLAG detection, and then to the initial value from timepoint 0. Values are mean ± SD from 3 experiments with 3 independent biological replicates. Two-way ANOVA with Dunnett's multiple comparisons to timepoint 0 (****$p < 0.0001$).