The lncRNA landscape of cardiac resident macrophages and identification of *Schlafenlnc* as a regulator of macrophage migratory function

Anne Dueck[1,2], Lara Althaus[1,2], Kathrin Heise[1,2], Dena Esfandyari[1,2], Seren Baygün[3], Ralf P. Brandes[4,5], Julien Gagneur[2,6,7,8], Nicolas Jaé[5,9], Percy Knolle[10], Matthias S. Leisegang[4,5], Lars Maegdefessel[2,11], Thomas Meitinger[2,7], Niklas Petzold[1,2], Deepak Ramanujam[1,2], Hendrik Sager[2,12], Christian Schulz[2,13], Evangelos Theodorakis[2,6,7,8], Anna Uzonyi[1], Tobias Weinberger[2,13], Michael Bader[14], Marc Schmidt-Supprian[3], Stefan Engelhardt[1,2]

[1] Institute of Pharmacology and Toxicology, Technical University Munich, Munich, Germany
[2] DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, Munich, Germany
[3] Institute of Experimental Hematology, Central Institute for Translation Cancer Research, School of Medicine, Technical University Munich, Munich, Germany
[4] Institute for Cardiovascular Physiology, Goethe University, Frankfurt, Germany
[5] DZHK (German Centre for Cardiovascular Research), partner site RheinMain, Frankfurt, Germany
[6] Institute of Informatics, Technical University Munich, Munich, Germany
[7] Institute of Human Genetics, Klinikum rechts der Isar, School of Medicine, Technical University of Munich, Munich, Germany
[8] Computational Health Center, Helmholtz Center Munich, Neuherberg, Germany
[9] Institute for Cardiovascular Regeneration, Goethe University Frankfurt, Frankfurt am Main, Germany
[10] Institute for Molecular Immunology, MRI, Technical University Munich, Munich, Germany
[11] Department of Vascular and Endovascular Surgery, Technical University Munich, Munich, Germany
[12] Department of Cardiology, German Heart Centre Munich, Technical University Munich, Munich, Germany
[13] Medizinische Klinik und Poliklinik I, Klinikum der Universität München, Ludwig-Maximilians-Universität, Munich, Germany
[14] Max-Delbrück-Center for Molecular Medicine (MDC), Berlin, Germany

Present address: Anna Uzonyi, Department of Molecular Genetics, Weizmann Institute of Science, Rehovot, Israel

Keywords: long non-coding RNA, cardiac resident macrophages, pressure overload, *Schlafenlnc*, single cell sequencing

ABSTRACT

Cardiac resident macrophages (crMPs) were recently shown to exert pivotal functions in cardiac homeostasis and disease, but the underlying molecular mechanisms are largely unclear. Long non-coding RNAs (lncRNAs) are increasingly recognized as important regulatory molecules in a number of cell types, but neither the identity nor the molecular mechanisms of lncRNAs in crMPs are known. Here, we have employed deep RNA-seq and single cell RNA sequencing to resolve the crMP lncRNA landscape from healthy and diseased murine myocardium. CrMPs express previously unknown and highly cell type-specific lncRNAs, among which one lncRNA, termed *Schlafenlnc*, was particularly abundant and enriched in crMPs. We found *Schlafenlnc* to be necessary for migration-associated gene expression in macrophages *in vitro* and *in vivo* and essential for their adhesion and migration. Collectively, our data provide a basis to the systematic characterization of lncRNAs in crMPs and establish *Schlafenlnc* as a critical regulator of macrophage migratory functions.

INTRODUCTION

Long non-coding RNAs (lncRNAs) are the largest group of non-coding RNA molecules, with 14,764 lncRNA genes currently annotated in mouse and 19,933 in humans (Gencode in October 2022). Although only a small fraction of these has been functionally characterized, there is strong evidence that lncRNAs, through a variety of mechanisms, adopt critical roles in disease, for example in cancer [1], metabolic, neurologic [2] and cardiovascular disease [3]. Macrophage lncRNAs in cardiovascular disease have been primarily characterized in the vasculature and within the context of atherosclerosis, e.g., formation of foam cells or polarization (reviewed in [4]). With regard to cardiac disease, the analysis of lncRNAs and their functions has largely focused on cardiomyocytes [5], endothelial cells [6] and fibroblasts [7] (reviewed in [3]). By contrast, very little is known about lncRNAs in cardiac resident macrophages (crMPs), although this cell type constitutes the largest population of immune cells in the myocardium during homeostasis and also determines cardiac disease processes (reviewed in [8]). CrMPs have in recent years been characterized with respect to their developmental origin [9,10], their composition [10,11] and their putative functions such as roles in cardiac conduction [12,13], hypertrophy [14] or lymph vessel growth [15]. Although it can be readily assumed that many of these functions are subject to lncRNA-mediated regulation, a comprehensive description of the lncRNome from crMPs or functional data on a crMP-specific lncRNA is missing thus far.

In this study, we established the first comprehensive analysis of the lncRNA portfolios of the major cardiac cell types in health and disease, with particular emphasis on macrophages. In mouse myocardium, we identified lncRNAs that are specific for, and define, this cell type. One crMP lncRNA that has previously not been analyzed and which we termed *Schlafenlnc*, stands out for its strong enrichment and high abundance in crMPs. Upon functional characterization of a *Schlafenlnc* knockout in a mouse model for left ventricular pressure overload, we found that this lncRNA regulates macrophage abundance in the heart and the ability of this cell type to exert chemotactic migration. We therefore propose that *Schlafenlnc* critically regulates the function of cardiac macrophages in disease.

RESULTS

lncRNAs are important determinants for the definition of specific cell types

To determine the long non-coding RNA (lncRNA) repertoire in the major cardiac cell types and its changes under disease conditions, we subjected mice to chronic left ventrivcular pressure overload (induced by transverse aortic constriction, TAC), an established model of cardiac inflammation that leads into hypertrophic and fibrotioc remodelling of the heart. At day 6 after surgery, we sacrificed the mice, isolated primary cells by Langendorff digestion [16], and performed droplet-based single cell RNA sequencing (scRNA-seq, Chromium platform) (Fig. 1A). This experimental approach analyzes all cardiac non-myocytes, while the number of cardiac myocytes is decreased due to size constraints of the method. Unsupervised clustering of both samples based on the expression of both coding and non-coding RNAs with the Seurat analysis tool (Seurat V3.2 [17], Gencode vM22 [18]) yielded a total of 20,010 cells after quality control. These comprised all major non-myocyte cardiac cell types (Seurat clustering resolution 0.3, Fig. 1B), including subpopulations of immune cells, fibroblasts, endothelial cells and mural cells (Fig. 1B). Differential gene expression analysis from these cell types revealed distinct sets of protein-coding genes that define each major cardiac cell type as individual clusters (Extended Data Fig. S1), in line with previous reports [19,20].

Despite the generally lower expression levels of lncRNAs in comparison to protein-coding genes [21–24], we were able to readily detect over 440 lncRNAs. Several of the most strongly expressed lncRNAs exhibited a very high level of cell type specificity (Fig. 1C and 1D), suggesting that cardiac cell types can be defined based on their lncRNA transcriptomes. As Fig. 1D shows, our data not only confirmed the previously reported expression of *Meg3* in fibroblasts [25], *H19* in endothelial cells [26] and *Mhrt* in cardiac myocytes [27], but also revealed lncRNAs that had hitherto not been reported to be cell type-specific in the heart. These lncRNAs may reasonably be presumed to potentially exert important roles in their specific cell type, prompting for functional characterization.

To validate the contribution of lncRNAs to cell type identity, we performed differential clustering of the scRNA-seq data sets with or without lncRNA genes, and tested for clustering precision. Independent of whether the top 3,000 variable genes defined by the Seurat pipeline, all annotated genes (n=55,487) or expressed genes (n=19,719, expression value >0) were used to perform clustering, the inclusion of lncRNA marker genes significantly improved clustering compared to protein coding genes alone (Fig. 1E-G). Compared to clustering based on protein-coding genes alone, this resulted in an overall increased sensitivity (indicated by an increased adjusted Rand Index value) and yielded additional subclusters of cell types. Specifically, the detection of clusters representing venous capillary endothelial cells, pericytes, Fabp4$^+$ macrophages and Notch2$^+$ fibroblasts was dependent on the inclusion of lncRNAs (resolution 0.65) (Fig. 1H, Extended Data Fig. S1B).

In conclusion, lncRNAs were shown to be a valuable determinant of defining clusters and fine-tuning cell type assignations.

As a means to approach lncRNAs of potential disease relevance, we next thought to assess the expression of lncRNAs during cardiac disease by comparing the datasets obtained from sham-treated vs. TAC-treated mice. Upon increasing the clustering resolution from 0.3 to 0.7, a more detailed and granular cell assignment of lncRNA genes to cellular subtypes was achieved (Extended Data Fig. S2A

and S2B). With this setup, we performed differential gene expression analysis for the sub-populations
of the major cell types in the heart, and subset the resulting list to lncRNAs (Fig. 2A-C, Extended Data
Fig. S2C). Some lncRNAs were regulated after TAC throughout all subpopulations of a cell type, e.g.,
Gm29686 in fibroblasts (Fig. 2A), whereas others were only significantly regulated within a defined
subpopulation, e.g., Gm49785 in the fibroblast subpopulation (FB6) (Fig. 2A). This subpopulation of
fibroblasts represents activated myofibroblasts (high expression of periostin), a cell type involved in
extracellular matrix secretion and immune cell recruitment [28]. Similarly, several lncRNAs displayed
dysregulation within cardiac endothelial cells (Fig. 2B) and cardiac macrophages (Fig. 2C), resolved to
the level of cellular subtypes.

Of particular interest are lncRNAs in macrophages of the myocardium, since no lncRNA had thus far
been identified or functionally characterized in this cardiac cell type. As a first approach to the potential
function of the lncRNAs we had identified as deregulated in crMPs in the TAC model, we correlated
these lncRNAs with biological processes of their overlapping neighboring genes (Extended Data Fig.
S2D). Since lncRNAs are often co-expressed and co-regulated with proximal genes in *cis* [29], we
speculated that the latter may give insight into the biological context within which these lncRNAs
function. For example, genes antisense to downregulated lncRNAs in the Mac1 cluster (Lyve1+
macrophages) after TAC are associated with programmed cell death processes and genes antisense
to downregulated lncRNAs in the Mac2 cluster (Cd74+ macrophages) after TAC were predominantly
linked to adhesion pathways (Extended Data Fig. S2D), suggesting that the reported de-regulated
lncRNAs might also be involved in the regulation of these processes.

Taken together, we report a comprehensive inventory of regulated lncRNA genes in the major non-
myocyte cell types of the heart during pressure overload. These lncRNA genes will be interesting
candidates to be studied in more detail.

Determination and annotation of the lncRNA landscape of cardiac resident macrophages

As the sequencing depth of current single cell sequencing methodology limits detection sensitivity, we
therefore complemented our scRNA-seq data by performing deep RNA-Seq on basal total murine
myocardium as well as crMPs that were isolated using well-established cell markers (Cd45, Cd64,
Cd11b and F4/80 [9,10]). These surface proteins collectively labelled essentially all crMPs in our scRNA-
seq data (Extended Data Fig. S3A), corroborating our purification strategy. Using a pipeline to identify
and quantify novel lncRNAs (Extended Data Fig. S3B), we identified 250 genes that had previously not
been annotated by Gencode vM22 [18]. The identified novel lncRNAs showed the same characteristics
as the known lncRNAs with respect to having no coding probability (true non-coding) and transcript
length (Extended Data Fig. S3C upper panel), but were more highly expressed in our data set than the
known lncRNA population (Extended Data Fig. S3C lower panel). RNA-seq from fractionation
experiments showed for known lncRNAs a distribution of 80% in the nucleus (and 20% to the
cytoplasm), the novel lncRNAs are slightly less nuclear in localization (70% nuclear, 30% cytoplasmic
(Extended Data Fig. S3D).
As a further approach to the function of the newly identified lncRNAs, we built on the observation that
the products of lncRNA genes and protein-coding genes which are in proximity of 5kb or less and in
antisense overlapping orientation interact in the same biological processes [30]. The distribution of the

interaction categories (e.g. "antisense overlapping") was comparable between novel and known
lncRNAs (Extended Data Fig. S3E). Notably, 15 novel lncRNA genes were found without any interaction,
categorizing them as novel long intergenic RNAs (lincRNA). For both known and novel lncRNAs we
found many well-expressed genes (> 5 transcripts per million reads, TPM) that are specific in cardiac
macrophages compared to total murine myocardium (containing myocytes, endothelial cells, fibroblasts,
and macrophages) (Fig. 3A).

Identification of lncRNA *Schlafenlnc* as an abundant and enriched macrophage lncRNA

In order to select lncRNA candidates for experimental validation, we further filtered the datasets from
scRNA-Seq data and RNA-Seq based on the following stringent criteria: low coding probability
according to CPAT score (true non-coding RNA < 44% coding probability), absolute expression > 25
TPM in macrophages, enrichment of $\log_2$FC >3 and FDR <0.05, and de-regulation in TAC (scRNA-seq
data) FC >0.25). This yielded 13 lncRNA candidates that are highly abundant, enriched and regulated
in cardiac macrophages (Fig. 3B).

We found one of these candidates, lncRNA *AI662270*, particularly interesting due to its genomic
association with a cluster of eight genes on chromosome 11 which encode members of the Schlafen
family of proteins (Fig. 3C). Schlafen proteins are found in human and mouse and known to regulate
important processes and functions such as immune cell development, invasion and migration [31].
AI662270, however, is a hitherto uncharacterized gene within this locus. The genomic location of
lncRNA *AI662270* in the Schlafen cluster led us to name the transcript *Schlafenlnc.*

The Schlafen locus of protein-coding genes is very similar between human and mouse in general,
yet the internal order of the proteins (and lncRNAs) is slightly different. A study by Ulitsky and colleagues
developed a pipeline to identify syntenic genes across multiple organisms, where lncRNAs are tested
on the relative location and orientation to orthologous protein-coding genes [32]. By comparing Gencode
annotations [18] of human (V30) and mouse (vM21) with their pipeline, the gene *ENSG00000267074* was
designated as the putative human *Schlafenlnc* homolog (Fig. 3C). It is expressed in various human cell
types of leukocyte origin (using the RNA-Get module Encode data sets [33], Extended Data Fig. S4A),
making it an attractive molecule for studying the observed phenotypes in human cells. Besides the
analysis of expression, we tested if the two lncRNAs share common potential binding partners using a
pretrained deep neural network (DNN) [34] on eCLIP-seq data [35]. Indeed, the mouse *Schlafenlnc* and
human *ENSG00000267074* contain three major binding modules of the same or similar proteins
(Extended Data Fig. S4B and S4C), supporting the hypothesis of conservation.

Murine *Schlafenlnc* was up-regulated after TAC in both macrophage populations, but even more
pronounced in Mac2 (MHCII-high, Cd74+) macrophages (Fig. 3D). Its strong expression made it readily
detectable by Northern blotting (Fig. 3E). Interrogation of the RNA-seq datasets from the nucleus and
the cytosol, as well as RNA *in situ* hybridization, showed that *Schlafenlnc* localizes to both compartments
(Fig. 3F and G, Extended Data Fig. S4D).

199
200
201

Schlafenlnc determines myocardial macrophage infiltration and polarization in response to pressure overload

To assess the phenotype of a macrophage-specific deficiency of *Schlafenlnc in vivo*, we generated a *Schlafenlnc*$^{-/-}$ mouse line by CRISPR/Cas9 methodology and examined these mice under basal and disease conditions. The entire gene body was targeted with specific sgRNAs, including the promoter region upstream of *Schlafenlnc* (determined by H3K4me3 signals in murine bone marrow-derived macrophages, ENCODE data set ENCSR000CFF[36]) to abrogate all transcription from that locus (Extended Data Fig. 5 A). Deletion was verified by PCR from genomic DNA (Extended Data Fig. 5 B). The early compensation stage of the TAC model is characterized by sterile inflammation and both an influx of monocytes as well as proliferation of resident macrophages. We therefore performed sham and TAC surgery (Fig. 4 A) and analyzed the heart phenotype and monocyte/macrophage numbers after 6 days (Fig. 4 B). The heart weight of *Schlafenlnc*-deficient mice was significantly reduced (Fig. 4 C) and heart function (comparing ejection fraction of sham and TAC mice) was significantly improved (Fig. 4 D). Furthermore, *Schlafenlnc*$^{-/-}$ hearts showed a substantial decrease in pressure overload-induced interstitial fibrosis (Fig. 4 E) and interestingly a significantly reduced monocyte/macrophage infiltration after TAC, as determined by immunostaining of left ventricular myocardial sections against the monocyte/macrophage marker Cd68 (Fig. 4 F).

To assess the composition of immune cells in more detail we next performed scRNA-seq of the CD45+ cell fraction isolated from the myocardium and blood (Fig. 5 A). As expected, macrophages are the dominant cell population within the heart whereas the blood contains more monocytes (Fig. 5 B). Activated macrophages and monocytes release signals in an auto- and paracrine fashion[19]. *Schlafenlnc*$^{-/-}$ mice showed decreased signaling between almost all macrophage and monocyte populations after 6 days of TAC (Fig. 5 C), with a significant decrease in inflammatory signaling pathways such as Osteopontin (Spp1) and Macrophage Inhibitory Factor (Mif) signaling (Fig. 5 D). Both Osteopontin and MIF are important factors in recruiting monocytes and other inflammatory cells from the blood after cardiac injury[37,38]. Performing RNA velocity analysis of cardiac leukocytes after TAC (Fig. 5 E) showed gene dynamics of *Schlafenlnc* WT Ly6C-high monocytes towards non-classical monocytes as well as a moderate dynamic from MHCII-high macrophages to Lyve1 macrophages. In *Schlafenlnc*$^{-/-}$, the dynamics between monocytes is abolished while transition dynamics from the monocyte-derived macrophages (MHCII high) to resident macrophages (Lyve1) is strongly increased. This analysis showed stronger polarization of cardiac macrophages towards the *Lyve1* macrophage population in the *Schlafenlnc*$^{-/-}$ heart which is generally believed to play a more anti-inflammatory role following cardiac damage.

Next, we asked which genes were changing globally in Lyve1$^+$ macrophages after TAC due to *Schlafenlnc* deletion. Among the genes that are de-regulated during TAC (WT TAC vs. sham), 64 genes were additionally strongly de-regulated in *Schlafenlnc*$^{-/-}$ vs. *Schlafenlnc* WT Lyve1$^+$ macrophages (|Log2FC|>0.5, FDR<0.05). 17 of these genes are classified as coding for secreted proteins (Fig. 5 F, categorization according to Human Protein Atlas[39]), for example, *Lgals3* (Galectin-3) and *Spp1* (Osteopontin), which are de-regulated in all macrophage populations (Extended Data Fig. 5 C). An

241 analysis of the associated biological processes of de-regulated genes in Lyve1[+] macrophages (Fig.5 F)
242 showed leukocyte chemotaxis and migration as the strongest impacted pathways (Fig. 5 G).

243 Taken together, *Schlafenlnc* deletion in immune cells of the heart altered cardiac macrophage response
244 to early pressure overload towards an anti-inflammatory phenotype and disease-associated phenotypes
245 such as fibrosis and heart weight were improved.

246 *Schlafenlnc* is required for macrophage adhesion and migration

To elucidate the function of *Schlafenlnc* in more detail and explore its potential role in migration
processes, we generated *Schlafenlnc*-deficient macrophages from Hoxb8-FL macrophage
progenitors[40,41] (Extended Data Fig. S6A). Gene deletion was validated by RNA-Seq (Extended Data
Fig. S6A, middle track). As a genomic deletion might have an impact on nearby genes due to a
mechanism in *cis* by the deleted lncRNA or due to the deletion of a regulatory element within the DNA,
the surrounding region of *Schlafenlnc* (0.5 Mbp in each direction) was analyzed for de-regulation
(Extended Data Fig. S6B). 9 of 64 genes were found to be moderately dysregulated (with |log2FC| > 1,
FDR < 0.05) in the vicinity of *Schlafenlnc* revealing that there was not a general activation or de-
activation of that locus by *Schlafenlnc*. To identify major perturbations on gene expression level,
Schlafenlnc KO monoclonal cell lines (2 clones, each measured in n=4) were subjected to RNA-Seq.
These samples were compared to control monoclonal cell lines (two clones with n=3 each) that were
generated using an unrelated sgRNA (sequenced as wildtype at the *Schlafenlnc* locus). We observed
2,660 genes significantly de-regulated (|log2FC| > 1, FDR < 0.05) (Fig. 6 A), among those *Schlafenlnc*
that dropped from 67 TPM to 0.33 TPM. Among the downregulated genes, we observed an enrichment
of genes involved in migration and adhesion similarly to our *in vivo* study (Fig. 6 B, e.g., *Mmp14* and
Lgals3). Lower amounts of *Lgals3* could not only be observed in our RNA-Seq data but were validated
on protein level (Galectin-3 is the gene product of *Lgals3*) by measurement of cell culture supernatant
using an enzyme-linked immunosorbent assay (ELISA) (Fig. 6 C). More generally, gene ontology
analysis of all significantly de-regulated genes implicated pathways such as regulation of cytokine
production, regulation of cell adhesion and chemotaxis that are promoted by *Schlafenlnc* (Fig. 6 D). To
validate these results, we performed an adhesion assay by rigorous washing (Fig. 6 E) as well as tested
the chemotactic migration ability of *Schlafenlnc* KO macrophages with a gradient of the chemokine C5a
(Fig. 6 F). Adhesion was diminished by approximately 23% (Fig. 6 E) and chemotaxis was about 50%
reduced (Fig. 6 F) in *Schlafenlnc* KO cells compared to wild type controls.
Together these data obtained on *Schlafenlnc*-deficient macrophages corroborate a cell autonomous
role for this lncRNA in macrophages, in line with its specific expression in this cell type. *Schlafenlnc* is
required for the activation of macrophage gene programs underlying cell adherence and migration
providing a mechanistic basis for the phenotype observed in vivo for the *Schlafenlnc*-deficient mice.

DISCUSSION

This study provides the first comprehensive lncRNome of cardiac resident macrophages. One of the most highly enriched and abundant lncRNAs in cardiac resident macrophages – lncRNA *AI662270*, which we termed *Schlafenlnc* – was shown to be involved in adhesion and migration. As for *Schlafenlnc*, a particular characteristic of lncRNAs in general is their high degree of cell type-specificity[21–23] which makes them important for processes such as development, differentiation of cell types, cellular fate and function (reviewed in [3]). This characteristic could therefore make lncRNAs the natural choice when describing cell population expression signatures, especially in single cell RNA sequencing (scRNA-seq). scRNA-seq allows for an unbiased approach to identify main- and sub-populations of cells, yet little focus has been set to expressly include lncRNA marker genes. Initially, methodical problems such as library preparation, sequencing depth or analysis software hampered the detection of lncRNAs in such data sets. LncRNAs are still not as readily measured as mRNAs due to their on average lower expression levels, but together with improved scRNA-seq depth and improved annotations, our data suggest that lncRNAs are underestimated markers of cell type definition. Emerging platforms measuring RNA transcripts in tissue such as spatial transcriptomics or *in situ* sequencing (reviewed in [42]) might be able to leverage that potential further.

Focusing on lncRNAs and their potential regulation in ventricular myocardium during disease, we could identify an extensive set of regulated lncRNAs for all major cell populations. In general, we find more up-regulated than down-regulated genes upon induction of disease, which we believe to be due to the detection limit of scRNA-seq concerning very lowly expressed genes [43]. LncRNAs more specific to and regulated in a subset of cell types might offer new possibilities for therapeutic intervention and to gain insight into regulatory mechanisms specific for defined cell types. As we have shown for the three macrophage clusters (Mac1 (Lyve1+), Mac2 (Cd74+) and transitional macrophages), the different sets of regulated lncRNAs are implicated in distinct gene ontology pathways, which points to distinct functions of these macrophage populations.

Cardiac resident macrophages have been shown to be key players in diseases such as heart failure and myocardial infarction (reviewed in [44–47]). In the mouse model of left ventricular pressure overload, crMPs are proliferating during the compensation phase (from 0 days to about 7-14 days). An infiltration of monocyte-derived MPs starts early in disease and increases in the late, decompensated phase [48]. The infiltration of monocyte-derived MPs is typically associated with a long-term detrimental cardiac function [48–55]. This recruitment of monocytes is at least to some degree dependent on crMPs[56], and suppression of this communication axis might be beneficial. Galectin-3, for example, is known to be increased in heart failure patients [57] and is associated with murine cardiac remodeling during pressure-overload [58]. It is well known to be involved in the monocyte as well as macrophage migration and infiltration in cardiovascular diseases [59–61]. Our combination of unbiased scRNA-seq measurements and in-depth coverage by RNA-Seq allowed us to identify the hitherto uncharacterized lncRNA *Schlafenlnc* as a regulator of macrophage adhesion and migration. *Schlafenlnc* is up-regulated after 6 days of TAC as well as other genes involved in adhesion and migration such as Galectin-3 and its deletion impacts Galectin-3 expression both *in vivo* and *in vitro*. Our quantitative assessment of macrophage/monocyte numbers (Cd68 staining) and the analysis in relative immune cell composition by scRNA-seq after 6 days of TAC point to a mechanism of defective signaling by *Schlafenlnc* KO crMPs to initiate recruitment

of inflammatory cells such as monocytes. Given its selective expression in crMPs, we believe that
targeting of *Schlafenlnc* warrants further exploration as a potential anti-inflammatory therapeutic
strategy.

Similar to its orthologue in mice, human *Schlafenlnc* is specific for immune cell types (Extended Data
Fig. S4A). In support of conserved function in immune cells, our protein interactor analysis of both
murine and human *Schlafenlnc* using deep learning showed similar binding modules for RNA binding
proteins (Extended Data Fig. S4B and S4C). Further studies will be required to dissect molecular
interactors for *Schlafenlnc* in mouse and human.

Taken together, our study demonstrates that lncRNAs act as critical contributors to the
transcriptome-based analysis of cell clusters in line with their crucial role in development and cell fate
determination [3], leading to the definition of cell identities. The combination of single cell sequencing and
RNA-Seq approaches led us to the identification of multiple novel candidate lncRNAs that warrant
further analysis in cardiac homeostasis and disease. One of the most abundant and macrophage-
enriched lncRNAs, *Schlafenlnc*, acts as a crucial regulator of macrophage adhesion and migration.

335
336

METHODS

<u>Experimental animals, surgery conditions</u>

Surgeries on mice were performed according to the guidelines and regulations of the responsible
authorities. Consent was given by the institutional review board of the Regierung of Oberbayern (ROB-
55.2-2532.Vet_02-19-194).

Wild type male mice of C57BL/6N background were purchased from Jackson Laboratories and
submitted to transverse aortic constriction (TAC) or sham procedure as reported previously [16] at an age
of 8-10 weeks.

To produce *Schlafenlnc* -knockout mice, two sgRNAs (5' cut site: 5'- GAGGTTCCATCAGTCTGACA-
3', 3' cut site: 5'-ACATGTCTTTGGTAGGAGCA-3') targeting the flanking region of the *Schlafenlnc* gene
in the mouse genome were designed and synthesized by IDT (Coralville, IA, USA). A mixture of Cas9
protein (IDT) and sgRNAs was electroporated into C57BL/6 mouse zygotes. Genomic DNA of newborn
mice was amplified using the primers Slfnlnc1 (5'-TTCCTTCAGAGGTCCATCAG-3') and Slfnlnc4 (5'-
AGCCAGTCTAAAGACTCTGG-3'). The appearance of a 360 bp band identified founder mice with a
complete loss of the *Schlafenlnc* gene. The animals were bred to homozygosity and used for the
experiments.

Echocardiography was performed as reported before[16].

<u>Primary cell isolation from adult murine heart</u>

Hearts of C57BL/6N mice were harvested at an age of 8-10 weeks and subjected to digestion by
perfusion via the coronary arteries using Collagenase II (Worthington, Lakewood, NJ) as described
previously [62]. Cell suspension was processed according to subsequent protocol/experiment.

 Single cell cloning and sequencing

Primary cells from a sham [16] and an adult mouse heart 6d after TAC were washed three times with PBS
to remove cell debris and passed successively through 100µm, 70µm and 30µm filters. The single cell
library was generated using the Chromium Single Cell 3' GEM, Library & Gel Bead Kit v2 (10X
Genomics) with a target of 10,000 cells per sample. The single cell library was cloned according to the
manufacturer's instructions. Sequencing was performed on a HiSeq4000 (Illumina).

Primary cells from the hearts from *Schlafenlnc*-knockout and littermate wildtype mice in basal condition
as well as after 6d TAC were passed through a 100µm filter and centrifuged for 1min at 100*xg* to pellet
cardiomyocytes. The supernatant (non-myocytes) was passed through a 40µm filter and washed twice
in magnetic-activated cell sorting (MACS) buffer (PBS + 0.5% BSA + 2mM EDTA) to result in 200µl cell
suspension. For the isolation of PBMCs, blood was diluted with 5 ml of MACS buffer and layered onto
5ml of Ficoll-Paque (17144002, Cytiva), followed by centrifugation at 400xg and 4°C for 30min in a
swinging rotor without a brake. The white layer containing the cells was aspirated, transferred to a 50ml
tube and mixed with 30ml of MACS buffer. After centrifugation for 5 min at 300xg and 4°C, the pellet
was washed once more with 30ml MACS buffer (200xg and 4°C for 5min) and resuspended in 200µl
MACS buffer. Cardiac and blood leukocytes (basal condition) were further purified using MACS, labelled
with cell hashing antibodies and isolated by FACS (all steps at 4°C): Cell suspensions were incubated
with Fc-block (1:50, BD Pharmingen 553142) for 10min and subsequently incubated with mouse Cd45
MicroBeads (Miltenyi Biotec) for 15min. Cells were washed with MACS buffer and MACS was performed
using an AutoMACS (Miltenyi Biotec) running the program *possel_s*. After elution, pelleted cells were
resuspended in MACS buffer containing anti-Cd45-FITC (1:1000, clone 30-F11, Biolegend 103108), PI
(1:100) and cell hashing antibodies (1:1000, #1 for heart, #2 for blood, Biolegend) and incubated for
30min protected from light. For TAC samples, PBMCs were purified as above, cardiac non-myocytes
were directly used for cell sorting. Both cell fractions were stained with cell hashing antibodies (1:1000,
#2 for heart, #3 for blood, Biolegend) and PI (1:100) and incubated for 30min. After washing, live cells
of all samples were sorted on a Sony cell sorter SH800S, heart and blood cells of each genotype and
condition were pooled and run on a Chromium controller using the Chromium Next GEM Single Cell 3'
v3.1 Library & Gel Bead Kit (10X Genomics) with a target of 10,000 cells per pool. Libraries were cloned
according to the 10x Genomics user guide with feature barcode technology for cell surface protein
(RevD) and Biolegend's protocol for amplification of the cell hashing libraries. Libraries were pooled
according to the recommendations of 10x genomics protocol and sequenced on a NovaSeq6000
(Illumina).

Purification of primary cells and cardiac macrophages by F4/80 MACS and cell sorting for RNA-seq

For a total myocardial cell sample, 100µl of the raw cell suspension of primary cells from adult wildtype
mouse hearts were washed twice with PBS and lysed in Trizol (ThermoFisher) for subsequent RNA
preparation. For the purification of cardiac macrophages, the cell suspension was filtered through cell
sieves of 70µm and 40µm of size and centrifuged at 100 *x g* for 3min at 4°C to completely remove
cardiac myocytes. All following steps were performed on ice/at 4°C. The supernatant (non-myocytes)
was washed twice in magnetic-activated cell sorting (MACS) buffer (PBS + 0.5% biotin-free BSA + 2mM

EDTA). MACS was performed in a two-step process using anti-F4/80-biotin coupled antibody and

Streptavidin microbeads (Miltenyi Biotec) on LS columns. The eluted macrophage fraction was

resuspended in a final volume of 200µl MACS buffer and incubated 10 min with Fc block (1:50). 200µl

of staining solution containing flow cytometry antibodies (anti-Cd45-PECy7, anti-Cd64-PEDazzle594,

anti-Cd11b-PE) was added and incubated for 30min in the dark. Cells were washed twice with MACS

buffer and sorted on a Biorad S3 cell sorter. Cd45+/Cd64+/Cd11+ macrophages were collected

(between 100k-170K cells), washed with PBS and lysed in Trizol (Invitrogen) for RNA preparation.

<u>Fibrosis staining</u>

Paraffin sections (8 µm) of ventricular myocardium were stained with Sirius red and Fast green. Images

of the whole heart were taken with a 10x objective using a AxioObserver.Z1 (Zeiss, Jena, Germany)

motorized scanning-stage microscopy (130 x 85; Märzhäuser, Wetzlar, Germany). Fibrosis of the left

ventricular myocardium was quantified as the ratio of signals from Sirius red to Fast green in each

section using the Metamorph software (Molecular Devices).

<u>Cd68 staining</u>

Cryosections (5 µm) of ventricular myocardium were fixed with methanol for 10 min at -20°C, washed

thoroughly with tap water and treated with acetone for 1 min at -20°C. Sections were air dried and

subsequently blocked with 5% goat serum in PBS for 1 h. Sections were stained with anti-mouse Cd68

(1:100) (AbD Serotec, MCA 1957) at 4°C overnight. The sections were washed three times with PBS +

0.05 % Tween 20 and stained with anti-rat AF564 (1:100) and Cytox Green (1:1000). Images were

acquired by confocal microscopy using 63x glycerol objective (Leica SP5) and analyzed using

Metamorph (Molecular Devices).

<u>Generation of knockout macrophages</u>

Conditionally immortalized hematopoietic progenitor cells were generated from a R26-Cas9-eGFP

transgenic mouse line [41] as previously described [40]. For inducing CRISPR-Cas9 mediated knockout in

Hoxb8-FL cells, cells were electroporated with an sgRNA pair targeting the *Schlafenlnc* locus (IDT: Alt-

R® CRISPR-Cas9 sgRNA, 5' cut site 5'-GAGGTTCCATCAGTCTGACA-3', 3' cut site 5'-

GGCTTGGAAGTAGAAAACCG-3'), as control, a single sgRNA was used (5'-

ACCCCTTCAGCTTCACCTCC-3'). Transfected cells were single-cell cloned by limiting dilution assay

into 96-well plates and screened for knockout clones. CRISPR-directed deletions on *Schlafenlnc* were

identified by a PCR-based strategy to amplify the genomic DNA of the clones. Single-cell clones carrying

the induced deletions were expanded and frozen for further use. To do this, extraction of genomic DNA

was performed with QuickExtract™ DNA Extraction Solution (Lucigen) according to the manufacturer's

instruction. Briefly, approximately 100,000 cells were lysed in 50µl of QuickExtract by incubation as

follows: 65°C – 15min, 68°C – 15min, 98°C – 10min. 50ng of genomic DNA was then subjected to PCR

amplification with two primer pairs to distinguish successful *Schlafenlnc* knockout clones from wildtype

or heterozygous knockout. *Schlafenlnc*_KO_forward 5'-CAGCCCTGAGTCAGAACATAC-3' and

*Schlafenlnc*_KO_reverse 5'-TCCTTAGGGTGCACATCTACT-3' were used for detection of a

successfully cut site with a resulting amplicon length of 480bp, *Schlafenlnc*_WT_forward 5'-

TCTGTCTCTACTCCCTCAACTC-3' and *SchlafenInc*_WT_reverse 5'-AAGGCATGTTCTGCTCCTAC-

3' were used for detection of the wildtype allele/uncut gene locus with a resulting amplicon length of

320bp.

Hoxb8-FL *SchlafenInc* KO cell culture and differentiation

Hoxb8 progenitor cells were cultured in RPMI 1640 supplemented with 10% FCS, 1% Pen/Strep, 50

ng/mL FLT3L, 50nM b-mercaptoethanol and 1nM b-estradiol. FLT3L supernatant was obtained from a

FLT3L-producing B16 melanoma cell line.

For differentiation of HoxB8 cells into macrophages, Hoxb8 progenitor cells were cultured in

differentiation medium consisting of the culturing medium described above without b-estradiol and

supplemented with 20% MCSF supernatant, which was obtained from a MCSF-producing L929 cell line.

Differentiation was accomplished after 6 days.

Generation of bone marrow-derived macrophages

Femurs and tibias from 6-12 week old C57BL/6N mice were prepared and bone marrow was isolated

by flushing the bones with R10 medium (RPMI medium, 10% FCS, 1% antibiotics) with a syringe. To

generate bone marrow derived macrophages, 3-4 mio. isolated bone marrow cells were seeded in a

10cm petri dish containing R10 medium supplemented with 20% MCSF supernatant. Medium was

added on day 3 and exchanged on day 6 after seeding, macrophages were fully differentiated on day 8

and used for experiments.

Cytoplasmic/nuclear fractionation

Cytoplasmic/nuclear fractionation was performed as described before [63]. Briefly, bone marrow-derived

macrophages (BMDMs) were washed two times with cold PBS and harvested. 10% of the cells was

transferred into a new tube and used as a sample for the total gene expression. Cells were centrifuged

at 200 *x g* for 10 minutes and the total expression sample was directly used for RNA isolation. The

fractionation sample was resuspended in 200µl lysis buffer A (10 mM Tris pH=8, 140 mM NaCl, 1.5 mM

MgCl2, 0.5% Nonidet P-40) and incubated for 5 minutes on ice. The lysate was centrifuged at 1000 *x g*

for 3 minutes at 4°C. The supernatant containing the cytoplasmic fraction was directly used for RNA

isolation through lysis with Trizol. The sediment containing the nuclear fraction was washed two times

with lysis buffer A and one time with lysis buffer B (Buffer A supplemented with 1% Tween-40, 0.5%

deoxycholic acid) and afterwards resuspended in Trizol for RNA isolation.

RNA cloning and sequencing

Libraries from primary cardiac macrophages and total myocardium as well as *SchlafenInc* KO and

control cell lines were generated using the TruSeq Stranded RNA Kit (Illumina) according to the

manufacturer's protocol, except for one step. Instead of using the SuperScriptII (ThermoFisher) as the

reverse transcriptase, SuperScriptIII (ThermoFisher) was used at a reaction temperature of 50°C.

Sequencing was performed on a HiSeq4000 platform (Illumina).

Libraries from fractionation samples were generated using the Stranded mRNA Prep, Ligation Kit

(Illumina) according to the manufacturer's instructions. Sequencing was performed on a NovaSeq6000

machine (Illumina).

RNA *in situ* hybridization (RNAscope)

Bone marrow derived macrophages were seeded on a removable 8-well chambers slide (Ibidi) and RNA

in situ hybridization was performed using the RNAscope Fluorescent Multiplex Reagent Kit (Advanced

Cell Diagnostics (ACD)) according to manufacturer's instructions. A custom designed RNA probe by

ACD was used for detection of *Schlafenlnc* (ENSMUST00000143673.2). Following the RNA *in situ*

hybridization, the slides were incubated with Phalloidin-AF647 (Abcam) for 60 minutes at room

temperature. Subsequently, slides were washed three times with PBS and mounted using ProLong™

Gold Antifade Mountant (Invitrogen). Images were acquired using the confocal microscope Leica TCS

SP II and the Leica Application Suite.

Galectin-3 ELISA

Cell culture supernatant of differentiated Hoxb8 macrophages was collected and the amount of soluble

Galectin-3 was quantitatively assessed using the Galectin-3 Mouse SimpleStep enzyme-linked immuno-

sorbent assay (ELISA) kit (Abcam) according to the manufacturer's instructions. The cell culture

supernatant was diluted 1:10 in sample diluent before use.

Chemotaxis

An IncuCyte® ClearView 96-Well Chemotaxis Plate (Sartorius) was coated on both sides of the

membrane using 50 µg/mL Matrigel (Corning) at 37°C for 30 minutes. After coating, plates were

incubated at room temperature for 30 minutes and afterwards washed with D-PBS. Differentiated HoxB8

macrophages were harvested and 2,000 cells were seeded in each well in differentiation medium

containing 0.5% FCS. Cells were allowed to settle at the bottom of the upper well for 30 minutes.

Subsequently, 100nM recombinant murine C5a (Pepro Tech) was added to the lower well of the

ClearView migration plate. Cells were imaged every 2 h for 72 hours using an Incucyte ZOOM

instrument and a 10x objective with chemotaxis plate settings. Chemotaxis was analyzed using the

IncuCyte ZOOM software.

Adhesion Assay

Ibidi black 96 well µ-Plate was coated with 20µg/mL Fibronectin (Sigma Aldrich) for 1 h at 37°C. 10,000

differentiated Hoxb8 macrophages were seeded in each well in differentiation medium and were

incubated for 2 hours at 37°C in a 5% CO_2 atmosphere. After the incubation, cells were stained using

NucBlue Live ReadyProbes reagent (Invitrogen) and following the manufacturers' instructions. Images

of the Nuclei were taken before and after removing the non-adherent cells by washing three times with

pre-warmed medium using an EVOS FL Auto 2 cell imaging system (Invitrogen). Cell nuclei were

counted automatically using the MetaMorph analysis software (Molecular Devices).

Northern blotting

Northern blotting was performed as reported previously [64]. Briefly, 10µg total RNA from fibroblasts
(NIH/3T3) and in vitro differentiated macrophages were run on a 1.2% formaldehyde agarose gel,
transferred to a Hybond-N+ membrane (Ge Healthcare) and UV-crosslinked. Probes were end-labelled
using [gamma-^{32}P]-ATP and purified using an Illustra ProbeQuant G-50 micro column (GE Healthcare).
A mixture of two different probes was used per target. Probes were incubated over night with a pre-
hybridized membrane, signal was accumulated using a phophoimager screen and detected using a
Cyclone   Plus   (PerkinElmer).   mGAPDH   probe1   5'-
CATGGACTGTGGTCATGAGCCCTTCCACAATGCCAAAGTTG-3',   mGAPDH   probe2   5'-
GAATTTGCCGTGAGTGGAGTCATACTGGAACATGTAGACCATGTAGTTGAGG-3',   Slfnlnc   probe1
5'-CAGCTCTGCGTCTCTGCTCCTGTTCAGGTCCATCTTTCATCTCCG-3',   Slfnlnc   probe2   5'-
CAGTCGGGGTTCTGCCTCTTGATGTGGTTGTGCTAACCGGTGGGAC-3'.

Quantitative real-time PCR

1 µg of total RNA was reverse transcribed with Protoscript II reverse transcriptase (NEB), both according
to the manufacturer's instructions using random hexamer primers. Quantitative real-time PCR (qPCR)
was carried out with primers listed below, using the FastStart universal SYBR Green master mix
(Roche). The sample volume of 12.5 µl contained 1x SYBR Green master mix, 5 pmol of each primer
and 10 ng of cDNA. PCR was performed using StepOnePlus Real-Time PCR systems (Applied
Biosystems). Slfnlnc_qPCR fwd 5'- ATGACTGGTCCTGGTTTTCG -3', Slfnlnc_qPCR fwd 5'-
TCTGCCTCTTGATGTGGTTG   -3',   mRPL32   fwd   (housekeeping   gene)   5'-
ACATCGGTTATGGGAGCAAC   -3',   mRPL32   rev   (housekeeping   gene)   5'-
GGGATTGGTGACTCTGATGG -3'.

Data analysis

Single cell sequencing data analysis
Raw reads generated by single cell RNA sequencing data from sham[16] and 6d after TAC (this study)
was aligned against the mouse genome (mm10, Gencode annotation vM22 [18]) using CellRanger 5.0.0
(10x Genomics). Cell that were called by CellRanger were subsequently analyzed using the R package
Seurat v3.2 [65]. After quality control (retaining cells with >500 or <3,600 detected genes, <10%
mitochondrial cell content), 7,913 cells of the sham condition and 13,803 of the TAC condition were
used for further downstream analysis. SCTransform [66] was used to normalize and scale the data as well
as remove unwanted variation (regressing out mitochondrial gene content and cell cycle score).
SCTransform was then applied to identify variable feature which were used to integrate both data sets
("IntegrateData" function). By using the ElbowPlot function on Principal Component Analysis results, the
number of relevant components was determined (set to 40) and used for embedding and visualization
by Uniform Manifold Approximation and Projection (UMAP). Seurat's "FindAllMarkers" function was
used for determination of cell population marker genes.
For the comparison of clusterings with adjusted Rand index score, gene sets were defined as follows:
all genes = all genes listed in the vM22 annotation of Gencode; protein coding genes = all genes with

gene biotype "protein coding" in vM22; lncRNAs = all genes in vM22 lncRNA annotation of Gencode
(gene biotype "lncRNA"+"TEC"); "expressed" gene sets = same definitions as above with filter for
average expression >0 in at least one cell cluster; all variable genes = gene list generated by Seurat's
"SelectIntegrationFeatures", default is 3,000 genes; subset of variable genes based on gene lists
defined above. Next, clusterings with resolutions from 0.01 to 1.00 with 0.01 increments were iteratively
calculated ("FindClusters" function) and cluster ID per cell was saved in a data frame. The adjusted
Rand Index is a measure for the degree of similarity between two clusterings, for example. The R
package mclust [67] with inbuilt function "adjustedRandIndex" was used to compare all gene sets to the
respective complete gene set (e.g. the subset of expressed lncRNAs to all expressed genes).
For calculating de-regulated genes after TAC, first Adaptively thresholded Low-Rank Approximation
(ALRA) [68] was performed in order to impute drop out gene expression values followed by using Seurat's
"FindMarkers" function to compare each cell population before and after TAC.
For the analysis of scRNA-seq data of *Schlafenlnc* KO and WT animals, raw reads from basal and 6d
after TAC from WT and KO were aligned against the mouse genome (mm10, Gencode annotation
vM25[18]) using CellRanger 6.0.1 (10x Genomics) and analyzed using the R package Seurat v3.2 [65].
Quality control was applied to basal (>200 or <5,000 detected genes, <10% mitochondrial cell content)
and TAC (>200 or <4,000 detected genes, <10% mitochondrial cell content) samples. SCTransform and
FindAllMarkers was applied as above to normalize, scale, integrate data and identify marker genes for
cell populations. The integrated object was subset on all cell clusters expressing Cd45 (Leukocytes).
Deregulated genes were calculated using Seurat's "FindMarkers" function to compare each cell
population between *Schlafenlnc* KO and WT. GO Term analysis of the deregulated genes was
performed using "DAVID 2021" [69].

Cell-cell communication analysis
A subset of cells of interest from blood and heart of *Schlafenlnc* KO and WT CD45+ (*Lyve1*
macrophages, *MHCII*-high macrophages, *Isg* macrophages, intermediate macrophages, classical
monocytes and non-classical monocytes) was used for analysis of secreted signaling pathways within
the CellChat database. The *Schlafenlnc* KO and WT objects were analyzed separately using CellChat[70]
and merged with "mergeCellChat". The differential number of interactions between the cell clusters in
Schlafenlnc KO vs. WT was analyzed and displayed using the CellChat function
"netVisual_diffInteraction" and the differential signaling pathways were assessed using "rankNet".

Estimating RNA velocity
To calculate the RNA velocity of single cells, the abundance of the spliced and unspliced RNA was
quantified by running Velocyto CLI (v0.17.5) on CellRanger output BAM files and the corresponding
gene annotation (GENCODE vM22). The resulting loom files were incorporated into the Seurat objects
using SeuratWrapper R package (v0.3.0), and after integration and clustering, the objects were
converted to AnnData format (h5ad) to be compatible for downstream analysis with the Python package
scVelo (v0.2.5). Using scVelo, cell-to-cell transition probabilities were estimated and RNA-velocity was
projected as streamlines on the UMAP embedding.

Identification of novel lncRNAs

Most steps of this pipeline were performed using the Galaxy platform [71]. Reads were mapped using RNA STAR aligner (Galaxy version 2.6.0b-2) [72] using a built-in index based on Gencode vM22 annotation and otherwise default parameters. StringTie (Galaxy Version 1.3.3.2) [73] was used to predict novel transcripts (strand information RF, reference file Gencode vM22, average read length 90). StringTie merge (Galaxy version 1.3.6) with default parameters was used to merge all new annotations together, including Gencode annotation vM22 as reference annotation. Novel genes were extracted by Bedtools Intersect tool (Galaxy version v2.29.0) [74], intersecting the merged annotation with the Gencode vM22 annotation. Transcripts were extracted with the "Extract features" tool (Galaxy 1.0.0) and analyzed for coding potential with FEELnc (Galaxy version 0.1.1.1) [30]. FEELnc produces a gene transfer format (gtf) annotation file containing only transcripts classified as long non-coding RNAs. "cat" command was used to merge this lncRNA annotation to Gencode vM22 reference annotation file to create reference annotation file with additional novel lncRNA genes. All RNA-Seq libraries were measured again against this new transcriptome using StringTie's "count" option. For visualizing coding probability, the python version of the Coding-Potential Assessment Tool (CPAT, v2.0.0) [75] was run on the fasta sequences of the novel lncRNAs (\$ cpat.py -g novel_lncRNAs_gffread.fasta -d Mouse_logitModel_CPAT.RData -x Mouse_Hexamer_CPAT.tsv -o output_novel_lncRNAs.txt). GraphPad Prism 8 was used to calculate the frequency distribution in percentages. For plotting transcript lengths, the length determined in the CPAT output was used and again transformed to frequency distribution in Prism. For the categorization of interactions between lncRNAs and other genes, the classifier output of FEELnc was used. Interactions with the same gene were discarded, for known lncRNAs only genes expressed predominantly in cardiac macrophages were considered ($\log_2$ fold change >1). To compute differential gene expression analysis, DESeq2 package was used (Galaxy version 2.11.40.2) [76], outputting also a normalized counts file. Normalized counts served as input for clustering and heatmap visualization by ClustVis R package (v. 0.0.0.9) [77].

RNA-Seq analysis of Hoxb8 SchlafenInc KO macrophages

Fastq reads were mapped using RNA STAR aligner (Galaxy version 2.6.0b-2) [72] using a built-in index based on Gencode vM22 annotation and otherwise default parameters. StringTie (Galaxy Version 1.3.3.2) [73] count option was used to quantify gene expression using Gencode vM22 as reference annotation. DESeq2 was used to calculate differential gene expression (Galaxy version 2.11.40.2) [76]. The normalized counts file was used for heatmap visualization with ln(x) transformation of the top 20 down-regulated genes (ClustVis R package [77]).

Neighboring genes analysis and Gene Ontology Analysis

Up- or downregulated lncRNAs during pressure overload (p-value<0.05) in either macrophage, fibroblast or endothelial cell populations were analyzed for neighboring genes using FEELnc (Galaxy Version 0.2, [30]) and filtered for antisense protein coding genes. Gene ontology analysis for these gene sets for each cell population was performed using the web-based tool "DAVID 2021" [69]. Neighboring genes of known and novel lncRNAs were also identified using FEELnc (Galaxy Version 0.2, [30]).

Biological processes associated with the differentially regulated genes in *SchlafenInc* KO macrophages (log2FC > 1 or < -1, FDR < 0.05) were determined using ClueGO application (v2.5.8) [78]

ran within Cytoscape (v3.8.2) [79] network analysis tool (p-value cutoff = 0.01, Min GO level = 4, Max GO

Level = 10, Number of Genes = 2, Min. Percentage = 4.0).

Kipoi RBP-eclip analysis (Protein interactors analysis)

Model training data: To access the binding probability of known RBPs on *Schlafenlnc*, a pretrained deep

neural network (DNN) [34] on eCLIP-seq data [35] was downloaded from the Kipoi repository [80]. In order to

test the binding probability of the 101 proteins on *Schlafenlnc*, a bed file containing the coordinates of

Schlafenlnc's 4 exons (ENSMUST00000143673.2 on Ensembl) was obtained. Then for each exon, we

took sliding windows of 101nt length resulting on a new bed6 file where each row is of 101nt length.

Kipoi builds on environments, making the sharing and loading of models easy. We started our analysis

by generating an environment with all the needed dependencies. Afterwards for each protein of interest,

we programmatically obtained the dataloader, load the fasta sequence for GRCm38.p6, load the

GENCODE annotation for mouse version vM22 [18], and a bed file with the pre-extracted 101nt length

sliding windows and ran the predict function with default arguments (considering all the genomic

landmarks also for non-protein coding genes). Finally, for each protein of interest a csv file was

generated, containing the following info for each 101nt sequence considered: the chromosome, the

starting position, the end position, a sequence ID, a score as in the bed6 file, the strand and the binding

score [0 (no binding probability), 1 (high binding probability)] of the protein of interest on that window.

<u>Software list</u>

Leica Application Suite Advanced Fluorescence, version 2.7.9723

Invitrogen EVOS FL Auto 2 Software

CellRanger, version 5.0.0

Seurat, version 3.2

RNA STAR aligner, version 2.6.0b-2

StringTie amd StringTie merge, version 1.3.3.2 and 1.3.6

Bedtools Intersect tool, version v2.29.0

"Extract features" tool, version 1.0.0

FEELnc, version 0.1.1.1 (Galaxy) and version 0.2

Coding-Potential Assessment Tool, CPAT, version 2.0.0

DESeq2, version 2.11.40.2

ClustVis R package, version 0.0.0.9

Cytoscape, version 3.8.2

ClueGO, version 2.5.8

MetaMorph Offline, version 7.10.1

Prism 8 for Mac OS, version 8.4.1

Adobe Illustrator for Mac OS, version 24.2.1

R studio, version 3.6.3

Microsoft Office for Mac

<u>Statistical Analysis</u>

All data on bar graphs are presented as mean ± standard error of the mean (SEM). The
nuclear/cytoplasmic distribution of lncRNAs of the BMDM fractionation experiments is reported as violin
plots with median and quartiles. The chemotaxis assay is represented as xy-plot with mean ± standard
error of the mean (SEM). The ARI calculations are displayed as boxplot with median and quartiles.
Statistical analysis was performed using GraphPad Prism 8. Data distribution and the difference
between variances were tested using the Kolmogorov–Smirnov. Depending on the distribution of data,
unpaired t-test or Mann–Whitney test were used to determine statistical significance. For chemotaxis
experiments, statistical significance was tested using repeated measures two-way ANOVA followed by
Sidak correction. Significance is depicted as $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$. The
detailed statistical analyses are provided as a Source Data file.

ACCESSION NUMBERS

All sequencing data sets can be accessed at GEO database under GSE210732.

ACKNOWLEDGEMENT

We thank Lucia Koblitz for expert support with primary cell preparations, Julia Kerler and Anton Bomhard
for expert assistance in animal surgeries and Dr. Elisabeth Graf for assistance with RNA-seq. We thank
Dr. Daniel Andergassen and Dr. Bernhard Laggerbauer for critical reading and helpful discussions.

FUNDING

This project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation)—Project- ID 403584255–TRR 267 (S.E., A.D., R.P.B., M.S.L., L.M., T.M., J.G.). S.E., A.D.,
L.A., D.E., K.H., N.P., H.S., C.S., T.W., J.G., E.T., H.B.S., T.M., N.J., M.S.L. and R.P.B. were supported
in the framework of the German Centre for Cardiovascular Research (DZHK), and the partner site in
Munich (MHA) (S.E., A.D., H.S., C.S., T.W., J.G., H.B.S., T.M.) projects 81X3600404 (A.D.),
81X2600245 (C.S.) and 81X2600246 (C.S.) or partner site RheinMain, Frankfurt (N.J., M.S.L. and
R.P.B.). S.E. and D.R. were supported by DFG through the Research Training Group GRK2338 and
the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-
Curie grant agreement no. 813716.

Author contributions

A.D. and S.E. designed the study and wrote the manuscript. A.D., L.A. D.E., S.B., E.T., K.H., N.P. and
A.U. performed the experiments or analyzed the resulting data. M.B. provided the *Schlafenlnc* knockout

mouse. R.P.B., J.G., P.K., M.S.L., L.M., T.M., D.R., H.B.S., M.S.-S., C.S. and T.W. provided
experimental models, instruments and crucial expertise. All authors read, commented on and approved
the manuscript.

CONFLICT OF INTEREST

None declared.

REFERENCES

1. Goodall, G. J. & Wickramasinghe, V. O. RNA in cancer. *Nat. Rev. Cancer* **21**, 22–36 (2021).

2. Aliperti, V., Skonieczna, J. & Cerase, A. Long non-coding rna (Lncrna) roles in cell biology,
neurodevelopment and neurological disorders. *Non-coding RNA* **7**, (2021).

3. Anderson, K. M. & Anderson, D. M. LncRNAs at the heart of development and disease. *Mamm.*
Genome **33**, 354–365 (2022).

4. Wysoczynski, M., Kim, J., Iv, J. B. M. & Uchida, S. Macrophage Long Non-Coding RNAs in
Pathogenesis of Cardiovascular Disease. (2020).

5. Klattenhoff, C. A. *et al.* Braveheart, a long noncoding RNA required for cardiovascular lineage
commitment. *Cell* **152**, 570–583 (2013).

6. Li, K. *et al.* A noncoding antisense RNA in tie-1 locus regulates tie-1 function in vivo. *Blood* **115**,
133–139 (2010).

7. Micheletti, R. *et al.* The long noncoding RNA Wisper controls cardiac fibrosis and remodeling.
Sci. Transl. Med. **9**, (2017).

8. Zaman, R. & Epelman, S. Resident cardiac macrophages: Heterogeneity and function in health
and disease. *Immunity* **55**, 1549–1563 (2022).

9. Gomez Perdiguero, E. *et al.* Tissue-resident macrophages originate from yolk-sac-derived
erythro-myeloid progenitors. *Nature* **518**, 547–551 (2015).

10. Epelman, S. *et al.* Embryonic and adult-derived resident cardiac macrophages are maintained
through distinct mechanisms at steady state and during inflammation. *Immunity* **40**, 91–104
(2014).

11. Dick, S. A. *et al.* Three tissue resident macrophage subsets coexist across organs with
conserved origins and life cycles. *Sci. Immunol.* **7**, 1–19 (2022).

12. Hulsmans, M. *et al.* Macrophages Facilitate Electrical Conduction in the Heart. *Cell* **169**, 510-
522.e20 (2017).

13. Grune, J. *et al.* Neutrophils incite and macrophages avert electrical storm after myocardial
infarction. *Nat. Cardiovasc. Res.* **1**, 649–664 (2022).

14. Fujiu, K. *et al.* A heart-brain-kidney network controls adaptation to cardiac stress through tissue
macrophage activation. *Nat. Med.* **23**, 611–622 (2017).

15. Cahill, T. J. *et al.* Tissue-resident macrophages regulate lymphatic vessel growth and patterning
in the developing heart. *Dev.* **148**, (2021).

16.   Ramanujam, D. *et al.* MicroRNA-21–Dependent Macrophage-to-Fibroblast Signaling
734   Determines the Cardiac Response to Pressure Overload. *Circulation* **143**, 1513–1525 (2021).

735   17.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic
736   data across different conditions, technologies, and species. *Nat. Biotechnol.* **36**, 411–420 (2018).

737   18.   Harrow, J. *et al.* GENCODE: The reference human genome annotation for the ENCODE project.
738   *Genome Res.* **22**, 1760–1774 (2012).

739   19.   Skelly, D. A. *et al.* Single-Cell Transcriptional Profiling Reveals Cellular Diversity and
740   Intercommunication in the Mouse Heart. *Cell Rep.* **22**, 600–610 (2018).

741   20.   McLellan, M. A. *et al.* High-Resolution Transcriptomic Profiling of the Heart during Chronic Stress
742   Reveals Cellular Drivers of Cardiac Fibrosis and Hypertrophy. *Circulation* 1448–1463 (2020)
743   doi:10.1161/CIRCULATIONAHA.119.045115.

744   21.   Mercer, T. R., Dinger, M. E., Sunkin, S. M., Mehler, M. F. & Mattick, J. S. Specific expression of
745   long noncoding RNAs in the mouse brain. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 716–721 (2008).

746   22.   Derrien, T. *et al.* The GENCODE v7 catalog of human long noncoding RNAs: Analysis of their
747   gene structure, evolution, and expression. *Genome Res.* **22**, 1775–1789 (2012).

748   23.   Cabili, M. *et al.* Integrative annotation of human large intergenic noncoding RNAs reveals global
749   properties and specific subclasses. *Genes Dev.* **25**, 1915–1927 (2011).

750   24.   Ulitsky, I. Evolution to the rescue: Using comparative genomics to understand long non-coding
751   RNAs. *Nat. Rev. Genet.* **17**, 601–614 (2016).

752   25.   Piccoli, M.-T. *et al.* Inhibition of the Cardiac Fibroblast-Enriched lncRNA Meg3 Prevents Cardiac
753   Fibrosis and Diastolic Dysfunction. *Circ. Res.* CIRCRESAHA.117.310624 (2017)
754   doi:10.1161/CIRCRESAHA.117.310624.

755   26.   Hofmann, P. *et al.* Long non-coding RNA H19 regulates endothelial cell aging via inhibition of
756   STAT3 signalling. *Cardiovasc. Res.* **115**, 230–242 (2019).

757   27.   Han, P. *et al.* A long noncoding RNA protects the heart from pathological hypertrophy. *Nature*
758   **514**, 102–106 (2014).

759   28.   Alexanian, M. & Haldar, S. M. The cardiac myofibroblast from supporting cast to leading role?
760   *Circ. Res.* **123**, 1258–1260 (2018).

761   29.   Guttman, M. *et al.* Chromatin signature reveals over a thousand highly conserved large non-
762   coding RNAs in mammals. *Nature* **458**, 223–227 (2009).

763   30.   Wucher, V. *et al.* FEELnc: A tool for long non-coding RNA annotation and its application to the
764   dog transcriptome. *Nucleic Acids Res.* **45**, 1–12 (2017).

765   31.   Liu, F., Zhou, P., Wang, Q., Zhang, M. & Li, D. The Schlafen family: complex roles in different
766   cell types and virus replication. *Cell Biol. Int.* **42**, 2–8 (2018).

767   32.   Hezroni, H. *et al.* Principles of Long Noncoding RNA Evolution Derived from Direct Comparison
768   of Transcriptomes in 17 Species. *Cell Rep.* **11**, 1110–1122 (2015).

769   33.   Luo, Y. *et al.* New developments on the Encyclopedia of DNA Elements (ENCODE) data portal.
770   *Nucleic Acids Res.* **48**, D882–D889 (2020).

771   34.   Avsec, Z., Barekatain, M., Cheng, J. & Gagneur, J. Modeling positional effects of regulatory
772   sequences with spline transformations increases prediction accuracy of deep neural networks.
773   *Bioinformatics* **34**, 1261–1269 (2018).

774   35.   Van Nostrand, E. L. *et al.* Robust transcriptome-wide discovery of RNA-binding protein binding
775   sites with enhanced CLIP (eCLIP). *Nat. Methods* **13**, 508–514 (2016).

776   36.   Dunham, I. *et al.* An integrated encyclopedia of DNA elements in the human genome. *Nature*
777   **489**, 57–74 (2012).

778   37.   Shi, L., Shi, L., Wang, X. & He, J. Regulatory Roles of Osteopontin in Production of Monocyte-
779   Origin MCP-1. *Cell Transplant.* **27**, 1185–1194 (2018).

780   38.   Gregory, J. L. *et al.* Macrophage Migration Inhibitory Factor Induces Macrophage Recruitment
781   via CC Chemokine Ligand 2. *J. Immunol.* **177**, 8072–8079 (2006).

782   39.   Uhlén, M. *et al.* The human secretome. *Sci. Signal.* **12**, 1–9 (2019).

783   40.   Redecke, V. *et al.* Hematopoietic progenitor cell lines with myeloid and lymphoid potential. *Nat.*
784   *Methods* **10**, 795–803 (2013).

785   41.   Platt, R. J. *et al.* CRISPR-Cas9 knockin mice for genome editing and cancer modeling. *Cell* **159**,
786   440–455 (2014).

787   42.   Marx, V. Method of the Year: spatially resolved transcriptomics. *Nat. Methods* **18**, 9–14 (2021).

788   43.   Svensson, V. *et al.* Power analysis of single-cell rnA-sequencing experiments. *Nat. Methods* **14**,
789   381–387 (2017).

790   44.   Rhee, A. J. & Lavine, K. J. New Approaches to Target Inflammation in Heart Failure: Harnessing
791   Insights from Studies of Immune Cell Diversity. *Annu. Rev. Physiol.* **82**, 1–20 (2020).

792   45.   Duncan, S. E. *et al.* Macrophage Activities in Myocardial Infarction and Heart Failure. *Cardiol.*
793   *Res. Pract.* **2020**, (2020).

794   46.   Moskalik, A., Niderla-Bielińska, J. & Ratajska, A. Multiple roles of cardiac macrophages in heart
795   homeostasis and failure. *Heart Fail. Rev.* (2021) doi:10.1007/s10741-021-10156-z.

796   47.   Peet, C., Ivetic, A., Bromage, D. I. & Shah, A. M. Cardiac monocytes and macrophages after
797   myocardial infarction. *Cardiovasc. Res.* **116**, 1101–1112 (2020).

798   48.   Liao, X. *et al.* Distinct roles of resident and nonresident macrophages in nonischemic
799   cardiomyopathy. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E4661–E4669 (2018).

800   49.   Leuschner, F. *et al.* Rapid monocyte kinetics in acute myocardial infarction are sustained by
801   extramedullary monocytopoiesis. *J. Exp. Med.* **209**, 123–137 (2012).

802   50.   Dutta, P. *et al.* Myocardial infarction activates CCR2+ hematopoietic stem and progenitor cells.
803   *Cell Stem Cell* **16**, 477–487 (2015).

804   51.   Dutta, P. *et al.* Myocardial infarction accelerates atherosclerosis. *Nature* **487**, 325–329 (2012).

805   52.   Aoki, S. *et al.* Elevated peripheral blood mononuclear cell count is an independent predictor of
806   left ventricular remodeling in patients with acute myocardial infarction. *J. Cardiol.* **57**, 202–207
807   (2011).

808   53.   Van Der Laan, A. M. *et al.* A proinflammatory monocyte response is associated with myocardial
809   injury and impaired functional outcome in patients with ST-segment elevation myocardial
810   infarction: Monocytes and myocardial infarction. *Am. Heart J.* **163**, 57-65.e2 (2012).

811   54.   Swirski, F. K. & Al., E. Identification of Splenic Reservoir. *Science (80-.).* **325**, 612–616 (2009).

812   55.   Frantz, S. & Nahrendorf, M. Cardiac macrophages and their role in ischaemic heart disease.
813   *Cardiovasc. Res.* **102**, 240–248 (2014).

56.   Bajpai, G. *et al.* Tissue Resident CCR2- and CCR2+ Cardiac Macrophages Differentially

Orchestrate Monocyte Recruitment and Fate Specification Following Myocardial Injury. *Circ.*
Res. **124**, 263–278 (2019).

57. Yakar Tülüce, S. *et al.* Galectin-3 levels in patients with hypertrophic cardiomyopathy and its
relationship with left ventricular mass index and function. *Anatol. J. Cardiol.* **16**, 344–348 (2016).

58. Frunza, O. *et al.* Myocardial Galectin-3 Expression Is Associated with Remodeling of the
Pressure-Overloaded Heart and May Delay the Hypertrophic Response without Affecting
Survival, Dysfunction, and Cardiac Fibrosis. *Am. J. Pathol.* **186**, 1114–1127 (2016).

59. Danella Polli, C. *et al.* Monocyte Migration Driven by Galectin-3 Occurs through Distinct
Mechanisms Involving Selective Interactions with the Extracellular Matrix. *ISRN Inflamm.* **2013**,
1–9 (2013).

60. Jia, W., Kidoya, H., Yamakawa, D., Naito, H. & Takakura, N. Galectin-3 accelerates M2
macrophage infiltration and angiogenesis in tumors. *Am. J. Pathol.* **182**, 1821–1831 (2013).

61. Papaspyridonos, M. *et al.* Galectin-3 is an amplifier of inflammation in atherosclerotic plaque
progression through macrophage activation and monocyte chemoattraction. *Arterioscler.*
Thromb. Vasc. Biol. **28**, 433–440 (2008).

62. Ramanujam, D., Sassi, Y., Laggerbauer, B. & Engelhardt, S. Viral vector-based targeting of miR-
21 in cardiac nonmyocyte cells reduces pathologic remodeling of the heart. *Mol. Ther.* **24**, 1939–
1948 (2016).

63. Hwang, H. W., Wentzel, E. A. & Mendell, J. T. A hexanucleotide element directs microRNA
nuclear import. *Science (80-.).* **315**, 97–100 (2007).

64. Ziegler, C. *et al.* The long non-coding RNA LINC 00941 and SPRR 5 are novel regulators of
human epidermal homeostasis . *EMBO Rep.* **20**, 1–13 (2019).

65. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21 (2019).

66. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression. *Genome Biol.* **20**, 1–15 (2019).

67. Scrucca, L., Fop, M., Murphy, T. B. & Raftery, A. E. Mclust 5: Clustering, classification and
density estimation using Gaussian finite mixture models. *R J.* **8**, 289–317 (2016).

68. Linderman, G. C., Zhao, J. & Kluger, Y. Zero-preserving imputation of scRNA-seq data using
low-rank approximation. *bioRxiv* 397588 (2018) doi:10.1101/397588.

69. Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional
annotation of gene lists (2021 update). *Nucleic Acids Res.* 1–6 (2022) doi:10.1093/nar/gkac194.

70. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat. Commun.* **12**,
1–20 (2021).

71. Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and collaborative biomedical
analyses: 2018 update. *Nucleic Acids Res.* **46**, W537–W544 (2018).

72. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

73. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nat. Biotechnol.* **33**, 290–295 (2015).

74. Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing genomic features.
Bioinformatics **26**, 841–842 (2010).

75. Wang, L. *et al.* CPAT: Coding-potential assessment tool using an alignment-free logistic

regression model. *Nucleic Acids Res.* **41**, 1–7 (2013).

76.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

77.   Metsalu, T. & Vilo, J. ClustVis: A web tool for visualizing clustering of multivariate data using

Principal Component Analysis and heatmap. *Nucleic Acids Res.* **43**, W566–W570 (2015).

78.   Bindea, G. *et al.* ClueGO: A Cytoscape plug-in to decipher functionally grouped gene ontology

and pathway annotation networks. *Bioinformatics* **25**, 1091–1093 (2009).

79.   Shannon, P. *et al.* Cytoscape: A software Environment for integrated models of biomolecular

interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

80.   Avsec, Ž. *et al.* The Kipoi repository accelerates community exchange and reuse of predictive

models for genomics. *Nat. Biotechnol.* **37**, 592–600 (2019).

Figure 1

Figure 1: LncRNA are important determinants for definition of specific cell types.

(A) Illustration for the measurement of lncRNA in murine myocardium and re-clustering analysis. (B) UMAP projection of murine myocardium as measured by single-cell sequencing (scRNA-seq). Cell populations are colored by type as indicated by the label. DC: dendritic cells, EC: endothelial cells, CM: cardiac myocytes. (C) Bubble plot showing lncRNA marker genes conserved across conditions. Bubble size corresponds to enrichment (logFC) and fill color to expression value (log10). Only values/bubbles with a significant enrichment (p-val < 0.05) are shown. (D) Feature plots showing mRNA and lncRNA markers for fibroblasts, endothelial cells (EC) and macrophages. (E), (F), (G) Adjusted Rand Index (ARI) calculation. In order to compare the clusterings with and without lncRNAs, the respective complete gene set was considered to yield the optimal clustering. We iterated the clustering in 0.01 steps with resolutions ranging from 0.10 to 1.00, testing for the similarity to optimal clustering using adjusted Rand index scoring. Here, the more similar a clustering would be to the "optimal" one the closer it will be to 1. (E) ARI calculation for the top variable genes (3,000 genes). (F) ARI calculation for all annotated genes (Gencode vM22 [18]). (G) ARI calculation for all expressed genes (expression value >0). pc: protein coding (H) Umap projection of murine myocardium with cell population clustering based on protein coding genes plus lncRNA genes (left panel) or only protein coding genes (right panel). The middle panel shows differentially assigned cell population clusters.

Figure 2: Dynamic changes of lncRNA expression in crMPs during pressure overload.

Bubble plot showing the top 5 up-regulated and top 5 down-regulated lncRNA genes after TAC for the sub-populations of the major cell types in the heart: (A) Fibroblasts (B) Endothelial cells and (C) Macrophages. Fill color corresponds to regulation in TAC vs. sham (lnFC TAC/sham) and bubble size corresponds to expression value (log10). Only values with a significant enrichment (p-val < 0.05) are shown. Some genes are among the top regulated genes in more than one cell type.

Figure 3: Determination and annotation of the lncRNA landscape of cardiac resident macrophages.

(A) Unsupervised clustering of known and novel lncRNAs using normalized counts values, rows were averaged across samples for clustering. Genes shown are expressed with a TPM value >5 and differentially regulated (enriched) with a q-val of <0.05. (B) Flow scheme for the identification of lncRNA candidates using the indicated filters. (C) The Schlafen family locus showing the expression in cardiac resident macrophages (accumulated plus and minus strand). Lower panel: Zoom in of the *Schlafenlnc* locus in mouse as well as human for comparison. (D) Expression of the lncRNA *Schlafenlnc* in Mac1 (*Lyve1+* macrophages) and Mac2 (MHCII high, *Cd74+* macrophages) after sham or TAC in the scRNA-seq data. (E) Northern Blot showing the expression of *Schlafenlnc* transcripts in bone marrow-derived macrophages in comparison to fibroblasts (NIH3T3 cell line). (F) Relative nuclear and cytoplasmic localization of *Malat1* (nuclear lncRNA), *Rpl32* (mRNA) and *Schlafenlnc* as determined by fractionation of macrophages followed by quantification through RNA-Seq. (G) Detection of *Schlafenlnc* (green) in macrophages by RNA *in situ* hybridization (RNAScope). Nuclei were stained with DAPI (blue), Phalloidin (red) was used for outlining the cellular body (Scale bars = 15μm).

Figure 4: *Schlafenlnc*-deficient mice show improved cardiac function and reduced inflammatory response after transverse aortic constriction (TAC).

(A) Experimental setup of cardiac phenotyping of *Schlafenlnc$^{-/-}$* and wildtype mice. (B) Quantification of normalized heart weight after TAC. HW= heart weight, TL= tibia length. C) Ejection fraction changes after 6 days TAC. (C) Left: Quantification of interstitial fibrosis by Sirius red/Fast green staining. Right: Representative images of interstitial fibrosis, scale bar = 200µm. (D) Representative images and quantification of macrophage/monocyte content by Cd68 staining in WT and *Schlafenlnc$^{-/-}$* hearts after 6d after TAC. White arrows label Cd68-positive cells, scale bar = 25µm.

Figure 5: scRNA-seq of *Schlafenlnc*[-/-] mice reveals macrophage polarization towards an anti-inflammatory phenotype after TAC.

(A) Experimental setup of scRNA-seq of *Schlafenlnc*[-/-] and wildtype mice. (B) Dimensional reduction (Umap) representation of combined measured cells with assigned cell types. (C) Cell interaction analysis of *Schlafenlnc* WT and KO mice (CD45+ cells from blood and heart) after TAC using CellChat[70]. (D) Differential signaling in *Schlafenlnc*[-/-] and WT macrophage and monocyte populations after TAC. (E) RNA Velocity of *Schlafenlnc*[-/-] and WT cells from heart after 6 days of TAC. (F) Comparison of de-regulated genes (q-val<0.05) in TAC vs. basal condition in WT compared to de-regulated genes after 6d TAC KO vs. WT in *Lyve1*[+] macrophages (G) Gene ontology terms of genes de-regulated in *Schlafenlnc*-deficient mice and after 6 days TAC.

Figure 6: *SchlafenInc* is required for the activation of macrophage gene programs for cell adhesion and migration.

(A) MA plot showing differentially expressed genes in *SchlafenInc*$^{-/-}$ vs. Ctrl macrophages. Significantly deregulated genes (FDR < 0.05) are shown in red. (B) Heatmap of an unsupervised clustering showing the top 20 downregulated genes in *SchlafenInc*$^{-/-}$ macrophages. (C) Measurement of Galectin-3 protein secretion by control and *SchlafenInc*$^{-/-}$ macrophages as determined by ELISA. The supernatant of one control cell line and three *SchlafenInc*$^{-/-}$ clones were used in n=3 independent experiments. (D) Functionally grouped gene ontology term analysis of biological processes associated with significantly downregulated genes (FDR < 0.05, log2FC < -1) in *SchlafenInc*$^{-/-}$ macrophages. (E) Adhesion assay with control and *SchlafenInc*$^{-/-}$ macrophages (Ctrl clones n=2, KO clones n=3; 4 independent experiments). (F) Chemotaxis assay against a gradient of the chemokine C5a with control and *SchlafenInc*$^{-/-}$ macrophages stimulated with 10ng/ml LPS overnight (Ctrl clones n=1, KO clones n=3; 3 independent experiments).