Trade-offs between baseline thermal tolerance and thermal tolerance plasticity are much less

common than it appears

Alex R. Gunderson

Department of Ecology and Evolutionary Biology, Tulane University, New Orleans, LA, USA

Running header: Constraints on plasticity

email: agunderson@tulane.edu

Keywords: acclimation, global warming, physiological ecology, thermal adaptation, phenotypic

flexibility

Abstract

Thermal tolerance plasticity is a core mechanism by which organisms can mitigate the effects of

climate change. As a result, there is a need to understand how variation in tolerance plasticity

arises. The Baseline Tolerance/Plasticity Trade-off Hypothesis (hereafter referred to as the

Trade-off Hypothesis, TOH) has recently emerged as a potentially powerful explanation. The

TOH posits that organisms with high baseline thermal tolerance have reduced thermal

tolerance plasticity relative to those with low baseline tolerance. Many studies have found

support for the TOH. However, this support must be regarded cautiously because the most

common means of testing the TOH can yield spurious 'trade-offs' due to regression to the

mean. I acquired data for 25 previously-published analyses that supported the TOH at the

intraspecific level and reanalyzed them after applying a method that adjusts plasticity estimates

for regression to the mean. Only four of the 25 analyses remained statistically significant after

adjustment, and effect size and variance explained decreased in all cases. The few data sets in

which support for the TOH was maintained after adjustment point to areas of future study, but

are too few to make generalizations at this point. In sum, regression to the mean has led to a

substantial overestimation of support for the TOH and must be accounted for in future tests of

the hypothesis.

Introduction

The capacity to physiologically adjust to temperature change through phenotypic plasticity is

thought to influence vulnerability to anthropogenic warming (Gunderson, Dillon, & Stillman,

2017; Huey et al., 2012; Morley, Peck, Sunday, Heiser, & Bates, 2019; Seebacher, White, &

Franklin, 2015; Somero, 2010). Uncovering how and why taxa differ in physiological plasticity is

therefore of great interest (Kelly, 2019; Pottier, Burke, Drobniak, Lagisz, & Nakagawa, 2021;

Rohr et al., 2018). Recently, the Trade-off Hypothesis (TOH) has emerged as a potentially

powerful mechanism to explain variation in the plasticity of thermal tolerance limits (Stillman,

2003; van Heerwaarden & Kellermann, 2020). This hypothesis posits that taxa with high

baseline levels of thermal tolerance will exhibit low levels of thermal tolerance plasticity. From

a mechanistic perspective, the TOH is usually discussed in terms of phenotypic constraint.

Thermal limits can only be so high, and therefore plasticity must be reduced as baseline

tolerance rises. The implications of this hypothesis, if and when true, are many, and touch not

just on global change biology but fundamental aspects of physiological ecology and evolution

(reviewed in van Heerwaarden & Kellermann, 2020).

Support for the TOH has been found in several studies at both the intraspecific and

interspecific levels (Barley et al., 2021; Comte & Olden, 2017; van Heerwaarden & Kellermann,

2020; see also Table 1). Nonetheless, these results must be regarded carefully. This is because

many tests of the TOH open the door for regression to the mean to generate spurious support

(van Heerwaarden & Kellermann, 2020). Baseline thermal tolerance values are used to calculate

plasticity, and then plasticity is modeled as a function of the baseline value. A common term is

present on both sides of the function, potentially leading to a negative relationship that is a

statistical artifact.

Approaches for identifying, avoiding, and/or correcting for regression to the mean exist,

but to date have rarely been applied to tests of the TOH. O'Neill, Davis, & MacMillan (2021)

tentatively found support for the TOH among *Drosophila melanogaster* individuals with respect

to cold tolerance, but from the correlation structure within their data, they inferred that the

relationship was probably artifactual. Deery et al. (2021) found initial support for the TOH at

the individual level among *Anolis* lizards with respect to heat hardening, but subsequently

conducted a data resampling analysis (Ghalambor et al., 2015; Jackson & Somers, 1991) and

concluded that the relationships were spurious. Finally, Gunderson & Revell (2022) used

simulations to show how pervasive regression to the mean is when testing the TOH at

macroevolutionary scales. They implemented a post-hoc method for adjusting plasticity data to

account for regression to the mean (Kelly & Price, 2005), and found that doing so reduced

support for the TOH in a previously-published dataset (Armstrong, Tanner, & Stillman, 2019;

Gunderson & Revell, 2022). In sum, studies that attempt to account for regression to the mean

in tests of the TOH have generally found that support for the hypothesis declines. How general

is this outcome?

Methods

I reanalyzed data from published studies that tested and found support for the TOH, focusing

on work conducted at the intraspecific level (see Gunderson & Revell (2022) for analysis of the

same issues at a macroevolutionary scale). I started with data from the studies summarized in

Table 1 of van Heerwaarden & Kellermann (2020), and added as many subsequently-published

data sets as I could identify. Data for some studies could not be obtained or the analysis

supporting the TOH did not allow for correction. In total, I reanalyzed 25 correlations between

baseline thermal tolerance and thermal tolerance plasticity representing 10 species (Table 1). I

first conducted correlation analyses on the original data ("unadjusted"), and then with plasticity

values adjusted to remove the effect of regression to the mean following the method

developed by Kelly & Price (2005; "adjusted"). Detailed explanation of the method can be found

Table 1. Summary of the data sets reanalyzed in the present study. "Unadjusted" correlation coefficients were calculated using raw plasticity data as in the original analyses, and "Adjusted" correlation coefficients were calculated after removing the effect of regression to the mean (Kelly and Price 2005). Correlation coefficients (r) in bold indicate statistically significant relationships at the α = 0.05 level. If a study conducted multiple analyses with different subsets of their data, the subset is indicated by the "Sample" column. The "Metric" definitions are as follows: *CTmax* = critical thermal maximum; *LD50* = temperature of 50% mortality; *CCRT* = chill coma recovery time; *CCO* = chill coma onset.

Clade	Species	Level	Metric	Sample	Unadjusted r	Adjusted r	Reference
Heat tolerance							
Fish	*Gasterosteus aculeatus*	Individual	CTmax	-	**-0.59**	0.02	Mottola et al. 2022
Fish	*Danio rerio*	Individual	CTmax	Females	**-0.70**	-0.09	Morgan et al. 2018
Fish	*D. rerio*	Individual	CTmax	Males	**-0.80**	0.02	Morgan et al. 2018
Reptile	*Anolis sagrei*	Individual	CTmax	-	**-0.55**	0.04	Deery et al. 2021
Reptile	*A. carolinensis*	Individual	CTmax	-	**-0.65**	0.00	Deery et al. 2021
Reptile	*Lampropholis coggeri*	Individual	CTmax	Observer 1	**-0.77**	**-0.49**	Phillips et al. 2016
Reptile	*L. coggeri*	Individual	CTmax	Observer 2	**-0.52**	-0.11	Phillips et al. 2016
Reptile	*Urosaurus ornatus*	Individual	CTmax		**-0.48**	-0.07	Glbert and Miles 2019
Insect	*Drosophila melanogaster*	Population	CTmax	-	**-0.73**	**-0.52**	van Heerwaarden et al. 2016
Crustacean	*Acartia tonsa*	Population	LD50	-	**-0.84**	-0.06	Sasaki and Dam 2019
Crustacean	*A. tonsa*	Population	LD50	-	**-0.97**	0.66	Sasaki and Dam 2020
Crustacean	*A. tonsa*	Selection line	LD50	CT pop., control	**-0.99**	0.54	Sasaki and Dam 2021
Crustacean	*A. tonsa*	Selection line	LD50	CT pop., warming	**-0.94**	-0.38	Sasaki and Dam 2021
Crustacean	*A. tonsa*	Selection line	LD50	SP pop., control	**-1.00**	0.84	Sasaki and Dam 2021
Crustacean	*A. tonsa*	Selection line	LD50	SP pop., warming	**-0.99**	**-0.99**	Sasaki and Dam 2021
Crustacean	*A. tonsa*	Selection line	LD50	PG pop., control	**-0.97**	0.13	Sasaki and Dam 2021
Crustacean	*A. tonsa*	Selection line	LD50	PG pop., warming	**-0.98**	**-0.97**	Sasaki and Dam 2021
Cold tolerance							
Insect	*D. melanogaster*	Female lines	CCRT	Low heat	**-0.81**	0.01	Noh et al. 2017
Insect	*D. melanogaster*	Female lines	CCRT	Mid heat	**-0.77**	-0.04	Noh et al. 2017
Insect	*D. melanogaster*	Female lines	CCRT	High heat	**-0.87**	0.02	Noh et al. 2017
Insect	*D. melanogaster*	Individuals	CCRT	15℃ Acclimated	**-0.68**	-0.09	O'Neill et al. 2021
Insect	*D. melanogaster*	Individuals	CCRT	25℃ Acclimated	**-0.62**	-0.03	O'Neill et al. 2021
Insect	*D. melanogaster*	Individuals	CCO	15℃ Acclimated	**-0.63**	0.00	O'Neill et al. 2021
Insect	*D. melanogaster*	Individuals	CCO	25℃ Acclimated	**-0.62**	0.01	O'Neill et al. 2021
Insect	*Aphaenogaster picea*	Populations	CCRT	-	**-0.88**	0.04	Nguyen et al. 2019

95   in Kelly & Price (2005) and Gunderson & Revell (2022). All analyses were conducted in R (R

96   Development Core Team, 2021). Data adjustment was done with a function I wrote in R

97   ("rttm.adj"), and the script can be found in the Supplementary Materials.

Results and Discussion

100   Two examples of what data look like before and after adjustment for regression to the mean

101   are shown in Figure 1. The first example (Figure 1 A, B) shows heat hardening data for

Figure 1. Examples of how tests of the Trade-off Hypothesis (TOH) are affected when plasticity values are adjusted to account for regression to the mean using the Kelly & Price (2005) method. A, B). Heat hardening data from the lizard *Anolis carolinensis* (Deery, Rej, Haro, & Gunderson, 2021). Support for the TOH was lost after data adjustment. C,D) Heat tolerance acclimation data from adult *Drosophila melanogaster* (van Heerwaarden, Kellermann, & Sgrò, 2016). Support for the TOH was maintained after data adjustment, though the effect size and variance explained decreased.

individuals of the lizard *Anolis carolinensis* (Deery et al., 2021). In the unadjusted data (Figure 1

A), there is a strong negative relationship between an individual's baseline heat tolerance limit

and how much their heat tolerance limit increases after heat shock (i.e., plasticity). However,

that relationship disappears after plasticity values are adjusted for regression to the mean

(Figure 1 B). The second example shows data on heat tolerance acclimation in *Drosophila*

melanogaster (van Heerwaarden et al., 2016). In this case, a negative relationship between

baseline tolerance and plasticity is maintained in both the unadjusted and adjusted data (Figure

1 C, D), though the effect of regression to the mean is still noticeable: the correlation coefficient

(and therefore variance explained) and the effect size (slope) are all smaller using the adjusted

versus the unadjusted plasticity data.

When this approach is applied across all the data sets, it becomes apparent that most

relationships between baseline thermal tolerance and thermal tolerance plasticity are spurious

due to regression to the mean (Table 1). Of the 25 results consistent with the TOH using

unadjusted data, only four remain statistically significant at the α = 0.05 level after adjustment

of plasticity values. Furthermore, correlation coefficients decrease in almost every case

whether statistical significance is maintained or not. The magnitude and sometimes direction of

effect also changes considerably after data adjustment. Slopes of the relationship between

baseline tolerance and tolerance plasticity for all data sets become less negative when using

adjusted versus unadjusted plasticity values (Figure 2).

Some of the regression to the mean effects demonstrated here are consistent with

conclusions drawn by the original authors using different approaches (e.g., Deery et al., 2021;

O'Neill et al., 2021; see above). However, most of the initial, unadjusted analyses were

Figure 2. Slopes for the relationship between baseline thermal tolerance and thermal tolerance plasticity for the data sets in Table 1. Intercepts were set a 0 so all relationships could easily be visualized together. *Left*: Slopes when thermal tolerance plasticity values are not adjusted for regression to the mean. *Right*: Slopes when thermal tolerance plasticity values are adjusted for regression to the mean. Solid lines: data sets for which significance was maintained at $\alpha = 0.05$ after data adjustment. Dashed lines: data sets for which statistical significance was not maintained after data adjustment.

interpreted as evidence consistent with the TOH. Comparing data sets for which support for the

TOH was and was not maintained after data adjustment is informative. Support for the TOH

was rarely maintained after data adjustment in individual-level studies (Table 1). The lone

example was with respect to heat hardening in the lizard *Lampropholis coggeri* (Phillips et al.,

2016). Interpretation of this result is not straightforward, however, because it is not consistent

with other data from the same species in the same study: two different observers measured

thermal tolerance values, and support for the TOH was maintained after adjustment with data

from one observer, but not the other (Table 1).

The other data sets for which support for the TOH was maintained came from a study of

trade-offs between developmental and adult plasticity in *Drosophila melanogaster* (van

Heerwaarden et al., 2016) and an experimental evolution study with the copepod *Acartia tonsa*

(Sasaki & Dam, 2021). In the *Drosophila* study, animals were exposed to different

developmental temperatures, after which these same animals had their capacity for adult

thermal acclimation measured. The authors found that development in warm conditions led to

increased baseline thermal tolerance, but lower capacity for adult acclimation (van

Heerwaarden et al., 2016). These data are consistent with a mechanistic link between

developmental and adult plasticity (Beaman, White, & Seebacher, 2016; van Heerwaarden et

al., 2016).

In the experimental evolution study with the copepod *A. tonsa*, control and high

temperature lines were established from three different source populations (Sasaki & Dam,

2021). In the initial analysis, there was an apparent trade-off between baseline thermal

tolerance and plasticity over time across all control and high temperature groups from all three

source populations. However, data adjustment shows that the relationship was artifactual in all

three of the control groups, but not in two of the three high temperature selection groups

(Table 1). In the third high temperature selection group, the correlation coefficient remained

relatively large and negative after data adjustment (-0.38), though statistical significance was

not maintained (Table 1). In sum, the data are consistent with a real evolutionary correlation

between baseline tolerance and plasticity as populations adapt to warmer conditions (Sasaki &

Dam, 2021). However, the generality of this conclusion is not clear even within this species, as

the TOH is not supported after data adjustment in a different study focused on natural

populations evolving under different thermal regimes (Sasaki & Dam, 2019; Table 1).

Conclusion

Due to regression to the mean, support for the Trade-off Hypothesis is much lower than it

appears at first glance (Table 1; see also Sgro et al. (2010) and van Heerwaarden, Lee,

Overgaard, & Sgro (2014), who report analyses that reject the TOH with unadjusted data).

Studies in which support for the TOH is maintained after data adjustment point to situations in

which trade-offs between baseline tolerance and plasticity may be most common. However, at

this point there are too few examples to make generalizations. More tests of the TOH are

certainly warranted. At the core, the inflated support for the TOH is a problem of identifying

the appropriate null expectation (Hertz, Huey, & Stevenson, 1993). A lack of association

between baseline thermal tolerance and thermal tolerance plasticity is not always the correct

null hypothesis based on the methodological approaches taken (Gunderson & Revell, 2022). By

recognizing such situations and accounting for them, we can more quickly develop a clear

picture of the forces that influence the evolution of thermal tolerance plasticity, and in what

situations different mechanisms are important. Doing so will greatly increase our understanding

of organismal responses to ongoing anthropogenic warming.

Acknowledgements

I would like to thank authors who provided data that were not publicly available. Eric Gangloff

provided helpful feedback on the manuscript.

Data availability

The data for this paper will be made available in Dryad.

References
180

Armstrong, E. J., Tanner, R. L., & Stillman, J. H. (2019). High heat tolerance is negatively
correlated with heat tolerance plasticity in nudibranch mollusks. *Physiological and*
Biochemical Zoology, 92(4), 430-444.

Barley, J. M., Cheng, B. S., Sasaki, M., Gignoux-Wolfsohn, S., Hays, C. G., Putnam, A. B., . . . Kelly,
M. (2021). Limited plasticity in thermally tolerant ectotherm populations: evidence for a
trade-off. *Proceedings of the Royal Society B, 288*(1958), 20210765.

Beaman, J. E., White, C. R., & Seebacher, F. (2016). Evolution of plasticity: Mechanistic link
between development and reversible acclimation. *Trends in Ecology & Evolution, 31*(3),
237-249.

Comte, L., & Olden, J. D. (2017). Evolutionary and environmental determinants of freshwater
fish thermal tolerance and plasticity. *Global Change Biology, 23*(2), 728-736.

Deery, S. W., Rej, J. E., Haro, D., & Gunderson, A. R. (2021). Heat hardening in a pair of *Anolis*
lizards: constraints, dynamics and ecological consequences. *Journal of Experimental*
Biology, 224(7), jeb240994.

Ghalambor, C. K., Hoke, K. L., Ruell, E. W., Fischer, E. K., Reznick, D. N., & Hughes, K. A. (2015).
Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in
nature. *Nature, 525*(7569), 372-375.

Gilbert, A. L., & Miles, D. B. (2019). Antagonistic responses of exposure to sublethal
temperatures: adaptive phenotypic plasticity coincides with a reduction in organismal
performance. *The American Naturalist, 194*(3), 344-355.

Gunderson, A. R., Dillon, M. E., & Stillman, J. H. (2017). Estimating the benefits of plasticity in
ectotherm heat tolerance under natural thermal variability. *Functional Ecology, 31*(8),
1529-1539.

Gunderson, A. R., & Revell, L. J. (2022). Testing for genetic assimilation with phylogenetic
comparative analysis: Conceptual, methodological, and statistical considerations.
Evolution, 76, 1942-1952.

Hertz, P. E., Huey, R. B., & Stevenson, R. (1993). Evaluating temperature regulation by field-
active ectotherms: the fallacy of the inappropriate question. *The American Naturalist,*
142, 796-818.

Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A., Jess, M., & Williams, S. E. (2012).
Predicting organismal vulnerability to climate warming: roles of behaviour, physiology
and adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences,*
367(1596), 1665-1679.

Jackson, D. A., & Somers, K. M. (1991). The spectre of 'spurious' correlations. *Oecologia, 86*(1),
147-151.

Kelly, C., & Price, T. D. (2005). Correcting for regression to the mean in behavior and ecology.
The American Naturalist, 166(6), 700-707.

Kelly, M. (2019). Adaptation to climate change through genetic accommodation and
assimilation of plastic phenotypes. *Philosophical Transactions of the Royal Society B,*
374(1768), 20180176.

Morgan, R., Finnøen, M. H., & Jutfelt, F. (2018). CT max is repeatable and doesn't reduce
growth in zebrafish. *Scientific Reports, 8*(1), 1-8.

Morley, S., Peck, L., Sunday, J., Heiser, S., & Bates, A. (2019). Physiological acclimation and
persistence of ectothermic species under extreme heat events. *Global Ecology and*
Biogeography, 28, 1018-1037.
Mottola, G., López, M. E., Vasemägi, A., Nikinmaa, M., & Anttila, K. (2022). Are you ready for the
heat? Phenotypic plasticity versus adaptation of heat tolerance in three-spined
stickleback. *Ecosphere, 13*(4), e4015.
Nguyen, A. D., Brown, M., Zitnay, J., Cahan, S. H., Gotelli, N. J., Arnett, A., & Ellison, A. M.
(2019). Trade-offs in cold resistance at the northern range edge of the common
woodland ant *Aphaenogaster picea* (Formicidae). *The American Naturalist, 194*(6), E151-
E163.
Noh, S., Everman, E. R., Berger, C. M., & Morgan, T. J. (2017). Seasonal variation in basal and
plastic cold tolerance: Adaptation is influenced by both long-and short-term phenotypic
plasticity. *Ecology and Evolution, 7*(14), 5248-5257.
O'Neill, E., Davis, H. E., & MacMillan, H. A. (2021). A lack of repeatability creates the illusion of a
trade-off between basal and plastic cold tolerance. *Proceedings of the Royal Society B,*
288(1964), 20212121.
Phillips, B. L., Munoz, M. M., Hatcher, A., Macdonald, S. L., Llewelyn, J., Lucy, V., & Moritz, C.
(2016). Heat hardening in a tropical lizard: geographic variation explained by the
predictability and variance in environmental temperatures. *Functional Ecology, 30*(7),
1161-1168.
Pottier, P., Burke, S., Drobniak, S. M., Lagisz, M., & Nakagawa, S. (2021). Sexual (in) equality? A
meta-analysis of sex differences in thermal acclimation capacity across ectotherms.
Functional Ecology, 35, 2663-2678.
R Development Core Team. (2021). R: a Language and Environment for Statistical Computing.
Vienna: The R Foundation for Statistical Computing.
Rohr, J. R., Civitello, D. J., Cohen, J. M., Roznik, E. A., Sinervo, B., & Dell, A. I. (2018). The
complex drivers of thermal acclimation and breadth in ectotherms. *Ecology Letters,*
21(9), 1425-1439.
Sasaki, M. C., & Dam, H. G. (2019). Integrating patterns of thermal tolerance and phenotypic
plasticity with population genetics to improve understanding of vulnerability to warming
in a widespread copepod. *Global Change Biology, 25*(12), 4147-4164.
Sasaki, M. C., & Dam, H. G. (2020). Genetic differentiation underlies seasonal variation in
thermal tolerance, body size, and plasticity in a short-lived copepod. *Ecology and*
Evolution, 10(21), 12200-12210.
Sasaki, M. C., & Dam, H. G. (2021). Negative relationship between thermal tolerance and
plasticity in tolerance emerges during experimental evolution in a widespread marine
invertebrate. *Evolutionary Applications, 14*(8), 2114-2123.
Seebacher, F., White, C. R., & Franklin, C. E. (2015). Physiological plasticity increases resilience
of ectothermic animals to climate change. *Nature Climate Change, 5*(1), 61-66.
Sgro, C. M., Overgaard, J., Kristensen, T. N., Mitchell, K. A., Cockerell, F. E., & Hoffmann, A. A.
(2010). A comprehensive assessment of geographic variation in heat tolerance and
hardening capacity in populations of *Drosophila melanogaster* from eastern Australia.
Journal of Evolutionary Biology, 23(11), 2484-2493.

Somero, G. (2010). The physiology of climate change: how potentials for acclimatization and
genetic adaptation will determine 'winners' and 'losers'. *The Journal of Experimental*
Biology, 213(6), 912-920.
Stillman, J. H. (2003). Acclimation capacity underlies susceptibility to climate change. *Science,*
301(5629), 65-65.
van Heerwaarden, B., & Kellermann, V. (2020). Does plasticity trade off with basal heat
tolerance? *Trends in Ecology & Evolution, 35*, 874-885.
van Heerwaarden, B., Kellermann, V., & Sgrò, C. M. (2016). Limited scope for plasticity to
increase upper thermal limits. *Functional Ecology, 30*, 1947-1956.
van Heerwaarden, B., Lee, R. F. H., Overgaard, J., & Sgro, C. M. (2014). No patterns in thermal
plasticity along a latitudinal gradient in *Drosophila simulans* from eastern Australia.
Journal of Evolutionary Biology, 27(11), 2541-2553.
278