Multiomic Profiling of Human Clonal Hematopoiesis Reveals Genotype and Cell-Specific

Inflammatory Pathway Activation

Authors:

J. Brett Heimlich*[1], Pawan Bhat*[2], Alyssa C. Parker [2], Matthew T. Jenkins[2], Caitlyn Vlasschaert[3],

Jessica Ulloa[4], Joseph C. Van Amburg[4], Chad R. Potts[5], Sydney Olson[4], Alexander J. Silver [2],

Ayesha Ahmad[4], Brian Sharber [4], Donovan Brown[5], Ningning Hu[4], Peter van Galen[6,7], Michael

R. Savona[5,8], Alexander G. Bick[+4], P. Brent Ferrell[+5]

Affiliations:

[1]Division of Cardiovascular Medicine, Department of Medicine, Vanderbilt University Medical

Center, Nashville, TN 37232

[2]Vanderbilt University School of Medicine, Nashville, TN 37232

[3]Department of Medicine, Queen's University, Kingston Ontario, Canada

[4]Division of Genomic Medicine, Department of Medicine, Vanderbilt University Medical Center,

Nashville, TN 37232

[5]Division of Hematology and Oncology, Department of Medicine, Vanderbilt University Medical

Center, Nashville, TN 37232

[6]Division of Hematology, Department of Medicine, Brigham and Women's Hospital, Boston, MA

02115 USA

[7]Ludwig Center at Harvard, Harvard Medical School, Boston, MA 02115, USA

[8]Vanderbilt-Ingram Cancer Center, Program in Cancer Biology, and Center for Immunobiology

Nashville, TN 37232

[+]Correspondence to Dr. Bick (alexander.bick@vumc.org) & Dr. Ferrell (brent.ferrell@vumc.org)

*Denotes equal contribution

Key points:

- CD14+ monocytes from clonal hematopoiesis patients stimulate inflammation through increased cytokine expression.

- T cells from clonal hematopoiesis are deficient in GIMAP expression, suggesting CH may impair T cell differentiation.

Abstract:

Clonal hematopoiesis (CH) is an age-associated phenomenon that increases risk for hematologic malignancy and cardiovascular disease. CH is thought to enhance disease risk through inflammation in the peripheral blood[1]. Here, we profile peripheral blood gene expression in 66,968 single cells from a cohort of 17 CH patients and 7 controls. Using a novel mitochondrial DNA barcoding approach, we were able to identify and separately compare mutant *TET2* and *DNMT3A* cells to non-mutant counterparts. We discovered the vast majority of mutated cells were in the myeloid compartment. Additionally, patients harboring *DNMT3A* and *TET2* CH mutations possessed a pro-inflammatory profile in CD14+ monocytes through previously unrecognized pathways such as galectin and macrophage Inhibitory Factor (MIF). We also found that T cells from CH patients, though mostly un-mutated, had decreased expression of GTPase of the immunity associated protein (GIMAP) genes, which are critical to T cell development, suggesting that CH may impair T cell function.

Introduction:

With age, hematopoietic stem cells acquire mutations in driver genes such as DNA Methyltransferase 3A (*DNMT3A)* and Tet Methylcytosine Dioxygenase 2 (*TET2),* resulting in a selective advantage and clonal hematopoiesis. CH is a risk factor not only for hematologic

malignancy but also for multiple diseases of aging including cardiovascular disease, kidney

disease, and osteoporosis[1–5]. While many epidemiological analyses consider CH as a single

entity, the literature reveals that significant associations are gene specific. For example, *TET2* is

more strongly associated with a pro-inflammatory disease mechanism across multiple forms of

cardiovascular disease[1,6] while *DNMT3A* CH is associated with heart failure[7,8] and

osteoporosis[5].

Much attention has focused on how CH mutations lead to skewed hematopoiesis in

hematopoietic stem and progenitor cells (HSPC)[9,10]. However, little attention has focused on the

peripheral compartment. Circulating immune cells with CH mutations are morphologically and

immunophenotypically similar to their non-mutated counterparts, making direct comparisons

difficult in primary human tissues. Whether primary cell intrinsic transcriptional changes or

secondary microenvironment effects, or both, drive pathological phenotypes is unknown.

Though transcriptional profiling of single cells has become routine, it remains difficult to extract

genotype and transcriptional data out of the same cell. Since *DNMT3A* and *TET2* CH blood

samples are a mixture of mutated and non-mutated cells, both genotyping and transcriptomic

sequencing modalities are necessary to delineate these cells. Cell-intrinsic consequences may

arise directly from the somatic mutation while extrinsic, indirect consequences may arise from

altered cell-cell interactions or secreted immune effectors. These phenomena could be

distinguished by identifying mutant and non-mutant cells from the same sample. Several

technologies have sought to close this gap by selectively amplifying the mRNA transcriptome

and using this to genotype cells[11–14]. This approach is effective in HSPCs that express *DNMT3A*

and *TET2*; however, genotyping is less efficient in cells that do not express these genes, such

as fully differentiated cells in the peripheral blood[12,13,15].

To overcome this, we combined single-cell RNA-sequencing (scRNA-seq) with cell-specific

mitochondrial DNA barcoding to simultaneously resolve single-cell DNA mutation status for

100% of cells[15]. Our analysis of 66,968 single cells from 17 individuals with *TET2* CH or

DNMT3A CH and 7 age-matched controls finds novel mechanisms of CH-driven inflammation

and enables direct comparison between peripheral CH mutated and wildtype cells across

individuals.

Methods:

Primary patient samples

All patients in this study consented to all study procedures under VUMC institutional review

board approved research protocols (IRBs #210022, #201583) in accordance with the Treaty of

Helsinki. Adult patients able to give consent were recruited from VUMC clinics who had known

CH mutations as a result of clinical evaluation or patients who were at risk of having a CH

mutation. All patients were confirmed to be without active hematological malignancy at the time

of enrollment. Fresh PBMCs were isolated using Ficoll separation. Following low-speed

centrifugation, pelleted cells were resuspended in freezing media (88% FBS and 12% dimethyl

sulfoxide (DMSO)) and placed in liquid nitrogen.

DNA extraction and CH variant calling

All enrolled patients underwent targeted sequencing to evaluate for the presence of CH

mutations. DNA was extracted using Qiagen Mini kits Cat #27104 according to manufacturer's

recommendations. We sequenced samples using a custom capture panel designed to tile

known CH genes, targeting 600x read depth coverage as previously described[16]. Somatic

mutations were called using publicly available methods in workflow description language in

Mutect2 on the Terra Platform (https://terra.bio/). A putative variant list was formulated and then

cross referenced with a list of known CH driver mutations[1]. Variants were then filtered for read

quality including sequencing depth and minimum alternate allele read depth.

scRNAseq library preparation

Cryopreserved PBMCs from CH patients and controls were thawed at 37□°C and washed with

complete RPMI (RPMI + 10%FBS + 1%PS, cRPMI) to remove freezing media. PBMCs from

each sample (500,000 cells) were plated. Cells were pooled following staining with unique

hashtag antibody oligonucleotide-conjugates (HTO) (30 minutes) staining (Biolegend, TotalSeq-

B). Pooled samples were immediately run on a 10x Chromium Controller after preparation with

a 10X Chromium 3' library preparation kit (10X Genomics) to create scRNA-seq libraries.

10X single-cell sequencing and data preparation

Next-generation 150-nt paired-end sequencing was conducted on an Illumina Novaseq6000

using the cDNA libraries produced by the 10X Chromium library (**Supplemental Table6**).

CellRanger Count (10X Genomics) was used to filter low quality reads and align to the GRCh38

reference genome using STAR as described elsewhere[17]. Resulting matrices from the

CellRanger pipeline were then converted into Seurat[18,19] objects. Demultiplexing was performed

using the HTODemux function in the R package Seurat, applying a positive quantile value of

0.99. Cells containing 15% or more reads mapping to the mitochondrial genome were filtered.

Similarly, we filtered cells with less than 250 genes and 500 UMIs respectively. Remaining

doublets were removed with the R package DoubletFinder[20], using the first 10 principal

components and a doublet formation rate of 7.5%[21]. One lane had an abnormally high number

of doublets, so a more stringent filter was applied for that lane using the DoubletFinder metric

pANN of 0.28. Batch correction was performed with the R package Harmony (v0.1.1). R

package Harmony (v0.1.1).

After performing dimensionality reduction with the function RunUMAP from Seurat and

calculating clusters with the function FindClusters (resolution = 0.75), cell type assignment was

performed using ScType (v1.0). Low confidence cell types were annotated manually.

Mitochondrial and ribosomal genes were removed.

Single-cell mitochondrial enrichment

Mitochondrial enrichment of 10x Genomics v3 3' cDNA was performed using primer sequences

as described in Miller, TE et al[22]. Briefly, 10X 3' cDNA was amplified using primers

encompassing the entire mitochondrial genome along with i5 indexes (**Supplemental Table 5)**.

The samples were pooled, mixed, and incubated with 1.0X AMPure XP Beads. Following

incubation, i7 indexes were added. The amplified cDNA library was purified with AMPure XP

beads and eluted in TE Buffer.

Single-cell mitochondrial DNA sequencing, read processing, and variant calling

MT-DNA processing and variant calling were carried out as previously published[15]. Briefly, fastq

files from MT-DNA enrichment were filtered for reads associated with low-frequency cell

barcodes (CB) and trimmed to remove the UMI and CB. Reads were aligned using STAR to

hg38. Next, we used maegatk to call variants across the mitochondrial genome[15,23]. Maegatk

calls mtDNA variants using combined CB from both scRNAseq and MAESTER enrichment for

variants with at least five supporting reads.

Identification and selection of informative single cell mtDNA variants

Using output from Maegatk, an allele frequency matrix was constructed using all possible

variants in the mitochondrial genome. Cell type annotations were transferred using the cell

barcodes from the RNA expression dataset. A table of MT variants was constructed based on

cell type enrichment for individual MT mutations. MT mutations enriched within monocytes and

absent or very low in lymphoid lineage cells were considered candidate markers of CH

mutations. Single-cell DNA sequencing was performed with Tapestri scDNAseq and colony

based single cell DNA sequencing to confirm candidate MT mutations alignment with CH

mutations.

Single-cell DNA sequencing via MissionBio Tapestri

Single-cell DNA sequencing was performed using the Tapestri platform. Cells were stained with

the BioLegend Total-Seq D Heme Oncology Panel and the Human TruStain FcX antibody.

Targeted DNA amplification was carried out using custom designed probes from MissionBio

(**Supplemental Table 8**). The amplified DNA was released from individual oil droplets using

Ampure XP beads. The final product was quantified using a Qubit fluorometer from

ThermoFisher and assessed for quality on an Agilent Bioanalyzer. Samples were pooled prior to

sequencing with a 25% spike-in of PhiX and run on a NovaSeq 6000 S4 flow cell from Illumina

to generate 150 bp paired-end reads. Sequencing was performed at the Vanderbilt

Technologies for Advanced Genomics (VANTAGE) sequencing core.

Pipeline processing and variant filtering for Tapestri single-cell DNA sequencing

Single-cell DNA samples were processed using the Tapestri Pipeline v1.8.4. Adapters were

trimmed and reads were aligned to the hg19 reference genome. Variants were called using

GATK 3.7 and filtered based on quality scores, read depth, and genotype frequency. Informative

variants were annotated, and cells were clustered based on their genotypes.

To annotate cell populations, unsupervised hierarchical clustering was performed on the

antibody-oligo conjugate (AOC) data. Reads were normalized and AOCs with low expression

were removed. Principal component analysis was conducted on the normalized AOCs, and the

first ten principal components were used for UMAP coordinate calculation. The resulting cells

were clustered using a k state of 100, and clusters with noisy AOC expression were eliminated.

The remaining clusters were annotated based on expert knowledge of surface marker

expression.

Differential expression analysis and pathway analysis

Differential gene expression was calculated using a pseudobulk-like approach in which

measurements from groups of similar cells were summed. Cells were separated by genotype

and cell type then clustered using the Python module Metacell-2 (v0.8.0). We followed standard

procedures from the Metacell-2 vignette. We excluded gene modules that had an average

correlation with cell cycling genes of 0.75 or greater. Differential expression was calculated for

genes that had at least 10 transcripts in at least 85% of metacells using the R package DESeq2.

We performed Wald tests of significance with Benjamini-Hochberg multiple testing correction.

Genes that were sex-specific and red-blood-cell-specific were removed. Pathway analysis was

performed on differential expression results using the function gseGO with ont = "ALL",

minGSSize = 50, maxGSSize = 800, nPermSimple = 10000 from the R package clusterProfiler

(v4.8.1).

Cell signaling interactions were predicted from single-cell RNA sequencing data with the R

package CellChat (v1.6.1). Genes with extremely high or low expression were removed with the

parameter trim = 0.1. Comparisons were restricted to cell types that had at least 25 cells with

the parameter min.cells = 25.

To compare mutant and wildtype monocyte cell states, we performed pairwise DGE analysis by

using the MAST method with false-discovery rate (FDR) correction. To reduce transcriptional

noise prior to DGE, we only included genes that were detected in at least 10 cells. We then

applied the Hurdle model from the MAST R package (v.1.24.1) and adjusted for the cellular

detection rate to determine significant differences in gene expression (threshold: absolute value

of the log fold-change coefficients > 0.25, FDR > 0.05).

Phospho-specific flow cytometry

Cryovials of cryopreserved cells from healthy donors and CH patients were thawed and washed

with 10 mL of cRPMI. The cells were stained for viability with AlexaFlour700 (Invitrogen,

cat#P10163) and counted. An aliquot of 500,000 cells were plated in 200 µL of media in a 96-

well plate and stimulated with 20 ng/mL of IL-6 (Peprotech) for 15 minutes. Cells were then fixed

with 1.6% PFA at room temperature and permeabilized with 150 µL of methanol at -80C for at

least 30 minutes. Cells were resuspended in 180 µL of PBS and fluorescence cell barcoding

performed as previously described[24] with serial dilutions of Pacific Blue (LifeTechnologies,

cat#P10163, PB) and Pacific Orange (Invitrogen, cat#P30253, PO) dyes for 30 minutes in the

dark at RT. Two concentrations of PB were prepared (20 and 4 µg/mL), while six concentrations

of PO were prepared (7.00, 2.99, 1.27, 0.54, 0.23, 0.10 µg/mL). Barcoding was quenched with

80 µL of cell staining media. The barcoded cells were then collected into a single tube and

stained with a cocktail of antibodies: CD33 PECy7 (5 µL per 100 µL stain, Biolegend,

cat#303434, clone WM53) and pSTAT3 AlexFlour488 (2.5 µL per 100 µL stain, Biolegend,

cat#651006, clone 13A3-1) for 30 minutes prior to acquisition on a BD 5-laser Fortessa flow

cytometer.

Results:

We used scRNA-seq with cell-specific mitochondrial DNA sequencing to resolve single-cell

genomic DNA mutation status and investigate pathological mechanisms of CH (**Fig. 1A,**

methods). Peripheral blood mononuclear cells (PBMCs) from 8 *TET2*, 9 *DNMT3A* and 7 age-

matched controls (ages 47-89) were selected from a prospective CH observational study which

was designed to capture patients at high risk for CH through a robust referral network (**Fig. 1A-**

B and Table 1).

To trace the effects of CH mutations on peripheral blood cell type proportions, we derived cell

type annotations based on known marker genes (**Fig. 1C-D, Supplemental Fig. 1).** There were

no significant differences in cell type proportions on routine clinical laboratories **(Supplemental**

Table 1). Notably, four patients had multiple CH mutations with concomitant cytopenias without

bone marrow dysplasia, meeting diagnostic criteria for clonal cytopenias of undetermined

significance (CCUS**)**[25].

To annotate mutant and non-mutant cells from the same sample, we combined single-cell

targeted amplicon sequencing (scDNAseq) and 3' RNA mitochondrial lineage tracing in our

TET2 and *DNMT3A* patients (**Fig. 2A**). PBMCs from both *TET2* and *DNMT3A* patients were

processed through the scDNAseq pipeline (Mission Bio) which captures known genomic CHIP

mutations and co-occurring mitochondrial variants. We also profiled the immunophenotype of

the samples by combining scDNAseq with oligo-conjugated antibodies to annotate cell

populations **(Fig 2B, 2E, and Supplemental Fig. 2-B).** In one patient with a known *TET2*

mutation at chr4:106157967 with 51% VAF **(Supplemental Fig 2, Supplemental Table 2)** our

scDNAseq analysis revealed a single mitochondrial variant (MT 7754G>C) that was concordant

with cells harboring the known *TET2* mutation, suggesting common lineage. We found 492 cells

that carried both the *TET2* mutation and the MT 7754G>C variant and 492 cells that carried

neither variant. We excluded a marginal number (n = 15) of cells where only the 7754G>C

variant was detected. The mature myeloid cell compartment was heavily enriched for both the

TET2 mutation and the mitochondrial variant **(Fig. 2C-D, Extended Data Fig. 3A)**. We repeated

the analysis for a sample with a known *DNMT3A* mutation with 24% VAF **(Fig 2F-G,**

Supplemental Fig 2). The concomitant genomic and mitochondrial variants (chr2:25470560

and MT 747A>G) were detected in most myeloid cells and a moderate proportion of

lymphocytes, consistent with previous knowledge regarding *DNMT3A* mutations in

hematopoiesis[26] (**Fig. 2F-G and Extended Data Fig. 3B**). There were 111 *DNMT3A* cells

where the 747A>G variant was not detected, indicating that 747A>G marks a cell population

that is subclonal to *DNMT3A*. We validated the co-occurrence of the CH variant and 747A>G

using primary template amplification of genomic DNA from single cell colonies (**Extended Data**

Fig. 3D, see Methods). Subsequently, we were able to use the MT DNA single nucleotide

variant (SNV) as an identifying 'barcode' for the mutant cell annotation, allowing for the partition

of mutant and wildtype populations within our scRNAseq data (**Fig. 2H-K**). In the corresponding

single- cell RNA dataset evaluating for the *TET2* sample, we found a significant myeloid bias

among cells identified with MT 7754G>C (log2(fold change) = 4.146, FDR = 0.001, **Fig 2H-I**). In

DNMT3A scRNAseq sample, we identified cells with 747A>G, finding a less severe monocytic

skew accounting for the lower relative VAF compared to the *TET2* sample (**Fig. 2J-K**).

We applied our mitochondrial lineage tracing *method* to a total of 4 *TET2* and 2 *DNMT3A*

patients to identify CH clones (**Supplemental Fig. 3).** We first performed differential gene

expression testing (DGE) on CD14+ monocytes comparing CH mutant cells to their wildtype

(WT) counterparts. This resulted in 70 differentially expressed genes in *TET2* mutants whereas

there were zero differentially expressed genes in *DNMT3A* mutants compared to WT

counterparts. We then evaluated mutant *TET2* and *DNMT3A* CD14+ monocytes against

unaffected control CD14+ monocytes. We identified 202 and 122 differentially expressed genes

(DEGs) in *TET2* and *DNMT3A* CD14+ monocytes, respectively (**Supplemental Table 7)**. There

were 12 overlapping DEGs when comparing mutant *TET2* CD14+ monocytes to WT cells and

when comparing to controls (**Supplemental Fig. 3**). Notable among these were inflammatory

mediators *CXCL1, CXCL3, and IL1B* (False Discovery Rate [FDR] < 1 x 10^{-20} for all

comparisons) **(Fig. 3A, Supplemental Table 7**). Top DE genes in mutant *DNMT3A* monocytes

compared to controls included C-C Motif Chemokine Ligand 4 (*CCL4*, FDR = 1.26 x 10^{-4}), C-C

Motif Chemokine Ligand 2 (*CCL2,* FDR = 2.78 x 10^{-12}), and C-C Motif Chemokine Ligand 7

(CCL7, FDR = 1.34 x 10^{-7}) (**Fig. 3B, Supplemental Table 7**). Pathway analysis showed

upregulation of leukocyte activation and cell adhesion in mutant *TET2* monocytes whereas

mutant *DNMT3A* monocytes had enrichment in regulation of cellular death pathways and

leukocyte migration (**Fig. 3C-D**).

Noting the increased expression of IL-1B, a prominent downstream mediator of the IL-6

pathway among mutant *TET2* monocytes, we sought to further evaluate whether signaling along

this axis was a cell intrinsic or cell extrinsic phenomenon. To do this, we employed

phosphospecific flow cytometry to measure response to IL-6 in high VAF *TET2* mutant

(*TET2*hi), low VAF *TET2* mutant (*TET2*lo), and controls. The basal pSTAT3+ monocyte

percentage was significantly higher in *TET2*hi monocytes compared to controls. All samples

showed some response to IL-6, while *TET2*hi monocytes had the highest proportion of

pSTAT3+ cells, significantly higher than both control and *TET2*lo samples, in response to IL-6

stimulation **(Fig. 3E-F)**. Notably, there was a linear increase in the proportion of pSTAT3+ cells

after IL-6 stimulation in accordance with increasing VAF (R = 0.77, p = 0.008), suggesting cell

intrinsic altered signaling among the mutant fraction (**Fig. 3G**).

We then queried whether there were also cell-extrinsic effects of CH mutations in our cohort as

has been recently reported.[27] To determine this, we compared grouped RNA expression profiles

from CD14+ monocytes between *DNMT3A* or *TET2* and control patients since this would

include both mutated and non-mutated cells. To reduce potential for false discovery from high

dropout rates, we partitioned our dataset into metacells[28] prior to performing DGE analysis. The

top DEGs among *TET2* patients included fibronectin 1 (*FN1*, adj p = 2.39 x 10^{-26}) and Fc Epsilon

Receptor II (*FCER2*, adj p = 2.4 x 10^{-14}), which encodes CD23. Both of these are important

components of monocyte adhesion[29,30] (**Fig 4A**). This contrasts with the top DEGs from

DNMT3A CD14+ monocytes which included interferon induced transmembrane protein 2

(*IFITM2*, adj p = 2.37 x 10^{-18}) and adhesion G protein-coupled receptor E5 (*ADGRE5*, adj p =

4.4 x 10^{-14}), genes involved in monocyte adhesion[31] and differentiation[32] (**Fig 4B**). While the

specific genes impacted were different between *TET2* and *DNMT3A* comparisons, the pathways

they converged on were similar. In general, genes related to immune responses and leukocyte

activation were upregulated whereas genes related to transport activity and endoplasmic

reticulum regulation were downregulated (**Fig. C-D).** Similarly, GSEA highlighted convergent

pathways between *TET2* and *DNMT3A* CD14+ monocytes including leukocyte activation,

regulation of leukocyte activation, and regulation of cell activation (**Fig. 2E-F**). Using CellChat[33]

to infer intercellular interactions in our scRNA-seq data, we found CD14+ monocytes from TET

samples exhibited enhanced signaling across IL-1, macrophage migration inhibitory factor

(*MIF*), and Galectin, all parts of the inflammatory signaling axis (**Fig. 2G**). CD14+ monocytes

from *DNMT3A* samples also exhibited enhanced IL-1 and galectin signaling and uniquely had

elevated integrin beta 2 (ITGB2) signaling **(Fig 2H, Supplemental Table 7**).

When evaluating signaling patterns between cell types, we noted increased signaling from both

CH CD14+ monocytes to T cells leading us to investigate the impact of CH on T cells **(Fig. 5A-**

B). We found that genes involved in T cell activation and immune response were highly

expressed in both *TET2* and *DNMT3A* samples compared to controls (**Fig. 5C-D,**

Supplemental Table 7). Each member of the GTPase of the immune associated nucleotide

binding protein (GIMAP) family, which plays a critical role in proper T and B cell

differentiation[34,35] was downregulated in CD4+ T cells and CD8+ T cells in both *TET2* and

DNMT3A samples (**Fig. 5E-F, Supplemental Fig 4**). *GIMAP1* and *GIMAP5*, which both result in

T cell deficiency when knocked out in mice[34], were significantly downregulated (adj p < 0.05,

log2(fold change) < -0.5) in each comparison, except for *GIMAP5* in *TET2* CD8+T cells which

had a p-value of 0.235.

Discussion:

Here, we present transcriptional profiling and characterization of *DNMT3A* and *TET2* CH in

human peripheral blood. By using a novel approach that integrates multimodal single-cell RNA

sequencing with scDNA sequencing to link mitochondrial mutations to somatic nuclear

mutations, we simultaneously resolve DNA mutational status and cell state. Our study revealed

CH mutation specific aberrations in cellular state allowing several conclusions.

First, we identified CD14+ monocytes as drivers of CH-associated inflammation in the

peripheral blood in both *TET2* and *DNMT3A* CH. Specifically, we found *TET2* CH mutant

CD14+ monocytes harbored important differences suggesting cell intrinsic mechanisms are

important to *TET2* phenotypes. In relation to non-mutant wildtype monocytes from *TET2*

patients, mutant CD14+ monocytes exhibited significant differences across important

inflammatory genes including *IL1B*, *CXCL3*, and *CXCL1*, a phenomenon that was not seen in

DNMT3A. Furthermore, intracellular monocyte signaling via STAT3 in response to IL-6 exhibited

a VAF dependent increase further supporting the notion that mutant *TET2* monocytes exhibit

cell intrinsic signaling patterns. These experiments suggest a precision medicine therapeutic for

TET2 CH. A recent analysis of the Canakinumab Anti-inflammatory Thrombosis Outcomes Trial

(CANTOS) found that IL-1B antagonist, Canakinumab[36], reduced cardiovascular risk in *TET2*

but not *DNMT3A* CH patients [2,37]. Our data provides a mechanistic rationale for a genotype-

specific approach to treat CH, a finding only possible with the ability to partition mutant and

wildtype cells from the same sample.

361 Second, collective differences between CD14+ monocytes from both *TET2* and DNMT3A and

362 controls identify novel gene targets and signaling pathways. Computationally-inferred outgoing

363 signaling in monocytes from *TET2* and *DNMT3A* patients indicated a notable increase in *MIF*

364 signaling. *MIF* resides as a pre-formed peptide in a variety of cell types and binds with its

365 receptors CXCR2 as well as CXCR4 to promote the recruitment of monocytes and T cells to

366 sites of tissue injury[38]. Recruitment of hyperinflammatory monocytes has been identified as the

367 initiating event in the development of atherosclerotic plaques[39]. *ADGRE5* which encodes CD97

368 had significantly higher expression among monocytes from *DNMT3A* patients. The protein

369 product of this gene promotes the adhesion and migration to sites of inflammation[40] and has

370 been associated with rheumatoid arthritis[41]. Therefore, MIF and *ADGRE5* may represent novel

371 targets in treating inflammation associated with *TET2* and *DNMT3A* CH.

373 Third, our study clarified the cell-extrinsic effects of CH-mutations in peripheral blood.

374 Comparison between T cells from CH samples and controls highlighted significant effects of CH

375 on both T cell differentiation and T cell activation. We observed consistent downregulation of the

376 GIMAP protein family in CD4+ and CD8+ T cells in both *TET2* and *DNMT3A* samples. Work in

377 mice has established that knockout of GIMAP proteins impairs development of T and B cells,

378 resulting in a relative T/B deficiency and a myeloid skew[34], similar to what is observed in CH.

379 GIMAP proteins are regulated together under the direction of the transcription factors RUNX1,

380 GATA3, and TAL1[42,43]. DNMT3A directly binds to RUNX1 and GATA3[42], and TAL1 expression

381 has been shown to be disrupted by knockout of both *TET2* and *DNMT3A*[44]. Further work

382 investigating the effect of *TET2* and *DNMT3A* mutations on GIMAP expression and subsequent

383 differentiation is warranted.

385 Our study has several limitations. First, while *TET2* and *DNMT3A* mutations make up

386 approximately 2/3 of all CH mutations, CH represents a diverse set of mutations in >70 genes.

These CH mutations are likely to have divergent effects from those we describe here. We also

binned samples with co-occurring mutations additional to *TET2* or *DNMT3A* mutations to

increase our sample set size, though this may introduce a source of variability. Second, we

cannot exclude that CH with small clones below our limit of detection are present in our control

samples. However, we would expect minimal pathological effect given the marginal size of

these clones. Third, a shortcoming of our work is the absence of neutrophils in PBMC samples.

Given the myeloid bias of CH mutations, it is likely that neutrophils also harbor mutations, and

so their functional consequences within the periphery require investigation. Finally, two of our

TET2 samples have co-occurring *SRSF2* mutations, which, as others have noted, are

unfortunately not able to be detected consistently with scDNA-seq. Based on VAF data, these

mutations are subclonal to TET2 in each case, this is due to GC content in the region[45]; thus,

we are unable to assess the effect of this mutation our samples on a single cell level. Future

efforts to improve single cell genotyping of this gene may help elucidate this.

Overall, our study provides mechanistic support for a genotype specific precision medicine

approach for future CH therapeutics.

Data and materials availability: All filtered count matrices and DGE tables are available on

Open Science Framework at https://osf.io/rac5w/. Seurat objects will be made available through

the Chan Zuckerberg Initiative database. All data analysis was completed using R (v4.1.2) on

the Terra.bio cloud platform. All R files used to generate the figures and tables will be made

publicly available on GitHub upon publication and can be made available for reviewers upon

request.

Acknowledgements

We thank Angela Jones for her assistance with sequencing efforts associated with this work.

A.G.B. is supported by a Burroughs Wellcome Foundation Career Award for Medical Scientists

and the NIH Director's Early Independence Award (DP5-OD029586). P.B.F is supported by a

NIH K23HL138291 and a Mark Foundation Endeavor Award. P.v.G. is supported by the Ludwig

Center at Harvard, the NIH (R00CA218832), Gilead Sciences, the Bertarelli Rare Cancers

Fund, the Starr Cancer Consortium, the William Guy Forbeck Research Foundation, and is an

awardee of the Glenn Foundation for Medical Research and American Federation for Aging

Research (AFAR) Grant for Junior Faculty. M.R.S. is supported by NIH 1R01CA262287 and

1U01OH012271, an LLS Clinical Scholar Award, the Biff Ruttenburg Foundation, the Adventure

Alle Fund, the Beverly and George Rawlings Research Directorship, the E.P. Evans MDS

Foundation.

Author contributions

J.B.H. and P.B. designed the study, facilitated data collection, conducted formal analysis and

interpretation of results, generated figures, prepared the original draft and edited the

manuscript.

A.C.P., C.V., and M.T.J. collected data, conducted formal analysis and interpretation of results,

generated figures, prepared the original draft and edited the manuscript.

J.U., C.R.P., S.O., and N.N.H. facilitated sample curation and data collection.

B.S. and A.A. provided analysis software.

P.V.G. provided resources, analysis software, and edited manuscript.

A.J.S. and M.R.S. facilitated sample curation, provided resources and project administration,

and edited the manuscript.

A.G.B. and P.B.F. conceived and supervised the study, provided funding for the study, provided

resources and project administration, conducted formal analysis and interpretation of results,

generated figures, prepared the original draft and edited the manuscript.

Competing Interests Statement

All unrelated to the present work: M.R.S. reports personal fees from AbbVie, BMS, CTI, Sierra

Oncology, Novartis, grants from Astex and Incyte, personal fees and other support from

Karyopharm, Ryvu, personal fees from Sierra Oncology, grants and personal fees from Takeda,

and TG Therapeutics outside the submitted work. P.B.F. reports research funding from Novartis.

A.G.B. is a scientific co-founder and has equity in TenSixteen Bio. All other authors declare that

they have no competing interests.

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to:

Alexander G. Bick

P. Brent Ferrell

Figure legends:

Fig. 1: Single-cell RNA sequencing reveals distinct cell type profiles in clonal hematopoiesis

A) Cells from 17 CH patients and 7 controls were processed via scRNAseq. Mutational lineage tracing was performed to link genomic variants to scRNAseq results. **B)** UpSet plot displaying CH mutations for each patient with accompanying dot plot showing variant allele frequency for each mutation. Color corresponds to mutational group. Dots correspond to mutations, so patients with multiple CH mutations have separate dots for each mutation. **C)** UMAP displaying cell type clusters, as defined by unsupervised clustering. Clusters were annotated using scType. **D)** Radar plot showing cell type proportions for controls, patients with *TET2* mutations, and patients with *DNMT3A* mutations.

Fig. 2: Single-cell DNA and RNA sequencing define cells carrying CH mutations. A) Schematic showing experimental design. **B)** UMAP showing cell types for cells from *TET2* patient CH-21-014. **C)** UMAP showing mutational status for cells from *TET2* patient CH-21-014. **D)** Stacked bar plot quantifying proportion of mutant cells per cell type for cells from *TET2* patient CH-21-014. **E)** UMAP showing cell types for cells from *DNMT3A* patient CH-20-046. **F)** UMAP showing mutational status for cells from *DNMT3A* patient CH-20-046. **G)** Stacked bar plot quantifying proportion of mutant cells per cell type for cells from *DNMT3A* patient CH-20-046. **H)** UMAP showing predicted mutational status for cells from *TET2* patient CH-21-014, based on presence of MT mutation 7754G>C. **I)** Stacked bar plot quantifying proportion of mutant cells per cell type for cells from *TET2* patient CH-21-014. **J)** UMAP showing predicted mutational status for cells from *DNMT3A* patient CH-20-046, based on presence of MT mutation 747A>G. **K)** Stacked bar plot quantifying proportion of mutant cells per cell types for cells from *DNMT3A* patient CH-20-046.

Fig. 3: CD14+ monocyte mutant vs control analysis highlights inflammatory cell-intrinsic

effects of CH mutations. A) Violin plots displaying expression of cytokines in control CD14+

monocytes and *TET2* mutant CD14+ monocytes. **B)** Violin plots displaying expression of

cytokines in control CD14+ monocytes and *DNMT3A* mutant monocytes. **C)** GSEA based on

results from differential expression analysis comparing *TET2* mutant CD14+ monocytes to

control CD14+ monocytes. **D)** GSEA based on results from differential expression analysis

comparing *DNMT3A* mutant CD14+ monocytes to control CD14+ monocytes. **E)** Quantification

of phospho-flow cytometry displaying pSTAT3(Y705)+ in the CD33+ gate of large VAF *TET2*

CH samples (>25%, n = 3) compared to both low VAF (<25%, n = 3) and control samples (n =

4) (** = $p < 0.01$ by ANOVA with Tukey's HSD). **F)** Same as in (E), with IL-6 stimulation

condition. **G)** Pearson correlation of %pSTAT3(Y705)+ in the CD33+ gate and VAF following IL-

6 stimulation. For samples with multiple mutations, the highest VAF value was selected for the

analysis.

Fig. 4: Genotype-grouped CD14+ monocyte vs control comparisons highlight

inflammatory cell-extrinsic effects of CH mutations. A) Violin plots displaying expression of

FN1 and *FCER2* in CD14+ monocytes from controls and CD14+ monocytes from patients with

TET2 mutations. **B)** Violin plots displaying expression of *IFITM2* and *ADGRE5* in CD14+

monocytes from controls and CD14+ monocytes from patients with *DNMT3A* mutations. **C)**

Volcano plot showing results of differential expression analysis comparing CD14+ monocytes

from patients with *TET2* mutations to CD14+ monocytes from controls, colored by biological

pathway. **D)** Volcano plot showing results of differential expression analysis comparing CD14+

monocytes from patients with *DNMT3A* mutations to CD14+ monocytes from controls, colored

by biological pathway. **E)** GSEA based on results from differential expression analysis

comparing CD14+ monocytes from patients with *TET2* mutations to CD14+ monocytes from

controls. **F)** GSEA based on results from differential expression analysis comparing CD14+

monocytes from patients with *DNMT3A* mutations to CD14+ monocytes from controls. **G)**

Heatmap showing predicted outgoing signaling from CD14+ monocytes from patients with *TET2*

mutations and from controls for pathways involved in inflammatory response and immune cell

migration, as determined by CellChat (* indicates p-value < 0.05/7). **H)** Heatmap showing

predicted outgoing signaling from CD14+ monocytes from patients with *DNMT3A* mutations and

from controls for pathways involved in inflammatory response and immune cell migration, as

determined by CellChat (* indicates p-value < 0.05/7).

Fig. 5: Genotype-grouped CD4+ and CD8+ T cell vs control comparisons highlight

alterations to T cell activation and differentiation in CH. A) Circle plot displaying predicted

differential interaction strength between CD14+ monocytes and relevant hematopoietic cells

from *TET2* samples compared to controls, as determined by CellChat. Line thickness

corresponds to differential interaction strength. Green color indicates increased signaling in

TET2 samples compared to controls. Grey indicates decreased signaling. **B)** Circle plot

displaying predicted differential interaction strength between CD14+ monocytes and relevant

hematopoietic cells from *DNMT3A* samples compared to controls, as determined by CellChat.

Line thickness corresponds to differential interaction strength. Red color indicates increased

signaling in *DNMT3A* samples compared to controls. Grey indicates decreased signaling. **C)**

Volcano plot showing results of differential expression analysis comparing CD4+ T cells from

patients with *TET2* (left) or *DNMT3A* (right) mutations to CD4+ T cells from controls, colored by

biological pathway. **D)** Volcano plot showing results of differential expression analysis

comparing CD8+ T cells from patients with *TET2* (left) or *DNMT3A* (right) mutations to CD8+ T

cells from controls, colored by biological pathway. **E)** Violin plots displaying expression of

GIMAP1 in CD4+ T cells and CD8+ T cells from controls and from patients with CH mutations.

F) Violin plots displaying expression of *GIMAP5* in CD4+ T cells and CD8+ T cells from controls

from patients with CH mutations.

Table 1. Demographic features of CH and control patients. Values are listed by counts and

by percentages for categorical variables and by mean and standard deviation for continuous

variables.

References

1. Jaiswal S, Natarajan P, Silver AJ, et al. Clonal Hematopoiesis and Risk of Atherosclerotic Cardiovascular Disease. *N Engl J Med*. 2017;377(2):111-121. doi:10.1056/NEJMoa1701719

2. Jaiswal S, Fontanillas P, Flannick J, et al. Age-Related Clonal Hematopoiesis Associated with Adverse Outcomes. *N Engl J Med*. 2014;371(26):2488-2498. doi:10.1056/NEJMoa1408617

3. Bick AG, Weinstock JS, Nandakumar SK, et al. Inherited causes of clonal haematopoiesis in 97,691 whole genomes. *Nature*. 2020;586(7831):763-768. doi:10.1038/s41586-020-2819-2

4. Zink F, Stacey SN, Norddahl GL, et al. Clonal hematopoiesis, with and without candidate driver mutations, is common in the elderly. *Blood*. 2017;130(6):742-752. doi:10.1182/blood-2017-02-769869

5. Kim PG, Niroula A, Shkolnik V, et al. *Dnmt3a* -mutated clonal hematopoiesis promotes osteoporosis. *Journal of Experimental Medicine*. 2021;218(12):e20211872. doi:10.1084/jem.20211872

6. Bhattacharya R, Zekavat SM, Haessler J, et al. Clonal Hematopoiesis Is Associated With Higher Risk of Stroke. *Stroke*. 2022;53(3):788-797. doi:10.1161/STROKEAHA.121.037388

7. Abplanalp WT, Cremer S, John D, et al. Clonal Hematopoiesis–Driver DNMT3A Mutations Alter Immune Cells in Heart Failure. *Circulation Research*. 2021;128(2):216-228. doi:10.1161/CIRCRESAHA.120.317104

8. Assmus B, Cremer S, Kirschbaum K, et al. Clonal haematopoiesis in chronic ischaemic heart failure: prognostic role of clone size for DNMT3A- and TET2-driver gene mutations. *European Heart Journal*. 2021;42(3):257-265. doi:10.1093/eurheartj/ehaa845

9. Arends CM, Galan-Sousa J, Hoyer K, et al. Hematopoietic lineage distribution and evolutionary dynamics of clonal hematopoiesis. *Leukemia*. 2018;32(9):1908-1919. doi:10.1038/s41375-018-0047-7

10. Izzo F, Lee SC, Poran A, et al. DNA methylation disruption reshapes the hematopoietic differentiation landscape. *Nat Genet*. 2020;52(4):378-387. doi:10.1038/s41588-020-0595-4

11. Nam AS, Kim KT, Chaligne R, et al. Somatic mutations and cell identity linked by Genotyping of Transcriptomes. *Nature*. 2019;571(7765):355-360. doi:10.1038/s41586-019-1367-0

12. Nam AS, Dusaj N, Izzo F, et al. Single-cell multi-omics of human clonal hematopoiesis reveals that DNMT3A R882 mutations perturb early progenitor states through selective hypomethylation. Published online January 16, 2022:2022.01.14.476225. doi:10.1101/2022.01.14.476225

13. Myers RM, Izzo F, Kottapalli S, et al. Integrated Single-Cell Genotyping and Chromatin Accessibility Charts JAK2V617F Human Hematopoietic Differentiation. Published online May 11, 2022:2022.05.11.491515. doi:10.1101/2022.05.11.491515

14. van Galen P, Hovestadt V, Wadsworth M, et al. Single-cell RNA-seq reveals AML
hierarchies relevant to disease progression and immunity. *Cell.* 2019;176(6):1265-
1281.e24. doi:10.1016/j.cell.2019.01.031

15. Miller TE, Lareau CA, Verga JA, et al. Mitochondrial variant enrichment from high-
throughput single-cell RNA sequencing resolves clonal populations. *Nat Biotechnol.*
Published online February 24, 2022:1-5. doi:10.1038/s41587-022-01210-8

16. Miller PG, Fell GG, Foy BH, et al. Clonal hematopoiesis of indeterminate potential and risk
of death from COVID-19. *Blood.* 2022;140(18):1993-1997. doi:10.1182/blood.2022018052

17. Guess T, Potts CR, Bhat P, et al. Distinct Patterns of Clonal Evolution Drive Myelodysplastic
Syndrome Progression to Secondary Acute Myeloid Leukemia. *Blood Cancer Discovery.*
2022;3(4):316-329. doi:10.1158/2643-3230.BCD-21-0128

18. Stuart T, Butler A, Hoffman P, et al. Comprehensive integration of single-cell data. *Cell.*
2019;177:1888-1902. doi:10.1016/j.cell.2019.05.031

19. Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data.
Cell. 2021;184(13):3573-3587.e29. doi:10.1016/j.cell.2021.04.048

20. *DoubletFinder: DoubletFinder Is a Suite of Tools for Identifying Doublets in Single-Cell RNA*
Sequencing Data.; 2022.

21. 10x Genomics. *Technical Note - Resolving Cell Types as a Function of Read Depth and*
Cell Number.; 2019.
https://assets.ctfassets.net/an68im79xiti/6gDArDPBTOg4IlkYEO2Sis/803be2286bba5ca67f
353e6baf68d276/CG000148_10x_Technical_Note_Resolving_Cell_Types_as_Function_of_
Read_Depth_Cell_Number_RevA.pdf

22. Miller TE, Lareau CA, Verga JA, et al. Mitochondrial variant enrichment from high-
throughput single-cell RNA sequencing resolves clonal populations. *Nat Biotechnol.*
2022;40(7):1030-1034. doi:10.1038/s41587-022-01210-8

23. Lareau CA, Ludwig LS, Muus C, et al. Massively parallel single-cell mitochondrial DNA
genotyping and chromatin profiling. *Nat Biotechnol.* 2021;39(4):451-461.
doi:10.1038/s41587-020-0645-6

24. Earl DC, Ferrell PB, Leelatian N, et al. Discovery of human cell selective effector molecules
using single cell multiplexed activity metabolomics. *Nat Commun.* 2018;9(1):39.
doi:10.1038/s41467-017-02470-8

25. DeZern AE, Malcovati L, Ebert BL. CHIP, CCUS, and Other Acronyms: Definition,
Implications, and Impact on Practice. *Am Soc Clin Oncol Educ Book.* 2019;39:400-410.
doi:10.1200/EDBK_239083

26. Buscarlet M, Provost S, Zada YF, et al. Lineage restriction analyses in CHIP indicate
myeloid bias for TET2 and multipotent stem cell origin for DNMT3A. *Blood.*
2018;132(3):277-280. doi:10.1182/blood-2018-01-829937

27. Abplanalp WT, Schuhmacher B, Cremer S, et al. Cell-intrinsic effects of clonal
hematopoiesis in heart failure. *Nat Cardiovasc Res.* Published online September 4, 2023.
doi:10.1038/s44161-023-00322-x

28. Ben-Kiki O, Bercovich A, Lifshitz A, Tanay A. Metacell-2: a divide-and-conquer metacell
algorithm for scalable scRNA-seq analysis. *Genome Biol.* 2022;23(1):100.
doi:10.1186/s13059-022-02667-1

29. Mosesson MW. The role of fibronectin in monocyte/macrophage function. *Prog Clin Biol*
Res. 1984;154:155-175.

30. Lecoanet-Henchoz S, Plater-Zyberk C, Graber P, et al. Mouse CD23 regulates monocyte
activation through an interaction with the adhesion molecule CD11b/CD18. *Eur J Immunol.*
1997;27(9):2290-2294. doi:10.1002/eji.1830270924

31. Friedman RL, Manly SP, McMahon M, Kerr IM, Stark GR. Transcriptional and
posttranscriptional regulation of interferon-induced gene expression in human cells. *Cell.*
1984;38(3):745-755. doi:10.1016/0092-8674(84)90270-8

32. Eichler W, Hamann J, Aust G. Expression characteristics of the human CD97 antigen.
Tissue Antigens. 1997;50(5):429-438. doi:10.1111/j.1399-0039.1997.tb02897.x

33. Jin S, Guerrero-Juarez CF, Zhang L, et al. Inference and analysis of cell-cell communication
using CellChat. *Nat Commun.* 2021;12(1):1088. doi:10.1038/s41467-021-21246-9

34. Saunders A, Webb LMC, Janas ML, et al. Putative GTPase GIMAP1 is critical for the
development of mature B and T lymphocytes. *Blood.* 2010;115(16):3249-3257.
doi:10.1182/blood-2009-08-237586

35. Nitta T, Nasreen M, Seike T, et al. IAN Family Critically Regulates Survival and
Development of T Lymphocytes. Marrack P, ed. *PLoS Biol.* 2006;4(4):e103.
doi:10.1371/journal.pbio.0040103

36. Svensson EC, Madar A, Campbell CD, et al. TET2-Driven Clonal Hematopoiesis and
Response to Canakinumab: An Exploratory Analysis of the CANTOS Randomized Clinical
Trial. *JAMA Cardiology.* 2022;7(5):521-528. doi:10.1001/jamacardio.2022.0386

37. Genovese G, Kähler AK, Handsaker RE, et al. Clonal Hematopoiesis and Blood-Cancer
Risk Inferred from Blood DNA Sequence. *New England Journal of Medicine.*
2014;371(26):2477-2487. doi:10.1056/NEJMoa1409405

38. Bernhagen J, Krohn R, Lue H, et al. MIF is a noncognate ligand of CXC chemokine
receptors in inflammatory and atherogenic cell recruitment. *Nat Med.* 2007;13(5):587-596.
doi:10.1038/nm1567

39. Pan JH, Sukhova GK, Yang JT, et al. Macrophage Migration Inhibitory Factor Deficiency
Impairs Atherosclerosis in Low-Density Lipoprotein Receptor-Deficient Mice. *Circulation.*
2004;109(25):3149-3153. doi:10.1161/01.CIR.0000134704.84454.D2

40. Leemans JC, Te Velde AA, Florquin S, et al. The Epidermal Growth Factor-Seven
Transmembrane (EGF-TM7) Receptor CD97 Is Required for Neutrophil Migration and Host

Defense. *The Journal of Immunology*. 2004;172(2):1125-1131.
doi:10.4049/jimmunol.172.2.1125

41. Hamann J, Wishaupt JO, van Lier RA, Smeets TJ, Breedveld FC, Tak PP. Expression of the
activation antigen CD97 and its ligand CD55 in rheumatoid synovial tissue. *Arthritis Rheum*.
1999;42(4):650-658. doi:10.1002/1529-0131(199904)42:4<650::AID-ANR7>3.0.CO;2-S

42. Liau WS, Tan SH, Ngoc PCT, et al. Aberrant activation of the GIMAP enhancer by
oncogenic transcription factors in T-cell acute lymphoblastic leukemia. *Leukemia*.
2017;31(8):1798-1807. doi:10.1038/leu.2016.392

43. Shiao YM, Chang YH, Liu YM, et al. Dysregulation of GIMAP genes in non-small cell lung
cancer. *Lung Cancer*. 2008;62(3):287-294. doi:10.1016/j.lungcan.2008.03.021

44. Izzo F, Lee SC, Poran A, et al. DNA methylation disruption reshapes the hematopoietic
differentiation landscape. *Nat Genet*. 2020;52(4):378-387. doi:10.1038/s41588-020-0595-4

45. Taylor J, Mi X, North K, et al. Single-cell genomics reveals the genetic and molecular bases
for escape from mutational epistasis in myeloid neoplasms. *Blood*. 2020;136(13):1477-
1486. doi:10.1182/blood.2020006868

Figure 1

Figure 2

a.

Single Cell DNA

Single Cell Genomic DNA sequencing

ID Co-occurring Mitochondrial DNA SNVs and CH mutations

Single Cell 3 'RNA

Single Cell 3' RNA and Mitochondrial RNA Sequencing

Use Mitochondrial DNA SNVs to ID CHIP cells in scRNAseq

b.
Cell types in TET2

- CD14+ Monocytes
- NK cells
- CD4+ T cells
- CD8+ T cells

c.
Mutations in TET2

- mTET2+7754G>C
- 7754G>C
- mDNMT3A+747A>G
- mDNMT3A
- WT

d.
TET2 Cell Type Proportions (DNA)

e.
Cell types in DNMT3A

f.
Mutations in DNMT3A

g.
DNMT3A Cell Type Proportions (DNA)

h.
TET2 indicated by 7754G>C

Mutant status
- Mutant
- Wildtype

i.
TET2 Cell Type Proportions indicated by 7754G>C

TET2 Mutant   Wildtype

j.
DNMT3A indicated by 747A>G

Mutant status
- Mutant
- Wildtype

k.
DNMT3A Cell Type Proportions indicated by 747A>G

DNMT3A Mutant   Wildtype

Figure 3

Figure 4

Figure 5

Table 1

	Control (N=7)	DNMT3A (N=9)	TET2 (N=8)
Sex			
Female	4 (57.1%)	2 (22.2%)	2 (25.0%)
Male	3 (42.9%)	7 (77.8%)	6 (75.0%)
Race			
White	7 (100%)	9 (100%)	6 (75.0%)
Asian	0 (0%)	0 (0%)	1 (12.5%)
Missing	0 (0%)	0 (0%)	1 (12.5%)
Age			
Mean (SD)	75 (14)	67 (15)	75 (8.8)
Missing	1 (14.3%)	0 (0%)	0 (0%)
Hx tobacco use	3 (42.9%)	5 (55.6%)	3 (37.5%)
Hx CAD	3 (42.9%)	3 (33.3%)	2 (25.0%)
Hx CHF	1 (14.3%)	1 (11.1%)	1 (12.5%)
Systolic BP			
Mean (SD)	130 (14)	120 (16)	130 (11)
Diastolic BP			
Mean (SD)	73 (16)	70 (9.1)	74 (11)
Heart rate			
Mean (SD)	70 (11)	78 (14)	68 (9.4)
Missing	0 (0%)	1 (11.1%)	0 (0%)