Prior Knowledge Biases the Perception of Body Postures

Qiu Han[*], Marco Gandolfo, Marius V. Peelen

Donders Institute for Brain, Cognition and Behaviour, Radboud University,
6525 HR, Nijmegen, The Netherlands

*Correspondence: qiu.han@donders.ru.nl

One-sentence summary:
In two perceptual judgment tasks, body postures were misremembered as lower, reflecting knowledge of gravity, and as more similar to possible postures, reflecting knowledge of biomechanical constraints.

Abstract

Body postures provide information about others' actions, intentions, and emotional states. However, little is known about how postures are represented in the brain's visual system. Considering our extensive visuomotor experience with body postures, we hypothesized that priors derived from this experience may systematically bias visual body posture representations. We examined two priors: gravity and biomechanical constraints. Gravity pushes lifted body parts downwards, while biomechanical constraints limit the range of possible postures (e.g., an arm raised far behind the head cannot go down further). We probed participants' (N = 246) memory of briefly presented postures using change discrimination and adjustment tasks. Results showed that lifted arms were misremembered as lower and as more similar to biomechanically plausible postures. Inverting the body stimuli eliminated both biases, implicating holistic body processing. Together, these findings show that knowledge shapes body posture representations, reflecting modulation from a combination of category-general and category-specific priors.

Keywords: Body; Postures; Prior; Bias; Bayesian theory; Expectation

Statement of relevance

The human visual system has built up prior knowledge about the environment that shapes our perception and memory. This knowledge includes general physical rules of the world, for example, gravity and motion laws, as well as specific properties of one category of objects. For example, we know that body parts can only move within certain ranges. In this study, we show that visual body representation is biased by both the general knowledge of gravity and the specific knowledge of biomechanical constraints: Lifted arms were remembered as lower and more similar to biomechanically possible postures. These effects suggest that our brain integrates priors of both the physical world and the specific object category to inform the processing of current stimuli.

Introduction

Body posture is an important social cue that provides information about others' emotions, intentions, and mental states. The pressure to quickly and accurately recognize bodies and their movements has resulted in humans' typically excellent performance in detecting and discriminating body posture and body motion (Neri et al., 1998; Reed et al., 2003; Stein et al., 2012; Thorat & Peelen, 2022), a skill that is supported by dedicated brain regions in visual cortex (Peelen & Downing, 2007). Owning a body ourselves, we also have extensive motor, tactile, and proprioceptive experience of a body and its dynamics (Berlucchi & Aglioti, 2010). Together, these sensory modalities provide us with rich knowledge of hierarchical limb structure, the possible range of movements of joints, and the effort required for executing specific body actions. For example, we know that lifting an arm requires more effort than lowering it; and that reaching the chest by hand is easier than reaching the back. In this study, we asked whether and how our knowledge of the body influences the perceptual representation of body postures.

Previous research has shown that perception is influenced by knowledge and expectations (Bruner & Postman, 1949; de Lange et al., 2018). Specifically, Bayesian accounts of perception propose that regularities of the environment are used to build an internal model of the world, which then informs perceptual inference (Pouget et al., 2013). These priors include the distribution of visual properties in natural scenes (Girshick et al., 2011; Weiss et al., 2002), basic physical principles of motion (Freyd & Finke, 1984), general gravity (Hubbard, 1990), and physical state (Hafri et al., 2022). Effects of priors typically manifest themselves by biasing the interpretation of new input in the direction of the prior, particularly when the input is ambiguous. For example, the last memory of a moving object is biased toward the motion direction, reflecting the prior that moving objects normally keep moving (Freyd & Finke, 1984). This representational momentum is larger for downward motion than upward motion, consistent with the omnipresent downward force of gravity (Hubbard, 1990). These results raise the possibility that body-specific priors similarly influence how we perceive and remember body postures.

We considered two priors that are relevant when constructing an internal model of the body. The first is the general force of gravity: We know that an object will fall if not supported, and likewise, a lifted arm will fall if no muscle force is used. If gravity knowledge affects body representations, we predict that a lifted arm will be represented as slightly lower than its actual position, as has been observed for unsupported objects (Bertamini, 1993; Freyd et al., 1988).

Second, we consider priors related to the biomechanical constraints that are specific to the body. Because of the physiological structure of the body, postures are confined to a limited range (Shiffrar & Freyd, 1990). For example, the elbow can bend inwards but not outwards. If knowledge of these constraints informs perception, the representation of a (nearly) impossible posture may be biased towards the nearest possible posture. Crucially, biomechanical constraints can counteract the effect of gravity: an arm raised in front of the head will fall but an arm raised behind the head can hardly fall lower (Fig. 1).

To test these hypotheses, we used two different perceptual judgment tasks involving four arm

postures subject to one or both of the two biases (Fig. 1). We predicted that lifted arms will generally be remembered as lower, towards the ground, reflecting a gravity-related bias. Furthermore, we predicted that lifted arms will be biased towards biomechanically possible postures. Specifically, biomechanical constraints limit further movement of the arm when the arm is raised behind the head, counteracting the gravity bias (Fig. 1b), and strengthening the gravity bias when the arm is behind the hip (Fig. 1d).

Fig. 1. Illustration of the hypothesis. The orange and blue arrows indicate the direction of the gravity and the biomechanical constraints, respectively; transparent arms indicate the predicted perceived arm positions according the two hypotheses. Here the arrows and the predicted arms only suggest the direction but not the extent of the effects. For the Upper-back posture (b), gravity and the biomechanical constraints point in opposite direction, potentially eliminating each other's influence. For the lower-back posture (d), gravity and the biomechanical constraints go in the same direction, their effects potentially adding up.

Open Practices Statement

All the codes, stimuli, and data related to the experiments reported here are available at https://osf.io/qmtkw/.

Experiment 1

We designed a change discrimination task to probe the existence of biases in body posture representation due to gravity and biomechanical constraints. Participants compared two sequential postures whose arm positions slightly differed, with the second arm posture being slightly higher or lower (Fig. 2a). In the absence of biases, participants should detect upward and downward changes equally well. Instead, if biases distort the representation of the first posture during the brief interval, we may observe that detecting a change in one direction is easier than a change in the other.

Method

Participants

The desired sample size was set to 60 for all experiments before testing. This was determined by a power analysis, revealing that a sample size of 52 was needed to detect an effect size of $d = 0.4$ with 80% power, as suggested recently (Brysbaert, 2019). We rounded this recommendation up to 60 to ensure enough power.

Participants were recruited through the SONA system in return for course credits. We required the subjects to be above 18 years old with normal or corrected-to-normal vision. Recruitment stopped when the sample size reached 60 after the exclusion of low-quality data (see Analysis section). Experiment 1 included two separate experiments for arm-upper postures (Experiment 1a) and arm-lower postures (Experiment 1b), recruiting 75 and 67 subjects, respectively. Out of 74/65 completed data, 14/5 were excluded, resulting a sample size of 60 for both Experiment 1a (49 females, 11 males; age: M = 20.1, range = [18, 36]) and Experiment 1b (48 females, 12 males, 2 other; age: M = 20.6, range = [18, 47]). For this and the following experiments, digital informed consent was obtained from all subjects. The procedures were approved by the University's Ethical committee (Ethics no.: ECSW2017-2306).

Stimuli

Body images were generated by rendering human models in DAZ studio 4.15 (Daz Productions, Inc). A female character and a male character were used. The characters were standing in profile, one arm lifted, the other leaning naturally on the hip. The lifted arm positions were categorized into four quadrants (Fig. 1): two directions (front and back) x two arm heights (upper and lower). In each quadrant, we designated the upper bound of that quadrant as the zero point, with larger angles meaning that the arm is lower, i.e., closer to the feet. The arm could be presented at an angle of 36, 39, 42, 45, 48, 51, or 54 degrees in each quadrant. Both left-facing and right-facing figures were generated to avoid possible confounds of visual hemifield, so that an arm in the front of the body could be equiprobably on both the left and right visual field. The lifted arm was always on the viewer's side (right arm lifted when facing right, left arm lifted when facing left) to avoid the arm being occluded by other body parts.

Mask images were grey-scale checkerboard images. Body images were 300 pixels wide, 480 pixels high. Size in degree depended on the online participant's screen resolution and the eye-to-screen distance. Mask images were 350 pixels wide, 525 pixels high. Masks were presented slightly larger than the body to achieve a better masking effect.

Procedure

Experimental procedures were programmed with JsPsych library (de Leeuw, 2015) and the psychophysics plugin (Kuroki, 2021). Experiments 1a and 1b tested the front and the back arm directions for upper postures (Experiment 1a) and lower postures (Experiment 1b). Data were aggregated for analysis. Experiment 1a also included a machine condition which was not relevant to the purpose of the current study (see Supplementary material 2).

In each trial, a fixation cross was first presented at the center for 800, 900, or 1000 ms, then a target body posture of either 36, 39, 42, 45, 48, 51, or 54 degrees in either quadrant was shown for 200 ms (Fig. 2a). Participants were instructed to remember the posture of the target and hold it in memory. Immediately after the target, a 1000-ms dynamic mask consisting of four consecutive checkerboard images (250 ms each) was shown to minimize aftereffects and/or apparent motion of the arm, after which the probe image appeared for 200 ms. Compared to the target, the arm in the probe would move upwards or downwards by an angle of 3, 6, or 9 degrees (equiprobable). The task was to judge whether the arm had moved up or down relative to the target. Participants indicated their choice by pressing F or J on the keyboard. The key-response mapping was counterbalanced across participants. Participants were asked to respond as accurately and as quickly as possible. The trial ended upon response or 4000 ms after the probe had disappeared.

Each combination of arm direction and angle difference included 24 trials, resulting in 288 trials in total, separated into four blocks. Angle difference, arm direction, and figure gender were completely interleaved while facing direction was kept identical within blocks to avoid extra effort for switching viewpoint between trials. The two left-facing and two right-facing blocks were in ABBA order, with about half of the subjects starting with a left-facing block and the others starting with a right-facing block. A practice session of 12 trials was delivered before the formal experiment. Feedback on the accuracy and mean response time across conditions were shown to the participant at the end of each block.

a. Experiment 1

| 800/ 900/ 1000 ms | 200 ms | 250 ms x 4 | 200 ms | Until response |

Target · Probe

+9
+6
+3
−3
−6
-9 Or

b. Experiments 2&3

| 800/ 900/ 1000 ms | 200 ms | 250 ms x 4 | Until keypress | Adjust | Confirm |

Target · Test

Fig. 2. Trial Procedures. a) Change discrimination task used in Experiments 1a and 1b. In the trial shown here, the target is 45 degrees in the Upper-front, and the probe moves -9 degrees (i.e., upwards). Participants indicated whether the arm had moved up or down. The up-down text screen is shown for illustration purposes. b) Adjustment task used in Experiments 2 and 3. The target posture was either 36, 39, 42, 48, 51, or 54 degrees within each quadrant. The starting posture in the test image was chosen randomly from 30 to 60 from the same quadrant as the target. Participants adjusted the arm to match the target angle, indicated by transparent arms, using the up-arrow and down-arrow keys.

Analysis

Responses with RTs < 250 ms (relative to probe onset) were excluded, as these most likely reflect anticipatory responses. For each participant, data quality was inspected by plotting the percentage of up responses for each angle difference level in each quadrant (Fig. S1a). Participants following task instructions should show an increase in the percentage of up responses as the angle difference decreases. That is, the more obvious that the arm moves up, the more likely people choose up, yielding a sigmoid curve. Some participants exhibited a flat or reversed curve, suggesting that they misunderstood the task or pressed randomly. These participants were detected using a slope index (Fig. S1b) of the difference between the mean up response percentage of the two most obvious moving-up levels (-9 and -6) and the mean of the two most obvious moving-down levels (9 and 6). Participants with a slope index below 0.2 in either the front or the back condition were excluded, resulting in 14 exclusions in Experiment 1a and 5 exclusions in Experiment 1b.

The perceptual biases of interest were quantified by the criterion (c) from signal detection theory, using the Psycho package in R (Makowski, 2018). Taking upward movement as the signal, a negative criterion means the subject was biased in favour of choosing up over down, suggesting upward changes were more noticeable. We calculated individual criterion in each condition from the hit rate (up response percentage when the arm actually moved up) and false alarm rate (up response percentage when the arm actually moved down):

$$C = -(z(\text{hit}) + z(\text{false alarm})) / 2$$

The criteria for all subjects were tested against zero using two-tailed t-tests. A comparison of the four postures was made using a mixed ANOVA with arm height (upper, Experiment 1a; versus lower, Experiment 1b) as a between-subject factor and arm direction as a within-subject factor. Results of d prime were also analyzed and presented in Supplementary material 1.

Results

Participants' responses were in line with a gravity bias (Fig. 3): The arm in the target posture was remembered as lower than its actual position, as indexed by a criterion that was significantly below zero for all postures (Upper-front: M = -0.25, 95% CI = [-0.32, -0.18], $t(59)$ = -7.08, $p < .001$, d = -0.91, BF_{10} = 5.75E6; Upper-back: M = -0.144, 95% CI = [-0.21, -0.07], $t(59)$ = -4.38, $p < .001$, d = -0.57, BF_{10}= 413; Lower-back: M = -0.11, 95% CI = [-0.19, -0.04], $t(59)$ = -2.96, p = .004, d = -0.38, BF_{10} = 7.14) except the Lower-front: M = -0.06, 95% CI = [-0.14, 0.02], $t(59)$ = -1.60, p = .114, d = -0.21, BF_{10} = 0.47.

Next, we combined Experiments 1a and 1b using a mixed ANOVA with arm height and arm direction as factors to test the presence of a biomechanical bias. As illustrated in Fig. 1, compared to the front, the bias in the back should be diminished by biomechanical constraints when in the upper quadrant (Fig. 1a vs. 1b), but strengthened when in the lower quadrant (Fig. 1c vs. 1d). We thus predicted an interaction between arm height and arm direction. We indeed found this interaction (Fig. 3): $F(1, 118)$ = 8.09, p = .005, η^2_p = 0.64. Specifically, for the upper postures, the gravity bias was stronger in the front than in the back, M = -0.108, 95% CI = [-0.03, -0.19], $t(59)$ = -2.74, p = .008, d = -0.35, BF_{10} = 4.17, indicating that the upward biomechanical constraint counteracted the gravity

bias. As predicted, the gravity bias for the Lower-front was numerically weaker than the gravity bias for the Lower-back, but this difference did not reach significance: M = 0.05, 95% CI = [-0.03, 0.13], $t(59) = 1.3$, $p = .197$, $d = 0.17$, $BF_{10} = 0.32$.

Fig. 3. Criterion results for the four conditions in Experiment 1.

***: $p < .001$, **: $p < .01$, *: $p < .05$, n.s.: not significant. Error bars denote 95% CI.

Experiment 2

The change discrimination task provided evidence for both gravity and biomechanical biases. We wondered whether these results would be specific to the change detection task, in which the two consecutive body postures may be perceived as part of an action. If so, the results could reflect biases in human action perception rather than biases in the static representation of the target posture. To address this, in Experiment 2 we tested whether the identified biases replicate in an adjustment task, in which participants reproduced the target posture.

Method

Participants
Experiment 2 adopted a within-subject design. The sample size was kept consistent with Experiment 1. 60 subjects (30 females, 30 males; age: M = 33.42, range = [21, 45]) were recruited through Prolific with monetary reward.

Procedure
Experiment 2 included the same four conditions as in Experiment 1. In each quadrant, six target angles (36, 39, 42, 48, 51, and 54) were used. Consistent with Experiment 1, a fixation and then the target posture was shown, followed by the mask. After the mask disappeared, the subjects were

instructed to press one of the left-arrow or right-arrow keys to show the test image for adjustment. The initial posture of the test was randomized between 30 and 60 degrees, but always in the same quadrant as the target. Participants then pressed up and down arrow keys to manipulate the arm of the test image to move upwards or downwards. After adjusting the arm to the remembered target position, participants pressed space to confirm their answer. If no response was made, the trial ended after 10 s. If the test image was not initiated within 3 s after the mask, a warning would be given.

All other factors, including figure gender and facing direction, were kept consistent with Experiment 1. Facing direction was blocked, and other factors were interleaved. Each angle in each quadrant was presented eight times, resulting in 48 trials for each quadrant, 192 trials in total. The trials were divided into four blocks, with the order manipulated as in Experiment 1. Two mini practice blocks were completed before the start of the experiment. Feedback on average absolute error was given at the end of each block.

Analysis

Trials in which participants pressed space before adjusting the test image and trials in which the test image was not initiated within 3 s after the mask were discarded. Trials with an absolute error larger than 15 degrees were also discarded. Altogether, this led to the rejection of 5.41% of the total number of trials. No participants were excluded.

The error was defined as the target angle minus the response angle. For example, a target of 48 degrees adjusted as 52 degrees would yield an error of -4 degrees. Negative values indicate that the posture was adjusted to be lower than it was actually shown. Errors of all trials in one quadrant were averaged to get the mean error for each quadrant of each participant. Group mean error data were tested in the same way as the criterion in Experiment 1, except that a repeated-measures ANOVA was used instead of a mixed ANOVA.

For visualization purposes, we computed two indexes that reflect the two hypothesized effects. The gravity bias was given by the overall adjustment error, averaged across the four conditions (Upper-front, Upper-Back, Lower-front, Lower-back). The biomechanical constraint was indexed by the difference between postures with vs without biomechanical constraint, averaged across upper and lower postures (the mean of (Upper-front – Upper-back) and (Lower-back - Lower-front); Fig. 4b).

Results

Fig. 4. Results of Experiment 2. a) Mean error of the four conditions. b) Bias Indexes for individual participants. On the top are the calculation methods for the two indexes. ***: $p < .001$, **: $p < .01$, *: $p < .05$, n.s.: not significant. Error bars denote 95% CI.

In the adjustment task, the direction and magnitude of biases are directly reflected in the direction and magnitude of the adjustment error. A negative error indicates that the target was remembered as lower than its actual position, reflecting a gravity bias. This was the case for all of the postures tested (Fig. 4a, Upper-front: M = -2.54, 95% CI = [-2.97, -2.11], $t(59)$ = -11.8, $p < .001$, d = -1.52, BF_{10} = 1.68E14; Upper-back: M = -1.89, 95% CI = [-2.34, -1.44], $t(59)$ = -8.46, $p < .001$, d = -1.09, BF_{10} = 9.86E8; Lower-back: M = -1.07, 95% CI = [-1.48, -0.66], $t(59)$ = -5.22, $p < .001$, d = -0.67, BF_{10} = 6.52E3) except for the lower-front (M = -0.09, 95% CI = [-0.35, 0.52], $t(59)$ = -0.39, $p = .694$, d = 0.05, BF_{10} = 0.15).

Also consistent with Experiment 1, a two-way repeated-measures ANOVA showed an interaction between arm height and arm direction, revealing the effect of biomechanical constraints, $F(1, 59)$ = 48.1, $p < .001$, η^2_p = 0.45. As in Experiment 1, for the upper postures, the gravity bias was stronger in the front than the back: M = -0.65, 95% CI = [-0.25, -1.05], $t(59)$ = -3.28, $p = .002$, d = -0.42, BF_{10} = 16.5. By contrast, for the lower postures, the gravity bias was stronger in the back than the front: M = -1.15, 95% CI = [-0.79, -1.52], $t(59)$ = -6.32, $p < .001$, d = -0.82, BF_{10} = 3.44E5.

Fig. 4b shows the bias indexes for individual participants. Taking the four postures together, the error caused by gravity was significantly different from zero, M = -1.35, 95% CI = [-1.58, -1.12], $t(59)$ = -11.61, $p < .001$, d = -1.50, BF_{10} = 8.50E13. The overall biomechanical constraint (M = -0.90, 95% CI = [-1.16, -0.64], $t(59)$ = -6.94, $p < .001$, d = -0.90, BF_{10} = 3.37E6, also reflected in the interaction in the ANOVA) was also highly consistent across individuals. These results confirm and extend the results of Experiment 1 using a different task, generalizing the effects to a scenario where no action

or motion is implied.

Experiment 3

A prominent feature of body perception is its susceptibility to inversion. Inversion has been shown to disrupt body and face perception more than other objects, which is believed to reflect the disrupted configural processing of bodies and faces (Reed et al., 2003; Yin, 1969; Gandolfo & Downing, 2020; Stein et al., 2012). In Experiment 3, we tested whether the effects of gravity and biomechanical constraints were contingent on holistic processing of bodies, using the inversion effect. Inverted bodies also serve as an ideal control for upright bodies since they are identical in terms of their low-level visual features.

Method

Participants
66 participants (21 females, 45 males; age: M = 30.04, range = [18, 45]) were recruited through Prolific in return for monetary reward. First, we recruited the intended sample size of 60, however, the number of participants starting with the upright vs inverted condition was not yet balanced when the sample size reached 60, therefore six more were recruited. Data from only the first 60 participants yielded highly similar results. Digital informed consent was obtained.

Procedure
The task was identical to Experiment 2 except for that both upright and inverted conditions were tested. The inverted body images were generated by vertically flipping the upright images. Half of the participants started with the upright condition and half with the inverted. Both upright and inverted conditions contained a left-facing block and a right-facing block, with the order randomized across participants but kept the same for upright and inverted conditions. Within each block, trials of different combinations of arm direction, arm height, within-quadrant angle, and figure identities were interleaved. Because of the inclusion of the inverted condition, the trial number for each angle was halved compared to Experiment 2. In total, Upper-front, Upper-back, Lower-front, and Lower-back all included 24 trials for the upright and 24 for the inverted condition.

Analysis
Data cleaning procedures were identical to Experiment 2, with an exclusion rate of 6.90% of the total number of trials. Adjustment errors for each quadrant were averaged for upright and inverted conditions separately. A three-way repeated-measures ANOVA with arm height, arm direction, and body orientation was conducted. Two bias indexes for upright and inverted conditions were calculated and compared with two-tailed t-tests to test whether inversion diminished one or both of the biases.

The coding of angles for the inverted condition still followed the upright condition (Fig. 1c), i.e. in a body-centered reference frame rather than a spatial reference frame, so that the two body orientations can be directly compared. Negative errors for an upright body indicate that the arm was adjusted as closer to the feet and thus closer to the lower part of the screen. Negative errors for an inverted body similarly indicate that the arm was adjusted as closer to the feet; however, because of

the inversion, this is now closer to the upper part of the screen.

Results

We found a significant three-way interaction between arm direction, arm height and body orientation, $F(1,65) = 7.15$, $p = .009$, $\eta^2_P = .10$, indicating that body orientation modulated the interaction between arm height and arm direction. Inspecting upright and inverted conditions separately (Fig. 5), the interaction of arm height and arm direction was significant for the upright body, replicating results from Experiment 2, $F(1,65) = 24.1$, $p < .001$, $\eta^2_P = .27$. In contrast, the inverted body did not show the interaction between arm height and arm direction, $F(1,65) = 1.91$, $p = .171$, $\eta^2_P = .029$. A main effect of body orientation ($F(1,65) = 46.9$, $p < .001$, $\eta^2_P = .42$) showed that inversion diminished the overall negative adjustment error, indicating a reduced gravity bias (Fig. 5).

The bias indexes provided a more direct description of the inversion effect. As shown in Fig. 5b, inversion significantly disrupted both gravity bias: M = 1.25, 95% CI = [0.88, 1.61], $t(65) = 2.67$, $p < .001$, $d = 0.84$, BF$_{10}$ = 3.52E6 and biomechanical constraints: M = 0.61, 95% CI = [0.15, 1.06], $t(65) = 2.67$, $p = .009$, $d = -0.33$, BF$_{10}$ = 3.54. The inversion effect excluded the possibility that these biases emerge from part-based processing of bodies.

Fig. 5. Results of Experiment 3. a) Mean error of the four conditions. Left: upright, Right: inverted. b) Bias Indexes for individual participants. On top are the calculation methods for the two indexes. ***: p < .001, **: p < .01, *: p < .05, n.s.: not significant. Error bars denote 95% CI.

Discussion

The current results demonstrate that knowledge of gravity and of biomechanical constraints jointly shape visual memory representations of human body postures. In three experiments, these effects were replicated both directly and conceptually. Importantly, the biases were absent when bodies were inverted, ruling out low-level visual confounds and indicating that the biases emerge from whole-body representations. Together, the two tasks we used excluded potential confounds of motion perception, response biases and local visual feature processing, demonstrating a top-down influence on static body representations.

One interpretation of the current findings is that they reflect visuomotor simulation. According to the simulation theory, a "rehearsal" in the motor system is involved in perceiving and predicting others' actions (Kilner et al., 2007; Zentgraf et al., 2011). Motor simulation has been linked to the mirror neuron system - visuomotor neurons that respond to both action execution and action observation (Buccino et al., 2004; Järveläinen et al., 2004). In the current experiments, participants may have memorized the postures partly through motor simulation, embodying the observed postures. A similar interpretation has previously been adopted to account for the finding of smaller representational momentum for biomechanically awkward arm movements (Wilson et al., 2010).

Alternatively, the biases observed here may reflect expectations based on our extensive visual experience of seeing posture transitions in other people. Indeed, the representational momentum effect referred to above is preserved in individuals born without upper limbs, i.e., in the absence of motor experience of arm movements (Vannuscorps & Caramazza, 2016). Furthermore, these individuals do not differ from controls in measures of action observation, action prediction, and mental imagery of hand postures, providing evidence against the simulation theory (Vannuscorps et al., 2012; Vannuscorps & Caramazza, 2016). Knowing whether the effects observed here are also present in individuals born without limbs will give us more insight into the contributions of visual and motor experience.

Irrespective of the visual or motor contribution, our results are well explained by Bayesian theories in which perception is the result of an interplay between sensory input and priors derived from an internal model (de Lange et al., 2018; Ma et al., 2022). This internal model is shaped by environmental statistics (Girshick et al., 2011) and serves to achieve optimal inference (Pouget et al., 2013). When input is more ambiguous, priors will have a stronger influence, such that we perceive what is most likely. In our task, a body posture had to be maintained in visual memory for a brief interval. The fidelity of the sensory information will be reduced during this interval, making the posture representation susceptible to the influence of priors. This account would also naturally explain the current finding that the gravity bias was stronger for the upper arm postures than the lower arm postures (see Fig. 3, 4a, 5a) because an internal model of postures would reflect the probabilities of posture transitions: during most arm-involved actions (e.g., walking, using tools), an arm in the lower-front is equally likely to go up or down, whereas an arm in the upper-front is more likely to go down. Analyzing posture transitions in large video databases would be informative to establish the probability of posture transitions during natural actions, so that these can be more directly linked to the biases observed here.

The current findings also raise new questions about the neural representation of body postures. Neuroimaging research on body perception has provided evidence for multiple cortical areas that are specifically engaged in body perception, including the extrastriate body area (EBA, Downing et al., 2001) and the fusiform body area (FBA, Peelen & Downing, 2005; Schwarzlose et al., 2005). Compared to the more extensively studied fusiform face area (FFA, Kanwisher et al., 1997), little is known about the representational structure of these areas (Downing & Peelen, 2011). It has been shown that the FFA represents faces in a face space centered around the average face, with distances from the center representing the deviation from the mean face (Leopold et al., 2001; Loffler et al., 2005). Based on the current results that knowledge of body structure informs posture perception, the body-selective areas may store an internal model of the body, including its constraints. Given the many combinations of body part postures, it would be advantageous for neurons to be tuned primarily to biomechanically possible postures. Indeed, previous work has shown that body representations in the EBA more strongly represent postures in commonly experienced visual field locations (Chan et al., 2010).

In sum, we show that body posture representation is biased towards the ground and towards biomechanically plausible postures, indicating an influence of both general knowledge of the world (gravity) and specific knowledge of the body. These findings may reflect the influence of an internal model of the body based on environmental statistics. By employing such an encoding scheme, the visual system can efficiently predict upcoming postures, a critical component for humans' ability to read others' actions, intentions, and social interactions (Quadflieg & Koldewyn, 2017).

Acknowledgments

We thank Paul Downing and Eelke Spaak for feedback on earlier versions of the manuscript and the Peelen Lab members for the helpful suggestions on experimental design during lab meetings.

Funding

This project has received funding from the China Scholarship Council (CSC), European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie (grant agreement No. 101033489), and European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 725970).

Author Contributions

Qiu Han: Conceptualization, Methodology, Software, Investigation, Formal Analysis, Validation, Visualization, Writing – Original Draft, Writing – Review & Editing, Funding Acquisition

Marco Gandolfo: Conceptualization, Methodology, Software, Validation, Writing – Review & Editing, Funding Acquisition

Marius Peelen: Conceptualization, Methodology, Validation, Writing – Review & Editing, Funding Acquisition

References

Berlucchi, G., & Aglioti, S. M. (2010). The body in the brain revisited. *Experimental Brain Research*, *200*(1), 25–35. https://doi.org/10.1007/s00221-009-1970-7

Bertamini, M. (1993). Memory for position and dynamic representations. *Memory & Cognition*, *21*(4), 449–457. https://doi.org/10.3758/BF03197176

Bruner, J. S., & Postman, L. (1949). On the Perception of Incongruity: A Paradigm. *Journal of Personality*, *18*(2), 206–223. https://doi.org/10.1111/j.1467-6494.1949.tb01241.x

Brysbaert, M. (2019). How many participants do we have to include in properly powered experiments? A tutorial of power analysis with reference tables. *Journal of Cognition*, *2*(1), Article 1. https://doi.org/10.5334/joc.72

Buccino, G., Binkofski, F., & Riggio, L. (2004). The mirror neuron system and action recognition. *Brain and Language*, *89*(2), 370–376. https://doi.org/10.1016/S0093-934X(03)00356-0

Chan, A. W.-Y., Kravitz, D. J., Truong, S., Arizpe, J., & Baker, C. I. (2010). Cortical representations of bodies and faces are strongest in commonly experienced configurations. *Nature Neuroscience*, *13*(4), 417–418. https://doi.org/10.1038/nn.2502

de Lange, F. P., Heilbron, M., & Kok, P. (2018). How Do Expectations Shape Perception? *Trends in Cognitive Sciences*, *22*(9), 764–779. https://doi.org/10.1016/j.tics.2018.06.002

de Leeuw, J. R. (2015). jsPsych: A JavaScript library for creating behavioral experiments in a Web browser. *Behavior Research Methods*, *47*(1), 1–12. https://doi.org/10.3758/s13428-014-0458-y

Downing, P. E., Jiang, Y., Shuman, M., & Kanwisher, N. (2001). A Cortical Area Selective for Visual Processing of the Human Body. *Science*, *293*(5539), 2470–2473. https://doi.org/10.1126/science.1063414

Downing, P. E., & Peelen, M. V. (2011). The role of occipitotemporal body-selective regions in person perception. *Cognitive Neuroscience*, *2*(3–4), 186–203. https://doi.org/10.1080/17588928.2011.582945

Freyd, J. J., & Finke, R. A. (1984). Representational momentum. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *10*(1), 126–132. https://doi.org/10.1037/0278-7393.10.1.126

Freyd, J. J., Pantzer, T. M., & Cheng, J. L. (1988). Representing statics as forces in equilibrium. *Journal of Experimental Psychology. General*, *117*(4), 395–407. https://doi.org/10.1037//0096-3445.117.4.395

Gandolfo, M., & Downing, P. E. (2020). Asymmetric visual representation of sex from human body shape. *Cognition*, *205*, 104436. https://doi.org/10.1016/j.cognition.2020.104436

Girshick, A. R., Landy, M. S., & Simoncelli, E. P. (2011). Cardinal rules: Visual orientation perception

reflects knowledge of environmental statistics. *Nature Neuroscience*, *14*(7), 926–932. https://doi.org/10.1038/nn.2831

Hafri, A., Boger, T., & Firestone, C. (2022). Melting Ice With Your Mind: Representational Momentum for Physical States. *Psychological Science*, *33*(5), 725–735. https://doi.org/10.1177/09567976211051744

Hubbard, T. L. (1990). Cognitive representation of linear motion: Possible direction and gravity effects in judged displacement. *Memory & Cognition*, *18*(3), 299–309. https://doi.org/10.3758/BF03213883

Järveläinen, J., Schürmann, M., & Hari, R. (2004). Activation of the human primary motor cortex during observation of tool use. *NeuroImage*, *23*(1), 187–192. https://doi.org/10.1016/j.neuroimage.2004.06.010

Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The Fusiform Face Area: A Module in Human Extrastriate Cortex Specialized for Face Perception. *Journal of Neuroscience*, *17*(11), 4302–4311. https://doi.org/10.1523/JNEUROSCI.17-11-04302.1997

Kilner, J. M., Friston, K. J., & Frith, C. D. (2007). The mirror-neuron system: A Bayesian perspective. *Neuroreport*, *18*(6), 619–623. https://doi.org/10.1097/WNR.0b013e3281139ed0

Kuroki, D. (2021). A new jsPsych plugin for psychophysics, providing accurate display duration and stimulus onset asynchrony. *Behavior Research Methods*, *53*(1), 301–310. https://doi.org/10.3758/s13428-020-01445-w

Leopold, D. A., O'Toole, A. J., Vetter, T., & Blanz, V. (2001). Prototype-referenced shape encoding revealed by high-level aftereffects. *Nature Neuroscience*, *4*(1), 89–94. https://doi.org/10.1038/82947

Loffler, G., Yourganov, G., Wilkinson, F., & Wilson, H. R. (2005). FMRI evidence for the neural representation of faces. *Nature Neuroscience*, *8*(10), 1386–1391. https://doi.org/10.1038/nn1538

Ma, W., Kording, K. P., & Goldreich, D. (2022). Bayesian Models of Perception and Action. *MIT Press*. https://mitpress.mit.edu/9780262047593/bayesian-models-of-perception-and-action/

Makowski, D. (2018). The psycho Package: An Efficient and Publishing-Oriented Workflow for Psychological Science. *Journal of Open Source Software*, *3*(22), 470. https://doi.org/10.21105/joss.00470

Neri, P., Morrone, M. C., & Burr, D. C. (1998). Seeing biological motion. *Nature*, *395*(6705), 894–896. https://doi.org/10.1038/27661

Peelen, M. V., & Downing, P. E. (2005). Selectivity for the Human Body in the Fusiform Gyrus. *Journal of Neurophysiology*, *93*(1), 603–608. https://doi.org/10.1152/jn.00513.2004

Peelen, M. V., & Downing, P. E. (2007). The neural basis of visual body perception. *Nature Reviews Neuroscience*, *8*(8), 636–648. https://doi.org/10.1038/nrn2195

Pouget, A., Beck, J. M., Ma, W. J., & Latham, P. E. (2013). Probabilistic brains: Knowns and unknowns. *Nature Neuroscience*, *16*(9), 1170–1178. https://doi.org/10.1038/nn.3495

Quadflieg, S., & Koldewyn, K. (2017). The neuroscience of people watching: How the human brain makes sense of other people's encounters. *Annals of the New York Academy of Sciences*, *1396*(1), 166–182. https://doi.org/10.1111/nyas.13331

Reed, C. L., Stone, V. E., Bozova, S., & Tanaka, J. (2003). The Body-Inversion Effect. *Psychological Science*, *14*(4), 302–308. https://doi.org/10.1111/1467-9280.14431

Schwarzlose, R. F., Baker, C. I., & Kanwisher, N. (2005). Separate Face and Body Selectivity on the Fusiform Gyrus. *Journal of Neuroscience*, *25*(47), 11055–11059. https://doi.org/10.1523/JNEUROSCI.2621-05.2005

Shiffrar, M., & Freyd, J. J. (1990). Apparent Motion of the Human Body. *Psychological Science*, *1*(4), 257–264. https://doi.org/10.1111/j.1467-9280.1990.tb00210.x

Stein, T., Sterzer, P., & Peelen, M. V. (2012). Privileged detection of conspecifics: Evidence from inversion effects during continuous flash suppression. *Cognition*, *125*(1), 64–79. https://doi.org/10.1016/j.cognition.2012.06.005

Thorat, S., & Peelen, M. V. (2022). Body shape as a visual feature: Evidence from spatially-global attentional modulation in human visual cortex. *NeuroImage*, *255*, 119207. https://doi.org/10.1016/j.neuroimage.2022.119207

Vannuscorps, G., & Caramazza, A. (2016). Typical action perception and interpretation without motor simulation. *Proceedings of the National Academy of Sciences*, *113*(1), 86–91. https://doi.org/10.1073/pnas.1516978112

Vannuscorps, G., Pillon, A., & Andres, M. (2012). Effect of biomechanical constraints in the hand laterality judgment task: Where does it come from? *Frontiers in Human Neuroscience*, *6*. https://doi.org/10.3389/fnhum.2012.00299

Weiss, Y., Simoncelli, E. P., & Adelson, E. H. (2002). Motion illusions as optimal percepts. *Nature Neuroscience*, *5*(6), Article 6. https://doi.org/10.1038/nn0602-858

Wilson, M., Lancaster, J., & Emmorey, K. (2010). Representational momentum for the human body: Awkwardness matters, experience does not. *Cognition*, *116*(2), 242–250. https://doi.org/10.1016/j.cognition.2010.05.006

Yin, R. K. (1969). Looking at upside-down faces. *Journal of Experimental Psychology*, *81*(1), 141–145. https://doi.org/10.1037/h0027474

Zentgraf, K., Munzert, J., Bischoff, M., & Newman-Norlund, R. D. (2011). Simulation during observation of human actions—Theories, empirical studies, applications. *Vision Research*, *51*(8), 827–835. https://doi.org/10.1016/j.visres.2011.01.007