1 **Distal regulatory sequences contribute to diversity in brain oxytocin**
2 **receptor expression patterns and social behavior**
3

4 Qi Zhang[1,2,3,*], Luis Augusto Eijy Nagai[4], Mina Tsukamoto[1], Lenin C. Kandasamy[1],
5 Kiyoshi Inoue[5], Maria F. Pires[1], Minsoo Shin[1], Yutaro Nagasawa[1], Tsetsegee Sambuu[1],
6 Sonoko Ogawa[6], Kenta Nakai[7], Shigeyoshi Itohara[2], Larry J Young[3,5,8]
7

[1]Laboratory for Social Neural Networks, Faculty of Human Sciences, University of
Tsukuba, Japan; [2]Laboratory for Behavioral Genetics, CBS, RIKEN, Japan; [3]Center for
Social Neural Networks, Faculty of Human Sciences, University of Tsukuba, Japan;
[4]Institute for Quantitative Biosciences, The University of Tokyo, Japan; [5]Center for
Translational Social Neuroscience, Emory National Primate Research Center, Emory
University, USA; [6]Laboratory of Behavioral Neuroendocrinology, Faculty of Human
Sciences, University of Tsukuba, Japan; [7]Laboratory of Functional Analysis in silico,
Human Genome Center, The University of Tokyo, Tokyo, Japan; [8]Department of
Psychiatry and Behavioral Science, Emory University, USA.
17

18 *Correspondence: zhangqi@human.tsukuba.ac.jp or qi.zhang@a.riken.jp;

19
20

21 Classification: Biological Science: Neuroscience
22

23 Keywords: oxytocin system, behavioral evolution, topologically associating domain
24 (TAD), bacterial artificial chromosome (BAC), brain, mammary gland, partner
25 preference, maternal care, evolvability.
26

27 The PDF file includes:
28 Main Text
29 Figures 1 to 4
30 Supplementary Figures 1to 6, supplementary Tables 1 and 2
31
32

Summary

The oxytocin receptor (OXTR) modulates social behaviors in a species-specific manner.
Remarkable inter- and intraspecies variation in brain OXTR distribution are associated
with diversity in social behavior. To test the causal effect of developmental variation of
OXTR expression on the diversity of social behaviors, and to investigate potential genetic
mechanisms underlying the phylogenetic plasticity in brain *Oxtr* expression, we
constructed BAC transgenic mice harboring the entire prairie vole *Oxtr* locus with the
entire surrounding intergenic regulatory elements. Eight independent "volized" *prairie*
vole-Oxtr (*pvOxtr*) mouse lines were obtained; remarkably, each line displayed a unique
pattern of brain expression distinct from mice and prairie voles. Four *pvOxtr* lines were
selected for further investigation. Despite robust differences in brain expression, *Oxtr*
expression in mammary tissue was conserved across lines. These results and
topologically associating domain (TAD) structure analysis suggest that *Oxtr* expression
patterns in brain, but not other tissues, involve contributions of distal regulatory elements
beyond our BAC construct. Moreover, "volized" mouse lines with different brain *Oxtr*
expression patterns showed differences in partner preference and maternal behaviors. We
speculate that transcriptional hypersensitivity to variable distal chromosomal sequences
through long-distance interactions with proximal regulatory elements may contribute to
"evolvability" of brain *Oxtr* expression. The "evolvability" of brain *Oxtr* expression
constitutes a transcriptional mechanism to generate variability in brain OXTR which,
through natural selection, can generate diversity in adaptive social behaviors while
preserving critical peripheral expression. Transcriptional lability of brain *OXTR*
expression may also contribute to variability in social phenotype in humans, including
psychiatric outcomes.

Introduction

Brain oxytocin receptors (OXTR) regulate a wide range of social behaviors including
social recognition, maternal care, social bonding, empathy related behaviors, and
aggression [1-3]. In contrast to sex steroid receptors, which have highly conserved brain
expression patterns across species, OXTR shows remarkable inter- and intra-species
variation in brain distribution [2, 4]. For example, monogamous prairie voles have a brain
OXTR distribution in cortex, striatum and amygdala that [4]is distinct from promiscuous
montane voles and laboratory rats and mice. Furthermore, there is robust individual
variation in brain OXTR expression among prairie voles that is associated with single
nucleotide polymorphisms (SNPs) in the OXTR locus (*Oxtr*), and this variation predicts

pair bonding behavior and resilience to early life social neglect [5-8]. Different species of

primates also differ in brain OXTR distribution [9], and SNPs in the human OXTR have

been linked to variation in social function [10, 11]. Thus, variation in OXTR distribution

in brain likely constitutes a mechanism for the emergence of diverse social traits,

potentially including psychiatric endophenotypes.

Acute overexpression or knockdown of OXTR in a specific brain region alters social

attachment and parental behaviors in monogamous prairie voles but not polygamous

meadow voles [7, 12, 13]. A recent report showed that social attachment can occur in

constitutive *Oxtr* knockout prairie voles [14], suggesting that developmental OXTR

signaling may be essential for behavioral dependency on OXTR later in life. The direct

cause-effect relationship between species-specific OXTR expression pattern and social

behaviors, either developmentally or in adulthood, is unclear.

The evolution of gene expression patterns could be mediated by *cis* (via linked

polymorphisms) or *trans* (through diffusible products of distal genes, e.g., transcription

factors) changes, as a result of adaptation to the environment [15-17]. *Cis*-regulatory

differences are more commonly responsible for adaptive evolution and interspecific

divergence [18]. To explore the transcriptional mechanisms giving rise to species-specific

Oxtr expression and social behavior, we created transgenic mice using a prairie vole *Oxtr*

(pv*Oxtr*) bacterial artificial chromosome (BAC). The BAC construct covers the full

coding sequence, introns and the entire intergenic region of the pv*Oxtr* with its

neighboring genes. We expected two possible results: (1) our BAC construct

accommodates all the critical promoter/enhancers elements responsible for species- and

tissue-specific expression, therefore the *pvOxtr* transgenic mice would express brain

pvOxtr in a prairie vole-like pattern; (2)The regulatory elements essential for species- and

tissue-specific expression of *pvOxtr* in our BAC construct are not complete, the *pvOxtr*

transgenic mice would express *pvOxtr* in a pattern different from either the prairie vole or

the mouse. In either case, we can address our question: do developmentally distinct

patterns of brain OXTR expression give rise to different social behaviors?

We obtained 8 lines of pv*Oxtr* mice with germline transmission, each "volized" (pv*Oxtr*)

mouse line, i.e., each different integration site, displayed a unique pattern of expression in

brain distinct from that of either wildtype mice or prairie voles. Intriguingly, *pvOxtr*

expression was conserved in mammary gland across different lines. Moreover, different

"volized" mouse lines showed differences in partner preference and maternal behaviors.

These results demonstrate that transcriptional redistribution of brain OXTR in space and

during development can support the emergence of variation in social preference and

parental behaviors, which natural selection could act upon and amplify in niches where
the novel behavior is adaptive. In addition, our results provide important clues to
understand the genetic mechanism underlying species differences in brain gene
expression.

Results

Creation of the pv*Oxtr-P2A-Cre* BAC transgenic mice

BAC vectors can accommodate dispersed cis-regulatory elements across large regions of
genome. Transgene expression from BAC constructs are generally resistant to insertion
position effects [19] due to both the large spans of insulating genetic material which
protect the transgene cassette from the influence of the chromosomal environment, and
through the inclusion of necessary regulatory elements such as enhancers, silencers, locus
control regions, and matrix attachment regions [20, 21]. Large-scale screening of CNS
gene expression in BAC transgenic mice by the GENSAT (Gene Expression Nervous
System Atlas) Project found that more than 85% of BACs express reproducibly in
multiple independent transgenic lines and reporter gene expression faithfully recapitulates
endogenous gene expression patterns [22-24]. Indeed, reproducible expression from
mouse BAC vectors has been achieved for more than 500 genes [22-24]. Those that don't
express reproducibly are influenced by variation in BAC copy number and insertion sites
[25]. BAC engineering has also been used to mediate cross-species transgene expression.
A large number of mouse models of human dominant neurodegenerative disorders have
been created using human BAC transgene approaches. These model mice successfully
express human transgenes in human-like patterns and recapitulate disease-like
phenotypes [26-30]. We therefore decided to use BAC approaches to create "volized"
Oxtr mouse lines.
The most important determinant for correct expression from BAC vectors is inclusion of
as much of the intergenic region surrounding a gene of interest within the BAC construct
as possible[25]. To create pv*Oxtr-P2A-Cre* BAC transgenic mouse line, we obtained a
BAC clone (GeneBank: DP001214.1) from a prairie vole BAC library (CHORI-232) [31,
32] that includes the entire *pvOxtr* locus (the coding sequence and all the introns) with the
full intergenic sequence upstream (150kb) and downstream of the *pvOxtr* coding region
(9kb) (Fig1 B). The BAC clone contains a portion of *Rad18,* minus the first 11 exons*,*
located upstream of *pvOxtr,* and *Cav3*, minus the first exon including the start codon,
downstream of *pvOxtr.* Therefore, there will be no interference of these two flanking
transgenes on behavior*,* and the proximal topological structure around *pvOxtr* locus is

preserved to the maximum extent. A P2A-NLS-Cre cassette was inserted in-frame just
before the *Oxtr* stop codon to ensure co-expression of *Cre* and *Oxtr* (Fig1 C). Transgenic
mice were generated by pronuclear injection using C57BL/6J zygotes. Proper integration
of the reporter into the BAC clone was confirmed using gene specific PCR assays and
sequencing (Fig1 C, D and data not shown). We obtained 11 founder mice carrying the
transgenes, which was verified by PCR with multiple primer sets (Fig1 C, E). One
founder line was sterile, and two founder lines did not show germ-line transmission.
Eight founder lines successfully produced offspring and stably transmitted the transgene
to at least 3 generations. We named these lines "Koi", meaning "love" in Japanese.

Each Koi line showed a unique expression pattern of transgene in brain, yet
conserved expression in mammary gland.

We systemically evaluated 8 Koi lines by crossing them with ROSA-26-NLS-*LacZ* mice,
a Cre reporter mouse line expressing nuclear-localized beta-galactosidase [33, 34]. Since
Lac-Z is only expressed in the nucleus of double positive (CRE+, LacZ+) cells, but not in
the neuronal processes, we could easily assess the expression pattern of our transgene. We
found that each Koi founder line displayed a unique pattern of expression in brain distinct
from that of wildtype mice or prairie voles (Fig2A-O, and supplementary Fig1). Almost
all Koi lines showed transgene expression in olfactory bulb, lateral septum (Fig2F-I
Supplementary Fig1B,E,H,K) and ventromedial nucleus of the hypothalamus (Fig2K-N
Supplementary Fig1C,F,I,L), though with different intensities, suggesting that these
regions constitute a stable core *Oxtr* expression network, as these regions also express
OXTR in mice and voles [35, 36]. Several Koi lines expressed the transgene in some
brain regions specific for vole *Oxtr* expression, including nucleus accumbens (NAc)
(Fig2A,B,D Supplementary Fig1A,G,J), prefrontal cortex (PFC) (Fig2A,B,C
Supplementary Fig1J), lateral amygdala (Fig2K,L,M Supplementary Fig1C,I,L) and deep
layers of cingulate cortex (Fig2A,B,C Supplementary Fig1C,L), which suggested that cis
regulatory elements in the BAC are capable of mediating the expression of *Oxtr* in the
reward and reinforcement circuitry of brain, albeit depending on the sequence/structure of
distal sequences >150 kb upstream and >21 kb downstream of the transcription start site.
None of the Koi lines exactly "mirrored" the vole-specific expression patten. Six of the
Koi lines expressed the transgene in broad areas of thalamus (Fig2K,L,N and
Supplementary Fig1C,I,L), reminiscent of the strong expression of V1a vasopressin
receptor (*Avpr1a*) in the thalamus [4]. Intriguingly, the expression of OXTR in mammary

glands showed similar pattern among all the Koi lines and *mouseOxtr(mOxtr)-Ires-Cre*
knock-in line (Fig2P-T and Supplementary Fig2).

Transgenes that are introduced by pronuclear injection typically integrate into a single
site of the genome as tandem concatemers [37]. BAC copy number variation and
different insertional loci can lead to distinct patterns of ectopic expression, and increased
BAC transgene copy numbers often correlate with increased BAC gene expression [25].
We then examined the relationship between transgene copy number and LacZ signal
using quantitative genomic PCR [38]. As expected, the copy number of WT,
heterozygous and homozygous *mOxtr-Ires-Cre* knock-in mice was 0, 1 and 2 respectively
(Figure2Z). The copy number of transgene in Koi lines are: Koi-1 (12), Koi-2 (2), Koi-3
(2), Koi-4 (1), Koi-5 (2), Koi-6 (4), Koi-7 (9), Koi-8 (3). Koi-6 and Koi-7 showed the
most limited expression of reporter gene, though the copy number of transgene were 4
and 9 respectively. Koi-4 showed strong expression of reporter gene in NAc, septum and
thalamus, though there was only 1 copy of transgene. Therefore, integration site rather
than copy number is the reason of the variant BAC transgene expression patterns across
our Koi lines.

Oxtr is localized to a boundary of large Topologically Associating Domains

The divergent brain expression patterns across our Koi lines suggest that distal elements
external to our BAC cassette may contribute to the variation in *Oxtr* expression.
Topologically associating domains (TADs) represent a key feature of hierarchical genome
organization comprising chromatin regions that frequently interact within the chromatin
domain while insulating regulatory interactions across TAD boundaries [39-42]. Hi-C
assays provide a genome-wide view of chromosome conformation in multiple layers,
including TADs [43, 44]. Utilizing publicly available 3D chromatin datasets from mouse
(Supplementary Table1), we discovered that *Oxtr* is consistently located at the boundary
region of TAD structures (approximately 2000kb) across several tissues and cell lines
(Supplementary Fig3A). Specifically, only in brain related samples (whole brain tissue,
neural progenitor cells, and cortex tissue), *Oxtr* is situated at the boundary of a large
inter-TAD interaction between two neighboring TADs, forming larger TAD structures
(approximately 4000kb) (Supplementary Fig3A). In contrast, the oxytocin peptide gene,
Oxt, which is expressed in a highly conserved brain pattern across vertebrates, lacks such
organized large TAD structures (Supplementary Fig3B). Analysis with the datasets from
human brain-related samples revealed that human *Oxtr* is also localized at the boundary
region of TAD structures, though the heatmap showed a clearly distinct pattern with that

of the mouse (Supplementary Table2; Supplementary Fig4A). Like in mouse, human *Oxt*
doesn't show typical TAD structures (Supplementary Fig4 B). Although the Hi-C data
was not generated from prairie vole tissues, it demonstrates that the TAD landscape is
dramatically different between *Oxtr* and *Oxt* and highlights that the 3D genomic structure
surrounding the *Oxtr* gene varies across species and tissues.

Adult brain OXTR binding patterns differ across Koi lines

To detect the distribution of OXTR protein in adult Koi lines without the interference of
endogenous mouse OXTR (mOXTR) signal, Koi lines were crossed with *mOxtr$^{-/-}$* mice
and pvOXTR binding was detected using receptor autoradiography. We focused on 4 Koi
lines: Koi-1, Koi-2, Koi-3 and Koi-4 based on the strong expression of reporter gene in
behaviorally relevant neuronal populations. Autoradiography demonstrated functional
pvOXTR in brain resembling to some extent Cre-induced reporter gene expression
(Figure 3). Brain OXTR binding of Koi-3 was consistent with Lac-Z expression in the
reporter line (Fig2C, H, M and Figure3 J-L). Interestingly, the OXTR binding in PFC of
Koi-3 was similar to that of prairie vole. In contrast, Koi-4 showed low OXTR binding in
NAc when compared with the high Lac-Z staining (Fig2D, Figure3M), consistent with
transient *pvOxtr* expression during development, a pattern previously reported in rat NAc
[45]. Cre-dependent reporter gene product reflects the cumulative expression history of
the transgene throughout development. Koi-2, the line which displayed the expression of
reporter genes in the broadest regions, did not show strong signal in autoradiography,
again suggesting developmental changes in transgene expression. It is interesting that
Layer IV of cortex in Koi-4 showed strong OXTR binding (Figure3 N and O), which is
highly likely to originate from the axon projections from the thalamus (Fig2N), as was
previously reported for NAc projections in voles [46]. Consistent with Lac-Z expression,
we do not see a clear relationship between OXTR binding and transgene copy number.
For example, Koi-1 showed low OXTR signal (Fig3 D, E, F) with the highest copy
number, while Koi-4 showed very strong OXTR signal at lateral septum and thalamus
(Fig3 N, O), with the lowest copy number. *Oxtr* expression level was further confirmed
with qRT-PCR (Supplementary Fig 5, 6). Though OXTR binding in NAc of Koi-4 was
barely detected using autoradiography, the more sensitive qRT-PCR revealed that adult
Koi-4 mice had the highest Oxtr mRNA signal in the striatum, several fold higher than
WT, consistent with the Lac-Z staining in the reporter line.

Although none of our Koi lines mirrored the endogenous OXTR expression in prairie
voles, we confirmed strong expression of *pvOxtr* in olfactory bulb of Koi-1line, PFC and

amygdala of Koi-3line, striatum and thalamus of Koi-4 lines, and the broad expression of
pvOxtr in Koi-2 line. As OXTR signaling in some of these regions are similar to that in
voles, albeit in separate lines, and postulated to be involved in vole social behavior [47],
we used these lines to address the question of whether developmental variation in *Oxtr*
transcription in brain can lead to variation in social behaviors.

Different Koi lines showed diverse behaviors in a Partner Preference Test

The partner preference test is widely used as a laboratory proxy for pair bonding in
prairie voles. In a pilot study, we found that Kio-4 female mice on a m$Oxt^{+/+}$ background
showed a preference for their mate relative to a novel male. We then focused on the
females of 4 lines (Koi-1, 2, 3, 4) based on transgene expression in PFC, BLA or NAc.
We bred the transgenic mice with *mOxtr*$^{-/-}$ mice (both on C57/B6 background) to generate
offspring expressing only pv*Oxtr*. Partner preference tests (PPT) were performed in
ovariectomized females after 21 days of cohabitation with 3 mating bouts elicited by
estrogen and progesterone injections on days 2, 9 and 16. We divided our analysis into 3
phases (Fig4A, B). In the habituation phase, the experimental subject was habituated to
the chamber for 5 minutes with both pen-boxes empty. In the recognition phase, the
partner stimulus animal was restricted in one pen-box, and a novel "stranger" male was
restricted in another pen-box, and the experimental female was allowed to explore the
arena for 10 minutes to test social novelty preference. In the preference phase, the
positions of the partner stimulus animal and the "stranger" stimulus animal were switched
to avoid position bias, and the female was allowed to explore the chamber for an
additional 20 minutes.

Mixed ANOVA was conducted to examine the effect of genotype and the stimulus box on
the area stay time during all the phases. In the habituation phase (Fig4C), there was no
statistically significant interaction between the effects of genotype and position of empty
pen-box on stay time, $F_{(4, 60)} = 0.69$, $p = 0.6$. There was no difference in position
preference for all the groups, $F_{(1,60)} = 0.03$, $p=0.87$. In the recognition phase (Fig4D),
there was a main effect of stimulus animal (i.e., partner or stranger) on stay time, the time
staying in close proximity to the partner was significantly shorter than the time stay with
stranger animal ($F_{(1,60)} = 15.38$, $p<0.001$). There was a main effect of genotype on social
stay time, $F_{(4,60)} = 2.62$, $p<0.05$, and a significant interaction between the effects of
genotype and stimulus animal on area stay time, $F_{(4, 60)} = 2.61$, $p <0.05$. Bonferroni
adjusted pairwise comparisons were applied to analyze the simple main effects. Koi-4 and
WT mice spent significantly longer time in "stranger" area than in "partner" area (p<0.05),

demonstrating a typical novelty preference. There was no significant difference between
the stay time with "stranger" and "partner" for Koi-1 (p=0.96), Koi-2 (p=0.49) and Koi-3
(p=0.09) mice. In the preference phase (Fig5E), conducted after both stimulus mice were
presumably familiar following the initial 10 min recognition phase, there was no main
effect of stimulus animal on stay time, F (1,60) =0.04, p=0.84. There was no main effect of
genotype on stay time either, F (4,60) =1.36, p=0.26. There was a statistically significant
interaction between the effects of genotype and stimulus animal on area stay time, F (4, 60)
= 4.64, p <0.01. Bonferroni adjusted pairwise comparisons revealed that Koi-2 mice spent
significantly longer time in "stranger" area than in "partner" area (p<0.05). In contrast, Koi-
4 mice spent significantly longer time in "partner" area than in "stranger" area (p<0.01).
There was no significant difference between the stay time with "stranger" and "partner" for
WT (p=0.99), Koi-1 p=0.16) or Koi-3 (p=0.45) mice.

Different Koi lines showed diverse behaviors in Pup Retrieval Test

Based on the abundant pvOXTR of Koi-3 and Koi-4 in reward-related brain regions
(Fig3J-O), and the role of OXTR in mediated maternal care in mice and voles [48-50], we
further tested the parental behaviors of these two lines in virgins (Fig4F, G). Mixed
ANOVA was conducted to examine the effect of genotype and pup number (1-3) on the
latency of pup retrieval. Mauchly's test showed that the sphericity was violated.
Therefore, the Greenhouse-Geisser correction was used for the repeated measures
ANOVA. There was a significant main effect of genotype on latency to retrieve, F (2,48)
=29.58, p<0.0001. Koi-4 took a significantly longer time to retrieve all the pups than WT
and Koi-3 (p<0.0001 for both). There was also a significant interaction between the
effects of genotype and retrieved pup number on the retrieval latency, F (2.40, 57.49) =
4.45, p <0.05. Post Hoc multiple comparisons with Bonferroni correction revealed that
Koi-4 took a significantly longer time to accomplish the 1st retrieval (p<0.001), the 2nd
retrieval (p<0.0001), and the 3rd retrieval (p<0.0001) when compare to WT and Koi-3.
And Koi-3 took a significantly shorter time to accomplish the 1st retrieval (p<0.05), the
2nd retrieval (p<0.01), and the 3rd retrieval (p<0.01) when compare to WT.

Crouching time was analyzed as well (Fig4F, H). There was a statistically significant
difference between groups as determined by one-way ANOVA (F (2,48) = 27.56, p
<0.0001). A Bonferroni post hoc test revealed that the crouching time of Koi-3 was
statistically significantly longer than WT (p<0.01) and Koi-4(p<0.0001). Additionally, the
crouching time of Koi-4 was statistically significantly shorter than WT (p<0.01).

Therefore, two Koi lines with different OXTR expression pattern demonstrated

quantitative differences in parental behavior tests.

Discussion

Diversity of developmental *Oxtr* expression patterns causes diversity of social

behaviors

The diverse OXTR expression of our Koi lines is reminiscent of the diverse OXTR

expression across different species. Species and individual differences in OXTR

distribution in the brain have been associated with variation in social behaviors [2, 4].

For instance, the monogamous prairie vole expresses high densities of *Oxtr* in the PFC,

NAc and BLA compared to non-monogamous vole species or mice. Pharmacological and

viral mediated siRNA manipulations of OXTR in adult vole PFC and NAc suggest that

OXTR in these regions play a role in parental care and pair bonding[12, 13, 48].

Furthermore, variation in prairie vole OXTR expression in brain are also associated with

individual differences in parental behavior, pair bonding, and resilience to neonatal

neglect[5, 8, 51, 52].   However, it is unclear whether inter- or intra-species variation in

brain OXTR signaling during development, due to variation in brain expression pattern,

is causally mediating diversity in social behaviors. We obtained expression in PFC, NAc

and BLA regions in our Koi mice, but in 3 separate lines. These Koi lines with distinct

brain OXTR expression pattern provided an opportunity to test the causal effect of

development variation of OXTR expression on diverse social behaviors developmentally.

In contrast to wildtype mice, the Koi-4 line showed a preference for their mating partners

over unfamiliar males, as is typically observed in prairie voles. Multiple studies showed

that NAc and PFC play important roles in mediating partner preference behavior in

prairie voles [53, 54]. Although the Koi-4 line showed robust transgene expressed in

these regions using the Lac-Z reporter mouse method, qPCR and receptor

autoradiography suggest more limited OXTR expression as adults, albeit several fold

higher than WT mice, suggesting transient changes in transgene expression during

development. Emerging research have shown that species differences exist in

developmental patterns of OXTR expression. For example, pre-weaning rats have a

transient peak OXTR specifically in NAc and cingulate cortex [55, 56], while mice have

a transient peak of OXTR throughout the entire neocortex [57]. Transient OXTR

expression may define sensitive periods for oxytocin to shape the brain development of

those brain regions in a social stimulus dependent way[58]. OXTR signaling in the NAc

during the first two weeks of life appear to shape adult partner preference behavior in

prairie voles [5]. The brain regions that display transient appearance of OXTR are not

conserved across species, and transient OXTR expression may help to shape activity-
dependent development and contribute to adult social behaviors in a species-specific
manner [59]. Partner preference formation and maintenance needs to recruit multi- and
cross-modality sensation, social recognition and social memory, selective attention,
arousal, and reward systems. We propose a model that the sufficient upregulation of the
OXTR expression in one or multiple of these systems can tune the network and boost
partner preference behavior. In addition, the maternal behavior in Koi-3 and Koi-4 lines
was enhanced and reduced, respectively. This result suggested that OXTR signaling
facilitates social partner preference and maternal care by coordinating activity across
different neural networks. A main conclusion of our study is that simply by redistributing
OXTR binding in space and during development, variation in social preference and
parental behaviors emerged, which conceivably could impart advantageous behaviors that
natural selection could act upon and amplify. Future study will determine how the
variation in OXTR distribution in our Koi lines affects neural circuits mediating social
behaviors.

A potential genetic mechanism contributing to diversity of brain OXTR expression
and social behavior

The sex steroid and oxytocin systems play important roles in modulating reproductive
and related social behaviors in vertebrates. Sexual behavior, e.g., the motivation to mate
and motor patterns, is highly conserved across vertebrate species. By contrast, social
behaviors, e.g., sociality, mating strategy (monogamy vs polygamy), parental and
alloparental behaviors are more nuanced and can vary dramatically across species in the
same genus or even between individuals of a single species. It is interesting, therefore,
that the distribution of steroid receptors is quite conserved among species [60], while that
of the OXTR is extremely diverse across and even within species [2, 4]. It is even more
interesting that although the distribution of OXTR is diverse, its ligand, oxytocin, is
expressed in a highly conserved neuroanatomical pattern in vertebrates [61]. Previous
work showed that three independent rat lines with puffer fish oxytocin BAC transgene
(containing 16 kb of 5' flanking regulatory sequence) expressed the *Fugu* oxytocin
transcript specifically in oxytocin neurons in rat [62]. Moreover, a similar transgenic
experiment transferring approximately 5 kb of the *Fugu* oxytocin already resulted in a
faithful expression of the fish genes in mouse oxytocin neurons [63]. These studies
suggest there is a remarkably conserved transcriptional regulatory machinery across
species, even between the *Fugu* and rodent oxytocin genes, despite being separated by

400 million years of evolution. Finally, 2.6 kb of the oxytocin promoter is sufficient to

faithfully express transgenes in oxytocin neurons when delivered using a viral vector[64].

Based on the principle of choosing a proper BAC clone to create BAC transgenic mice,

the most important determinant for correct expression from BAC vectors is inclusion of

as much as possible of the 5' and 3' intergenic region surrounding a gene of interest [23,

25, 65]. The *pvOxtr* BAC clone we chose carries the entire *pvOxtr* locus with the full

intergenic sequence upstream and downstream of *pvOxtr*. Though in some cases BAC

sequences don't include the whole TAD, they still contain the critical tissue-specific

control elements for the locus, and can confer reproducible, accurate expression patterns

of transgenes [66]. However, the 8 lines we obtained demonstrated remarkably variant

brain pvOXTR expression, distinct from both prairie vole and mouse. The different

expression pattern is not from the copy number of transgenes. Instead, it is most likely

due to the effects from different insertion sites of transgenes. These results are

reminiscent to the observation that virtually every rodent or primate species examined has

a unique distribution of OXTR binding in the brain. Our 3D genomic structure analysis

using open databases showed that the mouse *Oxtr* and human *OXTR* loci sit within large

TAD structures, while no typical TAD structure surrounds *Oxt* loci in either species.

Considering the size of TADs where *Oxtr* locates in brain related tissues both in mouse

and human (about 4000kb), certain regulatory elements (including insulator sequences)

essential for brain-specific expression of *pvOxtr* may lie outside of the BAC region. This

might explain why short promoter sequences suffice for faithful expression of oxytocin

across species and when employing viral vectors, but a 200kb BAC is not adequate for

precise, reproducible transcription regulation of *Oxtr*. This may also account for the

transcriptional hypersensitivity of *pvOxtr* to chromosomal position effect in brain.

It is also noteworthy that pv*Oxtr* expression in mammary gland were conserved across

different Koi lines. These results suggested that the promoter/enhancer/insulator elements

critical for mammary gland-specific expression of *pvOxtr* are encompassed by the BAC

construct, and the regulation of *pvOxtr* expression in mammary gland is highly resistant

to the position effect of integration sites. Our 3D genomic structure analysis using mouse

datasets showed that the chromatin landscape differs considerably between brain-related

tissues and other tissues. This is observed by an obvious increase of long-distance

interactions between neighboring TADs, forming a larger TAD structure in brain-related

tissues. We here propose a new genetic model of *Oxtr* expression regulatory machinery:

The sequence proximal to *Oxtr* loci within the BAC construct is sufficient to form a

complete regulatory unit. This unit is resistant to changes in interactions with distal

regulatory sequences and confer faithful expression of OXTR in peripheral tissues where
OXTR expression is essential for reproduction. In contrast, interacting regulatory
elements that govern brain region-specific expression of OXTR are dispersed over a large
sequence, extending far beyond the BAC construct. This makes OXTR expression in
brain susceptible to the chromosomal landscape, even at locations very distal to the gene
loci. This is well aligned with the notion that subtle changes in 3D chromatin structure
may result in substantial changes in local gene expression [67]. Such an organization
could offer increased opportunities for chromosomal variation across species to impart
evolvability (the capacity of an evolving system to generate or facilitate adaptive change)
of *Oxtr* expression in the brain. Consequently, this would allow oxytocin to modulate
novel neural circuits, which can then modify various facets of social behavior, leading to
the evolution of diversity in social behaviors. If the resultant behavior is adaptive, the
frequency of the mutation will increase in the population and the social behavioral profile
of the population may change. Future work applying Hi-C and ChIP approaches on
different brain regions and peripheral tissues from prairie vole, montane vole and our Koi
lines to see the chromatin structure and epigenetic landscape surrounding *Oxtr*, will be
very helpful to understand how TAD landscape and epigenetics influences *Oxtr* gene
expression and social behavior traits.

There is other experimental evidence supporting the notion that brain expression of *Oxtr*
is labile, sensitive to genetic variation. SNPs in the *Oxtr* intron explains 74% of the
variance in striatal *Oxtr* expression and social attachment in prairie voles [8] and it is
possible that these SNPs near *Oxtr* are linked to larger distal variations that affect brain
Oxtr expression. Different *Oxtr*-reporter (either lac-z, EGFP or Venus etc.,) mice created
from independent labs either by knock-in or BAC transgene methods showed different
expression pattern in brain as well [25, 35, 65], further supporting the ultra-sensitivity
and lability of *Oxtr* gene to subtle changes of distal cis sequence and local chromosomal
architecture. Additionally, SNPs in the human OXTR predict OXTR transcripts in a brain
regions-specific manner [68] as well as brain functional connectivity[69]. This
transcriptional ultra-sensitivity to genomic variation appears to be tissue specific and
mainly appears in brain, but not in mammary gland, which is responsible for essential
lactation in mammals.

Another interesting point is that *Oxtr* is localized at TAD boundary regions in mouse and
human. TAD boundaries are thought to be essential for normal genome function, given
their roles in defining regulatory territories along chromosomes and in preventing

unintended enhancer-promoter interactions between adjacent chromatin domains. Cross-
species multiple sequence alignments have revealed an enrichment of syntenic breaks at
TAD boundaries[70].   Future studies will explore synteny regions along the *Oxtr* TAD in
different species.

In summary, we created mouse lines showing different expression pattern of brain
OXTR using a BAC transgenic method. Our results suggest that complex, long-distance
interactions between proximal and distal cis regulatory elements contribute to region-
specific expression in brain, but not mammary tissue. The variability in transgene
expression pattern in brain depending on integration site suggest that *Oxtr* expression is
sensitive to large-scale chromosomal interactions, which may impart evolvability of *Oxtr*
expression patterns in the brain. Subtle variation in chromatin landscape of *Oxtr* gene
may alter long distance regulatory interactions, and thereby alter brain expression
patterns and lead to the emergence of novel social behaviors, which may allow organisms
to adapt to and survive variable environments. We speculate that this transcriptional
evolvability may contribute to inter- and intra-specific *Oxtr* expression patterns in brain.
We also confirmed for the first time the causal effect of developmental variation in brain
Oxtr distribution on the diversity of social behaviors. Our mice lines with different Oxtr
expression patterns display differences in partner preference and enhanced or reduced
parental behaviors. This research on the origins of diversity in social behaviors across
species may lead to conceptual understandings relevant for the development of treatments
for psychiatric disorders.

Acknowledgements:
This work was supported by the International Education and Research Laboratory
Program of University of Tsukuba to L.Y., the RIKEN Incentive Research Project
(100226201701100443) to Q.Z., the Brain Science Project, Center for novel science
initiatives, National institutes of natural sciences (BS291003) to Q.Z., the RIKEN Aging
Project (10026-201701100263-340120) to Q.Z., the JSPS Kakenhi Grant-in-Aid for
Young Scientists (17K18362) to Q.Z., the JSPS Kakenhi Grant-in-Aid for Challenging
Research (19K21807) to Q.Z., the JSPS Kakenhi Grant-in-Aid for Challenging Research
(2 2 K 1 9 4 7 8)to Q.Z., and the Takeda Medical Research Grant to Q.Z.. We thank Dr.
Katsuhiko Nishimori (Tohoku University and Fukushima Medical University, Japan) for
providing *mOxtr-Ires-Cre* knock-in line and *mOxtr^{-/-}* mutant line; Dr. Cary Lai (Indiana
University, USA), Dr. Takuji Iwasato(National Institute of Genetics, Japan) and Dr. Seiya

Mizuno(University of Tsukuba, Japan) for their advice on BAC vector construction; Dr. Volkhard Maeckel (CERN, Switzerland) for helping with image capturing; Dr. Yong Zhang (Tongji University, China) for the stimulating discussion about 3D chromatin structure, and we thank the Laboratory of Animal Resource Center of University of Tsukuba for pronuclear injection.

Author Contributions:

Q.Z. conceived the project, designed the study, led the teamwork, conducted transgene vector construction and drafted manuscript. Q.Z. and L.N. and K.N analyzed Hi-C datasets. Q.Z., L.K. and M.T. conducted histological analysis. Q.Z. and M.P. analyzed transgene copy number and gene expression level. Q.Z. and K.I. conducted autoradiography analysis. Q.Z., L.K., M.T. and M.P. performed behavioral test and analyzed the data. Y.N. and T.S. performed ovariectomy surgery. L.K., M.T., M.S. and M.P. performed genotyping and mouse maintenance. S.I. provided advice and facility for transgene vector construction. L.Y helped conceive the project, provided advice for the whole project and edited the manuscript. All authors discussed the manuscript.

Declaration of Interests:

The authors declare no competing interests.

Figure titles and Legends:

Fig.1 Construction of Koi lines. (A) Example of the remarkable species difference of brain OXTR distribution. OXTR receptor autoradiography in dorsal caudate putamen (CP) and nucleus accumbens shell (NAccSh) of rat, mouse, and prairie voleis (Adapted Froemke and Young[2]). (B) The *pvOxtr* BAC clone contains the entire sequence of *pvOxtr* gene (colored green), the complete intergenic sequence both upstream and downstream of *pvOxtr* gene, the 12th exon of *Rad18* (colored blue) and the 2nd exon of *Cav3* (colored purple). The yellow arrows indicate the direction of transcription. (C) Schematic diagram of the strategy to generate Koi mice. The P2A-NLS-Cre-Frt-Amp-Frt cassette was inserted in-frame right before the stop codon of *Oxtr*. And the ampicillin selection marker was deleted by 706-Flpe. PCR primers for verifying the targeted alleles are shown as red arrows. (D) Two correctly modified vector clones were verified by using PCR with p2 and p8. (E) Koi lines were verified by 5 pairs of primers. From Lane1 to Lane8 are 100bp marker, p3+p7, p2+p6, p2+p7, p4+p8, p2+p8, blank and 1kb marker.

Fig.2 Transgenic CRE mediated Lac-Z reporter gene expression in the brain and

mammary gland of Koi lines. X-gal staining of the brains and mammary glands are

from the double positive (*Cre+, LacZ+*) offspring from Koi-1 (A, F, K, P), Koi-2 (B, G,

L, Q), Koi-3 (G, H, M, R), Koi-4 (D, I, N, S) and *mOxtr-Ires-Cre* knock-in line (E, J, O,

T). U, V, M, X, Y are the magnified images of the areas within the yellow-border squares

from D, N, E, J, O, respectively, to show the distribution of low-density Lac-Z positive

cells in these areas. Scale bar=1mm (A-O), 500um (P-T) and 200um (U-Y). (Z) The

estimated copy number of transgene of all Koi lines. See Supplementary Figure 2 for

images of Koi-5-8 lines. PFC=prefrontal cortex, LS=lateral septum, BLA=basolateral

amgydala, VMH=ventromedial nucleus of the hypothalamus, Tha=thalamus.

Fig.3 The expression of OXTR in adult brain of Koi lines. Autoradiographs illustrate

the distribution of OXTR binding in adult WT (A-C), Koi-1 (D-F), Koi-2 (G-I), Koi-3 (J-

L), and Koi-4 (M-O). PFC=prefrontal cortex, LS=lateral septum, BLA=basolateral

amgydala, VMH=ventromedial nucleus of the hypothalamus, Tha=thalamus. Scale

bar=1mm.

Fig.4 Social behaviors of Koi lines. (A) Illustration of the partner preference test (PPT)

paradigm. The experimental mouse was placed in a chamber with an automatic

monitoring system, where it could freely explore two pen-boxes localized in two diagonal

corners. Habituation phase: two pen-boxes are empty. Recognition phase: a stimulating

partner mouse is placed into one pen-box and a stimulating stranger mouse is placed in

the other one. Preference phase: The positions of two stimulating mice are switched. (B)

Representative traces of one experimental mouse during the three phases. (C-E) The stay

time in the surrounding area of each pen box during three phases;(C) Habituation

phase;(D) Recognition phase; (E) Preference phase. Samples sizes for the PPT are WT

(n=12), Koi-1 (n=14), Koi-2 (n=13) Koi-3 (n=12) Koi-4 (n=14). All the experimental

mice are female. (F) The illustration of the procedure of the maternal behavior test. (G)

Pup retrieval latency and (H) crouching time were recorded and analyzed. Sample sizes

for the maternal behavior tests are WT (n=15), Koi-3 (n=18), Koi-4 (n=18). Significance

of interaction between independent factors: # $p<0.05$; # # $p<0.01$. Significance of main

effect or single main effect of genotype: ＊$p<0.05$; ＊＊$p<0.01$; ＊＊＊$p<0.001$; ＊＊

＊$p<0.0001$.(Illustrations created with Biorender.com)

Star Methods

pvOxtr-P2A-Cre **BAC vector construction**.

The vole BAC clone (GeneBank: DP001214.1) containing the *Oxtr* locus was obtained

from a prairie vole BAC library (CHORI-232) [31, 32]. This BAC clone carries the entire

pvOxtr locus with the full intergenic sequence upstream of 5' of *pvOxtr* (150kb) and

downstream of coding region of *pvOxtr* (9kb). The NLS (nuclear localization signal)-

Cre-Frt-Amp-Frt cassette was obtained from Dr. Takuji Iwasato, and was slightly

modified to introduce a P2A sequence [71]. The P2A sequence was introduced using

PCR with the following primers:

fw5'- <u>ATGGGAAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAGGCTGGAGA</u>

<u>CGTGGAGGAGAACCCTGGACCT</u>CTCGAAACTGACAGGAGAACCACC-3';

rv5'-AGACTGGAGTCCGCATAGCCCCCCTCCCCCGCCCCAGGCGCGGTGGGCC

AGGCAGGTGGCTCACCTTGACCAAGTTGCTGAAGTTCCTATTCC-3'. The P2A sequence is

underlined. And the resulting amplified cassette had the P2A upstream of the NLS-Cre.

Then the P2A-NLS-Cre-Frt-Amp-Frt cassette was inserted in-frame right before the stop

codon of Oxtr in the BAC clone using the Red/ET Recombineering kit (Gene Bridges

GmbH, Heidelberg Germany) after adding homologous arms by PCR with the following

primers: fw5'-CACCTTCGTCCTGAGTCGCCGCAGCTCCAGCCAGAGGAGCTG

CTCTCAACCATCTTCAGCAATGGGAAGCGGAGCTACTAAC-3';

rv5'-GAGGAGAGGGATACACACCAATAGGCACCTTATACAACTCCACGCAC

GGCCACCAGGGGCAGACTGGAGTCCGCATAGCC-3'. The ampicillin selection marker was

deleted with the 706-Flpe-induced recombination method (Gene Bridges Heidelberg

Germany). A correctly modified BAC clone was verified by using PCR at both the 5' and

3' junctions of the targeted insertion with the following primer pair : fr5'-

GCCTTCATCATCGCCATGCTCTT-3'(p2) and rv5'-GATGGCTGAGTG ACTGGCATCT-3'(p8);

The construct was further verified by sequencing using the following 7 primers specific

either for BAC vector or the p2A-NLS-Cre fragment:

5'- CCTGCAGCCAACTGGAGCTTC-3'(p1); 5'- CTTCCTTGGGCGCATTGACGTC-3'(p5);

5'-TACCTGTTTTGCCGGGTCAG-3(p4)；5'-GCCTTCATCATCGCCATGCTCTT-3'(p2)；

5'-CTGACCCGGCAAAACAGGTA-3'(p7)；   5'-TCCGGTTATTCAACTTGCACCATGC-3'(p6);

5'-GATGGCTGAGTGACTGGCATCT-3'(p8).

Animals.

***pvOxtr-P2A-Cre* BAC transgenic mouse lines (Koi lines)**. *pvOxtr-P2A-Cre* BAC vector

was digested with NotI for linearization and to remove the pTARBAC vector backbone.

The *Vole-Oxtr-P2A-Cre* BAC linearized fragment was purified using CL-4B sepharose

(Sigma-Aldrich), and injected into pronuclei of C57BL/6J zygotes. Mice carrying the

BAC transgene were identified and confirmed using PCR with 5 pairs of primers: fr5'-

TCGACCAGGTTCGTTCACTC-3'(p3) + p7; p2+ p6; p2 +p7; p4 +p8; p2+p8. 8 lines (Koi-1~Koi-

8) were confirmed with successful transgene inheritance. These lines were maintained on

a congenic C57BL/6J background by backcrossing with wildtype C57BL/6J for

assessments of BAC DNA copy numbers and initial transgene expression analysis.

The ***ROSA-26-NLS-LacZ* mouse** line [33] was obtained from Itohara Lab at RIKEN.

The ***mOxtr-Ires-Cre* knock-in** [72] and ***mOxtr*$^{-/-}$ mutant** [73] lines were obtained from

Dr. Katsuhiko Nishimori at Tohoku University, and both were backcrossed with

C57BL/6J for more than 10 generations. Each Koi line and *mOxtr-Ires-Cre* knock-in

mice were crossed with *ROSA-26-NLS-LacZ* mice to generate Cre$^+$, LacZ$^+$ double

positive mice for further gene expression analysis. Koi mice were then crossed with *Oxtr*$^{-/-}$

to generate mice in which the pvOXTR was expressed while endogenous mOXTR was

absent, and these mice were used for autoradiography, qRT-PCR and behavioral tests. All

mice were generated using continuously housed breeder pairs and P21 as the standard

weaning date. All the animal procedures were approved by University of Tsukuba Animal

Care and Use Committee. Mice were housed under constant temperature and light

condition (12 h light and 12 h dark cycle), and received food and water ad libitum.

Determination of transgene copy number

Custom Taqman® MGB probes were synthesized by oligoJp (Thermo Fisher, Japan) for

detecting the transgenic *Cre* and the mouse *Jun* gene (internal control). The following

primer pairs and probes were used: for the *Cre* assay, fr5'-

ATGACTGGGCACAACAGACAAT-3'; rv5'- CGCTGACAGCCGGAACAC-3'; probe:

5'-FAM- AAACATGCTTCATCGTCGGTC CGG-MGB-3'; for *Jun* assay, fr5'-

GAGTGCTAGCGGAGTCTTAACC-3'; rv5'-CTCCAGACGGCAGTGCTT-3'; probe:

5'-VIC-CTGAGCCCTCCTCCCC-MGB-3'. Real-time PCR was performed using an

Applied Biosystem7500. Transgene copy number was determined through absolute

quantification (standard curve method) following the protocol described in [38]. Briefly,

2 microliters (20 ng) of genomic DNA samples or copy number standards were analyzed

in a 20μL reaction volume with two primer-probe sets (*Cre, Jun*). The data from

transgenic samples were then compared to a standard curve of calibrator samples that are

generated by diluting purified BAC DNA (linearized) over a range of known

concentrations into wild-type mouse genomic DNA. In addition, no-template controls

were included in each experiment. All reactions were performed in triplicate.

Histology

Adult mice (3-4month old), 4 male and 4 female from each line/ generation were

analyzed for 3 generations. Mice were perfused intracardially with 4% formalin in 0.1M

sodium phosphate buffer. Brains were embedded in 2% agarose in 0.1M PB and cut into

100-m-thick sections with a Micro-slicer (Dosaka, Kyoto, Japan). Mammary glands were

isolated from virgin female mice (2month old), 2 mice/each line were analyzed. The brain

slices and whole mount mammary glands preparation were stained in X-gal solution (5

mM K3FeCN6, 5mMK4FeCN6,2mMMgCl2, 0.02% NP-40, 0.01% Na-deoxycholate, 1

mg/ml X-gal in 0.1M PB) at 37°C for 6 hours and then were stained with hematoxylin.

All experiments were done with positive and negative controls (Cre$^+$, LacZ$^-$) to monitor

the reliability of the X-gal staining. Images were caught by Nanozoomer 2.0-HT slide

scanner (Hamamatsu Photonics).

3D chromatin verification on Hi-C datasets of humans and mice

641 The 3D genome structure tracks were obtained from 4DNucleome Consortium

642 (https://www.4dnucleome.org/) where datasets were systematically reanalyzed following

643 recommended standard protocols. The datasets IDs and sample information are provided

644 in the supplementary material (Supplementary Table1,2). 3D Genome Browser[74] and

645 WashU Epigenome Browser[75] were employed for visualizations.

Receptor Autoradiography

OXTR autoradiography was performed as previously described [36]. Briefly, freshly

frozen brains were stored at -80°C. Coronal sections were cut in a cryostat and 20 μm

sections were collected and then stored at -80°C until use in autoradiography. Brain

sections were removed from -80°C storage and air dried, and then fixed for two minutes

with 0.1% paraformaldehyde in PBS at room temperature, and rinsed twice in 50

mM Tris buffer, pH 7.4, to remove endogenous OT. They were then incubated in 50 pM

125I-OVTA (2200 Ci/mmol; PerkinElmer; Boston, MA), a selective, radiolabeled OXTR

ligand, for one hour. Unbound 125I-OVTA was then washed away with Tris-MgCl$_2$

buffer (50 mM Tris plus 2% MgCl2, pH 7.4) and sections were air dried. Sections were

exposed to BioMax MR film (Kodak; Rochester, New York) for five days. Digital images

were obtained with a light box and a Cannon camera (Cannon 6D MarkII, Japan). The

brightness and contrast of representative images were equally adjusted for all

autoradiography images within a panel using Adobe Photoshop.

Quantification of *Oxtr* mRNA expression by qRT-PCR

Real-time quantitative polymerase chain reaction (RT-qPCR) using an Applied
Biosystems 7500 Real-Time PCR system was used to quantify *Oxtr* mRNA expression
levels in different brain regions of Koi lines. Total RNA of different brain regions was
isolated using the RNeasy extraction kit (Qiagen, Germantown, US), and cDNA was
synthesized after Deoxyribonuclease I (Invitrogen, Waltham, US) treatment by using
EvoScript Universal cDNA Master (Roche, Penzberg, Germany), and then quantitative
RT-PCR was performed with FastStart Universal SYBR Green Master (Roche, Penzberg,
Germany). The relative standard curve method was used to obtain the relative quantities
of *Oxtr* expression following the manual of Applied Biosystems. To allow the
comparation between Koi lines and WT mice, the following primer pair (which can detect
the expression of both *mOxtr* and *pvOxtr*) was used: fr5'-
GCCTTTCTTCTTCGTGCAGATG-3' and rv5′-ATGTAGATCCAGGGGTTGCAG-3′ In
addition, to specifically detect the expression of *pvOxtr*, the primer pair fr5′-
GCCTTTCTTCTTCGTGCAGATG-3 and rv5'-AAAGAGGTGGCCCGTGAAC-3′ was
used in the 2nd qRT-PCR experiment. GAPDH was used as the endogenous control for
both experiments and was amplified by the following primer pair: fr5'-
GGGTTCCTATAAATACGGACTGC-3' and rv5'- CCATTTGTCTACGGGACGA-3'.
Samples were analyzed in triplicates. A non-template control was performed to ensure
that there was no amplification of genomic DNA. All of the experimenters were blind to
the genotype of the subjects.

Partner preference test (PPT)

Subjects were housed with 4 age-matched and same-sex littermates until testing at
adulthood (2-5 months old). 12-14 homozygous? mice were tested for each group.
Ovariectomized females were paired and cohoused with a sexually experienced adult
wildtype male for 21 days before PPT test. The female was injected with estradiol
benzoate (10 µg and 5 µg at 48 h and 24 h before induced mating) and progesterone (500
µg at 4–7 h before induced mating) to ensure high sexual receptivity before mating.
Mating was induced on Day 2,9 and16. PPT data was analyzed using TimeSSI1 for social
interaction test system (O' Hara & Co., Ltd.). Briefly, the experimental subject was
placed in a chamber, in which two pen-boxes (diameter 8cm, height10cm) were located at
two diagonally opposite corners. The experimental animal was free to move throughout
the chamber and the time spent in close proximity to each pen-box (stay time) is recorded

using an automated mouse tracking system (Fig.4 A, B). In the habituation phase, the

experimental subject was allowed to explore the chamber for 5 minutes with both pen-

boxes empty. In the recognition phase, the partner stimulus animal was restricted in one

pen-box, and a novel "stranger" stimulus animal was restricted in another pen-box, and

the experimental subject was allowed to explore the chamber for 10 minutes. In the

preference phase, the positions of the partner stimulus animal and the "stranger" stimulus

animal were switched, and the experimental subject was allowed to explore the chamber

for 20 minutes. To assure unbiased design, pen-box assignments were counterbalanced

for the diagonal positions. All of the experimenters were blind to the genotype of the

subjects.

Parental behavior test

To examine maternal behavior, we performed pup retrieval test with virgin female mice

(2- to 4-months old, 15-18mice/line), following the method described in [76] with some

modifications. Briefly, 5 days prior to the behavioral testing, each subject was isolated

and housed in separate home cages in the test room. Nesting material (about 1.5 g of

cotton wool) was provided for nest making. The tests were conducted in the dark phase of

light/dark cycle (12 hours/12 hours). On the day of the test, each subject was placed in

the recording area and allowed 15 minutes for habituation. The stimulus pups (age: P2-

P3) were collected from a group of donor mothers immediately before the start of the

experiment. A retrieval test began with the placement of 3 stimulus pups on the side

farthest form the nest, and the female's behavior was recorded by an ARNAN 4 channel

security system for 15 minutes. Pup retrieval was defined as picking up the pup and

bringing it to the nest. If the subject did not complete the retrieval for 3 pups in 15

minutes, the video recording time was extended to 30 minutes. The latency to retrieve

each pup (1st, 2nd, 3rd) and the time spent crouching on the pups were time-stamped and

calculated manually. A latency of 1800 seconds was assigned if the pup-retrieval was not

completed in 30 minutes. Crouching was defined as the mouse supporting itself in a

lactation-position over the pups in the nest. Crouching time on a single pup, two pups,

and three pups were summed up as total crouching time. All of the

experimenters were blind to the genotype of the subjects.

Statistical analysis

All the statistical analyses were performed by SPSS21(IBM). Mixed ANOVA was used

for analyzing PPT data and pup retrieval data. Mauchly's test of sphericity was used to

test whether or not the assumption of sphericity was met in repeated

measures. Greenhouse-Geisser correction was applied when the sphericity was violated.

Two-way ANOVA was used for analyzing qRT-PCR region specific data across different

mouse lines. One-way ANOVA was used for analyzing crouching time in the parental

behavioral test. If there was a significant main effect of an independent factor, post hoc

test was used to do multiple comparisons. If there was a significant interaction between

within-subject factor and between-subject factor, post hoc pairwise comparisons were

applied to analyze the simple main effects. Bonferroni correction was used for all the post

hoc tests.

Supplemental information titles and legends:

Supplementary Fig.1 Transgenic CRE mediated Lac-Z reporter gene expression in

the brain of Koi lines. X-gal staining of the brains from the double positive (*Cre+*,

LacZ+) offspring of Koi-5 (A-C), Koi-6 (D-F), Koi-7 (G-I), Koi-8 (J-L). Scale=1mm

Supplementary Fig.2 Negative control of whole-mount mammary gland staining.

X-gal staining of a mammary gland from (*Cre-*, *LacZ*+) offspring from heterozygous Koi-

4 was shown. The lymph node showed some blue signal, indicating that the lymph node

could be stained nonspecifically. All the epithelium ducts did not show any blue signal,

verifying the specificity of duct labeling of X-gal staining. Scale=500um

Supplementary Fig.3 Heatmap of the chromatin contacts surrounding *Oxtr* and *Oxt*

across mouse tissues/cells from megabase-size visualization. (A) Heatmap of the

chromatin contacts surrounding *Oxtr* across mouse tissues/cells: *Oxtr* is located at a TAD

boundary region. The increase of interactions characterizes the TADs structure and the

interaction of two neighboring TADs can be more clearly observed in brain-related

samples. Blue dashed lines were prepared to help observing the TAD structures, and

black circles were prepared to help observing the long-distance interaction obtained from

Hi-C interaction matrices surrounding *Oxtr*. (B) Heatmap of the chromatin contacts

surrounding *Oxt* across mouse tissues/cells. No TAD structure was observed surrounding

Oxt locus.

Supplementary Fig.4 Heatmap of the chromatin contacts surrounding *OXTR* and

***OXT* across human tissues/cells from megabase-size visualization.**

(A) Heatmap of the chromatin contacts surrounding *OXTR* across human tissues/cells.
The human *OXTR* is also located at a TAD boundary region. Blue dashed lines were
prepared to help observing the large TAD structure obtained from Hi-C interaction
matrices. (B) Heatmap of the chromatin contacts surrounding *OXT* across human
tissues/cells. No typical TAD structure was observed surrounding *OXT* locus.
772

Supplementary Fig.5 Quantitative PCR analysis of *Oxtr* mRNA expression in
different brain regions revealed by common primer set recognizing both m*Oxtr* and
pv*Oxtr*. qPCR was used to analyzed Oxtr mRNA in olfactory bulb, prefrontal cortex,
striatum and thalamus from WT, Koi-1, Koi-2, Koi-3, Koi-4 and *mOxtr*$^{-/-}$ mutant mice,
n=4 for each genotype. Note that the level of *Oxtr* mRNA in *mOxtr*$^{-/-}$ mutant mice was
nondetectable (ND). The autoradiograph of OXTR binding in olfactory bulb from each
line were shown alongside the legend. There was a significant interaction between
position and genotype $F_{(12,64)} = 26.33$, $p < 0.0001$; Post hoc Bonferroni test revealed that
Oxtr expression level in olfactory bulb of Koi-1 was significantly higher than that of
other genotype groups ($p < 0.0001$ for all), and that of Koi-4 was significantly higher than
that of Koi-3 $p < 0.05$. And the striatum of Koi-4 expressed significantly higher *Oxtr* when
compared with WT and Koi-3($p < 0.05$ for both). The *Oxtr* expression level in thalamus of
Koi-4 was significantly higher than other genotype groups ($p < 0.0001$ when compared
with WT and Koi-3, $p < 0.001$ when compared with Koi-2, $p < 0.05$ when comparedg with
Koi-1.) Significance of interaction between genotype and brain region: # # # # $p < 0.0001$.
The significance of single main effect of genotype: ✶$p < 0.05$; ✶✶$p < 0.01$; ✶✶✶
$p < 0.001$; ✶✶✶✶$p < 0.0001$.
790

Supplementary Fig.6 Quantitative analysis of *Oxtr* mRNA expression in different
brain regions revealed by a primer set specifically recognizing pv*Oxtr*. The tissue of
olfactory bulb, prefrontal cortex, striatum and thalamus from WT, Koi-1, Koi-2, Koi-3
and Koi-4 mice were analyzed, n=4 for each genotype. Note that the level of *Oxtr*
mRNA in WT mice was barely detectable. There was a significant interaction between
position and genotype $F_{(9,48)} = 89.25$, $p < 0.0001$; Post Hoc Bonferroni test revealed that
Oxtr expression level in the olfactory bulb of Koi-1 was significantly higher than that of
other genotype groups ($p < 0.0001$ for all), and that of Koi-4 was significantly higher than
that of Koi-2 and Koi-3 ($p < 0.0001$ for both). The *Oxtr* expression level in prefrontal
cortex of Koi-3 was significantly higher than that of other groups ($p < 0.01$ when
comparing with Koi-1 and Koi-4; $p < 0.05$ when comparing with Koi-2). And the striatum
of Koi-4 expressed significantly higher *Oxtr* when comparing with Koi-2 and Koi-3

(p<0.01 for both). The *Oxtr* expression level in thalamus of Koi-4 was significantly

higher than that of other genotype groups (p<0.0001 for all). The significance of

interaction between genotype and brain region: # # # # p<0.0001. The significance of

single main effect of genotype: ★p<0.05; ★★p<0.01; ★★★p<0.001; ★★★★

p<0.0001.

Supplementary Table.1

The datasets IDs and sample information for mouse 3D chromatin verification

The database source of each mouse sample, analysis and visualization tool information,

and the region analyzed was listed.

Supplementary Table.2

The datasets IDs and sample information for human 3D chromatin verification

The database source of each human sample, analysis and visualization tool information,

and the region analyzed was listed.

References:

1.   Jurek, B., and Neumann, I.D. (2018). The Oxytocin Receptor: From Intracellular

Signaling to Behavior. Physiol Rev *98*, 1805-1908.

2.   Froemke, R.C., and Young, L.J. (2021). Oxytocin, Neural Plasticity, and Social Behavior.

Annu Rev Neurosci *44*, 359-381.

3.   Rigney, N., de Vries, G.J., Petrulis, A., and Young, L.J. (2022). Oxytocin, Vasopressin,

and Social Behavior: From Neural Circuits to Clinical Opportunities. Endocrinology *163*.

4.   Young, L.J., and Zhang, Q.I. (2021). On the Origins of Diversity in Social Behavior. The

Japanese Journal of Animal Psychology *doi: 10.2502/janip.71.1.4*.

5.   Barrett, C.E., Arambula, S.E., and Young, L.J. (2015). The oxytocin system promotes

resilience to the effects of neonatal isolation on adult social attachment in female prairie

voles. Transl Psychiatry *5*, e606.

6.   Ahern, T.H., Olsen, S., Tudino, R., and Beery, A.K. (2021). Natural variation in the

oxytocin receptor gene and rearing interact to influence reproductive and nonreproductive

social behavior and receptor binding. Psychoneuroendocrinology *128*, 105209.

7.   Ross, H.E., Freeman, S.M., Spiegel, L.L., Ren, X., Terwilliger, E.F., and Young, L.J.

(2009). Variation in oxytocin receptor density in the nucleus accumbens has differential

effects on affiliative behaviors in monogamous and polygamous voles. J Neurosci *29*, 1312-

1318.

8.   King, L.B., Walum, H., Inoue, K., Eyrich, N.W., and Young, L.J. (2016). Variation in the
Oxytocin Receptor Gene Predicts Brain Region-Specific Expression and Social
Attachment. Biol Psychiatry *80*, 160-169.

9.   Rogers Flattery, C.N., Coppeto, D.J., Inoue, K., Rilling, J.K., Preuss, T.M., and Young,
L.J. (2022). Distribution of brain oxytocin and vasopressin V1a receptors in chimpanzees
(Pan troglodytes): comparison with humans and other primate species. Brain Struct Funct
227, 1907-1919.

10.   Skuse, D.H., Lori, A., Cubells, J.F., Lee, I., Conneely, K.N., Puura, K., Lehtimäki, T.,
Binder, E.B., and Young, L.J. (2014). Common polymorphism in the oxytocin receptor
gene (*OXTR*) is associated with human social recognition skills. Proceedings of
the National Academy of Sciences *111*, 1987-1992.

11.   Theofanopoulou, C., Andirkó, A., Boeckx, C., and Jarvis, E.D. (2022). Oxytocin and
vasotocin receptor variation and the evolution of human prosociality. Comprehensive
Psychoneuroendocrinology *11*, 100139.

12.   Keebaugh, A.C., and Young, L.J. (2011). Increasing oxytocin receptor expression in the
nucleus accumbens of pre-pubertal female prairie voles enhances alloparental
responsiveness and partner preference formation as adults. Horm Behav *60*, 498-504.

13.   Keebaugh, A.C., Barrett, C.E., Laprairie, J.L., Jenkins, J.J., and Young, L.J. (2015). RNAi
knockdown of oxytocin receptor in the nucleus accumbens inhibits social attachment and
parental care in monogamous female prairie voles. Soc Neurosci *10*, 561-570.

14.   Berendzen, K.M., Sharma, R., Mandujano, M.A., Wei, Y., Rogers, F.D., Simmons, T.C.,
Seelke, A.M.H., Bond, J.M., Larios, R., Goodwin, N.L., et al. (2023). Oxytocin receptor is
not required for social attachment in prairie voles. Neuron *111*, 787-796.e784.

15.   Mack, K.L., Campbell, P., and Nachman, M.W. (2016). Gene regulation and speciation in
house mice. Genome Res *26*, 451-461.

16.   Osada, N., Miyagi, R., and Takahashi, A. (2017). Cis- and Trans-regulatory Effects on
Gene Expression in a Natural Population of Drosophila melanogaster. Genetics *206*,
2139-2148.

17.   Shi, X., Ng, D.W.K., Zhang, C., Comai, L., Ye, W., and Jeffrey Chen, Z. (2012). Cis- and
trans-regulatory divergence between progenitor species determines gene-expression
novelty in Arabidopsis allopolyploids. Nature Communications *3*, 950.

18.   Signor, S.A., and Nuzhdin, S.V. (2018). The Evolution of Gene Expression in cis and
trans. Trends Genet *34*, 532-544.

19.   Suster, M.L., Abe, G., Schouw, A., and Kawakami, K. (2011). Transposon-mediated BAC
transgenesis in zebrafish. Nature Protocols *6*, 1998-2021.

20.   Beil, J., Fairbairn, L., Pelczar, P., and Buch, T. (2012). Is BAC transgenesis obsolete?
State of the art in the era of designer nucleases. J Biomed Biotechnol *2012*, 308414.

21.   Bian, Q., and Belmont, A.S. (2010). BAC TG-EMBED: one-step method for high-level,
copy-number-dependent, position-independent transgene expression. Nucleic Acids
Research *38*, e127-e127.

22.   Schmidt, E.F., Kus, L., Gong, S., and Heintz, N. (2013). BAC transgenic mice and the
GENSAT database of engineered mouse strains. Cold Spring Harb Protoc *2013*.

23.   Gerfen, C.R., Paletzki, R., and Heintz, N. (2013). GENSAT BAC cre-recombinase driver
lines to study the functional organization of cerebral cortical and basal ganglia circuits.
Neuron *80*, 1368-1383.

24.   Heintz, N. (2004). Gene Expression Nervous System Atlas (GENSAT). Nature
Neuroscience *7*, 483-483.

25.   Gong, S., Doughty, M., Harbaugh, C.R., Cummins, A., Hatten, M.E., Heintz, N., and
Gerfen, C.R. (2007). Targeting Cre recombinase to specific neuron populations with
bacterial artificial chromosome constructs. J Neurosci *27*, 9817-9823.

26.   Ferrante, R.J. (2009). Mouse models of Huntington's disease and methodological
considerations for therapeutic trials. Biochimica et Biophysica Acta (BBA) - Molecular
Basis of Disease *1792*, 506-520.

27.   Shenoy, S.A., Zheng, S., Liu, W., Dai, Y., Liu, Y., Hou, Z., Mori, S., Tang, Y., Cheng, J.,
Duan, W., et al. (2022). A novel and accurate full-length HTT mouse model for
Huntington's disease. Elife *11*, e70217.

28.   Crook, Z.R., and Housman, D. (2011). Huntington's disease: can mice lead the way to
treatment? Neuron *69*, 423-435.

29.   Taguchi, T., Ikuno, M., Hondo, M., Parajuli, L.K., Taguchi, K., Ueda, J., Sawamura, M.,
Okuda, S., Nakanishi, E., Hara, J., et al. (2019). α-Synuclein BAC transgenic mice exhibit
RBD-like behaviour and hyposmia: a prodromal Parkinson's disease model. Brain *143*,
249-265.

30.   Yang, X.W., and Lu, X.-H. (2008). Chapter 19 - The Bac Transgenic Approach to Study
Parkinson's Disease in Mice. In Parkinson's Disease, R. Nass and S. Przedborski, eds.
(San Diego: Academic Press), pp. 247-268.

31.   McGraw, L.A., Davis, J.K., Lowman, J.J., ten Hallers, B.F.H., Koriabine, M., Young, L.J.,
de Jong, P.J., Rudd, M.K., and Thomas, J.W. (2010). Development of genomic resources
for the prairie vole (Microtus ochrogaster): construction of a BAC library and vole-mouse
comparative cytogenetic map. BMC Genomics *11*, 70.

32.   McGraw, L.A., Davis, J.K., Thomas, P.J., Young, L.J., and Thomas, J.W. (2012). BAC-
based sequencing of behaviorally-relevant genes in the prairie vole. PLoS One *7*, e29345.

33.   Zhang, Q., Sano, C., Masuda, A., Ando, R., Tanaka, M., and Itohara, S. (2016). Netrin-G1
regulates fear-like and anxiety-like behaviors in dissociable neural circuits. Scientific
Reports *6*, 28750.

34.   Kandasamy, L.C., Tsukamoto, M., Banov, V., Tsetsegee, S., Nagasawa, Y., Kato, M.,
Matsumoto, N., Takeda, J., Itohara, S., Ogawa, S., et al. (2021). Limb-clasping, cognitive
deficit and increased vulnerability to kainic acid-induced seizures in neuronal
glycosylphosphatidylinositol deficiency mouse models. Human Molecular Genetics *30*,
758-770.

35.   Newmaster, K.T., Nolan, Z.T., Chon, U., Vanselow, D.J., Weit, A.R., Tabbaa, M.,
Hidema, S., Nishimori, K., Hammock, E.A.D., and Kim, Y. (2020). Quantitative cellular-
resolution map of the oxytocin receptor in postnatally developing mouse brains. Nature
Communications *11*, 1885.

36.   Inoue, K., Ford, C.L., Horie, K., and Young, L.J. (2022). Oxytocin receptors are widely
distributed in the prairie vole (Microtus ochrogaster) brain: Relation to social behavior,
genetic polymorphisms, and the dopamine system. Journal of Comparative Neurology *530*,
2881-2900.

37.   Smirnov, A., Fishman, V., Yunusova, A., Korablev, A., Serova, I., Skryabin, B.V.,
Rozhdestvensky, T.S., and Battulin, N. (2019). DNA barcoding reveals that injected
transgenes are predominantly processed by homologous recombination in mouse zygote.
Nucleic Acids Research *48*, 719-735.

38.   Chandler, K.J., Chandler, R.L., Broeckelmann, E.M., Hou, Y., Southard-Smith, E.M., and
Mortlock, D.P. (2007). Relevance of BAC transgene copy number in mice: transgene copy
number variation across multiple transgenic lines and correlations with transgene integrity
and expression. Mamm Genome *18*, 693-708.

39.   Dekker, J., Rippe, K., Dekker, M., and Kleckner, N. (2002). Capturing chromosome
conformation. Science *295*, 1306-1311.

40.   Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S., and Ren, B.
(2012). Topological domains in mammalian genomes identified by analysis of chromatin
interactions. Nature *485*, 376-380.

41.   Rajderkar, S., Barozzi, I., Zhu, Y., Hu, R., Zhang, Y., Li, B., Alcaina Caro, A., Fukuda-
Yuzawa, Y., Kelman, G., Akeza, A., et al. (2023). Topologically associating domain
boundaries are required for normal genome function. Communications Biology *6*, 435.

42.   Beagan, J.A., and Phillips-Cremins, J.E. (2020). On the existence and functionality of
topologically associating domains. Nature Genetics *52*, 8-16.

43.   Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling,
A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner, M.O., et al. (2009). Comprehensive

Mapping of Long-Range Interactions Reveals Folding Principles of the Human Genome.
Science *326*, 289-293.

44.   Gibcus, Johan H., and Dekker, J. (2013). The Hierarchy of the 3D Genome. Molecular
Cell *49*, 773-782.

45.   Tribollet, E., Charpak, S., Schmidt, A., Dubois-Dauphin, M., and Dreifuss, J.J. (1989).
Appearance and transient expression of oxytocin receptors in fetal, infant, and
peripubertal rat brain studied by autoradiography and electrophysiology. J Neurosci *9*,
1764-1773.

46.   Inoue, K., Ford, C.L., Horie, K., and Young, L.J. Oxytocin receptors are widely distributed
in the prairie vole (Microtus ochrogaster) brain: Relation to social behavior, genetic
polymorphisms, and the dopamine system. Journal of Comparative Neurology *n/a*.

47.   Walum, H., and Young, L.J. (2018). The neural mechanisms and circuitry of the pair
bond. Nat Rev Neurosci *19*, 643-654.

48.   Olazábal, D.E., and Young, L.J. (2006). Oxytocin receptors in the nucleus accumbens
facilitate "spontaneous" maternal behavior in adult female prairie voles. Neuroscience *141*,
559-568.

49.   Carcea, I., Caraballo, N.L., Marlin, B.J., Ooyama, R., Riceberg, J.S., Mendoza Navarro,
J.M., Opendak, M., Diaz, V.E., Schuster, L., Alvarado Torres, M.I., et al. (2021). Oxytocin
neurons enable social transmission of maternal behaviour. Nature *596*, 553-557.

50.   Marlin, B.J., Mitre, M., D'Amour J, A., Chao, M.V., and Froemke, R.C. (2015). Oxytocin
enables maternal behaviour by balancing cortical inhibition. Nature *520*, 499-504.

51.   Olazábal, D.E., and Young, L.J. (2006). Species and individual differences in juvenile
female alloparental care are associated with oxytocin receptor density in the striatum and
the lateral septum. Horm Behav *49*, 681-687.

52.   Ophir, A.G., Gessel, A., Zheng, D.J., and Phelps, S.M. (2012). Oxytocin receptor density
is associated with male mating tactics and social monogamy. Horm Behav *61*, 445-453.

53.   Walum, H., and Young, L.J. (2018). The neural mechanisms and circuitry of the pair
bond. Nature Reviews Neuroscience *19*, 643-654.

54.   Amadei, E.A., Johnson, Z.V., Jun Kwon, Y., Shpiner, A.C., Saravanan, V., Mays, W.D.,
Ryan, S.J., Walum, H., Rainnie, D.G., Young, L.J., et al. (2017). Dynamic corticostriatal
activity biases social bonding in monogamous female prairie voles. Nature *546*, 297-301.

55.   Shapiro, L.E., and Insel, T.R. (1989). Ontogeny of oxytocin receptors in rat forebrain: a
quantitative study. Synapse *4*, 259-266.

56.   Tribollet, E., Charpak, S., Schmidt, A., Dubois-Dauphin, M., and Dreifuss, J. (1989).
Appearance and transient expression of oxytocin receptors in fetal, infant, and

peripubertal rat brain studied by autoradiography and electrophysiology. The Journal of
Neuroscience *9*, 1764-1773.

57.   Hammock, E.A., and Levitt, P. (2013). Oxytocin receptor ligand binding in embryonic
tissue and postnatal brain development of the C57BL/6J mouse. Front Behav Neurosci *7*,
195.

58.   Nardou, R., Lewis, E.M., Rothhaas, R., Xu, R., Yang, A., Boyden, E., and Dölen, G.
(2019). Oxytocin-dependent reopening of a social reward learning critical period with
MDMA. Nature *569*, 116-120.

59.   Hammock, E.A.D. (2015). Developmental Perspectives on Oxytocin and Vasopressin.
Neuropsychopharmacology *40*, 24-42.

60.   Young, L.J., and Crews, D. (1995). Comparative neuroendocrinology of steroid receptor
gene expression and regulation: Relationship to physiology and behavior. Trends
Endocrinol Metab *6*, 317-323.

61.   Ross, H.E., and Young, L.J. (2009). Oxytocin and the neural mechanisms regulating social
cognition and affiliative behavior. Front Neuroendocrinol *30*, 534-547.

62.   Venkatesh, B., Si-Hoe, S.L., Murphy, D., and Brenner, S. (1997). Transgenic rats reveal
functional conservation of regulatory controls between the Fugu isotocin and rat oxytocin
genes. Proc Natl Acad Sci U S A *94*, 12462-12466.

63.   Gilligan, P., Brenner, S., and Venkatesh, B. (2003). Neurone-specific expression and
regulation of the pufferfish isotocin and vasotocin genes in transgenic mice. J
Neuroendocrinol *15*, 1027-1036.

64.   Knobloch, H.S., Charlet, A., Hoffmann, Lena C., Eliava, M., Khrulev, S., Cetin, Ali H.,
Osten, P., Schwarz, Martin K., Seeburg, Peter H., Stoop, R., et al. (2012). Evoked Axonal
Oxytocin Release in the Central Amygdala Attenuates Fear Response. Neuron *73*, 553-
566.

65.   Gong, S., Zheng, C., Doughty, M.L., Losos, K., Didkovsky, N., Schambra, U.B., Nowak,
N.J., Joyner, A., Leblanc, G., Hatten, M.E., et al. (2003). A gene expression atlas of the
central nervous system based on bacterial artificial chromosomes. Nature *425*, 917-925.

66.   Gawenis, L.R., Hodges, C.A., McHugh, D.R., Valerio, D.M., Miron, A., Cotton, C.U., Liu,
J., Walker, N.M., Strubberg, A.M., Gillen, A.E., et al. (2019). A BAC Transgene
Expressing Human CFTR under Control of Its Regulatory Elements Rescues Cftr
Knockout Mice. Scientific Reports *9*, 11828.

67.   Xiao, J.Y., Hafner, A., and Boettiger, A.N. (2021). How subtle changes in 3D structure
can create large changes in transcription. Elife *10*, e64320.

68.   Lee, M., Lori, A., Langford, N.A., and Rilling, J.K. (2022). Enhanced endogenous oxytocin
signaling in the brain modulates neural responses to social misalignment and promotes

conformity in humans: A multi-locus genetic profile approach. Psychoneuroendocrinology
144, 105869.

69.   Hernandez, L.M., Lawrence, K.E., Padgaonkar, N.T., Inada, M., Hoekstra, J.N., Lowe,
J.K., Eilbott, J., Jack, A., Aylward, E., Gaab, N., et al. (2020). Imaging-genetics of sex
differences in ASD: distinct effects of OXTR variants on brain connectivity. Transl
Psychiatry *10*, 82.

70.   Vietri Rudan, M., Barrington, C., Henderson, S., Ernst, C., Odom, Duncan T., Tanay, A.,
and Hadjur, S. (2015). Comparative Hi-C Reveals that CTCF Underlies Evolution of
Chromosomal Domain Architecture. Cell Reports *10*, 1297-1309.

71.   Kim, J.H., Lee, S.R., Li, L.H., Park, H.J., Park, J.H., Lee, K.Y., Kim, M.K., Shin, B.A., and
Choi, S.Y. (2011). High cleavage efficiency of a 2A peptide derived from porcine
teschovirus-1 in human cell lines, zebrafish and mice. PLoS One *6*, e18556.

72.   Hidema, S., Fukuda, T., Hiraoka, Y., Mizukami, H., Hayashi, R., Otsuka, A., Suzuki, S.,
Miyazaki, S., and Nishimori, K. (2016). Generation of Oxtr cDNA(HA)-Ires-Cre Mice for
Gene Expression in an Oxytocin Receptor Specific Manner. J Cell Biochem *117*, 1099-
1111.

73.   Takayanagi, Y., Yoshida, M., Bielsky, I.F., Ross, H.E., Kawamata, M., Onaka, T.,
Yanagisawa, T., Kimura, T., Matzuk, M.M., Young, L.J., et al. (2005). Pervasive social
deficits, but normal parturition, in oxytocin receptor-deficient mice. Proceedings of the
National Academy of Sciences *102*, 16096-16101.

74.   Wang, Y., Song, F., Zhang, B., Zhang, L., Xu, J., Kuang, D., Li, D., Choudhary, M.N.K.,
Li, Y., Hu, M., et al. (2018). The 3D Genome Browser: a web-based browser for
visualizing 3D genome organization and long-range chromatin interactions. Genome Biol
19, 151.

75.   Li, D., Purushotham, D., Harrison, J.K., Hsu, S., Zhuo, X., Fan, C., Liu, S., Xu, V., Chen,
S., Xu, J., et al. (2022). WashU Epigenome Browser update 2022. Nucleic Acids Research
50, W774-W781.

76.   Stolzenberg, D.S., Stevens, J.S., and Rissman, E.F. (2012). Experience-facilitated
improvements in pup retrieval; evidence for an epigenetic effect. Horm Behav *62*, 128-
135.

1046

1047

1048

1049 **Fig.1 The construction of Koi lines**

1050

1051

Fig.2 Transgenic CRE mediated Lac-Z reporter gene expression in the brain and mammary gland of Koi lines.

1054

1055

1056   **Fig.3The expression of OXTR in the brain of Koi lines.**

1057

1058

1059   **Fig.4 Social behaviors of Koi lines.**

1060

Supplementary Materials:

1062

1063

Supplementary Fig.1Transgenic CRE mediated Lac-Z reporter gene expression in the brain of Koi lines.

1066

1067 **Supplementary Fig.2 Negative control of whole-mount mammary gland staining.**

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089
1090
1091

1092

1093

1094

Supplementary Fig.3 Heatmap of the chromatin contacts surrounding *Oxtr* (A) and *Oxt* (B) across mouse tissues/cells from megabase-size visualization.

1097

1098

1099

1100

1101

1102

1103 **Supplementary Fig.4 Heatmap of the chromatin contacts surrounding *OXTR* (C)**

1104 **and *OXT* (D) across human tissues from megabase-size visualization.**

1105

1106

1107 **Supplementary Fig.5 Quantitative analysis of *Oxtr* mRNA expression in different**

1108 **brain regions revealed by common primer set recognizing both *mOxtr* and *pvOxtr*.**

1109

Supplementary Fig.6 Quantitative analysis of *Oxtr* mRNA expression in different brain regions revealed by a primer set specifically recognizing pv*Oxtr*.

1132

	track	ID	Cell/tissue	Organism	Resolution	Restr. Enz.	Database	Database ID	Browser	Publication
Hi-C	1	Whole brain	whole brain	mm10	40 Kb	DNaseI	4DN	4DNFIQ48NYOW	WashU	Deng 2015
	2	NPC	NPC	mm10	40 Kb	DpnII	4DN	4DNFIHW8NTQX	WashU	Bonev 2017
	3	Cortex	cortex	mm10	40 Kb	MboI	4DN	4DNFII3JV8I1	WashU	Du 2017
	4	ESC	ESC	mm10	40 Kb	DpnII	4DN	4DNFIU8AF5ZY	WashU	Bonev 2017
	5	Hepatocyte	hepatocyte	mm10	40 Kb	HindIII	4DN	4DNFIFULDMGN	WashU	Schwarzer 2017
	6	Sperm	sperm	mm10	40 Kb	MboI	4DN	4DNFIN8F14CS	WashU	Du 2017
WashU	Fig	Regions	Genes							
	A	chr6:104149809-119528368	Oxtr							
	B	chr2:121804885-139358632	Oxt							

1133

1134

Supplementary Table.1

The datasets IDs and sample information for mouse 3D chromatin verification

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

	track	ID	Cell/tissue	Organism	Resolution	Database	Publication
Hi-C	1	Fetal brain	Fetal Brain GZ	hg19	10 Kb	3D genome	Won et al, 2016
	2	Hippocampus	Human hippocampus	hg19	40 Kb	3D genome	Schmitt et al., 2016
	3	Cortex	Cortex	hg19	40 Kb	3D genome	Schmitt et al., 2016
WashU	Fig	Regions	Genes				
	A	chr3:1831010-15559067	OXTR				
	B	chr20:0-15000000	OXT				

1148

1149

Supplementary Table.2

The datasets IDs and sample information for human 3D chromatin verification

1152