Cross-Species BAC Transgenesis Reveals Long-Range Regulation Drives

Variation in Brain Oxytocin Receptor Expression and Social Behaviors

Mina Tsukamoto [1], Luis AE Nagai[2], Kiyoshi Inoue[3], Lenin C. Kandasamy [1], Maria F.

Pires[1], Minsoo Shin[1], Yutaro Nagasawa[1], Tsetsegee Sambuu[1], Kenta Nakai[4], Shigeyoshi

Itohara[5], Larry J Young[3,6,7#], Qi Zhang[1,5,6,8*]

[1]Laboratory for Social Neural Networks, Faculty of Human Sciences, University of

Tsukuba, Japan305-8577; [2] Laboratory of Computational Genomics, Institute for

Quantitative Biosciences, The University of Tokyo, Japan113-0032; [3]Center for

Translational Social Neuroscience, Emory National Primate Research Center, Emory

University, USA30329; [4]Laboratory of Functional Analysis in silico, Human Genome

Center, The University of Tokyo, Tokyo, Japan108-8639; [5]Laboratory for Behavioral

Genetics, CBS, RIKEN, Japan351-0198; [6]Center for Social Neural Networks, Faculty of

Human Sciences, University of Tsukuba, Japan305-8577; [7]Department of Psychiatry and

Behavioral Science, Emory University, USA30329; [8] Institute for Glyco-core Research,

Tokai National Higher Education and Research System, Gifu University, Japan501-1193

[#]Deceased in March, 2024. **We dedicate this work to the memory of Dr. Larry J.**

Young, whose encouragement and unwavering support helped our team persevere

through challenges.

*Correspondence: zhang.qi.b3@f.gifu-u.ac.jp or zhangqi@human.tsukuba.ac.jp

or qi.zhang@a.riken.jp

Classification: Biological Science: Neuroscience

Keywords: oxytocin system, behavioral evolution, bacterial artificial chromosome

(BAC), brain, mammary gland, partner preference, maternal care, topologically

associating domain (TAD), evolvability.

The PDF file includes:

Main Text

Figures 1 to 7

Supplementary Figures 1to 3, supplementary Tables 1

Abstract

Although oxytocin (OXT) exhibits a highly conserved neuroanatomical pattern among

vertebrates, the distribution of OXT receptor (OXTR) in brain varies considerably across

species and is associated with species-typical social behavior. To investigate the genomic

basis of the phylogenetic plasticity in brain *Oxtr* expression and its social behavioral

consequences, we generated transgenic mice carrying a bacteria artificial chromosome

(BAC) harboring the entire prairie vole *Oxtr* locus and flanking intergenic regulatory

regions. We established eight independent "volized" mouse lines expressing prairie vole

Oxtr (pv*Oxtr*). Strikingly, despite conserved *Oxtr* expression in mammary gland of all

transgenic mouse lines, each line displayed a unique pattern of brain expression distinct

from both mice and prairie voles. Together with topologically associating domain (TAD)

structure analysis with mouse genome, our findings suggest that unlike *Oxt*, *Oxtr*

expression patterns in brain, involve contributions of distal regulatory elements beyond

the BAC insert. In contrast, *Oxtr* expression in peripheral tissues appears resistant to such

distal influences. Moreover, the "volized" mouse lines with different brain *Oxtr*

expression patterns showed differences in partner preference and maternal behaviors,

providing direct functional evidence that variation in brain *Oxtr* expression can drive

differences in social behaviors. We propose that brain *Oxtr* expression is transcriptionally

sensitive to long-range interactions with distal genomic elements, rendering it more

susceptible to diverse regulatory influences. This supports a model in which regulatory

flexibility facilitates the evolutionary diversification of social behavior, while maintaining

essential peripheral *Oxtr* expression.

Highlights

Unlike essential physiological phenomenon including feeding, sex, parturition and

lactation, which are conserved across mammalian species, there is extraordinary diversity

of social behaviors between and within species. Comparative studies across species suggest

that variation in brain oxytocin receptor expression may mediate diversity in social

behavior. Subtle variations in human oxytocin receptor sequence have been related to

psychiatric phenotypes. Our studies suggest that the oxytocin receptor gene is

hypersensitive to long-distance sequence variation, as revealed by differential expression

patterns when the BAC transgene was inserted at different genomic sites, leading to

variability in brain expression and social behavior. This research provides new insights into

the evolvability of genes producing diversity in social behaviors, which allows efficient

adaptation of animals to variable environment. Furthermore, our cross-species study

demonstrates that diversity in brain OXTR expression patterns leads to variation in mouse

social behaviors. The "volized" *Oxtr* mouse lines generated here present a valuable

resource for further exploring OXTR expression regulation and neural circuits/networks

mediating social behavioral variability.

Introduction

Brain oxytocin receptors (OXTR) regulate a wide range of social behaviors including

social recognition, maternal care, social bonding, empathy related behaviors, and

aggression (Froemke & Young, 2021; Jurek & Neumann, 2018; Rigney, de Vries,

Petrulis, & Young, 2022). In contrast to sex steroid receptors, which have highly

conserved brain expression patterns across species, OXTR shows remarkable inter- and

intra-species variation in brain distribution (Froemke & Young, 2021; Larry J. Young &

Zhang, 2021). For example, monogamous prairie voles have a brain OXTR distribution in

cortex, striatum and amygdala that is significantly distinct from promiscuous montane

voles and laboratory rats and mice(Larry J. Young & Zhang, 2021). Furthermore, there is

robust individual variation in brain OXTR expression among prairie voles that is

associated with single nucleotide polymorphisms (SNPs) in the OXTR locus (*Oxtr*), and

this variation predicts pair bonding behavior and resilience to early life social neglect

(Ahern, Olsen, Tudino, & Beery, 2021; Barrett, Arambula, & Young, 2015; King, Walum,

Inoue, Eyrich, & Young, 2016; Ross et al., 2009). Different species of primates also differ

in brain OXTR distribution (Rogers Flattery et al., 2022), and SNPs in the human OXTR

have been linked to variation in social function (Skuse et al., 2014; Theofanopoulou,

Andirkó, Boeckx, & Jarvis, 2022). Thus, variation in OXTR distribution in brain likely

constitutes a mechanism for the emergence of diverse social traits, potentially including

psychiatric endophenotypes.

The evolution of gene expression patterns could be mediated by *cis* (via linked

polymorphisms) or *trans* (through diffusible products of distal genes, e.g., transcription

factors) changes, as a result of adaptation to the environment (Mack, Campbell, &

Nachman, 2016; Osada, Miyagi, & Takahashi, 2017; Shi et al., 2012). *Cis*-regulatory

differences are more commonly responsible for adaptive evolution and interspecific

divergence (Signor & Nuzhdin, 2018). To explore the transcriptional mechanisms giving

rise to species-specific *Oxtr* expression and social behavior, we created transgenic mice

using a prairie vole *Oxtr* (pv*Oxtr*) bacterial artificial chromosome (BAC). The BAC

construct covers the full coding sequence, introns and the entire intergenic region of the

pv*Oxtr* with its neighboring genes. We expected that our BAC construct accommodates

all the critical regulatory sequences responsible for species- and tissue-specific

expression, so that pv*Oxtr* transgenic mice would express brain pv*Oxtr* in a prairie vole-

like pattern, largely independent of integration sites. Then we would examine if our

transgenic mouse could develop social behaviors similar to that of prairie vole.

We generated eight pv*Oxtr* mouse lines with successful germline transmission. Each line,

defined by a distinct BAC integration site, exhibited a unique brain-specific expression

pattern of *Oxtr*, distinct from both wildtype mice and prairie voles. Notably, *Oxtr*

expression in the mammary gland was conserved across all lines. This suggests that brain,

but not mammary gland, expression of *Oxtr* is shaped by distal regulatory elements

beyond the BAC insert. These findings imply that evolution has placed brain *Oxtr*

expression under more complex regulatory control, likely to accommodate the

diversification of social behaviors, whereas mammary gland expression is maintained by

simpler mechanisms supporting conserved lactation functions. Furthermore, behavioral

assays revealed line-specific differences in partner preference and maternal behaviors.

The 3D architecture of the genome, segregated into TADs demarcated by CCCTC-

binding factor (CTCF), crucially influences gene expression. Through the loop extrusion

model, CTCF enables interactions between distant regulatory elements, facilitating

precise gene regulation, which ultimately shapes an organism's phenotype and

behavior(Wendt et al., 2008). Hi-C data provide a genome-wide view of chromosome

conformation in multiple layers, such as compartments, TADs, and loops(Gibcus &

Dekker, 2013; Lieberman-Aiden et al., 2009). Our TAD analysis with Hi-C database

showed a clear difference exists between the 3D structures of *Oxt* and *Oxtr* gene, and a

clear difference exists between the 3D structures of brain tissue *Oxtr* and peripheral tissue

Oxtr, suggesting that Oxtr expression patterns in brain depends on contributions from

very distal sequences. These results provide important clues to understand the genetic

mechanism underlying species differences in brain gene expression.

Together, our data demonstrate that transcriptional redistribution of brain *Oxtr* can drive

variation in social and parental behaviors—variation that natural selection could act upon

in specific ecological contexts. Our results also offer insights into the genetic mechanisms

underlying species-specific brain gene expression.

Results

Creation of the pv*Oxtr-P2A-Cre* BAC transgenic mice

BAC vectors can accommodate dispersed cis-regulatory elements across large regions of

genome. Transgene expression from BAC constructs are generally resistant to insertion

position effects (Suster, Abe, Schouw, & Kawakami, 2011) due to both the large spans of

insulating genetic material which protect the transgene cassette from the influence of the

chromosomal environment, and through the inclusion of necessary regulatory elements

such as enhancers, silencers, locus control regions, and matrix attachment regions (Beil,

Fairbairn, Pelczar, & Buch, 2012; Bian & Belmont, 2010). Large-scale screening of CNS

gene expression in BAC transgenic mice by the GENSAT (Gene Expression Nervous

System Atlas) Project found that more than 85% of BACs express reproducibly in

multiple independent transgenic lines and reporter gene expression faithfully recapitulates

endogenous gene expression patterns (Gerfen, Paletzki, & Heintz, 2013; Heintz, 2004;

Schmidt, Kus, Gong, & Heintz, 2013). Indeed, reproducible expression from mouse BAC

vectors has been achieved for more than 500 genes (Gerfen et al., 2013; Heintz, 2004;

Schmidt et al., 2013). Few that don't express reproducibly are influenced by variation in

BAC copy number and insertion sites (S. Gong et al., 2007). BAC engineering has also

been used to mediate cross-species transgene expression. A large number of mouse

models of human dominant neurodegenerative disorders have been created using human

BAC transgene approaches. These model mice successfully express human transgenes in

human-like patterns and recapitulate disease-like phenotypes (Crook & Housman, 2011;

Ferrante, 2009; Shenoy et al., 2022; Taguchi et al., 2019; Yang & Lu, 2008). Previously

we constructed *Pkcd-BAC-Cre* transgenic mice and obtained independent lines with

reproducible expression patterns as well (Zhang et al., 2016)(Supplementary Fig.1).We

therefore decided to use BAC approaches to create "volized" *Oxtr* mouse lines.

To create pv*Oxtr-P2A-Cre* BAC transgenic mouse line, the prairie vole BAC clone we

used includes the entire pv*Oxtr* locus (the coding sequence and all the introns) with the

full intergenic sequence upstream (150kb) and downstream of the pv*Oxtr* coding region

(9kb) (Fig.1 B). The BAC clone contains a portion of *Rad18,* minus the first 11 exons*,*

located upstream of pv*Oxtr,* and *Cav3*, minus the first exon including the start codon,

downstream of pv*Oxtr.* Therefore, there will be no interference of these two flanking

transgenes on behavior analysis*,* and the proximal topological structure around pv*Oxtr*

locus is preserved to the maximum extent. A P2A-NLS-Cre cassette was inserted in-

frame just before the *Oxtr* stop codon to ensure co-expression of *Cre* and *Oxtr* (Fig.1 C).

Transgenic mice were generated by pronuclear injection using C57BL/6J zygotes. Proper

integration of the reporter into the BAC clone was confirmed using gene specific PCR

assays and sequencing (Fig.1 C, D and data not shown). We obtained 11 founder mice

carrying the transgenes, which was verified by PCR with multiple primer sets (Fig.1 C,

E). One founder line was sterile, and two founder lines did not show germ-line

transmission. Eight founder lines successfully produced offspring and stably transmitted

the transgene to at least 3 generations. We named these lines "Koi", meaning "love" in

Japanese.

Each Koi line showed a unique expression pattern of transgene in brain, yet

conserved expression in mammary gland.

We systemically evaluated 8 Koi lines by crossing them with ROSA-26-NLS-*LacZ* mice,

a Cre reporter mouse line expressing nuclear-localized beta-galactosidase (Kandasamy et

al., 2021; Zhang et al., 2016). Since Lac-Z is only expressed in the nucleus of double

positive (CRE+, LacZ+) cells, but not in the neuronal processes, we could easily assess

the expression pattern of our transgene. We found that each Koi founder line displayed a

unique pattern of expression in brain distinct from that of wildtype mice or prairie voles

(Fig.2 A-O, and supplementary Fig.2). Almost all Koi lines showed transgene expression

in olfactory bulb, lateral septum (Fig.2 F-I and Supplementary Fig.2 B,E,H,K) and

ventromedial nucleus of the hypothalamus (Fig.2 K-N and Supplementary Fig.2 C,F,I,L),

though with different intensities, suggesting that these regions constitute a stable core

Oxtr expression network, as these regions also express OXTR in mice and voles (Inoue,

Ford, Horie, & Young, 2022; Newmaster et al., 2020). Several Koi lines expressed the

transgene in some brain regions specific for vole *Oxtr* expression, including nucleus

accumbens (NAc) (Fig.2 A,B,D and Supplementary Fig.2 A,G,J), prefrontal cortex (PFC)

(Fig.2 A,B,C and Supplementary Fig.2 J), lateral amygdala (Fig.2 K,L,M and

Supplementary Fig.2 C,I,L) and deep layers of cingulate cortex (Fig.2A,B,C and

Supplementary Fig.2 C,L), which suggested that cis regulatory elements in the BAC are

capable of mediating the expression of *Oxtr* in the reward and reinforcement circuitry of

brain, albeit depending on the sequence/structure of distal sequences >150 kb upstream

and >21 kb downstream of the transcription start site. None of the Koi lines exactly

"mirrored" the vole-specific expression patten. Six of the Koi lines expressed the

transgene in broad areas of thalamus (Fig.2 K,L,N and Supplementary Fig.2 C,I,L),

reminiscent of the strong expression of V1a vasopressin receptor (*Avpr1a*) in the

thalamus (Larry J. Young & Zhang, 2021). Intriguingly, the expression of OXTR in

mammary glands showed similar pattern among all the Koi lines and *mouseOxtr(mOxtr)-*

Ires-Cre knock-in line (Fig.2 P-T and Supplementary Fig.3).

Transgenes that are introduced by pronuclear injection typically integrate into a single

site of the genome as tandem concatemers (Smirnov et al., 2019). BAC copy number

variation and different insertional loci can lead to distinct patterns of ectopic expression,

and increased BAC transgene copy numbers often correlate with increased BAC gene

expression (S. Gong et al., 2007). We then examined the relationship between transgene

copy number and LacZ signal using quantitative genomic PCR (Chandler et al., 2007). As

expected, the copy number of WT, heterozygous and homozygous *mOxtr-Ires-Cre* knock-

in mice was 0, 1 and 2 respectively (Figure.2 Z). The copy number of transgene in Koi

lines are: Koi-1 (12), Koi-2 (2), Koi-3 (2), Koi-4 (1), Koi-5 (2), Koi-6 (4), Koi-7 (9), Koi-

8 (3). Koi-6 and Koi-7 showed the most limited expression of reporter gene, though the

copy number of transgene were 4 and 9 respectively. Koi-4 showed strong expression of

reporter gene in NAc, septum and thalamus, though there was only 1 copy of transgene.

Therefore, integration site rather than copy number is the reason of the variant BAC

transgene expression patterns across our Koi lines.

Adult brain OXTR binding patterns differ across Koi lines

To detect the distribution of OXTR protein in adult Koi lines without the interference of

endogenous mouse OXTR (mOXTR) signal, Koi lines were crossed with *mOxtr*$^{-/-}$ line

twice sequentially to produce Koi: m$Oxtr^{-/-}$ mice, and pvOXTR binding was detected

using receptor autoradiography. We focused on analyzing 4 lines: Koi-1: m$Oxtr^{-/-}$, Koi-2:

m$Oxtr^{-/-}$, Koi-3: m$Oxtr^{-/-}$ and Koi-4: m$Oxtr^{-/-}$ based on the strong expression of reporter

gene in behaviorally relevant neuronal populations. Autoradiography demonstrated

functional pvOXTR in brain resembling to some extent Cre-induced reporter gene

expression (Fig.3). Brain OXTR binding of Koi-3: m$Oxtr^{-/-}$ was consistent with Lac-Z

expression in the reporter line (Fig.2 C, H, M and Figure.3 C, H, M). Interestingly, the

OXTR binding in PFC of Koi-3: m$Oxtr^{-/-}$ was similar to that of prairie vole (Figure.3 C).

In contrast, Koi-4: m$Oxtr^{-/-}$ showed low OXTR binding in NAc when compared with the

high Lac-Z staining (Fig.2 D and Figure.3 D). Koi-2: m$Oxtr^{-/-}$, the line which displayed

the expression of reporter genes in the broadest regions, did not show strong signal in

autoradiography. It is interesting that Layer IV of cortex in Koi-4: m$Oxtr^{-/-}$ showed

strong OXTR binding (Figure.3 I and N), which is highly likely to originate from the

axon projections from the thalamus (Fig.2 N), as was previously reported for NAc

projections in voles (Inoue et al., 2022). Consistent with Lac-Z expression, we do not see

a clear relationship between OXTR binding and transgene copy number. For example,

Koi-1: m$Oxtr^{-/-}$ showed rather low OXTR signal (Fig.3 A, F, K) with the highest copy

number, while Koi-4: m$Oxtr^{-/-}$ showed very strong OXTR signal at lateral septum and

thalamus (Fig.3 I, N), with the lowest copy number.

$Oxtr$ mRNA level in Koi-3: m$Oxtr^{-/-}$ and Koi-4: m$Oxtr^{-/-}$ was further investigated with

Fluorescent RNAscope $in situ$ hybridization (Fig.4). Consistent with Lac-Z expression in

the reporter line, Koi-3: m$Oxtr^{-/-}$ showed strong signal in PFC and Ctx (Fig.4 A, C), while

Koi-4: m$Oxtr^{-/-}$ showed prominent signal in NAc and Thalamus (Fig.4 B, D). qRT-PCR

analysis further revealed that adult Koi-4: m$Oxtr^{-/-}$ mice had the highest Oxtr mRNA

signal in the ventral striatum, several fold higher than WT, consistent with the Lac-Z

staining in the reporter line (Fig.5 A, B). The discrepancy of OXTR signal in NAc of

Koi-4: m$Oxtr^{-/-}$ between autoradiography and Lac-Z staining /RNAscope/qRT-PCR may,

therefore, be due to the translational regulation or differences in the sensitivity of the

detecting methods.

Although none of our Koi lines mirrored the endogenous OXTR expression in prairie

voles, we confirmed strong expression of pv$Oxtr$ in olfactory bulb of Koi-1line, PFC and

amygdala of Koi-3line, striatum and thalamus of Koi-4 lines, and the broad expression of

pv$Oxtr$ in Koi-2 line. As OXTR signaling in some of these regions are similar to that in

voles, albeit in separate lines, and postulated to be involved in vole social behavior (H.

Walum & L. J. Young, 2018), we used these lines to address the question of whether

variation in *Oxtr* transcription in brain can lead to variation in social behaviors.

Different Koi lines showed diverse behaviors in a Partner Preference Test

The partner preference test is widely used as a laboratory proxy for pair bonding in

prairie voles. In a pilot study, we found that Kio-4 female mice on a m$Oxt^{+/+}$ background

showed a preference for their mate relative to a novel male. We then focused on the

females of 4 lines (Koi-1, 2, 3, 4) based on transgene expression in PFC, BLA or NAc.

Again, we bred the Koi lines with m$Oxtr^{-/-}$ mice (both on C57/B6 background) to

generate Koi: m$Oxtr^{-/-}$ offspring expressing only pv$Oxtr$. Partner preference tests (PPT)

were performed in ovariectomized females after 21 days of cohabitation with 3 mating

bouts elicited by estrogen and progesterone injections on days 2, 9 and 16. We divided

our analysis into 3 phases (Fig.6 A, B). In the habituation phase, the experimental subject

was habituated to the chamber for 5 minutes with both pen-boxes empty. In the

recognition phase, the partner stimulus animal was restricted in one pen-box, and a novel

"stranger" male was restricted in another pen-box, and the experimental female was

allowed to explore the arena for 10 minutes to test social novelty preference. In the

preference phase, the positions of the partner stimulus animal and the "stranger" stimulus

animal were switched to avoid position bias, and the female was allowed to explore the

chamber for an additional 20 minutes.

Mixed ANOVA was conducted to examine the effect of genotype and the stimulus box on

the area stay time during all the phases. In the habituation phase (Fig.6 C), there was no

statistically significant interaction between the effects of genotype and position of empty

pen-box on stay time, $F_{(4, 60)} =0.69$, $p=0.6$. There was no difference in position preference

for all the groups, $F_{(1,60)} =0.03$, $p=0.87$. In the recognition phase (Fig.6 D), there was a

main effect of stimulus animal (i.e., partner or stranger) on stay time, the time staying in

close proximity to the partner was significantly shorter than the time stay with stranger

animal ($F_{(1,60)} =15.38$, $p=0.0003$). There was a main effect of genotype on social stay

time, $F_{(4,60)} =2.62$, $p=0.04$, and a significant interaction between the effects of genotype

and stimulus animal on area stay time, $F_{(4, 60)} =2.61$, $p=0.04$. Bonferroni adjusted

pairwise comparisons were applied to analyze the simple main effects. Koi-4: m$Oxtr^{-/-}$ and

WT mice spent significantly longer time in "stranger" area than in "partner" area ($p=0.0001$

for Koi-4: m$Oxtr^{-/-}$; $p=0.028$ for WT), demonstrating a typical novelty preference. There

was no significant difference between the stay time with "stranger" and "partner" for Koi-

1: m$Oxtr^{-/-}$ ($p=0.96$), Koi-2: m$Oxtr^{-/-}$ ($p=0.49$) and Koi-3: m$Oxtr^{-/-}$ ($p=0.09$) mice. In the

preference phase (Fig.6 E), conducted after both stimulus mice were presumably familiar

following the initial 10 min recognition phase, there was no main effect of stimulus animal

on stay time, F (1,60) =0.04, p=0.84. There was no main effect of genotype on stay time

either, F (4,60) =1.36, p=0.26. There was a statistically significant interaction between the

effects of genotype and stimulus animal on area stay time, F (4, 60) = 4.64, p=0.003.

Bonferroni adjusted pairwise comparisons revealed that Koi-2: m$Oxtr^{-/-}$ mice spent

significantly longer time in "stranger" area than in "partner" area (p=0.024). In contrast,

Koi-4: m$Oxtr^{-/-}$ mice spent significantly longer time in "partner" area than in "stranger"

area (p=0.002). There was no significant difference between the stay time with "stranger"

and "partner" for WT (p=0.99), Koi-1: m$Oxtr^{-/-}$ p=0.16) or Koi-3: m$Oxtr^{-/-}$ (p=0.45) mice.

Different Koi lines showed diverse behaviors in Pup Retrieval Test

Based on the abundant pvOXTR of Koi-3and Koi-4 in reward-related brain regions (Fig.3

C,D,H,I,M,N), and the role of OXTR in mediated maternal care in mice and voles

(Carcea et al., 2021; Marlin, Mitre, D'Amour J, Chao, & Froemke, 2015; Olazábal &

Young, 2006), we further tested the parental behaviors of these two lines in virgins (Fig.6

F, G). Mixed ANOVA was conducted to examine the effect of genotype and pup number

(1-3) on the latency of pup retrieval. Mauchly's test showed that the sphericity was

violated. Therefore, the Greenhouse-Geisser correction was used for the repeated

measures ANOVA. There was a significant main effect of genotype on latency to retrieve,

F (2,48) =29.58, p<0.0001. Koi-4: m*Oxtr*$^{-/-}$ took a significantly longer time to retrieve all

the pups than WT and Koi-3: m*Oxtr*$^{-/-}$ (p<0.0001 for both). There was also a significant

interaction between the effects of genotype and retrieved pup number on the retrieval

latency, F (2.40, 57.49) = 4.45, p =0.011. Post Hoc multiple comparisons with

Bonferroni correction revealed that Koi-4: m*Oxtr*$^{-/-}$ took a significantly longer time to

accomplish the 1st retrieval (p=0.0003 and p=0.0005), the 2nd retrieval (p<0.0001), and

the 3rd retrieval (p<0.0001) when compare to WT and Koi-3: m*Oxtr*$^{-/-}$. And Koi-3:

m*Oxtr*$^{-/-}$ took a significantly shorter time to accomplish the 1st retrieval (p=0.01), the 2nd

retrieval (p=0.006), and the 3rd retrieval (p=0.003) when compare to WT.

Crouching time was analyzed as well (Fig.6 F, H). There was a statistically significant

difference between groups as determined by one-way ANOVA (F (2,48) = 27.56, p

<0.0001). A Bonferroni post hoc test revealed that the crouching time of Koi-3: m*Oxtr*$^{-/-}$

was statistically significantly longer than WT (p=0.008) and Koi-4: m*Oxtr*$^{-/-}$ (p<0.0001).

Additionally, the crouching time of Koi-4: m*Oxtr*$^{-/-}$ was statistically significantly shorter

than WT (p=0.001). Therefore, two Koi lines with different OXTR expression pattern

demonstrated quantitative differences in parental behavior tests.

The TAD structure of brain *Oxtr* is markedly distinct from that of both *Oxt* and

peripheral *Oxtr*.

The divergent brain expression patterns across our Koi lines suggest that distal elements

external to our BAC cassette contribute to the variation in *Oxtr* expression. TADs

represent a key feature of hierarchical genome organization comprising regions of high

chromatin inter-connectivity, and TAD formation appears to be linked to transcription

regulation (Beagan & Phillips-Cremins, 2020; Dekker, Rippe, Dekker, & Kleckner, 2002;

Dixon et al., 2012; Rajderkar et al., 2023). TADs are generally conserved within the same

tissue type across different species. We therefore utilized publicly available 3D chromatin

datasets from mouse (Extended data Table7-1) and analyzed the TAD structure of *Oxtr*

and *Oxt*, both in brain related tissues/cells and peripheral tissues/cells. We discovered

that *Oxtr* is consistently located at the boundary region of TAD structures across several

tissues and cell lines (Fig.7 A). Specifically, only in brain related samples (whole brain

tissue, neural progenitor cells, and cortex tissue), *Oxtr* is situated at the boundary of a

large inter-TAD interaction between two neighboring TADs (Fig.7 A). In contrast, the

oxytocin peptide gene, *Oxt*, which is expressed in a highly conserved brain pattern across

vertebrates, lacks organized TAD structures (Fig.7 B). These findings demonstrate that

the TAD landscape differ dramatically between *Oxtr* and *Oxt*, and highlight that the 3D

genomic architecture surrounding the *Oxtr* gene varies between brain tissues/cells and

peripheral tissue/cells. There exists clear long-distance interaction in brain *Oxtr* gene

TAD structure. The presence of extensive long-distance chromatin interactions around

Oxtr in brain tissues may at least partially explain why its expression is hypersensitive to

distal regulatory sequences beyond the range of a BAC construct. Moreover, such large

inter-TAD interactions likely expose *Oxtr* regulation to a broader set of cis- and trans-

acting influences, rendering brain *Oxtr* expression inherently more variable across

individuals and species.

Discussion

Diversity of *Oxtr* expression patterns causes diversity of social behaviors

The diverse OXTR expression of our Koi lines is reminiscent of the diverse OXTR

expression across different species. The unique expression pattern of *Oxtr* in

monogamous prairie vole was mainly observed in PFC, NAc and BLA. We obtained

expression in all of these specific regions, but in 3 separate lines. These Koi lines with

distinct brain OXTR expression pattern provided an opportunity to test the effect of

variation of OXTR expression on diverse social behaviors.

It is intriguing that Koi-4 line showed preference to their mating partners but not

strangers, which is a behavior specific in prairie voles but not in mice. Multiple studies

suggested that NAc and PFC play important roles in mediating partner preference

behavior in prairie voles (Amadei et al., 2017; Hasse Walum & Larry J. Young, 2018).

Consistently, the Koi-4 line showed robust transgene expression in these brain regions.

Notably Koi-4 line also demonstrated strong *Oxtr* expression in the entire thalamus

region, which reminds us the distribution of *Avpr1a* (Phelps & Young, 2003). Given the

fact that OXTR and AVPR1A genes were derived from a duplication of a single ancestral

gene (Grinevich, Knobloch-Bollmann, Eliava, Busnelli, & Chini, 2016), it is not

surprising that the similar expression regulation machinery exists. Recent studies indeed

revealed the critical functions of different thalamic nuclei in social memory and cognition

(Chen et al., 2024; Kappel et al., 2022; Rangel, Baldo, & Canteras, 2018; Wolff & Vann,

2019). Partner preference formation and maintenance needs to recruit multi- and cross-

modality sensation, social recognition and social memory, selective attention, arousal,

and reward systems. We propose a model that the sufficient upregulation of the OXTR

expression in one or multiple of these systems can tune the network and boost partner

preference behavior. In addition, the maternal behavior in Koi-3 and Koi-4 lines was

enhanced and reduced, respectively. This result suggested that OXTR signaling facilitates

social partner preference and maternal care by coordinating activity across different

neural networks. A main conclusion of our study is that simply by redistributing OXTR

binding in space, variation in social preference and parental behaviors emerged, which

conceivably could impart advantageous behaviors that natural selection could act upon

and amplify. Future study will determine how the variation in OXTR distribution in our

Koi lines affects neural circuits mediating social behaviors.

A potential genetic mechanism contributing to diversity of brain OXTR expression

The sex steroid and oxytocin systems play important roles in modulating reproductive

and related social behaviors in vertebrates. Sexual behavior, e.g., the motivation to mate

and motor patterns, is highly conserved across vertebrate species. By contrast, social

behaviors, e.g., sociality, mating strategy (monogamy vs polygamy), parental and

alloparental behaviors are more nuanced and can vary dramatically across species in the

same genus or even between individuals of a single species. It is interesting, therefore,

that the distribution of steroid receptors is quite conserved among species (L. J. Young &

Crews, 1995), while that of the OXTR is extremely diverse across and even within

species (Froemke & Young, 2021; Larry J. Young & Zhang, 2021). It is even more

interesting that although the distribution of OXTR is diverse, its ligand, oxytocin, is

expressed in a highly conserved neuroanatomical pattern in vertebrates (Ross & Young,

2009). Previous work showed that three independent rat lines with puffer fish oxytocin

BAC transgene (containing 16 kb of 5' flanking regulatory sequence) expressed the *Fugu*

oxytocin transcript specifically in oxytocin neurons in rat (Venkatesh, Si-Hoe, Murphy, &

Brenner, 1997). Moreover, a similar transgenic experiment transferring approximately 5

kb of the *Fugu* oxytocin already resulted in a faithful expression of the fish genes in

mouse oxytocin neurons (Gilligan, Brenner, & Venkatesh, 2003). These studies suggest

there is a remarkably conserved transcriptional regulatory machinery across species, even

between the *Fugu* and rodent oxytocin genes, despite being separated by 400 million

years of evolution.

The GENSAT (Gene Expression Nervous System Atlas) Project demonstrated that most

BAC transgenes reliably recapitulate endogenous expression across independent lines,

provided that extensive flanking intergenic regions are included (Gerfen et al., 2013; S.

Gong et al., 2007; Shiaoching Gong et al., 2003; Heintz, 2004; Schmidt et al., 2013). Our

pv*Oxtr* clone contained the entire locus with full upstream and downstream intergenic

sequences, yet the transgenic lines we generated displayed strikingly divergent brain

expression patterns. These differences were not explained by transgene copy number, but

instead reflected strong insertion-site effects. This paradox contrasts with the general

reproducibility of BAC transgenics and echoes the observation that virtually every rodent

or primate species has a unique distribution of OXTR binding in the brain.

The 3D genomic structure analysis showed that the *Oxtr* loci sit within large TAD

structures, increasing its exposure to distal regulatory elements through an expanded

interaction landscape, whereas *Oxt* lacks organized TAD structures. This difference

might explain why short promoter sequences suffice for conserved *Oxt* expression across

species, while even a large BAC is insufficient for precise, reproducible transcription

regulation of *Oxtr*. The large TAD and extensive long-range interactions render *Oxtr*

especially sensitive to chromosomal context, accounting for the position effects we

observed.

It is also noteworthy that pv*Oxtr* expression in mammary gland (which is responsible for

essential lactation in mammals) was conserved across all Koi lines, indicating that the

essential cis-regulatory elements for peripheral expression are captured within the BAC

and are resistant to position effects. This conclusion is consistent with the chromatin

architecture we observed: the chromatin landscape differs considerably between brain-

related tissues and peripheral tissues. There exists an obvious increase of long-distance

interactions between neighboring TADs, forming a larger TAD structure in brain-related

tissues but not peripheral tissues. Moreover, critical peripheral regulatory sequences may

443 be clustered proximally around the locus, so that they fall entirely within the BAC. By

444 comparison, in brain tissues, regulatory elements are dispersed across a much broader

445 genomic interval, extending beyond the BAC. Together, the larger interaction landscape

446 and distributed regulatory architecture in brain tissues likely explain why BAC constructs

447 are sufficient for peripheral but not brain expression of *Oxtr*.

448 This is well aligned with broader principles of genome organization: subtle changes

449 within 3D chromatin structure may result in substantial changes in local gene expression

450 (Xiao, Hafner, & Boettiger, 2021). TADs also display increased SNPs and structure

451 variations (SVs) density and higher recombination rate compared to inter-TAD

452 regions(Beagan & Phillips-Cremins, 2020; Dekker et al., 2002; Dixon et al., 2012;

453 Rajderkar et al., 2023). In fact, SNPs in the *Oxtr* intron explains 74% of the variance in

454 striatal *Oxtr* expression and social attachment in prairie voles (King et al., 2016) and it is

455 possible that these SNPs near *Oxtr* are linked to variations of long-range chromosomal

456 interaction. Collectively, these evidences support the notion that brain *Oxtr* expression is

457 hyper-sensitive to genetic variation.

458 Such an organization may endow *Oxtr* with enhanced evolvability. Oxytocin is a secreted

459 neuromodulator that can diffuse along axons and through the extracellular space(Leng &

460 Ludwig, 2008; Mitre et al., 2016; Parmaksiz & Kim, 2025). Its functional effects depend

critically on the precise localization of its receptors, as the spatial distribution of oxytocin

receptors determines the neural circuits engaged and the behavioral outcomes elicited. By

making brain expression more reliant on distal regulatory sequences, *Oxtr* becomes

subject to a wider range of cis- and trans-acting influences, rendering its expression in the

brain inherently more variable across individuals and species. This variability allows

oxytocin signaling to recruit novel neural circuits and modulate diverse aspects of social

behavior, thereby contributing to the evolutionary diversification of sociality. Future

studies should aim to resolve the precise mechanisms underlying this variability—for

example, by identifying which distal regulatory elements physically interact with *Oxtr*

within brain-specific chromatin architectures.

Ideas and Speculation

In summary, our results revealed that region-specific expression of *Oxtr* in the brain

requires long-distance interactions between proximal and distal cis-regulatory elements.

In contrast, many other genes remained consistent across BAC-transgenic lines,

indicating that proximal regulatory sequences within the BAC were sufficient to drive

faithful expression in these tissues, and that this expression was resistant to positional

effects.

Our findings support a broader hypothesis: genes vary in their regulatory robustness and

evolutionary flexibility. We propose a framework dividing genes into two broad classes

based on their transcriptional architecture—rigid and flexible genes. Rigid genes—

perhaps comprising the majority of the genome—maintain core life-sustaining functions

(e.g., metabolism, motor function, development, reproduction) and are typically regulated

by compact proximal elements. Their expression is stable across species, and across

individuals within a same species. For such genes, BAC constructs often recapitulate

endogenous expression reliably—examples include CamKII, Protein Kinase C

Delta(Pkcd), and other genes characterized in large-scale projects like GENSAT.

In contrast, a subset of genes are flexible genes—such as neuromodulatory receptors like

Oxtr and potentially *Avpr1a* and *Drd4 (Dopamine receptor D4)*— possess inherently

more fragile or flexible transcriptional regulation. Their expression is highly sensitive to

multiple factors including the chromatin landscape and complex long-range regulatory

interactions. We hypothesize that this regulatory sensitivity is not a flaw but an

evolutionary feature—what we term transcriptional evolvability. Subtle genetic or

epigenetic changes, such as SNPs, structural variants, or chromosomal rearrangements,

may lead to variation in neuromodulator receptor expression, thereby altering the

modulation of preexisting neural circuits. Such changes could unmask cryptic behavioral

phenotypes or give rise to novel social traits without requiring changes to core neural

architecture. For example, caregiving circuits might evolve into pair bonding, empathy, or

romantic attachment, while defensive or aggressive circuits might become

recontextualized into behaviors like jealousy or mate guarding.

This model of transcriptional evolvability helps explain why *Oxtr* and *Avpr1a* expression

varies so markedly across and within species, while their ligands (*Oxt* and *Avp*) remain

stably expressed. The functional consequence of receptor variability is amplified by the

secreted and diffusive nature of these neuromodulators: even small shifts in receptor

localization can profoundly alter which neural circuits are engaged and which behaviors

are modulated. It also raises the possibility that the capacity for behavioral innovation is,

at least in part, encoded in the regulatory fragility of a small set of genes. Our model

carries profound implications. It suggests that the evolution of complex social behavior is

driven not only by neural circuitry or protein function, but by the regulatory architecture

of a small set of flexible genes. These genes serve as behavioral tuning knobs—sensitive

to the chromosomal environment, and capable of generating diverse phenotypes in

response to evolutionary or environmental pressures. This model also offers insight into

variability in human social phenotypes, including those observed in psychiatric

conditions such as autism or social anxiety disorders, where subtle changes in receptor

gene regulation may lead to major phenotypic shifts. Further research into the regulatory

logic of *Oxtr, Avpr1a*, and similar genes may reveal a generalizable mechanism by which

genome structure enables behavioral diversity and adaptation across the animal kingdom.

Materials and Methods

pvOxtr-P2A-Cre **BAC vector construction**.

The vole BAC clone (GeneBank: DP001214.1) containing the *Oxtr* locus was obtained

from a prairie vole BAC library (CHORI-232) (Lisa A. McGraw et al., 2010; L. A.

McGraw, Davis, Thomas, Young, & Thomas, 2012). The NLS (nuclear localization

signal)-Cre-Frt-Amp-Frt cassette was obtained from Dr. Takuji Iwasato, and was slightly

modified to introduce a P2A sequence (Kim et al., 2011). The P2A sequence was

introduced using PCR with the following primers:

 fw5'- ATGGGAAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAGGCTGGAGA

CGTGGAGGAGAACCCTGGACCTCTCGAAACTGACAGGAGAACCACC-3';

 rv5'-AGACTGGAGTCCGCATAGCCCCCCTCCCCCGCCCCAGGCGCGGTGGGCC

AGGCAGGTGGCTCACCTTGACCAAGTTGCTGAAGTTCCTATTCC-3'. The P2A sequence is

underlined. And the resulting amplified cassette had the P2A upstream of the NLS-Cre.

Then the P2A-NLS-Cre-Frt-Amp-Frt cassette was inserted in-frame right before the stop

codon of Oxtr in the BAC clone using the Red/ET Recombineering kit (Gene Bridges

GmbH, Heidelberg Germany) after adding homologous arms by PCR with the following

primers: fw5'-CACCTTCGTCCTGAGTCGCCGCAGCTCCAGCCAGAGGAGCTG

CTCTCAACCATCTTCAGCAATGGGAAGCGGAGCTACTAAC-3';

rv5'-GAGGAGAGGGATACACACCAATAGGCACCTTATACAACTCCACGCAC

GGCCACCAGGGGCAGACTGGAGTCCGCATAGCC-3'. The ampicillin selection marker was

deleted with the 706-Flpe-induced recombination method (Gene Bridges Heidelberg

Germany). A correctly modified BAC clone was verified by using PCR at both the 5' and

3' junctions of the targeted insertion with the following primer pair: fr5'-

GCCTTCATCATCGCCATGCTCTT-3'(p2) and rv5'-GATGGCTGAGTG ACTGGCATCT-3'(p8);

The construct was further verified by sequencing using the following 7 primers specific

either for BAC vector or the p2A-NLS-Cre fragment:

5'- CCTGCAGCCAACTGGAGCTTC-3'(p1); 5'- CTTCCTTGGGCGCATTGACGTC-3'(p5);

5'-TACCTGTTTTGCCGGGTCAG-3(p4)；5'-GCCTTCATCATCGCCATGCTCTT-3'(p2)；

5'-CTGACCCGGCAAAACAGGTA-3'(p7)； 5'-TCCGGTTATTCAACTTGCACCATGC-3'(p6);

5'-GATGGCTGAGTGACTGGCATCT-3'(p8).

Animals.

pv*Oxtr-P2A-Cre* BAC transgenic mouse lines (Koi lines). pv*Oxtr-P2A-Cre* BAC vector

was digested with NotI for linearization and to remove the pTARBAC vector backbone.

The *Vole-Oxtr-P2A-Cre* BAC linearized fragment was purified using CL-4B sepharose

(Sigma-Aldrich), and injected into pronuclei of C57BL/6J zygotes. Mice carrying the

BAC transgene were identified and confirmed using PCR with 5 pairs of primers: fr5'-

TCGACCAGGTTCGTTCACTC-3'(p3) + p7; p2+ p6; p2 +p7; p4 +p8; p2+p8. 8 lines (Koi-1~Koi-

8) were confirmed with successful transgene inheritance. These lines were maintained on

a congenic C57BL/6J background by backcrossing with wildtype C57BL/6J for

assessments of BAC DNA copy numbers and initial transgene expression analysis.

The **ROSA-26-NLS-LacZ mouse** line (Zhang et al., 2016) was obtained from Itohara Lab

at RIKEN.

The **mOxtr-Ires-Cre knock-in** (Hidema et al., 2016) and **mOxtr$^{-/-}$ mutant** (Takayanagi et

al., 2005) lines were obtained from Dr. Katsuhiko Nishimori at Tohoku University, and

both were backcrossed with C57BL/6J for more than 10 generations. Each Koi line and

mOxtr-Ires-Cre knock-in mice were crossed with *ROSA-26-NLS-LacZ* mice to generate

Cre$^+$, LacZ$^+$ double positive mice for further gene expression analysis. Female Koi mice

were crossed with male m*Oxtr*$^{-/-}$ to produce offspring with the genotype of pv*Oxtr-P2A-*

Cre(Koi): m*Oxtr*$^{+/-}$. These offspring were then bred to produce Koi: m*Oxtr*$^{-/-}$ mice

expressing pv*Oxtr* while lacking endogenous *mOxtr*, and these mice were used for

autoradiography, qRT-PCR and behavioral tests. All mice were generated using

continuously housed breeder pairs and P21 as the standard weaning date. All the animal

procedures were approved by University of Tsukuba Animal Care and Use Committee.

Mice were housed under constant temperature and light condition (12 h light and 12 h

dark cycle), and received food and water ad libitum.

Determination of transgene copy number

Custom Taqman® MGB probes were synthesized by oligoJp (Thermo Fisher, Japan) for

detecting the transgenic *Cre* and the mouse *Jun* gene (internal control). The following

primer pairs and probes were used: for the *Cre* assay, fr5'-

ATGACTGGGCACAACAGACAAT-3'; rv5'- CGCTGACAGCCGGAACAC-3'; probe:

5'-FAM- AAACATGCTTCATCGTCGGTC CGG-MGB-3'; for *Jun* assay, fr5'-

GAGTGCTAGCGGAGTCTTAACC-3'; rv5'-CTCCAGACGGCAGTGCTT-3'; probe:

5'-VIC-CTGAGCCCTCCTCCCC-MGB-3'. Real-time PCR was performed using an

Applied Biosystem7500. Transgene copy number was determined through absolute

quantification (standard curve method) following the protocol described in (Chandler et

al., 2007). Briefly, 2 microliters (20 ng) of genomic DNA samples or copy number

standards were analyzed in a 20μL reaction volume with two primer-probe sets (*Cre,*

Jun). The data from transgenic samples were then compared to a standard curve of

calibrator samples that are generated by diluting purified BAC DNA (linearized) over a

range of known concentrations into wild-type mouse genomic DNA. In addition, no-

template controls were included in each experiment. All reactions were performed in

triplicate.

Histology

Adult mice (3-4month old), 4 male and 4 female from each line/ generation were

analyzed for 3 generations. Mice were perfused intracardially with 4% formalin in 0.1M

sodium phosphate buffer. Brains were embedded in 2% agarose in 0.1M PB and cut into

100-m-thick sections with a Micro-slicer (Dosaka, Kyoto, Japan). Mammary glands were

isolated from virgin female mice (2month old), 2 mice/each line were analyzed. The brain

slices and whole mount mammary glands preparation were stained in X-gal solution (5

mM K3FeCN6, 5mMK4FeCN6,2mMMgCl2, 0.02% NP-40, 0.01% Na-deoxycholate, 1

mg/ml X-gal in 0.1M PB) at 37°C for 6 hours and then were stained with hematoxylin.

All experiments were done with positive and negative controls (Cre$^+$, LacZ$^-$) to monitor

the reliability of the X-gal staining. Images were caught by Nanozoomer 2.0-HT slide

scanner (Hamamatsu Photonics).

Receptor Autoradiography

OXTR autoradiography was performed as previously described (Inoue et al., 2022).

Briefly, freshly frozen brains were stored at -80°C. Coronal sections were cut in a cryostat

and 20 μm sections were collected and then stored at -80°C until use in autoradiography.

Brain sections were removed from -80°C storage and air dried, and then fixed for two

minutes with 0.1% paraformaldehyde in PBS at room temperature, and rinsed twice in 50

mM Tris buffer, pH 7.4, to remove endogenous OT. They were then incubated in 50 pM

125I-OVTA (2200 Ci/mmol; PerkinElmer; Boston, MA), a selective, radiolabeled OXTR

ligand, for one hour. Unbound 125I-OVTA was then washed away with Tris-MgCl$_2$

buffer (50 mM Tris plus 2% MgCl2, pH 7.4) and sections were air dried. Sections were

exposed to BioMax MR film (Kodak; Rochester, New York) for five days. Digital images

were obtained with a light box and a Cannon camera (Cannon 6D MarkII, Japan). The

brightness and contrast of representative images were equally adjusted for all

autoradiography images within a panel using Adobe Photoshop.

Fluorescent RNAscope *in situ* hybridization for *Oxtr* mRNA localization

RNAscope Multiplex Fluorescent Reagent Kit v2 (Advanced Cell Diagnostics, Newark,

California; Cat. No. 323270) was used according to the manufacturer's instructions.

Adjacent sections to those utilized for I-125 OVTA receptor binding autoradiography

were used. Briefly, sections were thawed and fixed in 10% neutral buffered formalin for

15 minutes followed by treatment with hydrogen peroxide for 10 minutes and RNAscope

Protease IV for 30 minutes. Probes specific to prairie vole *Oxtr* mRNA (Cat. No. 500721)

were then applied and allowed to hybridize for two hours at 40 °C. After three

amplification steps, hybridized probes were labelled with horseradish peroxidase (HRP),

and TSA Vivid Dye 570 fluorophore. Sections were counterstained with DAPI for nuclei

labeling and coverslipped with ProLong Gold Antifade Mountant (ThermoFisher

Scientific; Waltham, Massachusetts). Negative controls were performed using a bacterial

dapB probe (Cat. No. 310043), which did not exhibit any nonspecific labeling.

Fluorescent RNAscope images were acquired using a Keyence BZ-X710 microscope.

Whole sections were captured with 10x objective lens in two channels (red and blue), and

images were stitched together using Keyence BZX Analyzer Software. Brightness and

contrast of the images were adjusted using Photoshop (Adobe; San Jose, California).

Quantification of *Oxtr* mRNA expression by qRT-PCR

Real-time quantitative polymerase chain reaction (RT-qPCR) using an Applied

Biosystems 7500 Real-Time PCR system was used to quantify *Oxtr* mRNA expression

levels in different brain regions of Koi lines. Total RNA of different brain regions was

isolated using the RNeasy extraction kit (Qiagen, Germantown, US), and cDNA was

synthesized after Deoxyribonuclease I (Invitrogen, Waltham, US) treatment by using

EvoScript Universal cDNA Master (Roche, Penzberg, Germany), and then quantitative

RT-PCR was performed with FastStart Universal SYBR Green Master (Roche, Penzberg,

Germany). The relative standard curve method was used to obtain the relative quantities

of *Oxtr* expression following the manual of Applied Biosystems. To allow the

comparation between Koi lines and WT mice, the following primer pair (which can detect

the expression of both *mOxtr* and pv*Oxtr*) was used: fr5'-

GCCTTTCTTCTTCGTGCAGATG-3' and rv5′-ATGTAGATCCAGGGGTTGCAG-3′ In

addition, to specifically detect the expression of pv*Oxtr*, the primer pair fr5′-

GCCTTTCTTCTTCGTGCAGATG-3 and rv5'-AAAGAGGTGGCCCGTGAAC-3′ was

used in the 2nd qRT-PCR experiment. GAPDH was used as the endogenous control for

both experiments and was amplified by the following primer pair: fr5'-

GGGTTCCTATAAATACGGACTGC-3' and rv5'- CCATTTGTCTACGGGACGA-3'.

Samples were analyzed in triplicates. A non-template control was performed to ensure

that there was no amplification of genomic DNA. All of the experimenters were blind to

the genotype of the subjects.

Partner preference test (PPT)

Subjects were housed with 4 age-matched and same-sex littermates until testing at

adulthood (2-5 months old). 12-14 homozygous? mice were tested for each group.

Ovariectomized females were paired and cohoused with a sexually experienced adult

wildtype male for 21 days before PPT test. The female was injected with estradiol

benzoate (10 μg and 5 μg at 48 h and 24 h before induced mating) and progesterone (500

μg at 4–7 h before induced mating) to ensure high sexual receptivity before mating.

Mating was induced on Day 2,9 and16. PPT data was analyzed using TimeSSI1 for social

interaction test system (O' Hara & Co., Ltd.). Briefly, the experimental subject was

placed in a chamber, in which two pen-boxes (diameter 8cm, height10cm) were located at

two diagonally opposite corners. The experimental animal was free to move throughout

the chamber and the time spent in close proximity to each pen-box (stay time) is recorded

using an automated mouse tracking system (Fig.4 A, B). In the habituation phase, the

experimental subject was allowed to explore the chamber for 5 minutes with both pen-

boxes empty. In the recognition phase, the partner stimulus animal was restricted in one

pen-box, and a novel "stranger" stimulus animal was restricted in another pen-box, and

the experimental subject was allowed to explore the chamber for 10 minutes. In the

preference phase, the positions of the partner stimulus animal and the "stranger" stimulus

animal were switched, and the experimental subject was allowed to explore the chamber

for 20 minutes. To assure unbiased design, pen-box assignments were counterbalanced

for the diagonal positions. All of the experimenters were blind to the genotype of the

subjects.

Parental behavior test

To examine maternal behavior, we performed pup retrieval test with virgin female mice

(2- to 4-months old, 15-18mice/line), following the method described in (Stolzenberg,

Stevens, & Rissman, 2012) with some modifications. Briefly, 5 days prior to the

behavioral testing, each subject was isolated and housed in separate home cages in the

test room. Nesting material (about 1.5 g of cotton wool) was provided for nest making.

The tests were conducted in the dark phase of light/dark cycle (12 hours/12 hours). On

the day of the test, each subject was placed in the recording area and allowed 15 minutes

for habituation. The stimulus pups (age: P2-P3) were collected from a group of donor

mothers immediately before the start of the experiment. A retrieval test began with the

placement of 3 stimulus pups on the side farthest form the nest, and the female's behavior

was recorded by an ARNAN 4 channel security system for 15 minutes. Pup retrieval was

defined as picking up the pup and bringing it to the nest. If the subject did not complete

the retrieval for 3 pups in 15 minutes, the video recording time was extended to 30

minutes. The latency to retrieve each pup (1st, 2nd, 3rd) and the time spent crouching on

the pups were time-stamped and calculated manually. A latency of 1800 seconds was

assigned if the pup-retrieval was not completed in 30 minutes. Crouching was defined as

the mouse supporting itself in a lactation-position over the pups in the nest. Crouching

time on a single pup, two pups, and three pups were summed up as total crouching time.

All of the experimenters were blind to the genotype of the subjects.

3D chromatin verification on Hi-C datasets of humans and mice

The 3D genome structure tracks were obtained from 4DNucleome Consortium

(https://www.4dnucleome.org/) where datasets were systematically reanalyzed following

recommended standard protocols. The datasets IDs and sample information are provided

in the supplementary material (Supplementary Table1,2). 3D Genome Browser(Wang et

al., 2018) and WashU Epigenome Browser(Li et al., 2022) were employed for

visualizations.

Experimental design and statistical analysis

The sample sizes were determined by G*Power 3 program (Faul, Erdfelder, Lang, &

Buchner, 2007) based on pilot experiments. They are also within the range that is

generally accepted in the field. All experiments were performed and analyzed blind to the

genotypes and no data point was excluded. All reported sample numbers (n) represent

independent biological replicates that are the numbers of tested mice or tissue. All the

statistical analyses were performed by SPSS21(IBM). Mixed ANOVA was used for

analyzing PPT data and pup retrieval data. Mauchly's test of sphericity was used to test

whether or not the assumption of sphericity was met in repeated measures. Greenhouse-

Geisser correction was applied when the sphericity was violated. Two-way ANOVA was

used for analyzing qRT-PCR region specific data across different mouse lines. One-way

ANOVA was used for analyzing crouching time in the parental behavioral test. If there

was a significant main effect of an independent factor, post hoc test was used to do

multiple comparisons. If there was a significant interaction between within-subject factor

and between-subject factor, post hoc pairwise comparisons were applied to analyze the

simple main effects. Bonferroni correction was used for all the post hoc tests.

747   **Acknowledgements**

748   This work was supported by the RIKEN Incentive Research Project

749   (100226201701100443) to Q.Z., the Brain Science Project, Center for novel science

750   initiatives, National institutes of natural sciences (BS291003) to Q.Z., the RIKEN Aging

751   Project (10026-201701100263-340120) to Q.Z., the JSPS Kakenhi Grant-in-Aid for

752   Young Scientists (17K18362) to Q.Z., the JSPS Kakenhi Grant-in-Aid for Challenging

753   Research (19K21807) to Q.Z., the JSPS Kakenhi Grant-in-Aid for Challenging Research

754   (22K19478)to Q.Z., the JSPS Kakenhi Grant-in-Aid for Scientific Research (25K10198)

755   to Q.Z., the Takeda Medical Research Grant to Q.Z., the International Education and

756   Research Laboratory Program of University of Tsukuba to L.Y. and Q.Z.. We thank Dr.

757   Katsuhiko Nishimori (Tohoku University and Fukushima Medical University, Japan) for

758   providing *mOxtr*-Ires-Cre knock-in line and *mOxtr*-/- mutant line; Dr. Cary Lai (Indiana

759   University, USA), Dr. Takuji Iwasato(National Institute of Genetics, Japan) and Dr. Seiya

760   Mizuno(University of Tsukuba, Japan) for their advice on BAC vector construction; Dr.

761   Volkhard Maeckel (CERN, Switzerland) for helping with image capturing; Dr. Yong

762   Zhang (Tongji University, China) for the stimulating discussion about 3D chromatin

763   structure, and we thank the Laboratory of Animal Resource Center of University of

764   Tsukuba for pronuclear injection.

Author Contributions:

Q.Z. and L.Y. conceived and designed the study. Q.Z., M.T., L.N., K.I., L.K., M.P., Y.N.,

M.S., and T.S. performed experiments. Q.Z., M.T., L.N., K.I., L.K., M.P., and K.N.

analyzed data. S.I. provided guidance and facilities for transgene vector construction.

Q.Z. and L.Y. supervised the project and wrote the manuscript. All authors contributed to

discussion and approved the final version.

References:

Ahern, T. H., Olsen, S., Tudino, R., & Beery, A. K. (2021). Natural variation in the oxytocin

receptor gene and rearing interact to influence reproductive and nonreproductive social

behavior and receptor binding. *Psychoneuroendocrinology, 128*, 105209.

doi:10.1016/j.psyneuen.2021.105209

Amadei, E. A., Johnson, Z. V., Jun Kwon, Y., Shpiner, A. C., Saravanan, V., Mays, W. D., . . . Liu,

R. C. (2017). Dynamic corticostriatal activity biases social bonding in monogamous

female prairie voles. *Nature, 546*(7657), 297-301. doi:10.1038/nature22381

Barrett, C. E., Arambula, S. E., & Young, L. J. (2015). The oxytocin system promotes resilience to

the effects of neonatal isolation on adult social attachment in female prairie voles. *Transl*

Psychiatry, 5(7), e606. doi:10.1038/tp.2015.73

Beagan, J. A., & Phillips-Cremins, J. E. (2020). On the existence and functionality of topologically

associating domains. *Nature Genetics, 52*(1), 8-16. doi:10.1038/s41588-019-0561-1

Beil, J., Fairbairn, L., Pelczar, P., & Buch, T. (2012). Is BAC transgenesis obsolete? State of the

art in the era of designer nucleases. *J Biomed Biotechnol, 2012*, 308414.

doi:10.1155/2012/308414

Bian, Q., & Belmont, A. S. (2010). BAC TG-EMBED: one-step method for high-level, copy-

number-dependent, position-independent transgene expression. *Nucleic Acids Research,*

38(11), e127-e127. doi:10.1093/nar/gkq178

Carcea, I., Caraballo, N. L., Marlin, B. J., Ooyama, R., Riceberg, J. S., Mendoza Navarro, J. M., . . .

Froemke, R. C. (2021). Oxytocin neurons enable social transmission of maternal

behaviour. *Nature, 596*(7873), 553-557. doi:10.1038/s41586-021-03814-7

Chandler, K. J., Chandler, R. L., Broeckelmann, E. M., Hou, Y., Southard-Smith, E. M., &

Mortlock, D. P. (2007). Relevance of BAC transgene copy number in mice: transgene

copy number variation across multiple transgenic lines and correlations with transgene

integrity and expression. *Mamm Genome, 18*(10), 693-708. doi:10.1007/s00335-007-

9056-y

Chen, Z., Han, Y., Ma, Z., Wang, X., Xu, S., Tang, Y., . . . Zhan, Y. (2024). A prefrontal-thalamic

circuit encodes social information for social recognition. *Nature Communications, 15*(1),

1036. doi:10.1038/s41467-024-45376-y

Crook, Z. R., & Housman, D. (2011). Huntington's disease: can mice lead the way to treatment?

Neuron, 69(3), 423-435. doi:10.1016/j.neuron.2010.12.035

Dekker, J., Rippe, K., Dekker, M., & Kleckner, N. (2002). Capturing chromosome conformation.

Science, 295(5558), 1306-1311. doi:10.1126/science.1067799

Dixon, J. R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., . . . Ren, B. (2012). Topological

domains in mammalian genomes identified by analysis of chromatin interactions. *Nature,*

485(7398), 376-380. doi:10.1038/nature11082

Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G*Power 3: a flexible statistical power

analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods,*

39(2), 175-191. doi:10.3758/bf03193146

Ferrante, R. J. (2009). Mouse models of Huntington's disease and methodological considerations

for therapeutic trials. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease,*

1792(6), 506-520. doi:https://doi.org/10.1016/j.bbadis.2009.04.001

Froemke, R. C., & Young, L. J. (2021). Oxytocin, Neural Plasticity, and Social Behavior. *Annu Rev*

Neurosci, 44, 359-381. doi:10.1146/annurev-neuro-102320-102847

Gerfen, C. R., Paletzki, R., & Heintz, N. (2013). GENSAT BAC cre-recombinase driver lines to

study the functional organization of cerebral cortical and basal ganglia circuits. *Neuron,*

80(6), 1368-1383. doi:10.1016/j.neuron.2013.10.016

Gibcus, Johan H., & Dekker, J. (2013). The Hierarchy of the 3D Genome. *Molecular Cell, 49*(5),

773-782. doi:https://doi.org/10.1016/j.molcel.2013.02.011

Gilligan, P., Brenner, S., & Venkatesh, B. (2003). Neurone-specific expression and regulation of

the pufferfish isotocin and vasotocin genes in transgenic mice. *J Neuroendocrinol, 15*(11),

1027-1036. doi:10.1046/j.1365-2826.2003.01090.x

Gong, S., Doughty, M., Harbaugh, C. R., Cummins, A., Hatten, M. E., Heintz, N., & Gerfen, C. R.

(2007). Targeting Cre recombinase to specific neuron populations with bacterial artificial

chromosome constructs. *J Neurosci, 27*(37), 9817-9823. doi:10.1523/jneurosci.2707-

07.2007

Gong, S., Zheng, C., Doughty, M. L., Losos, K., Didkovsky, N., Schambra, U. B., . . . Heintz, N.

(2003). A gene expression atlas of the central nervous system based on bacterial artificial

chromosomes. *Nature, 425*(6961), 917-925. doi:10.1038/nature02033

Grinevich, V., Knobloch-Bollmann, H. S., Eliava, M., Busnelli, M., & Chini, B. (2016). Assembling

the Puzzle: Pathways of Oxytocin Signaling in the Brain. *Biol Psychiatry, 79*(3), 155-164.

doi:10.1016/j.biopsych.2015.04.013

Heintz, N. (2004). Gene Expression Nervous System Atlas (GENSAT). *Nature Neuroscience,*

7(5), 483-483. doi:10.1038/nn0504-483

Hidema, S., Fukuda, T., Hiraoka, Y., Mizukami, H., Hayashi, R., Otsuka, A., . . . Nishimori, K.

(2016). Generation of Oxtr cDNA(HA)-Ires-Cre Mice for Gene Expression in an

Oxytocin Receptor Specific Manner. *J Cell Biochem, 117*(5), 1099-1111.

doi:10.1002/jcb.25393

Inoue, K., Ford, C. L., Horie, K., & Young, L. J. (2022). Oxytocin receptors are widely distributed

in the prairie vole (Microtus ochrogaster) brain: Relation to social behavior, genetic

polymorphisms, and the dopamine system. *Journal of Comparative Neurology, 530*(16),

2881-2900. doi:https://doi.org/10.1002/cne.25382

Jurek, B., & Neumann, I. D. (2018). The Oxytocin Receptor: From Intracellular Signaling to

Behavior. *Physiological reviews, 98*(3), 1805-1908. doi:10.1152/physrev.00031.2017

Kandasamy, L. C., Tsukamoto, M., Banov, V., Tsetsegee, S., Nagasawa, Y., Kato, M., . . . Zhang,

Q. (2021). Limb-clasping, cognitive deficit and increased vulnerability to kainic acid-

induced seizures in neuronal glycosylphosphatidylinositol deficiency mouse models.

Human Molecular Genetics, 30(9), 758-770. doi:10.1093/hmg/ddab052

Kappel, J. M., Förster, D., Slangewal, K., Shainer, I., Svara, F., Donovan, J. C., . . . Larsch, J.

(2022). Visual recognition of social signals by a tectothalamic neural circuit. *Nature,*

608(7921), 146-152. doi:10.1038/s41586-022-04925-5

Kim, J. H., Lee, S. R., Li, L. H., Park, H. J., Park, J. H., Lee, K. Y., . . . Choi, S. Y. (2011). High

cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in human cell lines,

zebrafish and mice. *PloS one, 6*(4), e18556. doi:10.1371/journal.pone.0018556

King, L. B., Walum, H., Inoue, K., Eyrich, N. W., & Young, L. J. (2016). Variation in the Oxytocin

Receptor Gene Predicts Brain Region-Specific Expression and Social Attachment. *Biol*

Psychiatry, 80(2), 160-169. doi:10.1016/j.biopsych.2015.12.008

Leng, G., & Ludwig, M. (2008). Neurotransmitters and peptides: whispered secrets and public

announcements. *J Physiol, 586*(23), 5625-5632. doi:10.1113/jphysiol.2008.159103

Li, D., Purushotham, D., Harrison, J. K., Hsu, S., Zhuo, X., Fan, C., . . . Wang, T. (2022). WashU

Epigenome Browser update 2022. *Nucleic Acids Research, 50*(W1), W774-W781.

doi:10.1093/nar/gkac238

Lieberman-Aiden, E., van Berkum, N. L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., . . .

Dekker, J. (2009). Comprehensive Mapping of Long-Range Interactions Reveals Folding

Principles of the Human Genome. *Science, 326*(5950), 289-293.

doi:doi:10.1126/science.1181369

Mack, K. L., Campbell, P., & Nachman, M. W. (2016). Gene regulation and speciation in house

mice. *Genome Res, 26*(4), 451-461. doi:10.1101/gr.195743.115

Marlin, B. J., Mitre, M., D'Amour J, A., Chao, M. V., & Froemke, R. C. (2015). Oxytocin enables

maternal behaviour by balancing cortical inhibition. *Nature, 520*(7548), 499-504.

doi:10.1038/nature14402

McGraw, L. A., Davis, J. K., Lowman, J. J., ten Hallers, B. F. H., Koriabine, M., Young, L. J., . . .

Thomas, J. W. (2010). Development of genomic resources for the prairie vole (Microtus

ochrogaster): construction of a BAC library and vole-mouse comparative cytogenetic map.

BMC Genomics, 11(1), 70. doi:10.1186/1471-2164-11-70

McGraw, L. A., Davis, J. K., Thomas, P. J., Young, L. J., & Thomas, J. W. (2012). BAC-based

sequencing of behaviorally-relevant genes in the prairie vole. *PloS one, 7*(1), e29345.

doi:10.1371/journal.pone.0029345

Mitre, M., Marlin, B. J., Schiavo, J. K., Morina, E., Norden, S. E., Hackett, T. A., . . . Froemke, R.

C. (2016). A Distributed Network for Social Cognition Enriched for Oxytocin Receptors. *J*

Neurosci, 36(8), 2517-2535. doi:10.1523/jneurosci.2409-15.2016

Newmaster, K. T., Nolan, Z. T., Chon, U., Vanselow, D. J., Weit, A. R., Tabbaa, M., . . . Kim, Y.

(2020). Quantitative cellular-resolution map of the oxytocin receptor in postnatally

developing mouse brains. *Nature Communications, 11*(1), 1885. doi:10.1038/s41467-

020-15659-1

Olazábal, D. E., & Young, L. J. (2006). Oxytocin receptors in the nucleus accumbens facilitate

"spontaneous" maternal behavior in adult female prairie voles. *Neuroscience, 141*(2), 559-

568. doi:10.1016/j.neuroscience.2006.04.017

Osada, N., Miyagi, R., & Takahashi, A. (2017). Cis- and Trans-regulatory Effects on Gene

Expression in a Natural Population of Drosophila melanogaster. *Genetics, 206*(4), 2139-

2148. doi:10.1534/genetics.117.201459

Parmaksiz, D., & Kim, Y. (2025). Navigating Central Oxytocin Transport: Known Realms and

Uncharted Territories. *Neuroscientist, 31*(3), 234-261. doi:10.1177/10738584241268754

Phelps, S. M., & Young, L. J. (2003). Extraordinary diversity in vasopressin (V1a) receptor

distributions among wild prairie voles (Microtus ochrogaster): patterns of variation and

covariation. *J Comp Neurol, 466*(4), 564-576. doi:10.1002/cne.10902

Rajderkar, S., Barozzi, I., Zhu, Y., Hu, R., Zhang, Y., Li, B., . . . Pennacchio, L. A. (2023).

Topologically associating domain boundaries are required for normal genome function.

Communications Biology, 6(1), 435. doi:10.1038/s42003-023-04819-w

Rangel, M. J., Baldo, M. V. C., & Canteras, N. S. (2018). Influence of the anteromedial thalamus

on social defeat-associated contextual fear memory. *Behavioural Brain Research, 339*,

269-277. doi:https://doi.org/10.1016/j.bbr.2017.10.038

Rigney, N., de Vries, G. J., Petrulis, A., & Young, L. J. (2022). Oxytocin, Vasopressin, and Social

Behavior: From Neural Circuits to Clinical Opportunities. *Endocrinology, 163*(9).

doi:10.1210/endocr/bqac111

Rogers Flattery, C. N., Coppeto, D. J., Inoue, K., Rilling, J. K., Preuss, T. M., & Young, L. J.

(2022). Distribution of brain oxytocin and vasopressin V1a receptors in chimpanzees (Pan

troglodytes): comparison with humans and other primate species. *Brain Struct Funct,*

227(5), 1907-1919. doi:10.1007/s00429-021-02369-7

Ross, H. E., Freeman, S. M., Spiegel, L. L., Ren, X., Terwilliger, E. F., & Young, L. J. (2009).

Variation in oxytocin receptor density in the nucleus accumbens has differential effects on

affiliative behaviors in monogamous and polygamous voles. *J Neurosci, 29*(5), 1312-1318.

doi:10.1523/jneurosci.5039-08.2009

Ross, H. E., & Young, L. J. (2009). Oxytocin and the neural mechanisms regulating social

cognition and affiliative behavior. *Front Neuroendocrinol, 30*(4), 534-547.

doi:10.1016/j.yfrne.2009.05.004

Schmidt, E. F., Kus, L., Gong, S., & Heintz, N. (2013). BAC transgenic mice and the GENSAT

database of engineered mouse strains. *Cold Spring Harb Protoc, 2013*(3).

doi:10.1101/pdb.top073692

Shenoy, S. A., Zheng, S., Liu, W., Dai, Y., Liu, Y., Hou, Z., . . . Li, C. (2022). A novel and accurate

full-length HTT mouse model for Huntington's disease. *eLife, 11*, e70217.

doi:10.7554/eLife.70217

Shi, X., Ng, D. W. K., Zhang, C., Comai, L., Ye, W., & Jeffrey Chen, Z. (2012). Cis- and trans-

regulatory divergence between progenitor species determines gene-expression novelty in

Arabidopsis allopolyploids. *Nature Communications, 3*(1), 950. doi:10.1038/ncomms1954

Signor, S. A., & Nuzhdin, S. V. (2018). The Evolution of Gene Expression in cis and trans. *Trends*

Genet, 34(7), 532-544. doi:10.1016/j.tig.2018.03.007

Skuse, D. H., Lori, A., Cubells, J. F., Lee, I., Conneely, K. N., Puura, K., . . . Young, L. J. (2014).

Common polymorphism in the oxytocin receptor gene (<i>OXTR</i>) is associated with

human social recognition skills. *Proceedings of the National Academy of Sciences, 111*(5),

1987-1992. doi:doi:10.1073/pnas.1302985111

Smirnov, A., Fishman, V., Yunusova, A., Korablev, A., Serova, I., Skryabin, B. V., . . . Battulin, N.

(2019). DNA barcoding reveals that injected transgenes are predominantly processed by

homologous recombination in mouse zygote. *Nucleic Acids Research, 48*(2), 719-735.

doi:10.1093/nar/gkz1085

Stolzenberg, D. S., Stevens, J. S., & Rissman, E. F. (2012). Experience-facilitated improvements in

pup retrieval; evidence for an epigenetic effect. *Horm Behav, 62*(2), 128-135.

doi:10.1016/j.yhbeh.2012.05.012

Suster, M. L., Abe, G., Schouw, A., & Kawakami, K. (2011). Transposon-mediated BAC

transgenesis in zebrafish. *Nature Protocols, 6*(12), 1998-2021.

doi:10.1038/nprot.2011.416

Taguchi, T., Ikuno, M., Hondo, M., Parajuli, L. K., Taguchi, K., Ueda, J., . . . Takahashi, R.

(2019). α-Synuclein BAC transgenic mice exhibit RBD-like behaviour and hyposmia: a

prodromal Parkinson's disease model. *Brain, 143*(1), 249-265. doi:10.1093/brain/awz380

Takayanagi, Y., Yoshida, M., Bielsky, I. F., Ross, H. E., Kawamata, M., Onaka, T., . . . Nishimori,

K. (2005). Pervasive social deficits, but normal parturition, in oxytocin receptor-deficient

mice. *Proceedings of the National Academy of Sciences, 102*(44), 16096-16101.

doi:doi:10.1073/pnas.0505312102

Theofanopoulou, C., Andirkó, A., Boeckx, C., & Jarvis, E. D. (2022). Oxytocin and vasotocin

receptor variation and the evolution of human prosociality. *Comprehensive*

Psychoneuroendocrinology, 11, 100139.

doi:https://doi.org/10.1016/j.cpnec.2022.100139

Venkatesh, B., Si-Hoe, S. L., Murphy, D., & Brenner, S. (1997). Transgenic rats reveal functional

conservation of regulatory controls between the Fugu isotocin and rat oxytocin genes.

Proc Natl Acad Sci U S A, 94(23), 12462-12466. doi:10.1073/pnas.94.23.12462

Walum, H., & Young, L. J. (2018). The neural mechanisms and circuitry of the pair bond. *Nat Rev*

Neurosci, 19(11), 643-654. doi:10.1038/s41583-018-0072-6

Walum, H., & Young, L. J. (2018). The neural mechanisms and circuitry of the pair bond. *Nature*

Reviews Neuroscience, 19(11), 643-654. doi:10.1038/s41583-018-0072-6

Wang, Y., Song, F., Zhang, B., Zhang, L., Xu, J., Kuang, D., . . . Yue, F. (2018). The 3D Genome

Browser: a web-based browser for visualizing 3D genome organization and long-range

chromatin interactions. *Genome Biol, 19*(1), 151. doi:10.1186/s13059-018-1519-9

Wendt, K. S., Yoshida, K., Itoh, T., Bando, M., Koch, B., Schirghuber, E., . . . Peters, J. M. (2008).

Cohesin mediates transcriptional insulation by CCCTC-binding factor. *Nature,*

451(7180), 796-801. doi:10.1038/nature06634

Wolff, M., & Vann, S. D. (2019). The Cognitive Thalamus as a Gateway to Mental

Representations. *J Neurosci, 39*(1), 3-14. doi:10.1523/jneurosci.0479-18.2018

Xiao, J. Y., Hafner, A., & Boettiger, A. N. (2021). How subtle changes in 3D structure can create

large changes in transcription. *eLife, 10*, e64320. doi:10.7554/eLife.64320

Yang, X. W., & Lu, X.-H. (2008). Chapter 19 - The Bac Transgenic Approach to Study

Parkinson's Disease in Mice. In R. Nass & S. Przedborski (Eds.), *Parkinson's Disease* (pp.

247-268). San Diego: Academic Press.

Young, L. J., & Crews, D. (1995). Comparative neuroendocrinology of steroid receptor gene

987   expression and regulation: Relationship to physiology and behavior. *Trends Endocrinol*

988   *Metab, 6*(9-10), 317-323. doi:10.1016/1043-2760(95)00175-1

989   Young, L. J., & Zhang, Q. I. (2021). On the Origins of Diversity in Social Behavior. *The Japanese*

990   *Journal of Animal Psychology, doi: 10.2502/janip.71.1.4*. doi:doi: 10.2502/janip.71.1.4

991   Zhang, Q., Sano, C., Masuda, A., Ando, R., Tanaka, M., & Itohara, S. (2016). Netrin-G1 regulates

992   fear-like and anxiety-like behaviors in dissociable neural circuits. *Scientific Reports, 6*(1),

993   28750. doi:10.1038/srep28750

1000

1001

1002

1003

Figure titles and Legends:

Fig.1 Construction of Koi lines. (A) Example of the remarkable species difference of

brain OXTR distribution. OXTR receptor autoradiography in dorsal caudate putamen

(CP) and nucleus accumbens shell (NAccSh) of rat, mouse, and prairie voleis (Adapted

Froemke and Young(Froemke & Young, 2021)). (B) The pv*Oxtr* BAC clone contains the

entire sequence of pv*Oxtr* gene (colored green), the complete intergenic sequence both

upstream and downstream of pv*Oxtr* gene, the 12^{th} exon of *Rad18* (colored blue) and the

2^{nd} exon of *Cav3* (colored purple). The yellow arrows indicate the direction of

transcription. (C) Schematic diagram of the strategy to generate Koi mice. The P2A-

NLS-Cre-Frt-Amp-Frt cassette was inserted in-frame right before the stop codon of *Oxtr*.

And the ampicillin selection marker was deleted by 706-Flpe. PCR primers for verifying

the targeted alleles are shown as red arrows. (D) Two correctly modified vector clones

were verified by using PCR with p2 and p8. (E) Koi lines were verified by 5 pairs of

primers. From Lane1 to Lane8 are 100bp marker, p3+p7, p2+p6, p2+p7, p4+p8, p2+p8,

blank and 1kb marker.

1019

Fig.2 Transgenic CRE mediated Lac-Z reporter gene expression in the brain and

mammary gland of Koi lines. X-gal staining of the brains and mammary glands are

from the double positive (*Cre+*, *LacZ+*) offspring from Koi-1 (A, F, K, P), Koi-2 (B, G,

L, Q), Koi-3 (G, H, M, R), Koi-4 (D, I, N, S) and *mOxtr-Ires-Cre* knock-in line (E, J, O,

T). The *mOxtr-Ires-Cre* KI line used contains Cre (and an HA tag) inserted in the third

exon, right after the start codon. Note that regardless of the insertion site of Cre in the

mouse genome DNA or pvOxtr-BAC transgene, *Oxtr* expression visualized by Lac-Z

staining in the mammary gland was comparable between the mOxtr-Ires-Cre KI and Koi

lines. This contrasts with the diverse expression patterns of *Oxtr* expression in the brain.

U, V, M, X, Y are the magnified images of the areas within the yellow-border squares

from D, N, E, J, O, respectively, to show the distribution of low-density Lac-Z positive

cells in these areas. Scale bar=1mm (A-O), 500um (P-T) and 200um (U-Y). (Z) The

estimated copy number of transgene of all Koi lines. See Supplementary Figure 2 for

images of Koi-5-8 lines. PFC=prefrontal cortex, LS=lateral septum, BLA=basolateral

amygdala, VMH=ventromedial nucleus of the hypothalamus, Tha=thalamus.

1035

Fig.3 The expression of OXTR in adult brain of Koi lines. Autoradiographs illustrate

the distribution of OXTR binding in adult Koi-1: m*Oxtr*$^{-/-}$ (A, F, K), Koi-2: m*Oxtr*$^{-/-}$ (B,

G, L), Koi-3: m*Oxtr*$^{-/-}$ (C, H, M), Koi-4: m*Oxtr*$^{-/-}$ (D, I, N) and wild type (WT) (E, J, O).

All Koi mice used for this analysis were pv*Oxtr-P2A-Cre*: m*Oxtr*$^{-/-}$ mice in which the

pvOXTR was expressed while endogenous mOXTR was absent. PFC=prefrontal cortex,

LS=lateral septum, BLA=basolateral amgydala, VMH=ventromedial nucleus of the

hypothalamus, Tha=thalamus. Scale bar=1mm.

1043

Fig.4 mRNA expression of *Oxtr* in the adult brain of Koi lines. *Oxtr* mRNA (red) was

labeled with in situ hybridization against DAPI for nuclei labelling (purple). (A) and (C)

show clear expression of *Oxtr* in the PFA and Ctx in Koi-3: m*Oxtr*$^{-/-}$. (B) and(D) show

clear expression of *Oxtr* in the NAc and Tha in Koi-4: m*Oxtr*$^{-/-}$. All Koi mice used for this

analysis were pv*Oxtr-P2A-Cre*: m*Oxtr*$^{-/-}$ mice, in which the pvOXTR was expressed

while endogenous mOXTR was absent. PFC=prefrontal cortex, NAc= Nucleus

accumbens, Ctx=Cortex, Tha=thalamus. Scale bar=400um for A and B, 500um for C and

D.

1052

Fig.5 Quantitative PCR analysis of *Oxtr* mRNA expression in different brain regions

of Koi lines. (A) A primer set recognizing both m*Oxtr* and pv*Oxtr* was used to analyze

Oxtr mRNA in olfactory bulb, prefrontal cortex, ventral striatum and thalamus from WT,

Koi-1: m*Oxtr*$^{-/-}$, Koi-2: m*Oxtr*$^{-/-}$, Koi-3: m*Oxtr*$^{-/-}$, Koi-4: m*Oxtr*$^{-/-}$ and *mOxtr*$^{-/-}$ mutant

mice, n=4 for each genotype. Note that the level of *Oxtr* mRNA in *mOxtr*$^{-/-}$ mutant mice

was nondetectable (ND). The autoradiograph of OXTR binding in olfactory bulb from

each line were shown alongside the legend. There was a significant interaction between

position and genotype $F_{(12,64)} = 26.33$, $p < 0.0001$; Post hoc Bonferroni test revealed that

Oxtr expression level in olfactory bulb of Koi-1: m*Oxtr*$^{-/-}$ was significantly higher than

that of other genotype groups ($p < 0.0001$ for all), and that of Koi-4: m*Oxtr*$^{-/-}$ was

significantly higher than that of Koi-3: m*Oxtr*$^{-/-}$ $p < 0.05$. And the striatum of Koi-4:

m*Oxtr*$^{-/-}$ expressed significantly higher *Oxtr* when compared with WT and Koi-3: m*Oxtr*$^{-/-}$

($p < 0.05$ for both). The *Oxtr* expression level in thalamus of Koi-4: m*Oxtr*$^{-/-}$ was

significantly higher than other genotype groups ($p < 0.0001$ when compared with WT and

Koi-3: m*Oxtr*$^{-/-}$, $p < 0.001$ when compared with Koi-2: m*Oxtr*$^{-/-}$, $p < 0.05$ when compared

with Koi-1: m*Oxtr*$^{-/-}$) (B) A primer set specifically recognizing pv*Oxtr*. was used to

analyze *Oxtr* mRNA in WT, Koi-1: m*Oxtr*$^{-/-}$, Koi-2: m*Oxtr*$^{-/-}$, Koi-3: m*Oxtr*$^{-/-}$ and Koi-4:

m*Oxtr*$^{-/-}$ mic, n=4 for each genotype. Note that the level of *Oxtr* mRNA in WT mice was

barely detectable. There was a significant interaction between position and genotype F

$_{(9,48)} = 89.25$, $p < 0.0001$; Post Hoc Bonferroni test revealed that *Oxtr* expression level in

the olfactory bulb of Koi-1: m*Oxtr*$^{-/-}$ was significantly higher than that of other genotype

groups ($p < 0.0001$ for all), and that of Koi-4: m*Oxtr*$^{-/-}$ was significantly higher than that of

Koi-2: m*Oxtr*$^{-/-}$ and Koi-3: m*Oxtr*$^{-/-}$ ($p < 0.0001$ for both). The *Oxtr* expression level in

prefrontal cortex of Koi-3: m$Oxtr^{-/-}$ was significantly higher than that of other groups

(p<0.01 when comparing with Koi-1: m$Oxtr^{-/-}$ and Koi-4: m$Oxtr^{-/-}$; p<0.05 when

comparing with Koi-2: m$Oxtr^{-/-}$). And the striatum of Koi-4: m$Oxtr^{-/-}$ expressed

significantly higher $Oxtr$ when comparing with Koi-2: m$Oxtr^{-/-}$ and Koi-3: m$Oxtr^{-/-}$

(p<0.01 for both). The $Oxtr$ expression level in thalamus of Koi-4: m$Oxtr^{-/-}$ was

significantly higher than that of other genotype groups (p<0.0001 for all). All Koi mice

used for this analysis were pv$Oxtr$-P2A-Cre: m$Oxtr^{-/-}$ mice, in which the pvOXTR was

expressed while endogenous mOXTR was absent. The significance of interaction

between genotype and brain region: # # # # p<0.0001. The significance of single main

effect of genotype: ＊p<0.05; ＊＊p<0.01; ＊＊＊p<0.001; ＊＊＊＊p<0.0001.

1086

Fig.6 Social behaviors of Koi lines. (A) Illustration of the partner preference test (PPT)

paradigm. The experimental mouse was placed in a chamber with an automatic

monitoring system, where it could freely explore two pen-boxes localized in two diagonal

corners. Habituation phase: two pen-boxes are empty. Recognition phase: a stimulating

partner mouse is placed into one pen-box and a stimulating stranger mouse is placed in

the other one. Preference phase: The positions of two stimulating mice are switched. (B)

Representative traces of one experimental mouse during the three phases. (C-E) The stay

1094 time in the surrounding area of each pen box during three phases;(C) Habituation

1095 phase;(D) Recognition phase; (E) Preference phase. Samples sizes for the PPT are wild

1096 type(WT) (n=12), Koi-1: m$Oxtr^{-/-}$ (n=14), Koi-2: m$Oxtr^{-/-}$ (n=13) Koi-3: m$Oxtr^{-/-}$ (n=12)

1097 Koi-4 : m$Oxtr^{-/-}$ (n=14). All the experimental mice are female. (F) The illustration of the

1098 procedure of the maternal behavior test. (G) Pup retrieval latency and (H) crouching time

1099 were recorded and analyzed. Sample sizes for the maternal behavior tests are WT (n=15),

1100 Koi-3: m$Oxtr^{-/-}$ (n=18), Koi-4: m$Oxtr^{-/-}$ (n=18). Significance of interaction between

1101 independent factors: # $p<0.05$; # # $p<0.01$. Significance of main effect or single main

1102 effect of genotype: ＊$p<0.05$; ＊＊$p<0.01$; ＊＊＊$p<0.001$; ＊＊＊＊

1103 $p<0.0001$.(Illustrations created with Biorender.com). All Koi mice used for this analysis

1104 were pv$Oxtr$-$P2A$-Cre: m$Oxtr^{-/-}$ mice in which the pvOXTR was expressed while

1105 endogenous mOXTR was absent.

1106

1107 **Fig.7 Heatmap of the chromatin contacts surrounding *Oxtr* and *Oxt* across mouse**

1108 **tissues/cells from megabase-size visualization.** (A) Heatmap of the chromatin contacts

1109 surrounding *Oxtr* across mouse tissues/cells: *Oxtr* is located at a TAD boundary region.

1110 The increase of interactions characterizes the TADs structure and the interaction of two

1111 neighboring TADs can be more clearly observed in brain-related samples. Blue dashed

1112   lines were prepared to help observing the TAD structures, and black circles were prepared

1113   to help observing the long-distance interaction obtained from Hi-C interaction matrices

1114   surrounding *Oxtr*. (B) Heatmap of the chromatin contacts surrounding *Oxt* across mouse

1115   tissues/cells. No TAD structure was observed surrounding *Oxt* locus.

1116

Figures

Fig.1 The construction of Koi lines

Fig.2 Transgenic CRE mediated Lac-Z reporter gene expression in the brain and mammary gland of Koi line

Fig.3The expression of OXTR in the brain of Koi lines.

Koi-3:*mOxtr-/-*   **Koi-4:*mOxtr-/-***

Fig.4 mRNA expression of *Oxtr* in the adult brain of Koi lines.

Fig.5 Quantitative PCR analysis of *Oxtr* mRNA expression in different brain regions of Koi lines.

Fig.6 Social behaviors of Koi lines.

Fig.7 Heatmap of the chromatin contacts surrounding *Oxtr* and *Oxt* across mouse tissues/cells from megabase-size visualization.

Supplementary Information

Supplementary Fig.1 Expression pattern of the *Pkcd-Cre* BAC transgene.

All five independent *Pkcd-Cre* BAC transgenic mouse lines (A–E) exhibited expression

patterns that closely resembled the endogenous *Pkcd* expression, as shown by in situ

hybridization (F). The *Pkcd-Cre* BAC construct was generated using the same strategy as

that employed for the Koi lines, incorporating the entire intergenic regions both upstream

and downstream of the *Pkcd* gene.

Supplementary Fig.2 Transgenic CRE mediated Lac-Z reporter gene expression in the brain of Koi lines. X-gal staining of the brains from the double positive (*Cre+*, *LacZ+*) offspring of Koi-5 (A-C), Koi-6 (D-F), Koi-7 (G-I), Koi-8 (J-L). Scale bar=1mm

Supplementary Fig.3 Negative control of whole-mount mammary gland staining.

X-gal staining of a mammary gland from (*Cre*-, *LacZ*+) offspring from heterozygous Koi-

4 was shown. The lymph node showed some blue signal, indicating that the lymph node

could be stained nonspecifically. All the epithelium ducts did not show any blue signal,

verifying the specificity of duct labeling of X-gal staining. Scale=500um

	track	ID	Cell/tissue	Organism	Restr. Enz.	Database	Database ID	Browser	Publication
Hi-C	1	Whole brain	whole brain	mm10	DNaseI	4DN	4DNFIQ48NYOW	WashU	Deng 2015
		https://4dn-open-data-public.s3.amazonaws.com/fourfront-webprod/wfoutput/60d98cc6-4583-4ebe-9c5a-77fc421ef560/4DNFIQ48NYOW.hic							
	2	NPC	NPC	mm10	DpnII	4DN	4DNFIHW8NTQX	WashU	Bonev 2017
		https://4dn-open-data-public.s3.amazonaws.com/fourfront-webprod/wfoutput/219497b5-3f35-473b-9d4e-e1cf21c69561/4DNFIHW8NTQX.hic							
	3	Cortex	cortex	mm10	MboI	4DN	4DNFII3JV8I1	WashU	Du 2017
		https://4dn-open-data-public.s3.amazonaws.com/fourfront-webprod/wfoutput/268b7d52-9655-474c-9467-8ba31bb2195c/4DNFII3JV8I1.hic							
	4	ESC	ESC	mm10	DpnII	4DN	4DNFIU8AF5ZY	WashU	Bonev 2017
		https://4dn-open-data-public.s3.amazonaws.com/fourfront-webprod/wfoutput/c2eaf9bf-9584-4cec-8685-bd74038a6c01/4DNFIU8AF5ZY.hic							
	5	Hepatocyte	hepatocyte	mm10	HindIII	4DN	4DNFIFULDMGN	WashU	Schwarzer 2017
		https://4dn-open-data-public.s3.amazonaws.com/fourfront-webprod/wfoutput/57ecaa79-5c9c-49c7-b841-0a4bf3117555/4DNFIFULDMGN.hic							
	6	Sperm	sperm	mm10	MboI	4DN	4DNFIN8F14CS	WashU	Du 2017
		https://4dn-open-data-public.s3.amazonaws.com/fourfront-webprod/wfoutput/27f54fcb-54fe-41a4-b25a-2f8944c89044/4DNFIN8F14CS.hic							

	Fig	Regions	Genes
WashU	A	chr6:104149809-119528368	Oxtr
	B	chr2:121804885-139358632	Oxt

Supplementary Table.1

The datasets IDs and sample information for mouse 3D chromatin verification

The database source of each mouse sample, analysis and visualization tool information, and the region analyzed was listed.