Network topology in brain tumor patients with and without structural

epilepsy- a prospective MEG-study

Barbara Ladisich[1,2#], Stefan Rampp [3#], Eugen Trinka[4,5,6], Nathan Weisz [4,5,7], Christoph

Schwartz[1], Theo Kraus[8], Camillo Sherif[2], Franz Marhold[2], Gianpaolo Demarchi[5,7]

[1]Department of Neurosurgery, Christian Doppler University Hospital, Paracelsus Medical University, Salzburg,

Austria

[2]Department of Neurosurgery, University Hospital St.Poelten, Karl Landsteiner University of Health Sciences,

Austria

[3]Department of Neurosurgery, University Hospital Erlangen, Germany

[4]Department of Neurology, Christian Doppler University Hospital, Paracelsus Medical University, Center for

Cognitive Neuroscience Salzburg, Member of the European Reference Network, EpiCARE, Salzburg, Austria

[5]Neuroscience Institute, Christian Doppler University Hospital, Salzburg Austria

[6]Karl Landsteiner Institute of Neurorehabilitation and Space Neurology, Salzburg, Austria

[7]Center for Cognitive Neuroscience & Department of Psychology, Paris Lodron University Salzburg, Austria

[8]Institute of Pathology, University Hospital Salzburg, Paracelsus Medical University, Salzburg, Austria

[#]Contributed equally

Corresponding author:

Barbara Ladisich

barbara.ladisich@pmu.ac.at

Disclosures

This project was partly supported by research funding of Österreichische Krebshilfe Salzburg.
There is no conflict related to the content of this study.

Abbreviations and acronyms

BF: Bayes factor

BM: brain metastasis

CC: clustering coefficient

DNET: dsyembryoplastic neuroepthelial tumor

EEG: Electroencephalography

FC: Functional connectivity

FDR: False discovery rate

fMRI: Functional magnetic resonance imaging

GBT: Glial brain tumor

HC: Healthy control

HGG: High-grade glioma

HPI: Head position indicator

IDH: Isocitrate dehydrogenase

L: Shortest path length

LGG: Low-grade glioma

MEG: Magnetoencephalography

MRI: Magnetic resonance imaging

ND: Node degree

NT: network topology

PA: Patient

PCNSL: Primary CNS lymphoma

PSD: Power spectral density

PNSE: Patients without symptomatic epilepsy

PSE: Patients with symptomatic epilepsy

RBC: Rank biserial coefficient

SSS: Signal space separation

SSP: Signal space projection

wb: Whole brain

Abstract

It has been proposed that functional connectivity (FC) and network topology (NT) are altered

in patients with glial brain tumors. So far there is no consensus on the pattern of these

changes, furthermore data on FC in patients with brain metastases (BMs) as well as on the

presence and absence of tumor-related epilepsy is lacking.

We aimed to analyze preoperative NT of newly diagnosed, singular, supratentorial glial brain

tumors (GBTs) and BMs with and without structural epilepsy.

FC and NT derived from resting state magnetoencephalography (MEG) were compared between patients (PAs) and matched healthy controls (HCs), between GBTs and BMs as well as between patients with and without structural epilepsy. We investigated all patients, who met our inclusion criteria from February 2019 to March 2021. Clinical data was collected from patients' electronic medical charts. We analyzed whole brain (wb) connectivity in six frequency bands, calculated three different network topological parameters (node degree (ND), shortest path length (L), clustering coefficient (CC)) and performed a stratification, where differences in the power were to be found. For data analysis we used Fieldtrip, Brain Connectivity Matlab toolboxes and in-house built scripts.

We consecutively included 41 patients (21 men), mean age 60.1ys (range 23- 82), who were operated on at our institution. Tumor histology included high-grade gliomas (n=18), low-grade gliomas (n=4), dysembryoplastic neuroepithelial tumor (DNET, n=1), BMs (n=14) and others (n=4). Statistical analysis revealed a significant decrease for wb ND in patients compared to healthy controls in every frequency range analyzed at the descriptive and corrected level (p_{1-30Hz}=0.002, p_{γ}=0.002, p_{β}=0.002, p_{α}=0.002, p_{θ}=0.024, p_{δ}=0.002). Furthermore, at the descriptive level, we found a significant augmentation for wb CC (p_{1-30Hz}=0.031, p_{δ}=0.013) in PAs compared to HCs, which did not persist the false discovery rate (FDR) correction. There were no differences in the networks of glial brain tumors and metastases identified. However, we found a significant increase in wb CC in patients with structural epilepsy (p_{θ}= 0.048), and significantly lower wb ND (p_{α}= 0.039) at the uncorrected level.

Our data suggests that network topology is altered in brain tumor patients, which is in line with previous studies. Tumor histology per se might not influence the brain's functional network, however, tumor-related epilepsy seems to do so. Longitudinal studies and in-depth analysis of possible factors and confounders might be helpful to further substantiate these findings.

Keywords: magnetoencephalography, resting-state, network topology, brain tumor, brain metastasis, epilepsy

Introduction

Several symptoms observed in patients with brain tumors, such as attention deficits or

reduced psychomotor speed, affecting quality of life, cannot be explained solely by the

location of the lesion but might be explained by changes in the underlying neuronal network

(Bartolomei et al., 2006).

The neuronal network is often characterized by means of functional connectivity and network

topology (Bullmore and Sporns, 2009; Stam et al., 2009; C.J. Stam, 2010; Stam and van

Straaten, 2012). Functional connectivity describes how information is transmitted locally as

well as between different areas and is defined by statistical interdependencies of signals

resulting from brain activity (Guggisberg et al., 2008; Stam et al., 2007). These signals can be

measured by functional neuroimaging techniques such as functional magnetic resonance

imaging (fMRI), or neurophysiological procedures, such as electroencephalography (EEG) or

magnetoencephalography (MEG).

Network topology describes how the elements defining the network are arranged and how

they interact. It is based on a mathematical graph theory that applies to various networks. This

model has been increasingly applied in the fields of cognitive and clinical neuroscience,

showing that functional and neuronal networks have so called "small-world" properties

(Douw et al., 2011; Stam et al., 2009). Small-world networks are characterized by short path

lengths (L) and high clustering coefficient (CC) (Barahona and Pecora, 2002). Douw et al.

calculated a small-world coefficient (ratio between CC and L) from resting-state MEG and

performed neuropsychological testing in healthy controls and found that higher small-

worldness significantly correlated with better cognitive performance (Douw et al., 2011). A

short path length indicates that nodes of a network relate to each other via short paths and

high clustering describes that the probability of several nodes to form connections is high

(Boccaletti et al., 2014). The node degree is another network topological parameter, which is

an absolute value that specifies the number of connections of one single node (Vogel et al.,

2021).

Among adult intraparenchymal brain tumors, brain metastases are the most common,

followed by gliomas (Nayak et al., 2012). Even though prior studies already investigated the

functional network of patients with glial brain tumors (GBTs), non-glial lesions and

meningiomas, brain metastases (BMs) – to the best of our knowledge- have not yet been

investigated in regard to their impact on the brain's network (Bartolomei et al., 2006; Bosma

et al., 2009; Derks et al., 2019, 2021; Douw et al., 2013; Semmel et al., 2021; van Dellen et

al., 2012; van Nieuwenhuizen et al., 2018; Wang et al., 2010; Yang et al., 2022).

Even though low-grade gliomas (LGGs) and high-grade gliomas (HGGs) both derive from

neuroepithelial tissue, they are different in terms of malignancy/ -clinical outcome, growth

patterns, molecular signature, and protein expression (Louis et al., 2016). In contrast, BMs are

secondary neoplasms that derive from completely different cell lines. Tumors from the lung,

prostate, breast, melanoma, renal cell carcinoma most commonly spread to the brain (Nayak

et al., 2012). The growth pattern of brain metastases is distinct from glial brain tumors, which

can be visualized radiologically and intraoperatively. High-grade, as well as low-grade

gliomas mostly infiltrate the brain parenchyma, whereas BMs usually only displace the

surrounding tissue.

The authors of previous studies grasped to identify reasons for network changes in brain

tumor patients. They proposed that histology and tumor grade may explain and directly

impact the extent of network changes (Derks et al., 2019; van Dellen et al., 2012). It has been

previously described that LGGs have a different influence on the network characteristics than

HGGs, namely decreased synchronizability and reduced global integration (van Dellen et al.,

2012). Further it has been found that patients with non-glial lesions, LGGs and HGGs show

altered functional connectivity and impaired network topology, especially in the theta band,

compared to healthy controls (Douw et al., 2010; van Dellen et al., 2012). These changes

were also associated with impaired cognition and weren't found to be explicable by the

tumors' anatomical location.

Another mechanism for network changes in brain tumor patients might be structural

(symptomatic) epilepsy. Tumors causing structural epilepsy might have an even greater

negative impact on the functional network compared to those that do not. It has been

suggested that LGGs have a greater impact on network topology than HGGs (Bosma et al.,

2009; Douw et al., 2013, 2010; van Dellen et al., 2012). This corresponds to the fact that

LGGs also cause structural epilepsy more often than HGGs. Douw et al. further investigated

tumor related epilepsy and found it to be interdependent with network characteristics and even

to be related to protein expression (Douw et al., 2013). Previous MEG-studies on epilepsy

found increased functional connectivity for focal and generalized epilepsy, as well as

evidence for an increase in seizure frequency by augmented dissociation of the functional

network (Douw et al., 2015; Li Hegner et al., 2018). So far, however there has not yet been

any investigation published that compared the network topology of patients with versus

without tumor related epilepsy.

Network changes, i.e., could be the cause or consequence of diseases. It might be lesions, like

brain tumors, or pathological processes, like in Alzheimer's/ Parkinson's Disease or Epilepsy,

that provoke alterations in functional networks (Douw et al., 2015; Engels et al., 2017; Hyder

et al., 2021). Conversely, the distortion of the network could also generate the symptoms, or

rather the diseases themselves.

Compared to the different impact of low-grade gliomas and high-grade gliomas on functional

connectivity, we hypothesized that brain metastases would interact even more differently with

their environment, respectively within the brain's functional network, in comparison to the

group of glial brain tumors.

And since previous studies found the FC in epilepsy to be clearly impaired, (Douw et al.,

2015; Li Hegner et al., 2018; Rampp et al., 2021; Vogel et al., 2021) we now aimed to further

investigate the effect of tumor-related epilepsy on NT.

Study objective

The main objective of this study was to investigate whether network topology is altered in

patients with brain tumors and whether histology or the presence of structural epilepsy have

roles in these alterations.

We also aimed to confirm results from previous studies by applying current and most

common network parameters and intended to add further information to the understanding of

(pathological) network topology by investigating brain metastases.

We hypothesized that there is a relevant difference in network topology of patients with brain

tumors compared to healthy controls, that BMs have a different influence on the brain's

network compared to GBTs and that structural epilepsy relevantly influenced network

topology compared to patients without symptomatic seizures.

Methods

Study population

We included all consecutive adult patients who were referred to our department for

neurosurgical resection or biopsy of a single, supratentorial tumor with MRI-radiological

suspicion of either being a glial brain tumor or brain metastasis from 4[th] of February 2019 to

21[st] of March 2021.

We excluded patients who had undergone prior neurosurgery or radiotherapy of the brain,

patients with unrelated psychiatric, neurological, endocrine comorbidities or implanted (non-

titanium) metallic devices.

We matched every patient with a healthy control (HC) in gender and age (+/-5 years). The

control group consisted of staff of the University Hospital Salzburg, as well as of students

from the Paris-Lodron-University, Salzburg, Austria.

Both groups underwent a single 20-minute resting-state MEG recording. For co-registration

with MEG data, a template MRI was used for HCs, whereas patients' T1- pre- and post-

gadolinium or T2- sequences were employed for the tumor group.

Clinical data was retrieved from patients' electronic medical charts. In particular, the presence

of structural epilepsy was recorded. In most patients with structural epilepsy, tumors had been

initially detected due to the occurrence of seizures. Whenever this was unclear, an EEG was

performed, and a neurological consultation was conducted.

The patient cohort was divided into two subgroups: glial brain tumors (GBTs) and brain

metastases (BMs), and patients with structural epilepsy (PSEs) and patients without structural

epilepsy (PNSEs).

Ethics and dissemination

The study was reviewed and approved by the local ethics committee prior to recruitment

(415-E/2412/6-2018). All eligible subjects gave their written informed consent to participate

in this study. The Declaration of Helsinki was respected when conducting this study.

MEG recordings

We recorded the brains' magnetic signal at 1000 Hz (hardware filters: 0.1–330 Hz) in a

standard passive magnetically shielded room (AK3b, Vacuumschmelze, Germany) using a

whole head MEG (Neuromag Triux, MEGIN Oy, Finland). Signals are captured by 102

magnetometers and 204 orthogonally placed planar gradiometers at 102 different positions.

All subjects wore nonmetallic clothes and had 5 head position indicator (HPI) coils attached

on the glabella, forehead, as well as preauricularly bilaterally. They all underwent head shape

acquisition prior to the recording, and the head shape points were later used for coregistering

the anatomical MRI scan. During the measurement, subjects were seated and instructed to

keep their eyes open. We periodically checked if subjects stayed awake via a camera in the

magnetically shielded room. Every measurement consisted of one block of 20 minutes.

Data analysis

To preprocess the data, we employed a signal space separation (SSS) algorithm that is implemented in the Maxfilter program (version 2.2.15), provided by the MEG manufacturer (Taulu and Simola, 2006). Here external noise from the MEG signal is removed (mainly 16.6 Hz train mains supply, and 50 Hz plus harmonics) and data is realigned to a common standard head position (*-trans default* Maxfilter parameter). This is performed across different blocks based on the measured head position at the beginning of each block using the five coils that have been placed before the measurement.

We used Brainstorm (Version 181008 (08-Oct-2018)) to further preprocess the data (Tadel et al., 2011). At first, we applied a high-pass filter at 0.1 Hz (6th order zero-phase Butterworth filter) to the continuous data. Subsequently, for signal space projection (SSP), continuous data were chunked in 10 s blocks (Uusitalo and Ilmoniemi, 1997). The resulting data was manually scrutinized to identify remaining eye blinks, eye movements, heartbeat and the 16 and 2/3 Hz train power supply artifacts and to create appropriate projectors. Finally, the artifactual components were projected out and the continuous data was segmented in 2s chunks. Then, data was again visually scrutinized to find bad epochs that were not corrected by the SSS and SSP combination, and further removed. An average of 15.4 (range 0- 152.6) trials, corresponding to approximately 0.5 minutes were therefore removed, leaving roughly 19 minutes of artifact free data per subject (range 15-20 min) that was used for further analysis.

The subsequent steps were then performed using the *fieldtrip* toolbox (git version 20180124) (Oostenveld et al., 2011). At first, the data was downsampled to 300 Hz, to speed up the following computations of the source space connectivity and graph theoretical values.

Then the coordinate frame of the MEG data was coregistered to a template MRI scan in the case of healthy controls, and to T1-contrast enhanced or T2 anatomical images (depending on best visualization of the tumor) in patients, using the right and left preauricular points, nasion and the head shape points measured during preparation. A single shell source model was generated based on the segmented anatomical scan and a per subject source model with resolution 10mm was created (Nolte, 2003). Next, the data was Fourier transformed for each of the frequency bands of interest ('broadband'=1-30 Hz, gamma=25-40 Hz, beta=15-25, alpha=8-15 Hz and theta=4-7, delta=1-4) was computed and used for a partial canonical coherence beamformer, with a noise covariance regularization factor of 5% (Oostenveld et al., 2011).

From the resulting source activity, for each position of the source model the component of

the dipole moment projected onto the dominant direction was taken and subsequently used to

compute the imaginary part of the coherence as an all-to-all whole brain voxel-wise

connectivity metric. The 95th percentile of coherence values was used as a threshold for the

construction of the adjacency matrix for computing node degree (ND), clustering coefficient

(CC), shortest path length (L). All source, connectivity and network analyses were carried on

using the fieldtrip toolbox (git version 20180124) (Oostenveld et al., 2011).

To minimize the spurious connectivity effects driven purely by the power difference, we

decided to stratify the data to be sent to source space for connectivity and network analysis.

For selecting the strata, two big pools containing the power spectral densities values were

computed overall 2s blocks on the magnetometers, for the patients and for the healthy controls

respectively. Then, the fieldtrip function *ft_stratify* (with the parameters method = 'histogram',

equalbinavg = 'yes', numbin = 20) was employed to select the trials and equalize the two

power spectral distributions, per each frequency band of interest.

Surgery and histopathology

All decisions on neurosurgical treatment (consisting of either microsurgical tumor resection or

biopsy) were agreed upon in the interdisciplinary neuro-oncological tumor board. Patients

gave their written, informed consent prior to surgery.

Histopathological examinations (frozen section, formalin-fixed and paraffin embedded tissue,

immunohistochemical and molecular genetic analysis) were performed according to the 2016

World Health Organization Classification of Tumours of the central nervous system, the

current edition available during the inclusion period/ histological sampling (Louis et al.,

2016).

Statistical analysis

For statistical analysis we used Jamovi, Version 1 and R studio, Version 2022.02.1 (Şahin and Aybek, 2019; Wickham et al., 2019). As a dependent variable the maximum value of each network measure, per subject/patient and per frequency band of interest, was taken, that is node degree (ND), clustering coefficient (CC) and shortest path (L). We tested for differences in age in the patient population by means of Student's t-test and Welch's test. We designed a contingency table for one subgroup (PSE vs PNSE) -for which it was applicable- and calculated Fisher's exact test. For the comparison between the groups 1) PAs vs HCs, 2) GBTs vs BMs and 3) PSEs vs PNSEs Shapiro Wilk test for normality was calculated. When the normality assumption was met, we performed independent Student's T-tests, when it was violated, Mann-Whitney U tests were employed. Consequently, effect sizes were calculated by Cohen's d or Rank biserial correlation (RBC), depending on the test employed. As customary, for p-values lower than 0.05 the null hypothesis was rejected, and the test was considered significant. We additionally provide the Bayes Factors (BF) to inform on the relative levels of evidence for the null/alternative hypotheses. At last, we corrected for multiple comparisons by calculating the false discovery rate (FDR).

Methods prior to statistical analysis are roughly summarized by a graphical depiction in Figure 1.

Figure 1: Summary of data collection and analysis. Patients with glial brain tumors and brain metastases were consecutively included and matched with healthy controls. Presence of structural epilepsy was recorded. MEG examination was performed several days preoperatively, whole brain connectivity was analyzed, resultant analysis of network topological parameters

301 (node degree, shortest path length, clustering coefficient) was performed with Fieldtrip, Brain Connectivity Matlab toolboxes
302 and in-house built scripts.

Results

Clinical data

A total of 42 patients were initially enrolled in the study. One patient diagnosed with acute

disseminated encephalomyelitis by stereotactic biopsy had to be excluded from further

analysis. Thus, the final study population consisted of 41 patients (21 men), as well as 41

healthy controls, who were matched in gender and age. Tumor histology included glial brain

tumors (n=22), dysembryoplastic neuroepithelial tumor (DNET) (n=1), brain metastases

(n=14), meningiomas (n=2) and primary CNS lymphomas (PCNSL) (n=2). The patient with

DNET was investigated together with patients who had glial brain tumors, so that the final

group of GBTs included 23 subjects.

None of the patients had undergone prior tumor treatment, besides corticosteroids or

antiepileptogenic drugs at the time of MEG measurement.

Mean age in patients was 60.5years (+/- 15.2ys). T-Test and Welch's Test revealed that there

was no significant difference in age between patients with GBTs and brain metastases

(p=0.260 and p=0.195 respectively).

A total of 27 patients were suffering from structural epilepsy: 15 with GBTs, ten with BMs

and two with primary CNS lymphomas. Fisher's exact test revealed no significant difference

in the analysis of the contingency table.

Patient characteristics are shown in Table 1, 2 and 3.

Network characteristics

We computed the resting state power spectral density (PSD) analysis for patients (PAs) and

healthy controls (HCs) that revealed significant differences (p<0.001, RBC= 0.65, BF_{10}=907).

Therefore, to minimize the spurious connectivity effects driven purely by the power

difference, we decided to stratify the data to be sent to source space for connectivity and

network analysis. In the literature, there seems to be no standard way to summarize the whole

brain network parameters, being mean, median, or maximum. Here, for simplicity, we report

only the maximum values of the whole brain network parameters, but the usage of the mean

and median delivers similar results (Hillebrand and Stam, 2019).

Figure 2: T-test or Mann-Whitney U test comparison (depending on normality distribution) between patients (red) and healthy controls (blue) revealed significantly lower node degree in patients at the corrected level (p_{1-30Hz} =0.0015, $p\gamma$=0.0024, $p\beta$=0.0015, $p\alpha$=0.0015, p_θ=0.024, p_δ=0.0015).

We found a highly significant lower node degree at the uncorrected level in patients compared to healthy controls (p_{1-30Hz} <0.001, d=-0.86, BF_{10}=110.80; $p\gamma$=0.002, d=-0.72, BF_{10}=19.35; $p\beta$<0.001, d=-1.13, BF_{10}=6458.72; $p\alpha$<0.001, d=-0.99, BF_{10}=767.93; p_θ=0.024, RBC=0.29, BF_{10}=2.59; p_δ<0.001, RBC=0.42, BF_{10}=27.26) After the correction for multiple comparison, we still found a highly significant result in all six frequency bands as shown in Figure 1. Further analysis at the uncorrected level demonstrated significantly higher clustering coefficient (CC) (p_{1-30Hz}=0.031, RBC=0.49, BF_{10}=1.83; p_δ=0.013, d= 0.32, BF_{10}=9.07) in patients compared to healthy controls. However, the latter results of higher CC in patients showed a tendency but did not stand the correction (p_{1-30Hz}=0.093, p_δ=0.078).

For the comparison of the two tumor groups (glial brain tumors (GBTs) vs. brain metastases (BMs), we excluded the four patients with primary CNS lymphomas and meningiomas, respectively leaving 23 patients in the cohort of GBTs and 14 patients in the cohort of BMs. (Table 2). There was no discrepancy in power between the two groups (p=0.676, RBC= 0.09, BF_{10}= 0.41) and in further network analysis we didn't find a significant difference in any of the parameters and frequency bands.

We also compared patients with structural epilepsy against patients without structural

epilepsy. For this analysis, we investigated all 41 patients- 27 patients with versus 14 patients

without structural epilepsy (Table 3). Here no power spectral difference between the groups

was found (p=0.091, RBC=0.33, BF_{10}= 0.66), so that no further stratification was needed.

There was significantly higher maximum clustering (p_θ= 0.048, RBC=0.38, BF_{10}= 1.12), but

significantly lower maximum degrees (p_α= 0.039, RBC=0.40, BF_{10}= 1.09) at the uncorrected

level in PSE. However, after calculating the false discovery rate the two results did not stand

the correction (clustering coefficient: p_θ= 0.288; node degree: p_α= 0.095).

Results are listed in Tables 4-7 as well as Figures 2 and 3.

363
364 **Figure 3:** Representative coherence plot thresholded at 75% of the maximum respective coherence of a patient and its
365 matched control: on the left is patient 2, who was diagnosed of a metastasis in the right parietal lobe, on the right is its
366 matched control depicted.

Discussion

It was previously described that different kinds of neurological disorders, such as brain

tumors lead to deviations in the brain's functional network (Bartolomei et al., 2006; Cho et

al., 2021; De Baene et al., 2017; Derks et al., 2021, 2019; D'Souza et al., 2021; Douw et al.,

2013; Huang et al., 2014; Hyder et al., 2021; Na et al., 2018; van Dellen et al., 2012; Wang et

al., 2010).

In this prospective study we investigated 41 patients with brain tumors and 41 matched

healthy controls. We were able to identify significant differences in the network topology of

patients compared to healthy controls, as well as between patients with and without structural

epilepsy. Unexpectedly, we found no differences in the functional networks of patients with

glial brain tumors compared to brain metastases.

Comparison of patients and healthy controls

In the comparison of patients and healthy controls we found a significantly lower node degree

in patients in all frequencies analyzed. This result is also consistent with the definition of node

degree since the impact of the tumor on the functional network will impair the formation of

connections and therefore lead to lower degree networks (Vogel et al., 2021).

We found an increased clustering coefficient in brain tumor patients at the descriptive level,

but this result did not stand up to correction for multiple testing.

The clustering coefficient is a value that is defined by the probability that neighbors of a hub

are connected to another. It is an essential characteristic of the small-world concept that

describes an optimally functioning network (Bullmore and Sporns, 2009; Douw et al., 2011;

Stam et al., 2009). This result appeared unusual at first sight, since an intact network is

assumed to have high clustering and we would have expected these parameters to be

decreased in a pathologically disturbed network, as it is the case in a patient with a brain

tumor. Several previously conducted studies have also described increased clustering in brain

tumor patients (Bosma et al., 2009; van Dellen et al., 2012; Derks et al., 2021). Bosma et al.

found a higher clustering coefficient within the theta band in patients with low-grade gliomas

compared to healthy controls (HCs) (Bosma et al., 2009). They contextualized this result in

the setting of other neurological diseases such as major depression, autism disorders or

Alzheimer's disease, for which previous investigations have found similar network changes.

Also, Derks et al. described higher global clustering in glioma patients compared to HCs.

They further investigated this finding by differentiating various factors (e.g., age, histology,

local and global clustering) but despite this, could not find any relation to increased clustering

of any of these aspects (Derks et al., 2021). As such, it was concluded that the probable cause

of the increased clustering lay with diffuse, whole brain alterations in these patients.

The augmentation might also be an expression of reactive changes of the network to the

tumor. This assumption seems at least conclusive from a clinical point of view since lesions

also provoke morphological and immunochemical reactions in the surrounding parenchyma

that can be displayed radiologically and by the laboratory.

Comparison of patients with glial brain tumors and brain metastases

We further performed two analyses solely within the patient cohort. The first one between the

patients with glial brain tumors (GBTs) versus brain metastases (BMs). For this we excluded

four patients, due to their histological diagnoses- two showed PCNSL, the other two had

meningiomas. The latter two patients were primarily included since both had an oncological

history and from a neuroradiological point of view it had not been possible to rule out a BM.

There was one patient, in which we had the initial suspicion of a LGG, which then turned out

to be a DNET. Although DNETs are a distinct tumor entity, and do not belong to the group of

gliomas, we kept this case included within our cohort of GBTs, because of clinically similar

behavior.

Since data on the impact of BMs on the brain's functional network is lacking, we chose to

investigate those patients in this context. We expected metastases to show different patterns of

network changes, due to the completely different type of tumorigenesis and growth patterns.

However, this hypothesis cannot be supported by our data since no difference in the

functional networks of GBTs and BMs was found, - neither in the variables nor the

frequencies analyzed.

Previous studies had found different network patterns in HGGs and LGGs, which had led to

the conclusion that histology plays an important role in network changes (Derks et al., 2019;

van Dellen et al., 2012). Derks et al. even found significant differences in the

immunohistochemical profile of tumors with lower alpha functional connectivity (FC) in

IDH-wildtype gliomas compared to IDH-mutated gliomas. Nonetheless they did not find

differences in FC in the theta band between the two groups, and there was also no difference

in the alpha band in the comparison of classic histological grading between grade II/III and IV

(Derks et al., 2019).

Our results might be limited by the fact that the two tumor groups were not exactly equally

distributed. This is due to the prospective setting of this study and the fact that single

(supratentorial) brain metastases are rare.

Still, it is a noticeable finding, and if metastases and gliomas are indeed comparable regarding

their impact on network topology, then the main factors may be the mass effect or growth

rate, rather than histology. Metastases usually show less invasive growth patterns compared to

intrinsic brain tumors but may cause substantially more mass effect due to common extensive

peritumoral edema. Therefore, if these tumors do not lead to different changes in the

functional network, histology might not be an explanation for the disturbed network.

Comparison of patients with and without structural epilepsy

Another possible mechanism of the distortion of the functional network could be the presence

of seizures. We therefore also conducted a comparison between patients with and without

structural epilepsy. Here, we found significantly higher clustering (CC) and lower node

degree (ND) in PSEs at the uncorrected level. After calculating the FDR these results did not

retain significance.

However, we still want to discuss the finding of the lower ND in patients with structural

epilepsy, since it stands in contrast to previous studies that presented data from patients with

drug resistant focal epilepsies (Rampp et al., 2021; Vogel et al., 2021). In their data functional

connectivity and node degree was augmented in patients with epilepsy compared to healthy

controls. One explanation for this contradiction might be that tumor-related structural epilepsy

has a different impact on the network than non-tumor-related focal epilepsy. Patients with

epilepsy mostly suffer from their disease for many years, in contrast to patients with brain

tumors who are primarily structural with seizures and shortly after receive diagnosis and

treatment for the lesion. In the situation of long-lasting epilepsy, with many events of

seizures, the network might distort and adapt completely differently than in structural

epilepsy.

In this study, we applied graph theoretical measures since they have proven to be a valuable

tool to characterize the network topology of the brain (Bullmore and Sporns, 2009; Heimans

and Reijneveld, 2012; Reijneveld et al., 2007; Stam and Reijneveld, 2007). Some of the most

current network topology parameters are clustering coefficient, path length and node degree

(Boccaletti et al., 2014; Bosma et al., 2009; van Dellen et al., 2012; Vogel et al., 2021; Wang

et al., 2010). Until now there is only one review on the topic by Semmel et al., who found that

the methodology is often not comparable and that the results are sometimes even

contradictory between different publications (Semmel et al., 2021).

Therefore, we also carried out functional connectivity and network topology calculations in

source space, compared to other previous studies that computed the connectivity measures in

sensor space only. It is crucial to measure in source space, especially in patients with

morphological brain pathologies (Stufflebeam, 2011).

Whenever we found significant results in a network topological parameter, the increase/

decrease compared to healthy controls was concordant in the other frequency bands. To us,

this supports the reliability of the results.

On the contrary, another study found, for example, an increase of synchronization and

clustering in theta, but a decrease in beta (Bosma et al., 2009). They argued that there was

more small world organization in lower frequency bands than in higher ones.

It seems that this aspect needs further investigation.

Strengths and limitations

This is a prospective, experimental, single-center study that aims to further increase the

neurophysiological understanding of how brain tumors disturb the brain's functional network.

The study population might be small but is comparable to other studies (Bartolomei et al.,

2006; Bosma et al., 2009; Derks et al., 2021, 2019; van Dellen et al., 2012; van

Nieuwenhuizen et al., 2018; Wang et al., 2010). Due to the prospective setting, and because

we included patients in a consecutive fashion, subjects were not exactly equally distributed in

the two groups (glial brain tumors vs brain metastases and patients with vs without structural

epilepsy). On the one hand, this hampers the comparableness between groups, but on the other

side we can provide higher data quality and therefore higher significance.

We put a focus on rigorous methodology for best possible comparability with previous and

future studies. Therefore, we only investigated patients with a newly diagnosed, singular,

supratentorial mass lesion, suspected to be a primary brain tumor or brain metastases. Further

we arranged MEG measurement for all patients a few days, and up to a maximum of two

weeks prior to surgery. In this way, we aimed to prevent possible influences of previous

treatment, the presence of new lesions or additional tumor growth on the results of network

analysis. A defined timing of MEG with respect to treatment can help to define standards in

network analysis.

To ensure the significance of our results, we corrected for multiple comparisons, as we

compared 6 different frequency bands for every network parameter. After this, the

comparison of node degree in patients and healthy controls was still highly significant.

Admittedly the increased clustering coefficient in patients was not found to be persistently

significant, but there was still a noticeable tendency to be seen. The corrected results of the

comparison between patients with and without structural epilepsy were not found to be

significant. Nevertheless, we mentioned these results, because they differ from previous

publications and we expect this distinction to be an interesting aspect for future investigations.

So far, there exist only seven other studies on the topic of functional connectivity and network

topology measured by MEG in brain tumors (Bartolomei et al., 2006; Bosma et al., 2009;

Derks et al., 2021, 2019; van Dellen et al., 2012; van Nieuwenhuizen et al., 2018; Wang et al.,

2010). Although previous studies tried to draw conclusions from the influence of histology on

network alterations, BMs have not been investigated, though, from a pathohistological point

of view, they differ the most from primary brain tumors. We aimed to close this gap through

our analysis, however our results did not support the role of histology in network disruption.

Conclusion

We provide data of a prospective study on patients with brain tumors with and without

structural epilepsy compared to healthy controls. We found significant differences in several

network parameters in the broad band, as well as in all five different frequency bands. We did

not find differences in the functional network of primary brain tumors compared to brain

metastases, but there were significant differences at the descriptive level between patients

suffering from structural epilepsy compared to those that did not.

Originating from the results of this study the extent of network disruption in patients with

brain tumors seems to be influenced more by the presence of structural epilepsy, than by the

histology.

We advocate further prospective studies on even larger cohorts of patients. Especially the

differentiation of patients with structural epilepsy, patients suffering from epilepsy and the

comparison to healthy controls could be essential.

Appendices

Supplementary information

Tables

Characteristic	GBT	BM	Meningioma	PCNSL	HC
N	23	14	2	2	41
Mean age (years)	58.2	64.1	46	75.8	60.8
Gender					
Male	14	6	0	1	21
Female	9	8	2	1	20

Table 1: BM= brain metastases, HC=healthy controls, GBT= glial brain tumors, N= number, PCNSL= primary CNS lymphoma.

Lesion type		N	Mean age	Male	Female
Glial brain tumors	HGG	18	60	11	7
	LGG	4	46	3	1
	DNET	1	21	0	1
Brain metastases	Lung	6	68	3	3
	Other	3	59	1	2
	Breast	2	53	0	2
	Renal	2	68	1	1
	Melanoma	1	70	1	0

Table 2: DNET= dysembryoplastic neuroepithelial tumor (this patient was analyzed together with the GBTs), HGG= high-grade glioma, LGG= low-grade glioma.

	N with structural epilepsy	N without structural epilepsy
GBTs	15	8
BMs	10	4
Meningiomas	2	0
PCNSL	0	2

Table 3: Presence of structural epilepsy in the patient cohort. Fisher's exact test revealed no significant difference for the contingency table (p= 0.287). GBT= glial brain tumors; BMs= brain metastases; PCNSL= primary CNS lymphoma.

Frequency band	Network characteristic	Max	
		PA	HC
Broadband	ND	227*	287*
	CC	0.236*	0.217*
	L	2.70	2.69
Gamma	ND	302*	391*
	CC	0.270	0.265
	L	2.78	2.75
Beta	ND	288*	397*
	CC	0.281	0.269
	L	2.83	2.82
Alpha	ND	300*	398*
	CC	0.326	0.314
	L	2.96	2.96
Theta	ND	348*	399*
	CC	0.372	0.334
	L	3.09	3.07
Delta	ND	367*	454*
	CC	0.389*	0.338*
	L	3.14	3.20

544 **Table 4:** C= clustering coefficient; HCs= healthy controls; L= shortest path length; ND= node degree; PAs= patients.
545 Maximum values are given. *Values that were found to be significant in later analysis.

Frequency band	Network characteristics	p-value	Effect size	BF$_{10}$
Broadband*	ND*	<0.001	d=-085	110.80
	CC*	0.031	d=0.49	1.83
	L	0.623	d=0.11	0.26
Gamma*	ND*	0.002	d=-0.72	19.35
	CC	0.875	RBC=0.02	0.23
	L	0.523	d=0.14	0.28
Beta*	ND*	<0.001	d=-1.12	6458.72
	CC	0.479	d=0.15	0.29
	L	0.711	d=0.08	0.24
Alpha*	ND*	<0.001	d=-0.99	767.93
	CC	0.705	RBC=0.05	0.30
	L	0.796	RBC=0.03	0.23
Theta*	ND*	0.024	RBC=0.29	2.59
	CC	0.08	d=0.39	0.89
	L	0.717	d=0.08	0.24
Delta*	ND*	<0.001	RBC=0.42	27.26
	CC*	0.013	RBC=0.32	9.07
	L	0.218	d=-0.27	0.45

548 **Table 5:** BF10= Bayes factor$_{10}$; C= clustering coefficient; d= Cohen's d; HCs= healthy controls; L= shortest path length;
549 ND= node degree; PAs= patients; RBC= Rank biserial correlation; *Values that were found to be significantly different
550 between groups.

Frequency band	Network characteristic	Max	
		PSE	**PNSE**
Broadband	**ND**	210	250
	CC	0.214	0.215
	L	2.70	2.68
Gamma	**ND**	299	351
	CC	0.260	0.273
	L	2.87	2.80
Beta	**ND**	291	343
	CC	0.282	0.289
	L	2.90	2.91
Alpha*	**ND***	306*	354*
	CC	0.325	0.340
	L	2.99	2.97
Theta*	**ND**	370	373
	CC*	0.400*	0.361*
	L	3.19	3.15
Delta	**ND**	342	390
	CC	0.357	0.363
	L	3.16	3.11

553 **Table 6:** C= clustering coefficient; L= shortest path length; ND= node degree; PSEs= patients with structural epilepsy;
554 PNSEs= patients without structural epilepsy. Maximum values are given. *Values that were found to be significant in later
555 analysis.

Frequency band	Network characteristic	p-value	Effect size	BF_{10}
Broad band	**ND**	0.055	RBC=0.37	2.22
	CC	0.935	d=-0.02	0.32
	L	0.488	RBC=0.14	0.37
Gamma	**ND**	0.079	d=0.60	1.12
	CC	0.581	d=-0.18	0.36
	L	0.148	d=0.49	0.74
Beta	**ND**	0.078	d=-0.60	1.12
	CC	0.881	RBC=0.03	0.33
	L	0.923	d=-0.03	0.32
Alpha*	**ND***	0.039*	RBC=0.40	1.09
	CC	0.470	d=-0.24	0.39
	L	0.712	d=0.12	0.34
Theta*	**ND**	0.956	RBC=0.01	0.32
	CC*	0.048*	RBC=0.38	1.12
	L	0.494	d=0.23	0.38
Delta	**ND**	0.067	RBC=0.35	0.66
	CC	0.786	d=-0.09	0.33
	L	0.694	RBC=0.08	0.41

558 **Table 7:** C= clustering coefficient; L= shortest path length; ND= node degree; PSEs= patients with structural epilepsy;
559 PNSEs= patients without structural epilepsy; BF= Bayes Factor$_{10}$; RBC= Rank biserial correlation (effect size); *Values that
560 were found to be significantly different between groups.

Acknowledgements

To all my colleagues, friends and family, and the students from Lodron University who took their time to participate in this study as healthy controls.

To Ciara O'Sullivan for providing language help.

To Österreichische Krebshilfe Salzburg for their research funding, that partly supported this project. The financial support was used for data collection and analysis. There was no involvement in writing of the report, or in the decision to submit the article for publication

Data and code availability

The MATLAB scripts used to perform the analysis are available at one author's (GD) repository (github.com/gdemarchi). Down sampled (to 100 Hz) version of the healthy controls raw data will be available at OSF.io upon final publication. The sharing of the patients' data must be instead discussed upon request on a per-case basis with the CDK/PMU Salzburg Ethics committee.

References

Barahona, M., Pecora, L.M., 2002. Synchronization in Small-World Systems. Phys. Rev. Lett. 89, 054101. https://doi.org/10.1103/PhysRevLett.89.054101

Bartolomei, F., Bosma, I., Klein, M., Baayen, J.C., Reijneveld, J.C., Postma, T.J., Heimans, J.J., van Dijk, B.W., de Munck, J.C., de Jongh, A., Cover, K.S., Stam, C.J., 2006. Disturbed functional connectivity in brain tumour patients: Evaluation by graph analysis of synchronization matrices. Clinical Neurophysiology 117, 2039–2049. https://doi.org/10.1016/j.clinph.2006.05.018

Bartolomei, F., Bosma, I., Klein, M., Baayen, J.C., Reijneveld, J.C., Postma, T.J., Heimans, J.J., van Dijk, B.W., de Munck, J.C., de Jongh, A., Cover, K.S., Stam, C.J., 2006. How do brain tumors alter functional connectivity? A magnetoencephalography study. Ann Neurol. 59, 128–138. https://doi.org/10.1002/ana.20710

Boccaletti, S., Bianconi, G., Criado, R., del Genio, C.I., Gómez-Gardeñes, J., Romance, M., Sendiña-Nadal, I., Wang, Z., Zanin, M., 2014. The structure and dynamics of multilayer networks. Physics Reports 544, 1–122. https://doi.org/10.1016/j.physrep.2014.07.001

Bosma, I., Reijneveld, J.C., Klein, M., Douw, L., van Dijk, B.W., Heimans, J.J., Stam, C.J., 2009. Disturbed functional brain networks and neurocognitive function in low-grade glioma patients: a graph theoretical analysis of resting-state MEG. Nonlinear Biomed Phys 3, 9. https://doi.org/10.1186/1753-4631-3-9

Bullmore, E., Sporns, O., 2009. Complex brain networks: graph theoretical analysis of
structural and functional systems. Nat Rev Neurosci 10, 186–198.
https://doi.org/10.1038/nrn2575

Cho, N.S., Peck, K.K., Gene, M.N., Jenabi, M., Holodny, A.I., 2021. Resting-state functional
MRI language network connectivity differences in patients with brain tumors: exploration of
the cerebellum and contralesional hemisphere. Brain Imaging and Behavior.
https://doi.org/10.1007/s11682-021-00498-5

De Baene, W., Rutten, G.J.M., Sitskoorn, M.M., 2017. The Temporal Pattern of a Lesion
Modulates the Functional Network Topology of Remote Brain Regions. Neural Plasticity
2017, 1–11. https://doi.org/10.1155/2017/3530723

Derks, J., Kulik, S., Wesseling, P., Numan, T., Hillebrand, A., van Dellen, E., de Witt Hamer,
P.C., Geurts, J.J.G., Reijneveld, J.C., Stam, C.J., Klein, M., Douw, L., 2019. Understanding
cognitive functioning in glioma patients: The relevance of IDH☐mutation status and
functional connectivity. Brain Behav 9, e01204. https://doi.org/10.1002/brb3.1204

Derks, J., Kulik, S.D., Numan, T., de Witt Hamer, P.C., Noske, D.P., Klein, M., Geurts,
J.J.G., Reijneveld, J.C., Stam, C.J., Schoonheim, M.M., Hillebrand, A., Douw, L., 2021.
Understanding Global Brain Network Alterations in Glioma Patients. Brain Connectivity 11,
865–874. https://doi.org/10.1089/brain.2020.0801

Douw, L., de Groot, M., van Dellen, E., Aronica, E., Heimans, J.J., Klein, M., Stam, C.J.,
Reijneveld, J.C., Hillebrand, A., 2013. Local MEG networks: The missing link between
protein expression and epilepsy in glioma patients? NeuroImage 75, 195–203.
https://doi.org/10.1016/j.neuroimage.2013.02.067

Douw, L., Schoonheim, M.M., Landi, D., van der Meer, M.L., Geurts, J.J.G., Reijneveld, J.C.,
Klein, M., Stam, C.J., 2011. Cognition is related to resting-state small-world network
topology: an magnetoencephalographic study. Neuroscience 175, 169–177.
https://doi.org/10.1016/j.neuroscience.2010.11.039

Douw, L., van Dellen, E., de Groot, M., Heimans, J.J., Klein, M., Stam, C.J., Reijneveld, J.C.,
2010. Epilepsy is related to theta band brain connectivity and network topology in brain tumor
patients. BMC Neuroscience 11, 103. https://doi.org/10.1186/1471-2202-11-103

D'Souza, S., Hirt, L., Ormond, D.R., Thompson, J.A., 2021. Retrospective analysis of
hemispheric structural network change as a function of location and size of glioma. Brain
Communications 3, fcaa216. https://doi.org/10.1093/braincomms/fcaa216

Bartolomei, F., Bosma, I., Klein, M., Baayen, J.C., Reijneveld, J.C., Postma, T.J., Heimans,
J.J., van Dijk, B.W., de Munck, J.C., de Jongh, A., Cover, K.S., Stam, C.J., 2006. Disturbed
functional connectivity in brain tumour patients: Evaluation by graph analysis of
synchronization matrices. Clinical Neurophysiology 117, 2039–2049.
https://doi.org/10.1016/j.clinph.2006.05.018

Boccaletti, S., Bianconi, G., Criado, R., del Genio, C.I., Gómez-Gardeñes, J., Romance, M.,
Sendiña-Nadal, I., Wang, Z., Zanin, M., 2014. The structure and dynamics of multilayer
networks. Physics Reports 544, 1–122. https://doi.org/10.1016/j.physrep.2014.07.001

Bosma, I., Reijneveld, J.C., Klein, M., Douw, L., van Dijk, B.W., Heimans, J.J., Stam, C.J.,
2009. Disturbed functional brain networks and neurocognitive function in low-grade glioma
patients: a graph theoretical analysis of resting-state MEG. Nonlinear Biomed Phys 3, 9.
https://doi.org/10.1186/1753-4631-3-9

De Baene, W., Rutten, G.J.M., Sitskoorn, M.M., 2017. The Temporal Pattern of a Lesion
Modulates the Functional Network Topology of Remote Brain Regions. Neural Plasticity
2017, 1–11. https://doi.org/10.1155/2017/3530723

Derks, J., Kulik, S., Wesseling, P., Numan, T., Hillebrand, A., van Dellen, E., de Witt Hamer,
P.C., Geurts, J.J.G., Reijneveld, J.C., Stam, C.J., Klein, M., Douw, L., 2019. Understanding
cognitive functioning in glioma patients: The relevance of IDH mutation status and
functional connectivity. Brain Behav 9, e01204. https://doi.org/10.1002/brb3.1204

Derks, J., Kulik, S.D., Numan, T., de Witt Hamer, P.C., Noske, D.P., Klein, M., Geurts,
J.J.G., Reijneveld, J.C., Stam, C.J., Schoonheim, M.M., Hillebrand, A., Douw, L., 2021.
Understanding Global Brain Network Alterations in Glioma Patients. Brain Connectivity 11,
865–874. https://doi.org/10.1089/brain.2020.0801

Douw, L., de Groot, M., van Dellen, E., Aronica, E., Heimans, J.J., Klein, M., Stam, C.J.,
Reijneveld, J.C., Hillebrand, A., 2013. Local MEG networks: The missing link between
protein expression and epilepsy in glioma patients? NeuroImage 75, 195–203.
https://doi.org/10.1016/j.neuroimage.2013.02.067

Douw, L., DeSalvo, M.N., Tanaka, N., Cole, A.J., Liu, H., Reinsberger, C., Stufflebeam,
S.M., 2015. Dissociated multimodal hubs and seizures in temporal lobe epilepsy. Ann Clin
Transl Neurol 2, 338–352. https://doi.org/10.1002/acn3.173

Douw, L., Schoonheim, M.M., Landi, D., van der Meer, M.L., Geurts, J.J.G., Reijneveld, J.C.,
Klein, M., Stam, C.J., 2011. Cognition is related to resting-state small-world network
topology: an magnetoencephalographic study. Neuroscience 175, 169–177.
https://doi.org/10.1016/j.neuroscience.2010.11.039

Engels, M.M.A., van der Flier, W.M., Stam, C.J., Hillebrand, A., Scheltens, Ph., van Straaten,
E.C.W., 2017. Alzheimer's disease: The state of the art in resting-state
magnetoencephalography. Clinical Neurophysiology 128, 1426–1437.
https://doi.org/10.1016/j.clinph.2017.05.012

Hyder, R., Jensen, M., Højlund, A., Kimppa, L., Bailey, C.J., Schaldemose, J.L., Kinnerup,
M.B., Østergaard, K., Shtyrov, Y., 2021. Functional connectivity of spoken language
processing in early-stage Parkinson's disease: An MEG study. NeuroImage: Clinical 32,
102718. https://doi.org/10.1016/j.nicl.2021.102718

Stam, C.J., de Haan, W., Daffertshofer, A., Jones, B.F., Manshanden, I., van Cappellen van
Walsum, A.M., Montez, T., Verbunt, J.P.A., de Munck, J.C., van Dijk, B.W., Berendse,
H.W., Scheltens, P., 2009. Graph theoretical analysis of magnetoencephalographic functional
connectivity in Alzheimer's disease. Brain 132, 213–224.
https://doi.org/10.1093/brain/awn262

Stam, C.J., Reijneveld, J.C., 2007. Graph theoretical analysis of complex networks in the
brain. Nonlinear Biomedical Physics 1. https://doi.org/10.1186/1753-4631-1-3

van Dellen, E., Douw, L., Hillebrand, A., Ris-Hilgersom, I.H.M., Schoonheim, M.M.,

Baayen, J.C., De Witt Hamer, P.C., Velis, D.N., Klein, M., Heimans, J.J., Stam, C.J.,

Reijneveld, J.C., 2012. MEG Network Differences between Low- and High-Grade Glioma

Related to Epilepsy and Cognition. PLoS ONE 7, e50122.

https://doi.org/10.1371/journal.pone.0050122

Wang, H., Douw, L., Hernández, J.M., Reijneveld, J.C., Stam, C.J., Van Mieghem, P., 2010.

Effect of tumor resection on the characteristics of functional brain networks. Phys. Rev. E 82,

021924. https://doi.org/10.1103/PhysRevE.82.021924

Guggisberg, A.G., Honma, S.M., Findlay, A.M., Dalal, S.S., Kirsch, H.E., Berger, M.S.,

Nagarajan, S.S., 2008. Mapping functional connectivity in patients with brain lesions. Ann

Neurol. 63, 193–203. https://doi.org/10.1002/ana.21224

Heimans, J.J., Reijneveld, J.C., 2012. Factors affecting the cerebral network in brain tumor

patients. J Neurooncol 108, 231–237. https://doi.org/10.1007/s11060-012-0814-7

Hillebrand, A., Stam, C.J., 2019. Recent Developments in MEG Network Analysis, in: Supek,

S., Aine, C.J. (Eds.), Magnetoencephalography. Springer International Publishing, Cham, pp.

631–645. https://doi.org/10.1007/978-3-030-00087-5_12

Huang, Q., Zhang, R., Hu, X., Ding, S., Qian, J., Lei, T., Cao, X., Tao, L., Qian, Z., Liu, H.,

2014. Disturbed Small-World Networks and Neurocognitive Function in Frontal Lobe Low-

Grade Glioma Patients. PLoS ONE 9, e94095. https://doi.org/10.1371/journal.pone.0094095

Hyder, R., Jensen, M., Højlund, A., Kimppa, L., Bailey, C.J., Schaldemose, J.L., Kinnerup,

M.B., Østergaard, K., Shtyrov, Y., 2021. Functional connectivity of spoken language

processing in early-stage Parkinson's disease: An MEG study. NeuroImage: Clinical 32,

102718. https://doi.org/10.1016/j.nicl.2021.102718

Li Hegner, Y., Marquetand, J., Elshahabi, A., Klamer, S., Lerche, H., Braun, C., Focke, N.K.,

2018. Increased Functional MEG Connectivity as a Hallmark of MRI-Negative Focal and

Generalized Epilepsy. Brain Topogr 31, 863–874. https://doi.org/10.1007/s10548-018-0649-4

Louis, D.N., Perry, A., Reifenberger, G., von Deimling, A., Figarella-Branger, D., Cavenee,

W.K., Ohgaki, H., Wiestler, O.D., Kleihues, P., Ellison, D.W., 2016. The 2016 World Health

Organization Classification of Tumors of the Central Nervous System: a summary. Acta

Neuropathologica 131, 803–820. https://doi.org/10.1007/s00401-016-1545-1

Na, S., Li, L., Crosson, B., Dotson, V., MacDonald, T.J., Mao, H., King, T.Z., 2018. White

matter network topology relates to cognitive flexibility and cumulative neurological risk in

adult survivors of pediatric brain tumors. NeuroImage: Clinical 20, 485–497.

https://doi.org/10.1016/j.nicl.2018.08.015

Nayak, L., Lee, E.Q., Wen, P.Y., 2012. Epidemiology of Brain Metastases. Curr Oncol Rep

14, 48–54. https://doi.org/10.1007/s11912-011-0203-y

Nolte, G., 2003. The magnetic lead field theorem in the quasi-static approximation and its use

for magnetoencephalography forward calculation in realistic volume conductors. Phys. Med.

Biol. 48, 3637–3652. https://doi.org/10.1088/0031-9155/48/22/002

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., 2011. FieldTrip: Open Source Software

for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data.

Computational Intelligence and Neuroscience 2011, 1–9. https://doi.org/10.1155/2011/156869

Rampp, S., Rössler, K., Hamer, H., Illek, M., Buchfelder, M., Doerfler, A., Pieper, T.,
Hartlieb, T., Kudernatsch, M., Koelble, K., Peixoto-Santos, J.E., Blümcke, I., Coras, R., 2021.
Dysmorphic neurons as cellular source for phase-amplitude coupling in Focal Cortical
Dysplasia Type II. Clinical Neurophysiology 132, 782–792.
https://doi.org/10.1016/j.clinph.2021.01.004

Reijneveld, J.C., Ponten, S.C., Berendse, H.W., Stam, C.J., 2007. The application of graph
theoretical analysis to complex networks in the brain. Clinical Neurophysiology 118, 2317–
2331. https://doi.org/10.1016/j.clinph.2007.08.010

Şahin, M., Aybek, E., 2019. Jamovi: An Easy to Use Statistical Software for the Social
Scientists. International Journal of Assessment Tools in Education 670–692.
https://doi.org/10.21449/ijate.661803

Semmel, E.S., Quadri, T.R., King, T.Z., 2021. Graph Theoretical Analysis of Brain Network
Characteristics in Brain Tumor Patients: A Systematic Review. Neuropsychol Rev.
https://doi.org/10.1007/s11065-021-09512-5

Stam, C.J., 2010. Characterization of anatomical and functional connectivity in the brain: A
complex networks perspective. International Journal of Psychophysiology 77, 186–194.
https://doi.org/10.1016/j.ijpsycho.2010.06.024

Stam, Cornelis Jan, 2010. Emergence of modular structure in a large-scale brain network with
interactions between dynamics and connectivity. Front. Comput. Neurosci. 4.
https://doi.org/10.3389/fncom.2010.00133

Stam, C.J., de Haan, W., Daffertshofer, A., Jones, B.F., Manshanden, I., van Cappellen van
Walsum, A.M., Montez, T., Verbunt, J.P.A., de Munck, J.C., van Dijk, B.W., Berendse,
H.W., Scheltens, P., 2009. Graph theoretical analysis of magnetoencephalographic functional
connectivity in Alzheimer's disease. Brain 132, 213–224.
https://doi.org/10.1093/brain/awn262

Stam, C.J., Nolte, G., Daffertshofer, A., 2007. Phase lag index: Assessment of functional
connectivity from multi channel EEG and MEG with diminished bias from common sources.
Hum. Brain Mapp. 28, 1178–1193. https://doi.org/10.1002/hbm.20346

Stam, C.J., van Straaten, E.C.W., 2012. The organization of physiological brain networks.
Clinical Neurophysiology 123, 1067–1087. https://doi.org/10.1016/j.clinph.2012.01.011

Stufflebeam, S.M., 2011. Clinical Magnetoencephalography for Neurosurgery. Neurosurgery
Clinics of North America 22, 153–167. https://doi.org/10.1016/j.nec.2010.11.006

Tadel, F., Baillet, S., Mosher, J.C., Pantazis, D., Leahy, R.M., 2011. Brainstorm: A User-
Friendly Application for MEG/EEG Analysis. Computational Intelligence and Neuroscience
2011, 1–13. https://doi.org/10.1155/2011/879716

Taulu, S., Simola, J., 2006. Spatiotemporal signal space separation method for rejecting
nearby interference in MEG measurements. Phys. Med. Biol. 51, 1759–1768.
https://doi.org/10.1088/0031-9155/51/7/008

Uusitalo, M.A., Ilmoniemi, R.J., 1997. Signal-space projection method for separating MEG or

EEG into components. Med. Biol. Eng. Comput. 35, 135–140.
https://doi.org/10.1007/BF02534144

van Dellen, E., Douw, L., Hillebrand, A., Ris-Hilgersom, I.H.M., Schoonheim, M.M.,
Baayen, J.C., De Witt Hamer, P.C., Velis, D.N., Klein, M., Heimans, J.J., Stam, C.J.,
Reijneveld, J.C., 2012. MEG Network Differences between Low- and High-Grade Glioma
Related to Epilepsy and Cognition. PLoS ONE 7, e50122.
https://doi.org/10.1371/journal.pone.0050122

van Nieuwenhuizen, D., Douw, L., Klein, M., Peerdeman, S.M., Heimans, J.J., Reijneveld,
J.C., Stam, C.J., Hillebrand, A., 2018. Cognitive functioning and functional brain networks in
postoperative WHO grade I meningioma patients. J Neurooncol 140, 605–613.
https://doi.org/10.1007/s11060-018-2987-1

Vogel, S., Kaltenhäuser, M., Kim, C., Müller-Voggel, N., Rössler, K., Dörfler, A., Schwab,
S., Hamer, H., Buchfelder, M., Rampp, S., 2021. MEG Node Degree Differences in Patients
with Focal Epilepsy vs. Controls—Influence of Experimental Conditions. Brain Sciences 11,
1590. https://doi.org/10.3390/brainsci11121590

Wang, H., Douw, L., Hernández, J.M., Reijneveld, J.C., Stam, C.J., Van Mieghem, P., 2010a.
Effect of tumor resection on the characteristics of functional brain networks. Phys. Rev. E 82,
021924. https://doi.org/10.1103/PhysRevE.82.021924

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund,
G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K.,
Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo,
K., Yutani, H., 2019. Welcome to the Tidyverse. JOSS 4, 1686.
https://doi.org/10.21105/joss.01686

Yang, J., Kudulaiti, N., Chen, Z., Gao, L., Hameed, N.U.F., Feng, R., Lu, S., 2022. Within
and Beyond the Visual Cortex: Brain Tumors Induce Highly Sensitive Plasticity of Visual
Processing in Whole-Brain Neural Functional Networks. Cerebral Cortex bhab492.
https://doi.org/10.1093/cercor/bhab492