Erp and Rev adhesins of the Lyme disease spirochete's ubiquitous cp32

prophages assist the bacterium during vertebrate infection

Brian Stevenson[1,2] and Catherine A. Brissette[3]

1. Department of Microbiology, Immunology, and Molecular Genetics, University of

Kentucky, Lexington, KY, USA

2. Department of Entomology, University of Kentucky, Lexington, KY, USA

3. Department of Biomedical Sciences, University of North Dakota, Grand Forks, ND,

USA

* Addresses for correspondence:

Brian Stevenson, Department of Microbiology, Immunology, and Molecular Genetics,

University of Kentucky College of Medicine, Lexington, Kentucky, 40536-0298,

Phone: 1+859-257-9358, Email: brian.stevenson@uky.edu

Catherine A. Brissette, Department of Biomedical Sciences, School of Medicine and

Health Sciences, University of North Dakota, Grand Forks, North Dakota, 58203-9061,

Phone: 1+701-777-6412, Email: catherine.brissette@und.edu

<u>Abstract</u>

Almost all spirochetes in the genus *Borrelia* (sensu lato) naturally contain multiple

variants of closely related prophages. In the Lyme disease borreliae, these prophages are

maintained as circular episomes that are called cp32s (<u>c</u>ircular <u>p</u>lasmid <u>32</u>kb). The cp32s

of Lyme agents are particularly unique in that they encode two distinct families of

lipoproteins, Erp and Rev, that are expressed on the bacteria's outer surface during

infection of vertebrate hosts. All identified functions of those outer surface proteins

involve interactions between the spirochetes and host molecules: Erp proteins bind

plasmin(ogen), laminin, glycosaminoglycans, and/or components of complement, and

Rev proteins bind fibronectin. Thus, cp32 prophages provide their bacterial hosts with

surface proteins that can enhance infection processes, thereby facilitating their own

survival. Horizontal transfer via bacteriophage particles increases spread of beneficial

alleles and creates diversity among Erp and Rev proteins.

Introduction

Members of the spirochete genus *Borrelia* (sensu lato) persist through infectious

cycles between blood-feeding arthropods and vertebrates, and include agents of

infections that plague humans, domestic animals, and wildlife (1, 2). These include Lyme

disease, caused by *Borrelia burgdorferi* sensu lato, and tick-borne relapsing fever, caused

by *B. hermsii, B. turicatae,* and others. Although a proposal has been raised to divide the

genus into *Borrelia* for relapsing fever agents and *Borreliella* for Lyme disease agents, that

idea is controversial (3-8). For the purpose of this review, we use the unified genus name

Borrelia, in part because of the shared property among Lyme disease and relapsing fever

spirochetes of near-universal natural colonization with multiple, closely-related cp32

prophages (9-13) (Figs. 1 and 2). Of particular significance to this review, the cp32s of

both Lyme disease and relapsing fever borreliae carry *mlp* genes, while *erp* and *rev* genes

appear to be restricted to the Lyme disease borreliae. The significance of those genes will

be discussed further below. For simplicity, we also use the name *Borrelia burgdorferi* to

encompass all genospecies of Lyme disease spirochetes, *B. burgdorferi* sensu lato.

Borrelia species all have multipartite genomes, consisting of a ca. one megabase

main chromosome and numerous smaller replicons that are generally referred to as

"plasmids" (14-16). Some borrelial "plasmids" carry essential genes, and might be better

considered to be additional chromosomes (15-20). Another uncommon feature of

borrelial genomes is a preponderance of linear DNAs, with the main chromosome and

many of the smaller replicons being linear elements with closed hairpin telomeres (15, 16,

18, 21-23). Lyme disease spirochetes and many other borreliae also maintain circular

DNA replicons (24-26).

During the mid-1990s, four research groups published evidence of multiple,

homologous loci in *B. burgdorferi* (26-30). Our group pursued the genomic locations of

those loci and discovered that each related locus is carried on a different circular replicon

of approximately 32 kb in size, and that a single bacterial cell can contain numerous

different variants of those replicons (26). Based on the sizes and circular nature of the

replicons, they were designated "cp32"s (circular plasmid 32kb). The majority of

nucleotides that comprise *B. burgdorferi* cp32s are identical to each other, differing

significantly at only three loci: one that is responsible for plasmid maintenance and two

that encode outer-surface lipoproteins (Fig. 1) (12). The highly repetitive nature of cp32s

initially prevented assembly of their sequences, such that the initial, 1997 report of the

genome of *B. burgdorferi* type strain B31 did not include the native cp32 plasmids (15).

Genomics analysts subsequently used the cp32s as learning tools to refine sequence

assembly algorithms, and the entire genome sequence of strain B31, including all cp32s

and other repetitive sequences, was finally published in 2000 (16).

The different cp32s within a single bacterium appear to be compatible with each

other due to their encoding two maintenance proteins, ParA and BpaB, that are distinct

for each cp32 within a cell (30-35). Yet, *parA-bpaB* locus sequences are conserved between

strains (13, 31, 34). Extensive characterization of cp32 sequences from Lyme disease

spirochetes that were collected around the world revealed twelve apparent clades of cp32

parA-bpaB pairings (13, 34, 36). Additionally, some strains contain small plasmids of

approximately 9 kb that were derived from cp32s and which carry only a *bpaB* allele (15,

16, 37). The absence of a *parA* gene from those cp9 plasmids raises intriguing questions

about plasmid maintenance in *Borrelia* species.

The conserved size and sequence of *B. burgdorferi* cp32s, along with genes for

proteins that resemble bacteriophage portals and holins, led to an early hypothesis that

they might be prophage genomes (9, 38). Cultures of *B. burgdorferi* have been observed to

clear spontaneously, or when exposed to 1-methyl-3-nitroso-nitroguanadine (MNNG),

with production of bacteriophage particles that were named ϕBB1 (10, 39). Molecular

analyses of those particles revealed that they contained cp32 DNA, and that they were

capable of transducing nucleic acids between *B. burgdorferi* cells (40, 41). A subsequent

study revealed that MNNG can induce transcription of a long, multigene cp32 operon

that presumably encodes bacteriophage structural proteins (42). Intriguingly, even

though MNNG is a mutagen, *B. burgdorferi* lacks an SOS system (15), so the means

through which this chemical induces bacteriophage production is not yet known.

B. burgdorferi cp32-encoded outer-surface Erp lipoproteins.

Two antigenic outer-surface proteins of *B. burgdorferi* strain N40 were reported in

1994 and named OspE and OspF (<u>o</u>uter <u>s</u>urface <u>p</u>roteins <u>E</u> and <u>F</u>) (43). Shortly afterward,

we found that *B. burgdorferi* N40 expresses different levels of those two proteins under

different culture conditions (44). Screening of a lambda library of *B. burgdorferi* type strain

B31 DNA revealed several distinct loci in that strain that were closely related to, but

distinct from, the strain N40 *ospE-ospF* locus (26). As none of the strain B31 loci could be

definitively called "*ospE-ospF*", they were designated "*erp*" (Osp<u>E</u>F-<u>r</u>elated <u>p</u>rotein).

Further analyses revealed that various clonal cultures of *B. burgdorferi* B31 contain up to

9 different *erp* loci (9, 16) (Table 1). Strain N40 contains 6 *erp* loci, including the initially-

described *ospE-ospF* (13, 34, 36, 43, 45, 46) (Table 1). Researchers who investigated other

strains discovered additional *erp* genes, and gave them a variety of names such as *p21,*

bbk2.10, bbk2.11, pG and *elp* (28, 45, 47-49) (Table 1). To avoid unnecessary confusion, we

use the names originally applied to *erp* genes and Erp proteins, and urge other authors to

also retain names that have been in common use for the past several decades. Structures

have been determined for some Erp proteins, primarily alleles that occur in the B31 type

strain (50-53).

109   Considering *erp* genes and their proteins as a unified group is supported by their

all occupying an allelic location on cp32s, their conserved promoter and operator DNA

sequences, simultaneous co-expression in culture and during infection processes,

possession of conserved leader polypeptides and initial residues of their mature

lipoproteins, and localization of the mature lipoproteins on the outside of the borrelial

outer membrane (12, 54). Evolutionary analyses indicated that *erp* genes arose from a

single sequence that encodes the leader polypeptide and the 5 amino acid sorting

sequence, while the remainder of the open reading frames descended from a large

number of independent origins that frequently undergo novel recombination events (34).

118   Dendrograms of known *erp* sequences show at least three branches (31, 34, 55),

which led to a suggestion that *erp* genes could be divided and given three different group

names: *ospE*, *ospF*, and *elp* (55). In addition to variations in nucleotide sequences, the

encoded proteins generally differ considerably in size, with OspE-types being small

(approximately 20 kDa), OspF-types being medium sized (approximately 25-30 kDa), and

Elp-types being large (approximately 40 kDa) (34). We do not find the OspE/OspF/Elp

nomenclature to be particularly helpful, since functions are not always conserved within

each group, and there are no consistent differences in gene expression or operon

structures. In addition, due to frequent natural recombination events having occurred

between *erp* genes, some extended nucleotide sequences are found in members of all three

groups (34). For example, some "*elp*" alleles share greater similarities with "*ospF*" genes

(as high as 34%) than they do with other "*elp*" genes (as low as 17%) (34). Recombination

events have also resulted in alleles that do not fit in the tripartite scheme (34). We

previously published extensive analyses of *erp* open reading frames, operon structures,

promoter sequences, and both intragenic and intergenic recombination, and refer readers

to that publication for deeper evolutionary discussions (34).

Erp lipoproteins are localized to the borrelial outer surface (43, 56, 57). The strain

B31 Erp proteins are all simultaneously co-expressed, consistent with their genes

possessing conserved promoter and operator sequences (58-60). Differences in gene or

protein expression have been reported for two other strains, but the reasons for those

apparent variations were not investigated (45, 57, 61). The mechanism by which *B.*

burgdorferi controls transcription of *erp* operons has been well studied, and shown to

involve interactions between the BpaB repressor, BpuR co-repressor, and EbfC

antirepressor proteins that bind to specific sites adjacent to *erp* promoters. There is

evidence that some *erp* operons may be affected by an alternative sigma factor, RpoS (62,

63). For an extensive review of the mechanisms by which *erp* transcription is regulated,

please see our recent review on that topic (64).

B. burgdorferi expresses little to none of any Erp protein during colonization of

unfed ticks (45, 59, 65). As the tick begins to feed on the blood of a vertebrate host, the

bacteria induce production of their entire repertoire of Erp proteins (59). Indirect

fluorescence analyses (IFAs) of skin samples at the sites of tick attachment revealed that

almost all transmitted *B. burgdorferi* produced detectable levels of every examined Erp

protein (59). Quantitative reverse transcription PCR (qRT-PCR) and promoter fusions

with *gfp* revealed that *erp* genes remain induced throughout persistent mammalian

infection (59, 66, 67). Levels of antibodies against Erp proteins, including IgM, remain

elevated during prolonged infection of mice (59, 66-68). The persistence of IgM antibodies

against Erp proteins may reflect *B. burgdorferi*'s ability to manipulate host immune

responses (69-72). That phenomenon may also explain how *B. burgdorferi* is able to persist

extracellularly despite having its outer surface covered with Erp, Rev, and other antigenic

proteins. IFAs of naive ticks feeding on infected mice showed that *B. burgdorferi* continue

to produce Erp proteins during acquisition by feeding ticks, which ceases soon after

detachment (59) (59).

Numerous functions have been identified for Erp proteins, all of which involve

interactions with vertebrate host tissues:

Several Erp proteins, including ErpA, ErpC, and ErpP of strain B31 and OspE of

strain N40, exhibit high affinities for complement factor H and the related proteins FHL-

1 and the FHR group (73-83). Binding factor H and FHL-1 to *B. burgdorferi* via Erp proteins

confers resistance to the alternative pathway of complement activation. The exact

functions of FHR proteins are unclear, but they also appear to be involved with

complement control (84). Erp proteins with different sequences show varying affinities

to factor H molecules of different potential vertebrate host species, and may thus play

roles in expanding the host range of Lyme disease spirochetes (85, 86).

The ErpB, ErpM and ErpQ proteins of strain B31 (confusingly called ElpB, ElpM

and ElpQ in the publications) were recently demonstrated to bind complement factors

C1s and C1r (87, 88). Adhesion of those host proteins onto the borrelial surface appears

to afford protection from the classical arm of complement.

The ErpX protein of *B. burgdorferi* B31 binds human laminin (89). This may help

the spirochete target and/or adhere to laminin-rich tissues. The laminin-binding domain

of ErpX was mapped to a central, disordered domain that is almost entirely comprised of

charged amino acids (89). Of note, other strain B31 Erp proteins with amino acid

sequences that are similar to ErpX (others in the "Elp" group: ErpB, ErpM and ErpQ,

Table 2) do not detectably bind human laminin (89).

The *B. burgdorferi* B31 ErpG and six similar proteins from other strains bind

heparan sulfate, and promote attachment to glial cells (90). A phage display library,

injected into mice and screened for potential *B. burgdorferi* adhesins, found clones of B31

ErpG, ErpK, and ErpL, and strain 297 Bbk2.10 in mouse heart, bladder, and tibiotarsal

joint tissues (91). Those four proteins share some sequence similarities (Table 2) (34).

The *B. burgdorferi* B31 ErpA, ErpC, and ErpP proteins bind host plasminogen and

its proteolytically-active form, plasmin (92). Plasmin bound to Erps and other surface

proteins facilitates *B. burgdorferi* migration through solid tissues in vitro, a characteristic

that is hypothesized to also enable dissemination during vertebrate infection (93-102). It

is notable that the Erp proteins that bind plasmin(ogen) also bind complement factor H

(see above).

Due to the surface localization and antigenicity of Erp proteins, there have been

several investigations into their use as protective vaccines or for serological diagnosis of

B. burgdorferi infection. However, vaccination of mice with the OspE, OspF, or P21

proteins of strain N40 did not provide substantial protection against infection by the same

strain (45, 65). Several recombinant Erp proteins have been examined for serological

diagnosis of human infection, with mixed results (103-106).

***B. burgdorferi* cp32-encoded outer-surface Rev and Mlp lipoproteins.**

The cp32 replicons of Lyme disease and relapsing fever borreliae carry a distinct

mono- or bicistronic *mlp* (<u>m</u>ulticopy <u>l</u>ipo<u>p</u>rotein) operon (Figs. 1 and 2) (11, 16, 29). In the

majority of Lyme disease spirochete cp32s, the *mlp* locus is preceded by a distinct gene,

bdr, which has its own promoter (16, 29) (Fig. 1). The characterized cp32 of the relapsing

fever spirochete *B. hermsii* contains a single *mlp* gene, without an adjacent *bdr* (11).

Curiously, all cp32s of Lyme disease and relapsing fever borrelia carry a second *bdr* allele

adjacent to the episomal replication locus (9, 16, 30) (Fig. 1). Bdr proteins are produced

during vertebrate infection, but their function(s) have yet to be discovered (68, 107-115).

For unknown reasons, some *B. burgdorferi* cp32s do not contain a *bdr* adjacent to

the *mlp* locus, but instead carry a distinct gene named *revA* (so named because such genes

are arranged <u>rev</u>erse to *mlp*) (16, 29). The *revA* ORF and promoter sequences are very

different from those of *mlp* genes, implying that *mlp* and *revA* loci are genetically

unrelated (9, 16, 29). In the *B. burgdorferi* type strain, B31, cp32-1 and cp32-6 each contain

a *revA* gene adjacent to their *mlp*, while all other B31 cp32s contain only a *mlp* gene (16).

Mature RevA is a lipoprotein and localizes to the *B. burgdorferi* outer surface (116).

Additionally, some truncated cp32 derivatives, such as B31 cp9, carry a similar gene

named *revB* (15, 16, 116). The two *revA* orthologs of strain B31 encode an identical RevA

protein, while that strain's RevB protein shares only 28% amino acid identity with RevA.

Other strains encode RevA and RevB proteins with varying degrees of differences (29).

B. burgdorferi in unfed ticks do not produce detectable levels of *revA* mRNA, but

do produce elevated transcripts during mouse infection (117). Human patients and

experimentally infected mice produce antibodies against RevA (118-120). In culture, *revA*

transcription is induced by changes in temperature or pH that correspond to changing

conditions encountered by the bacteria during tick feeding and mammalian infection

(116). Despite its antigenicity and surface localization, vaccination with RevA did not

protect mice from *B. burgdorferi* infection (121).

B. burgdorferi B31 RevA binds the 70kDa N-terminal region of fibronectin via a

domain in the RevA N-terminus (117). This binding was not affected by salt or heparin,

suggesting the interaction of these two proteins involves more than ionic interactions.

The *B. burgdorferi* RevB protein also binds to fibronectin, despite having limited sequence

identity to RevA (117). Polyclonal antibodies against RevA did not block adherence of

fibronectin to wild type *B. burgdorferi*, although those antibodies did prevent fibronectin

adherence to *B. burgdorferi* that were deleted of another fibronectin-binding protein,

BBK32 (122, 123). Similarly, binding of wild-type *B. burgdorferi* to HUVECs or human

neuroglial cells was not blocked by anti-RevA monoclonal antibodies (124). Those results

underscore the functional redundancy of *B. burgdorferi* fibronectin-binding proteins (125).

RevA also binds mammalian laminin, although to a significantly lesser extent than to

fibronectin (117).

A *B. burgdorferi* mutant unable to produce RevA was found to be defective in

colonization of cardiac tissue (123). That mutant also caused increased pathology of

tibiotarsal joints and enhanced CCL2 production. Increased inflammation in the absence

of RevA protein suggests a role for that surface protein in immune evasion, perhaps by

masking a more immunogenic antigen.

Precise functions have not yet been ascribed to Mlp proteins. The cp32 prophages

of both Lyme disease and relapsing fever *Borrelia* spp. carry *mlp* genes, which suggests

that the proteins serve functions that are conserved among borreliae (9, 11, 16, 29, 126)

(Fig. 2). As with *erp* and *rev* genes, there is often considerable sequence diversity among

mlp genes within and between *B. burgdorferi* isolates (16, 126, 127). The patterns of

regulated expression and antigenicity of *B. burgdorferi* Mlp proteins imply that they serve

functions within vertebrate hosts (127-132). A recombinant Mlp protein was found to

adhere to human brain microvascular endothelial cells (133). The structure of the

MlpA/P28 protein that is encoded by B31 cp32-1 has been solved, and was noted to

contain a structural motif that is similar to one seen in a lipid-binding protein (16, 127,

134).

Horizontal transfer and recombination

Almost all examined Lyme disease spirochete isolates naturally contain 6-12

distinct cp32s, each with an *erp* and a *mlp* / *revA* locus. As noted above, relationships

between cp32s can be discriminated by their *parA-bpaB* maintenance loci. Using that as

reference, there is clear evidence of recombination having occurred between cp32s, and

exchange of cp32 DNA between bacteria (13, 31, 34, 36).

B. burgdorferi type strain B31 naturally carries three identical *erpAB* operons on

cp32-1, cp32-5, and cp32-8 (9, 26). In contrast, those cp32s of other strains possess distinct

erp operons (31, 32, 34). The cp32-1 and cp32-6 of strain B31 carry *revA* genes that are

nearly identical in sequence, and which encode identical RevA proteins, while related

plasmids of other strains have different genes at that locus (16, 117, 120). Two other

recombination events are also evident in strain B31: cp32-4 has an fragment of a cp32 *sbbP*

gene inserted into *erpH*, such that a complete lipoprotein cannot be produced, and cp32-

3 has an unrelated gene, *bapA*, inserted 3′ of *erpG* (9, 26). No other B31 cp32 carries a *bapA*

gene, although paralogous genes, named *eppA*, are found on some cp9 replicons (135-

137).

Despite cp32s having nearly identical sequences, and multiple cp32s residing

within individual bacteria, there are no known examples of recombination occurring

during cultivation. Recombination does not take place during mammalian infection, or,

at least, not among bacteria of the same strain (138). We hypothesize that genetic transfer

and recombination occur when *Borrelia* spirochetes are within ticks.

Additional evidence of horizontal transfer of cp32 DNAs in nature was obtained

from sequencing genomes of multiple isolates (34). *B. burgdorferi* strains B31 and BL206

contain almost identical repertoires of cp32s, the only differences being presence of a

cp32-1 in only B31 and a cp32-11 in only BL206. Intriguingly, B31 cp32-1 and BL206 cp32-

11 carry an identical *erpAB* locus. Two other strains, 297 and Sh-2-82, contain cp32

repertoires that are identical to each other, with the addition of a cp32-8 in only Sh-2-82.

The cp32-8 *erp* locus of Sh-2-82 is identical to the cp32-8 *erpAB* locus of strain B31,

implying that a cp32-8/φBB1 was exchanged between ancestors of those two strains.

Strain B31 was isolated from a tick on Shelter Island, New York, in 1981, and Sh-2-82 was

isolated from another tick on the same island in 1982 (139, 140). It is also notable that

BL206, which is a near clone of B31, was isolated several years later from the blood of a

Lyme disease patient in Westchester County, New York (141), and that 297, which is a

near clone of Sh-2-82, was isolated from the cerebrospinal fluid of another patient (142).

Transduction of cp32 DNA between different strains of *B. burgdorferi* has been

observed in culture (40). The mechanisms that induce φBB1 particles to form are not yet

known. As noted above, *B. burgdorferi* does not possess an SOS system (15). The timing

of bacteriophage production is also unknown. All *Borrelia* species are obligate parasites

of arthropods and vertebrates, so triggers for φBB1 would be limited to conditions

experienced within those hosts. Ticks can acquire multiple strains of *B. burgdorferi*, which

would reside together in the tick's midgut, leading us to propose that the midguts of

feeding ticks are likely locations for DNA exchange. Tick feeding is a time of rapid

replication for *B. burgdorferi*, which appears to be used by the spirochetes as an important

cue that transmission is imminent (64, 143). That would also be an appropriate time for

bacteria to exchange DNA and diversity their repertoires of Erp, Rev, and other surface

proteins that interact with vertebrate hosts.

At least two advantages appear to be conferred upon *B. burgdorferi* by exchanging

and recombining *erp* and *rev* genes. First, both groups of proteins are expressed early

during vertebrate infection, are located on the bacterial outer surface, and are antigenic.

Reservoir vertebrates are generally fed upon by numerous ticks, all of which might be

infected. Thus, Lyme disease spirochetes are under selective pressure to diversify

sequences of early, surface-borne antigens, to facilitate infection of animals that have

previously been exposed. A second advantage to Erp and Rev variability among

individual bacteria stems from their functions as adhesins, and from the fact that the ticks

that transmit Lyme borreliae may not be specific in their choices of hosts. For example, in

the northeastern USA, *Ixodes scapularis* ticks feed on birds, mice, and other mammals

(144). Since a *B. burgdorferi* cannot "know" in advance what species of vertebrate its tick

vector will feed upon, it would be advantageous for a bacterium to enter its new host

while bristling with a wide variety of adhesins that differ in relative affinities for ligands

of different hosts, with the likelihood that some Erp, Rev, and other adhesins will be

adequate for the bacteria's survival.

Exceptional *Borrelia*

To date, only two exceptions are known to the generalization that *Borrelia* species

naturally carry cp32s. The agent of avian spirochetosis, *Borrelia anserina*, is the type

species of the genus, and lacks cp32 DNA (11). Since Lyme disease and relapsing fever

Borrelia species carry cp32s, their absence from *B. anserina* suggests that either φBB1

entered in the genus after the split of *B. anserina* from the other species, *B. anserina* has a

means to exclude cp32s/φBB1, or cp32s encode properties that are beneficial to Lyme

disease and relapsing fever borreliae, but not to avian spirochetosis borreliae. The second

known example of a cp32-deficient *Borrelia* is strain Far04, a Lyme disease spirochete that

was isolated from a puffin that was collected on the Faroe Islands, Denmark (13, 145, 146).

The densely packed puffin nests on those islands are homes to *Ixodes uriae* ticks, which

preferentially feed on seabirds. Although Far04 is a single example, it is possible that *I.*

uriae ticks on the islands primarily feed on puffins, to the extent that some circulating

borreliae have lost the need for the variability of adhesin compositions afforded by cp32

prophages. Clearly, additional sampling and analyses are needed to test that hypothesis.

Future research directions.

Erp and Rev lipoproteins localize to the borrelial outer surface, and are known to

interact with a variety of vertebrate host proteins that protect against immune responses

and facilitate tissue invasion and colonization. Noting that Erp and Rev proteins were

discovered during the mid-1990s, and that adherence of complement C1r and C1s was

first reported in 2022, we consider it likely that other functions remain to be discovered.

Additional questions include: Why do the cp32s of Lyme disease spirochetes encode Erp

and Rev proteins, while those of relapsing fever spirochetes appear to lack those genes?

What are the functions of the Mlp outer surface proteins, which are encoded on cp32s of

both Lyme disease and relapsing fever borreliae? Why does *B. anserina*, the agent of avian

spirochetosis, lack cp32s? When do borreliae produce ɸBB1 bacteriophages during their

natural infectious cycles? There is genetic evidence of DNA transfer of other genes

encoding borrelial antigens, such as outer surface protein C (OspC) (147, 148) – might

that occur by generalized transduction via ɸBB1?

Concluding statements

The cp32 prophages / ɸBB1 bacteriophages have evidently co-evolved with many

species of *Borrelia*. They have acquired a number of different genes that confer benefits to

their bacterial hosts, which thereby enhance their own survival. These bacteriophages

have adopted a primarily lysogenic lifestyle, with a limited degree of bacteriophage

particle production and host cell lysis. The *parA* and *bpaB* maintenance genes have

diversified extensively, to the extent that large numbers of different cp32s can be

maintained in a single bacterial cell. This provides both extensive repertoires of *erp*, *rev*,

and other genes for horizontal exchange and enables the host bacteria to express wide

varieties of surface adhesins during vertebrate infection.

Acknowledgements

We thank the many colleagues from our laboratories and throughout the world

who have contributed to the studies described in this review. Research in our laboratories

is currently funded by NIH grants R01AI144126, 3R01AI144126-03S1, and R21AI147139

to Brian Stevenson and R01AI58304 to Catherine Brissette.

Figure 1. Schematic of the open reading frames of a representative *B. burgdorferi* cp32, the

cp32-1 of type strain B31. Arrowheads indicate direction of open reading frame

transcription. *B. burgdorferi* cp32s contain two loci that encode lipoproteins that are

expressed on the bacterial outer surface: a mono- or bicistronic locus that encodes Erp

proteins, and a separate locus that consists of either divergently-transcribed *revA* and *mlp*

genes, or a *mlp* operon adjacent to a *bdr* gene. B31 cp32-1 and cp32-6 each carry a *revA*

gene, while the other cp32s of that strain instead carry a *bdr* gene adjacent to their *mlp*

locus. Borreliae are apparently able to maintain numerous, different cp32s in a single cell

due to diversity among the *parA-bpaB* (<u>b</u>orrelial *par<u>B</u>*-analog) locus. Other open reading

frames show up to 100% conservation between different cp32s and *B. burgdorferi* strains,

shown as white. Two conserved genes, PF-50 and PF-57, are located adjacent to the *parA-*

bpaB loci of all cp32s and are necessary for replicon maintenance (149-151). The *parS*

segregation site appears to be associated with the 5′ end of *PF-57* (our unpublished results

and (149-151). Conserved genes *ssbP* and *nucP* encode a single-stranded DNA binding

protein and nuclease, respectively (35). Bacteriophage structural and assembly proteins

appear to be encoded from a single operon that extends from 3′ of the *erp* locus to the

holin-encoding genes (16, 38, 42, 152).

PF-57
PF-50
parA
bpaB
bdr
nucP
ssbP
}parS ?
mlpA
revA
holins
erpA
erpB
phage
morphogenesis?

Strain B31 cp32-1
30749 bp

Figure 2. Alignment of the open reading frames of *B. burgdorferi* B31 cp32-1 and a

composite cp32 of *B. hermsii* HS-1 that was assembled from sequenced contigs. Directions

of transcription are indicated by arrows. The *erp*, *revA* and *mlp* genes are indicated by

hatching. ORFs that are nearly identical between cp32s of *B. burgdorferi* and *B. hermsii* are

indicated by white boxes. Note that the plasmid replication genes, including *parA* and

bpaB, are located in reversed directions on the two different cp32 types. Adapted from

(11).

Table 1: The cp32 DNA elements and associated *erp* loci that naturally occur in the three

best-characterized Lyme disease spirochete isolates: *B. burgdorferi* strains B31, N40, and

297. Plasmid nomenclature is standardized according to segregation locus type, such that

all plasmids designated "cp32-1" contain a similar segregation locus, etc. All known cp32

segregation loci fall into 12 clades. The first *B. burgdorferi* strain to be completely

characterized, B31, contains 2 distinct cp32 prophages that possess identical maintenance

loci, cp32-2 and cp32-7, and define segregation locus type 32-2/7. As a result, the

numbering scheme for the 12 groups goes up to cp32-13. The published names of *erp*

genes are listed. A variety of naming schemes were applied to these genes by their

discoverers, particularly for strain 297. Additional *erp* genes of strain N40 that were

discovered by our group were given allele numbers only (e.g. *erp22, erp23*, etc.). Other

strain's *erp* genes have been designated either by an allele number (with a different

number for each gene) or by genome sequencing ORF numbers, which are too numerous

to list here but examples are described in (34).

cp32 segregation locus type	*B. burgdorferi* strain		
	B31 [a]	N40 [d]	297 [e]
cp32-1	*erpA1, erpB1* (or *erpB2*) [b]		*ospE1, elpB1-1*
cp32-2/7	*erpC, erpD* *erpL, erpM* [c]	*ospE, ospF*	*elpA2*
cp32-3	*erpG*		*ospF*

cp32-4	*erpH, erpY*	*erp23, erp24*	*elpA1*
cp32-5	*erpA5, erpB5* (previously *erpI, erpJ*)	*erp25*	*ospE5, elpB1-5*
cp32-6	*erpK*		*bbk2.10*
cp32-8	*erpA8, erpB8* (previously *erpN, erpO*)		
cp32-9	*erpP, erpQ* [c]	*p21, erp22*	*p21, elpB2*
cp32-10	*erpX*	*erp26*	
cp32-11			*bbk2.11*
cp32-12		*erp27*	*ospE12, elpB1-12*
cp32-13			

a. The *erp* loci of strain B31 were first identified and named by the authors and their

research colleagues (9, 16, 26). Strain B31 was one of the first organisms to have its

genome completely sequenced, but the sequenced culture lacks cp32-2 and cp32-5, so the

complete sequences of those prophages remains uncertain. Prophages cp32-1, cp32-5, and

cp32-8 carry identical *erp* loci, which are distinguished by allele numbers. The *erp* loci of

cp32-1, cp32-5, and cp32-8 encode identical proteins, which are collectively called ErpA

and ErpB. At some time in the past, the B31 cp32-10 underwent integration into an

unrelated linear replicon, which created an approximately 56kb linear plasmid that is

named lp56.

b. Two alleles of *erpB* have been identified in different cultures of strain B31. The *erp* locus

that was first identified in a high-passage, non-infectious culture contains a premature

stop codon within the *erpB* gene, and thus encodes a truncated protein (26). Subsequent

analyses of infectious cultures of B31 revealed a full-length *erpB* open reading frame,

which was designated allele *erpB*2 (9, 16).

c. Two recent publications designated ErpB as "ElpB", ErpM as "ElpM" and ErpQ as

"ElpQ" (87, 88).

d. Two *erp* loci of strain N40, *ospEF* and *p21*, were named by Fikrig and colleagues (43,

45). The remainder of the N40 genes were discovered and named by the authors and their

coworkers (34). The cp32-2/7 prophage of strain N40 is naturally truncated to an

approximately 18kb circular replicon, cp18 (46).

e. The *erp* loci of strain 297 were discovered and named by Radolf and colleagues (28, 32,

55). Our research group later found that the cp32-1, cp32-5, and cp32-12 prophages each

contain an identical *erp* locus, *ospE-elpB1*, which we differentiated by additional allele

numbers (34). The cp32-2/7 and cp32-9 plasmids of strain 297 are naturally truncated,

and have been named cp18-a and cp18-2, respectively (32).

Table 2. The Erp proteins of the three best-characterized *B. burgdorferi* strains, B31, N40,

and 297, sorted according to the tripartite OspE/OspF/Elp scheme. Please note that even

though some proteins of strain 297 proteins were assigned the same names as N40

proteins, those proteins are distinct from each other. That is, N40 OspE is not identical to

297 OspE, N40 P21 is not identical to 297 P21, and N40 OspF is not identical to 297 OspF.

The strain 297 nomenclature is further complicated in that the 297 OspE protein is more

similar to the N40 P21 protein than to the N40 OspE protein, and the 297 P21 protein is

more similar to the N40 OspE protein (34).

	"OspE" "small" (69% average pairwise identity) [a]	"OspF" "medium" (44% average pairwise identity) [a]	"Elp" "large" (40% average pairwise identity) [a]
B. burgdorferi B31	ErpA ErpC ErpH ErpP	ErpG ErpL ErpK ErpY	ErpB ErpD ErpM ErpQ ErpX
B. burgdorferi N40	OspE P21	OspF Erp23 Erp25 Erp27	Erp22 Erp24 Erp26
B. burgdorferi 297	OspE P21	OspF Bbk2.10 Bbk2.11	ElpA1 ElpA2 ElpB1

			ElpB2

a. Average pairwise identities of genes and proteins of 21 analyzed Lyme disease *Borrelia* strains (34). That study found that some alleles that fall into the OspF group due to their sizes share greater sequence identity with members of the Elp group than they do with other members of the OspF group, and vice versa. Those similarities and differences appear to be due to frequent recombination between alleles of different groups.

References

1.   Trevisan G, Cinco M, Trevisini S, di Meo N, K. C, Ruscio M, Forgione P, Bonin S.
2021. Borreliae Part 1: *Borrelia* Lyme group and echidna-reptile group. Biology
10:1036.

2.   Trevisan G, Cinco M, Trevisini S, di Meo N, Ruscio M, Forgione P, Bonin S. 2021.
Borreliae Part 2: *Borrelia* relapsing fever group and unclassified *Borrelia* Biology
10:1117.

3.   Adeolu M, Gupta RS. 2014. A phylogenomic and molecular marker based proposal
for the division of the genus *Borrelia* into two genera: the emended genus *Borrelia*
containing only the members of the relapsing fever *Borrelia*, and the genus
Borreliella gen. nov. containing the members of the Lyme disease *Borrelia* (*Borrelia*
burgdorferi sensu lato complex). Antonie Van Leeuwenhoek 105:1049-1072.

4.   Margos G, Marosevic D, Cutler S, Derdakova M, Diuk-Wasser M, Emler S, Fish D,
Gray J, Hunfeld KP, Jaulhac B, Kahl O, S. K, Kraiczy P, Lane RS, Lienhard R,
Lindgren PE, Ogden N, Ornstein K, Rupprecht T, Schwartz I, Sing A, Straubinger
RK, Strle F, Voordouw M, Rizzoli A, Stevenson B, Fingerle V. 2016. There is
inadequate evidence to support the division of the genus *Borrelia*. Int J Syst Evol
Microbiol:doi: 10.1099/ijsem.0.00171.

5.   Margos G, Gofton A, Wibberg D, Dangel A, Marosevic D, Loh S-M, Oskam C,
Fingerle V. 2018. The genus *Borrelia* reloaded. PLoS ONE 13:e0208432.

6.   Gupta RS. 2019. Distinction between *Borrelia* and *Borreliella* is more robustly
supported by molecular and phenotypic characteristics than all other
neighbouring prokaryotic genera: Response to Margos' et al. "The genus Borrelia
reloaded" (PLoS ONE 13(12): e0208432). PLoS ONE 14:e0221397.

7.   Stevenson B, Fingerle V, Wormser G, Margos G. 2019. Public health and patient
safety concerns merit retention of Lyme borreliosis-associated spirochetes within
the genus *Borrelia*, and rejection of the genus novum *Borreliella*. Ticks Tick Borne
Dis 10:1-4.

8.   Margos G, Castillo-Ramirez S, Cutler S, Dessau RB, Eikeland R, Estrada-Peña A,
Gofton A, Graña-Miraglia L, Hunfeld KP, Krause A, Lienhard R, Lindgren PE,
Oskam C, Rudolf I, Schwartz I, Sing A, Stevenson B, Wormser GP, Fingerle V.
2020. Rejection of the name *Borreliella* and all proposed species comb. nov. placed
therein. Int J Syst Evol Microbiol 70:3577-3581.

9.   Casjens S, van Vugt R, Tilly K, Rosa PA, Stevenson B. 1997. Homology throughout
the multiple 32-kilobase circular plasmids present in Lyme disease spirochetes. J
Bacteriol 179:217-227.

10. Eggers CH, Samuels DS. 1999. Molecular evidence for a new bacteriophage of
Borrelia burgdorferi. J Bacteriol 181:7308-7313.

11. Stevenson B, Porcella SF, Oie KL, Fitzpatrick CA, Raffel SJ, Lubke L, Schrumpf ME,
Schwan TG. 2000. The relapsing fever spirochete *Borrelia hermsii* contains multiple,
antigen-encoding circular plasmids that are homologous to the cp32 plasmids of
Lyme disease spirochetes. Infect Immun 68:3900-3908.

12. Stevenson B, Zückert WR, Akins DR. 2001. Repetition, conservation, and variation:
the multiple cp32 plasmids of *Borrelia* species, p 87-100. *In* Saier MH, García-Lara
J (ed), The Spirochetes: Molecular and Cellular Biology. Horizon Press, Oxford.

13. Casjens SR, Gilcrease EB, Vujadinovic M, Mongodin EF, Luft BJ, Schutzer SE,
Fraser CM, Qiu WG. 2017. Plasmid diversity and phylogenetic consistency in the
Lyme disease agent *Borrelia burgdorferi*. BMC Genomics 18:165.

14. Barbour AG. 1988. Plasmid analysis of *Borrelia burgdorferi*, the Lyme disease agent.
J Clin Microbiol 26:475-478.

15. Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O,
Ketchum KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb J-F,
Fleischmann RD, Richardson D, Peterson J, Kerlavage AR, Quackenbush J,
Salzberg S, Hanson M, van Vugt R, Palmer N, Adams MD, Gocayne J, Weidmann
J, Utterback T, Watthey L, McDonald L, Artiach P, Bowman C, Garland S, Fujii C,
Cotton MD, Horst K, Roberts K, Hatch B, Smith HO, Venter JC. 1997. Genomic
sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*. Nature 390:580-586.

16. Casjens S, Palmer N, van Vugt R, Huang WM, Stevenson B, Rosa P, Lathigra R,
Sutton G, Peterson J, Dodson RJ, Haft D, Hickey E, Gwinn M, White O, Fraser C.
2000. A bacterial genome in flux: the twelve linear and nine circular
extrachromosomal DNAs of an infectious isolate of the Lyme disease spirochete
Borrelia burgdorferi. Mol Microbiol 35:490-516.

17. Purser JE, Norris SJ. 2000. Correlation between plasmid content and infectivity in
Borrelia burgdorferi. Proc Natl Acad Sci USA 97:13865-13870.

18. Kobryn K, Chaconas G. 2002. ResT, a telomere resolvase encoded by the Lyme
disease spirochete. Mol Cell 9:195-201.

19. Purser JE, Lawrenz MB, Caimano MJ, Howell JK, Radolf JD, Norris SJ. 2003. A
plasmid-encoded nicotinamidase (PncA) is essential for infectivity of *Borrelia*
burgdorferi in a mammalian host. Mol Microbiol 48:753-764.

20. Byram R, Stewart PE, Rosa P. 2004. The essential nature of the ubiquitous 26-
kilobase circular replicon of *Borrelia burgdorferi*. J Bacteriol 186:3561-3569.

21. Barbour AG, Garon CF. 1987. Linear plasmids of the bacterium *Borrelia burgdorferi*
have covalently closed ends. Science 237:409-411.

22. Ferdows MS, Barbour AG. 1989. Megabase-sized linear DNA in the bacterium
Borrelia burgdorferi, the Lyme disease agent. Proc Natl Acad Sci USA 86:5969-5973.

23.   Casjens S, Huang WM. 1993. Linear chromosomal physical and genetic map of
533   *Borrelia burgdorferi*, the Lyme disease agent. Mol Microbiol 8:967-980.

534   24.   Simpson WJ, Garon CF, Schwan TG. 1990. Analysis of supercoiled circular
535   plasmids in infectious and non-infectious *Borrelia burgdorferi*. Microb Pathog 8:109-
536   118.

537   25.   Marconi RT, Samuels DS, Garon CF. 1993. Transcriptional analyses and mapping
538   of the *ospC* gene in Lyme disease spirochetes. J Bacteriol 175:926-932.

539   26.   Stevenson B, Tilly K, Rosa PA. 1996. A family of genes located on four separate 32-
540   kilobase circular plasmids in *Borrelia burgdorferi* B31. J Bacteriol 178:3508-3516.

541   27.   Zückert WR, Filipuzzi-Jenny E, Meister-Turner J, Stålhammar-Carlemalm M,
542   Meyer J. 1994. Repeated DNA sequences on circular and linear plasmids of *Borrelia*
543   *burgdorferi* sensu lato, p 253-260. *In* Axford JS, Rees DHE (ed), Lyme Borreliosis.
544   Plenum Press, New York.

545   28.   Akins DR, Porcella SF, Popova TG, Shevchenko D, Baker SI, Li M, Norgard MV,
546   Radolf JD. 1995. Evidence for in vivo but not in vitro expression of a *Borrelia*
547   *burgdorferi* outer surface protein F (OspF) homologue. Mol Microbiol 18:507-520.

548   29.   Porcella SF, Popova TG, Akins DR, Li M, Radolf JD, Norgard MV. 1996. *Borrelia*
549   *burgdorferi* supercoiled plasmids encode multi-copy tandem open reading frames
550   and a lipoprotein gene family. J Bacteriol 178:3293-3307.

551   30.   Zückert WR, Meyer J. 1996. Circular and linear plasmids of Lyme disease
552   spirochetes have extensive homology: characterization of a repeated DNA
553   element. J Bacteriol 178:2287-2298.

554   31.   Stevenson B, Casjens S, Rosa P. 1998. Evidence of past recombination events
555   among the genes encoding the Erp antigens of *Borrelia burgdorferi*. Microbiology
556   144:1869-1879.

557   32.   Caimano MJ, Yang X, Popova TG, Clawson ML, Akins DR, Norgard MV, Radolf
558   JD. 2000. Molecular and evolutionary characterization of the cp32/18 family of
559   supercoiled plasmids in *Borrelia burgdorferi* 297. Infect Immun 68:1574-1586.

560   33.   Burns LH, Adams CA, Riley SP, Jutras BL, Bowman A, Chenail AM, Cooley AE,
561   Haselhorst LA, Moore AM, Babb K, Fried MG, Stevenson B. 2010. BpaB, a novel
562   protein encoded by the Lyme disease spirochete's cp32 prophages, binds to *erp*
563   Operator 2 DNA. Nucleic Acids Res 38:5443-5455.

564   34.   Brisson D, Zhou W, Jutras BL, Casjens S, Stevenson B. 2013. Distribution of cp32
565   prophages among Lyme disease-causing spirochetes and natural diversity of their
566   lipoprotein-encoding *erp* loci. Appl Env Microbiol 79:4115-4128.

567   35.   Chenail AM, Jutras BL, Adams CA, Burns LH, Bowman A, Verma A, Stevenson B.
568   2012. *Borrelia burgdorferi* cp32 BpaB modulates expression of the prophage NucP

nuclease and SsbP single-stranded DNA-binding protein. J Bacteriol 194:4570-
4578.

36. Stevenson B, Miller JC. 2003. Intra- and interbacterial genetic exchange of Lyme
disease spirochete *erp* genes generates sequence identity amidst diversity. J Mol
Evol 57:309-324.

37. Dunn JJ, Buchstein SR, Butler L-L, Fisenne S, Polin DS, Lade BN, Luft BJ. 1994.
Complete nucleotide sequence of a circular plasmid from the Lyme disease
spirochete, *Borrelia burgdorferi*. J Bacteriol 176:2706-2717.

38. Damman CJ, Eggers CH, Samuels DS, Oliver DB. 2000. Characterization of *Borrelia*
burgdorferi BlyA and BlyB proteins: a prophage-encoded holin-like system. J
Bacteriol 182:6791-6797.

39. Eggers CH. 2022. Phage-mediated genetic manipulation of the Lyme disease
spirochete *Borrelia burgdorferi* J Vis Exp 187:doi: 10.3791/64408.

40. Eggers CH, Kimmel BJ, Bono JL, Elias AF, Rosa P, Samuels DS. 2001. Transduction
by ØBB-1, a bacteriophage of *Borrelia burgdorferi*. J Bacteriol 183:4771-4778.

41. Eggers CH, Gray CM, Preisig AM, Glenn DM, Pereira J, Ayers RW, Alshahrani M,
Acabbo C, Becker MR, Bruenn KN, T. C, Jendras TM, Shepley AB, Moeller JT. 2016.
Phage-mediated horizontal gene transfer of both prophage and heterologous
DNA by φBB-1, a bacteriophage of *Borrelia burgdorferi*. Pathog Dis 74:pii: ftw107.

42. Zhang H, Marconi RT. 2005. Demonstration of cotranscription and 1-methyl-3-
nitroso-nitroguanidine induction of a 30-gene operon of *Borrelia burgdorferi*:
evidence that the 32-kilobase circular plasmids are prophages. J Bacteriol 187:7985-
7995.

43. Lam TT, Nguyen T-PK, Montgomery RR, Kantor FS, Fikrig E, Flavell RA. 1994.
Outer surface proteins E and F of *Borrelia burgdorferi*, the agent of Lyme disease.
Infect Immun 62:290-298.

44. Stevenson B, Schwan TG, Rosa PA. 1995. Temperature-related differential
expression of antigens in the Lyme disease spirochete, *Borrelia burgdorferi*. Infect
Immun 63:4535-4539.

45. Das S, Barthold SW, Stocker Giles S, Montgomery RR, Telford SR, Fikrig E. 1997.
Temporal pattern of *Borrelia burgdorferi p21* expression in ticks and the mammalian
host. J Clin Invest 99:987-995.

46. Stevenson B, Casjens S, van Vugt R, Porcella SF, Tilly K, Bono JL, Rosa P. 1997.
Characterization of cp18, a naturally truncated member of the cp32 family of
Borrelia burgdorferi plasmids. J Bacteriol 179:4285-4291.

47. Suk K, Das S, Sun W, Jwang B, Barthold SW, Flavell RA, Fikrig E. 1995. *Borrelia*
burgdorferi genes selectively expressed in the infected host. Proc Natl Acad Sci USA
92:4269-4273.

48. Wallich R, Brenner C, Kramer MD, Simon MM. 1995. Molecular cloning and
immunological characterization of a novel linear-plasmid-encoded gene, *pG*, of
Borrelia burgdorferi expressed only in vivo. Infect Immun 63:3327-3335.

49. Sung SY, Lavoie CP, Carlyon JA, Marconi RT. 1998. Genetic divergence and
evolutionary instability in *ospE*-related members of the upstream homology box
gene family in *Borrelia burgdorferi* sensu lato complex isolates. Infect Immun
66:4656-4668.

50. Bhattacharjee A, Oeemig JS, Kolodziejczyk R, Meri T, Kajander T, Lehtinen MJ,
Iwaï H, Jokiranta TS, Goldman A. 2013. Structural basis for complement evasion
by Lyme disease pathogen *Borrelia burgdorferi*. J Biol Chem 288:18685-18695.

51. Caesar JJ, Johnson S, Kraiczy P, Lea SM. 2013. ErpC, a member of the complement
regulator-acquiring family of surface proteins from *Borrelia burgdorferi*, possesses
an architecture previously unseen in this protein family. Acta Crystallogr Sect F
Struct Biol Cryst Commun 69:624-628.

52. Brangulis K, Petrovskis I, Kazaks A, Akopjana I, Tars K. 2015. Crystal structures
of the Erp protein family members ErpP and ErpC from *Borrelia burgdorferi* reveal
the reason for different affinities for complement regulator factor H. Biochim
Biophys Acta 1854:349-355.

53. Kolodziejczyk R, Mikula KM, Kotila T, Postis VLG, Jokiranta TS, Goldman A, Meri
T. 2017. Crystal structure of a tripartite complex between C3dg, C-terminal
domains of factor H and OspE of *Borrelia burgdorferi*. PLoS ONE 12:e0188127.

54. Stevenson B, Bykowski T, Cooley AE, Babb K, Miller JC, Woodman ME, von
Lackum K, Riley SP. 2006. The Lyme disease spirochete Erp lipoprotein family:
structure, function and regulation of expression, p 354-372. *In* Cabello FC, Godfrey
HP, Hulinska D (ed), Molecular Biology of Spirochetes. IOS Press, Amsterdam.

55. Akins DR, Caimano MJ, Yang X, Cerna F, Norgard MV, Radolf JD. 1999. Molecular
and evolutionary analysis of *Borrelia burgdorferi* 297 circular plasmid-encoded
lipoproteins with OspE- and OspF-like leader peptides. Infect Immun 67:1526-
1532.

56. El-Hage N, Babb K, Carroll JA, Lindstrom N, Fischer ER, Miller JC, Gilmore RD,
Mbow ML, Stevenson B. 2001. Surface exposure and protease insensitivity of
Borrelia burgdorferi Erp (OspEF-related) lipoproteins. Microbiology 147:821-830.

57. Hefty PS, Jolliff SE, Caimano MJ, Wikel SK, Akins DR. 2002. Changes in the
temporal and spatial patterns of outer surface lipoprotein expression generate
population heterogeneity and antigenic diversity in the Lyme disease spirochete,
Borrelia burgdorferi. Infect Immun 70:3468-3478.

58. El-Hage N, Stevenson B. 2002. Simultaneous coexpression of *Borrelia burgdorferi*
Erp proteins occurs through a specific, *erp* locus-directed regulatory mechanism. J
Bacteriol 184:4536-4543.

59.   Miller JC, von Lackum K, Babb K, McAlister JD, Stevenson B. 2003. Temporal
647   analysis of *Borrelia burgdorferi* Erp protein expression throughout the mammal-tick
648   infectious cycle. Infect Immun 71:6943-6952.

649   60.   Babb K, McAlister JD, Miller JC, Stevenson B. 2004. Molecular characterization of
650   *Borrelia burgdorferi erp* promoter/operator elements. J Bacteriol 186:2745-2756.

651   61.   Hefty PS, Jolliff SE, Caimano MJ, Wikel SK, Radolf JD, Akins DR. 2001. Regulation
652   of OspE-related, OspF-related, and Elp lipoproteins of *Borrelia burgdorferi* strain
653   297 by mammalian host-specific signals. Infect Immun 69:3618-3627.

654   62.   Eggers CH, Caimano MJ, Radolf JD. 2004. Analysis of promoter elements involved
655   in the transcription initiation of RpoS-dependent *Borrelia burgdorferi* genes. J
656   Bacteriol 186:7390-7402.

657   63.   Eggers CH, Caimano MJ, Radolf JD. 2006. Sigma factor selectivity in *Borrelia
658   burgdorferi*: RpoS recognition of the *ospE/ospF/elp* promoters is dependent on the
659   sequence of the -10 region. Mol Microbiol 59:1859-1875.

660   64.   Stevenson B, Krusenstjerna AC, Castro-Padovani TN, Savage CR, Jutras BL, Saylor
661   TC. 2022. The consistent tick-vertebrate infectious cycle of the Lyme disease
662   spirochete enables *Borrelia burgdorferi* to control protein expression by monitoring
663   its physiological status. J Bacteriol 204:e0060621.

664   65.   Nguyen T-PK, Lam TT, Barthold SW, Telford SR, Flavell RA, Fikrig E. 1994. Partial
665   destruction of *Borrelia burgdorferi* within ticks that engorged on OspE- or OspF-
666   immunized mice. Infect Immun 62:2079-2084.

667   66.   McDowell JV, Sung SY, Price G, Marconi RT. 2001. Demonstration of the genetic
668   stability and temporal expression of select members of the Lyme disease
669   spirochete OspF protein family during infection in mice. Infect Immun 69:4831-
670   4838.

671   67.   Miller JC, von Lackum K, Woodman ME, Stevenson B. 2006. Detection of *Borrelia
672   burgdorferi* gene expression during mammalian infection using transcriptional
673   fusions that produce green fluorescent protein. Microb Pathog 41:43-47.

674   68.   Barbour AG, Jasinskas A, Kayala MA, Davies DH, Steere AC, Baldi P, Felgner PL.
675   2008. A genome-wide proteome array reveals a limited set of immunogens in
676   natural infections of humans and white-footed mice with *Borrelia burgdorferi*. Infect
677   Immun 76:3374-3389.

678   69.   Elsner RA, Hastey CJ, Baumgarth N. 2015. CD4+ T cells promote antibody
679   production but not sustained affinity maturation during *Borrelia burgdorferi*
680   infection. Infect Immun 83:48-56.

681   70.   Elsner RA, Hastey CJ, Olsen KJ, Baumgarth N. 2015. Suppression of long-lived
682   humoral immunity following *Borrelia burgdorferi* Infection. PLoS Pathog
683   11:e1004976.

684   71.   Tracy KE, Baumgarth N. 2017. *Borrelia burgdorferi* manipulates innate and adaptive
685   immunity to establish persistence in rodent reservoir hosts. Front Immunol 8:116.

686   72.   Bockenstedt LK, Wooten RM, Baumgarth N. 2021. Immune response to *Borrelia*:
687   lessons from Lyme disease spirochetes. Curr Issues Mol Biol 42:145-190.

688   73.   Hellwage J, Meri T, Heikkilä T, Alitalo A, Panelius J, Lahdenne P, Seppälä IJT, Meri
689   S. 2001. The complement regulatory factor H binds to the surface protein OspE of
690   *Borrelia burgdorferi*. J Biol Chem 276:8427-8435.

691   74.   Alitalo A, Meri T, Lankinen H, Seppälä I, Lahdenne P, Hefty PS, Akins D, Meri S.
692   2002. Complement inhibitor factor H binding to Lyme disease spirochetes is
693   mediated by inducible expression of multiple plasmid-encoded outer surface
694   protein E paralogs. J Immunol 169:3847-3853.

695   75.   Kraiczy P, Hellwage J, Skerka C, Kirschfink M, Brade V, Zipfel PF, Wallich R. 2003.
696   Immune evasion of *Borrelia burgdorferi*: mapping of a complement inhibitor factor
697   H-binding site of BbCRASP-3, a novel member of the Erp protein family. Eur J
698   Immunol 33:697-707.

699   76.   Metts MS, McDowell JV, Theisen M, Hansen PR, Marconi RT. 2003. Analysis of the
700   OspE determinants involved in binding of factor H and OspE-targeting antibodies
701   elicited during *Borrelia burgdorferi* infection. Infect Immun 71:3587-3596.

702   77.   Kraiczy P, Hartmann K, Hellwage J, Skerka C, Brade V, Zipfel PF, Wallich R,
703   Stevenson B. 2004. Immunological characterization of the complement regulator
704   factor H-binding CRASP and Erp proteins of *Borrelia burgdorferi*. Int J Med
705   Microbiol 293 S37:152-157.

706   78.   Jószi M, Oppermann M, Lambris JD, Zipfel PF. 2007. The C-terminus of
707   complement factor H is essential for host cell protection. Mol Immunol 44:2697-
708   2706.

709   79.   Haupt K, Kraiczy P, Wallich R, Brade V, Skerka C, Zipfel PF. 2008. FHR-1, an
710   additional human plasma protein, binds to complement regulator-acquiring
711   surface proteins of *Borrelia burgdorferi*. Int J Med Microbiol 298, S1:287-291.

712   80.   Kenedy MR, Akins DR. 2011. The OspE-related proteins inhibit complement
713   deposition and enhance serum resistance of *Borrelia burgdorferi*, the Lyme disease
714   agent. Infect Immun 79:1451-1457.

715   81.   Kraiczy P, Stevenson B. 2013. Complement regulator-acquiring surface proteins of
716   *Borrelia burgdorferi*: Structure, function and regulation of gene expression. Ticks
717   Tick Borne Dis 4:26-34.

718   82.   Nowak TA, Lown LA, Marcinkiewicz AL, Sürth V, Kraiczy P, Burke R, Lin YP.
719   2022. Outer surface protein E (OspE) mediates *Borrelia burgdorferi* sensu stricto
720   strain-specific complement evasion in the eastern fence lizard, *Sceloporus*
721   *undulatus*. Ticks Tick Borne Dis 14:102081.

83.   Seling A, Siegel C, Fingerle V, Jutras BL, Brissette CA, Skerka C, Wallich R, Zipfel
PF, Stevenson B, Kraiczy P. 2010. Functional characterization of *Borrelia spielmanii*
outer surface proteins that interact with distinct members of the human factor H
protein family and with plasminogen. Infect Immun 78:39-48.

84.   Zipfel PF, Skerka C, Hellwage J, Jokiranta ST, Meri S, Brade V, Kraiczy P, Noris M,
Remuzzi G. 2002. Factor H family proteins: on complement, microbes and human
diseases. Biochem Soc Trans 30:971-978.

85.   Stevenson B, El-Hage N, Hines MA, Miller JC, Babb K. 2002. Differential binding
of host complement inhibitor factor H by *Borrelia burgdorferi* Erp surface proteins:
a possible mechanism underlying the expansive host range of Lyme disease
spirochetes. Infect Immun 70:491-497.

86.   Hovis KM, Tran E, Sundy CM, Buckles E, McDowell JV, Marconi RT. 2006.
Selective binding of *Borrelia burgdorferi* OspE paralogs to factor H and serum
proteins from diverse animals: possible expansion of the role of OspE in Lyme
disease pathogenesis. Infect Immun 74:1967-1972.

87.   Pereira MJ, Wager B, Garrigues RJ, Gerlach E, Quinn JD, Dowdell A, Osburne MS,
Zückert WR, Kraiczy P, Garcia BL, Leong JM. 2022. Lipoproteome screening of the
Lyme disease agent identifies novel inhibitors of antibody-mediated complement
killing. Proc Natl Acad Sci USA 119:e2117770119.

88.   Garrigues RJ, Thomas S, Leong JM, Garcia BL. 2022. Outer surface lipoproteins
from the Lyme disease spirochete exploit the molecular switch mechanism of the
complement protease C1s. J Biol Chem 298:102557.

89.   Brissette CA, Verma A, Bowman A, Cooley AE, Stevenson B. 2009. The *Borrelia
burgdorferi* outer-surface protein ErpX binds mammalian laminin. Microbiology
155:863-872.

90.   Lin Y-P, Bhowmick R, Coburn J, Leong JM. 2015. Host cell heparan sulfate
glycosaminoglycans are ligands for OspF-related proteins of the Lyme disease
spirochete. Cell Microbiol 17:464-476.

91.   Antonara S, Chafel RM, LaFrance M, Coburn J. 2007. *Borrelia burgdorferi* adhesins
identified using *in vivo* phage display. Mol Microbiol 66:262-276.

92.   Brissette CA, Haupt K, Barthel D, Cooley AE, Bowman A, Skerka C, Wallich R,
Zipfel PF, Kraiczy P, Stevenson B. 2009. *Borrelia burgdorferi* infection-associated
surface proteins ErpP, ErpA, and ErpC bind human plasminogen. Infect Immun
77:300-306.

93.   Fuchs H, Wallich R, Simon MM, Kramer MD. 1994. The outer surface protein A of
the spirochete *Borrelia burgdorferi* is a plasmin(ogen) receptor. Proc Natl Acad Sci
USA 91:12594-12598.

94.   Klempner MS, Noring R, Epstein MP, McCloud B, Hu R, Limentani SA, Rogers
RA. 1995. Binding of human plasminogen and urokinase-type plasminogen

activator to the Lyme disease spirochete, *Borrelia burgdorferi*. J Infect Dis 171:1258-
1265.

95.   Coleman JL, Sellati TJ, Testa JE, Kew RR, Furie MB, Benach JL. 1995. *Borrelia*
burgdorferi binds plasminogen, resulting in enhanced penetration of endothelial
monolayers. Infect Immun 63:2478-2484.

96.   Hu LT, Perides G, Noring R, Klempner MS. 1995. Binding of human plasminogen
to *Borrelia burgdorferi*. Infect Immun 63:3491-3496.

97.   Coleman JL, Gebbia JA, Piesman J, Degen JL, Bugge TH, Benach JL. 1997.
Plasminogen is required for efficient dissemination of *B. burgdorferi* in ticks and for
enhancement of spirochetemia in mice. Cell 89:1111-1119.

98.   Coleman JL, Roemer EJ, Benach JL. 1999. Plasmin-coated *Borrelia burgdorferi*
degrades soluble and insoluble components of the mammalian extracellular
matrix. Infect Immun 67:3929-3936.

99.   Hallström T, Haupt K, Kraiczy P, Hortschansky P, Wallich R, Skerka C, Zipfel PF.
2010. Complement regulator-acquiring surface protein 1 of *Borrelia burgdorferi*
binds to human bone morphogenic protein 2, several extracellular matrix proteins,
and plasminogen. J Inf Dis 202:490-498.

100. Floden AM, Watt JA, Brissette CA. 2011. *Borrelia burgdorferi* enolase is a surface-
exposed plasminogen binding protein. PLoS One 6:e27502.

101. Önder Ö, Humphrey PT, McOmber B, Korobova F, Francella N, Greenbaum DC,
Brisson D. 2012. OspC is potent plasminogen receptor on surface of *Borrelia*
burgdorferi. J Biol Chem 287:16860-16868.

102. Koenigs A, Hammerschmidt C, Jutras BL, Pogoryelov D, Barthel D, Skerka C,
Kugelstadt D, Wallich R, Stevenson B, Zipfel PF, Kraiczy P. 2013. BBA70 of *Borrelia*
burgdorferi is a novel plasminogen-binding protein. J Biol Chem 288:25229-25243.

103. Magnarelli LA, Ijdo JW, Padula SJ, Flavell RA, Fikrig E. 2000. Serologic diagnosis
of Lyme borreliosis by using enzyme-linked immunosorbent assays with
recombinant antigens. J Clin Microbiol 38:1735-1739.

104. Miller JC, El-Hage N, Babb K, Stevenson B. 2000. *Borrelia burgdorferi* B31 Erp
proteins that are dominant immunoblot antigens of animals infected with isolate
B31 are recognized by only a subset of human Lyme disease patient sera. J Clin
Microbiol 38:1569-1574.

105. Hefty PS, Brooks CS, Jett AM, White GL, Wikel SK, Kennedy RC, Akins DR. 2002.
OspE-related, OspF-related, and Elp lipoproteins are immunogenic in baboons
experimentally infected with *Borrelia burgdorferi* and in human Lyme disease
patients. J Clin Microbiol 40:4256-4265.

106. Panelius J, Meri T, Seppälä I, Eholuoto M, Alitalo A, Meri S. 2008. Outer surface
protein E antibody response and its effect on complement factor H binding to
OspE in Lyme borreliosis. Microbes Infect 10:135-142.

107. Zückert WR, Meyer J, Barbour AG. 1999. Comparative analysis and
immunological characterization of the *Borrelia* Bdr protein family. Infect Immun
67:3257-3266.

108. Carlyon JA, Roberts DM, Marconi RT. 2000. Evolutionary and molecular analyses
of the *Borrelia bdr* super gene family: delineation of distinct sub-families and
demonstration of the genus wide conservation of putative functional domains,
structural properties and repeat motifs. Microb Pathogen 28:89-105.

109. Carlyon JA, Roberts DM, Theisen M, Sadler C, Marconi RT. 2000. Molecular and
immunological analyses of the *Borrelia turicatae* Bdr protein family. Infect Immun
68:2369-2373.

110. Roberts DM, Carlyon JA, Theisen M, Marconi RT. 2000. The *bdr* gene families of
the Lyme disease and relapsing fever spirochetes: potential influence on biology,
pathogenesis, and evolution. Emerging Infect Dis 6:110-122.

111. Roberts DM, Theisen M, Marconi RT. 2000. Analysis of the cellular localization of
Bdr paralogs in *Borrelia burgdorferi*, a causative agent of Lyme disease: evidence for
functional diversity. J Bacteriol 182:4222-4226.

112. Zückert WR, Barbour AG. 2000. Stability of *Borrelia burgdorferi bdr* loci in vitro and
in vivo. Infect Immun 68:1727-1730.

113. Roberts DM, Caimano M, McDowell J, Theisen M, Holm A, Orff E, Nelson D,
Wikel S, Radolf J, Marconi RT. 2002. Environmental regulation and differential
production of members of the Bdr protein family of *Borrelia burgdorferi*. Infect
Immun 70:7033-7041.

114. Zhang H, Raji A, Theisen M, Hansen PR, Marconi RT. 2005. BdrF2 of Lyme disease
spirochetes is coexpressed with a series of cytoplasmic proteins and is produced
specifically during early infection. J Bacteriol 187:175-184.

115. Baum E, Grosenbaugh DA, Barbour AG. 2014. Diversity of antibody responses to
Borrelia burgdorferi in experimentally infected beagle dogs. Clin Vaccine Immunol
21:838-846.

116. Carroll JA, El-Hage N, Miller JC, Babb K, Stevenson B. 2001. *Borrelia burgdorferi*
RevA antigen is a surface-exposed outer membrane protein whose expression is
regulated in response to environmental temperature and pH. Infect Immun
69:5286-5293.

117. Brissette CA, Bykowski T, Cooley AE, Bowman A, Stevenson B. 2009. *Borrelia*
burgdorferi RevA antigen binds host fibronectin. Infect Immun 77:2802-2812.

118.   Gilmore RD, Mbow ML. 1998. A monoclonal antibody generated by antigen
835   inoculation via tick bite is reactive to the *Borrelia burgdorferi* Rev protein, a member
836   of the 2.9 gene family locus. Infect Immun 66:980-986.

837   119.   Nowalk AJ, Gilmore RD, Carroll JA. 2006. Serologic proteome analysis of *Borrelia*
838   *burgdorferi* membrane-associated proteins. Infect Immun 74:3864-3873.

839   120.   Brissette CA, Rossmann E, Bowman A, Cooley AE, Riley SP, Hunfeld K-P, Bechtel
840   M, Kraiczy P, Stevenson B. 2010. The borrelial fibronectin-binding protein RevA is
841   an early antigen of human Lyme disease. Clin Vac Immunol 17:274-280.

842   121.   Floden AM, Gonzalez T, Gaultney RA, Brissette CA. 2013. Evaluation of RevA, a
843   fibronectin-binding protein of *Borrelia burgdorferi*, as a potential vaccine candidate
844   for Lyme disease. Clin Vaccine Immunol 20:892-899.

845   122.   Probert WS, Johnson BJB. 1998. Identification of a 47 kDa fibronectin-binding
846   protein expressed by *Borrelia burgdorferi* isolate B31. Mol Microbiol 30:1003-1015.

847   123.   Byram R, Gaultney RA, Floden AM, Hellekson C, Stone BL, Bowman A, Stevenson
848   B, Johnson BJB, Brissette CA. 2015. *Borrelia burgdorferi* RevA significantly affects
849   pathogenicity and host response in the mouse model of Lyme disease. Infect
850   Immun 83:3675-3683.

851   124.   Schmit VL, Patton TG, Gilmore RD. 2011. Analysis of *Borrelia burgdorferi* surface
852   proteins as determinants in establishing host cell interactions. Front Microbiol
853   2:141.

854   125.   Brissette CA, Gaultney RA. 2014. That's my story, and I'm sticking to it—an update
855   on *B. burgdorferi* adhesins. Front Cell Infect Microbiol 4:41, doi:
856   10.3389/fcimb.2014.00041.

857   126.   Yang X, Popova TG, Hagman KE, Wikel SK, Schoeler GB, Caimano MJ, Radolf JD,
858   Norgard MV. 1999. Identification, characterization, and expression of three new
859   members of the *Borrelia burgdorferi* Mlp (2.9) lipoprotein gene family. Infect Immun
860   67:6008-6018.

861   127.   Porcella SF, Fitzpatrick CA, Bono JL. 2000. Expression and immunological analysis
862   of the plasmid-borne *mlp* genes of *Borrelia burgdorferi* strain B31. Infect Immun
863   68:4992-5001.

864   128.   Yang X, Goldberg MS, Popova TG, Schoeler GB, Wikel SK, Hagman KE, Norgard
865   MV. 2000. Interdependence of environmental factors influencing reciprocal
866   patterns of gene expression in virulent *Borrelia burgdorferi*. Mol Microbiol 37:1470-
867   1479.

868   129.   Yang XF, Hübner A, Popova TG, Hagman KE, Norgard MV. 2003. Regulation of
869   expression of the paralogous Mlp family in *Borrelia burgdorferi*. Infect Immun
870   71:5012-5020.

871   130.   Yang X, Popova TG, Goldberg MS, Norgard MV. 2001. Influence of cultivation
872   media on genetic regulatory patterns in *Borrelia burgdorferi*. Infect Immun 69:4159-
873   4163.

874   131.   Xu Y, Bruno JF, Luft BJ. 2008. Profiling the humoral immune response to Borrelia
875   burgdorferi infection with protein microarrays. Microb Pathog 45:403-407.

876   132.   Caimano MJ, Groshong AM, Belperron A, Mao J, Hawley KL, Luthra A, Graham
877   DE, Earnhart CG, Marconi RT, Bockenstedt LK, Blevins JS, Radolf JD. 2019. The
878   RpoS gatekeeper in *Borrelia burgdorferi*: an invariant regulatory scheme that
879   promotes spirochete persistence in reservoir hosts and niche diversity. Front
880   Microbiol 10:1923.

881   133.   Tkáčová Z, Borszéková Pulzová L, Mochnáčová E, Jiménez-Munguía I, Bhide K,
882   Mertinková P, Majerová P, Kulkarni A, Kováč A, Bhide M. 2020. Identification of
883   the proteins of *Borrelia garinii* interacting with human brain microvascular
884   endothelial cells. Ticks Tick Borne Dis 11:101451.

885   134.   Fridmanis J, Otikovs M, Brangulis K, Tārs K, Jaudzems K. 2021. Solution NMR
886   structure of *Borrelia burgdorferi* outer surface lipoprotein BBP28, a member of the
887   mlp protein family. Proteins 89:588-594.

888   135.   Champion CI, Blanco DR, Skare JT, Haake DA, Giladi M, Foley D, Miller JN, Lovett
889   MA. 1994. A 9.0-kilobase-pair circular plasmid of *Borrelia burgdorferi* encodes an
890   exported protein: evidence for expression only during infection. Infect Immun
891   62:2653-2661.

892   136.   Miller JC, Bono JL, Babb K, El-Hage N, Casjens S, Stevenson B. 2000. A second
893   allele of *eppA* in *Borrelia burgdorferi* strain B31 is located on the previously
894   undetected circular plasmid cp9-2. J Bacteriol 182:6254-6258.

895   137.   Miller JC, Stevenson B. 2003. Immunological and genetic characterization of
896   *Borrelia burgdorferi* BapA and EppA proteins. Microbiology 149:1113-1125.

897   138.   Stevenson B. 2002. *Borrelia burgdorferi erp* (*ospE*-related) gene sequences remain
898   stable during mammalian infection. Infect Immun 70:5307-5311.

899   139.   Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. 1982.
900   Lyme disease - a tick-borne spirochetosis? Science 216:1317-1319.

901   140.   Schwan TG, Burgdorfer W, Schrumpf ME, Karstens RH. 1988. The urinary
902   bladder, a consistent source of *Borrelia burgdorferi* in experimentally infected white-
903   footed mice (*Peromyscus leucopus*). J Clin Microbiol 26:893-895.

904   141.   Wang G, Ojaimi C, Iyer R, Saksenberg V, McClain SA, Wormser GP, Schwartz IA.
905   2001. Impact of genotypic variation of *Borrelia burgdorferi* sensu stricto on kinetics
906   of dissemination and severity of disease in C3H/HeJ mice. Infect Immun 69:4303-
907   4312.

142.   Steere AC, Grodzicki RL, Kornblatt AN, Craft JE, Barbour AG, Burgdorfer W,
Schmid GP, Johnson E, Malawista SE. 1983. The spirochetal etiology of Lyme
disease. N Engl J Med 308:733-740.

143.   Jutras BL, Chenail AM, Stevenson B. 2013. Changes in bacterial growth rate govern
expression of the *Borrelia burgdorferi* OspC and Erp infection-associated surface
proteins. J Bacteriol 195:757-764.

144.   Goethert HK, Mather TN, Buchthal J, Telford SR. 2021. Retrotransposon-based
blood meal analysis of nymphal deer ticks demonstrates spatiotemporal diversity
of *Borrelia burgdorferi* and *Babesia microti* reservoirs. Appl Environ Microbiol
87:e02370-20.

145.   Gylfe Å, Olsen B, Strasevicius D, Marti Ras N, Weihe P, Noppa L, Ostberg Y,
Baranton G, Bergström S. 1999. Isolation of Lyme disease *Borrelia* from puffins
(*Fratercula arctica*) and seabird ticks (*Ixodes uriae*) on the Faeroe Islands. J Clin
Microbiol 37:890-896.

146.   Casjens SR, Mongodin EF, Qiu WG, Dunn JJ, Luft BJ, Fraser-Liggett CM, Schutzer
EE. 2011. Whole-genome sequences of two *Borrelia afzelii* and two *Borrelia garinii*
Lyme disease agent isolates. J Bacteriol 193:6995-6996.

147.   Stevenson B, Barthold SW. 1994. Expression and sequence of outer surface protein
C among North American isolates of *Borrelia burgdorferi*. FEMS Microbiol Lett
124:367-372.

148.   Jauris-Heipke S, Liegl G, Preac-Mursic V, Robler D, Schwab E, Soutschek E, Will
G, Wilske B. 1995. Molecular analysis of genes encoding outer surface protein C
(OspC) of *Borrelia burgdorferi* sensu lato: relationship to *ospA* genotype and
evidence of lateral gene exchange of *ospC*. J Clin Microbiol 33:1860-1866.

149.   Stewart PE, Thalken R, Bono JL, Rosa P. 2001. Isolation of a circular plasmid region
sufficient for autonomous replication and transformation of infectious *Borrelia*
burgdorferi. Mol Microbiol 39:714-721.

150.   Eggers CH, Caimano MJ, Clawson ML, Miller WG, Samuels DS, Radolf JD. 2002.
Identification of loci critical for replication and compatibility of a *Borrelia*
burgdorferi cp32 plasmid and use of a cp32-based shuttle vector for the expression
of fluorescent reporters in the Lyme disease spirochaete. Mol Microbiol 43:281-295.

151.   Stewart PE, Chaconas G, Rosa P. 2003. Conservation of plasmid maintenance
functions between linear and circular plasmids in *Borrelia burgdorferi*. J Bacteriol
185:3202-3209.

152.   Guina T, Oliver DB. 1997. Cloning and analysis of a *Borrelia burgdorferi* membrane-
interactive protein exhibiting haemolytic activity. Mol Microbiol 24:1201-1213.
944