An ESCRT-dependent pathway coordinates Nuclear and Cytoplasmic Spatial Protein Quality Control at Nuclear Vacuolar Junctions

Emily M. Sontag [1,*,†,£], Fabián Morales-Polanco [1,£], Jian-Hua Chen[2], Gerry McDermott[2], Patrick T. Dolan[1,‡], Dan Gestaut[1], Mark A. Le Gros[2,3], Carolyn Larabell[2,3] and Judith Frydman[1,*]

[1] Department of Biology, Stanford University, Stanford, CA 94305, USA

[2] Department of Anatomy, School of Medicine, University of California San Francisco, San Francisco, CA 94158, USA

[3] Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

[*] Correspondence to: emily.sontag@marquette.edu; jfrydman@stanford.edu,

[£] These authors contributed equally

[†] Current address: Department of Biological Sciences, Marquette University, Milwaukee, WI 53233, USA

[‡] Current address: Quantitative Virology and Evolution Unit, National Institute of Allergy and Infectious Diseases, Bethesda, MD 20892, USA

Abstract.

Effective Protein Quality Control (PQC), essential for cellular health, relies on spatial

sequestration of misfolded proteins into defined inclusions. Here we elucidate the coordination of

nuclear and cytoplasmic spatial PQC. While cytoplasmic misfolded proteins concentrate in a

cytoplasmic, perinuclear Juxta Nuclear Quality control compartment (JUNQ), nuclear misfolded

proteins sequester into a perinucleolar IntraNuclear Quality control compartment (INQ). Particle

tracking reveals the INQ and JUNQ converge to face each other across the nuclear envelope at a

site proximal to the Nuclear-Vacuolar Junction (NVJ) marked by perinuclear ESCRT-II/-III

protein Chm7. Strikingly, this ESCRT-dependent convergence facilitates VPS4-dependent

vacuolar clearance of misfolded cytoplasmic and nuclear proteins, the latter entailing extrusion of

nuclear INQ into the vacuole. We propose perinuclear ESCRT coordinates spatial PQC at nuclear-

vacuolar contacts to facilitate vacuolar clearance of nuclear and cytoplasmic misfolded proteins.

Introduction

Misfolded proteins can acquire toxic conformations that disrupt essential cellular processes [1-6] leading to human diseases ranging from neurodegeneration to cancer [7-10]. Accordingly, cellular protein homeostasis (proteostasis) is maintained by a network of chaperones and clearance factors that promote protein quality control (PQC) [11-15]. One fundamental PQC strategy is to spatially sequester misfolded proteins into distinctly localized membrane-less compartments, likely to remove damaging conformers from the cellular milieu and concentrate them for more efficient refolding or clearance through either the ubiquitin-proteasome system (UPS) or endo-lysosomal pathways [11, 16-18]. Misfolded cytoplasmic proteins partition into distinct inclusions depending on their aggregation state. Insoluble amyloid proteins are sequestered into the Insoluble Protein Deposit (herein the IPOD) [4, 11, 19]. Soluble misfolded proteins are sequestered into small, dynamic, ER-associated structures called Q-bodies [17] which, during sustained stress, coalesce along the endomembrane system [20] into a Juxtanuclear Quality Control compartment (herein the JUNQ). Spatial PQC through the IPOD and JUNQ pathways is evolutionarily conserved and is present in yeast, worms and mammalian cells [11, 19]. Spatial PQC is also observed in the nucleus, as nuclear misfolded proteins are sequestered in a membrane-less intranuclear quality control compartment (herein the INQ), proximal to the nucleolus [18, 21, 22].

The relationship and overall coordination between spatial PQC of cytoplasmic and nuclear proteins is not well understood. Many studies showed cytoplasmic misfolded proteins being concentrated into the cytoplasmic JUNQ and cleared through an ER-anchored cytoplasmic UPS pathway [11, 19, 23, 24]. It has also been proposed that clearance of cytoplasmic misfolded proteins requires import into the nucleus, whereby the JUNQ would form an inclusion identical to the INQ [18].

Given the importance of protein quality control for cell survival and its relevance to understand misfolding diseases, we here examined the logic and coordination of spatial PQC for nuclear and cytoplasmic misfolded proteins. We unequivocally demonstrate the cytoplasmic localization of the JUNQ and nuclear localization of the INQ. We further show the JUNQ and INQ home to opposite sides of the nuclear envelope, at a site proximal to NVJ, which facilitates the extrusion of the INQ for clearance into the vacuole in an ESCRT- and VPS4-dependent manner. Our study uncovers a surprising degree of spatial coordination between nuclear and cytoplasmic PQC inclusions providing an avenue for vacuolar clearance of misfolded proteins.

RESULTS

Organelle-specific PQC compartments sequester nuclear and cytoplasmic misfolded proteins

To define the relationship and interplay between nuclear and cytoplasmic spatial PQC (Fig. 1a), we utilized two validated models of protein misfolding: a temperature-sensitive variant of Firefly Luciferase (herein LuciTs) that misfolds at 37 °C, and the Von Hippel-Lindau (VHL) tumor suppressor, which is constitutively unfolded in yeast cells [25, 26]. We restricted their subcellular localization to either nucleus or cytoplasm by inclusion of a Nuclear Localization Signal (NLS) or a Nuclear Export Signal (NES), respectively. The expression of these PQC substrates was first selectively induced by growth in galactose at the permissive temperature, and then repressed by glucose exchange, which allowed measuring the kinetics of their clearance or spatial PQC (Fig. 1b). As previously described, both NLS- and NES-LuciTs and NLS- and NES-VHL are degraded

upon misfolding and are stabilized by proteasomal inhibition with either MG132 or Bortezomib [27]

(Extended Data Fig. 1a and Ref [27]).

To initiate LuciTs misfolding and monitor the fate of the misfolded species, the cells were shifted

to 37 °C and evaluated using time-resolved fluorescence microscopy (Fig. 1b). Under folding-

permissive conditions, NLS- and NES-GFP-LuciTs were diffusely localized in the nucleus or

cytoplasm, respectively (Time 0, Fig 1 c, d), but following the shift to 37 °C, both nuclear (NLS)

and cytoplasmic (NES) LuciTs were rapidly concentrated into dynamic puncta in their respective

cellular locations (Fig. 1c, d, and Supplemental video 1).

Over the course of the experiment, nuclear NLS-LuciTs coalesced into one intranuclear INQ

inclusion (Time 30 min, Fig. 1c). In contrast, cytoplasmic NES-LuciTs puncta coalesced into two

cytoplasmic inclusions, corresponding to the JUNQ and IPOD (Time 30 min, Fig. 1d), as

previously observed for Ubc9Ts and VHL [11]. Of note, the NLS- and NES- LuciTs variants formed

inclusions in both the absence or presence of proteasome inhibitor MG132 (Figure 1c, d),

indicating that spatial sequestration into PQC compartments is not a consequence of proteasomal

inhibition.

We used structured illumination (SIM) super-resolution microscopy to better define the nuclear

and cytoplasmic location and morphology of PQC inclusions. Cells expressing LuciTs variants

were incubated for 2 hrs at 37 °C, while cells expressing VHL variants were grown and fixed at

both 30 and 37 °C. The nuclear envelope was visualized by immunostaining of nuclear pore

protein Nsp1 (Nups, yellow) and DNA was visualized using Hoechst (blue) (Fig. 1e-h). The INQ,

formed by nuclear misfolded protein, localized inside the nucleus in a pocket between the nuclear

envelope (delineated by Nups) and the DNA (Fig. 1e and Supplemental video 2 for NLS-LuciTs

at 37 °C, Fig. 1f for NLS-VHL at 30 °C Extended Data Fig. 1c for NLS-VHL at 37 °C). Line

intensity profile analyses confirmed the intranuclear localization of this INQ (Fig. 1e, f, Extended

Data Fig. 1c). Importantly, SIM and line intensity profile analyses unequivocally established that

the PQC inclusions formed by cytoplasmic misfolded proteins NES-LuciTs and NES-VHL reside

outside of the nucleus: one inclusion is adjacent to the nuclear envelope as expected for the JUNQ,

while the other inclusion is closer to the cell periphery as expected for the IPOD (Fig. 1g-h,

Supplemental video 2, Extended Data Fig. 1d). Similar results were obtained with NLS- and NES-

VHL at both 30 °C and 37 °C indicating PQC inclusion localization is not temperature-dependent

(Extended Data Fig. 1c-d). Consistent with their distinct localization, cytoplasmic and nuclear

misfolded variants also exhibit differential toxicity to cells; cytoplasmic misfolded proteins were

slightly more toxic, albeit less so than a polyglutamine expanded exon1 variant of Huntingtin

(herein mHTT) associated with Huntington's disease (Extended Data Fig. 1b). The increased NES-

LuciTs toxicity was only observed at 37 °C when it is misfolded.

We conclude that nuclear and cytoplasmic misfolded proteins are spatially sequestered in PQC

inclusions located in their cognate compartment, yielding the INQ and JUNQ respectively. These

results resonate with findings that UPS degrades nuclear and cytoplasmic misfolded proteins

through different chaperones and ubiquitination enzymes [27], indicating the nucleus and

cytoplasm have compartment specific PQC machineries.

Proteasome inhibition disrupts nucleocytoplasmic transport

Most commonly used PQC reporters (e.g., Ubc9Ts, VHL, Cpy*, and Luciferase, etc.), are smaller

than the 200 kDa nuclear permeability barrier [28-30]. To examine if passive diffusion across nuclear

pores could explain why some cytoplasmic misfolded proteins can form intranuclear inclusions,

we utilized a temperature-sensitive variant of Ubc9 (Ubc9Ts) [11, 17, 18]. Upon Ubc9Ts unfolding at

37 °C for 2 hours under conditions where the proteasome is active, SIM super-resolution

microscopy showed Ubc9Ts is sequestered solely in two cytoplasmic compartments – the

perinuclear JUNQ, clearly outside the nuclear pore boundary and the IPOD in the cell periphery

(Fig. 2a, Supplemental Video 3). Treatment at 37 °C with added proteasome inhibitor MG132,

increased the fraction of cells containing Ubc9Ts inclusions, but in addition to the two cytoplasmic

inclusions as above, some cells (around 13%) had an additional Ubc9Ts inclusion inside the

nucleus (Fig. 2b, Extended Data Fig. 2a, Supplemental Video 3). Similar results were obtained for

another PQC substrate, VHL (Extended Data Fig. 2b). Interestingly, line intensity profile analyses

of both Ubc9Ts and VHL showed that the nuclear INQ and cytoplasmic JUNQ inclusions formed

in close proximity but separated by nuclear pores (Fig. 2b), These results confirmed that, absent

an NLS or NES localization tag, cytoplasmic proteins that can passively diffuse into the nucleus,

primarily form cytoplasmic inclusions, but under conditions of proteasome inhibition can also

form an additional distinct nuclear INQ in a small fraction of cells.

To reconcile the finding that spatial sequestration into a nuclear or a cytoplasmic PQC inclusion

depends on a misfolded protein's subcellular location when they misfold (Fig. 1a) with the fact

that MG132 leads to a small fraction of cells having a nuclear Ubc9Ts and VHL inclusion, we

hypothesized that prolonged proteasome inhibition may alter nucleocytoplasmic trafficking and

decrease protein export from the nucleus. Indeed, toxic misfolded proteins, such as the

polyglutamine expanded mHTT, are shown to impair nucleocytoplasmic transport [31, 32].

We examined the effect of MG132 on nucleocytoplasmic transport using a previously

characterized NLS- and NES-tagged shuttling GFP (sGFP) that migrates between the nucleus and

cytoplasm [31, 33] (Fig. 2c, d). As a control, we observed toxic mHTT expression caused nuclear

retention of sGFP as described (Fig. 2c, d). Of note, proteasome inhibition with MG132 caused

sGFP nuclear retention to the same extent as mHTT (Fig. 2c, d). This experiment was performed

in leptomycin B-sensitive crm^{T539C} cells, which permitted the use of leptomycin B (LMB) as a

positive control for blocked nuclear export (Fig. 2c, d), but similar results were obtained using WT

BY4741 yeast cells (not shown). Our finding that proteasome inhibition impairs nuclear export to

an extent comparable to toxic mHTT may explain the increased retention of misfolded protein

Ubc9Ts in the nucleus. Future studies should examine the basis for reducing nuclear export when

proteostasis is impaired [31, 34, 35].

To directly assess if cytoplasmic misfolded protein clearance requires their import to the nucleus,

we examined Ubc9Ts clearance in cells carrying a temperature-sensitive mutation in nuclear pore

complex (NPC) protein Nup116 (*nup116-5)*. At 37 °C, the *nup116-5* mutation seals nuclear pores

over the cytoplasmic face of NPCs blocking any transport into and out of the nucleus [36] (Fig. 2f,

i). WT and *nup116-5* cells expressing Ubc9Ts were shifted to 37 °C for 1 hour prior to the glucose-

repression to ensure nuclear pores become completely sealed prior to measuring Ubc9Ts clearance

rates (Figure. 2e, *ii*). If degradation of misfolded Ubc9Ts requires import into the nucleus, it should

be abrogated in *nup116-5* cells at 37 °C. We did not observe any significant differences in the rate

and the extent of Ubc9Ts degradation in *nup116-5* cells, suggesting that transport into the nucleus

is not required for its clearance (Figure. 2f, *ii-iii*; Extended Data Fig. 2c).

We next examined whether nuclear proteasomes participate in clearance of a cytoplasmic

misfolded protein. To this end, we used *sts1-2* cells that carry a mutation in Sts1, a protein required

for import of proteasomes into the nucleus and facilitate the degradation of nuclear proteins. At 37

°C, nuclear localization of proteasomes is lost in *sts1-2* cells [37-40] (Fig. 2g panel *i*). However,

preventing nuclear localization of proteasomes had no effect on the clearance of misfolded

Ubc9Ts, consistent with cytoplasmic degradation (Fig. 2g, panel *ii-iii*; Extended Data Fig. 2d).

These experiments indicate that import of misfolded proteins and proteosomes into the nucleus is

dispensable to degrade misfolded Ubc9Ts.

INQ and JUNQ home into adjoined locations on opposite sides of the nuclear envelope

Given that nuclear and cytoplasmic Ubc9Ts inclusions in MG132-treated cells localized to

opposite sides of the nuclear envelope (Fig. 2b), we next sought to define the spatial relationship

between INQ and JUNQ in live cells. To this end, NLS- LuciTs and NES- LuciTs were co-

expressed in WT yeast cells at 25 °C for 6 hr before repressing their synthesis with glucose. Cells

were shifted to 37 °C to unfold both proteins simultaneously, and inclusion formation was

monitored in real-time by live cell time-lapse fluorescence microscopy and particle tracking

analysis (Fig. 3a, b; Extended Data Fig. 3a; Supplemental Videos 4, 5). Spatial sequestration of

the NLS- and NES- misfolded proteins into motile condensates occurred with similar kinetics

independently in the nucleus and the cytoplasm. Once formed, puncta moved dynamically toward

each other, with nuclear and cytoplasmic inclusions converging on a specific location proximal to

the nuclear envelope (Fig. 3a). Of note, even after homing to this location, the two inclusions

remain distinct and separated; however, they co-migrated around the nuclear periphery as if

tethered to each other across the nuclear envelope. Particle tracking analyses measuring the

location and distance of NLS- and NES-LuciTs inclusions in 2D space showed the nuclear and

cytoplasmic inclusions homed into adjoining locations between 10 and 20 minutes after shifting

to 37 °C, and then continued moving in a coordinated manner throughout the duration of the

imaging experiment (Fig. 3b; Extended Data Fig. 3a; Supplemental video 5). While the distance

between these two inclusions decreased rapidly during homing, once they came together they

maintained a largely stable short distance, suggestive of a tether, throughout their co-migration for

the remainder of the experiment. Similar results were obtained if the fluorescent markers in NLS-

and NES-LuciTs were reversed (not shown) as well as for NLS- and NES-VHL (Extended Data

Fig. 3b). These observations, supporting our findings with Ubc9Ts, suggest a novel mechanism

coordinating convergence of nuclear and cytoplasmic PQC compartments.

The spatial relationship between the INQ and the JUNQ was confirmed using SIM super-resolution

microscopy. Cells co-expressing NLS-LuciTs and NES-LuciTs were incubated at 37 °C for 2 hrs

and then fixed and immunostained to detect nuclear pores (Fig. 3c). SIM combined with line

intensity profile analysis confirmed the INQ and JUNQ adjoined each other on opposite sides of

the nuclear envelope, separated by the nuclear pores (Fig. 3c; Supplemental video 6).

Nuclear pores play a role in INQ-JUNQ convergence at the nuclear periphery

To understand the nature of the signal bringing together the INQ and JUNQ, we examined their

subcellular co-localization with specific nuclear and perinuclear structures. The INQ was near the

nucleolus but not in direct contact, suggesting the nucleolus is not driving this homing mechanism

(Extended Data Fig. 3c). We examined the relationship of the INQ-JUNQ homing signal with two

complexes that localize to a defined perinuclear location: the spindle pole body [41] (SPB) and the

LINC complex [42], both of which contain protein Mps3 spanning both layers of the nuclear envelope

(Extended Data Fig. 3d). Co-staining with Mps3 showed neither SPB nor LINC co-localize with

the INQ-JUNQ perinuclear site, suggesting they do not participate in the INQ-JUNQ homing

mechanism (Extended Data Fig. 3d).

To investigate the role of nuclear pores in INQ/JUNQ homing, we exploited the finding that

deletion of nuclear pore protein Nup120 causes nuclear pores to remain functional but become

clustered [43, 44]. We thus compared INQ and JUNQ formation and localization in WT and *nup120Δ*

cells. In *nup120Δ* cells, the JUNQ always forms next to the clustered nuclear pores (Fig. 3d, panel

i) while the INQ did not localize proximal to the nuclear pores, but instead was found anywhere

within the nucleus (Fig. 3d, panel *ii*). Surprisingly, if we co-expressed NLS- and NES-LuciTs in

nup120Δ cells, the INQ now homed to the clustered nuclear pores opposite the JUNQ (Fig. 3d,

panel *iii*). This finding suggests formation of the JUNQ at the cytoplasmic side of the nuclear pores

communicates a signal for INQ recruitment to this location (Fig. 3e).

We next asked if the intrinsically disordered phenylalanine-glycine (FG) repeats in the central

nuclear pore channel, which form hydrogels capable of amyloid-like interactions [45], facilitate INQ-

JUNQ homing. Using live-cell time-lapse microscopy we examined the localization of the INQ

and JUNQ in *nupΔFG* cells, carrying deletions in the FG repeat tracts of six nuclear pore proteins

that removes approximately 70% of the FG content from the nuclear pore [46]. We found that the

INQ and JUNQ still localized to the same perinuclear location in *nupΔFG* cells suggesting the

high percentage of FG-repeats are dispensable for homing (Extended Data Fig. 3e). Although, we

cannot exclude the possibility that that the remaining 30% of FG content is still sufficient to direct

convergence of the INQ and JUNQ to the proximity of nuclear pores.

Cryo-soft X-ray tomography reveals the architecture of PQC inclusions in intact cells

To examine the native cellular context of PQC compartments we used cryogenic Soft X-ray

Tomography (cryo-SXT). This synchrotron-based imaging approach can visualize and quantify

the ultrastructure of intact, unstained, cryo-preserved cells (Fig. 4a, Extended Data Fig. 4a, b) [47,

48]. cryo-SXT uses the characteristic linear absorption of X-rays by different subcellular organelles

to produce projection images of cellular ultrastructure to a few tens nanometers of spatial

resolution [49]. To derive the information content needed to study misfolded protein inclusions we

initially used cryo-SXT in correlation with cryo-fluorescence microscopy (Fig. 4b,i). The ability

to absorb X-rays varies with the type and density of biomolecules present in different

compartments, yielding characteristic linear absorption coefficients (LAC) (Fig. 4b, ii) [50-52]. Since

chaperone Hsp104 localizes to both the JUNQ and IPOD upon heat shock, while aggregation-

prone mHTT only localizes to the IPOD [11, 20], we visualized these PQC compartments by

expressing mHTT-ChFP in cells with chromosomally tagged HSP104-GFP [53]. The inclusions

were formed by 1 hr incubation at 37 °C and cells were flash frozen in capillary tubes and directly

imaged using cryo-SXT correlated with cryo-fluorescence microscopy (Fig. 4b, i). The LAC

values were used to identify the exact location of specific compartments and organelles in X-ray

tomograms of intact frozen yeast cells (Fig. 4b, ii), yielding a 3D reconstruction of the PQC

inclusions in the context of cellular architecture (Fig. 4c, Supplemental Video 7). Cryo-

fluorescence identified the PQC compartments, with Hsp104-only marking the JUNQ and co-

localized Hsp104 and mHTT marking the IPOD. This analysis revealed that protein inclusions

have a characteristic LAC that can be used to examine their location and size without the need of

fluorescence microscopy. Of note, this correlated fluorescence-cryo-SXT analyses confirmed both

the JUNQ and the IPOD are cytoplasmic, as previously described (Fig. 4c, Supplemental Video 7

and [11]).

We next used cryo-SXT to directly detect inclusions formed by NLS- and NES-LuciTs at 37 °C

through their characteristic LAC values. NES-LuciTs formed small, cytoplasmic, punctate Q-

bodies that could be seen congregating at a cytoplasmic, juxtanuclear location to form the JUNQ

(Fig. 4d, Supplemental Video 8). Thus, cryo-SXT indicates the JUNQ is not the homogeneous

large compartment suggested by diffraction-limited fluorescence imaging, but rather consists of

multiple dense "cores" that resemble Q-bodies. This conclusion, also supported by the irregular

JUNQ morphology in SIM and cryo-EM imaging (Fig. 1, Fig 5 and [54]), suggests the JUNQ is in

fact a collection of Q-bodies that congregate at a perinuclear location proximal to the nuclear

envelope (Fig. 4d, inset *ii*; Supplemental video 8).

Cryo-SXT analysis also revealed a link between cytoplasmic PQC and mitochondria. Individual

Q-bodies can associate with the surface of mitochondria (Fig. 4e) and the JUNQ and IPOD appear

surrounded by a mitochondrial cage (Fig. 4d, *i*). Confocal fluorescence microscopy of cells

expressing mito-GFP supported the mitochondrial association with Q-bodies (Extended Data Fig.

4d). However, PQC compartments still formed in cells with disruptions in mitochondrial fission

or fusion (Extended Data Fig. 4e) indicating mitochondrial structure does not direct PQC

compartment formation. Since previous studies placed Q-bodies on the ER membrane [17, 20], which

cannot be visualized by cryo-SXT, it is possible that Q-bodies are proximal to ER-mitochondrial

contact sites. Alternatively, the proximity to mitochondria may serve to place energy-intensive

PQC compartments in areas of the cell enriched in ATP production [55].

Cryo-SXT also confirmed that the intranuclear INQ is distinct from but resides near the nucleolus

(Fig. 4g). This conclusion is supported by confocal fluorescence microscopy with nucleolar protein

Nsr1 (Extended Data Fig. 3c, 4c and [18, 31]). The proximity of the nucleolus to the INQ is consistent

with the nucleolus' role in nuclear folding and assembly [21, 56-58] and suggest a possible link to the

INQ's function in nuclear PQC [56, 59].

Strikingly, our cryo-SXT imaging revealed an unexpected link of both INQ and JUNQ with the

vacuole. The entire JUNQ site was surrounded by nuclear-vacuolar inter-organelle contact sites

(Figure. 4f, red arrows; Supplemental Video 8). Remarkably, Q-bodies within the JUNQ area were

nestled within invaginations in the vacuolar membrane, suggestive of direct vacuolar engulfment

and raising the idea that the vacuole promotes JUNQ clearance (Figure. 4f, blue arrows;

Supplemental Video 8). The INQ was also proximal to the vacuolar side of the nucleus, indicating

both INQ and JUNQ are steered to the vicinity of nuclear-vacuolar contact sites (Fig. 4g). The

nucleus-vacuole contact sites are formed by a tethering Nuclear-Vacuolar Junction (NVJ) complex

between nuclear Nvj1 and vacuolar Vac8 (Fig. 4h and [60]). Consistent with the idea that the INQ

and JUNQ converge to the vicinity of NVJs, all nuclear pores in *nup120Δ* cells cluster at the

vacuole-contacting side of the nucleus [44].

Nuclear-Vacuolar Junctions facilitate clearance of JUNQ and INQ

The role of NVJs remains poorly understood, but it has been implicated in lipid droplet biogenesis,

nuclear envelope autophagy, amino acid metabolism and a specialized form of autophagy called

Piecemeal Microautophagy of the Nucleus (PMN), where portions of the nucleus invaginate into

the vacuole [61-66]. We assessed the co-localization of the JUNQ and the NVJ by expressing NES-

LuciTs or Ubc9Ts in cells carrying a chromosomally tagged Nvj1-GFP [53]. Live-cell time-lapse

microscopy revealed transient and dynamic interactions between JUNQ and Nvj1, whereby both

puncta came in contact and then separated (Fig. 5a; Extended Data Fig. 5a; e.g. observe 9 min to

11 min in Supplemental Video 9).

The architecture of perinuclear spatial PQC and the close association of the INQ and JUNQ in

relation to the NVJ were next examined using SIM super-resolution microscopy (Fig. 5b-e;

Supplemental Videos 10, 11). The NVJ, visualized via Nvj1 imaging, forms a basket-shaped

structure contouring the Nuclear Envelope and connected to the vacuole (Fig. 5d). In these super-

resolution images, the JUNQ, visualized by NES-LuciTs, was proximal to the cytoplasmic side of

the NVJ as observed by Cryo-SXT (Fig. 5b). Of note, the Q-body "assembly" substructure of the

JUNQ was evident in these images. On the other hand, the nuclear INQ, visualized by NLS-

LuciTs, was nestled between the concave nuclear side of the NVJ and the nuclear DNA (Fig. 5c).

Strikingly, we were able to observe in some cells the INQ being extruded through the NVJ to enter

the vacuole (Fig. 5e; Supplemental Video 12). These experiments suggest dynamic tethering of

the INQ and JUNQ to the NVJ vicinity account for the homing of these inclusions as well as the

slight fluctuations in distance between these inclusions observed in particle tracking experiments

in Fig. 3.

We next tested if NVJs play a role in INQ and JUNQ clearance using *nvj1Δ* or *vac8Δ* yeast cells.

Live-cell time course analyses provided an unexpected insight into NVJ function. In *nvj1Δ* cells,

nuclear and diffuse NLS-LuciTs formed an initial nuclear inclusion, but during the time-course,

some cells showed the INQ being egressed from the nucleus into the cytoplasm (Fig. 5f).

Strikingly, some of the egressed NLS-LuciTs puncta ended up co-localizing with NES-LuciTs

(Fig. 5f, inset; Supplemental Video 13; white arrowheads). Similar results were obtained when

cycloheximide was added together to the temperature shift (Extended Data Fig. 5b) to ensure all

the NLS-LuciTs is in the nucleus. We observed NLS-LuciTs extruded from the nucleus in the

njv1Δ but not the WT cells, confirming NVJ disruption leads to nuclear extrusion of the INQ.

Confocal microscopy in *nvj1Δ* and *vac8Δ* cells confirmed that NVJ disruption led to extrusion of

nuclear NLS-LuciTs into the cytosol, where it ended up co-localizing with NES-LuciTs puncta

(Fig. 5g; white arrowheads). Of note, deleting the NJVs does not disrupt INQ and JUNQ

convergence, as the INQ and JUNQ are still juxtaposed in both *nvj1Δ* and *vac8Δ* cells (Fig. 5g).

We next examined if deletion of the NVJ affects clearance of PQC inclusions. Indeed, we observed

dramatic increases in the fraction of cells containing NLS and NES-LuciTs inclusions in

nvj1Δ and *vac8Δ* cells compared to WT cells (Fig. 5h, panels *i, ii*; Extended Data Fig. 5c,d), as

well as the levels of misfolded protein (Extended Data Fig. 5e). MG132 addition led to a

synergistic enhancement in the number of cells with NES-LuciTs inclusions, suggesting the NVJ

and the proteasome represent distinct cytoplasmic clearance pathways (Extended Data Fig. 5c).

The synergy effect for NLS-LuciTs inclusions was marginal, perhaps because the higher fraction

of cells with INQs poses a ceiling effect to our measurements. Alternatively, the NVJ may play a

greater role in the clearance of nuclear protein inclusions (Extended Data Fig. 5b). Further

supporting a role in misfolded protein clearance, NVJ deletion sensitizes cells to proteotoxic stress

(Fig. 5i). Both *nvj1Δ* and *vac8Δ* cells are sensitive to treatment with proline analog AZC, which

induces widespread misfolding of newly translated proteins (Fig. 5i).

The findings that the INQ and JUNQ converge on nuclear-vacuolar contact sites and that the NVJ

is required for efficient clearance of nuclear and cytoplasmic inclusions define nuclear-vacuolar

contacts as cellular hubs for spatial protein quality control.

ESCRT proteins are required for INQ-JUNQ homing and clearance at the vicinity of NVJs

The relationship observed between spatial PQC and the nuclear-vacuolar contact sites led us to

examine the role of the perinuclear ESCRT-II/III protein Chm7 (CHMP7 in mammals) which

participates in nuclear envelope sealing and nuclear pore quality control [67-69] (Fig. 6a, *i*).

In yeast, Chm7 normally looks diffuse throughout the cytoplasm (Extended Data Fig. 6a) and

becomes associated with a unique site in the nuclear envelope upon activation. The active

conformation of Chm7 can be induced by deleting auto-inhibitory helices in the ESCRT-III

domain [herein Chm7$_{OPEN}$] [69, 70]. Strikingly, activated Chm7$_{OPEN}$ localized to a single site in the

nuclear envelope in close proximity with both INQ and JUNQ (Fig. 6a, *ii*), suggesting that this

perinuclear ESCRT protein marks the site of INQ and JUNQ convergence for vacuolar delivery.

We next examined whether loss of Chm7 impacts INQ and JUNQ subcellular localization. In

chm7Δ yeast cells, both INQ and JUNQ still formed and migrated to the perinuclear region, but

they no longer converged to the same location (Fig. 6b. These results reveal a novel role for the

Chm7 ESCRT protein as recruiting nuclear and cytoplasmic PQC compartments to a specific

location on the nuclear envelope, either for further clearance or terminal sequestration.

Since Chm7 is an ESCRT II/III protein, we next examined the role of other ESCRT proteins in

spatial PQC: the ESCRT-I Vps23 and the membrane bound phosphatidylinositol 3-kinase (PI3K)

complexVps34-Vps15. Vps23 is a component of the ESCRT-I complex that binds to ubiquitinated

cargo proteins and mediates their vacuolar transport [71-73]. The Vps34-Vps15 complex is essential

for protein sorting to the vacuole and has been shown to localize to nuclear pores at NVJs [74-77].

We co-expressed NLS- and NES-LuciTs in either WT, *vps23Δ*, *vps34Δ*, and *vps15Δ* cells, and

followed their fate by inducing unfolding at 37 °C. While INQ and JUNQ formed in both WT and

ESCRT mutants, the spatial relationship between INQ and JUNQ was disrupted in the ESCRT

deletions. All three ESCRT mutants showed similar aberrant localization phenotypes which

included cells in which the INQ and JUNQ were no longer homed as well as cells displaying

cytoplasmic egress of the INQ (Fig. 6c, Extended Data Fig. 6b). Strikingly, in a subset of *vps23Δ*

and *vps34Δ* cells we observed a possible intermediate for nuclear egress of the INQ, whereby the

NLS-LuciTs inclusion was fully enveloped by the nuclear membrane in what appears to be a

budding event (Fig. 6c, Extended Data Fig. 6b). These data suggest that the ESCRT pathway plays

a role in homing of INQ and JUNQ to the NVJ as well as extrusion of the INQ, likely to mediate

vacuolar delivery and clearance. Furthermore, when this process is perturbed, the INQ can

sometimes bud into the cytoplasm.

We next asked if this perinuclear ESCRT pathway promotes vacuolar clearance of the INQ and

JUNQ. Since vacuolar engulfment often requires the AAA-ATPase Vps4, which functions by

disassembling ESCRT-III complexes [78, 79] (Fig. 6d), we examined the effect of deleting Vps4 on

INQ and JUNQ clearance as described above. Indicating that Vps4 is required for clearance of

both INQ and JUNQ, *vps4Δ* cells contained more NLS- and NES-LuciTs inclusions than WT

controls (Fig. 6e). Strikingly, NLS-LuciTs also egressed into the cytoplasm in some *vps4Δ* cells

(Fig. 6f), supporting the notion Vps4 mediates INQ budding into the vacuole. Together, these

results indicate that nuclear misfolded proteins sequestered in the INQ are homed to the nuclear

side of NVJ to be extruded out of the nucleus and into the vacuole in an ESCRT family protein-

dependent manner.

Vacuolar clearance of proteins in INQ and JUNQ

To further examine vacuolar targeting of nuclear and cytoplasmic inclusions in yeast we utilized

the pH sensitive protein sensor Keima (Fig. 7a). Keima has a pH-sensitive excitation spectrum,

with a maximum of 440 nm at neutral pH and a maximum of 558 nm in acidic environments such

as the vacuole [80, 81], whereas its emission spectrum has a pH-insensitive maximum at 620 nm [80].

Since Keima in yeast has a weak emission intensity and is easily bleached [81], we used N-terminal

tagging with two consecutive Keima moieties to visualize NLS- and NES-LuciTs (herein

2xKeima-LuciTS; Fig. 7a, b, schematic). In the folded state, both NLS-LuciTS and NES-LuciTS

proteins were diffusely located in the nucleus and the cytoplasm respectively, with the excitation

wavelength characteristic of a neutral pH environment. Upon misfolding at 37 °C, both NLS-

LuciTs-Keima and NES-LuciTs-Keima formed the nuclear INQ and cytoplasmic JUNQ

respectively maintaining the neutral pH environment maximum emission after 404 nm excitation,

suggesting that inclusion formation takes place initially in their respective subcellular location.

However, continued incubation at 37 °C yielded regions of characteristic acidic environment

emission at 620 nm following 558 nm excitation appearing in a time-dependent manner for both

inclusions (Fig. 7c, d, Extended Data Fig. 7a, b). The kinetics of acidification of the JUNQ was

slower than for the INQ, consistent with its slower clearance kinetics (Extended Data Fig. 1a). For

the INQ, the red shift appears to initiate at the nuclear envelope boundary on what seems to be

almost full co-localization, with an elongated structure observed extending from the inclusion and

entering the vacuole on some cells (Fig. 7c). The slower degradation kinetics of the JUNQ coupled

with the low quantum yield and quick photobleaching of Keima precluded continuous live-

imaging of the JUNQ, but an imaging time course of different regions on the same slide showed

also areas of almost full colocalization and also multiple red only foci arising proximal to the

JUNQ (Fig. 7d; Extended Data Fig. 7a-b). Of note, these experiments were carried out under

conditions of ongoing proteasome degradation indicating that under these experimental conditions

vacuolar targeting seems to occur in parallel to UPS-mediated clearance of misfolded proteins.

The contribution of vacuolar and proteasomal pathways in the clearance of misfolded nuclear and

cytoplasmic proteins was independently assessed using immunoblot analyses (Fig. 7e-f). Briefly,

GFP-tagged NLS- and NES-LuciTs were expressed under permissive conditions, followed by

glucose repression, as described in Fig 1. Following unfolding, the levels of remaining luciferase

were examined after 2 and 4 hrs at 37 °C by immunoblot analyses probing for the GFP moiety.

The contribution of proteasomal clearance was assessed by addition of 50 µM Bortezomib [27, 82]

(Bz) at T=0, while the contribution of vacuolar clearance was assessed using cells deleted for Pep4,

required for maturation of vacuolar proteinases, supplemented with 1mM PMSF to inhibit vacuolar

serine proteases [83]. As expected, the levels of full length nuclear and cytoplasmic luciferase

decreased as a function of time, and were partially stabilized by proteasomal inhibition (Figure 7

e,f; Extended Data Fig. 7c-e). Inhibition of vacuolar proteases in the *pep4Δ* cells stabilized both

the nuclear and cytoplasmic full-length proteins [84, 85](Figure 7c-e; Extended Data Fig. 7c, e).

Interestingly, we also observed a range of GFP-positive degradation products in the WT and +Bz

cells that disappear when the vacuolar proteases were inhibited (Figure 7g), which is a hallmark

of vacuolar targeting of GFP-tagged proteins [64, 65, 83, 86-89]. Quantification of the intensity of

degradation fragments relative to initial full-length protein confirmed the reduction in clearance

when vacuolar proteases are inhibited (Fig. 7g). Moreover, a longer exposure reveals fewer bands

and a completely different banding pattern between WT and protease inhibited cells (Extended

Data Fig. 7c).

The finding that inhibiting vacuolar proteases stabilizes NLS-LuciTs supports the existence of a

pathway to transfer nuclear PQC substrates to the vacuole. To independently corroborate this, we

next carried out live-cell time resolved imaging of the dynamic clearance of the INQ through the

vacuole (Fig. 7g). The INQ was visualized via NLS-LuciTs-GFP and the vacuole using the styryl

dye FM4-64 commonly used to visualize endocytosis and vacuolar membrane dynamics [90] (Fig.

7g; Supplemental Video 14). Since GFP fluorescence is quenched by the acidic vacuolar

environment, NLS-LuciTS[84, 85, 91, 92] (Fig. 7g; supplemental video 14; Extended Data Fig. 7d). In

WT cells, the INQ is opposed to the vacuole at the start of the time course and appears to be pulled

into the vacuole as a function of time. Fascinatingly, when the nuclear-vacuolar contact is

disrupted in *nvj1Δ,* the INQ is unable to interact with and enter the vacuole and moves dynamically

without becoming degraded (Fig. 7g, second row; Supplemental Video 15). Strikingly, in the vast

majority of *vps4Δ* cells, the INQ looks still proximal to the vacuole but does not get cleared,

supporting a role for Vps4-mediated membrane scission for INQ vacuolar engulfment (Fig 7g,

third row; Supplemental Video 16).

DISCUSSION

Spatial sequestration of misfolded proteins into membrane-less inclusions is an integral step in

PQC. Here, provide insights into the logic of spatial PQC. We demonstrate that cytoplasmic

misfolded proteins are sequestered into a cytoplasmic perinuclear JUNQ compartment, while

nuclear misfolded proteins are sequestered into a nuclear peri-nucleolar INQ compartment (Figure

8a, b). Previous studies identified distinct nuclear and cytoplasmic PQC pathways that target

proteins that misfold in these compartments for ubiquitin-dependent proteasomal degradation in

the nucleus and cytoplasm respectively [27, 93] (Figure 8b). Supporting the concept that nuclear and

cytoplasmic proteins are cleared in the compartment where they misfold, these studies identified

distinct ubiquitin ligases and ubiquitin linkage requirements in the nucleus and cytoplasm that

promote PQC [27, 94]. We find spatial sequestration is also compartment specific, with nuclear and

cytoplasmic inclusions forming in the nucleus and cytoplasm independently. These distinct PQC

compartments converge to face each other across the nuclear envelope at a site proximal to the

Nuclear-Vacuolar Junctions (NVJ) (Figure 8c). The homing process seems to be coordinated by

nuclear pores and a perinuclear ESCRT pathway involving ESCRT II/III protein Chm7 (Figure

8c). We further propose that proteins sequestered in both JUNQ and INQ undergo Vps4-dependent

vacuolar degradation (Figure 8c, inset).

Cryo-SXT and super-resolution imaging indicate the JUNQ is not a homogeneous inclusion, but

instead consists of an assembly of dynamic membrane-less PQC inclusions called Q-bodies[54, 95],

which form in a chaperone-dependent manner early during misfolding. Previously shown to be

proximal to the ER, we find they are also close to mitochondria[2,9]. It will be of interest to assess

how Q-bodies congregate to form the JUNQ, but perhaps the pathway is akin to the Syntaxin 5-

dependent pathway described to traffic Hsp104-associated aggregates [20]. Notably, the Q-bodies

and JUNQ exhibit a distinct linear absorption coefficient in cryo-SXT data (Figure 4b-f),

suggesting they may be liquid-like phase-separated entities [54].

Spatial sorting of misfolded proteins into inclusions is proposed to serve to sequester toxic species

away from the cellular milieu and enhance cellular fitness [17, 20, 96]. The convergence of INQ and

JUNQ to nuclear-vacuolar contacts may additionally serve to connect spatial PQC to vacuolar

clearance, thus providing an alternate route to proteasomal degradation. Of note, vacuolar protein

clearance of proteins in both INQ and JUNQ is seen without proteasome inhibitors, as shown using

Keima fusions, as well as by immunoblot analyses in *pep4Δ* or *vps4Δ* cells (Figures 1c-d, 2a, c-g,

5h, 7c-g). It thus appears that proteasomal and vacuolar clearance do not necessarily function

hierarchically for spatial PQC. This raises interesting questions for future studies, including

whether they can act on the same pool of PQC substrates, or select their substrates based on distinct

chaperone and/or ubiquitin signals [93].

Our observation that the INQ can be extruded from the nucleus via the NVJ, indicates a path to

clear the nuclear compartment from misfolded proteins even when the proteasome is overloaded

or otherwise impaired. The physical tether provided by NVJs previously shown to mediate

engulfment of regions of the nuclear membrane into the vacuole, similarly mediates egress of the

INQ into the vacuole. This could explain how NVJ disruption leads to INQ extrusion into the

cytoplasm. The perinuclear ESCRT-II/III protein Chm7 (CHMP7 in mammals), is involved in

this process. Chm7 participates in recruiting nuclear and cytoplasmic PQC compartments to a

unique location on the nuclear envelope. While Chm7 is normally diffuse throughout the

cytoplasm (Extended Data Fig. 5e). Upon activation Chm7 becomes associated with a unique site

in the nuclear envelope upon activation. While the active conformation can be induced by deleting

auto-inhibitory helices in the ESCRT-III domain [69], we did not yet identify the mode of activation

of Chm7 by PQC inclusions. CHMP7 was previously shown to function in remodeling nuclear

envelopes [67], clearance of misassembled nuclear pores [68], egress of herpes virus from the nucleus

to the cytosol [97], as well as nuclear envelope sealing and nuclear pore quality control [69]. Our data

provides an intriguing link between protein quality control and nuclear envelope quality control.

Future studies should determine whether additional forms of microautophagy, such as

nucleophagy, ERphagy, piecemeal microautophagy of the nucleus, or selective autophagy of

nuclear pore complexes participate in clearance of these PQC compartments.

The role for nuclear pores, nuclear-vacuolar contacts and the nuclear envelope in communicating

nucleocytoplasmic protein quality control decisions between the nucleus, the cytoplasm and the

endolysosomal system places spatial PQC in the larger context of cellular organization and inter-

organellar communication [98]. This idea is particularly intriguing given the link between nuclear

pore dysfunction and aging, lamin progeria mutants and neurodegenerative diseases such as

Huntington's disease, Amyotrophic Lateral Sclerosis and Frontal Temporal Dementia, which are

all characterized by nuclear pore dysfunction and altered nucleocytoplasmic trafficking [31, 34, 35, 99-

102]. Our findings raise the possibility that compromising nucleocytoplasmic trafficking leads to

impaired misfolded protein clearance contributing to further cellular dysfunction. A better

understanding of these pathways may provide fundamental insights into the critical biological

process of protein quality control and point to avenues to ameliorate a large range of human

disorders.

MAIN REFERENCES

1. Hartl, F.U., Bracher, A. & Hayer-Hartl, M. Molecular chaperones in protein folding and
proteostasis. *Nature* **475**, 324-332 (2011).
2. Needham, P.G., Guerriero, C.J. & Brodsky, J.L. Chaperoning Endoplasmic Reticulum-
Associated Degradation (ERAD) and Protein Conformational Diseases. *Cold Spring*
Harb Perspect Biol **11** (2019).
3. Balchin, D., Hayer-Hartl, M. & Hartl, F.U. In vivo aspects of protein folding and quality
control. *Science* **353**, aac4354 (2016).
4. Sontag, E.M., Samant, R.S. & Frydman, J. Mechanisms and Functions of Spatial Protein
Quality Control. *Annu Rev Biochem* (2017).
5. Pohl, C. & Dikic, I. Cellular quality control by the ubiquitin-proteasome system and
autophagy. *Science* **366**, 818-822 (2019).
6. Finley, D. & Prado, M.A. The Proteasome and Its Network: Engineering for Adaptability.
Cold Spring Harb Perspect Biol **12** (2020).
7. Alberti, S. & Hyman, A.A. Are aberrant phase transitions a driver of cellular aging?
Bioessays (2016).
8. Ano Bom, A.P. *et al.* Mutant p53 aggregates into prion-like amyloid oligomers and
fibrils: implications for cancer. *J Biol Chem* **287**, 28152-28162 (2012).
9. Douglas, P.M. & Dillin, A. Protein homeostasis and aging in neurodegeneration. *J Cell*
Biol **190**, 719-729 (2010).
10. De Mattos, E.P. *et al.* Protein Quality Control Pathways at the Crossroad of
Synucleinopathies. *J Parkinsons Dis* **10**, 369-382 (2020).
11. Kaganovich, D., Kopito, R. & Frydman, J. Misfolded proteins partition between two
distinct quality control compartments. *Nature* **454**, 1088-1095 (2008).
12. Dillin, A. & Cohen, E. Ageing and protein aggregation-mediated disorders: from
invertebrates to mammals. *Philos Trans R Soc Lond B Biol Sci* **366**, 94-98 (2011).
13. Balch, W.E., Morimoto, R.I., Dillin, A. & Kelly, J.W. Adapting proteostasis for disease
intervention. *Science* **319**, 916-919 (2008).
14. Reichmann, D., Voth, W. & Jakob, U. Maintaining a Healthy Proteome during Oxidative
Stress. *Mol Cell* **69**, 203-213 (2018).
15. Morales-Polanco, F., Lee, J.H., Barbosa, N.M. & Frydman, J. Cotranslational
Mechanisms of Protein Biogenesis and Complex Assembly in Eukaryotes. *Annu Rev*
Biomed Data Sci **5**, 67-94 (2022).

16.   Specht, S., Miller, S.B., Mogk, A. & Bukau, B. Hsp42 is required for sequestration of
544   protein aggregates into deposition sites in Saccharomyces cerevisiae. *J Cell Biol* **195**,
545   617-629 (2011).
546   17.   Escusa-Toret, S., Vonk, W.I. & Frydman, J. Spatial sequestration of misfolded proteins
547   by a dynamic chaperone pathway enhances cellular fitness during stress. *Nat Cell Biol*
548   (2013).
549   18.   Miller, S.B. *et al.* Compartment-specific aggregases direct distinct nuclear and
550   cytoplasmic aggregate deposition. *EMBO J* **34**, 778-797 (2015).
551   19.   Weisberg, S.J. *et al.* Compartmentalization of superoxide dismutase 1 (SOD1G93A)
552   aggregates determines their toxicity. *Proc Natl Acad Sci U S A* **109**, 15811-15816 (2012).
553   20.   Babazadeh, R. *et al.* Syntaxin 5 Is Required for the Formation and Clearance of Protein
554   Inclusions during Proteostatic Stress. *Cell Rep* **28**, 2096-2110 e2098 (2019).
555   21.   Nollen, E.A. *et al.* Dynamic changes in the localization of thermally unfolded nuclear
556   proteins associated with chaperone-dependent protection. *Proc Natl Acad Sci U S A* **98**,
557   12038-12043 (2001).
558   22.   Yasuda, S. *et al.* Stress- and ubiquitylation-dependent phase separation of the
559   proteasome. *Nature* **578**, 296-300 (2020).
560   23.   Polling, S. *et al.* Misfolded polyglutamine, polyalanine, and superoxide dismutase 1
561   aggregate via distinct pathways in the cell. *J Biol Chem* **289**, 6669-6680 (2014).
562   24.   Park, J.H. *et al.* Amyotrophic lateral sclerosis-related mutant superoxide dismutase 1
563   aggregates inhibit 14-3-3-mediated cell survival by sequestration into the JUNQ
564   compartment. *Hum Mol Genet* **26**, 3615-3629 (2017).
565   25.   McClellan, A.J., Scott, M.D. & Frydman, J. Folding and quality control of the VHL
566   tumor suppressor proceed through distinct chaperone pathways. *Cell* **121**, 739-748
567   (2005).
568   26.   Gupta, R. *et al.* Firefly luciferase mutants as sensors of proteome stress. *Nat Methods* **8**,
569   879-884 (2011).
570   27.   Samant, R.S., Livingston, C.M., Sontag, E.M. & Frydman, J. Distinct proteostasis circuits
571   cooperate in nuclear and cytoplasmic protein quality control. *Nature* **563**, 407-411
572   (2018).
573   28.   Popken, P., Ghavami, A., Onck, P.R., Poolman, B. & Veenhoff, L.M. Size-dependent
574   leak of soluble and membrane proteins through the yeast nuclear pore complex. *Mol Biol
575   Cell* **26**, 1386-1394 (2015).
576   29.   Amm, I. & Wolf, D.H. Molecular mass as a determinant for nuclear San1-dependent
577   targeting of misfolded cytosolic proteins to proteasomal degradation. *FEBS Lett* **590**,
578   1765-1775 (2016).
579   30.   Timney, B.L. *et al.* Simple rules for passive diffusion through the nuclear pore complex.
580   *J Cell Biol* **215**, 57-76 (2016).
581   31.   Woerner, A.C. *et al.* Cytoplasmic protein aggregates interfere with nucleocytoplasmic
582   transport of protein and RNA. *Science* **351**, 173-176 (2016).
583   32.   Park, S.H. *et al.* PolyQ Proteins Interfere with Nuclear Degradation of Cytosolic Proteins
584   by Sequestering the Sis1p Chaperone. *Cell* **154**, 134-145 (2013).
585   33.   Neuber, A. *et al.* Nuclear export receptor Xpo1/Crm1 is physically and functionally
586   linked to the spindle pole body in budding yeast. *Mol Cell Biol* **28**, 5348-5358 (2008).
587   34.   Grima, J.C. *et al.* Mutant Huntingtin Disrupts the Nuclear Pore Complex. *Neuron* **94**, 93-
588   107 e106 (2017).

35.   Gasset-Rosa, F. *et al.* Polyglutamine-Expanded Huntingtin Exacerbates Age-Related
Disruption of Nuclear Integrity and Nucleocytoplasmic Transport. *Neuron* **94**, 48-57 e44
(2017).
36.   Wente, S.R. & Blobel, G. A temperature-sensitive NUP116 null mutant forms a nuclear
envelope seal over the yeast nuclear pore complex thereby blocking nucleocytoplasmic
traffic. *J Cell Biol* **123**, 275-284 (1993).
37.   Chen, L. *et al.* Sts1 plays a key role in targeting proteasomes to the nucleus. *J Biol Chem*
286, 3104-3118 (2011).
38.   Takeda, K. & Yanagida, M. Regulation of nuclear proteasome by Rhp6/Ubc2 through
ubiquitination and destruction of the sensor and anchor Cut8. *Cell* **122**, 393-405 (2005).
39.   Tatebe, H. & Yanagida, M. Cut8, essential for anaphase, controls localization of 26S
proteasome, facilitating destruction of cyclin and Cut2. *Curr Biol* **10**, 1329-1338 (2000).
40.   Romero-Perez, L., Chen, L., Lambertson, D. & Madura, K. Sts1 can overcome the loss of
Rad23 and Rpn10 and represents a novel regulator of the ubiquitin/proteasome pathway.
J Biol Chem **282**, 35574-35582 (2007).
41.   Lord, C.L. & Wente, S.R. Nuclear envelope-vacuole contacts mitigate nuclear pore
complex assembly stress. *J Cell Biol* **219** (2020).
42.   Crisp, M. *et al.* Coupling of the nucleus and cytoplasm: role of the LINC complex. *J Cell*
Biol **172**, 41-53 (2006).
43.   Aitchison, J.D., Blobel, G. & Rout, M.P. Nup120p: a yeast nucleoporin required for NPC
distribution and mRNA transport. *J Cell Biol* **131**, 1659-1675 (1995).
44.   Heath, C.V. *et al.* Nuclear pore complex clustering and nuclear accumulation of poly(A)+
RNA associated with mutation of the Saccharomyces cerevisiae RAT2/NUP120 gene. *J*
Cell Biol **131**, 1677-1697 (1995).
45.   Ader, C. *et al.* Amyloid-like interactions within nucleoporin FG hydrogels. *Proc Natl*
Acad Sci U S A **107**, 6281-6285 (2010).
46.   Strawn, L.A., Shen, T., Shulga, N., Goldfarb, D.S. & Wente, S.R. Minimal nuclear pore
complexes define FG repeat domains essential for transport. *Nat Cell Biol* **6**, 197-206
(2004).
47.   Parkinson, D.Y. *et al.* Nanoimaging cells using soft X-ray tomography. *Methods Mol*
Biol **950**, 457-481 (2013).
48.   Le Gros, M.A. *et al.* Biological soft X-ray tomography on beamline 2.1 at the Advanced
Light Source. *J Synchrotron Radiat* **21**, 1370-1377 (2014).
49.   Larabell, C.A. & Nugent, K.A. Imaging cellular architecture with X-rays. *Curr Opin*
Struct Biol **20**, 623-631 (2010).
50.   Le Gros, M.A., McDermott, G., Uchida, M., Knoechel, C.G. & Larabell, C.A. High-
aperture cryogenic light microscopy. *J Microsc* **235**, 1-8 (2009).
51.   Ekman, A. *et al.* Putting Molecules in the Picture: Using Correlated Light Microscopy
and Soft X-Ray Tomography to Study Cells, in *Synchrotron Light Sources and Free-*
Electron Lasers: Accelerator Physics, Instrumentation and Science Applications. (eds. E.
Jaeschke, S. Khan, J.R. Schneider & J.B. Hastings) 1-32 (Springer International
Publishing, Cham; 2019).
52.   Ekman, A.A. *et al.* Mesoscale imaging with cryo-light and X-rays: Larger than molecular
machines, smaller than a cell. *Biol Cell* **109**, 24-38 (2017).
53.   Huh, W.K. *et al.* Global analysis of protein localization in budding yeast. *Nature* **425**,
686-691 (2003).

54.   Wu, G.H. *et al.* Multi-scale 3D Cryo-Correlative Microscopy for Vitrified Cells.
636   *Structure* **28**, 1231-1237.e1233 (2020).

637   55.   Schuler, M.H. *et al.* Miro1-mediated mitochondrial positioning shapes intracellular
638   energy gradients required for cell migration. *Mol Biol Cell* **28**, 2159-2169 (2017).

639   56.   Frottin, F. *et al.* The nucleolus functions as a phase-separated protein quality control
640   compartment. *Science* **365**, 342-347 (2019).

641   57.   Velazquez, J.M. & Lindquist, S. hsp70: nuclear concentration during environmental
642   stress and cytoplasmic storage during recovery. *Cell* **36**, 655-662 (1984).

643   58.   Falahati, H., Pelham-Webb, B., Blythe, S. & Wieschaus, E. Nucleation by rRNA Dictates
644   the Precision of Nucleolus Assembly. *Curr Biol* **26**, 277-285 (2016).

645   59.   Azkanaz, M. *et al.* Protein quality control in the nucleolus safeguards recovery of
646   epigenetic regulators after heat shock. *Elife* **8** (2019).

647   60.   Pan, X. *et al.* Nucleus-vacuole junctions in Saccharomyces cerevisiae are formed through
648   the direct interaction of Vac8p with Nvj1p. *Mol Biol Cell* **11**, 2445-2457 (2000).

649   61.   Kohlwein, S.D. *et al.* Tsc13p is required for fatty acid elongation and localizes to a novel
650   structure at the nuclear-vacuolar interface in Saccharomyces cerevisiae. *Mol Cell Biol* **21**,
651   109-125 (2001).

652   62.   Levine, T.P. & Munro, S. Dual targeting of Osh1p, a yeast homologue of oxysterol-
653   binding protein, to both the Golgi and the nucleus-vacuole junction. *Mol Biol Cell* **12**,
654   1633-1644 (2001).

655   63.   Kvam, E. & Goldfarb, D.S. Structure and function of nucleus-vacuole junctions: outer-
656   nuclear-membrane targeting of Nvj1p and a role in tryptophan uptake. *J Cell Sci* **119**,
657   3622-3633 (2006).

658   64.   Kvam, E. & Goldfarb, D.S. Nucleus-vacuole junctions and piecemeal microautophagy of
659   the nucleus in S. cerevisiae. *Autophagy* **3**, 85-92 (2007).

660   65.   Roberts, P. *et al.* Piecemeal microautophagy of nucleus in Saccharomyces cerevisiae.
661   *Mol Biol Cell* **14**, 129-141 (2003).

662   66.   Krick, R. *et al.* Piecemeal microautophagy of the nucleus requires the core
663   macroautophagy genes. *Mol Biol Cell* **19**, 4492-4505 (2008).

664   67.   von Appen, A. *et al.* LEM2 phase separation promotes ESCRT-mediated nuclear
665   envelope reformation. *Nature* **582**, 115-118 (2020).

666   68.   Thaller, D.J. & Patrick Lusk, C. Fantastic nuclear envelope herniations and where to find
667   them. *Biochem Soc Trans* **46**, 877-889 (2018).

668   69.   Webster, B.M. *et al.* Chm7 and Heh1 collaborate to link nuclear pore complex quality
669   control with nuclear envelope sealing. *EMBO J* **35**, 2447-2467 (2016).

670   70.   Thaller, D.J. *et al.* An ESCRT-LEM protein surveillance system is poised to directly
671   monitor the nuclear envelope and nuclear transport system. *Elife* **8** (2019).

672   71.   Babst, M., Odorizzi, G., Estepa, E.J. & Emr, S.D. Mammalian tumor susceptibility gene
673   101 (TSG101) and the yeast homologue, Vps23p, both function in late endosomal
674   trafficking. *Traffic* **1**, 248-258 (2000).

675   72.   Katzmann, D.J., Babst, M. & Emr, S.D. Ubiquitin-dependent sorting into the
676   multivesicular body pathway requires the function of a conserved endosomal protein
677   sorting complex, ESCRT-I. *Cell* **106**, 145-155 (2001).

678   73.   Herrador, A., Léon, S., Haguenauer-Tsapis, R. & Vincent, O. A mechanism for protein
679   monoubiquitination dependent on a trans-acting ubiquitin-binding domain. *J Biol Chem*
680   **288**, 16206-16211 (2013).

74. Stack, J.H., Herman, P.K., Schu, P.V. & Emr, S.D. A membrane-associated complex
containing the Vps15 protein kinase and the Vps34 PI 3-kinase is essential for protein
sorting to the yeast lysosome-like vacuole. *EMBO J* **12**, 2195-2204 (1993).
75. Herman, P.K. & Emr, S.D. Characterization of VPS34, a gene required for vacuolar
protein sorting and vacuole segregation in Saccharomyces cerevisiae. *Mol Cell Biol* **10**,
6742-6754 (1990).
76. Burda, P., Padilla, S.M., Sarkar, S. & Emr, S.D. Retromer function in endosome-to-Golgi
retrograde transport is regulated by the yeast Vps34 PtdIns 3-kinase. *J Cell Sci* **115**,
3889-3900 (2002).
77. Slessareva, J.E., Routt, S.M., Temple, B., Bankaitis, V.A. & Dohlman, H.G. Activation
of the phosphatidylinositol 3-kinase Vps34 by a G protein alpha subunit at the endosome.
Cell **126**, 191-203 (2006).
78. Henne, W.M., Stenmark, H. & Emr, S.D. Molecular mechanisms of the membrane
sculpting ESCRT pathway. *Cold Spring Harb Perspect Biol* **5** (2013).
79. Hurley, J.H. & Hanson, P.I. Membrane budding and scission by the ESCRT machinery:
it's all in the neck. *Nat Rev Mol Cell Biol* **11**, 556-566 (2010).
80. Katayama, H., Kogure, T., Mizushima, N., Yoshimori, T. & Miyawaki, A. A sensitive
and quantitative technique for detecting autophagic events based on lysosomal delivery.
Chem Biol **18**, 1042-1052 (2011).
81. Bialecka-Fornal, M., Makushok, T. & Rafelski, S.M. A Review of Fluorescent Proteins
for Use in Yeast. *Methods Mol Biol* **1369**, 309-346 (2016).
82. Sitron, C.S., Park, J.H., Giafaglione, J.M. & Brandman, O. Aggregation of CAT tails
blocks their degradation and causes proteotoxicity in S. cerevisiae. *PLoS One* **15**,
e0227841 (2020).
83. Schuck, S., Gallagher, C.M. & Walter, P. ER-phagy mediates selective degradation of
endoplasmic reticulum independently of the core autophagy machinery. *J Cell Sci* **127**,
4078-4088 (2014).
84. Haupts, U., Maiti, S., Schwille, P. & Webb, W.W. Dynamics of fluorescence fluctuations
in green fluorescent protein observed by fluorescence correlation spectroscopy. *Proc Natl
Acad Sci U S A* **95**, 13573-13578 (1998).
85. Shinoda, H., Shannon, M. & Nagai, T. Fluorescent Proteins for Investigating Biological
Events in Acidic Environments. *Int J Mol Sci* **19** (2018).
86. Klionsky, D.J. *et al.* Guidelines for the use and interpretation of assays for monitoring
autophagy (4th edition). *Autophagy* **17**, 1-382 (2021).
87. Eising, S. *et al.* A lysosomal biogenesis map reveals the cargo spectrum of yeast vacuolar
protein targeting pathways. *J Cell Biol* **221** (2022).
88. Waite, K.A., De-La Mota-Peynado, A., Vontz, G. & Roelofs, J. Starvation Induces
Proteasome Autophagy with Different Pathways for Core and Regulatory Particles. *J Biol
Chem* **291**, 3239-3253 (2016).
89. Kvam, E. & Goldfarb, D.S. Nvj1p is the outer-nuclear-membrane receptor for oxysterol-
binding protein homolog Osh1p in Saccharomyces cerevisiae. *J Cell Sci* **117**, 4959-4968
(2004).
90. Vida, T.A. & Emr, S.D. A new vital stain for visualizing vacuolar membrane dynamics
and endocytosis in yeast. *J Cell Biol* **128**, 779-792 (1995).
91. Thorn, K. Genetically encoded fluorescent tags. *Mol Biol Cell* **28**, 848-857 (2017).

92. Costantini, L.M. *et al.* A palette of fluorescent proteins optimized for diverse cellular
environments. *Nat Commun* **6**, 7670 (2015).
93. Schneider, K.L. *et al.* Using reporters of different misfolded proteins reveals differential
strategies in processing protein aggregates. *J Biol Chem* **298**, 102476 (2022).
94. Amm, I., Sommer, T. & Wolf, D.H. Protein quality control and elimination of protein
waste: the role of the ubiquitin-proteasome system. *Biochim Biophys Acta* **1843**, 182-196
(2014).
95. Eisele, F. *et al.* An Hsp90 co-chaperone links protein folding and degradation and is part
of a conserved protein quality control. *Cell Rep* **35**, 109328 (2021).
96. Arrasate, M., Mitra, S., Schweitzer, E.S., Segal, M.R. & Finkbeiner, S. Inclusion body
formation reduces levels of mutant huntingtin and the risk of neuronal death. *Nature* **431**,
805-810 (2004).
97. Kumar, B. *et al.* ESCRT-I Protein Tsg101 Plays a Role in the Post-macropinocytic
Trafficking and Infection of Endothelial Cells by Kaposi's Sarcoma-Associated
Herpesvirus. *PLoS Pathog* **12**, e1005960 (2016).
98. Gottschling, D.E. & Nystrom, T. The Upsides and Downsides of Organelle
Interconnectivity. *Cell* **169**, 24-34 (2017).
99. Zhang, K. *et al.* The C9orf72 repeat expansion disrupts nucleocytoplasmic transport.
Nature **525**, 56-61 (2015).
100. Fallini, C., Khalil, B., Smith, C.L. & Rossoll, W. Traffic jam at the nuclear pore: All
roads lead to nucleocytoplasmic transport defects in ALS/FTD. *Neurobiol Dis* **140**,
104835 (2020).
101. Shang, J. *et al.* Aberrant distributions of nuclear pore complex proteins in ALS mice and
ALS patients. *Neuroscience* **350**, 158-168 (2017).
102. Chou, C.C. *et al.* TDP-43 pathology disrupts nuclear pore complexes and
nucleocytoplasmic transport in ALS/FTD. *Nat Neurosci* **21**, 228-239 (2018).

FIGURE TITLES AND LEGENDS

Figure 1: The INQ and JUNQ are separate nuclear and cytoplasmic PQC compartments.

(a) Schematic showing nuclear and cytoplasmic misfolded proteins could be sequestered into

separate compartments or the cytoplasmic misfolded proteins could be imported into the nucleus

and sequestered into the nuclear PQC compartment INQ. (b) Experimental schematic.

Temperature sensitive Luciferase is properly folded at 25 °C. Heat shock at 37 °C leads to

unfolding and sequestration of the protein. Degradation of the unfolded and sequestered proteins

can be blocked by treatment with the proteasome inhibitor MG132. (c, d) Live-cell time-lapse

fluorescence microscopy of WT cells expressing NLS-LuciTs (c) or NES-LuciTs (d) at 37 °C,

treated with DMSO (top) or 100μM MG132 (bottom). Representative still frames at the times

shown. Scale bars are 1μm. (e, f) Representative SIM images of WT cells expressing NLS-LuciTs

(e) or NLS-VHL (f) after 2 hr at 37 °C and treated with 100μM MG132. NLS-LuciTs and NLS-

VHL are shown in green, nuclear pores in gold, and Hoechst counterstain in blue. Scale bars are

1μm. Line intensity profiles indicate relative locations of subcellular compartments to Nups and

DNA. (g, h) Representative SIM images of WT cells expressing NES-LuciTs (g) and NES-VHL

(h) after 2 hr at 37 °C and treated with 100μM MG132. NES-LuciTs and NES-VHL are shown in

purple, nuclear pores in gold, and Hoechst counterstain in blue. Scale bars are 1μm. Line intensity

profiles indicate relative locations of subcellular compartments to Nups and DNA.

Figure 2: Nuclear entry of misfolded proteins is not required for clearance.

(a) Representative SIM images of WT cells expressing Ubc9Ts-GFP after 2h at 37 °C treated with DMSO (a) or 100µM MG132 (b). Ubc9Ts-GFP is shown in green, nuclear pores in gold, and Hoechst counterstain in blue. Scale bars are 1µm. Line intensity profiles indicate relative locations of subcellular compartments to Nups and DNA. (c) Representative confocal fluorescence microscopy images of cells expressing sGFP with DMSO (control), 100µM MG132 treatment, co-expression of mHTT97QΔP or 200nM Leptomycin B treatment. Ratio of cytoplasmic:nuclear fluorescence is shown in d. Kruskal-Wallis test with Dunn's multiple comparisons test was performed using Prism. Adjusted p value of No Treatment vs. MG132 is 0.0014, No Treatment vs. 97QΔP is 0.0027 and No Treatment vs. LMB is <0.0001. A minimum of 500 cells per condition from 3 biologically independent experiments were analyzed. All scale bars are 1µm. (e) Schematic illustrating the clearance of Ubc9Ts in WT yeast. (*ii*) Timeline of treatments for clearance measurements with shift to 37 °C 60 minutes before initiation of the measurements. (f) Schematic illustrating the *nup116-5* yeast have sealed nuclear pores at 37 °C, thus blocking nucleocytoplasmic trafficking. (*ii*) Densitometric quantification of Western blot bands (*iii*) measuring the amount of Ubc9Ts-EGFP remaining in shut-off experiment of WT vs *nup116*-5 cells relative to $t = 0$ (mean ± s.e.m. from three biologically independent experiments) fitted with a one-phase decay non-linear fit regression line. (g) Schematic illustrating the *sts1-2* yeast do not translocate proteasomes to the nucleus at 37 °C. (*ii*) Densitometric quantification of Western blot bands (iii) measuring the amount of Ubc9Ts-EGFP remaining in shut-off experiment of WT vs sts1-2 cells relative to t = 0 (mean ± s.e.m. from three biologically independent experiments) fitted with a one-phase decay non-linear fit regression line.

Figure 3: INQ and JUNQ home to similar location on each side of the nuclear envelope via a

cytoplasmic signal linked to nuclear pores.

WT cells co-expressing NLS-GFP-LuciTs and NES-DsRed-LuciTs were shifted to 37 °C, treated

with 100μM MG132 and monitored by live cell time-lapse fluorescence. Representative still

frames at the times shown. Scale bar is 1μm. (b) Graph of the distance between the INQ and JUNQ

compartments by particle tracking of inclusions from cell shown in (a) over the time course of the

experiment. The slight variations of tethered distance between 60 and 90 minutes is likely due to

the relative migration of one of the inclusions around a cellular structure as this fluctuation can be

seen at different times post-tethering in other cells (not shown). (c) Representative SIM image

taken of cells co-expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 2 hr at 37 °C and

treated with 100μM MG132. NLS-fusion proteins are shown in green, NES-fusion proteins in

purple, nuclear pores in gold, and Hoechst counterstain in blue. Scale bar is 1μm. Line intensity

profiles indicate relative locations of subcellular compartments and Nups. (d) Representative

confocal microscopy images taken of WT and 3 separate *nup120Δ* yeast cells expressing NES-

LuciTs (*i*), NLS-LuciTs (*ii*) or co-expressing NLS- and NES-LuciTs (*iii*) after 2 hr at 37 °C. NLS-

LuciTs is shown in green, NES-LuciTs in purple, nuclear pores in gold, and Hoechst counterstain

in blue. Scale bar is 1μm. Schematics on the right summarize the findings of the data. (e) Schematic

summarizing the data that the JUNQ localizes to the nuclear pores. A signal is transmitted through

or near the nuclear pores to recruit the INQ to the same location, resulting in the homing of the 2

compartments.

Figure 4: INQ resides near the nucleolus, JUNQ is surrounded by mitochondria, and both compartments home into the Nuclear-Vacuolar Junction.

(a) Unfixed, intact yeast cells are frozen in a capillary tube and imaged by fluorescence microscopy and then X-ray tomography. The linear absorption coefficients are measured from the tomography data and used to identify the subcellular compartments and organelles. The fluorescence images are then correlated to the X-ray tomography data and 3D reconstructions of the subcellular components are generated to determine the cellular context of the PQC compartments. (b) WT cells co-expressing Hsp104-GFP from the endogenous locus and HTT97QP-ChFP after 30 mins at 37 °C. (left) X-ray tomograms without and with fluorescence overlay to indicate sites of Hsp104-GFP positive inclusions and HTT97QP-ChFP inclusion. (right) Table of Linear Absorption Coefficient (LAC) values from the X-ray tomograms used to annotate the images and generate a 3D reconstruction shown in panel c. (c) 3D reconstruction of fluorescence correlated X-ray tomograms showing the JUNQ residing 450nm outside the barrier of the nucleus as defined by the location of the nuclear envelope in the X-ray data (inset, top) and the IPOD marked by mHTT (inset, bottom). JUNQ is shown in green, IPOD in orange, nucleus in yellow, mitochondria in cyan, vacuoles in gray, and lipid droplets in white. Scale bar on X-ray 3D reconstruction is 1μm, insets are 0.5μm. (d) 3D reconstruction of X-ray tomograms from cell expressing NES-LuciTs after 90 mins heat shock at 37 °C and treated with 100μM MG132. The Q-bodies can be seen to coalesce into the JUNQ compartment just outside the nuclear boundary (inset ii). The Q-bodies also interact with vacuoles and the site of coalescence into the JUNQ compartment is surrounded by a mitochondrial cage (inset i). Q-bodies and IPOD are shown in purple, nucleus in yellow, mitochondria in cyan, vacuoles in gray, and lipid droplets in white. Scale bar on X-ray 3D reconstruction is 1μm, insets are 0.5μm. (e) Q-bodies can be seen interacting with mitochondria

in a separate cell expressing NES-LuciTs. Q-bodies are shown in purple, nucleus in yellow,

nucleolus in gold, mitochondria in cyan, and vacuoles in gray. Scale bar on X-ray 3D

reconstruction is 1μm, inset is 0.5μm. (f) Same 3D reconstruction of X-ray tomograms from cell

expressing NES-LuciTs after 90 mins heat shock at 37 °C as shown in d. Insets are from the same

cell rotated 180°. Blue arrows indicate sites where Q-bodies are directly interacting with the

vacuoles. Red arrows indicate sites of nuclear-vacuolar junctions. Scale bar on X-ray 3D

reconstruction is 1μm, insets are 0.5μm. (g) 3D reconstruction of X-ray tomograms from cell

expressing NLS-LuciTs after 90 mins heat shock at 37 °C and treated with 100μ$\underline{M}$ MG132. The

INQ resides 400nm from the nucleolus (inset). INQ is shown in green, nucleus in yellow, nucleolus

in gold, mitochondria in cyan, and vacuoles in gray. Scale bar on X-ray 3D reconstruction is 1μm,

inset is 0.5μm. (h) Schematic illustrating the novel findings of the association between

mitochondria and all 3 cytoplasmic PQC compartments, the proximity of the INQ to the nucleolus,

the interaction between Q-bodies and the vacuole, and the hypothesized location of the INQ and

JUNQ at the nuclear vacuolar junction. The inset highlights two of the component proteins of the

NVJ, Nvj1 and Vac8, that are required for Piecemeal Microautophagy of the Nucleus.

Figure 5: JUNQ and INQ converge at the Nuclear-Vacuolar Junction to facilitate clearance.

(a) Endogenously tagged Nvj1-GFP yeast expressing NES—DsRed-LuciTs were shifted to 37 °C

and monitored by live cell time-lapse fluorescence microscopy for the times shown. White

arrowheads indicate locations of Nvj1 punctum while yellow arrowheads indicate NES-LuciTs

punctum. Scale bar is 1μm. Representative SIM images of WT yeast co-expressing NES-DsRed-

LuciTs (b) or NLS-DsRed-LuciTs (c) and Nvj1-sfGFP after 2 hr incubation at 37 °C with 100μM

MG132. Panel *ii* of both (b) and (c) are insets to better visualize the relative location of the JUNQ

and INQ to the Nvj1. Panel *iii* of both (b) and (c) are additional cells to illustrate other phenotypes

seen in the experiment. NES-DsRed-LuciTs is shown in purple, NLS-DsRed-LuciTs is shown in

green, Nvj1-sfGFP is shown in yellow, and DNA is shown in blue. Scale bars are 1 μm. (d)

Representative confocal image of WT yeast expressing Nvj1-sfGFP (shown in yellow) stained

with FM4-64 vacuolar dye (shown in cyan). Scale bar is 1μm. (e) Representative SIM image of

WT yeast cell co-expressing NLS-DsRed-LuciTs (shown in green) with Nvj1-sfGFP (shown in

yellow) after 2 hr incubation at 37 °C with 100μM MG132. The NLS-LuciTs can be seen extruding

through the NVJ toward the vacuole. Scale bar is 1μm. (f) WT (top) and *nvj1Δ* (bottom) cells co-

expressing NLS-LuciTs and NES-LuciTs were shifted to 37 °C, treated with 100μM MG132 and

monitored by live cell time-lapse fluorescence microscopy for the times shown. White arrowheads

indicate inclusions of NLS-LuciTs pulled into the cytoplasm. Scale bars are 1μm. (g)

Representative confocal images of WT, *nvj1Δ*, and *vac8Δ* yeast co-expressing NLS-EGFP-LuciTs

and NES-DsRed-LuciTs after 2 hr at 37 °C and treated with 100μM MG132. NLS-EGFP-LuciTs

is shown in green, NES-DsRed-LuciTs in purple, nuclear pores in gold, and Hoechst counterstain

in blue. White arrows indicate cytoplasmic localization of NLS-LuciTs. Scale bar is 1μm. (h)

Quantitation of the percentage of cells containing inclusions in WT, *nvj1Δ*, and *vac8Δ* yeast co-

expressing NLS-EGFP-LuciTs (*i*) and NES-DsRed-LuciTs (*ii*) after 2 hr incubation at 37 °C with

100μM MG132. A minimum of 500 cells per condition from 3 biologically independent

experiments were counted and unpaired Student's t-tests were performed comparing the deletion

strains to WT. P values were adjusted using two-stage linear step-up procedure of Benjamini,

Krieger, and Yekutieli with a false discovery rate (Q) of 5%. Adjusted P value for NLS-LuciTs:

WT vs *nvj1Δ* is 0.0205, WT vs *vac8Δ* is 0.0406. Adjusted P value for NES-LuciTs: WT vs *nvj1Δ*

is 0.013, WT vs *vac8Δ* is 0.0008. (i) Drop tests showing serial dilutions of WT, *nvj1Δ*, and *vac8Δ*

yeast at 30 and 37 °C with no treatment and 0.5mg/mL AZC treatment after 72 h growth at

indicated temperature. *nvj1Δ* and *vac8Δ* yeast have a growth defect when subjected to proteostatic

stress by AZC treatment.

Figure 6: ESCRT-mediated extrusion from the nucleus and clearance.

(a) The ESCRT-II/-III protein Chm7 has been shown to play a role in clearance of defective nuclear

pores and nuclear membrane quality control, therefore, it may be involved in the homing of INQ

and JUNQ to the NVJ. (*ii*) Representative confocal images of WT yeast co-expressing $Chm7_{OPEN}$-

EGFP and either NLS-DsRed-LuciTs (top) or NES-DsRed-LuciTs (bottom) after 2 hr at 37 °C and

treated with 100μM MG132. $Chm7_{OPEN}$ is shown in teal, NLS-DsRed-LuciTs in green, NES-

DsRed-LuciTs in purple, nuclear pores in gold and Hoechst counterstain in blue. Arrows indicate

locations of puncta for each protein. Scale bar is 1μm. (b) Representative confocal images of WT

and *chm7Δ* yeast co-expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 120 minutes at

37 °C and treated with 100μM MG132. NLS-EGFP-LuciTs is shown in green, NES-DsRed-

LuciTs in purple, nuclear pores in gold, and Hoechst counterstain in blue. Arrows indicate

locations of puncta for each protein. Scale bar is 1μm. (c) ESCRT-family proteins Vps23, Vps34,

and Vps15 may be involved in clearance of INQ and JUNQ. Representative confocal images of

WT and *vps23Δ* , *vps34Δ,* and *vps15Δ* yeast co-expressing NLS-EGFP-LuciTs and NES-DsRed-

LuciTs after 2 hr at 37 °C and treated with 100μ<u>M</u> MG132. NLS-EGFP-LuciTs is shown in green,

NES-DsRed-LuciTs in purple, nuclear pores in gold, and Hoechst counterstain in blue. Insets show

the budding INQ encapsulated by nuclear pores. Scale bars are 1μm. (d) ESCRTIII proteins and

the ATPase Vps4 are known to remodel membranes and could be involved in vacuolar import of

PQC compartments. (e) Quantitation of the percentage of cells containing inclusions of NLS- or

NES-LuciTs in WT and *vps4Δ* yeast co-expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs

after 2 hr incubation at 37 °C with 100μ<u>M</u> MG132. A minimum of 300 cells per condition from 2

biologically independent experiments were counted and Student's T tests were performed

comparing the deletion strains to WT. P value for WT vs *vps4Δ* NLS-LuciTs is 0.0007, and WT

vs *vps4Δ* NES-LuciTs is 0.0018. (f) Representative confocal images of WT and *vps4Δ* yeast co-

expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 2 hr at 37 °C and treated with 100μ<u>M</u>

MG132. NLS-EGFP-LuciTs is shown in green, NES-DsRed-LuciTs in purple, nuclear pores in

gold, and Hoechst counterstain in blue. Green arrows indicate cytoplasmic localization of NLS-

LuciTs. Scale bar is 1μm.

Figure 7: Vacuolar clearance of the INQ and JUNQ

(a) Possible routes of entry into the vacuole for the INQ and JUNQ. (b) Schematic of the LuciTs-Keima experiments. (c) WT cells expressing NLS-2xKeima-LuciTs after 2 hr incubation at 37 °C with 100μM MG132. Over time, fluorescence is seen with excitation in the 589nm channel indicating the NLS-LuciTs has encountered an acidic environment. Inset shows the transition from green to red and a structure leaving the inclusion that is fully red. Scale bars are 5 uM. (d) Representative images of WT cells expressing NES-2xKeima-LuciTs after 2 hr incubation at 37 °C with 100μM MG132. Over time, fluorescence is seen with excitation in the 589nm channel indicating the NLS-LuciTs has encountered an acidic environment. Insets show the transition from green to red and a structure leaving the inclusion that is fully red. Scale bars are 5 uM. Representative Western Blots of NLS-GFP-LuciTs (e) and NES-GFP-LuciTs (f) in WT and *pep4Δ* yeast. *pep4Δ* yeast were also treated with 1mM PMSF to completely inhibit vacuolar proteases. A decrease in fragments can be seen in both the NLS- (e) and NES- (f) blots. (g) Densitometric quantification of Western blot bands measuring the amount of full-length LuciTs-GFP remaining in shut-off experiment as well as the fragments generated during clearance. The ratio of fragment intensity to full-length protein is shown in the graph (mean ± s.e.m. from three biologically independent experiments). One-way ANOVA was performed using Prism software followed by Dunnett's multiple comparisons test. Adjusted P value for WT vs *pep4Δ* NLS-LuciTs is 0.0018, and WT vs pep*4Δ* NES-LuciTs is 0.0220. (h) WT, *nvj1Δ*, and *vps4Δ* yeast expressing NLS-GFP-LuciTs were treated with 8uM of FM4-64 and incubated for 2hr at 37 °C with 100μM MG132. Cells were imaged every 5 mins for 90 mins. Scale bar is 1μm.

Figure 8: Principles of Nuclear and Cytoplasmic spatial protein quality control Model for vacuolar targeting and clearance of nuclear and cytoplasmic protein inclusions in yeast. (a) Upon heat shock, cytoplasmic and nuclear LuciTs locally misfold, and recruit chaperones and other PQC factors that facilitate the subcellular spatial sequestration on distinct protein quality control inclusions: (b) the Intranuclear Quality Control compartment (INQ) in the nucleus and Q-bodies in the cytoplasm.(c) Q-bodies then coalesce to form the Juxtanuclear Quality Control compartment (JUNQ), and misfolded proteins located in the INQ and the JUNQ can be degraded by the UPS system in the nucleus and the cytoplasm respectively c.ii). (d) Alternatively, under conditions of limited proteasomal activity, INQ and JUNQ can converge on the periphery of the nuclear envelope, a mechanism mediated by the nuclear pores (homing)(e) The homing mechanism could represent a coordinated way to target both inclusions for clearance at the vacuole. (e.ii) Vacuolar mediated clearance of both INQ and JUNQ. ESCRT-family proteins are involved in organizing the INQ and JUNQ for vacuolar degradation.

METHODS

Time resolved live cell imaging. BY4741 Cells were grown overnight in raffinose media followed by 4–6 h at 28 °C in galactose media to induce the expression. Cells were then immobilized on concanavalin A-coated coverslips. Samples were washed in glucose media and kept in the same medium by sealing the coverslips to slides with vacuum grease. When indicated, cells were treated 10 min before sealing with 100 uMof MG132 (Sigma) before sealing. Images were taken every 15 s for 60 min at 37 °C (or room temperature (25 °C)) using a Zeiss Axio Observer.Z1 inverted microscope equipped with X-cite 120 LED light source (Lumen Dynamics), HE GFP/Cy3/DAPI

shift free filter sets (Zeiss), a Plan-Apochromat 100x/1.4 oil DIC M27 objective (Zeiss) and a

digital Axiocam MRm camera (Zeiss) controlled with the Zen blue software.

Cycloheximide chase of preformed inclusions. Cells were grown overnight in raffinose media,

back diluted to an OD600 of 0.1 in a 10 ml of galactose media and grown at 25 °C for 5 hours to

induce expression of both NLS and NES constructs. Cells were then resuspended in 10 ml of

glucose media at 100 μM MG132 and incubated for additional 15 min 25 °C and then switched to

37 °C for 30 minutes to ensure inclusion formation before CHX chase.   Cells were then

immobilized on concanavalin deltaT culture dish (Fischer Scientific), washed with prewarmed

glucose media and incubated with glucose media supplemented with 100 μM MG132 and

50mg/ml of cycloheximide (CHX). Images were taken every 25 s for 120 min at 37 °C as

mentioned above.

FM4-64 vacuole labeling. 10 ml of cells were grown on galactose media to a final OD600 of 0.8

at 25°C. Vacuolar dye was added by resuspending the cells on 2 ml of fresh galactose media plus

FM4-64 to a final concentration of 8uM, and incubated 30 min at 25°C. For imaging, cells were

then resuspended in glucose media at 100uM MG132 and incubated 20 min at 37°C to induce

inclusion formation. Cells were adhered on a concanavalin deltaT culture dish and imaged as

mentioned above.

Immunostaining. As in the live-cell imaging assay, cells were grown overnight in raffinose media

followed by 4–6 h at 28 °C in galactose media to induce the expression. Then, the cells were heat-

shocked at 37 °C in glucose media for 1 h before fixation in 4% paraformaldehyde for 15 mins at

37 °C followed by methanol fixation for 20 mins at -20 °C. Immunostaining proceeded as in [1].

Briefly, samples were resuspended in sorbitol buffer and spheroplasted in Zymolyase (Zymo) for

30 mins at room temperature and further solubilized in 0.1% Triton X-100 for 10 mins. Antibodies

are diluted in Buffer WT (1% nonfat dry milk 0.5 mg/ml BSA 200 mM NaCl 50 mM HEPES–

KOH (pH 7.5) 1 mM NaN3 0.1% Tween-20) at a 1:500 dilution and incubated 2 hr at room

temperature or overnight at 4 °C. Samples are then washed in Buffer WT followed by incubation

in secondary antibodies at a 1:1000 dilution for 2 hr at room temperature. Cells were immobilized

on poly-lysine coated coverslips and mounted in Prolong Diamond mounting media (with DAPI

where indicated) (Thermo-Fisher). Nanobodies against GFP and RFP were conjugated to Alexa

Fluor 488 and Alexa Fluor 568 respectively and used to amplify the signal from the fluorescent

proteins during the SIM measurements. Anti-Nsp1 antibody from EnCor Biotechnology was used

to visualize nuclear pores and Anti-Nsr1 antibody from Abcam was used for staining the nucleolus.

Confocal microscopy. Confocal microscopy on immunostained samples was performed on a

Zeiss LSM 700 inverted confocal microscope with 405, 488, 555 and 639 nm laser lines with Basic

Filterset for LSM700 (Zeiss), a Plan-Apochromat 100x/1.4 oil DIC WD=0.17 objective (Zeiss)

and a digital Axiocam MRm camera (Zeiss) controlled with the Zen black software. Confocal data

sets were deconvolved when indicated using the Zen software with a fast-iterative function.

Fluorescence was quantified where indicated using ImageJ software (NIH). The sGFP intensity

measurements, were performed as described previously [2]. Briefly, the raw integrated density

(herein indicated as intensity) was measured for the nucleus and the whole cell. The intensity value

of the nucleus was then divided by the area. The cytoplasmic intensity was calculated by

subtracting the nuclear intensity from the whole cell intensity and the cytoplasmic area by

subtracting the nuclear area from the whole cell area. The cytoplasmic intensity was then divided

by the cytoplasmic area. Finally, the ratio of cytoplasmic to nuclear was calculated by dividing the

cytoplasmic intensity per area by the nuclear intensity per area. Statistical analysis was performed

using Prism (GraphPad).

Super-resolution microscopy. Structured illumination microscopy (SIM) on immunostained

samples was performed using an OMX V4 Blaze system (Applied Precision, GE Healthcare)

equipped with a 100×/1.40 NA PlanApo oil-immersion objective (Olympus), 405, 488, 568 and

647 nm diode lasers with standard filter sets, and 3 emCCD cameras (Photometrics Evolve 512).

SIM data were acquired with a Z-distance of 125 nm and with 15 images per plane (five phases,

three angles). The raw data was computationally reconstructed using a Wiener (high-frequency)

filter setting of between 0.001 and 0.003 and channel- specific OTFs employing the softWoRx

version 6.5.2 software package (GE Healthcare) to obtain a super-resolution image stack. 3D

reconstructions and line intensity profiles were generated in Volocity version 6.3 (Perkin Elmer).

To construct *NVJ1-sfGFP* strain with integrated sfGFP fusion at the C-terminal, sf*GFP* was

knocked-in via standard PCR-based homologous recombination. The *sfGFP* cassette was

amplified from plasmid GTL-g (Addgene plasmid #81099) with oligonucleotides

5'TAGATGCACAAGTGAACACTGAACAAGCATACTCTCAACCATTTAGATACGCGGC
CGCTCTAGAACTA and

5' TCGCACCTCGTTGTAAGTGACGATGATAACCGAGATGACGGAAATATAGTACA

ATGGAAAAACGCCAGCAACG. The cassette was amplified, separated by electrophoresis, gel
purified and transformed into yeast cells using the lithium acetate method. Correct transformants
were verified by standard PCR using the oligonucleotides NVJ1-VF:
5'GGATACCAGAACAACCTCTTC, NVJ1-VR: 5' ATGCCCGGCGTTATATATTGC, and
LEU2-VF: 5' AGCACGAGCCTCCTTTACCT.

Keima cloning. The NLS-Keima-luciTS plasmid was constructed by replacing the GFP moeity

with yeast optimized Keima using a cassette amplified from plasmid pFA6a-link-yomKeima-

Kan (Addgene plasmid #44902) with the oligonucleotides

5'ATGGCTTCTCCTAAGAAGAAACGTAAAGTTATGGTTTCTGTGATCGCTAAACAAAT

GAC and

5'CCATGCAAGCTTGCGCGGATCCGCGCCCTAATAGAGAATGTCTTGCGATAGC.

Similarly, the NES-Keima-luciTS plasmid was created replacing the GFP moity using the

olinucleotides

5'CTCGCACTTAAGTTCGCCGGTTTAGACCTGATGGTTTCTGTGATCGCTAAACAAAT

GAC and

5'CCATGCAAGCTTGCGCGGATCCGCGCCCTAATAGAGAATGTCTTGCGATAGC. To

construct the NLS and NES-2xKeima-luciTS, an additional Keima moeity was created by

inserting a second keima cassette amplified using the oligonucleotides

5'CCATGCAAGCTTGCGCGGATCCGCGCCCTAATAGAGAATGTCTTGCGATAGC and

5'TTCCATGCAAGCTTGCGCGGATCCGCGCCCTAATAGAGAATGTCTTGCGA.

Live cell LeicaSp8/sSTED microscopy of Keima-tagged inclusions. Cells were grown as for

time resolve live fluorescent microscopy, adhered on concanavalin deltaT culture dish and

examined at <u>37 °C for 90 minutes</u>. Representative cells were collected every 5 minutes on a

Leica TCS SP8 inverted sSTED microscope equipped with a 100x/1.40 APO objective and using

the following detection mirror settings: Keima 590-650nm. Cells were sequentially excited using

404nm and 568 nm laser. One representative middle slide was acquired, and images were

deconvolved and background subtracted using Huygens Professional (Scientific Volume

Imaging).

Xray-tomography:

Preparation of Specimen Mounting Capillaries. Thin-walled glass capillaries were pulled as

described in [3]. After being cut to length, capillaries were dipped in poly-L-lysine (0.01% solution,

Tissue Culture Grade, Sigma Aldrich, St. Louis, MO). Fiducial markers, used to align soft x-ray

projection images to a common axis, were generated by briefly immersing the capillaries a solution

of 100 nm gold nanoparticles (EMGC100, BBI International, Cardiff, CF14 5DX, UK), rinsing

and then drying them in air.

Specimen capillaries for use in correlated imaging experiments were prepared as above and then

dipped in a suspension of red polystyrene microspheres (FluoSpheres Carboxylate-Modified

Microspheres, 0.2 µm, Dark Red Fluorescent (660 excitation/ 680 emission), Life Technologies,

Invitrogen). The polystyrene microspheres were used fiducial markers for the alignment of

fluorescence images to each other and for the co-alignment of soft x-ray and cryo- fluorescence

tomographic reconstructions.

Specimen Mounting. 1 ml of yeast cells in mid-log phase growth (OD_{600} ~0.5) were pelleted and

re-suspended in ~50µl media. 1µl of this suspension was transferred into a specimen capillary

using a standard micropipette. The capillary contents were immediately cryopreserved by rapid

plunging into liquid propane at 165°C using a custom apparatus. Cryopreserved specimens were

transferred into storage boxes using a home-built cryo-transfer device and stored in a Dewar held

at 77K [3].

Imaging Cells in the Cryogenic Confocal Light Microscope (CLM). Specimen capillaries were

transferred cryogenically to a home-built cryogenic spinning disk confocal fluorescence

microscope [4]. Confocal scanning and detection were achieved using a commercial dual spinning

disk head (CSU-X1, Yokogawa, Tokyo, Japan). During imaging, the illumination wavelength was

selected with an acousto-optical tunable filter (AOTF) using an integrated system (Andor Laser

Combiner, Model LC-501A).

Experimental measurement of the CLM point spread function. The point spread function of

the cryogenic confocal light microscope was calculated by the methods described in [5].

Acquisition of Cryogenic Fluorescence Tomography data. Cryopreserved specimen capillaries were cryo-transferred to the CLM and aligned with respect to the rotation axis. Through-focus, or "z-stacks" were taken using a step size of 0.78 μm. For fluorescence tomography, the capillary was rotated about an axis normal to the objective lens using a motorized goniometer driven by custom LabView code (National Instruments Corporation, Austin, TX). Each tomographic data series consisted of 37 through-focus stacks (0° through 360° measured at 10° increments). Once measurements were completed, the capillary was cryo-transferred back to liquid nitrogen storage using the cryo-transfer device [5].

Alignment of through-focus fluorescence stacks using fluorescence fiducials. Well-isolated fluorescent fiducials were chosen manually from the raw fluorescence stacks and a three-dimensional centroid algorithm refined the bead position to sub-pixel accuracy using the software package Amira (FEI). The fiducial coordinates were used to write a new image stack in MATLAB containing a spherically-symmetrical representation of the fiducials, which we termed the "fiducial model". Six parameters were optimized for each alignment: three translations and three rotations (a rigid affine transformation). By limiting the search space to optimize correspondence between bead pairs that had been visually verified to be true fiducial correspondences, this alignment strategy was relatively fast and robust. The entire fluorescence fiducial alignment optimization process was visualized in Amira. Visualization was helpful for inspecting the results and troubleshooting rare cases where the optimization settled into a local maximum that was obviously not the real solution.

Fluorescence tomographic reconstruction and deconvolution. Preprocessed, aligned through-focus stacks were reconstructed into a single object by simply summing the through-focus datasets (Amira). The effective PSF for tomographic imaging in a specimen capillary was used as the

convolution kernel to deconvolve the fluorescence reconstruction. We used an iterative function

for the deconvolution. The algorithm found the fluorophore distribution that was most likely,

assuming the PSF was spatially invariant, and that image formation followed Poisson statistics.

Soft x-ray data collection and reconstruction. Specimens were imaged using a transmission soft

x-ray microscope, operated by the National Center for X-ray Tomography (NCXT), at the

Advanced Light Source of Lawrence Berkeley National Laboratory and reconstructed according

to previously published protocols.

Correlated Imaging. Co-alignment of cryogenic fluorescence and soft x-ray tomographic

reconstructions using joint fiducials

Joint fiducials - red fluorescent polystyrene microspheres - visible in both light- and soft x-ray

images were used to guide overlay of the two data types. The alignment process was optimized by

converting the raw fluorescence into a bead model and aligning the fluorescent bead model to the

x-ray reconstruction of the beads using an iterative optimization of the overlay.

Identification of JUNQ/IPOD in soft x-ray reconstructions. Segmentation is the process of

computationally isolating, visualizing, and quantifying specific cellular components in a

tomographic reconstruction. Each voxel in an SXT reconstruction is a direct measurement of the

soft x-ray Linear Absorption Coefficient (LAC) at the corresponding location in the cell.

Attenuation of the illumination by the specimen follows Beer's Law. Hence, the LAC values for

identical sized voxels depends solely on the concentration and composition of biomolecules

present. Consequently, water has an order of magnitude lower LAC than carbon-containing

molecules, such as lipids and proteins. LAC values for homogeneous solutions of isolated

biomolecules can also be calculated using tables of known absorption coefficients. For example,

pure water in the form of ice has a calculated LAC of 0.109 μm^{-1} whereas a model protein with the

chemical composition $C_{94}H_{139}N_{24}O_{31}S$ was calculated to have a theoretical LAC of 1.35μm^{-1} [6]. In

practice, most of the voxels in a 50 nm resolution SXT reconstruction of a cell will contain a

heterogeneous mixture of biomolecules. Using SXT data acquired at a single wavelength it is not

possible to distinguish the precise chemical species present. However, at this level of spatial

resolution organelles and other sub-cellular structures are sufficiently similar in their biochemical

composition to allow them to be readily identified from the surrounding cell contents. The

relatively high water content in vacuoles makes them readily distinguishable from organelles with

a greater density of biomolecules, for example, nuclei and mitochondria. Even relatively small

variations in organelle LAC can be distinguished, for example, the boundaries between nuclei and

nucleoli are very clear, as is the distinction between euchromatin and heterochromatin domains in

the nucleus. LACs were calculated, and cells segmented using the protocols described in complete

detail in [3].

Segmentation of the Hsp104-GFP reconstructions, in particular the assignment of the volumes as

JUNQ and IPOD, were guided by correlative fluorescence signal. In other strains, JUNQ and/or

IPOD were assigned based on LAC values established in the Hsp104-GFP study.

Western blot. Yeast cultures were grown to mid- log-phase in synthetic complete medium (yeast

nitrogen base, ammonium sulfate, amino acids, 2% raffinose) lacking uracil. Cultures were then

diluted to an OD_{600} of 0.2 in galactose medium (synthetic complete medium with 2% galactose

and 2% raffinose) and grown to an OD_{600} of 0.8. Cultures were then pelleted and resuspended in

glucose medium (synthetic complete medium with 2% glucose) and incubated at either 30 or 37

°C as indicated for the duration of the shut off. Where indicated, cultures were shifted to 37 °C for

1 hour prior to exchange into glucose medium and incubated at 37 °C for the duration of the

glucose shut-off. Cells were pelleted, proteins were extracted with urea and SDS, and protein

concentrations were measured with the bicinchoninic acid protein assay kit (Thermo Fisher

Scientific). Equal amounts of total protein were resolved on Tris-glycine gels (Invitrogen) and

transferred onto nitrocellulose membranes. GFP was detected with mouse anti-GFP antibodies

(Roche), GAPDH was detected with rabbit anti-GAPHD (Genetex), and PGK was detected with

rabbit anti-PGK antibodies (Thermo Fisher Scientific). Secondary antibodies were IRDye 800CW

donkey anti-mouse or IRDye 680RD donkey anti-rabbit and were imaged using the LiCOR

scanner and Image Studio Software.

To assess vacuolar protein degradation, the same procedure as above was followed but the *pep4D*

yeast were treated with 1mM PMSF (Research Products International) during the 37C incubation

step. The samples were then collected and lysed as above. GFP and GAPDH were detected with

the same primary antibodies, but the secondary antibodies were goat anti-mouse HRP conjugated

antibody (Promega Cat #W4021) and goat anti-rabbit HRP conjugated antibody (Promega Cat #

W4011). Blots were detected using Clarity Western ECL Substrate (Bio-Rad Cat# 1705061) and

imaged using the GE Amersham Imager 600.

Drop test. Yeast cultures were grown to OD_{600} of 0.6 in YPD or raffinose synthetic complete

medium and then diluted to OD_{600} of 0.1. Four 10-fold serial dilutions were performed in H_2O.

~2uL of diluted cultures were spotted onto YPD, galactose, or glucose plates using the pin plater.

Plates were incubated at indicated temperatures and imaged every 24 hours for 72 hours.

Nanobody purification and labeling. GFP-nanobody was expressed in BL21 Rosetta2 pLysS

cells (Novagen), and DsRed-nanobody (LAM4) was expressed in ArticExpress cells (Agilent

Technologies) overnight at 17 °C and 12 °C respectively. Cells were pelleted (4000XG), washed

with PBS containing 1 mM PMSF and pelleted again. Pellets were resuspended in column buffer

(500 mM NaCl and 50 mM HEPES pH 8 or 50 mM NaHCO$_3$ pH 8.3) with complete protease

inhibitors, PMSF, benzonase and 10 mM imidazole (Lysis buffer) and lysed using an emulsiflex.

Lysates were cleared at 20,000XG for 30 minutes. Nanobodies were purified using nickel resin

by passing cleared lysate over the resin, washing with lysis buffer, followed by column buffer, and

elution with column buffer containing 300 mM imidazole. Elution was concentrated using an

Amicon ultra 3 kD MWCO to less than 2 mL, and further purified by size exclusion using an

SDX75 column (GE Healthcare) equilibrated with 500 mM NaCl, 50 mM NaHCO$_3$, 1 mM DTT.

Labeling was performed by the addition of dyes at a molar ratio of 5:1 dye:nanobody using Alexa

488 NHS (Thermo Fisher Scientific) for the GFP nanobody and Alexa 568 NHS (Thermo Fisher

Scientific) for the DsRed nanobody. Labeling reactions were incubated 1 hr at room temperature

and quenched by addition of Tris-HCl pH 8 to a final concentration of 100 mM. Labeled nanobody

was separated from free dye by size exclusion using an SDX75 column.

GFP-nanobody cloning. pDG402, a vector for expression of GFP-nanobody-TEV-6XHis was

generated by PCR amplification of the GFP nanobody using the 5' primer (XbaI GFP nanobody)

AATCTAGAATTTTGTTTAACTTTAAGAAGG and 3' primer (GS link TEV His nanobody)

AAGGATCCTTATCAGTGATGATGGTGGTGATGGGACCCAGATCCACCCTGGAAGTAT

AAATTCTCACCCGAACCTCCGGATGAGGAGACGGTGACCTGGG. The PCR product

was cloned into the pST39 backbone [7] using XbaI and BamHI.

METHODS REFERENCES

1   Brickner, D. G., Light, W. & Brickner, J. H. Quantitative localization of chromosomal
loci by immunofluorescence. *Methods Enzymol* **470**, 569-580, doi:10.1016/S0076-
6879(10)70022-7 (2010).

2   Woerner, A. C. *et al.* Cytoplasmic protein aggregates interfere with nucleocytoplasmic
transport of protein and RNA. *Science* **351**, 173-176, doi:10.1126/science.aad2033 (2016).

3   Parkinson, D. Y. *et al.* Nanoimaging cells using soft X-ray tomography. *Methods Mol*
Biol **950**, 457-481, doi:10.1007/978-1-62703-137-0_25 (2013).

4   Le Gros, M. A., McDermott, G., Uchida, M., Knoechel, C. G. & Larabell, C. A. High-
aperture cryogenic light microscopy. *J Microsc* **235**, 1-8, doi:10.1111/j.1365-2818.2009.03184.x
(2009).

5   Le Gros, M. A. *et al.* Biological soft X-ray tomography on beamline 2.1 at the Advanced
Light Source. *J Synchrotron Radiat* **21**, 1370-1377, doi:10.1107/s1600577514015033 (2014).

6   Weiss, D. *et al.* Computed tomography of cryogenic biological specimens based on X-ray
microscopic images. *Ultramicroscopy* **84**, 185-197, doi:10.1016/s0304-3991(00)00034-6 (2000).

7   Tan, S. A modular polycistronic expression system for overexpressing protein complexes

in Escherichia coli. *Protein Expr Purif* **21**, 224-234, doi:10.1006/prep.2000.1363 (2001).

ACKNOWLEDGMENTS

We thank support of NIH (GM05643319 and AG054407 to JF; F32NS086253 to ES) and Way

Klingler Startup Funds from Marquette University (ES). F.M-P. was supported by The Pew Trusts

in the Biomedical Sciences postdoctoral Award (00034104). Cryo-SXT data were acquired at the

Advanced Light Source of Lawrence Berkeley National Laboratory, a U.S. DOE Office of Science

User Facility. The National Center for X-ray Tomography is supported by NIH (P41GM103445)

and DOE BER (DE-AC02-5CH11231). CAL and MAL were supported by the Gordon and Betty

Moore Foundation Award #3497. The Stanford Cell Sciences Imaging Facility is supported in part

by Award Number 1S10OD01227601 from the National Center for Research Resources (NCRR).

We thank Jon Mulholland and Youngbin Lim from the CSIF for training on the SIM. We are

grateful to Michael Rosbash (Brandeis University), Susan Wente (Vanderbilt University), Kiran

Madura (Rutgers University), and Patrick Lusk (Yale School of Medicine) for providing yeast

strains and Karsten Weis (ETH Zurich), Jodi Nunnari (University of California, Davis), and

Michael P. Rout (Rockefeller University) for plasmids. We thank Katharine Ullman (University

of Utah), Adam Frost (Altos Lab) and Lisbeth Veenhoff (University of Groningen) for discussions

and advice. We are grateful to Carson Trail for support in microscopy data analysis, Margaret

Wangeline for assisting with the 2xKeima cloning, Joan Steffan (UC Irvine), Kelly Rainbolt

(Stanford University) for discussions and comments on the manuscript, Felipe Serrano for assisting

on Figure 8, and the Frydman lab for advice and discussions.

AUTHOR CONTRIBUTIONS

JF and ES conceived the project; ES and F. M-P carried out all experiments; J-HC collected and

processed cryo-SXT data; GM carried out cryo-SXT data analysis and modelling; MLG

performed cryo-fluorescence data acquisition and correlation with cryo-SXT data and built the

microscope used for these experiments. PD analyzed particle tracking data and assisted on

statistical analyses. DG cloned the NLS- and NES- luciferase and VHL plasmids. DG and F. M-

P generated, purified and labeled the GFP and RFP nanobodies. ES, F. M-P and JF wrote the

MS. All authors commented on the final version. JF directed the project.

DECLARATION OF INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information is available for this paper.

Correspondence and requests for materials should be addressed to Judith Frydman

jfrydman@stanford.edu

EXTENDED DATA FIGURE LEGENDS

1234

Figure ED1: Spatial sequestration occurs during different types of stress with different

client proteins.

(a) Western blot analyses of Gal Shut-off assays showing the clearance of NLS-LuciTs (top) and

NES-LuciTs (bottom) with and without proteasome impairment by 50u$\underline{M}$ Bortezomib. (b) Drop

test of W303 yeast expressing model proteins without heat shock at 30C (left), with heat shock at

37 °C (middle), and without expression of the plasmids (right). (c) Representative Structured

Illumination super-resolution microscopy images taken of cells expressing NLS-VHL (left) or

NES-VHL (right) after 120 minutes at 37 °C and treated with 100μ$\underline{M}$ MG132. NLS-LuciTs is

shown in green, NES-LuciTs in purple, nuclear pores in gold and Hoechst counterstain in blue.

Scale bars are 1μm.

1245

Figure ED2: The effect of blocking nucleocytoplasmic transport on Ubc9Ts clearance.

(a) Quantitation of the percentage of cells containing nuclear or cytoplasmic inclusions in WT

yeast expressing Ubc9Ts-EGFP after 120 minutes at 37 °C with and without treatment with 100μM

MG132. A minimum of 500 cells per condition from 3 biologically independent experiments were

counted and Student's t-tests were performed comparing the WT yeast without MG132 treatment

to WT yeast with MG132 treatment using Prism software. P values were adjusted using two-stage

linear step-up procedure of Benjamini, Krieger and Yekutieli with a Q of 5%. Adjusted P value

for nuclear no MG132 vs. +MG132 is 0.0035 and cytoplasmic no MG132 vs. +MG132 is 0.0011.

(b) Representative Structured Illumination super-resolution microscopy images taken of cells

expressing GFP-VHL after 2 hr at 37 °C with DMSO (left) or with 100μM MG132 (right)

treatment. VHL is shown in green, nuclear pores in gold, and Hoechst counterstain in blue. Scale

bars are 1μm. (c) (left) Schematic illustrating the *nup116-5* yeast have sealed nuclear pores at 37

°C, thus blocking nucleocytoplasmic trafficking. (right) There is no significant difference in half-

life between WT and *nup116-5* yeast. (d) (left) Schematic illustrating the *sts1-2* yeast do not

translocate proteasomes to the nucleus at 37 °C. (right) There is no significant difference in half-

life between WT and *sts1-2* yeast. (c and d) Levels of EGFP were quantified from 3 biologically

independent experiments. Half-life was calculated using Prism.

1263

Figure ED3: INQ-JUNQ homing does not occur at the LINC, nucleolus, or involve FG repeats of the nuclear pore central channel.

(a) Graph of the X-Y positions of the INQ and JUNQ compartments by particle tracking of inclusions from cell shown in Figure 2a over the time course of the experiment. (b) Representative confocal image taken of cells co-expressing NLS-EGFP-VHL and NES-DsRed-VHL after 2 hr at 37 °C and treated with 100μM MG132. NLS-fusion proteins are shown in green, NES-fusion proteins in purple, nuclear pores in gold, and Hoechst counterstain in blue. Scale bar is 1μm. (c) Representative confocal fluorescence microscopy images taken of cells co-expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs (left) after 2 hr at 37 °C and treated with 100μM MG132. NLS-LuciTs is shown in green, NES-LuciTs in purple, nucleolus (Nsr1) in gold and Hoechst counterstain in blue. (right) Line intensity profile showing distance between nucleolus and homed INQ/JUNQ. Scale bars are 1μm. (d) (left) schematic of Mps3 component of LINC complex linking inner and outer nuclear membranes. (right) Representative widefield fluorescence microscopy images taken of cells co-expressing endogenously tagged Mps3-EGFP and NES-DsRed-LuciTs after 120 minutes at 37 °C with and without treatment with 100μM MG132. White arrowheads indicate locations of Mps3 puncta while yellow arrowheads indicate NES-LuciTs puncta. Scale bars are 1μm. (e) WT (top) and *nupΔFG* (bottom) cells co-expressing NLS-LuciTs and NES-LuciTs were shifted to 37 °C and monitored by live cell time-lapse fluorescence microscopy for the times shown. Scale bars are 1μm.

Figure ED4: Detailed representation of the cryo-SXT workflow and interactions between

mitochondria and cytoplasmic PQC compartments.

(a) Optical path through the specimen. Key: COL, cryogenic objective lens; SS, specimen stage;

SP, specimen port; MG, motorized goniometer; CIM, cryogenic immersion fluid; CCL, low

magnification cryogenic objective; CS, cryogenic specimen; CIE, cryogenic imaging

environment; AP, adapter port; AW, a heated, angled anti-reflection window. (b) Alignment of

fluorescence and soft x-ray tomographic data using fiducial markers. (c) A representative confocal

image of the spatial relationship between the INQ and nucleolus. NLS-LuciTs (INQ) is shown in

green, nucleolus in gold, and Hoechst counterstain in blue. Scale bar is 1μm. (d) The interaction

between mitochondria and cytoplasmic inclusions is also seen by fluorescence confocal

microscopy in a representative image of a cell co-expressing mito-GFP and NES-RFP-LuciTs.

NES-LuciTs is shown in purple, mitochondria in cyan, and Hoechst counterstain in blue. Scale bar

is 1μm. (e) Representative confocal fluorescence microscopy images taken of WT, fission mutants

(*dnm1Δ* and *fis1Δ*) and fusion mutant (*fzo1Δ* and *ugo1Δ*) cells expressing mito-GFP and NES-

DsRed-LuciTs after 120 minutes at 37 °C and treated with 100μM MG132. Mito-GFP is shown

in cyan, NES-LuciTs in purple, and Hoechst counterstain in blue. Scale bars are 1μm.

1300

Figure ED5: NVJ -mediated clearance of misfolded proteins.

(a) Endogenously tagged Nvj1-GFP yeast expressing Ubc9Ts-ChFP were shifted to 37 °C and

monitored by live cell time-lapse fluorescence microscopy for the times shown. White arrowheads

indicate locations of Nvj1 puncta while yellow arrowheads indicate Ubc9Ts-ChFP puncta. Scale

bar is 1μm. (b) WT (top) and *nvj1Δ* (bottom) cells co-expressing NLS-LuciTs and NES-LuciTs

were treated with 100μM MG132 and shifted to 37 °C for 30 mins to preform inclusions. Cells

were then placed in media containing 50mg/ml cycloheximide (CHX) and 100μM MG132 at 37

°C and monitored by live cell time-lapse fluorescence microscopy for the times shown. Scale bars

are 1μm. (c,d) Quantitation of the percentage of cells containing cytoplasmic inclusions in WT,

nvj1Δ, and *vac8Δ* yeast co-expressing NLS-EGFP-LuciTs (c) and NES-DsRed-LuciTs (d) after 2

hr at 37 °C with and without treatment with 100μ<u>M</u> MG132. A minimum of 500 cells per condition

from 3 biologically independent experiments were counted and unpaired Student's t-tests were

performed comparing the deletion strains to WT with or without MG132 treatment were compared.

P values were adjusted using two-stage linear step-up procedure of Benjamini, Krieger and

Yekutieli with a Q of 5%. (c) %. Adjusted P value for WT vs *nvj1Δ* no MG132 is 0.0205, WT vs

nvj1Δ +MG132 is 0.2357, WT vs *vac8Δ* no MG132 is 0.0406, WT vs *vac8Δ* +MG132 is 0.8974,

and WT no MG132 vs WT +MG132 is 0.0239. (d) Adjusted P value for WT vs *nvj1Δ* no MG132

is 0.013, WT vs *nvj1Δ* +MG132 is 0.013, WT vs *vac8Δ* no MG132 is 0.0008, WT vs *vac8Δ*

+MG132 is 0.0447, and WT no MG132 vs WT +MG132 is 0.0005. (e) Levels of EGFP after 1

hour at 37 °C were measured from Quantitative Western blots (mean ± s.e.m. from three

biologically independent experiments). WT and nvj1Δ yeast were compared using an unpaired

Student's t-test resulting in a P value of 0.0043 for WT vs nvj1Δ NLS-LuciTs and NES-LuciTs P

value is 0.0183. Statistics were performed using Prism.

Figure ED6: ESCRT involvement in the clearance of misfolded proteins.

(a) Representative confocal images of WT yeast co-expressing Chm7-EGFP and either NLS-
EGFP-LuciTs (left) or NES-DsRed-LuciTs (right) after 120 minutes at 37 °C and treated with
100μM MG132. Chm7 is shown in teal and remains diffuse throughout the cell, NLS-EGFP-
LuciTs in green, NES-DsRed-LuciTs in purple, nuclear pores in gold and Hoechst counterstain in
blue. Scale bar is 1μm. (b) Representative confocal images of WT and *vps23Δ*,
vps34Δ, and *vps15Δ* yeast co-expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 2 hr at
37 °C and treated with 100μM MG132. NLS-EGFP-LuciTs is shown in green, NES-DsRed-
LuciTs in purple, nuclear pores in gold, and Hoechst counterstain in blue. Insets show the budding
INQ encapsulated by nuclear pores. Scale bars are 1μm. Same data as shown in Figure 6c, but with
the green channel separated to clearly detail the colocalization with the cytoplasmic protein.

Figure ED7: Vacuole-mediated clearance of INQ and JUNQ.

(a) Representative images of WT cells expressing NES-2xKeima-LuciTs after 2 hr incubation at 37
°C with 100μM MG132. Over time, fluorescence is seen with excitation in the 589nm channel
indicating the NLS-LuciTs has encountered an acidic environment. Insets show the transition from
green to red and a structure leaving the inclusion that is fully red. Scale bars are 5 uM. Same data
shown in Figure 7d, but with more time points and a larger field of view in the images. (b) WT cells
expressing NES-2xKeima-LuciTs after 85 min incubation at 37 °C with 100μM MG132. (c) Longer
exposure of the blot shown in Figure 7f to highlight the difference in the number and pattern of the
GFP bands in the WT vs *pep4Δ* cells. (d) Levels of EGFP at time 0 were measured from
Quantitative Western blots such as those shown in Figure 7e, f (mean ± s.e.m. from three
biologically independent experiments). WT and *pep4Δ* yeast were compared using an unpaired
Student's t-test without reaching statistical significance. (e) WT yeast expressing NLS-GFP-LuciTs

were treated with 8uM of FM4-64 and incubated for 2hr at 37 °C with 100μM MG132. Cells were

imaged every 30 sec for 90 mins. Scale bar is 1μm. Same data shown in Figure 7h, but only WT

and with more timepoints during the entry into the vacuole.

1350

Legends to movies

Movie 1. Live-cell time-lapse fluorescence microscopy of WT cells expressing NLS-LuciTs (left)

or NES-LuciTs (right) at 37 °C, treated with 100μM MG132. Same data shown in stills in Figure

1c (left), d (right).

Movie 2. Dynamic representation of the 3D reconstructions of the data shown in Figure 1e (left),g

(right). Videos were created in Volocity.

Movie 3. Dynamic representation of the 3D reconstructions of the data shown in Figure 2a (left),

b (right). Videos were created in Volocity.

Movie 4. Live-cell time-lapse fluorescence microscopy of the data shown in Figure 3a.

Movie 5. Dynamic representation of the data shown in Figure 3b.

Movie 6. Dynamic representation of the 3D reconstruction of the data shown in Figure 3c. Video

was created in Volocity.

Movie 7. Dynamic representation of the 3D reconstruction shown in Figure 4c. Video was created

in Amira.

Movie 8. Dynamic representation of the 3D reconstruction shown in Figure 4d,f. Video was

created in Amira.

Movie 9. Live-cell time-lapse fluorescence microscopy of the data shown in Figure 5a.

Movie 10. Dynamic representation of the 3D reconstruction of the data shown in Figure 5c. Video

was created in Volocity.

Movie 11. Dynamic representation of the 3D reconstruction of the data shown in Figure 5b. Video

was created in Volocity.

Movie 12. Live-cell time-lapse fluorescence microscopy of the data shown in Figure 5f shown at

5 frames per second.

Movie 12b. Live-cell time-lapse fluorescence microscopy of the data shown in Figure 5f shown at

2 frames per second.

Movie 12c. Live-cell time-lapse fluorescence microscopy of the data shown in Figure 5f shown at

2 frames per second. Only the 488nm channel is shown in greyscale.

Movie 13. Dynamic representation of the 3D reconstruction of the data shown in Figure 5e. Video

was created in Volocity.

Movie 14. Live-cell time-lapse fluorescence microscopy of the WT cell data shown in Figure 5h.

Movie 15. Live-cell time-lapse fluorescence microscopy of the *nvj1Δ* cell data shown in Figure

5h.

Movie 16. Live-cell time-lapse fluorescence microscopy of the *vps4Δ* cell data shown in Figure

5h.

a Misfolding → Sequestration

i JUNQ / INQ or *ii* INQ

b LuciferaseTs: Nuclear (NLS) vs Cytoplasmic (NES)

Folded (25 °C) → Gal Shut-off Unfold (37 °C) → Unfolded → Sequestration (Nuc vs Cyto?) → MG132 ⊣ Degraded

c NLS-GFP-LuciTs, Time (min) at 37 °C

min 0 5 15 30 60 120

-MG132

+MG132

d NES-GFP-LuciTs, Time (min) at 37 °C

min 0 5 15 30 60 120

-MG132

+MG132

e *i.* NLS-LuciTs, 37 °C — INQ

ii. A→B: INQ, DNA, Nups

NLS-LuciTs: INQ
NES-LuciTs: JUNQ IPOD
Nups: Nuclear Pores
DNA: Nucleus

f *i.* NLS-VHL, 30 °C — INQ, A→B

ii. A→B: Nups, DNA, INQ

NLS-VHL: INQ
NES-VHL: JUNQ IPOD
Nups: Nuclear Pores
DNA: Nucleus

g *i.* NES-LuciTs, 37 °C — JUNQ, IPOD

ii. A→B: Nups, DNA, JUNQ

iii. C→D: IPOD, DNA, Nups

h *i.* NES-VHL, 30 °C — JUNQ, A→B

ii. A→B: JUNQ, Nups, DNA

a NLS-LuciTs + NES-LuciTs; Time (min) at 37 °C

0' 5' 15' 30' 60' 90'

Puncta Formation *Homing* *Co-migration*

b INQ–JUNQ Distance (nm)

Puncta Formation

INQ — Distance — JUNQ

Nups

Homing Nups

Co-migration Nups

Time at 37 °C (min)

c NLS-LuciTs (INQ)
NES-LuciTs (JUNQ)
Nups
DNA

INQ Nups JUNQ

A B

d

i. NES-LuciTs (JUNQ) Nups DNA

WT *nup120Δ* *nup120Δ*

Nups

ii. NLS-LuciTs (INQ) Nups DNA

WT *nup120Δ*

iii. NLS-LuciTs INQ NES-LuciTs JUNQ Nups DNA

WT *nup120Δ*

e JUNQ Nups INQ

a

Frozen, unfixed intact cells → Fluorescence microscopy → X-ray Tomography → Measure LAC → Correlate X-ray/ fluorescence → 3D reconstruction

b *i.*

Cells
Hsp104-GFP + Htt97QP-ChFP

X-ray Tomogram   Fluorescence overlay

ii.

Organelles	LAC
Cytosol	0.43
Nucleus	0.42
Mitochondria	0.50
Vacuoles	0.38
Lipid droplet	0.67
Fluorescent PQC Inclusion	0.47

c

3D reconstruction

JUNQ
450nm

IPOD

d

NES-LuciTs at 37 °C (JUNQ)

i.   *ii.*

JUNQ

e

NES-LuciTs

Mitochondria
Vacuole
Nucleus
Lipid droplets

f

180°

Mitochondria   Vacuole   Nucleus   NES-LuciTs

g

NLS-LuciTs at 37 °C (INQ)

400nm   INQ

Nucleolus   Nucleus
NLS-Luci   Vacuole

h

Mitochondria
Q-bodies

Nucleolus

IPOD   JUNQ   INQ   Nucleus
NVJ
Vacuole

JUNQ ?   Nucleus
INQ ?
Vac8   Nvj1
Vacuole

a *i.* Chm7 inactive
ESCRT-II /-III
Chm7 active (open)
JUNQ
INQ
Nucleus

ii. Chm7 (open) NLS-LuciTs
Chm7 (open) NES-LuciTs

b Co-express
NLS-LuciTs + NES-LuciTs
WT
chm7Δ

c ESCRT I Vps23 *Ub-binding*
Vps34-Vps15 *PI3K*
Chm7
INQ
Nu
Vac
NVJ
INQ

WT   *vps23Δ*   *vps34Δ*   *vps15Δ*

INQ-JUNQ not homed
Cytoplasmic INQ
Budding INQ

NLS-LuciTs
NES-LuciTs
Nups
DNA

d ESCRTIII
V
VPS4
V

e Cells with Inclusions
WT vs *vps4Δ*
WT
vps4Δ
% Cells with Inclusions
NLS-LuciTs   NES-LuciTs
***
**

f Co-express NLS-LuciTs + NES-LuciTs
WT   *vps4Δ*
Merge   NLS-LuciTs   Merge

INQ   JUNQ   Nuclear Pores   DNA

a

WT   + Bortezomib

NLS-LuciTs
GAPDH
Time (min)   0   30   60   90   120   0   30   60   90   120

NES-LuciTs
GAPDH
Time (hr)   0   1   2   4   0   1   2   4

Time after shift to 37°C

b

	Galactose 30 °C	Galactose 37 °C	Glucose
GFP			
LuciTs			
NLS-LuciTs			
NES-LuciTs			
Htt97QΔP			

c

VHL 37 °C +MG132

NLS-VHL
Nups
DNA

INQ   DNA   Nups

d

VHL 37 °C +MG132

NES-VHL
Nups
DNA

JUNQ   Nups   DNA

a

Ubc9Ts

b

37 °C

-MG132   +MG132

GFP-VHL   Nups   DNA

c

nup116-5

Ubc9Ts Half Life (37 °C)

d

sts1-2

Ubc9Ts Half Life (37 °C)

a

EMCCD-LW   EMCCD-SW   Laser input   Spectrometer input   AL   MG   LM-CCD

RL   DM   MFW   Spinning disk   TL   MDT

External surface of capillary

CS

COL   CCL

CIM

b

a Alignment Using Fluorescence and Gold Fiducials

CFT reconstruction and SXT co-alignment fiducials
(200 nm red polystyrene)

SXT reconstruction alignment fiducials
(100 nm gold)

Specimen

b Alignment Using Joint Fiducials

SXT reconstruction alignment fiducials
(100 nm gold)

Joint Fiducials
(500 nm polystyrene)

Internal surface of capillary

Specimen

c

NLS-LuciTs
Nucleolus
DNA

d

NES-LuciTs
Mitochondria
DNA

e

WT	Mitochondrial Fission mutants		Mitochondrial Fusion mutants	
	dnm1Δ	*fis1Δ*	*fzo1Δ*	*ugo1Δ*

Mito-GFP   **NES-LuciTs**   **DNA**

a

Nvj1-GFP + Ubc9Ts-ChFP; Time (min) at 37°C

b

NLS-LuciTs + NES-LuciTs; Time (min) in CHX

c Nuclear NLS-LuciTs

d Cytoplasmic NES-LuciTs

e LuciTs after 60 mins at 37C

a

b

a

NES-2xKeima-LuciTs at 37°C

Time at 37 °C (min)

| | 0 | 15 | 30 | 40 | 50 | 60 | 80 | 90 |

Ex 404

Ex 558

Merge

b

NES-2xKeima-LuciTs 85 min chase at 37°C

Ex 404 · Ex 558 · Merge

c

d

Normalized T0 Intensity

WT · pep4Δ · WT · pep4Δ
NLS-LuciTs · NES-LuciTs

e

WT Chase Time at 37 °C (min) · NLS-LuciTs · Vacuole

30:00 · 40:00 · 45:00 · 45:30 · 46:00 · 46:30

47:00 · 47:30 · 48:00 · 49:00 · 50:00 · 60:00

Nr.	NAME	GENOTYPE	SOURCE AND COMMENTS	
SUPPLEMENTARY TABLE S1. Genotypes and sources of yeast strains used in this study				
ESY 1	BY4741			
ESY 104	W303			
ESY 105	CRM1		WT strain for crm1-T539C	from Michael Rosbash
ESY 106	crm1-T539C		has nuclear export deficiency with leptomycin B treatment	from Michael Rosbash
ESY 95	Nup116-5	nup116-5::HIS3	SWY027	from Susan Wente
ESY 96	STS1		WT strain for sts1-2	from Kiran Madura
ESY 97	sts1-2		temperature sensitive mutation that blocks proteasome import into the nucleus	from Kiran Madura
ESY 113	*dnm1*	BY4741 Mata	Deletion collection	
ESY 119	*fis1*	BY4741 Mata	Deletion collection	
ESY 122	*fzo1*	BY4741 Mata	Deletion collection	
ESY 123	*ugo1*	BY4741 Mata	Deletion collection	
ESY 121	Mps3-GFP	BY4741 Mata	GFP collection	
ESY 127	*nup120*	BY4741 Mata	Deletion collection	
ESY 94	*nupΔFG*		SWY3042	from Susan Wente
ESY 14	Nvj1-GFP	BY4741 Mata	GFP collection	
ESY 4	*nvj1*	BY4741 Mata	Deletion collection	
ESY 3	*vac8*	BY4741 Mata	Deletion collection	
ESY 150	*vps4*	BY4741 Mata	Deletion collection	
ESY177	Chm7OPEN-GFP	W303, chm7OPEN-(1-369)-GFP::HIS3	DTCPL413	From Patrick Lusk
ESY183	*Chm7*	W303, chm7Δ::hphMX6		From Patrick Lusk
ESY182	Chm7-GFP	W303, CHM7-GFP::HIS3		From Patrick Lusk
FMP1X	Nvj1-sfGFP	BY4741 Mata	NVJ1 GFP C-terminally tagged	This study

Nr.	NAME	SOURCE AND COMMENTS
SUPPLEMENTARY TABLE S2. Plasmids used in this study		
ES p331	NLS-GFP-LuciTs	This study
ES p357	NES-DsRed-LuciTs	This study
ES p325	NLS-GFP-VHL	This study and (Samant et al. 2018)
ES p355	NES-DsRed-VHL	This study and (Samant et al. 2018)
ES p20	Ubc9Ts-GFP	(Escusa-Toret et al. 2013)
ES p562	Ubc9Ts-ChFP	(Escusa-Toret et al. 2013)
ES p144	GFP-VHL	(Escusa-Toret et al. 2013)
ES p536	pRS NLS-GFP2-NES	(Stade et al., 1997)
ES p57	mHTT 97QP-ChFP	(Escusa-Toret et al. 2013)
ES p520	vt100 GFP	(Westermann et al. 2000)
FMP 1X	NLS-Keima-LuciTs	This study
FMP 1Y	NLS-2xKeima-LuciTs	This study
FMP 2X	NES-Keima-LuciTs	This study
FMP 2Y	NES-2xKeima-LuciTs	This study