An ESCRT-dependent pathway of Nuclear and Cytoplasmic Spatial PQC is coordinated at

Nuclear Vacuolar Junctions

Emily M. Sontag [1,*,†,£], Fabián Morales-Polanco [1,£], Jian-Hua Chen[2], Gerry McDermott[2], Patrick

T. Dolan[1,‡], Dan Gestaut[1], Mark A. Le Gros[2,3], Carolyn Larabell[2,3] and Judith Frydman[1,*]

[1] Department of Biology, Stanford University, Stanford, CA 94305, USA

[2] Department of Anatomy, School of Medicine, University of California San Francisco, San

Francisco, CA 94158, USA

[3] Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National

Laboratory, Berkeley, CA 94720, USA

[*] Correspondence to: emily.sontag@marquette.edu; jfrydman@stanford.edu,

£ Equal Contribution

† Current address: Department of Biological Sciences, Marquette University, Milwaukee, WI

53233, USA

‡ Current address: Quantitative Virology and Evolution Unit, National Institute of Allergy and

Infectious Diseases, Bethesda, MD 20892, USA

Abstract. Spatial sequestration of misfolded proteins is an integral step in the process of cellular

Protein Quality Control (PQC). is essential for cellular health. One key PQC strategy relies on

spatial sequestration of misfolded proteins into protein inclusions. Herein we define the

coordination of nuclear and cytoplasmic spatial PQC. While cytoplasmic misfolded proteins

concentrate in a cytoplasmic, perinuclear Juxta Nuclear Quality control (JUNQ) compartment,

nuclear misfolded proteins sequester into a perinucleolar IntraNuclear Quality control (INQ)

compartment. Particle tracking analysis reveals that INQ and JUNQ converge to localize to face

each other across the nuclear envelope at a site proximal to the Nuclear-Vacuolar Junctions (NVJ).

Strikingly, this convergence is coordinated by a perinuclear ESCRT pathway involving nuclear

envelope-localized ESCRT-II/-III protein Chm7. Indeed, deleting the NVJ or ATPase Vsp4,

required for membrane invagination into the vacuole, impairs clearance of inclusions.

Furthermore, deleting other ESCRT components led to cytoplasmic accumulation of nuclear

misfolded proteins. Our findings uncover an alternative role of perinuclear ESCRT and the nuclear

envelope in coordinating spatial PQC and clearance of nuclear and cytoplasmic misfolded proteins

into the vacuole.

Introduction

Misfolded proteins can acquire toxic conformations that disrupt essential cellular processes [1-6] leading to a growing number of diseases, including neurodegeneration and cancer [7-10]. Accordingly, cellular protein homeostasis (proteostasis) is maintained by a network of chaperones and clearance factors that promote protein quality control (PQC) [11-15]. One fundamental PQC strategy is to spatially sequester misfolded proteins into distinctly localized membrane-less compartments, likely to remove damaging conformers from the cellular milieu and concentrate them for more efficient refolding or clearance through either the ubiquitin-proteasome system (UPS) or endolysosomal pathways [11, 16-18]. Misfolded cytoplasmic proteins partition into distinct inclusions depending on their aggregation state. Insoluble amyloid proteins are sequestered into the Insoluble Protein Deposit (herein the IPOD) [4, 11, 19]. Soluble misfolded proteins are sequestered into small, dynamic, ER-associated structures called Q-bodies [17] which, during sustained stress, coalesce along the endomembrane system [20] into a Juxtanuclear Quality Control compartment (herein the JUNQ). Spatial PQC through the IPOD and JUNQ pathways is evolutionarily conserved and is present in yeast, worms and mammalian cells [11, 19]. Spatial PQC is also observed in the nucleus, as nuclear misfolded proteins are sequestered in a membrane-less intranuclear quality control compartment (herein the INQ), proximal to the nucleolus [18, 21, 22].

The relationship and overall coordination between spatial PQC of cytoplasmic and nuclear proteins is not well understood. While many studies showed cytoplasmic misfolded proteins being concentrated into the cytoplasmic JUNQ and cleared through an ER-anchored cytoplasmic UPS pathway [11, 19, 23, 24], it's also been proposed that clearance of cytoplasmic misfolded proteins requires import into the nucleus, whereby the JUNQ would form an inclusion identical to the INQ [18].

Given the importance of protein quality control for cell survival and its relevance to understand

misfolding diseases, we examined the logic and coordination of spatial PQC for nuclear and

cytoplasmic misfolded proteins. We unequivocally demonstrate the cytoplasmic localization of the

JUNQ and nuclear localization of the INQ.

Of note, the JUNQ and INQ usually home to opposite sides of the nuclear envelope, at a site

proximal to NVJ, which facilitates the extrusion of the INQ for clearance into the vacuole in an

ESCRT- and VPS4-dependent manner. Here we propose the surprising degree of spatial

coordination between nuclear and cytoplasmic PQC inclusions providing an avenue for vacuolar

clearance of misfolded proteins.

RESULTS

Organelle-specific PQC compartments sequester nuclear and cytoplasmic misfolded

proteins

To define the relationship and interplay between nuclear and cytoplasmic spatial PQC (Fig. 1a),

we utilized two validated models of protein misfolding: a temperature-sensitive variant of Firefly

Luciferase (herein LuciTs) that misfolds at 37 °C, and the Von Hippel-Lindau (VHL) tumor

suppressor, which is constitutively unfolded in yeast cells [25, 26]. We restricted their subcellular

localization to either nucleus or cytoplasm by inclusion of a Nuclear Localization Signal (NLS) or

a Nuclear Export Signal (NES), respectively. The expression of these PQC substrates was first

selectively induced by growth in galactose, and then repressed by glucose addition, which allowed

measuring the kinetics of their clearance or spatial PQC (Fig. 1b). As previously described, both

NLS- and NES-LuciTs and NLS- and NES-VHL are degraded upon misfolding and are stabilized

by proteasomal inhibition with either MG132 or Bortezomib [27] (Extended Data Fig. 1a).

To initiate LuciTs misfolding and monitor the fate of the misfolded species, the cells were shifted

to 37 °C and evaluated using Time-resolved fluorescence microscopy (Fig. 1b). Under folding-

permissive conditions, NLS- and NES-GFP-LuciTs were diffusely localized in the nucleus or

cytoplasm, respectively (Time 0, Fig 1 c, d), but following the 37 °C shift, both nuclear (NLS) and

cytoplasmic (NES) LuciTs were rapidly concentrated into dynamic puncta in their respective

cellular locations (Fig. 1c, d, and Supplemental video 1).

Over the course of the experiment, nuclear NLS-LuciTs coalesced into one intranuclear INQ

inclusion (Time 30, Fig. 1c). In contrast, cytoplasmic NES-LuciTs puncta coalesced into two

cytoplasmic inclusions, corresponding to the JUNQ and IPOD (Time 30, Fig. 1d), as previously

observed for Ubc9Ts and VHL [11]. Of note, the NLS- and NES- LuciTs variants formed inclusions

in both absence or presence of proteasome inhibitor MG132 (Figure 1c, d), indicating that spatial

sequestration into PQC compartments is not a consequence of proteasomal inhibition.

We used structured illumination (SIM) super-resolution microscopy to better define the nuclear

and cytoplasmic location and morphology of these PQC inclusions. Cells expressing LuciTs

variants were incubated for 2 hrs at 37 °C, while cells expressing VHL variants were grown and

fixed at both 30 and 37 °C. The nuclear envelope was visualized by immunostaining of nuclear

pore protein Nsp1 (Nups, yellow) and DNA was visualized using Hoechst (blue) (Fig. 1e-h). The

INQ, formed by nuclear misfolded protein, localized inside the nucleus in a pocket between the

nuclear envelope (delineated by Nups) and the DNA (Fig. 1e and Supplemental video 2 for NLS-

LuciTs at 37 °C, Fig. 1f for NLS-VHL at 30 °C Extended Data Fig. 1c for NLS-VHL at 37 °C).

Line intensity profile analyses confirmed the intranuclear localization of this INQ (Fig. 1e, f,

Extended Data Fig. 1c). Importantly, SIM and line intensity profile analyses unequivocally

established that the PQC inclusions formed by cytoplasmic misfolded proteins NES-LuciTs and

NES-VHL reside outside of the nucleus: one inclusion is adjacent to the nuclear envelope as

expected for the JUNQ, while the other inclusion is closer to the cell periphery as expected for the

IPOD (Fig. 1g-h, Supplemental video 2, Extended Data Fig. 1d). Similar results were obtained

with NLS- and NES-VHL at both 30 °C and 37 °C indicating PQC inclusion localization is not

temperature-dependent (Extended Data Fig. 1c-d). Consistent with their distinct localization,

cytoplasmic and nuclear misfolded variants also exhibit differential toxicity to cells; cytoplasmic

misfolded proteins were slightly more toxic, albeit less so than a polyglutamine expanded exon1

variant of Huntingtin (herein mHTT) associated with Huntington's disease (Extended Data Fig.

1b). The increased NES-LuciTs toxicity, only observed at 37 °C when it is misfolded, may be

associated with the slower clearance kinetics of the cytoplasmic misfolded variant, which is much

slower than that of misfolded NLS-LuciTs (Extended Data Fig. 1a).

We conclude that nuclear and cytoplasmic misfolded proteins are spatially sequestered in PQC

inclusions located in their cognate compartment, yielding the INQ and JUNQ respectively.

Together with findings that UPS degrades nuclear and cytoplasmic misfolded proteins through

different chaperones and ubiquitination enzymes [27], our data indicate that the nucleus and the

cytoplasm   have   compartment   specific   PQC   machineries.

Proteasome inhibition disrupts nucleocytoplasmic transport

Most commonly used PQC reporters (e.g. Ubc9Ts, VHL, Cpy*, and Luciferase, etc.), are smaller

than the 200 kDa nuclear permeability barrier [28-30]. To examine if passive diffusion across nuclear

pores could lead cytoplasmic misfolded proteins to form intranuclear inclusions, we utilized

temperature-sensitive variant of Ubc9 (Ubc9Ts) [11, 17, 18]. Upon Ubc9Ts unfolding at 37 °C for 2

hours under conditions where the proteasome is active, SIM super-resolution microscopy showed

Ubc9Ts is sequestered exclusively in two cytoplasmic compartments – the perinuclear JUNQ,

clearly outside the nuclear pore boundary and the IPOD in the cell periphery (Fig. 2a,

Supplemental Video 3). When cells were treated at 37 °C for 2 hr with added proteasome inhibitor

MG132, Ubc9Ts also formed the two cytoplasmic inclusions as above, but some cells (around

13%) had an additional Ubc9Ts inclusion inside the nucleus (Fig. 2b, Extended Data Fig. 2a,

Supplemental Video 3). Of note, MG132 treatment led to a general increase in cells containing

inclusions in both nucleus and cytoplasm (Extended Data Fig. 2a). Similar results were obtained

for another PQC substrate, VHL (Extended Data Fig. 2b). Interestingly, line intensity profile

analyses of both Ubc9Ts and VHL showed that their nuclear INQ and cytoplasmic JUNQ

inclusions formed in close proximity but separated by nuclear pores (Fig. 2b), These results

confirmed that cytoplasmic proteins that can passively diffuse into the nucleus, primarily form

cytoplasmic inclusions, but also suggested that absent an exogenous NLS or NES localization tag,

they can form an additional distinct nuclear INQ in a small fraction of cells under conditions of

proteasome inhibition.

To reconcile the finding that spatial sequestration into a nuclear or a cytoplasmic PQC inclusion

depends on a misfolded protein's subcellular location when they misfold (Fig. 1a) with the fact

that MG132 leads to a small fraction of cells having a nuclear Ubc9Ts and VHL inclusion, we

hypothesized that prolonged proteasome inhibition may alter nucleocytoplasmic trafficking and

decrease protein export from the nucleus. Indeed, toxic misfolded proteins that affect proteostasis,

such as the polyglutamine expanded mHTT, can affect nucleocytoplasmic transport [31, 32].

We examined the effect of MG132 on nucleocytoplasmic transport using an NLS- and NES-tagged

shuttling GFP (sGFP) that migrates between the nucleus and cytoplasm [31, 33] (Fig. 2c, d). As

described, toxic mHTT expression caused sGFP to be retained in the nucleus (Fig. 2c, d). Of note,

proteasome inhibition with MG132 caused sGFP nuclear retention to the same extent as mHTT

(Fig. 2c, d). This experiment was performed in leptomycin B-sensitive crm^{T539C} cells, which

permitted the use of leptomycin B (LMB) as a positive control for blocked nuclear export (Fig. 2c,

d), but similar results were obtained using WT BY4741 yeast cells (not shown).

Our finding that proteasome inhibition impairs nuclear export similar to expression of toxic mHTT

may explain the increased retention of misfolded protein Ubc9Ts in the nucleus. Future studies

should examine the basis for reducing nuclear export when proteostasis is impaired [31, 34, 35].

To directly examine if clearance of cytoplasmic proteins requires their import to the nucleus, we

examined Ubc9Ts clearance in *nup116-5 yeast* cells that. carry a temperature-sensitive mutation

in the nuclear pore complex (NPC) protein Nup116. At 37 °C, the *nup116-5* mutation seals nuclear

pores over the cytoplasmic face of NPCs blocking any transport into and out of the nucleus [36] (Fig.

2f, *i*). WT and *nup116-5* cells expressing Ubc9Ts were shifted to 37 °C for 1 hour prior to the

glucose-repression to ensure nuclear pores become completely sealed before start measuring

Ubc9Ts clearance (Figure. 2e, *ii*). If degradation of misfolded Ubc9Ts requires import into the

nucleus, it should be abrogated in *nup116-5* cells at 37 °C. We did not observe any significant

differences in the rate and the extent of Ubc9Ts degradation in *nup116-5* cells, suggesting that

transport into the nucleus is not required for its clearance (Figure. 2f, *ii-iii*; Extended Data Fig. 2c).

We next examined whether nuclear proteasomes participate in the clearance of a cytoplasmic

misfolded protein. To this end, we used *sts1-2* yeast cells that carry a mutation in Sts1, a protein

required for import of proteasomes into the nucleus and facilitate the degradation of nuclear

proteins. At 37 °C, nuclear localization of proteasomes is lost in *sts1-2* cells [37-40] (Fig. 2g panel *i*).

However, preventing nuclear localization of proteasomes had no effect on the clearance of

misfolded Ubc9Ts, consistent with cytoplasmic degradation (Fig. 2g, panel *ii-iii*; Extended Data

Fig. 2d). Therefore, together these experiments suggest that import of misfolded proteins and

proteosomes into the nucleus is dispensable to degrade misfolded Ubc9Ts.

INQ and JUNQ home into adjoined locations on opposite sides of the nuclear envelope

Given that nuclear and cytoplasmic Ubc9Ts inclusions in cells treated with MG132 appeared to

localize to opposite sides of the nuclear envelope (Fig. 2b), we next sought to define the spatial

relationship between INQ and JUNQ in live cells.

To this extent, NLS- LuciTs and NES- LuciTs were co-expressed in WT yeast cells at 25 °C for 6

hr before repressing their synthesis with glucose. Cells were then shifted to 37 °C to unfold both

proteins simultaneously, and live cell time-lapse fluorescence microscopy and particle tracking

analysis were performed to monitor inclusion formation in real-time (Fig. 3a, b; Extended Data

Fig. 3a; Supplemental Videos 4, 5). Spatial sequestration of the NLS- and NES- misfolded proteins

into motile condensates occurred with similar kinetics independently in the nucleus and the

cytoplasm. We then noticed that in some cells, the puncta moved dynamically toward each other,

with both nuclear and cytoplasmic inclusions converging on a specific location that was proximal

to the nuclear envelope (Fig. 3a). Of note, even after homing to this location, the two inclusions

remain distinct and separated; however, they co-migrated around the nuclear periphery as if

tethered to each other across the nuclear envelope. Particle tracking analyses measuring the

location and distance of NLS- and NES-LuciTs inclusions in 2D space showed the nuclear and

cytoplasmic inclusions homed into adjoining locations between 10 and 20 minutes after shifting

to 37 °C, and then continued moving in a coordinated fashion throughout the duration of the

imaging experiment (Fig. 3b; Extended Data Fig. 3a; Supplemental video 5). While the distance

between these two inclusions decreased rapidly during homing, once they came together they

maintained a largely stable short distance, suggestive of a tether, throughout their co-migration for

the remainder of the experiment. Similar results were obtained if the fluorescent markers in NLS-

and NES-LuciTs were reversed (not shown) as well as for NLS- and NES-VHL (Extended Data

Fig. 3b). These observations, supporting our findings with Ubc9Ts, suggest a novel mechanism

coordinating convergence of nuclear and cytoplasmic PQC compartments.

We confirmed the spatial relationship between the INQ and the JUNQ using SIM super-resolution

microscopy. Cells co-expressing NLS-LuciTs and NES-LuciTs were incubated at 37 °C for 2 hrs

and then fixed and immunostained to detect nuclear pores (Fig. 3c). SIM combined with line

intensity profile analysis confirmed the INQ and JUNQ adjoined each other on opposite sides of

the nuclear envelope, separated by the nuclear pores (Fig. 3c; Supplemental video 6).

206   Nuclear pores play a role in INQ-JUNQ convergence at the nuclear periphery

To understand the nature of the signal bringing together the INQ and JUNQ, we examined their

subcellular co-localization with specific nuclear and perinuclear structures. The INQ was near the

nucleolus but not in direct contact, suggesting the nucleolus is not driving this homing mechanism

(Extended Data Fig. 3c). We also examined the relationship of the INQ-JUNQ homing signal with

two complexes that localize at a defined perinuclear location: the spindle pole body (SPB) and the

LINC complex, both of which contain protein Mps3 spanning both layers of the nuclear envelope

(Extended Data Fig. 3d). Co-staining with Mps3 showed neither SPB nor LINC co-localize with

the INQ-JUNQ perinuclear site, suggesting they do not participate in the INQ-JUNQ homing

mechanism (Extended Data Fig. 3d).

To investigate the role of nuclear pores in INQ/JUNQ homing, we exploited the finding that

deletion of nuclear pore protein Nup120 causes nuclear pores to remain functional but become

clustered [41, 42]. We thus compared INQ and JUNQ formation and localization in WT and *nup120Δ*

cells. In *nup120Δ* cells, the JUNQ always forms next to the clustered nuclear pores (Fig. 3d, panel

i) while the INQ did not localize proximal to the nuclear pores, but instead was found anywhere

within the nucleus (Fig. 3d, panel *ii*). Surprisingly, if we co-expressed NLS- and NES-LuciTs in

nup120Δ cells, the INQ now homed to the clustered nuclear pores opposite the JUNQ (Fig. 3d,

panel *iii*). This finding suggests formation of the JUNQ at the cytoplasmic side of the nuclear pores

communicates a signal for INQ recruitment to this location (Fig. 3e).

We then reasoned that the intrinsically disordered phenylalanine-glycine (FG) repeats in the

central nuclear pore channel, which form hydrogels capable of amyloid-like interactions [43], could

facilitate INQ-JUNQ homing. Using live-cell time-lapse microscopy we examined the localization

of the INQ and JUNQ in *nupΔFG* cells, carrying deletions in the FG repeat tracts of six nuclear

pore proteins that removes approximately 70% of the FG content from the nuclear pore [44]. We

found that the INQ and JUNQ still localized to the same perinuclear location in *nupΔFG* cells

suggesting the high percentage of FG-repeats are dispensable for homing (Extended Data Fig. 3e).

Although, we cannot exclude the possibility that that the remaining 30% of FG content is still

sufficient to direct convergence of the INQ and JUNQ to the proximity of nuclear pores.

Cryo-soft X-ray tomography reveals the architecture of PQC inclusions in intact cells

To examine the native cellular context of PQC compartments, we used cryogenic Soft X-ray

Tomography (cryo-SXT), a synchrotron-based imaging approach that visualizes and quantifies the

ultrastructure of intact, unstained, cryo-preserved cells (Fig. 4a, Extended Data Fig. 4a, b) [45, 46].

This method uses the characteristic linear absorption of X-rays by different subcellular organelles

to produce projection images of cellular ultrastructure to a few tens nanometers of spatial

resolution [47]. To derive the information content needed to study misfolded protein inclusions we

initially used cryo-SXT in correlation with cryo-fluorescence microscopy (Fig. 4b,i). The ability

to absorb X-rays varies with the type and density of biomolecules present in different

compartments, yielding characteristic linear absorption coefficients (LAC) (Fig. 4b, ii) [48-50]. Since

chaperone Hsp104 localizes to both the JUNQ and IPOD upon heat shock, while aggregation-

prone mHTT only localizes to the IPOD [11, 20], we visualized these PQC compartments by

expressing mHTT-ChFP in cells with chromosomally tagged HSP104-GFP [51]. The inclusions

were formed by 1 hr incubation at 37 °C and cells were flash frozen in capillary tubes and directly

imaged using cryo-SXT correlated with cryo-fluorescence microscopy (Fig. 4b, i). The LAC

values were used to identify the exact location of specific compartments and organelles in X-ray

tomograms of intact frozen yeast cells (Fig. 4b, ii), yielding a 3D reconstruction of the PQC

inclusions in the context of cellular architecture (Fig. 4c, Supplemental Video 7). Cryo-

fluorescence identified the PQC compartments, with Hsp104-only marking the JUNQ and co-

localized Hsp104 and mHTT marking the IPOD. This analysis revealed that protein inclusions

have a characteristic LAC that can be used to examine their location and size without the need of

fluorescence microscopy. Of note, this correlated fluorescence-cryo-SXT analyses confirmed both

the JUNQ and the IPOD are cytoplasmic, as previously described (Fig. 4c, Supplemental Video 7

and [11]).

We next used cryo-SXT to directly detect inclusions formed by NLS- and NES-LuciTs at 37 °C

through their characteristic LAC values. NES-LuciTs formed small, cytoplasmic, punctate Q-

bodies that could be seen congregating at a cytoplasmic, juxtanuclear location to form the JUNQ

(Fig. 4d, Supplemental Video 8). Thus, cryo-SXT indicates the JUNQ is not the homogeneous

large compartment suggested by diffraction-limited fluorescence imaging, but rather consists of

multiple dense "cores" that resemble Q-bodies. This conclusion, also supported by the irregular

JUNQ morphology in SIM and cryo-EM imaging (Fig. 1, Fig 5 and [52]), suggests the JUNQ is in

fact a collection of Q-bodies that congregate at a perinuclear location proximal to the nuclear

envelope (Fig. 4d, inset *ii*; Supplemental video 8).

Cryo-SXT analysis also revealed a link between cytoplasmic PQC and mitochondria. Individual

Q-bodies can associate with the surface of mitochondria (Fig. 4e) and the JUNQ and IPOD appear

surrounded by a mitochondrial cage (Fig. 4d, *i*). Confocal fluorescence microscopy of cells

expressing mito-GFP supported the mitochondrial association with Q-bodies (Extended Data Fig.

4d). However, PQC compartments still formed in cells with disruptions in mitochondrial fission

or fusion (Extended Data Fig. 4e) indicating mitochondrial structure does not direct PQC

compartment formation. Since previous studies placed Q-bodies on the ER membrane [17, 20], which

cannot be visualized by cryo-SXT, it is possible that Q-bodies are proximal to ER-mitochondrial

contact sites. Alternatively, the proximity to mitochondria may serve to place energy-intensive

PQC compartments in areas of the cell enriched in ATP production [53].

Cryo-SXT also confirmed that the intranuclear INQ is distinct from but resides near the nucleolus

(Fig. 4g). This conclusion is supported by confocal fluorescence microscopy with nucleolar protein

Nsr1 (Extended Data Fig. 3c, 4c and [18, 31]). The proximity of the nucleolus to the INQ is consistent

with the nucleolus' role in nuclear folding and assembly [21, 54-56] and suggest a possible link to the

INQ's function in nuclear PQC [54, 57].

Strikingly, our cryo-SXT imaging revealed an unexpected link of both INQ and JUNQ with the

vacuole. The entire JUNQ site was surrounded by nuclear-vacuolar inter-organelle contact sites,

suggesting the idea that the vacuole promotes JUNQ clearance (Figure. 4f, red arrows;

Supplemental Video 8). Remarkably, Q-bodies in the JUNQ area were nestled within

invaginations in the vacuolar membrane, further suggestive of direct vacuolar engulfment (Figure.

4f, blue arrows; Supplemental Video 8). Of note, the INQ was also proximal to the vacuolar side

of the nucleus, indicating both INQ and JUNQ are steered to the vicinity of nuclear-vacuolar

contact sites (Fig. 4g). The nucleus-vacuole contact sites are formed by a tethering complex

between nuclear Nvj1 and vacuolar Vac8 called the Nuclear-Vacuolar Junction (NVJ) (Fig. 4h and

[58]). Consistent with the idea that the INQ and JUNQ converge to the vicinity of NVJs, all nuclear

pores in *nup120Δ* cells cluster at the vacuole-contacting side of the nucleus [42].

Nuclear-Vacuolar Junctions facilitate clearance of JUNQ and INQ

The role of NVJs remains poorly understood, but it has been implicated in lipid droplet biogenesis,

nuclear envelope autophagy, amino acid metabolism and a specialized form of autophagy called

Piecemeal Microautophagy of the Nucleus (PMN), where portions of the nucleus invaginate into

the vacuole [59-64]. We assessed the co-localization of the JUNQ and the NVJ by expressing NES-

LuciTs or Ubc9Ts in cells carrying a chromosomally tagged Nvj1-GFP [51]. Live-cell time-lapse

microscopy revealed transient and dynamic interactions between JUNQ and Nvj1, whereby both

puncta came in contact and then separated (Fig. 5a; Extended Data Fig. 5a; e.g. observe 9 min to

11 min in Supplemental Video 9).

The architecture of perinuclear spatial PQC and the close association of the INQ and JUNQ in

relation to the NVJ were next examined using SIM super-resolution microscopy (Fig. 5b-e;

Supplemental Videos 10, 11). The NVJ, visualized via Nvj1 imaging, forms a basket-shaped

structure contouring the Nuclear Envelope and connected to the vacuole (Fig. 5d). In these super-

resolution images, the JUNQ, visualized by NES-LuciTs, was proximal to the cytoplasmic side of

the NVJ as observed by Cryo-SXT (Fig. 5b). Of note, the Q-body "assembly" substructure of the

JUNQ was evident in these images. On the other hand, the nuclear INQ, visualized by NLS-

LuciTs, was nestled between the concave nuclear side of the NVJ and the nuclear DNA (Fig. 5c).

Strikingly, we were able to observe in some cells the INQ being extruded through the NVJ to enter

the vacuole (Fig. 5e; Supplemental Video 12). These experiments suggest dynamic tethering of

the INQ and JUNQ to the NVJ vicinity account for the homing and slight fluctuations in distance

between these inclusions observed in particle tracking experiments in Fig. 3.

We then tested if NVJs play a role in INQ and JUNQ clearance using *nvj1Δ* or *vac8Δ* yeast cells.

Live-cell time course analyses provided an unexpected insight into NVJ function. In *nvj1Δ* cells,

nuclear and diffuse NLS-LuciTs formed an initial nuclear inclusion, but during the time-course,

some cells showed the INQ being egressed from the nucleus into the cytoplasm (Fig. 5f).

Strikingly, some of the egressed NLS-LuciTs puncta ended up co-localizing with NES-LuciTs

(Fig. 5f, inset; Supplemental Video 13; white arrowheads). Similar results were obtained when

cycloheximide was added together to the temperature shift (Extended Data Fig. 5b) to ensure all

the NLS-LuciTs is in the nucleus. We observed NLS-LuciTs extruded from the nucleus in the

njv1Δ but not the WT cells, confirming NVJ disruption leads to nuclear extrusion of the INQ.

Confocal microscopy in *nvj1Δ* and *vac8Δ* cells confirmed that NVJ disruption led to extrusion of

nuclear NLS-LuciTs into the cytosol, where it ended up co-localizing with NES-LuciTs puncta

(Fig. 5g; white arrowheads). Of note, deleting the NJVs does not disrupt INQ and JUNQ

convergence, as the INQ and JUNQ are still juxtaposed in both *nvj1Δ* and *vac8Δ* cells (Fig. 5g).

We next examined if deletion of the NVJ affects clearance of PQC inclusions. Indeed, we observed

dramatic increases in the fraction of cells containing NLS and NES-LuciTs inclusions in

nvj1Δ and *vac8Δ* cells compared to WT cells (Fig. 5h, panels *i, ii*; Extended Data Fig. 5c,d), as

well as the levels of misfolded protein (Extended Data Fig. 5e). MG132 addition led to a

synergistic enhancement in the number of cells with NES-LuciTs inclusions, suggesting the NVJ

and the proteasome represent distinct cytoplasmic clearance pathways (Extended Data Fig. 5c).

The synergy effect for NLS-LuciTs inclusions was marginal, perhaps because the higher fraction

of cells with INQs poses a ceiling effect to our measurements, or because the NVJ plays a greater

role in the clearance of nuclear protein inclusions (Extended Data Fig. 5b). Further supporting a

role in misfolded protein clearance, NVJ deletion sensitizes cells to proteotoxic stress (Fig. 5i).

Both *nvj1Δ* and *vac8Δ* cells are sensitive to treatment with proline analog AZC, which induces

widespread misfolding of newly translated proteins (Fig. 5i).

The findings that the INQ and JUNQ converge on nuclear-vacuolar contact sites and that the NVJ

is required for efficient clearance of nuclear and cytoplasmic inclusions define the NVJ as a

cellular center for spatial protein quality control.

ESCRT proteins are required for INQ-JUNQ homing and clearance at the vicinity of NVJs

The relationship observed between spatial PQC and the nuclear-vacuolar contact sites led us to

examine the role of the perinuclear ESCRT-II/III protein Chm7 (CHMP7 in mammals) which

participates in nuclear envelope sealing and nuclear pore quality control [65-67] (Fig. 6a, *i*). In yeast,

Chm7 is normally diffuse throughout the cytoplasm (Extended Data Fig. 6a),and becomes

associated with a unique site in the nuclear envelope upon activation.

The active conformation of Chm7 can be induced by deleting auto-inhibitory helices in the

ESCRT-III domain [herein Chm7$_{OPEN}$] [67, 68]. Strikingly, activated Chm7$_{OPEN}$ localized to a single

site in the nuclear envelope in close proximity with both INQ and JUNQ (Fig. 6a, *ii*), suggesting

that this perinuclear ESCRT protein marks the site of INQ and JUNQ convergence for vacuolar

delivery.

We next examined the impact the absence of Chm7 had on INQ and JUNQ subcellular

localization. In *chm7Δ* yeast cells, both INQ and JUNQ still formed and migrated to the

perinuclear region, but they no longer converged to the same location (Fig. 6b), suggesting a novel

role for the Chm7 ESCRT protein as recruiting nuclear and cytoplasmic PQC compartments to the

proper location on the nuclear envelope for further clearance for terminal sequestration.

Since Chm7 is an ESCRT II/III protein, we next examined the role of other ESCRT proteins in

spatial PQC: the ESCRT-I Vps23 and the membrane bound phosphatidylinositol 3-kinase (PI3K)

Vps34-Vps15. Vps23 is a component of the ESCRT-I complex that binds to ubiquitinated cargo

proteins and mediates their vacuolar transport [69-71]. The Vps34-Vps15 complex is essential for

protein sorting to the vacuole and has been shown to localize to nuclear pores at NVJs [72-75]. We

co-expressed NLS- and NES-LuciTs in WT, *vps23Δ*, *vps34Δ*, *and vps15Δ* cells, and followed their

fate by inducing unfolding at 37 °C. We observed INQ and JUNQ formation in both WT and

ESCRT mutants. However, the spatial relationship between INQ and JUNQ was disrupted in all

three ESCRT mutants, which showed similar aberrant phenotypes (Fig. 6c, Extended Data Fig.

6b). We observed that in most of the ESCRT mutant cells, the INQ and JUNQ were no longer

homed and/or displayed cytoplasmic egress of the INQ (Fig. 6c, Extended Data Fig. 6b).

Strikingly, in a subset of *vps23Δ* and *vps34Δ* cells we observed a possible intermediate of the

nuclear egress of the INQ, whereby the NLS-LuciTs inclusion was fully enveloped by the nuclear

membrane in what appears to be a budding event (Fig. 6c, Extended Data Fig. 6b). These data

suggest that the ESCRT pathway plays a role in homing of INQ and JUNQ to the NVJ and in the

extrusion of the INQ, likely to mediate vacuolar delivery and clearance. When this process is

perturbed, as in the NVJ deletion or the ESCRT mutants, the INQ can bud into the cytoplasm.

Our analyses suggest the NVJ tethers nuclear-vacuolar contacts whereby a perinuclear ESCRT

pathway promotes vacuolar clearance of the INQ and JUNQ. Since vacuolar engulfment often

requires the AAA-ATPase Vps4, which functions by disassembling ESCRT-III complexes [76, 77]

(Fig. 6d), we next examined the effect of deleting Vps4 on INQ and JUNQ clearance as described

above. The *vps4Δ* cells contained more NLS- and NES-LuciTs inclusions than WT controls (Fig.

6e) indicating that Vps4 is required for clearance of both INQ and JUNQ. Strikingly, NLS-LuciTs

also egressed into the cytoplasm in some *vps4Δ* cells (Fig. 6f), indicating Vps4 mediates INQ

budding into the vacuole. Together, these results indicate that nuclear misfolded proteins

sequestered in the INQ are homed to the nuclear side of NVJ to be extruded out of the nucleus and

into the vacuole in an ESCRT family protein-dependent manner.

Vacuolar clearance of INQ and JUNQ

To directly examine vacuolar targeting of nuclear and cytoplasmic inclusions in yeast we utilized

the pH sensitive protein sensor Keima (Fig. 7a). Keima has a fixed emission spectrum that does

not change with pH with a maximum peak at 620 nm [78] , whilst its excitation spectrum changes

according to pH. A short wavelength spectrum with a maximum of 440 nm is predominant in a

neutral cellular environment, while a longer wavelength of 586 nm is predominant in an acidic

environment [78, 79]. Considering that Keima in yeast has been reported to have a weak emission

intensity, and to be easily bleached [79], we N-terminally tagged NLS- and NES-LuciTs with two

consecutive Keima moieties (herein 2xKeima-LuciTS; Fig. 7a, b, schematic). In the folded state,

both NLS-LuciTS and NES-LuciTS proteins were diffusely located in the nucleus and the

cytoplasm respectively with 620 nm emission observed mostly under 404 nm excitation

wavelength characteristic of a neutral pH environment. Upon misfolding at 37 °C, both NLS-

LuciTs-Keima and NES-LuciTs-Keima formed the nuclear INQ and cytoplasmic JUNQ

respectively maintaining maximum emission after 404 nm excitation, suggesting that inclusion

formation takes place initially at their respective subcellular location on a neutral pH environment.

However, upon continued incubation at 37 °C, emission at 620 nm following 558 nm excitation

starts appearing for both inclusions, characteristic of an acidic environment like the vacuole (Fig.

7c, d, Extended Data Fig. 7a, b). The kinetics of acidification of the JUNQ seemed to be slower

than for the INQ, consistent with its slower clearance kinetics (Extended Data Fig. 1a). For the

INQ, the red shift appears to initiate at the nuclear envelope boundary with an elongated structure

observed extending from the inclusion and entering the vacuole (Fig. 7c). The slower degradation

kinetics of the JUNQ coupled with the low quantum yield and quick photobleaching of Keima

precluded continuous live-imaging of the JUNQ, but an imaging time course of different regions

on the same slide showed multiple red foci arising proximal to the JUNQ (Fig. 7d; Extended Data

Fig. 7a-b). Of note, these experiments were carried out under conditions of ongoing proteasome

degradation indicating vacuolar targeting occurs in parallel to UPS-mediated clearance of

misfolded proteins.

The contribution of vacuolar and proteasomal pathways in the clearance of misfolded nuclear and

cytoplasmic proteins was independently assessed using immunoblot analyses (Fig. 7e-f). Briefly,

GFP-tagged NLS- and NES-LuciTs were expressed under permissive conditions, followed by

glucose repression, as described in Fig 1. Following unfolding, the levels of remaining luciferase

were examined after 2 and 4 hrs at 37 °C by immunoblot analyses probing for the GFP moiety.

The contribution of proteasomal clearance was assessed by addition of 50 uM Bortezomib [27, 80]

(Bz) at T=0, while the contribution of vacuolar clearance was assessed using cells deleted for Pep4,

required for maturation of vacuolar proteinases, supplemented with 1mM PMSF to inhibit vacuolar

serine proteases [81]. As expected, the levels of full length nuclear and cytoplasmic luciferase

decreased as a function of time, and were partially stabilized by proteasomal inhibition (Figure 7

e,f; Extended Data Fig. 7c-e). Inhibition of vacuolar proteases in the *pep4Δ* cells stabilized both

the nuclear and cytoplasmic full-length prote[82, 83]ins (Figure 7c-e; Extended Data Fig. 7c,e).

Interestingly, we also observed a range of GFP-positive degradation products in the WT and +Bz

cells that disappear when the vacuolar proteases were inhibited (Figure 7g), which is a hallmark

of vacuolar targeting of GFP-tagged proteins [62, 63, 81, 84-87]. Quantification of the intensity of

degradation fragments relative to initial full-length protein confirmed the reduction in clearance

when vacuolar proteases are inhibited (Fig. 7g). A longer exposure reveals fewer bands and a

completely different banding pattern between WT and protease inhibited cells (Extended Data Fig.

7c).

The finding that inhibiting vacuolar proteases stabilizes NLS-LuciTs supports the existence of a

pathway to transfer nuclear PQC substrates to the vacuole. To independently corroborate this, we

next carried out live-cell time resolveimaging of the dynamic clearance of the INQ through the

vacuole (Fig. 7g). The INQ was visualized via NLS-LuciTs-GFP and the vacuole using the styryl

dye FM4-64 commonly used to visualize endocytosis and vacuolar membrane dynamics [88] (Fig.

7g; Supplemental Video 14). Since GFP fluorescence is quenched by the acidic vacuolar

environment, NLS-LuciTS[82, 83, 89, 90] (Fig. 7g; supplemental video 14; Extended Data Fig. 7d). In

WT cells, the INQ is opposed to the vacuole at the start of the time course and appears to be pulled

into the vacuole as a function of time. Fascinatingly, when the nuclear-vacuolar contact is

disrupted in *nvj1Δ,* the INQ is unable to interact with and enter the vacuole and moves dynamically

without becoming degraded (Fig. 7g, second row; Supplemental Video 15). Strikingly, in the vast

majority of *vps4Δ* cells, the INQ looks still proximal to the vacuole but does not get cleared,

supporting a role for Vps4-mediated membrane scission for INQ vacuolar engulfment (Fig 7g,

third row; Supplemental Video 16).

DISCUSSION

Spatial sequestration of misfolded proteins into membrane-less inclusions is an integral step in

PQC. Here, we demonstrate that the cytoplasm and the nucleus form distinct compartments of

spatial PQC and that both follow a path for clearance to the vacuole when the proteasome is

overburdened. We observed that, while cytoplasmic misfolded proteins are sequestered into a

cytoplasmic perinuclear JUNQ compartment, the nuclear misfolded proteins are sequestered into

a nuclear peri-nucleolar INQ compartment. Strikingly, we find the JUNQ and INQ sometimes

converged to face each other across the nuclear envelope at a site proximal to the Nuclear-Vacuolar

Junctions (NVJ). The homing process is coordinated by nuclear pores and a perinuclear ESCRT

pathway involving ESCRT II/III protein Chm7 likely to undergo Vps4-dependent vacuolar

degradation (See integrative model in **Figure 8)**. Our experiments provide insights into logic of

spatial PQC. Upon misfolding, it is well known that misfolded proteins are cleared by independent

pathways in the nucleus and cytoplasm [27]. We show that under situations of limited proteosomal

activity, nuclear and cytoplasmically localized proteins form protein inclusions as a means to

provide an alternate specific clearance pathway. Our data support the literature demonstrating

compartment specific clearance [27,91]. Previous work from our lab has shown that there are distinct

ubiquitin linkages for the nucleus and cytoplasm [27]. Here we show that in addition to these distinct

proteostasis machineries for the UPS, the nucleus and cytoplasm form inclusions independently

and communicate to facilitate clearance through the vacuole.

Our experiments also demonstrated that the JUNQ is a conglomerate of Q-bodies as seen in [52, 92].

This finding is particularly intriguing as it alludes to these membrane-less inclusions behaving like

liquid-liquid phase separated droplets (Q-bodies) coming together to form a larger structure

(JUNQ). Notably, the Q-bodies were visible in the cryo-SXT data (Figure 4d-f) and had a distinct

linear absorption coefficient (Figure 4b) to allow them to be visualized without further

manipulation such as staining. This finding makes cryo-SXT a particularly useful tool for

examining phase separated material in the future.

Many of our experiments were done in the absence of proteasome inhibitor (Figures 1c-d, 2a, c-g,

5h, 7c-g). These experiments are critical for our understanding of sequestration as part of normal

PQC. Both the Keima (Figure 7c, d) and Western blot data from *vps4Δ* cells (Figure 7e-g) confirm

that the vacuolar protein clearance mechanism is active even when the proteasome is active (Figure

7c-g). This indicates that clearance through the vacuole is not only activated during times of

proteasomal stress as a compensatory mechanism for the loss of proteasome activity.

We observe convergence of INQ and JUNQ to the nuclear-vacuolar region (Figure 3a-c). We posit

this could represent a way to provide an important alternative clearance route upon conditions of

proteasome overload. In particular, it offers a path to clear the nuclear compartment from

misfolded proteins through extrusion of the INQ from the nucleus via the NVJ (Figure 5c-f). As

such, we observe a strong interaction between NVJ1 and the INQ by SIM and 3D reconstruction

(Figure 5c, e). The physical tether provided by NVJs mediates engulfment of regions of the nuclear

membrane into the vacuole. A similar mechanism mediating egress of the INQ into the vacuole

could explain how NVJ disruption leads to INQ extrusion into the cytoplasm.

Previous studies described a perinuclear ESCRT-II/III protein Chm7 (CHMP7 in mammals),

linked to remodeling nuclear envelopes [65], clearance of misassembled nuclear pores [66], egress of

herpes virus from the nucleus to the cytosol [93], as well as nuclear envelope sealing and nuclear

pore quality control [67]. We hypothesized that Chm7 participated in the homing of the INQ and the

JUNQ. In yeast, Chm7 is normally diffuse throughout the cytoplasm (Extended Data Fig. 5e) [67].

Upon activation Chm7 becomes associated with a unique site in the nuclear envelope (Figure 6a-

b). This active conformation can be induced by deleting auto-inhibitory helices in the ESCRT-III

domain [67] [herein Chm7$_{OPEN}$]. Strikingly, activated Chm7$_{OPEN}$ localized to a single site in the

nuclear envelope (Fig. 6a, panel *ii*; b) that co-localized with both INQ and JUNQ (Fig. 6a, panel

ii; b). These findings suggest this perinuclear ESCRT marks the site of INQ and JUNQ homing

and connects nuclear and cytoplasmic spatial PQC to vacuolar delivery. We next examined INQ

and JUNQ localization in *chm7Δ* cells. Both INQ and the JUNQ still formed and migrated to the

perinuclear region in *chm7Δ* cells, but they no longer converged to the same location (Fig. 6b).

These experiments indicate a novel role for the Chm7 ESCRT protein in recruiting nuclear and

cytoplasmic PQC compartments to the proper location on the nuclear envelope for further

clearance or terminal sequestration.

We also observe egress of the INQ from the nucleus into the cytoplasm when the nucleus becomes

untethered from the vacuole in *nvj1Δ* or *vac8Δ* mutants (Fig. 5f, g) or when vacuolar engulfment

is blocked in the *vps23Δ, vps34Δ, vps15Δ,* and *vps4Δ* mutants (Fig. 6c, f). Further, we do not see

the INQ enter the vacuole in *nvj1Δ* or *vps4Δ* mutants (Fig. 7h). Previous studies described a

perinuclear ESCRT-II/III protein Chm7 (CHMP7 in mammals), linked to remodeling nuclear

envelopes, clearance of misassembled nuclear pores, egress of herpes virus from the nucleus to the

cytosol, as well as nuclear envelope sealing and nuclear pore quality control [67]. We hypothesized

that Chm7 participates in the homing of the INQ and the JUNQ to aid in clearance via the vacuole.

This data is very intriguing and may provide a link between protein quality control and nuclear

envelope quality control. Future studies should determine whether additional forms of

microautophagy, such as nucleophagy, ERphagy, piecemeal microautophagy of the nucleus, or

selective autophagy of nuclear pore complexes participate in clearance of these PQC

compartments.

Our analyses place spatial PQC in the larger context of cellular organization and inter-organellar

communication [94] and uncover a role for nuclear pores and the nuclear envelope in communicating

nucleocytoplasmic protein quality control decisions between the nucleus, the cytoplasm and the

endolysosomal system. This idea is particularly intriguing given the link between nuclear pore

dysfunction and aging, lamin progeria mutants and neurodegenerative diseases such as

Huntington's disease, Amyotrophic Lateral Sclerosis and Frontal Temporal Dementia, which are

all characterized by nuclear pore dysfunction and altered nucleocytoplasmic trafficking [31, 34, 35, 95-

98]. Our findings raise the possibility that compromising nucleocytoplasmic trafficking leads to

impaired misfolded protein clearance contributing to further cellular dysfunction. A better

understanding of these pathways may provide fundamental insights into the critical biological

process of protein quality control and point to avenues to ameliorate a large range of human

disorders.

MAIN REFERENCES

1.   Hartl, F.U., Bracher, A. & Hayer-Hartl, M. Molecular chaperones in protein folding and

proteostasis. *Nature* **475**, 324-332 (2011).

2. Needham, P.G., Guerriero, C.J. & Brodsky, J.L. Chaperoning Endoplasmic Reticulum-

Associated Degradation (ERAD) and Protein Conformational Diseases. *Cold Spring*

Harb Perspect Biol **11** (2019).

3. Balchin, D., Hayer-Hartl, M. & Hartl, F.U. In vivo aspects of protein folding and quality

control. *Science* **353**, aac4354 (2016).

4. Sontag, E.M., Samant, R.S. & Frydman, J. Mechanisms and Functions of Spatial Protein

Quality Control. *Annu Rev Biochem* (2017).

5. Pohl, C. & Dikic, I. Cellular quality control by the ubiquitin-proteasome system and

autophagy. *Science* **366**, 818-822 (2019).

6. Finley, D. & Prado, M.A. The Proteasome and Its Network: Engineering for Adaptability.

Cold Spring Harb Perspect Biol **12** (2020).

7. Alberti, S. & Hyman, A.A. Are aberrant phase transitions a driver of cellular aging?

Bioessays (2016).

8. Ano Bom, A.P. *et al.* Mutant p53 aggregates into prion-like amyloid oligomers and

fibrils: implications for cancer. *J Biol Chem* **287**, 28152-28162 (2012).

9. Douglas, P.M. & Dillin, A. Protein homeostasis and aging in neurodegeneration. *J Cell*

Biol **190**, 719-729 (2010).

10. De Mattos, E.P. *et al.* Protein Quality Control Pathways at the Crossroad of

Synucleinopathies. *J Parkinsons Dis* **10**, 369-382 (2020).

11. Kaganovich, D., Kopito, R. & Frydman, J. Misfolded proteins partition between two

distinct quality control compartments. *Nature* **454**, 1088-1095 (2008).

12. Dillin, A. & Cohen, E. Ageing and protein aggregation-mediated disorders: from

invertebrates to mammals. *Philos Trans R Soc Lond B Biol Sci* **366**, 94-98 (2011).

13.   Balch, W.E., Morimoto, R.I., Dillin, A. & Kelly, J.W. Adapting proteostasis for disease

intervention. *Science* **319**, 916-919 (2008).

14.   Reichmann, D., Voth, W. & Jakob, U. Maintaining a Healthy Proteome during Oxidative

Stress. *Mol Cell* **69**, 203-213 (2018).

15.   Morales-Polanco, F., Lee, J.H., Barbosa, N.M. & Frydman, J. Cotranslational

Mechanisms of Protein Biogenesis and Complex Assembly in Eukaryotes. *Annu Rev*

Biomed Data Sci **5**, 67-94 (2022).

16.   Specht, S., Miller, S.B., Mogk, A. & Bukau, B. Hsp42 is required for sequestration of

protein aggregates into deposition sites in Saccharomyces cerevisiae. *J Cell Biol* **195**,

617-629 (2011).

17.   Escusa-Toret, S., Vonk, W.I. & Frydman, J. Spatial sequestration of misfolded proteins

by a dynamic chaperone pathway enhances cellular fitness during stress. *Nat Cell Biol*

(2013).

18.   Miller, S.B. *et al.* Compartment-specific aggregases direct distinct nuclear and

cytoplasmic aggregate deposition. *EMBO J* **34**, 778-797 (2015).

19.   Weisberg, S.J. *et al.* Compartmentalization of superoxide dismutase 1 (SOD1G93A)

aggregates determines their toxicity. *Proc Natl Acad Sci U S A* **109**, 15811-15816 (2012).

20.   Babazadeh, R. *et al.* Syntaxin 5 Is Required for the Formation and Clearance of Protein

Inclusions during Proteostatic Stress. *Cell Rep* **28**, 2096-2110 e2098 (2019).

21.   Nollen, E.A. *et al.* Dynamic changes in the localization of thermally unfolded nuclear

proteins associated with chaperone-dependent protection. *Proc Natl Acad Sci U S A* **98**,

12038-12043 (2001).

22. Yasuda, S. *et al.* Stress- and ubiquitylation-dependent phase separation of the

proteasome. *Nature* **578**, 296-300 (2020).

23. Polling, S. *et al.* Misfolded polyglutamine, polyalanine, and superoxide dismutase 1

aggregate via distinct pathways in the cell. *J Biol Chem* **289**, 6669-6680 (2014).

24. Park, J.H. *et al.* Amyotrophic lateral sclerosis-related mutant superoxide dismutase 1

aggregates inhibit 14-3-3-mediated cell survival by sequestration into the JUNQ

compartment. *Hum Mol Genet* **26**, 3615-3629 (2017).

25. McClellan, A.J., Scott, M.D. & Frydman, J. Folding and quality control of the VHL

tumor suppressor proceed through distinct chaperone pathways. *Cell* **121**, 739-748

(2005).

26. Gupta, R. *et al.* Firefly luciferase mutants as sensors of proteome stress. *Nat Methods* **8**,

879-884 (2011).

27. Samant, R.S., Livingston, C.M., Sontag, E.M. & Frydman, J. Distinct proteostasis circuits

cooperate in nuclear and cytoplasmic protein quality control. *Nature* **563**, 407-411

(2018).

28. Popken, P., Ghavami, A., Onck, P.R., Poolman, B. & Veenhoff, L.M. Size-dependent

leak of soluble and membrane proteins through the yeast nuclear pore complex. *Mol Biol*

Cell **26**, 1386-1394 (2015).

29. Amm, I. & Wolf, D.H. Molecular mass as a determinant for nuclear San1-dependent

targeting of misfolded cytosolic proteins to proteasomal degradation. *FEBS Lett* **590**,

1765-1775 (2016).

30. Timney, B.L. *et al.* Simple rules for passive diffusion through the nuclear pore complex.

J Cell Biol **215**, 57-76 (2016).

31.   Woerner, A.C. *et al.* Cytoplasmic protein aggregates interfere with nucleocytoplasmic

transport of protein and RNA. *Science* **351**, 173-176 (2016).

32.   Park, S.H. *et al.* PolyQ Proteins Interfere with Nuclear Degradation of Cytosolic Proteins

by Sequestering the Sis1p Chaperone. *Cell* **154**, 134-145 (2013).

33.   Neuber, A. *et al.* Nuclear export receptor Xpo1/Crm1 is physically and functionally

linked to the spindle pole body in budding yeast. *Mol Cell Biol* **28**, 5348-5358 (2008).

34.   Grima, J.C. *et al.* Mutant Huntingtin Disrupts the Nuclear Pore Complex. *Neuron* **94**, 93-

107 e106 (2017).

35.   Gasset-Rosa, F. *et al.* Polyglutamine-Expanded Huntingtin Exacerbates Age-Related

Disruption of Nuclear Integrity and Nucleocytoplasmic Transport. *Neuron* **94**, 48-57 e44

(2017).

36.   Wente, S.R. & Blobel, G. A temperature-sensitive NUP116 null mutant forms a nuclear

envelope seal over the yeast nuclear pore complex thereby blocking nucleocytoplasmic

traffic. *J Cell Biol* **123**, 275-284 (1993).

37.   Chen, L. *et al.* Sts1 plays a key role in targeting proteasomes to the nucleus. *J Biol Chem*

286, 3104-3118 (2011).

38.   Takeda, K. & Yanagida, M. Regulation of nuclear proteasome by Rhp6/Ubc2 through

ubiquitination and destruction of the sensor and anchor Cut8. *Cell* **122**, 393-405 (2005).

39.   Tatebe, H. & Yanagida, M. Cut8, essential for anaphase, controls localization of 26S

proteasome, facilitating destruction of cyclin and Cut2. *Curr Biol* **10**, 1329-1338 (2000).

40.   Romero-Perez, L., Chen, L., Lambertson, D. & Madura, K. Sts1 can overcome the loss of

Rad23 and Rpn10 and represents a novel regulator of the ubiquitin/proteasome pathway.

J Biol Chem **282**, 35574-35582 (2007).

41.   Aitchison, J.D., Blobel, G. & Rout, M.P. Nup120p: a yeast nucleoporin required for NPC

distribution and mRNA transport. *J Cell Biol* **131**, 1659-1675 (1995).

42.   Heath, C.V. *et al.* Nuclear pore complex clustering and nuclear accumulation of poly(A)+

RNA associated with mutation of the Saccharomyces cerevisiae RAT2/NUP120 gene. *J*

Cell Biol **131**, 1677-1697 (1995).

43.   Ader, C. *et al.* Amyloid-like interactions within nucleoporin FG hydrogels. *Proc Natl*

Acad Sci U S A **107**, 6281-6285 (2010).

44.   Strawn, L.A., Shen, T., Shulga, N., Goldfarb, D.S. & Wente, S.R. Minimal nuclear pore

complexes define FG repeat domains essential for transport. *Nat Cell Biol* **6**, 197-206

(2004).

45.   Parkinson, D.Y. *et al.* Nanoimaging cells using soft X-ray tomography. *Methods Mol*

Biol **950**, 457-481 (2013).

46.   Le Gros, M.A. *et al.* Biological soft X-ray tomography on beamline 2.1 at the Advanced

Light Source. *J Synchrotron Radiat* **21**, 1370-1377 (2014).

47.   Larabell, C.A. & Nugent, K.A. Imaging cellular architecture with X-rays. *Curr Opin*

Struct Biol **20**, 623-631 (2010).

48.   Le Gros, M.A., McDermott, G., Uchida, M., Knoechel, C.G. & Larabell, C.A. High-

aperture cryogenic light microscopy. *J Microsc* **235**, 1-8 (2009).

49.   Ekman, A. *et al.* Putting Molecules in the Picture: Using Correlated Light Microscopy

and Soft X-Ray Tomography to Study Cells, in *Synchrotron Light Sources and Free-*

Electron Lasers: Accelerator Physics, Instrumentation and Science Applications. (eds. E.

Jaeschke, S. Khan, J.R. Schneider & J.B. Hastings) 1-32 (Springer International

Publishing, Cham; 2019).

50.   Ekman, A.A. *et al.* Mesoscale imaging with cryo-light and X-rays: Larger than molecular

647   machines, smaller than a cell. *Biol Cell* **109**, 24-38 (2017).

648   51.   Huh, W.K. *et al.* Global analysis of protein localization in budding yeast. *Nature* **425**,

649   686-691 (2003).

650   52.   Wu, G.H. *et al.* Multi-scale 3D Cryo-Correlative Microscopy for Vitrified Cells.

651   *Structure* **28**, 1231-1237.e1233 (2020).

652   53.   Schuler, M.H. *et al.* Miro1-mediated mitochondrial positioning shapes intracellular

653   energy gradients required for cell migration. *Mol Biol Cell* **28**, 2159-2169 (2017).

654   54.   Frottin, F. *et al.* The nucleolus functions as a phase-separated protein quality control

655   compartment. *Science* **365**, 342-347 (2019).

656   55.   Velazquez, J.M. & Lindquist, S. hsp70: nuclear concentration during environmental

657   stress and cytoplasmic storage during recovery. *Cell* **36**, 655-662 (1984).

658   56.   Falahati, H., Pelham-Webb, B., Blythe, S. & Wieschaus, E. Nucleation by rRNA Dictates

659   the Precision of Nucleolus Assembly. *Curr Biol* **26**, 277-285 (2016).

660   57.   Azkanaz, M. *et al.* Protein quality control in the nucleolus safeguards recovery of

661   epigenetic regulators after heat shock. *Elife* **8** (2019).

662   58.   Pan, X. *et al.* Nucleus-vacuole junctions in Saccharomyces cerevisiae are formed through

663   the direct interaction of Vac8p with Nvj1p. *Mol Biol Cell* **11**, 2445-2457 (2000).

664   59.   Kohlwein, S.D. *et al.* Tsc13p is required for fatty acid elongation and localizes to a novel

665   structure at the nuclear-vacuolar interface in Saccharomyces cerevisiae. *Mol Cell Biol* **21**,

666   109-125 (2001).

60.   Levine, T.P. & Munro, S. Dual targeting of Osh1p, a yeast homologue of oxysterol-

binding protein, to both the Golgi and the nucleus-vacuole junction. *Mol Biol Cell* **12**,

1633-1644 (2001).

61.   Kvam, E. & Goldfarb, D.S. Structure and function of nucleus-vacuole junctions: outer-

nuclear-membrane targeting of Nvj1p and a role in tryptophan uptake. *J Cell Sci* **119**,

3622-3633 (2006).

62.   Kvam, E. & Goldfarb, D.S. Nucleus-vacuole junctions and piecemeal microautophagy of

the nucleus in S. cerevisiae. *Autophagy* **3**, 85-92 (2007).

63.   Roberts, P. *et al.* Piecemeal microautophagy of nucleus in Saccharomyces cerevisiae.

Mol Biol Cell **14**, 129-141 (2003).

64.   Krick, R. *et al.* Piecemeal microautophagy of the nucleus requires the core

macroautophagy genes. *Mol Biol Cell* **19**, 4492-4505 (2008).

65.   von Appen, A. *et al.* LEM2 phase separation promotes ESCRT-mediated nuclear

envelope reformation. *Nature* **582**, 115-118 (2020).

66.   Thaller, D.J. & Patrick Lusk, C. Fantastic nuclear envelope herniations and where to find

them. *Biochem Soc Trans* **46**, 877-889 (2018).

67.   Webster, B.M. *et al.* Chm7 and Heh1 collaborate to link nuclear pore complex quality

control with nuclear envelope sealing. *EMBO J* **35**, 2447-2467 (2016).

68.   Thaller, D.J. *et al.* An ESCRT-LEM protein surveillance system is poised to directly

monitor the nuclear envelope and nuclear transport system. *Elife* **8** (2019).

69.   Babst, M., Odorizzi, G., Estepa, E.J. & Emr, S.D. Mammalian tumor susceptibility gene

101 (TSG101) and the yeast homologue, Vps23p, both function in late endosomal

trafficking. *Traffic* **1**, 248-258 (2000).

70.   Katzmann, D.J., Babst, M. & Emr, S.D. Ubiquitin-dependent sorting into the

multivesicular body pathway requires the function of a conserved endosomal protein

sorting complex, ESCRT-I. *Cell* **106**, 145-155 (2001).

71.   Herrador, A., Léon, S., Haguenauer-Tsapis, R. & Vincent, O. A mechanism for protein

monoubiquitination dependent on a trans-acting ubiquitin-binding domain. *J Biol Chem*

288, 16206-16211 (2013).

72.   Stack, J.H., Herman, P.K., Schu, P.V. & Emr, S.D. A membrane-associated complex

containing the Vps15 protein kinase and the Vps34 PI 3-kinase is essential for protein

sorting to the yeast lysosome-like vacuole. *EMBO J* **12**, 2195-2204 (1993).

73.   Herman, P.K. & Emr, S.D. Characterization of VPS34, a gene required for vacuolar

protein sorting and vacuole segregation in Saccharomyces cerevisiae. *Mol Cell Biol* **10**,

6742-6754 (1990).

74.   Burda, P., Padilla, S.M., Sarkar, S. & Emr, S.D. Retromer function in endosome-to-Golgi

retrograde transport is regulated by the yeast Vps34 PtdIns 3-kinase. *J Cell Sci* **115**,

3889-3900 (2002).

75.   Slessareva, J.E., Routt, S.M., Temple, B., Bankaitis, V.A. & Dohlman, H.G. Activation

of the phosphatidylinositol 3-kinase Vps34 by a G protein alpha subunit at the endosome.

Cell **126**, 191-203 (2006).

76.   Henne, W.M., Stenmark, H. & Emr, S.D. Molecular mechanisms of the membrane

sculpting ESCRT pathway. *Cold Spring Harb Perspect Biol* **5** (2013).

77.   Hurley, J.H. & Hanson, P.I. Membrane budding and scission by the ESCRT machinery:

it's all in the neck. *Nat Rev Mol Cell Biol* **11**, 556-566 (2010).

78.   Katayama, H., Kogure, T., Mizushima, N., Yoshimori, T. & Miyawaki, A. A sensitive

713   and quantitative technique for detecting autophagic events based on lysosomal delivery.

714   *Chem Biol* **18**, 1042-1052 (2011).

715   79.   Bialecka-Fornal, M., Makushok, T. & Rafelski, S.M. A Review of Fluorescent Proteins

716   for Use in Yeast. *Methods Mol Biol* **1369**, 309-346 (2016).

717   80.   Sitron, C.S., Park, J.H., Giafaglione, J.M. & Brandman, O. Aggregation of CAT tails

718   blocks their degradation and causes proteotoxicity in S. cerevisiae. *PLoS One* **15**,

719   e0227841 (2020).

720   81.   Schuck, S., Gallagher, C.M. & Walter, P. ER-phagy mediates selective degradation of

721   endoplasmic reticulum independently of the core autophagy machinery. *J Cell Sci* **127**,

722   4078-4088 (2014).

723   82.   Haupts, U., Maiti, S., Schwille, P. & Webb, W.W. Dynamics of fluorescence fluctuations

724   in green fluorescent protein observed by fluorescence correlation spectroscopy. *Proc Natl*

725   *Acad Sci U S A* **95**, 13573-13578 (1998).

726   83.   Shinoda, H., Shannon, M. & Nagai, T. Fluorescent Proteins for Investigating Biological

727   Events in Acidic Environments. *Int J Mol Sci* **19** (2018).

728   84.   Klionsky, D.J. *et al.* Guidelines for the use and interpretation of assays for monitoring

729   autophagy (4th edition). *Autophagy* **17**, 1-382 (2021).

730   85.   Eising, S. *et al.* A lysosomal biogenesis map reveals the cargo spectrum of yeast vacuolar

731   protein targeting pathways. *J Cell Biol* **221** (2022).

732   86.   Waite, K.A., De-La Mota-Peynado, A., Vontz, G. & Roelofs, J. Starvation Induces

733   Proteasome Autophagy with Different Pathways for Core and Regulatory Particles. *J Biol*

734   *Chem* **291**, 3239-3253 (2016).

87.   Kvam, E. & Goldfarb, D.S. Nvj1p is the outer-nuclear-membrane receptor for oxysterol-

binding protein homolog Osh1p in Saccharomyces cerevisiae. *J Cell Sci* **117**, 4959-4968

(2004).

88.   Vida, T.A. & Emr, S.D. A new vital stain for visualizing vacuolar membrane dynamics

and endocytosis in yeast. *J Cell Biol* **128**, 779-792 (1995).

89.   Thorn, K. Genetically encoded fluorescent tags. *Mol Biol Cell* **28**, 848-857 (2017).

90.   Costantini, L.M. *et al.* A palette of fluorescent proteins optimized for diverse cellular

environments. *Nat Commun* **6**, 7670 (2015).

91.   Amm, I., Sommer, T. & Wolf, D.H. Protein quality control and elimination of protein

waste: the role of the ubiquitin-proteasome system. *Biochim Biophys Acta* **1843**, 182-196

(2014).

92.   Eisele, F. *et al.* An Hsp90 co-chaperone links protein folding and degradation and is part

of a conserved protein quality control. *Cell Rep* **35**, 109328 (2021).

93.   Kumar, B. *et al.* ESCRT-I Protein Tsg101 Plays a Role in the Post-macropinocytic

Trafficking and Infection of Endothelial Cells by Kaposi's Sarcoma-Associated

Herpesvirus. *PLoS Pathog* **12**, e1005960 (2016).

94.   Gottschling, D.E. & Nystrom, T. The Upsides and Downsides of Organelle

Interconnectivity. *Cell* **169**, 24-34 (2017).

95.   Zhang, K. *et al.* The C9orf72 repeat expansion disrupts nucleocytoplasmic transport.

Nature **525**, 56-61 (2015).

96.   Fallini, C., Khalil, B., Smith, C.L. & Rossoll, W. Traffic jam at the nuclear pore: All

roads lead to nucleocytoplasmic transport defects in ALS/FTD. *Neurobiol Dis* **140**,

104835 (2020).

97.   Shang, J. *et al.* Aberrant distributions of nuclear pore complex proteins in ALS mice and

759   ALS patients. *Neuroscience* **350**, 158-168 (2017).

760   98.   Chou, C.C. *et al.* TDP-43 pathology disrupts nuclear pore complexes and

761   nucleocytoplasmic transport in ALS/FTD. *Nat Neurosci* **21**, 228-239 (2018).

FIGURE TITLES AND LEGENDS

Figure 1: The INQ and JUNQ are separate nuclear and cytoplasmic PQC compartments.

(a) Schematic showing nuclear and cytoplasmic misfolded proteins could be sequestered into

separate compartments or the cytoplasmic misfolded proteins could be imported into the nucleus

and sequestered into the nuclear PQC compartment INQ. (b) Experimental schematic.

Temperature sensitive Luciferase is properly folded at 25 °C. Heat shock at 37 °C leads to

unfolding and sequestration of the protein. Degradation of the unfolded and sequestered proteins

can be blocked by treatment with the proteasome inhibitor MG132. (c, d) Live-cell time-lapse

fluorescence microscopy of WT cells expressing NLS-LuciTs (c) or NES-LuciTs (d) at 37 °C,

treated with DMSO (top) or 100μ<u>M</u> MG132 (bottom). Representative still frames at the times

shown. Scale bars are 1μm. (e, f) Representative SIM images of WT cells expressing NLS-

LuciTs (e) or NLS-VHL (f) after 2 hr at 37 °C and treated with 100μ<u>M</u> MG132. NLS-LuciTs

and NLS-VHL are shown in green, nuclear pores in gold, and Hoechst counterstain in blue. Scale

bars are 1μm. Line intensity profiles indicate relative locations of subcellular compartments to

Nups and DNA. (g, h) Representative SIM images of WT cells expressing NES-LuciTs (g) and

NES-VHL (h) after 2 hr at 37 °C and treated with 100μ<u>M</u> MG132. NES-LuciTs and NES-VHL

are shown in purple, nuclear pores in gold, and Hoechst counterstain in blue. Scale bars are 1μm.

Line intensity profiles indicate relative locations of subcellular compartments to Nups and DNA.

Figure 2: Nuclear entry of misfolded proteins is not required for clearance. .

(a) Representative SIM images of WT cells expressing Ubc9Ts-GFP after 2h at 37 °C treated

with DMSO (a) or 100µM MG132 (b). Ubc9Ts-GFP is shown in green, nuclear pores in gold,

and Hoechst counterstain in blue. Scale bars are 1µm. Line intensity profiles indicate relative

locations of subcellular compartments to Nups and DNA. (c) Representative confocal

fluorescence microscopy images of cells expressing sGFP with DMSO (control), 100µM MG132

treatment, co-expression of mHTT97QΔP or 200nM Leptomycin B treatment. Ratio of

cytoplasmic:nuclear fluorescence is shown in d. Kruskal-Wallis test with Dunn's multiple

comparisons test was performed using Prism. Adjusted p value of No Treatment vs. MG132 is

0.0014, No Treatment vs. 97QΔP is 0.0027 and No Treatment vs. LMB is <0.0001. A minimum

of 500 cells per condition from 3 biologically independent experiments were analyzed. All scale

bars are 1µm. (e) Schematic illustrating the clearance of Ubc9Ts in WT yeast. (*ii*) Timeline of

treatments for clearance measurements with shift to 37 °C 60 minutes before initiation of the

measurements. (f) Schematic illustrating the *nup116-5* yeast have sealed nuclear pores at 37 °C,

thus blocking nucleocytoplasmic trafficking. (*ii*) Densitometric quantification of Western blot

bands (*iii*) measuring the amount of Ubc9Ts-EGFP remaining in shut-off experiment of WT vs

nup116-5 cells relative to $t = 0$ (mean ± s.e.m. from three biologically independent experiments)

fitted with a one-phase decay non-linear fit regression line. (g) Schematic illustrating the *sts1-2*

yeast do not translocate proteasomes to the nucleus at 37 °C. (*ii*) Densitometric quantification of

Western blot bands (iii) measuring the amount of Ubc9Ts-EGFP remaining in shut-off

experiment of WT vs sts1-2 cells relative to $t = 0$ (mean ± s.e.m. from three biologically

independent experiments) fitted with a one-phase decay non-linear fit regression line.

Figure 3: INQ and JUNQ home to similar location on each side of the nuclear envelope via a cytoplasmic signal linked to nuclear pores.

WT cells co-expressing NLS-GFP-LuciTs and NES-DsRed-LuciTs were shifted to 37 °C, treated with 100µ$\underline{M}$ MG132 and monitored by live cell time-lapse fluorescence. Representative still frames at the times shown. Scale bar is 1µm. (b) Graph of the distance between the INQ and JUNQ compartments by particle tracking of inclusions from cell shown in (a) over the time course of the experiment. The slight variations of tethered distance between 60 and 90 minutes is likely due to the relative migration of one of the inclusions around a cellular structure as this fluctuation can be seen at different times post-tethering in other cells (not shown). (c) Representative SIM image taken of cells co-expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 2 hr at 37 °C and treated with 100µ$\underline{M}$ MG132. NLS-fusion proteins are shown in green, NES-fusion proteins in purple, nuclear pores in gold, and Hoechst counterstain in blue. Scale bar is 1µm. Line intensity profiles indicate relative locations of subcellular compartments and Nups. (d) Representative confocal microscopy images taken of WT and 3 separate *nup120Δ* yeast cells expressing NES-LuciTs (*i*), NLS-LuciTs (*ii*) or co-expressing NLS- and NES-LuciTs (*iii*) after 2 hr at 37 °C. NLS-LuciTs is shown in green, NES-LuciTs in purple, nuclear pores in gold, and Hoechst counterstain in blue. Scale bar is 1µm. Schematics on the right summarize the findings of the data. (e) Schematic summarizing the data that the JUNQ localizes to the nuclear pores. A signal is transmitted through or near the nuclear pores to recruit the INQ to the same location, resulting in the homing of the 2 compartments.

Figure 4: INQ resides near the nucleolus, JUNQ is surrounded by mitochondria, and both

compartments home into the Nuclear-Vacuolar Junction.

(a) Unfixed, intact yeast cells are frozen in a capillary tube and imaged by fluorescence

microscopy and then X-ray tomography. The linear absorption coefficients are measured from

the tomography data and used to identify the subcellular compartments and organelles. The

fluorescence images are then correlated to the X-ray tomography data and 3D reconstructions of

the subcellular components are generated to determine the cellular context of the PQC

compartments. (b) WT cells co-expressing Hsp104-GFP from the endogenous locus and

HTT97QP-ChFP after 30 mins at 37 °C. (left) X-ray tomograms without and with fluorescence

overlay to indicate sites of Hsp104-GFP positive inclusions and HTT97QP-ChFP inclusion.

(right) Table of Linear Absorption Coefficient (LAC) values from the X-ray tomograms used to

annotate the images and generate a 3D reconstruction shown in panel c. (c) 3D reconstruction of

fluorescence correlated X-ray tomograms showing the JUNQ residing 450nm outside the barrier

of the nucleus as defined by the location of the nuclear envelope in the X-ray data (inset, top)

and the IPOD marked by mHTT (inset, bottom). JUNQ is shown in green, IPOD in orange,

nucleus in yellow, mitochondria in cyan, vacuoles in gray, and lipid droplets in white. Scale bar

on X-ray 3D reconstruction is 1μm, insets are 0.5μm. (d) 3D reconstruction of X-ray tomograms

from cell expressing NES-LuciTs after 90 mins heat shock at 37 °C and treated with 100μM

MG132. The Q-bodies can be seen to coalesce into the JUNQ compartment just outside the

nuclear boundary (inset ii). The Q-bodies also interact with vacuoles and the site of coalescence

into the JUNQ compartment is surrounded by a mitochondrial cage (inset i). Q-bodies and IPOD

are shown in purple, nucleus in yellow, mitochondria in cyan, vacuoles in gray, and lipid

droplets in white. Scale bar on X-ray 3D reconstruction is 1μm, insets are 0.5μm. (e) Q-bodies

can be seen interacting with mitochondria in a separate cell expressing NES-LuciTs. Q-bodies

are shown in purple, nucleus in yellow, nucleolus in gold, mitochondria in cyan, and vacuoles in

gray. Scale bar on X-ray 3D reconstruction is 1μm, inset is 0.5μm. (f) Same 3D reconstruction of

X-ray tomograms from cell expressing NES-LuciTs after 90 mins heat shock at 37 °C as shown

in d. Insets are from the same cell rotated 180°. Blue arrows indicate sites where Q-bodies are

directly interacting with the vacuoles. Red arrows indicate sites of nuclear-vacuolar junctions.

Scale bar on X-ray 3D reconstruction is 1μm, insets are 0.5μm. (g) 3D reconstruction of X-ray

tomograms from cell expressing NLS-LuciTs after 90 mins heat shock at 37 °C and treated with

100μM MG132. The INQ resides 400nm from the nucleolus (inset). INQ is shown in green,

nucleus in yellow, nucleolus in gold, mitochondria in cyan, and vacuoles in gray. Scale bar on X-

ray 3D reconstruction is 1μm, inset is 0.5μm. (h) Schematic illustrating the novel findings of the

association between mitochondria and all 3 cytoplasmic PQC compartments, the proximity of the

INQ to the nucleolus, the interaction between Q-bodies and the vacuole, and the hypothesized

location of the INQ and JUNQ at the nuclear vacuolar junction. The inset highlights two of the

component proteins of the NVJ, Nvj1 and Vac8, that are required for Piecemeal Microautophagy

of the Nucleus.

Figure 5: JUNQ and INQ converge at the Nuclear-Vacuolar Junction to facilitate

clearance.

(a) Endogenously tagged Nvj1-GFP yeast expressing NES—DsRed-LuciTs were shifted to 37

°C and monitored by live cell time-lapse fluorescence microscopy for the times shown. White

arrowheads indicate locations of Nvj1 punctum while yellow arrowheads indicate NES-LuciTs

punctum. Scale bar is 1μm. Representative SIM images of WT yeast co-expressing NES-DsRed-

LuciTs (b) or NLS-DsRed-LuciTs (c) and Nvj1-sfGFP after 2 hr incubation at 37 °C with

100μM MG132. Panel *ii* of both (b) and (c) are insets to better visualize the relative location of

the JUNQ and INQ to the Nvj1. Panel *iii* of both (b) and (c) are additional cells to illustrate other

phenotypes seen in the experiment. NES-DsRed-LuciTs is shown in purple, NLS-DsRed-LuciTs

is shown in green, Nvj1-sfGFP is shown in yellow, and DNA is shown in blue. Scale bars are 1

μm. (d) Representative confocal image of WT yeast expressing Nvj1-sfGFP (shown in yellow)

stained with FM4-64 vacuolar dye (shown in cyan). Scale bar is 1μm. (e) Representative SIM

image of WT yeast cell co-expressing NLS-DsRed-LuciTs (shown in green) with Nvj1-sfGFP

(shown in yellow) after 2 hr incubation at 37 °C with 100μM MG132. The NLS-LuciTs can be

seen extruding through the NVJ toward the vacuole. Scale bar is 1μm. (f) WT (top) and *nvj1Δ*

(bottom) cells co-expressing NLS-LuciTs and NES-LuciTs were shifted to 37 °C, treated with

100μM MG132 and monitored by live cell time-lapse fluorescence microscopy for the times

shown. White arrowheads indicate inclusions of NLS-LuciTs pulled into the cytoplasm. Scale

bars are 1μm. (g) Representative confocal images of WT, *nvj1Δ*, and *vac8Δ* yeast co-expressing

NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 2 hr at 37 °C and treated with 100μM MG132.

NLS-EGFP-LuciTs is shown in green, NES-DsRed-LuciTs in purple, nuclear pores in gold, and

Hoechst counterstain in blue. White arrows indicate cytoplasmic localization of NLS-LuciTs.

Scale bar is 1μm. (h) Quantitation of the percentage of cells containing inclusions in WT, *nvj1Δ*,

and *vac8Δ* yeast co-expressing NLS-EGFP-LuciTs (*i*) and NES-DsRed-LuciTs (*ii*) after 2 hr

incubation at 37 °C with 100μM MG132. A minimum of 500 cells per condition from 3

biologically independent experiments were counted and unpaired Student's t-tests were

performed comparing the deletion strains to WT. P values were adjusted using two-stage linear

step-up procedure of Benjamini, Krieger, and Yekutieli with a false discovery rate (Q) of 5%.

Adjusted P value for NLS-LuciTs: WT vs *nvj1Δ* is 0.0205, WT vs *vac8Δ* is 0.0406. Adjusted P

value for NES-LuciTs: WT vs *nvj1Δ* is 0.013, WT vs *vac8Δ* is 0.0008. (i) Drop tests showing

serial dilutions of WT, *nvj1Δ*, and *vac8Δ* yeast at 30 and 37 °C with no treatment and 0.5mg/mL

AZC treatment after 72 h growth at indicated temperature. *nvj1Δ* and *vac8Δ* yeast have a growth

defect when subjected to proteostatic stress by AZC treatment.

Figure 6: ESCRT-mediated extrusion from the nucleus and clearance.

(a) The ESCRT-II/-III protein Chm7 has been shown to play a role in clearance of defective

nuclear pores and nuclear membrane quality control, therefore, it may be involved in the homing

of INQ and JUNQ to the NVJ. (*ii*) Representative confocal images of WT yeast co-expressing

Chm7$_{OPEN}$-EGFP and either NLS-DsRed-LuciTs (top) or NES-DsRed-LuciTs (bottom) after 2 hr

at 37 °C and treated with 100μM MG132. Chm7$_{OPEN}$ is shown in teal, NLS-DsRed-LuciTs in

green, NES-DsRed-LuciTs in purple, nuclear pores in gold and Hoechst counterstain in blue.

Arrows indicate locations of puncta for each protein. Scale bar is 1μm. (b) Representative

confocal images of WT and *chm7Δ* yeast co-expressing NLS-EGFP-LuciTs and NES-DsRed-

LuciTs after 120 minutes at 37 °C and treated with 100μM MG132. NLS-EGFP-LuciTs is shown

in green, NES-DsRed-LuciTs in purple, nuclear pores in gold, and Hoechst counterstain in blue.

Arrows indicate locations of puncta for each protein. Scale bar is 1μm. (c) ESCRT-family

proteins Vps23, Vps34, and Vps15 may be involved in clearance of INQ and JUNQ.

Representative confocal images of WT and *vps23Δ* , *vps34Δ,* and *vps15Δ* yeast co-expressing

NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 2 hr at 37 °C and treated with 100μM MG132.

NLS-EGFP-LuciTs is shown in green, NES-DsRed-LuciTs in purple, nuclear pores in gold, and

Hoechst counterstain in blue. Insets show the budding INQ encapsulated by nuclear pores. Scale

bars are 1μm. (d) ESCRTIII proteins and the ATPase Vps4 are known to remodel membranes

and could be involved in vacuolar import of PQC compartments. (e) Quantitation of the

percentage of cells containing inclusions of NLS- or NES-LuciTs in WT and *vps4Δ* yeast co-

expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 2 hr incubation at 37 °C with

100μM MG132. A minimum of 300 cells per condition from 2 biologically independent

experiments were counted and Student's T tests were performed comparing the deletion strains

to WT. P value for WT vs *vps4Δ* NLS-LuciTs is 0.0007, and WT vs *vps4Δ* NES-LuciTs is

0.0018. (f) Representative confocal images of WT and *vps4Δ* yeast co-expressing NLS-EGFP-

LuciTs and NES-DsRed-LuciTs after 2 hr at 37 °C and treated with 100μM MG132. NLS-

EGFP-LuciTs is shown in green, NES-DsRed-LuciTs in purple, nuclear pores in gold, and

Hoechst counterstain in blue. Green arrows indicate cytoplasmic localization of NLS-LuciTs.

Scale bar is 1μm.

Figure 7: Vacuolar clearance of the INQ and JUNQ

(a) Possible routes of entry into the vacuole for the INQ and JUNQ. (b) Schematic of the LuciTs-Keima experiments. (c) WT cells expressing NLS-2xKeima-LuciTs after 2 hr incubation at 37 °C with 100μM MG132. Over time, fluorescence is seen with excitation in the 589nm channel indicating the NLS-LuciTs has encountered an acidic environment. Inset shows the transition from green to red and a structure leaving the inclusion that is fully red. Scale bars are 5 uM. (d) Representative images of WT cells expressing NES-2xKeima-LuciTs after 2 hr incubation at 37 °C with 100μM MG132. Over time, fluorescence is seen with excitation in the 589nm channel indicating the NLS-LuciTs has encountered an acidic environment. Insets show the transition from green to red and a structure leaving the inclusion that is fully red. Scale bars are 5 uM. Representative Western Blots of NLS-GFP-LuciTs (e) and NES-GFP-LuciTs (f) in WT and *pep4Δ* yeast. *pep4Δ* yeast were also treated with 1mM PMSF to completely inhibit vacuolar proteases. A decrease in fragments can be seen in both the NLS- (e) and NES- (f) blots. (g) Densitometric quantification of Western blot bands measuring the amount of full-length LuciTs-GFP remaining in shut-off experiment as well as the fragments generated during clearance. The ratio of fragment intensity to full-length protein is shown in the graph (mean ± s.e.m. from three biologically independent experiments). One-way ANOVA was performed using Prism software followed by Dunnett's multiple comparisons test. Adjusted P value for WT vs *pep4Δ* NLS-LuciTs is 0.0018, and WT vs pep*4Δ* NES-LuciTs is 0.0220. (h) WT, *nvj1Δ,* and *vps4Δ* yeast expressing NLS-GFP-LuciTs were treated with 8uM of FM4-64 and incubated for 2hr at 37 °C with 100μM MG132. Cells were imaged every 5 mins for 90 mins. Scale bar is 1μm.

Figure 8: Principles of Nuclear and Cytoplasmic spatial protein quality control Model for vacuolar targeting and clearance of nuclear and cytoplasmic protein inclusions in yeast. (a) Upon heat shock, cytoplasmic and nuclear LuciTs locally misfold, and recruit chaperones and other PQC factors that facilitate the subcellular spatial sequestration on distinct protein quality control inclusions: (b) the Intranuclear Quality Control compartment (INQ) in the nucleus and Q-bodies in the cytoplasm.(c) Q-bodies then coalesce to form the Juxtanuclear Quality Control compartment (JUNQ), and misfolded proteins located in the INQ and the JUNQ can be degraded by the UPS system in the nucleus and the cytoplasm respectively c.ii). (d) Alternatively, under conditions of limited proteasomal activity, INQ and JUNQ can converge on the periphery of the nuclear envelope, a mechanism mediated by the nuclear pores (homing)(e) The homing mechanism could represent a coordinated way to target both inclusions for clearance at the vacuole. (e.ii) Vacuolar mediated clearance of both INQ and JUNQ. ESCRT-family proteins are involved in organizing the INQ and JUNQ for vacuolar degradation.

METHODS

Time resolved live cell imaging. Cells were grown overnight in raffinose media followed by 4–6 h at 28 °C in galactose media to induce the expression. Cells were then immobilized on concanavalin A-coated coverslips. Samples were washed in glucose media and kept in the same medium by sealing the coverslips to slides with vacuum grease. When indicated, cells were treated 10 min before sealing with 100 uMof MG132 (Sigma) before sealing. Images were taken every 15 s for 60 min at 37 °C (or room temperature (25 °C)) using a Zeiss Axio Observer.Z1 inverted microscope equipped with X-cite 120 LED light source (Lumen Dynamics), HE GFP/Cy3/DAPI shift free filter sets (Zeiss), a Plan-Apochromat 100x/1.4 oil DIC M27 objective (Zeiss) and a digital Axiocam MRm camera (Zeiss) controlled with the Zen blue software.

Cycloheximide chase of preformed inclusions. Cells were grown overnight in raffinose media,

back diluted to an OD600 of 0.1 in a 10 ml of galactose media and grown at 25 °C for 5 hours to

induce expression of both NLS and NES constructs. Cells were then resuspended in 10 ml of

glucose media at 100 uM MG132 and incubated for additional 15 min 25 °C and then switched to

37 °C for 30 minutes to ensure inclusion formation before CHX chase. Cells were then

immobilized on concanavalin deltaT culture dish (Fischer Scientific), washed with prewarmed

glucose media and incubated with glucose media supplemented with 100 uM MG132 and 50mg/ml

of cycloheximide (CHX). Images were taken every 25 s for 120 min at 37 °C as mentioned above.

FM4-64 vacuole labeling. 10 ml of cells were grown on galactose media to a final OD600 of 0.8

at 25°C. Vacuolar dye was added by resuspending the cells on 2 ml of fresh galactose media plus

FM4-64 to a final concentration of 8uM, and incubated 30 min at 25°C. For imaging, cells were

then resuspended in glucose media at 100uM MG132 and incubated 20 min at 37°C to induce

inclusion formation. Cells were adhered on a concanavalin deltaT culture dish and imaged as

mentioned above.

Immunostaining. As in the live-cell imaging assay, cells were grown overnight in raffinose media

followed by 4–6 h at 28 °C in galactose media to induce the expression. Then, the cells were heat-

shocked at 37 °C in glucose media for 1 h before fixation in 4% paraformaldehyde for 15 mins at

37 °C followed by methanol fixation for 20 mins at -20 °C. Immunostaining proceeded as in [1].

Briefly, samples were resuspended in sorbitol buffer and spheroplasted in Zymolyase (Zymo) for

30 mins at room temperature and further solubilized in 0.1% Triton X-100 for 10 mins. Antibodies

are diluted in Buffer WT (1% nonfat dry milk 0.5 mg/ml BSA 200 mM NaCl 50 mM HEPES–

KOH (pH 7.5) 1 mM NaN3 0.1% Tween-20) at a 1:500 dilution and incubated 2 hr at room

temperature or overnight at 4 °C. Samples are then washed in Buffer WT followed by incubation

in secondary antibodies at a 1:1000 dilution for 2 hr at room temperature. Cells were immobilized

on poly-lysine coated coverslips and mounted in Prolong Diamond mounting media (with DAPI

where indicated) (Thermo-Fisher). Nanobodies against GFP and RFP were conjugated to Alexa

Fluor 488 and Alexa Fluor 568 respectively and used to amplify the signal from the fluorescent

proteins during the SIM measurements. Anti-Nsp1 antibody from EnCor Biotechnology was used

to visualize nuclear pores and Anti-Nsr1 antibody from Abcam was used for staining the nucleolus.

Confocal microscopy. Confocal microscopy on immunostained samples was performed on a

Zeiss LSM 700 inverted confocal microscope with 405, 488, 555 and 639 nm laser lines with Basic

Filterset for LSM700 (Zeiss), a Plan-Apochromat 100x/1.4 oil DIC WD=0.17 objective (Zeiss)

and a digital Axiocam MRm camera (Zeiss) controlled with the Zen black software. Confocal data

sets were deconvolved when indicated using the Zen software with a fast-iterative function.

Fluorescence was quantified where indicated using ImageJ software (NIH). The sGFP intensity

measurements, were performed as described previously [2]. Briefly, the raw integrated density

(herein indicated as intensity) was measured for the nucleus and the whole cell. The intensity value

of the nucleus was then divided by the area. The cytoplasmic intensity was calculated by

subtracting the nuclear intensity from the whole cell intensity and the cytoplasmic area by

subtracting the nuclear area from the whole cell area. The cytoplasmic intensity was then divided

by the cytoplasmic area. Finally, the ratio of cytoplasmic to nuclear was calculated by dividing the

cytoplasmic intensity per area by the nuclear intensity per area. Statistical analysis was performed

using Prism (GraphPad).

Super-resolution microscopy. Structured illumination microscopy (SIM) on immunostained

samples was performed using an OMX V4 Blaze system (Applied Precision, GE Healthcare)

equipped with a 100×/1.40 NA PlanApo oil-immersion objective (Olympus), 405, 488, 568 and

647 nm diode lasers with standard filter sets, and 3 emCCD cameras (Photometrics Evolve 512).

SIM data were acquired with a Z-distance of 125 nm and with 15 images per plane (five phases,

three angles). The raw data was computationally reconstructed using a Wiener (high-frequency)

filter setting of between 0.001 and 0.003 and channel- specific OTFs employing the softWoRx

version 6.5.2 software package (GE Healthcare) to obtain a super-resolution image stack. 3D

reconstructions and line intensity profiles were generated in Volocity version 6.3 (Perkin Elmer).

To construct *NVJ1-sfGFP* strain with integrated sfGFP fusion at the C-terminal, sf*GFP* was

knocked-in via standard PCR-based homologous recombination. The *sfGFP* cassette was

amplified from plasmid GTL-g (Addgene plasmid #81099) with oligonucleotides

5'TAGATGCACAAGTGAACACTGAACAAGCATACTCTCAACCATTTAGATACGCGGC
CGCTCTAGAACTA and

5' TCGCACCTCGTTGTAAGTGACGATGATAACCGAGATGACGGAAATATAGTACA

ATGGAAAAACGCCAGCAACG. The cassette was amplified, separated by electrophoresis, gel
purified and transformed into yeast cells using the lithium acetate method. Correct transformants
were verified by standard PCR using the oligonucleotides NVJ1-VF:
5'GGATACCAGAACAACCTCTTC, NVJ1-VR: 5' ATGCCCGGCGTTATATATTGC, and
LEU2-VF: 5' AGCACGAGCCTCCTTTACCT.

Keima cloning. The NLS-Keima-luciTS plasmid was constructed by replacing the GFP moeity

with yeast optimized Keima using a cassette amplified from plasmid pFA6a-link-yomKeima-

Kan (Addgene plasmid #44902) with the oligonucleotides

5'ATGGCTTCTCCTAAGAAGAAACGTAAAGTTATGGTTTCTGTGATCGCTAAACAAAT

GAC and

5'CCATGCAAGCTTGCGCGGATCCGCGCCCTAATAGAGAATGTCTTGCGATAGC.

Similarly, the NES-Keima-luciTS plasmid was created replacing the GFP moity using the

olinucleotides

5'CTCGCACTTAAGTTCGCCGGTTTAGACCTGATGGTTTCTGTGATCGCTAAACAAAT

GAC and

5'CCATGCAAGCTTGCGCGGATCCGCGCCCTAATAGAGAATGTCTTGCGATAGC. To

construct the NLS and NES-2xKeima-luciTS, an additional Keima moeity was created by

inserting a second keima cassette amplified using the oligonucleotides

5'CCATGCAAGCTTGCGCGGATCCGCGCCCTAATAGAGAATGTCTTGCGATAGC and

5'TTCCATGCAAGCTTGCGCGGATCCGCGCCCTAATAGAGAATGTCTTGCGA.

Live cell LeicaSp8/sSTED microscopy of Keima-tagged inclusions. Cells were grown as for

time resolve live fluorescent microscopy, adhered on concanavalin deltaT culture dish and

examined at 37 °C for 90 minutes. Representative cells were collected every 5 minutes on a

Leica TCS SP8 inverted sSTED microscope equipped with a 100x/1.40 APO objective and using

the following detection mirror settings: Keima 590-650nm. Cells were sequentially excited using

404nm and 568 nm laser. One representative middle slide was acquired, and images were

deconvolved and background subtracted using Huygens Professional (Scientific Volume

Imaging).

Xray-tomography:

Preparation of Specimen Mounting Capillaries. Thin-walled glass capillaries were pulled as

described in [3]. After being cut to length, capillaries were dipped in poly-L-lysine (0.01% solution,

Tissue Culture Grade, Sigma Aldrich, St. Louis, MO). Fiducial markers, used to align soft x-ray

projection images to a common axis, were generated by briefly immersing the capillaries a solution

of 100 nm gold nanoparticles (EMGC100, BBI International, Cardiff, CF14 5DX, UK), rinsing

and then drying them in air.

Specimen capillaries for use in correlated imaging experiments were prepared as above and then

dipped in a suspension of red polystyrene microspheres (FluoSpheres Carboxylate-Modified

Microspheres, 0.2 µm, Dark Red Fluorescent (660 excitation/ 680 emission), Life Technologies,

Invitrogen). The polystyrene microspheres were used fiducial markers for the alignment of

fluorescence images to each other and for the co-alignment of soft x-ray and cryo- fluorescence

tomographic reconstructions.

Specimen Mounting. 1 ml of yeast cells in mid-log phase growth (OD_{600} ~0.5) were pelleted and

re-suspended in ~50µl media. 1µl of this suspension was transferred into a specimen capillary

using a standard micropipette. The capillary contents were immediately cryopreserved by rapid

plunging into liquid propane at 165°C using a custom apparatus. Cryopreserved specimens were

transferred into storage boxes using a home-built cryo-transfer device and stored in a Dewar held

at 77K [3].

Imaging Cells in the Cryogenic Confocal Light Microscope (CLM). Specimen capillaries were

transferred cryogenically to a home-built cryogenic spinning disk confocal fluorescence

microscope [4]. Confocal scanning and detection were achieved using a commercial dual spinning

disk head (CSU-X1, Yokogawa, Tokyo, Japan). During imaging, the illumination wavelength was

selected with an acousto-optical tunable filter (AOTF) using an integrated system (Andor Laser

Combiner, Model LC-501A).

Experimental measurement of the CLM point spread function. The point spread function of

the cryogenic confocal light microscope was calculated by the methods described in [5].

Acquisition of Cryogenic Fluorescence Tomography data. Cryopreserved specimen capillaries

were cryo-transferred to the CLM and aligned with respect to the rotation axis. Through-focus, or

"z-stacks" were taken using a step size of 0.78 μm. For fluorescence tomography, the capillary

was rotated about an axis normal to the objective lens using a motorized goniometer driven by

custom LabView code (National Instruments Corporation, Austin, TX). Each tomographic data

series consisted of 37 through-focus stacks (0° through 360° measured at 10° increments). Once

measurements were completed, the capillary was cryo-transferred back to liquid nitrogen storage

using the cryo-transfer device [5].

Alignment of through-focus fluorescence stacks using fluorescence fiducials. Well-isolated

fluorescent fiducials were chosen manually from the raw fluorescence stacks and a three-

dimensional centroid algorithm refined the bead position to sub-pixel accuracy using the software

package Amira (FEI). The fiducial coordinates were used to write a new image stack in MATLAB

containing a spherically-symmetrical representation of the fiducials, which we termed the "fiducial

model". Six parameters were optimized for each alignment: three translations and three rotations

(a rigid affine transformation). By limiting the search space to optimize correspondence between

bead pairs that had been visually verified to be true fiducial correspondences, this alignment

strategy was relatively fast and robust. The entire fluorescence fiducial alignment optimization

process was visualized in Amira. Visualization was helpful for inspecting the results and

troubleshooting rare cases where the optimization settled into a local maximum that was obviously

not the real solution.

Fluorescence tomographic reconstruction and deconvolution. Preprocessed, aligned through-

focus stacks were reconstructed into a single object by simply summing the through-focus datasets

(Amira). The effective PSF for tomographic imaging in a specimen capillary was used as the

convolution kernel to deconvolve the fluorescence reconstruction. We used an iterative function

for the deconvolution. The algorithm found the fluorophore distribution that was most likely,

assuming the PSF was spatially invariant, and that image formation followed Poisson statistics.

Soft x-ray data collection and reconstruction. Specimens were imaged using a transmission soft

x-ray microscope, operated by the National Center for X-ray Tomography (NCXT), at the

Advanced Light Source of Lawrence Berkeley National Laboratory and reconstructed according

to previously published protocols.

Correlated Imaging. Co-alignment of cryogenic fluorescence and soft x-ray tomographic

reconstructions using joint fiducials

Joint fiducials - red fluorescent polystyrene microspheres - visible in both light- and soft x-ray

images were used to guide overlay of the two data types. The alignment process was optimized by

converting the raw fluorescence into a bead model and aligning the fluorescent bead model to the

x-ray reconstruction of the beads using an iterative optimization of the overlay.

Identification of JUNQ/IPOD in soft x-ray reconstructions. Segmentation is the process of

computationally isolating, visualizing, and quantifying specific cellular components in a

tomographic reconstruction. Each voxel in an SXT reconstruction is a direct measurement of the

soft x-ray Linear Absorption Coefficient (LAC) at the corresponding location in the cell.

Attenuation of the illumination by the specimen follows Beer's Law. Hence, the LAC values for

identical sized voxels depends solely on the concentration and composition of biomolecules

present. Consequently, water has an order of magnitude lower LAC than carbon-containing

molecules, such as lipids and proteins. LAC values for homogeneous solutions of isolated

biomolecules can also be calculated using tables of known absorption coefficients. For example,

pure water in the form of ice has a calculated LAC of 0.109 μm^{-1} whereas a model protein with

the chemical composition $C_{94}H_{139}N_{24}O_{31}S$ was calculated to have a theoretical LAC of $1.35\mu m^{-1}$

[6]. In practice, most of the voxels in a 50 nm resolution SXT reconstruction of a cell will contain a

heterogeneous mixture of biomolecules. Using SXT data acquired at a single wavelength it is not

possible to distinguish the precise chemical species present. However, at this level of spatial

resolution organelles and other sub-cellular structures are sufficiently similar in their biochemical

composition to allow them to be readily identified from the surrounding cell contents. The

relatively high water content in vacuoles makes them readily distinguishable from organelles with

a greater density of biomolecules, for example, nuclei and mitochondria. Even relatively small

variations in organelle LAC can be distinguished, for example, the boundaries between nuclei and

nucleoli are very clear, as is the distinction between euchromatin and heterochromatin domains in

the nucleus. LACs were calculated, and cells segmented using the protocols describe in complete

detail in [3].

Segmentation of the Hsp104-GFP reconstructions, in particular the assignment of the volumes as

JUNQ and IPOD, were guided by correlative fluorescence signal. In other strains, JUNQ and/or

IPOD were assigned based on LAC values established in the Hsp104-GFP study.

Western blot. Yeast cultures were grown to mid- log-phase in synthetic complete medium (yeast

nitrogen base, ammonium sulfate, amino acids, 2% raffinose) lacking uracil. Cultures were then

diluted to an OD_{600} of 0.2 in galactose medium (synthetic complete medium with 2% galactose

and 2% raffinose) and grown to an OD_{600} of 0.8. Cultures were then pelleted and resuspended in

glucose medium (synthetic complete medium with 2% glucose) and incubated at either 30 or 37

°C as indicated for the duration of the shut off. Where indicated, cultures were shifted to 37 °C for

1 hour prior to exchange into glucose medium and incubated at 37 °C for the duration of the

glucose shut-off. Cells were pelleted, proteins were extracted with urea and SDS, and protein

concentrations were measured with the bicinchoninic acid protein assay kit (Thermo Fisher

Scientific). Equal amounts of total protein were resolved on Tris-glycine gels (Invitrogen) and

transferred onto nitrocellulose membranes. GFP was detected with mouse anti-GFP antibodies

(Roche), GAPDH was detected with rabbit anti-GAPHD (Genetex), and PGK was detected with

rabbit anti-PGK antibodies (Thermo Fisher Scientific). Secondary antibodies were IRDye 800CW

donkey anti-mouse or IRDye 680RD donkey anti-rabbit and were imaged using the LiCOR

scanner and Image Studio Software.

To assess vacuolar protein degradation, the same procedure as above was followed but the *pep4D*

yeast were treated with 1mM PMSF (Research Products International) during the 37C incubation

step. The samples were then collected and lysed as above. GFP and GAPDH were detected with

the same primary antibodies, but the secondary antibodies were goat anti-mouse HRP conjugated

antibody (Promega Cat #W4021) and goat anti-rabbit HRP conjugated antibody (Promega Cat #

W4011). Blots were detected using Clarity Western ECL Substrate (Bio-Rad Cat# 1705061) and

imaged using the GE Amersham Imager 600.

Drop test. Yeast cultures were grown to OD_{600} of 0.6 in YPD or raffinose synthetic complete

medium and then diluted to OD_{600} of 0.1. Four 10-fold serial dilutions were performed in H_2O.

~2uL of diluted cultures were spotted onto YPD, galactose, or glucose plates using the pin plater.

Plates were incubated at indicated temperatures and imaged every 24 hours for 72 hours.

Nanobody purification and labeling. GFP-nanobody was expressed in BL21 Rosetta2 pLysS

cells (Novagen), and DsRed-nanobody (LAM4) was expressed in ArticExpress cells (Agilent

Technologies) overnight at 17 °C and 12 °C respectively. Cells were pelleted (4000XG), washed

with PBS containing 1 mM PMSF and pelleted again. Pellets were resuspended in column buffer

(500 mM NaCl and 50 mM HEPES pH 8 or 50 mM $NaHCO_3$ pH 8.3) with complete protease

inhibitors, PMSF, benzonase and 10 mM imidazole (Lysis buffer) and lysed using an emulsiflex.

Lysates were cleared at 20,000XG for 30 minutes. Nanobodies were purified using nickel resin

by passing cleared lysate over the resin, washing with lysis buffer, followed by column buffer, and

elution with column buffer containing 300 mM imidazole. Elution was concentrated using an

Amicon ultra 3 kD MWCO to less than 2 mL, and further purified by size exclusion using an

SDX75 column (GE Healthcare) equilibrated with 500 mM NaCl, 50 mM NaHCO$_3$, 1 mM DTT.

Labeling was performed by the addition of dyes at a molar ratio of 5:1 dye:nanobody using Alexa

488 NHS (Thermo Fisher Scientific) for the GFP nanobody and Alexa 568 NHS (Thermo Fisher

Scientific) for the DsRed nanobody. Labeling reactions were incubated 1 hr at room temperature

and quenched by addition of Tris-HCl pH 8 to a final concentration of 100 mM. Labeled nanobody

was separated from free dye by size exclusion using an SDX75 column.

GFP-nanobody cloning. pDG402, a vector for expression of GFP-nanobody-TEV-6XHis was

generated by PCR amplification of the GFP nanobody using the 5' primer (XbaI GFP nanobody)

AATCTAGAATTTTGTTTAACTTTAAGAAGG and 3' primer (GS link TEV His nanobody)

AAGGATCCTTATCAGTGATGATGGTGGTGATGGGACCCAGATCCACCCTGGAAGTAT

AAATTCTCACCCGAACCTCCGGATGAGGAGACGGTGACCTGGG. The PCR product

was cloned into the pST39 backbone [7] using XbaI and BamHI.

METHODS REFERENCES

1   Brickner, D. G., Light, W. & Brickner, J. H. Quantitative localization of chromosomal
loci by immunofluorescence. *Methods Enzymol* **470**, 569-580, doi:10.1016/S0076-
6879(10)70022-7 (2010).

2   Woerner, A. C. *et al.* Cytoplasmic protein aggregates interfere with nucleocytoplasmic
transport of protein and RNA. *Science* **351**, 173-176, doi:10.1126/science.aad2033 (2016).

3   Parkinson, D. Y. *et al.* Nanoimaging cells using soft X-ray tomography. *Methods Mol
Biol* **950**, 457-481, doi:10.1007/978-1-62703-137-0_25 (2013).

4   Le Gros, M. A., McDermott, G., Uchida, M., Knoechel, C. G. & Larabell, C. A. High-
aperture cryogenic light microscopy. *J Microsc* **235**, 1-8, doi:10.1111/j.1365-2818.2009.03184.x
(2009).

5   Le Gros, M. A. *et al.* Biological soft X-ray tomography on beamline 2.1 at the Advanced
Light Source. *J Synchrotron Radiat* **21**, 1370-1377, doi:10.1107/s1600577514015033 (2014).

6   Weiss, D. *et al.* Computed tomography of cryogenic biological specimens based on X-ray
microscopic images. *Ultramicroscopy* **84**, 185-197, doi:10.1016/s0304-3991(00)00034-6 (2000).

7   Tan, S. A modular polycistronic expression system for overexpressing protein complexes

in Escherichia coli. *Protein Expr Purif* **21**, 224-234, doi:10.1006/prep.2000.1363 (2001).

ACKNOWLEDGMENTS

We thank support of NIH (GM05643319 and AG054407 to JF; F32NS086253 to ES) and Way

Klingler Startup Funds from Marquette University (ES). F.M-P. was supported by The Pew Trusts

in the Biomedical Sciences postdoctoral Award (00034104). Cryo-SXT data were acquired at the

Advanced Light Source of Lawrence Berkeley National Laboratory, a U.S. DOE Office of Science

User Facility. The National Center for X-ray Tomography is supported by NIH (P41GM103445)

and DOE BER (DE-AC02-5CH11231). CAL and MAL were supported by the Gordon and Betty

Moore Foundation Award #3497. The Stanford Cell Sciences Imaging Facility is supported in part

by Award Number 1S10OD01227601 from the National Center for Research Resources (NCRR).

We thank Jon Mulholland and Youngbin Lim from the CSIF for training on the SIM. We are

grateful to Michael Rosbash (Brandeis University), Susan Wente (Vanderbilt University), Kiran

Madura (Rutgers University), and Patrick Lusk (Yale School of Medicine) for providing yeast

strains and Karsten Weis (ETH Zurich), Jodi Nunnari (University of California, Davis), and

Michael P. Rout (Rockefeller University) for plasmids. We thank Katharine Ullman (University

of Utah), Adam Frost (Altos Lab) and Lisbeth Veenhoff (University of Groningen) for discussions

and advice. We are grateful to Carson Trail for support in microscopy data analysis, Margaret

Wangeline for assisting with the 2xKeima cloning, Joan Steffan (UC Irvine), Kelly Rainbolt

(Stanford University) for discussions and comments on the manuscript, Felipe Serrano for assisting

on Figure 8, and the Frydman lab for advice and discussions.

AUTHOR CONTRIBUTIONS

JF, ES conceived the project; ES and F. M-P carried out all experiments; J-HC collected and

processed cryo-SXT data; GM carried out cryo-SXT data analysis and modelling; MLG

performed cryo-fluorescence data acquisition and correlation with cryo-SXT data and built the

microscope used for these experiments. PD analyzed particle tracking data and assisted on

statistical analyses. DG cloned the NLS- and NES- luciferase and VHL plasmids. DG and F. M-

P generated, purified and labeled the GFP and RFP nanobodies. ES, F. M-P and JF wrote the

MS. All authors commented on the final version. JF directed the project.

DECLARATION OF INTERESTS

1239 The authors declare no competing interests.

ADDITIONAL INFORMATION

1241 Supplementary information is available for this paper.

1242 Correspondence and requests for materials should be addressed to Judith Frydman

1243 jfrydman@stanford.edu

EXTENDED DATA FIGURE LEGENDS

1245

Figure ED1: Spatial sequestration occurs during different types of stress with different

client proteins.

(a) Western blot analyses of Gal Shut-off assays showing the clearance of NLS-LuciTs (top) and

NES-LuciTs (bottom) with and without proteasome impairment by 50u$\underline{M}$ Bortezomib. (b) Drop

test of W303 yeast expressing model proteins without heat shock at 30C (left), with heat shock at

37 °C (middle), and without expression of the plasmids (right). (c) Representative Structured

Illumination super-resolution microscopy images taken of cells expressing NLS-VHL (left) or

NES-VHL (right) after 120 minutes at 37 °C and treated with 100μ$\underline{M}$ MG132. NLS-LuciTs is

shown in green, NES-LuciTs in purple, nuclear pores in gold and Hoechst counterstain in blue.

Scale bars are 1μm.

1256

Figure ED2: The effect of blocking nucleocytoplasmic transport on Ubc9Ts clearance.

(a) Quantitation of the percentage of cells containing nuclear or cytoplasmic inclusions in WT

yeast expressing Ubc9Ts-EGFP after 120 minutes at 37 °C with and without treatment with

100μM MG132. A minimum of 500 cells per condition from 3 biologically independent

experiments were counted and Student's t-tests were performed comparing the WT yeast without

MG132 treatment to WT yeast with MG132 treatment using Prism software. P values were

adjusted using two-stage linear step-up procedure of Benjamini, Krieger and Yekutieli with a Q

of 5%. Adjusted P value for nuclear no MG132 vs. +MG132 is 0.0035 and cytoplasmic no

MG132 vs. +MG132 is 0.0011. (b) Representative Structured Illumination super-resolution

microscopy images taken of cells expressing GFP-VHL after 2 hr at 37 °C with DMSO (left) or

with 100μM MG132 (right) treatment. VHL is shown in green, nuclear pores in gold, and

Hoechst counterstain in blue. Scale bars are 1μm. (c) (left) Schematic illustrating the *nup116-5*

yeast have sealed nuclear pores at 37 °C, thus blocking nucleocytoplasmic trafficking. (right)

There is no significant difference in half-life between WT and *nup116-5* yeast. (d) (left)

Schematic illustrating the *sts1-2* yeast do not translocate proteasomes to the nucleus at 37 °C.

(right) There is no significant difference in half-life between WT and *sts1-2* yeast. (c and d)

Levels of EGFP were quantified from 3 biologically independent experiments. Half-life was

calculated using Prism.

1275

Figure ED3: INQ-JUNQ homing does not occur at the LINC, nucleolus, or involve FG

repeats of the nuclear pore central channel.

(a) Graph of the X-Y positions of the INQ and JUNQ compartments by particle tracking of

inclusions from cell shown in Figure 2a over the time course of the experiment. (b)

Representative confocal image taken of cells co-expressing NLS-EGFP-VHL and NES-DsRed-

VHL after 2 hr at 37 °C and treated with 100μM MG132. NLS-fusion proteins are shown in

green, NES-fusion proteins in purple, nuclear pores in gold, and Hoechst counterstain in blue.

Scale bar is 1μm. (c) Representative confocal fluorescence microscopy images taken of cells co-

expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs (left) after 2 hr at 37 °C and treated with

100μM MG132. NLS-LuciTs is shown in green, NES-LuciTs in purple, nucleolus (Nsr1) in gold

and Hoechst counterstain in blue. (right) Line intensity profile showing distance between

nucleolus and homed INQ/JUNQ. Scale bars are 1μm. (d) (left) schematic of Mps3 component

of LINC complex linking inner and outer nuclear membranes. (right) Representative widefield

fluorescence microscopy images taken of cells co-expressing endogenously tagged Mps3-EGFP

and NES-DsRed-LuciTs after 120 minutes at 37 °C with and without treatment with 100μM

MG132. White arrowheads indicate locations of Mps3 puncta while yellow arrowheads indicate

NES-LuciTs puncta. Scale bars are 1μm. (e) WT (top) and *nupΔFG* (bottom) cells co-expressing

NLS-LuciTs and NES-LuciTs were shifted to 37 °C and monitored by live cell time-lapse

fluorescence microscopy for the times shown. Scale bars are 1μm.

1295

Figure ED4: Detailed representation of the cryo-SXT workflow and interactions between

mitochondria and cytoplasmic PQC compartments.

(a) Optical path through the specimen. Key: COL, cryogenic objective lens; SS, specimen stage;

SP, specimen port; MG, motorized goniometer; CIM, cryogenic immersion fluid; CCL, low

magnification cryogenic objective; CS, cryogenic specimen; CIE, cryogenic imaging

environment; AP, adapter port; AW, a heated, angled anti-reflection window. (b) Alignment of

fluorescence and soft x-ray tomographic data using fiducial markers. (c) A representative

confocal image of the spatial relationship between the INQ and nucleolus. NLS-LuciTs (INQ) is

shown in green, nucleolus in gold, and Hoechst counterstain in blue. Scale bar is 1μm. (d) The

interaction between mitochondria and cytoplasmic inclusions is also seen by fluorescence

confocal microscopy in a representative image of a cell co-expressing mito-GFP and NES-RFP-

LuciTs. NES-LuciTs is shown in purple, mitochondria in cyan, and Hoechst counterstain in blue.

Scale bar is 1μm. (e) Representative confocal fluorescence microscopy images taken of WT,

fission mutants (*dnm1Δ* and *fis1Δ*) and fusion mutant (*fzo1Δ* and *ugo1Δ*) cells expressing mito-

GFP and NES-DsRed-LuciTs after 120 minutes at 37 °C and treated with 100μM MG132. Mito-

GFP is shown in cyan, NES-LuciTs in purple, and Hoechst counterstain in blue. Scale bars are

1μm.

1313

Figure ED5: NVJ -mediated clearance of misfolded proteins.

(a) Endogenously tagged Nvj1-GFP yeast expressing Ubc9Ts-ChFP were shifted to 37 °C and

monitored by live cell time-lapse fluorescence microscopy for the times shown. White

arrowheads indicate locations of Nvj1 puncta while yellow arrowheads indicate Ubc9Ts-ChFP

puncta. Scale bar is 1μm. (b) WT (top) and *nvj1Δ* (bottom) cells co-expressing NLS-LuciTs and

NES-LuciTs were treated with 100μM MG132 and shifted to 37 °C for 30 mins to preform

inclusions. Cells were then placed in media containing 50mg/ml cycloheximide (CHX) and

100μM MG132 at 37 °C and monitored by live cell time-lapse fluorescence microscopy for the

times shown. Scale bars are 1μm. (c,d) Quantitation of the percentage of cells containing

cytoplasmic inclusions in WT, *nvj1Δ*, and *vac8Δ* yeast co-expressing NLS-EGFP-LuciTs (c) and

NES-DsRed-LuciTs (d) after 2 hr at 37 °C with and without treatment with 100μM MG132. A

minimum of 500 cells per condition from 3 biologically independent experiments were counted

and unpaired Student's t-tests were performed comparing the deletion strains to WT with or

without MG132 treatment were compared. P values were adjusted using two-stage linear step-up

procedure of Benjamini, Krieger and Yekutieli with a Q of 5%. (c) %. Adjusted P value for WT

vs *nvj1Δ* no MG132 is 0.0205, WT vs *nvj1Δ* +MG132 is 0.2357, WT vs *vac8Δ* no MG132 is

0.0406, WT vs *vac8Δ* +MG132 is 0.8974, and WT no MG132 vs WT +MG132 is 0.0239. (d)

Adjusted P value for WT vs *nvj1Δ* no MG132 is 0.013, WT vs *nvj1Δ* +MG132 is 0.013, WT vs

vac8Δ no MG132 is 0.0008, WT vs *vac8Δ* +MG132 is 0.0447, and WT no MG132 vs WT

+MG132 is 0.0005. (e) Levels of EGFP after 1 hour at 37 °C were measured from Quantitative

Western blots (mean ± s.e.m. from three biologically independent experiments). WT and nvj1Δ

yeast were compared using an unpaired Student's t-test resulting in a P value of 0.0043 for WT

vs nvj1Δ NLS-LuciTs and NES-LuciTs P value is 0.0183. Statistics were performed using Prism.

1337

Figure ED6: ESCRT involvement in the clearance of misfolded proteins.

(a) Representative confocal images of WT yeast co-expressing Chm7-EGFP and either NLS-

EGFP-LuciTs (left) or NES-DsRed-LuciTs (right) after 120 minutes at 37 °C and treated with

100μM MG132. Chm7 is shown in teal and remains diffuse throughout the cell, NLS-EGFP-

LuciTs in green, NES-DsRed-LuciTs in purple, nuclear pores in gold and Hoechst counterstain

in blue. Scale bar is 1μm. (b) Representative confocal images of WT and *vps23Δ* ,

vps34Δ, and *vps15Δ* yeast co-expressing NLS-EGFP-LuciTs and NES-DsRed-LuciTs after 2 hr

at 37 °C and treated with 100μM MG132. NLS-EGFP-LuciTs is shown in green, NES-DsRed-

LuciTs in purple, nuclear pores in gold, and Hoechst counterstain in blue. Insets show the

budding INQ encapsulated by nuclear pores. Scale bars are 1μm. Same data as shown in Figure

6c, but with the green channel separated to clearly detail the colocalization with the cytoplasmic

protein.

Figure ED7: Vacuole-mediated clearance of INQ and JUNQ.

(a) Representative images of WT cells expressing NES-2xKeima-LuciTs after 2 hr incubation at

37 °C with 100μM MG132. Over time, fluorescence is seen with excitation in the 589nm channel

indicating the NLS-LuciTs has encountered an acidic environment. Insets show the transition

from green to red and a structure leaving the inclusion that is fully red. Scale bars are 5 uM. Same

data shown in Figure 7d, but with more time points and a larger field of view in the images. (b)

WT cells expressing NES-2xKeima-LuciTs after 85 min incubation at 37 °C with 100μM

MG132. (c) Longer exposure of the blot shown in Figure 7f to highlight the difference in the

number and pattern of the GFP bands in the WT vs *pep4Δ* cells. (d) Levels of EGFP at time 0

were measured from Quantitative Western blots such as those shown in Figure 7e,f (mean ±

s.e.m. from three biologically independent experiments). WT and *pep4Δ* yeast were compared

using an unpaired Student's t-test without reaching statistical significance. (e) WT yeast

expressing NLS-GFP-LuciTs were treated with 8uM of FM4-64 and incubated for 2hr at 37 °C

with 100μM MG132. Cells were imaged every 30 sec for 90 mins. Scale bar is 1μm. Same data

shown in Figure 7h, but only WT and with more timepoints during the entry into the vacuole.

1365

1366

Legends to movies

Movie 1. Live-cell time-lapse fluorescence microscopy of WT cells expressing NLS-LuciTs

(left) or NES-LuciTs (right) at 37 °C, treated with 100μM MG132. Same data shown in stills in

Figure 1c (left), d (right).

Movie 2. Dynamic representation of the 3D reconstructions of the data shown in Figure 1e

(left),g (right). Videos were created in Volocity.

Movie 3. Dynamic representation of the 3D reconstructions of the data shown in Figure 2a (left),

b (right). Videos were created in Volocity.

Movie 4. Live-cell time-lapse fluorescence microscopy of the data shown in Figure 3a.

Movie 5. Dynamic representation of the data shown in Figure 3b.

Movie 6. Dynamic representation of the 3D reconstruction of the data shown in Figure 3c. Video

was created in Volocity.

Movie 7. Dynamic representation of the 3D reconstruction shown in Figure 4c. Video was

created in Amira.

Movie 8. Dynamic representation of the 3D reconstruction shown in Figure 4d,f. Video was

created in Amira.

Movie 9. Live-cell time-lapse fluorescence microscopy of the data shown in Figure 5a.

Movie 10. Dynamic representation of the 3D reconstruction of the data shown in Figure 5c.

Video was created in Volocity.

Movie 11. Dynamic representation of the 3D reconstruction of the data shown in Figure 5b.

Video was created in Volocity.

Movie 12. Live-cell time-lapse fluorescence microscopy of the data shown in Figure 5f.

Movie 13. Dynamic representation of the 3D reconstruction of the data shown in Figure 5e.

Video was created in Volocity.

Video 14. Live-cell time-lapse fluorescence microscopy of the WT cell data shown in Figure 5h.

Video 15. Live-cell time-lapse fluorescence microscopy of the *nvj1Δ* cell data shown in Figure

5h.

Video 16. Live-cell time-lapse fluorescence microscopy of the *vps4Δ* cell data shown in Figure

5h.

a Misfolding → Sequestration

i JUNQ / INQ / or *ii* INQ

b LuciferaseTs: Nuclear (NLS) vs Cytoplasmic (NES)

Folded (25 °C) → Gal Shut-off / Unfold (37 °C) → Unfolded → Sequestration (Nuc vs Cyto?) → MG132 ⊣ Degraded

c NLS-GFP-LuciTs, Time (min) at 37 °C
min   0   5   15   30   60   120
-MG132
+MG132

d NES-GFP-LuciTs, Time (min) at 37 °C
min   0   5   15   30   60   120
-MG132
+MG132

e *i.* NLS-LuciTs, 37 °C — INQ
ii. A–B: INQ, DNA, Nups

NLS-LuciTs: INQ
NES-LuciTs: JUNQ IPOD
Nups: Nuclear Pores
DNA: Nucleus

f *i.* NLS-VHL, 30 °C — INQ, A–B
ii. A–B: Nups, INQ, DNA

NLS-VHL: INQ
NES-VHL: JUNQ IPOD
Nups: Nuclear Pores
DNA: Nucleus

g *i.* NES-LuciTs, 37 °C — JUNQ, IPOD
ii. A–B: Nups, DNA, JUNQ
iii. C–D: IPOD, DNA, Nups

h *i.* NES-VHL, 30 °C — JUNQ, A–B
ii. A–B: JUNQ, Nups, DNA

a NLS-LuciTs + NES-LuciTs; Time (min) at 37 °C

0' 5' 15' 30' 60' 90'

Puncta Formation *Homing* *Co-migration*

b Puncta Formation

INQ — Distance — JUNQ

Homing

Co-migration

INQ–JUNQ Distance (nm)

Time at 37 °C (min)

c NLS-LuciTs (INQ)
NES-LuciTs (JUNQ)
Nups
DNA

INQ Nups JUNQ

A B

d
i. WT nup120Δ nup120Δ
NES-LuciTs (JUNQ)
Nups
DNA
Nups

ii. WT nup120Δ
NLS-LuciTs (INQ)
Nups
DNA

iii. WT nup120Δ
NLS-LuciTs INQ
NES-LuciTs JUNQ
Nups
DNA

e JUNQ Nups INQ

a

Frozen, unfixed intact cells → Fluorescence microscopy → X-ray Tomography → Measure LAC → Correlate X-ray/fluorescence → 3D reconstruction

b *i.*

Cells
Hsp104-GFP + Htt97QP-ChFP

X-ray Tomogram

Fluorescence overlay

ii.

Organelles	LAC
Cytosol	0.43
Nucleus	0.42
Mitochondria	0.50
Vacuoles	0.38
Lipid droplet	0.67
Fluorescent PQC Inclusion	0.47

c

3D reconstruction

JUNQ
450nm

IPOD

d

NES-LuciTs at 37 °C (JUNQ)

i

ii
JUNQ

e

NES-LuciTs

Mitochondria
Vacuole
Nucleus
Lipid droplets

f

Mitochondria   Vacuole   Nucleus   NES-LuciTs

180°

g

NLS-LuciTs at 37 °C (INQ)

400nm   INQ

Nucleolus   Nucleus
NLS-Luci   Vacuole

h

Mitochondria
Q-bodies
Nucleolus
IPOD
JUNQ
INQ
Nucleus
NVJ
Vacuole

JUNQ ?
Nucleus
INQ ?
Nvj1
Vac8
Vacuole

a Nvj1-GFP + NES-LuciTs; Time (min) at 37°C

8:15 | 8:15 | 8:30 | 8:45 | 9:00 | 9:15 | 9:30

9:45 | 10:00 | 10:15 | 10:30 | 10:45 | 11:00 | 11:00

b *i.* NES-LuciTs / Nvj1 / DNA   *ii.* JUNQ / Nu / Nvj1   *iii.* JUNQ / Nvj1 / Nu / IPOD

d Nu / Nvj1 / Vac

c *i.* NLS-LuciTs / Nvj1 / DNA   *ii.* Nu / INQ / Nvj1   *iii.* Nvj1 / INQ

e Nu / Nvj1 / INQ / Vac

f Time (min) NLS-LuciTs + NES-LuciTs; Time (min) at 37 °C

WT, PMN   0 | 5 | 15 | 30 | 60 | 90

nvj1Δ?   WT

    nvj1Δ

g NLS-LuciTs / NES-LuciTs / Merge

WT

nvj1Δ

vac8Δ

INQ / JUNQ / IPOD / Nuclear Pores

h Nuclear NLS-LuciTs — % Cells with Inclusions — WT, *nvj1Δ* **, *vac8Δ* *

Cytoplasmic NES-LuciTs — % Cells with Inclusions — WT, *nvj1Δ* **, *vac8Δ* ***

i YPD — WT, *nvj1Δ*, *vac8Δ*

0.5 mg/mL AZC — WT, *nvj1Δ*, *vac8Δ*

a

WT   + Bortezomib

NLS-LuciTs

GAPDH

Time (min)   0   30   60   90   120   0   30   60   90   120

NES-LuciTs

GAPDH

Time (hr)   0   1   2   4   0   1   2   4

Time after shift to 37°C

b

	Galactose 30 °C	Galactose 37 °C	Glucose
GFP			
LuciTs			
NLS-LuciTs			
NES-LuciTs			
Htt97QΔP			

c

VHL 37 °C +MG132

NLS-VHL
Nups
DNA

INQ   DNA

Nups

d

VHL 37 °C +MG132

NES-VHL
Nups
DNA

Nups

JUNQ   DNA

a Ubc9Ts

b 37 °C

-MG132   +MG132

GFP-VHL   Nups   DNA

no MG132   +MG132

Percent of cells with puncta

Nuclear   Cytoplasmic

c *nup116-5*

Ubc9Ts Half Life (37 °C)

Time (min)

WT   *nup116-5*

d *sts1-2*

Ubc9Ts Half Life (37 °C)

Time (min)

WT   *sts1-2*

a

EMCCD-LW   EMCCD-SW   Laser input   Spectrometer input   AL   MG   LM-CCD

RL   DM   MFW   Spinning disk   TL   MDT

External surface of capillary

CS

COL   CCL

CIM

b

a Alignment Using Fluorescence and Gold Fiducials

CFT reconstruction and SXT co-alignment fiducials
(200 nm red polystyrene)

SXT reconstruction alignment fiducials
(100 nm gold)

Specimen

b Alignment Using Joint Fiducials

SXT reconstruction alignment fiducials
(100 nm gold)

Joint Fiducials
(500 nm polystyrene)

Internal surface of capillary

Specimen

c

NLS-LuciTs
Nucleolus
DNA

d

NES-LuciTs
Mitochondria
DNA

e

WT	Mitochondrial Fission mutants		Mitochondrial Fusion mutants	
	dnm1Δ	*fis1Δ*	*fzo1Δ*	*ugo1Δ*

Mito-GFP   **NES-LuciTs**   **DNA**

a   **Nvj1-GFP** + **Ubc9Ts-ChFP**; Time (min) at 37°C

b   **NLS-LuciTs** + **NES-LuciTs;** Time (min) in CHX

c Nuclear NLS-LuciTs

d Cytoplasmic NES-LuciTs

e LuciTs after 60 mins at 37C

a

b

a

NES-2xKeima-LuciTs at 37°C

Time at 37 °C (min): 0 15 30 40 50 60 80 90

Ex 404

Ex 558

Merge

b

NES-2xKeima-LuciTs 85 min chase at 37°C

Ex 404   Ex 558   Merge

c

d

NLS-LuciTs   NES-LuciTs

e

WT Chase Time at 37 °C (min)   NLS-LuciTs   Vacuole

Nr.	NAME	SOURCE AND COMMENTS
SUPPLEMENTARY TABLE S2. Plasmids used in this study		
ES p331	NLS-GFP-LuciTs	This study
ES p357	NES-DsRed-LuciTs	This study
ES p325	NLS-GFP-VHL	This study and (Samant et al. 2018)
ES p355	NES-DsRed-VHL	This study and (Samant et al. 2018)
ES p20	Ubc9Ts-GFP	(Escusa-Toret et al. 2013)
ES p562	Ubc9Ts-ChFP	(Escusa-Toret et al. 2013)
ES p144	GFP-VHL	(Escusa-Toret et al. 2013)
ES p536	pRS NLS-GFP2-NES	(Stade et al., 1997)
ES p57	mHTT 97QP-ChFP	(Escusa-Toret et al. 2013)
ES p520	vt100 GFP	(Westermann et al. 2000)
FMP 1X	NLS-Keima-LuciTs	This study
FMP 1Y	NLS-2xKeima-LuciTs	This study
FMP 2X	NES-Keima-LuciTs	This study
FMP 2Y	NES-2xKeima-LuciTs	This study

Nr.	NAME	GENOTYPE	SOURCE AND COMMENTS	
SUPPLEMENTARY TABLE S1. Genotypes and sources of yeast strains used in this study				
ESY 1	BY4741			
ESY 104	W303			
ESY 105	CRM1		WT strain for crm1-T539C	from Michael Rosbash
ESY 106	crm1-T539C		has nuclear export deficiency with leptomycin B treatment	from Michael Rosbash
ESY 95	Nup116-5	nup116-5::HIS3	SWY027	from Susan Wente
ESY 96	STS1		WT strain for sts1-2	from Kiran Madura
ESY 97	sts1-2		temperature sensitive mutation that blocks proteasome import into the nucleus	from Kiran Madura
ESY 113	*dnm1*	BY4741 Mata	Deletion collection	
ESY 119	*fis1*	BY4741 Mata	Deletion collection	
ESY 122	*fzo1*	BY4741 Mata	Deletion collection	
ESY 123	*ugo1*	BY4741 Mata	Deletion collection	
ESY 121	Mps3-GFP	BY4741 Mata	GFP collection	
ESY 127	*nup120*	BY4741 Mata	Deletion collection	
ESY 94	*nupΔFG*		SWY3042	from Susan Wente
ESY 14	Nvj1-GFP	BY4741 Mata	GFP collection	
ESY 4	*nvj1*	BY4741 Mata	Deletion collection	
ESY 3	*vac8*	BY4741 Mata	Deletion collection	
ESY 150	*vps4*	BY4741 Mata	Deletion collection	
ESY177	Chm7OPEN-GFP	W303, chm7OPEN-(1-369)-GFP::HIS3	DTCPL413	From Patrick Lusk
ESY183	*Chm7*	W303, chm7Δ::hphMX6		From Patrick Lusk
ESY182	Chm7-GFP	W303, CHM7-GFP::HIS3		From Patrick Lusk
FMP1X	Nvj1-sfGFP	BY4741 Mata	NVJ1 GFP C-terminally tagged	This study