

Metabolomic comparison using *Streptomyces* spp. as a factory of secondary metabolites

Rene Flores Clavo ^{1,2*}, Alana Kelyene Pereira ³, Nataly Ruiz Quiñones ², Jonas Henrique Costa ³, Taícia Pacheco Fill ³ and Fabiana Fantinatti Garboggini ^{4,*}

- ¹ Graduate Program in Genetics and Molecular Biology, Institute of Biology, State University of Campinas (UNICAMP), street Monteiro Lobato, 255 – Campinas, SP, Brazil; renefloresclavo@gmail.com (R.F.C)
- ² Center for Research and Innovation in Multidisciplinary Active Sciences -CIICAM, street Pasaje Real 174, J.L. O, Lambayeque, Peru; nruiq@gmail.com (N.R.Q)
- ³ Institute of Chemistry (IQ) State University of Campinas (UNICAMP), University City Zeferino Vaz - Barão Geraldo, Campinas, SP, Brazil; alanakelyene@gmail.com (A.K.P); j161206@dac.unicamp.br (J.H.C); taicia@gmail.com (T.P. F)
- ⁴ Pluridisciplinary Center for Chemical, Biological and Agricultural Research (CPQBA), State University of Campinas (UNICAMP), Paulínia, SP, Brazil; fabianaf@cpqba.unicamp.br (F.F.G)
- * Correspondence: renefloresclavo@gmail.com; fabianaf@cpqba.unicamp.br

Abstract: Understanding extremophiles and their usefulness in biotechnology involves studying their habitat, physiology and biochemical adaptations, as well as their ability to produce biocatalysts, in environments that are still poorly explored. In northwestern Peru, which has saline lagoons of marine origin Pacific Ocean, the other site is from the coast of Brazil of the Atlantic Ocean. Both environments are considered extreme. The objective of the present work was to compare two different strains isolated from these extreme environments at the metabolic level using molecular network methodology through the Global Natural Products Molecular Social Network (GNPS). In our study, the MS/MS spectra from the network were compared with GNPS spectral libraries, where the metabolites were annotated. Differences were observed in the molecular network presented in the two strains of *Streptomyces* spp. coming from these two different environments. Within the annotated compounds from marine bacteria, the metabolites characterized for *Streptomyces* sp. B-81 from Peruvian marshes were lobophorins A (1) and H (2), as well as divergolides A (3), B (4) and C (5). *Streptomyces* sp. 796.1 produced different compounds, such as glucopiericidin A (6) and dehydro-piericidin A1a (7). The search for new metabolites in underexplored environments may therefore reveal new metabolites with potential application in different areas of biotechnology.

Keywords: Genome mining; marine environments; molecular networking; bacterial extremophiles; secondary metabolites

Citation: Flores Clavo, R.; Kelyene Pereira, A.; Ruiz Quiñones, N.; Henrique Costa, J.; Pacheco Fill, T.; Fantinatti Garboggini, F. Metabolomic comparison using *Streptomyces* spp as a factory of secondary metabolites. *xxxxxxx*, 20, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Last-name

Received: date

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Approximately 22,500 biologically active substances are obtained from microorganisms, 45% of which are represented by Actinomycetes and 70% of which are *Streptomyces* metabolites; however, information on substances isolated from microorganisms inhabiting saline environments is scarce [1,2,3]. The search for these microorganisms has been mainly associated with the production of antibiotics and antitumor substances [4]. Halophilic and halotolerant strains show heterogeneous physiological characteristics for different genera because these bacteria can synthesize secondary metabolites to cope with the high salinity and extreme temperature conditions of their environments [5,6]. These extreme conditions favor the development of metabolic competitiveness for the production of enzymes, providing adaptation to the high salinity of the environment [4]. The production of bioactive molecules from actinobacteria other than *Streptomyces* allows extending the search for new molecules and interactions. During the last five years, constant

isolation efforts of microorganisms from various environmental niches have resulted in the recovery of thousands of members of the actinobacteria affiliated with different taxonomic classes [5]. The members of the genus *Streptomyces* stand out as producers of the largest amount of secondary metabolites already described [6]. Various studies have revealed an approach combining LC–MS/MS and molecular networking as a rapid analytical method for the identification of compounds [7]; this microbial group has a unique ability to produce new products, mainly antibiotics [8,9].

[8] Investigated the metabolic profiling of a cultured endophyte strain (*Streptomyces* sp. HKI0576) by HPLC–MS and revealed a complex metabolome, *Streptomyces* sp. 12A35 was isolated from deep sea sediment collected from the South China Sea and showed promising antibacterial activities attributed to the spirorotronate antibiotics lobophorins B, F, H, and I from a marine origin [9]; Piericidins and Glucopiericidins A and B isolated from the culture broth of *Streptomyces pactum* S48727 [10], a new natural product of the lobophorin family designated lobophorin K, from cultures of the marine actinobacteria *Streptomyces* sp. M-207 [11]; A cryptic BGC for type I polyketides was activated by metabolic engineering methods, enabling the discovery of a known compound, lobophorin CR4, of deep-sea-derived *Streptomyces olivaceus* SCSIO T05 [12].

A study reported in Salar de Huasco in the Atacama Desert, considered a polyextreme environment in Chile, [13] described that species of the genus *Streptomyces* were dominant, and a preliminary study [14] present in the salt ponds of Bayovar and Morrope in Peru showed physicochemical characteristics common to extremophilic bacteria and ample potential to produce bioactive compounds. These extreme conditions encourage the development of metabolic competitiveness for the production, for example, of enzymes [15]. The application of various targeted studies on the biosynthesis of natural products allows a better understanding of the full potential of the producing microorganisms, thus increasing the return on investment in the search for new compounds [16].

In this study, we used a molecular networking approach for the rapid detection of molecules from two strains of *Streptomyces* spp. obtained from extreme environments: Saloons northwestern of Perú and Cabo Frio in Brazil (Figure 1). The detection method, based on UHPLC–MS/MS combined with data analysis using Global Natural Product Social Molecular Networking (GNPS), led to the annotation of very different compounds produced by each of the *Streptomyces* spp.




Figure 1. The geographical distribution of *Streptomyces* spp. location with a description of growth in solid medium. (1) Sediment of a Bayovar saline lagoon located in northwestern Peru and (2) starfish in Cabo Frio (RJ) in Brazil.

47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78

79
80
81
82
83

2. Results and Discussion

2.1. Phylogenetic analysis of 16S rRNA from two *Streptomyces* spp.

The 16S rRNA sequences of the two marine *Streptomyces* species analyzed in this study, *Streptomyces* sp. B-81 collected in 2015 in Peru and *Streptomyces* sp. 796.1 collected in Brazil 2012, were determined and compared by MEGA11.0 software. The phylogenetic tree obtained from the two *Streptomyces* strains was mainly divided into two characteristic branches. The samples mentioned before were constructed against 29 *Streptomyces* strains according to EZBioCloud and are presented in Figure 2. The strain obtained from the Peruvian saline lagoon was determined to be *Streptomyces* sp. B-81 (MW562807), whose phylogenetically related sequence identity (1345 bp 92.7%, gene sequence identity) groups to the type strain *S. olivaceus* NRRL B-3009^T with 99.93%, *S. pactum* NBRC 13433^T and bootstrap value 97% (Fig. 2 and Table S1 and Table S2). However, for the strain isolated from a marine invertebrate in Brazil, *Streptomyces* sp. 796.1 (MG654686) sequence identity (1370 bp 93.4%, gene sequence identity similarity) is phylogenetically related to the type strains *S. buecheriae* AC541^T with 99.70%, *S. youssoufiensis* X4^T and *S. zagrosensis* HM 1154^T with 99.48%, and *Streptomyces philanthi triangulum*^T (DQ3752) with 99.45% gene sequence identity similarity and a bootstrap value of 85% (Figure 2 and Tables S1 and S2). Subsequently, each part was further subdivided, and most strains were well separated by clustering under each species, generating indications that they may be new species. Existing aquatic ecosystems are relatively unexplored but have been reported to harbor a great diversity of microorganisms, which may be reflected in their broad biosynthetic potential [17].




Figure 2. Phylogenetic tree of *Streptomyces* spp. strains B-81 and 796.1. The evolutionary history was inferred using the neighbor-joining method [18]. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches. The tree is drawn to scale, with branch lengths similar to evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Kimura 2-parameter method [19] and are in units of the number of base substitutions per site. The analysis involved 32 nucleotide sequences. All positions containing gaps and missing data were eliminated. There were a total of 1498 positions in the final dataset. Evolutionary analyses were conducted in MEGA11 [20]. Bar, 0.02 substitutions per nucleotide, *Pseudomonas humanensis* LV^T (JX545210) was used as the outgroup.

2.2. MS/MS-Based Molecular Networks of *Streptomyces* spp.

The metabolites of the crude extracts obtained from each of the *Streptomyces* spp. samples were prepared and analyzed by UHPLC–MS/MS, and the resulting data were analyzed through multivariate data analyses to compare both strains using LC–MS (Fig. 3). The data showed a clear separation between both *Streptomyces* spp.

strains. In particular, using principal component analysis (PCA), *Streptomyces* sp. B-81 was separated from *Streptomyces* sp. 796.1 by PC1 and PC3, explaining ~ 40% of the variance (Fig. 3A). Another multivariate analysis, partial least squares-discriminant analysis (PLS-DA), showed clear differentiation between both *Streptomyces* strains using components 1 and 2 (Fig. 3B), and chemotaxonomic classification based on *Streptomyces* using liquid chromatography-electrospray ionization-tandem mass spectrometry (LC-ESI-MS/MS) combined with multivariate statistical analysis demonstrated that metabolite-based chemotaxonomic classification is an effective tool for distinguishing *Streptomyces* spp. and for determining their species-specific metabolites [21].





Figure 3. Multivariate data analysis of LC-MS/MS: (A). Principal component analysis (PCA) and (B). partial least square-discriminant analysis (PLS-DA) comparing both *Streptomyces* strains. *Streptomyces* sp. B81 is in pink, *Streptomyces* sp. 796.1, in green, the culture medium R2A, in yellow and the solvent blank is in white.

Mass spectrometric data obtained from these analyses were then used to generate molecular networking coupled to in silico tools (Networking annotation propagation (NAP) (<https://ccms-ucsd.github.io/GNPSDocumentation/nap/>) and MolNetEnhancer (<https://ccms-ucsd.github.io/GNPSDocumentation/molnetenhancer/>)). Data obtained from these analyses were then used to generate more detailed molecular networking (Fig. 4A). Our analyses have presented several chemical classes, such as lipids and lipid-like molecules, organic nitrogen organoheterocyclic compounds, organic acids, organosulfur compounds, polyketides, and benzenoid compounds (Fig. 4B).

Microorganisms in the environment can produce a wide range of secondary metabolites that are natural products with diverse chemical structures and that will perform a variety of functions acting as antibiotics, antitumour agents, cholesterol-lowering agents, etc. [22]. According to the LC-MS/MS analysis, the number of ions can be classified according to the strain source. In all 1005 detected ions, ~ 30% are only from *Streptomyces* sp. B-81 and ~ 64,5% from *Streptomyces* sp. 796.1 (Fig. 4C). Thus, compounds of the same chemical class are grouped in the same cluster,

demonstrating a broad biomolecule profile for *Streptomyces* sp. B-81 (pink), including the production of lobophorins A (1) and H (2), annotated based on their accurate masses, with specific masses of 1157.6379 (1) and 753.4320 (2), respectively; metabolites were annotated as hits in the GNPS database or manually by accurate mass analyses, which showed mass errors below 0.10 ppm (Table 1). The cluster of the GNPS database indicated the production of lobophorin A by *Streptomyces* sp. B-81 (MW562807) strain, and fragmentation profile with typical fragments at 97.06, 183.11, 517.29 m/z (Fig. S1), the metabolite lobophorin H presented the precursor at m/z 753.4320 (Fig S2), and compounds that belong to the divergolide family, including divergolides A (3), B (4) and C (5), based on their precursor ions with m/z values of 554.2750 (3), 536.2650 (4), and 550.2440 (5), respectively. The fragmentation profile with typical fragments at divergolide C showed the main fragments m/z 182.08, 332.22, 398.32 (S3 – S5 Figs). In a previous study of Bayovar's saloons *Streptomyces* sp. B-81 (MW562807) demonstrated the most promising activity, and six biomolecules (Cholic Acid, Lobophorin A, B, E, K and six compounds of type Furano) [14]. The MS/MS spectra of the network metabolites were compared with the GNPS spectral libraries and identified for *Streptomyces* sp. 796.1, two substances were annotated as glucopiericidin A (6) and dehydropiericidin A1a (7), with precursor ions at m/z 578.333 (7) and 414.26 (8), which were identified based on their fragmentation profiles with typical fragments at m/z 161.13, 182.09, 398.26 and 135.11, 182.09, 330.20, respectively (Figs S6-S7) (Fig. 4D and Table 1). In a study carried out on the marine sediment of Saint Peter and Saint Paul's archipelago in Brazil, dereplication analysis of the extracts showed the presence of several compounds identified as piericidins A and C as well as glucopiericidin A, produced by *Streptomyces* sp. BRA-199 [23]



195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212

Figure 4. (A) MolNetEnhancer of *Streptomyces* strains LC–MS/MS data. (B) Pie chart based on the chemical classes found on the LC–MS/MS data. (C) Euler diagram based on the ion distribution of both strains. (D) Classical molecular networking of both *Streptomyces* strains.

Table 1. Mass error (0,10 ppm) of observed and calculated m/z values of known compounds found in clusters based on MS/MS data obtained from UHPL–HRMS.

N	GNPS NEW-ID	Compound	Molecular Formula [M+H] ⁺	m/z theoretical	m/z observed	error (ppm)	Strain
1	5653	Lobophorin A	C61H93N2O19	1157.6367	1157.6379	1.032275	B-81
2	5178	Lobophorin H	C42H61N2O10	753.4320	753.43200	0.096890	B-81
3	4097	Divergolide A	C31H40NO8	554.2748	554.27500	0.281449	B-81
4	3981	Divergolide B	C31H38NO7	536.2642	536.26500	1.344486	B-81
5	4068	Divergolide C	C31H36NO8	550.2435	550.24400	0.828724	B-81
6	4214	Glucopiericidin A	C31H48NO9	578.3323	578.3330	1.108359	796.1
7	3145	Dehydropiericidin A1a	C25H36NO4	414.263885	414.2650	2.691521	796.1

Error (0,10 ppm) = observed m/z – calculated m/z .

Ding and colleagues in 2018 took an approach combining LC–MS/MS and molecular networks as a rapid analytical method to detect six compounds. Aplysiatoxins from marine cyanobacteria were described with potential new analogs. Another study of MS/MS data obtained from fractions on the public GNPS platform succeeded in dereplicating new glycolipopeptides called characellides, cyanocobalamin and poecillastrins of a deep-sea tetractinellid sponge, *Characella pachastrelloides* [24,25]. Initial chemical investigation of the extract obtained from *Streptomyces* spp. indicated the production of compounds 1-7. Each node within the clusters represents an ionized metabolite, and dereplication was performed by comparison with the m/z values of known lobophorins, divergolides and piericidin-related analogs (Table 2). Lobophorins A and B were discovered by Jiang et al. (1999) [26] to Belize onboard the Columbus research ship. Lobophorin C and D were isolated from a new actinomycete strain named # CNC-837, identified as *Streptomyces carnosus* strain AZS17, isolated from the coastal waters of the East China Sea [27]. Two polyketides, spiro-tetronate classified as lobophorins H and I, were isolated from *Streptomyces* sp. 1053U.I.1a.3b [28], as well as the other substances of the Lobophorin family (Table 2). The family of compounds named Divergolides isolated from the stem of the mangrove tree *Streptomyces* sp. HKI0576 grown by HPLC–MS revealed a complex metabolome [8]. Four ansamycins, named divergolides T–W, isolated from the mangrove-derived actinomycete *Streptomyces* sp. KFD18 [29], a substance named 3'-deoxytalopiericidin A1 or glucopiericidin A, was isolated from *Streptomyces* sp. soil sample collected in Gotemba City, Shizuoka, Japan [30], 13-hydroxyglucopiericidin A and the analog Glucopiericidin A were isolated from the fermentation broth of *Streptomyces* sp. OM-5689 [31] (Table 2).

Table 2. A list of known compound families related to *Streptomyces* spp. with compound bioactivity with m/z values.

Compound	Molecular Formula [M+H] ⁺	Mass Molecular	Mass calculated	Origins	Reference
Lobophorin A	C61H93N2O19	1157.636705	1157.6379	<i>Streptomyces</i> sp. CNC-837	B-81, [26]
Lobophorin B	C61H91N2O21	1187.610884	1187.6114	<i>Streptomyces</i> sp. CNC-837	[26], [14]
Lobophorin C	C61H90N2O21	1186.603059	1209.5958	<i>Streptomyces</i> sp. AZS 17	[27]
Lobophorin D	C61H93N2O19	1157.636705	1157.6379	<i>Streptomyces</i> sp. AZS 17	[27]
Lobophorin E	C61H91N2O20	1171.615970	1171.6165	<i>Streptomyces</i> sp. SCSIO 01127	[32]

Lobophorin F	C54H78N2O17	1026.529500	1049.5198	<i>Streptomyces</i> sp. SCSIO 01127	[32]
Lobophorin G	C63H94N2O20	1198.639445	1199.6472	<i>Streptomyces</i> sp. MS100061	[33]
Lobophorin H	C42H61N2O10	753.432073	753.43200	<i>Streptomyces</i> sp. 1053U.I.1a.3b	B-81, [28]
Lobophorin I	C61H89N2O21	1185.596209	1185.5958	<i>Streptomyces</i> sp. 1053U.I.1a.3b	[28]
Lobophorin K	C61H93N2O20	1173.631620	1173.6322	<i>Streptomyces</i> sp. M-207	[34]
Lobophorin CR1	C61H92NO20	1144.617647	1180.6032	<i>Streptomyces</i> sp. 7790_N4	[35]
Lobophorin CR2	C61H90N2O22	1202.597974	1247.5702	<i>Streptomyces</i> sp. 7790_N4	[35]
Lobophorin CR3	C61H90N2O23	1218.592889	1241.5832	<i>Streptomyces</i> sp. 7790_N4	[35]
Lobophorin CR4	C52H77N2O15	969.531846	-	<i>Streptomyces olivaceus</i> SCSIO T05	[36]
Lobophorin L	C54H80N2O16	1013.5618	1013.5581	<i>Streptomyces</i> sp. 4506	[37]
Lobophorin M	C48H70N2O13	882.487242	883.4951	<i>Streptomyces</i> sp. 4506	[37]
Lobophorin N	C61H92N2O18	1141.64336	1141.6418	<i>Streptomyces</i> sp. HDN1844000	[38]
Divergolide A	C31H39NO8	554.274844	554.27500	<i>Streptomyces</i> sp. HKI0576	B-81, [8]
Divergolide B	C31H37NO7	536.264279	536.26500	<i>Streptomyces</i> sp. HKI0576	B-81, [8]
Divergolide C	C31H33NO8	550.243544	550.24400	<i>Streptomyces</i> sp. HKI0576	B-81, [8]
Divergolide D	C31H35NO8	535.232645	535.2326	<i>Streptomyces</i> sp. HKI0576	[8]
Divergolide T	C31H37NO7	536.264300	536.2641	<i>Streptomyces</i> sp. KFD18	[29]
Divergolide U	C31H37NO8	550.244600	550.2438	<i>Streptomyces</i> sp. KFD18	[29]
Divergolide V	C31H37NO7	536.264000	536.2640	<i>Streptomyces</i> sp. KFD18	[29]
Divergolide W	C31H37NO7	534.249000	534.2490	<i>Streptomyces</i> sp. KFD18	[29]
Piericidin A-A1	C25H ₃₇₋₃₉ NO4	401.268636	401.2686	<i>Streptomyces</i> sp. 16-22	[39]
Piericidin B-B1	C26H39NO4	415.284286	-	<i>Streptomyces mobaraensis</i>	[40]
Piericidins B1N-oxide	C26H39NO5	431.279201	446.2897	<i>Streptomyces</i> sp. MJ288-OF3	[41]
Piericidin C7	C28H41NO5	457.294851	472.3062	<i>Streptomyces</i> sp. YM14-060	[42]
Piericidin C8	C29H43NO5	471.310501	486.3223	<i>Streptomyces</i> sp. YM14-060	[42]
Glucopiericidin A	C31H48NO9	578.332359	578.3330	<i>Streptomyces</i> spp.	796.1, [30]
Dehydropiericidin A	C25H36NO4	414.263885	414.2650	<i>Streptomyces</i> spp.	796.1, [31]
13-hydroxyglucopiericidin A	C31H47NO10	579.316374	593.3163	<i>Streptomyces</i> sp. OM-5689	[43]
Glucopiericidin C	C30H45NO8	533.310895	533.3108	<i>Streptomyces</i> sp. B811	[44]
JBIR-02	C27H39NO3	411.289372	426.2979	<i>Streptomyces</i> sp. ML55	[45]

3. Materials and Methods

245

3.1. Biological Material

246

The microorganisms used in this study were preserved by deep freezing at -80°C in 30% glycerol in the research collection of the Microbial Resources Division, Pluridisciplinary Center for Chemical, Biological and Agricultural Research (CPQBA) State University of Campinas (UNICAMP), Campinas, SP, Brazil. (<https://www.cpqba.unicamp.br/>) and collection of CIICAM Research Center (www.ciicam.com). Two bacteria belonging to the genus *Streptomyces* were identified as *Streptomyces* sp. 796.1 (MG654686), isolated from a starfish in Cabo Frio (RJ) in Brazil in 2012, *Streptomyces* sp. B-81 (MW562807) was isolated from the sediment of a Bayovar saline lagoon located in northwestern Peru in 2015.

247

248

249

250

251

252

253

254

255

The isolates were reactivated using R2A medium (KASVI) with artificial sea-
water (ASW) ($\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ 10.83 g. L^{-1} , $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ 1.51 g. L^{-1} , $\text{SrCl}_2 \cdot 6\text{H}_2\text{O}$ 0.02 g. L^{-1} ,
 NaCl 23.93 g. L^{-1} , Na_2SO_4 4.01 g. L^{-1} , KCl 0.68 g. L^{-1} , NaHCO_3 0.20 g. L^{-1} , KBr 0.098 g.
 L^{-1} , H_3BO_3 0.03 g. L^{-1}) with 5% NaCl and incubated at 28°C for 15 days. After incuba-
tion, the colonies verified for purity were collected from the plate and used for the
extraction of genomic DNA.

3.2. DNA isolation and PCR amplification of bacteria

The extraction of genomic DNA was performed using the protocol established
by Van Soolingen et al. (1991) adapted to the conditions of the laboratory
(<https://dx.doi.org/10.17504/protocols.io.bs4ncqw>). First, quantification of genomic
DNA was performed by comparing the λ DNA at different concentrations in a 1%
TBE 1X agarose gel submitted to electrophoresis for 20 min at 5 V per cm^2 . Then, the
DNA was kept at -20°C until the amplification of the 16S rRNA gene was performed
by polymerase chain reaction (PCR). The specifications of this protocol have been
previously described (dx.doi.org/10.17504/protocols.io.brrmm546).

3.3. Sequencing and Phylogenetic Analysis of 16

The samples were purified using a minicolumns GFX PCR DNA & gel band pu-
rification kit (GE Healthcare Bio-Sciences AB Uppsala, Sweden) according to a pre-
viously described protocol (<https://dx.doi.org/10.17504/protocols.io.brzpm75n>). The
samples were sequenced using an ABI3500XL Series automatic sequencer (Applied
Biosystems Foster City, California, USA) in the Laboratory of the Division of Micro-
bial Resources, Chemical, Biological and Agricultural Pluridisciplinary Research
Center (CPQBA), University of Campinas (UNICAMP), Paulínia, São Paulo, Brazil.
Sequencing reactions were performed using the Big Dye Terminator Cycle Sequenc-
ing Ready Reaction Kit (Applied Biosystems) according to the manufacturer's modi-
fied by the author's protocol (<https://dx.doi.org/10.17504/protocols.io.brzpm75n>).

Partial sequences of the 16S ribosomal RNA gene obtained from each isolate
were assembled into a contig using BioEdit 7.0. The sequences of organisms were
added to the EZBioCloud 16S Database (<https://www.ezbiocloud.net/>) using the
"Identify" service, and species assignment was based on the closest hit 16S rRNA
gene sequences retrieved from the database and related to the unknown organism
gene were selected for alignment in the Clustal W program. Phylogenetic analyses
were performed using the MEGA version 11.0 program, and the evolutionary dis-
tance matrix was calculated using the Kimura-2 model parameters. Then, the phylo-
genetic tree was constructed from the evolutionary distances calculated by the neigh-
bor-joining method with bootstrap values based on 1000 resamples.

3.4. Production of Crude Extracts of *Streptomyces* spp

The production of crude ethyl acetate extracts of bacterial isolates was per-
formed in culture media R2A broth (Himedia ref. 1687) for *Streptomyces* spp. B-81 and
796.1 strains as described in a previously published protocol
(dx.doi.org/10.17504/protocols.io.br2cm8aw). *Streptomyces* sp. metabolites were ex-
tracted with 1000 μL of methanol in an ultrasonic bath for 40 minutes, dried under
inert conditions and analyzed by ultrahigh pressure liquid chromatography–mass
spectrometry (UHPLC–MS) in a Thermo Scientific QExactive® Hybrid Quadrupole-
Orbitrap Mass Spectrometer.

3.5. LC–MS/MS data acquisition

Analyses were performed in the positive mode with a range of m/z 133-2000, a
capillary voltage of 3.4 kV, inlet capillary temperature of 280°C , and S-lens 100 V. A
Thermo Scientific column Accucore C18 2.6 μm (2.1 mm x 100 mm) was used as the

stationary phase, and 5 μL of the sample was injected. The mobile phase was composed of 0.1% formic acid and acetonitrile, and the eluent profile followed a molar ratio of 95/5 up to 2/98 within 10 min, held for 5 min, up to 95/5 within 1.2 min, and held for 8.8 min. The total run time was 25 min for each run, and the flow rate was 0.2 mL \times min⁻¹. The data were initially processed with Xcalibur software (version 3.0.63) developed by Thermo Fisher Scientific.

3.6. LC-MS/MS data analysis and metabolite annotation

Raw data, blanks, and media controls were converted to.mzML using MSConvert software (<http://proteowizard.sourceforge.net>). Next, a molecular network for *Streptomyces* sp. strain metabolites was created using the online workflow (<https://ccms-ucsd.github.io/GNPSDocumentation/networking/>) on the Global Natural Products Social Molecular Networking (GNPS) website platform (<http://gnps.ucsd.edu>) using the Classical Molecular Networking (CMN) tool. For network creation, the data were filtered by removing all MS/MS fragment ions within ± 17 Da of the precursor m/z. MS/MS spectra were filtered by choosing only the top 6 fragment ions in the ± 50 Da window throughout the spectrum. The precursor ion mass tolerance was set to 0.02 Da and an MS/MS fragment ion tolerance of 0.02 Da. A network was then created where edges were filtered with a cosine score above 0.65 and more than six (6) matched peaks.

Furthermore, edges between two nodes were kept in the network if and only if each of the nodes appeared in each other's respective top 10 most similar nodes. Finally, the maximum size of a molecular family was set to 100, and the lowest-scoring edges were removed from molecular families until the molecular family size was below this threshold. The spectra in the network were then searched against GNPS' spectral libraries. The library spectra were filtered in the same manner as the input data. All matches kept between network spectra and library spectra were required to have a score above 0.65 and at least six (6) matched peaks [46]. The resulting molecular networking is available at:

(<https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=b7a29b5a78864223bde01fb8af0c5eff>). To perform the multivariate data analysis, the LC-MS data were processed into MZmine online using GNPS Dashboard (<https://ccms-ucsd.github.io/GNPSDocumentation/lcms-dashboard/>). The job is available at:

<https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=88a21c1aabb04a1287348300b711ebfd>. The quantification table was submitted to the MetaboAnalyst 5.0 platform (<https://www.metaboanalyst.ca/>) for statistical analysis.

The nodes (MS/MS spectra) originating from R2A culture media and blanks (methanol) were filtered from the original network to enable visualization of metabolites. Finally, the final spectral network (.cys) was uploaded in Cytoscape 3.8 to obtain better visualization and editing.

4. Conclusions

In this study, we sought to classify two *Streptomyces* species based on their secondary metabolites. The dendrograms based on metabolites and 16S rRNA exhibited the same patterns of species discrimination and may be new species. The multivariate data analyses to compare both strains using LC-MS showed a clear separation between both *Streptomyces* spp. The MS/MS spectra of the network metabolites were compared with the GNPS spectral libraries, and differences in the molecular network presented in *Streptomyces* sp. B-81 isolated from the Bayovar saline lagoon located in northwestern Peru were lobophorins A (1) and H (2), as well as divergolides A (3), B

(4) and C (5); *Streptomyces* sp. 796.1 isolated in Cabo Frio (RJ) in Brazil dereplicated two compounds, glucopiericidin A (6) and dehydropiericidin A1a (7).

Supplementary Materials: The following supporting information can be downloaded at www.mdpi.com/xxx/s1. Phylogenetic data of *Streptomyces* sp. B-81 Accession number (MW562807), and *Streptomyces* sp. 796.1 Accession number (MG654686); MS/MS match between GNPS database lobophorin A (1), in the GNPS database or manually by accurate determination of which ion precursor lobophorin H (2) and ions precursor divergolides A (3), B (4) and C (5) from *Streptomyces* sp. B-81 extract and MS/MS match between GNPS database with Glucopiericidin A (6) and Dehydropiericidin A1a (7).

Author Contributions: R.F.C. performed the extraction of *Streptomyces* sp. B-81 sample, the isolation, purification, and structure elucidation of compounds, and constructed the molecular networks A.K.P. wrote and paper; N.R.Q. performed the isolation and production of *Streptomyces* sp. 796.1 extracts. A.K.P. and J.H.C performed the UHPLC–MS/MS analysis and Thermo Scientific QExactive® Hybrid Quadrupole-Orbitrap Mass Spectrometer and constructed the molecular networks; T.P.F and F.F.G edited and vetted the paper. All authors read and corrected the paper.

Funding: This study has been financed by the Concytec-World Bank Project "Improvement and Expansion of the Services of the National Science Technology and Technological Innovation System" 8682-PE, through its executing unit ProCiencia [contract number 190-2018] and "The postgraduate programs in Genetics and Molecular Biology and CNPQ.

Acknowledgments: This work was supported by the Biotechnology Department of the Center for Research and Innovation in Multidisciplinary Active Sciences -CIICAM and Pluridisciplinary Center for Chemical, Biological and Agricultural Research (CPQBA), State University of Campinas (UNICAMP), Paulínia, SP, Brazil.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

- Hayakawa, Y; Shirasaki, S; Shiba, S; Kawasaki, T; Matsuo, Y; Adachi, K., & Sishuri, Y. Piericidins C7 and C8, new cytotoxic antibiotics produced by a marine *Streptomyces* sp. *J. Antibiot.* **2007**, *60*, 196–200. doi:10.1038/ja.2007.22. [PubMed]
- Valliappan, K; Sun, W; Li, Z. Marine actinobacteria associated with marine organisms and their potentials in producing pharmaceutical natural products. *Appl. Microbiol. Biotechnol.* **2014**, *98*, 7365–7377. doi:10.1007/s00253-014-5954-6. [PubMed]
- Binayke, A; Ghorbel, S; Hmidet, N; Raut, A; Gunjal, A; Uzgare, A,... & Nawani, N. Analysis of diversity of actinomycetes from arid and saline soils at Rajasthan, India. *Environ. Sustain.* **2018**, 1–10. doi:10.1007/s42398-018-0003-5. [Crossref]
- Kiyoon Kim; Sandipan Samaddar; Poulami Chatterjee; Ramasamy Krishnamoorthy; Sunyoung Jeon, T.S. Structural and functional responses of microbial community with respect to salinity levels in a coastal reclamation land. *Appl. Soil. Ecol.* **2019**, *137*, 96–105. doi:10.1016/j.apsoil.2019.02.011. [Crossref]
- Berdi, J. Bioactive microbial metabolites. *Journal of antibiotics.* **2015**, *58*, 1, 1-26. doi:10.1038/ja.2005.1. [Crossref] [PubMed]
- Kamjam, M; Sivalingam, P; Deng, Z; Hong, K. Deep sea actinomycetes and their secondary metabolites. *Front. Microbiol.* **2017**, *8*, 1–9. doi:10.3389/fmicb.2017.00760. [Crossref] [PubMed]
- Tangerina, M.M.P, Furtado, L.C, Leite, V.M.B, Bauermeister, A; Velasco-Alzate, K; Jimenez, P.C.... & Pena Ferreira, M.J. Metabolomic study of marine *Streptomyces* sp.: Secondary metabolites and the production of potential anticancer compounds. *PLoS One.* **2021**, *15*, 1–19. doi:10.1371/journal.pone.0244385. [Crossref] [PubMed]
- Ding, L; Maier, A; Fiebig, H.H; Görls, H; Lin, W.H; Peschel, G., & Hertweck, C. Divergolides A-D from a mangrove endophyte reveal an unparalleled plasticity in ansa-macrolide biosynthesis. *Angewandte Chemie.* **2011**, *50*, 1630–1634. doi:10.1002/anie.201006165. [Crossref] [PubMed]
- Hua-Qi, Pan; Song-Ya, Zhang; Nan, Wang; Zhan-Lin Li; Hui-Ming; Hua J-CH and Wang, S.J. "New spirotetronate antibiotics, lobophorins H and I, from a South China Sea-derived *Streptomyces* sp. 12A35. *Marine Drugs.* **2013**. 3891–3901. doi:10.3390/md11103891. [Crossref] [PubMed]
- Masaru Matsumoto; Kin-Ichi Mogi; Katsuhiko Nagaoka; Seiji Ishizeki, R.K and T.N. New Piericidin Glucosides, Glucopiericidins A and B. *J. Antibiot.* **1987**, 149–156. doi:10.7164/antibiotics.40.149. [Crossref] [PubMed]
- Braña, A.F; Sarmiento-Vizcaíno, A; Osset, M; Pérez-Victoria, I; Martín, J; De Pedro, N.... & Blanco, G. Lobophorin K, a new natural product with cytotoxic activity produced by *Streptomyces* sp. M-207 associated with the deep-sea coral *Lophelia pertusa*. *Marine Drugs.* **2017**. doi:10.3390/md15050144. [Crossref] [PubMed]

12. Zhang, C; Ding, W; Qin, X; Ju, J. Genome Sequencing of *Streptomyces olivaceus* SCSIO T05 and Activated Production of Lobophorin CR4 via Metabolic Engineering and Genome Mining. *Mar. Drugs*. **2019**, *17*, 1–11. doi:10.3390/md17100593. [\[Crossref\]](#) [\[PubMed\]](#)
13. Carlos, Cortés-Albayay; Silver, J; Imhoff, J.F; Asenjo, J.A; Andrews, B; Nouioui, I., & Dorador, C. The Polyextreme Ecosystem, Salar de Huasco at the Chilean Altiplano of the Atacama Desert Houses Diverse *Streptomyces* spp. with Promising Pharmaceutical Potentials. *Diversity*. **2019**, *11*, 1–12. doi:10.3390/d11050069. [\[Crossref\]](#)
14. Flores Clavo, R; Ruiz Quiñones, N; Hernández-Tasco, A.J; Jose Salvador, M; Tasca Gois Ruiz, A.L; A. L, De Oliveira Braga; L.E....& Fantinatti Garboggini, F. Evaluation of antimicrobial and antiproliferative activities of Actinobacteria isolated from the saline lagoons of northwestern Peru. *PLoS One*. **2021**. 1–18. doi: <https://doi.org/10.1371/journal.pone.0240946>. [\[Crossref\]](#) [\[PubMed\]](#)
15. Lee, S.Y; Kim, H.U; Chae, T.U; Cho, J.S; Kim, J.W; Shin, J.H....& Jang, Y.S. A comprehensive metabolic map for production of biobased chemicals. *Nat. Catal*. **2019**, *2*, 18–33. doi:10.1038/s41929-018-0212-4. [\[Crossref\]](#)
16. Du Chao. Multiomics studies of the control of growth and antibiotic production of *Streptomyces*. (Doctoral dissertation Leiden University). **2020**, 1–176. [\[Crossref\]](#)
17. Cuadrat, R.R.C; Ionescu, D; Dávila, A.M.R; Grossart, H.P. Recovering genomics clusters of secondary metabolites from lakes using genome-resolved metagenomics. *Front. Microbiol*. **2018**, *9*, 1–13. doi:10.3389/fmicb.2018.00251. [\[Crossref\]](#) [\[PubMed\]](#)
18. Saitou, N; Nei, M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol*. **1987**, *4*, 406–425. doi:10.1093/oxfordjournals.molbev.a040454. [\[Crossref\]](#) [\[PubMed\]](#)
19. Kimura, M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol Evol*. **1980**, *16*, 111–120. doi:10.1007/BF01731581. [\[Crossref\]](#) [\[PubMed\]](#)
20. Tamura K; Stecher, G; Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Mol. Biol. Evol*. **2021**, *38*, 3022–3027. doi:10.1093/molbev/msab120. [\[Crossref\]](#) [\[PubMed\]](#)
21. Lee, M.Y; Kim, H.Y; Lee, S; Kim, J.G; Suh, J.W; Lee, C.H. Metabolomics-based chemotaxonomic classification of *Streptomyces* spp. and its correlation with antibacterial activity. *J. Microbiol. Biotechnol*. **2015**, *25*, 1265–1274. doi:10.4014/jmb.1503.03005. [\[Crossref\]](#) [\[PubMed\]](#)
22. Chen, R; Wong, H; Burns, B. New Approaches to Detect Biosynthetic Gene Clusters in the Environment. *Medicines*. **2019**, *6*, 32. doi:10.3390/medicines6010032. [\[Crossref\]](#) [\[PubMed\]](#)
23. Ferreira, E.G; Torres, M da C.M; da Silva A.B; Colares, L.L.F; Pires, K; Lotufo, T.M.C.... & Jimenez, P.C. Prospecting Anticancer Compounds in Actinomycetes Recovered from the Sediments of Saint Peter and Saint Paul’s Archipelago, Brazil. *Chem Biodivers*. **2016**, *13*, 1149–1157. doi:10.1002/cbdv.201500514. [\[Crossref\]](#) [\[PubMed\]](#)
24. Ding, C.Y.G; Pang, L.M; Liang, Z.X; Goh, K.K.K; Glukhov, E; Gerwick, W.H; Tan, T.L. MS/MS-Based molecular networking approach for the detection of aplysiatoxin-Related compounds in environmental marine cyanobacteria. *Mar Drugs*. **2018**, *16*, 1–15. doi:10.3390/md16120505. [\[Crossref\]](#) [\[PubMed\]](#)
25. Afoullouss, S; Sanchez, A.R; Jennings, L.K; Kee, Y; Allcock, A.L; Thomas O.P. Unveiling the Chemical Diversity of the Deep-Sea Sponge *Characella pachastrelloides*. *Marine Drugs*. **2022**, *20*, 1-52. doi:10.3390/md20010052. [\[Crossref\]](#) [\[PubMed\]](#)
26. Zhi-Dong Jiang, Paul R. Jensen and WF. Lobophorins A and B, new antiinflammatory macrolides produced by a tropical marine bacterium. *Bioorganic & Medicinal Chemistry Letters*. **1999**, *9* (14), 2003–2006. [\[Crossref\]](#) [\[PubMed\]](#)
27. Rong-Bian Wei, Tao Xi, Jing Li, Ping Wang, Fu-Chao Li Y-CL and Qin, S. Lobophorin C and D, New Kijanamicin Derivatives from a Marine Sponge-Associated Actinomycetal Strain AZS17. *Marine Drugs*. **2011**, *9* (3), 359–368. doi:10.3390/md9030359. [\[Crossref\]](#) [\[PubMed\]](#)
28. Lin, Z; Koch, M; Pond, C.D; Mabeza, G; Seronay, R.A; Concepcion, G.P....& Schmidt, E.W. Structure and activity of lobophorins from a turrid mollusk-associated *Streptomyces* sp. *Journal of Antibiotics*. **2014**, *67*(1), 121–126. doi:10.1038/ja.2013.115. [\[Crossref\]](#) [\[PubMed\]](#)
29. Li-Man Zhou, Fan-Dong Kong, Qing-Yi Xie, Qing-Yun Ma, Zhong Hu, You-Xing Zhao and Luo, D.Q. Divergolides T–W with Apoptosis Inducing Activity from the Mangrove Derived Actinomycete *Streptomyces* sp. KFD18. *Marine Drugs*. **2019**, *17*(4), 1–9. doi:10.3390/md17040219. [\[Crossref\]](#) [\[PubMed\]](#)
30. Iwasaki, H; Kamisango, K.I; Kuboniwa, H; Sasaki, H; Matsubara, S. 3'-Deoxytalopiericidin a1 a novel analog of antitumor antibiotics from oligotroph. *J. Antibiot*. **1991**, *44*, 451–452. doi:10.7164/antibiotics.44.451. [\[Crossref\]](#) [\[PubMed\]](#)
31. Irohara Mori; Shinji Funayama YS; Omura, K.K and Omura, S. A new antibiotic, 13-Hydroxyglucopiericidin A isolation, structure, elucidation and biological characteristics. *J. Antibiot*. **1990**, XLIII, 1329–1331. [\[Crossref\]](#) [\[PubMed\]](#)
32. Niu, S; Li, S; Chen, Y; Tian, X; Zhang, H; Zhang, G., ...& Zhang, C. Lobophorins E and F, new spirotetronate antibiotics from a South China Sea-derived *Streptomyces* sp. SCSIO 01127. *J. Antibiot*. **2011**, *64*(11), 711–716. doi:10.1038/ja.2011.78. [\[Crossref\]](#) [\[PubMed\]](#)
33. Chen, C; Wang, J; Guo, H; Hou; Yang, N; Ren, B....& Zhang, L. Three antimycobacterial metabolites identified from a marine-derived *Streptomyces* sp. MS100061. *Applied microbiology and biotechnology*. **2013**, *97*(9), 3885–3892. doi:10.1007/s00253-012-4681-0. [\[Crossref\]](#) [\[PubMed\]](#)
34. Patricia, G; Cruz, Andrew M. Fribley; Justin R. Miller; Martha J. Larsen; Pamela J. Schultz; Renju T. Jacob; Giselle Tamayo-Castillo; Randal J. Kaufman and Sherman, D.H. Novel Lobophorins inhibit Oral Cancer Cell Growth and Induce Atf4 and

- chop-Dependent Cell Death in Murine Fibroblasts. *ACS Medicinal Chemistry Letters*. **2015**, 6(8), 877-881. doi:10.1021/acsmchemlett.5b00127. [[Crossref](#)] [[PubMed](#)]
35. Zhang, C; Ding, W; Qin, X; and Ju, J. Genome Sequencing of *Streptomyces olivaceus* SCSIO T05 and Activated Production of Lobophorin CR4 via Metabolic Engineering and Genome Mining CR4. *Marine Drugs*. **2019**, 17(10), 1–11. doi:10.3390/md17100593. [[Crossref](#)] [[PubMed](#)]
36. Luo, M; Tang, L; Dong, Y; Huang, H; Deng, Z; Sun, Y. Antibacterial natural products lobophorin L and M from the marine-derived *Streptomyces* sp. 4506. *Nat. Prod. Res.* **2021**, 35(24), 1–7. doi:10.1080/14786419.2020.1797730. [[Crossref](#)] [[PubMed](#)]
37. Li, X; Yi-min, CH; Ya-xin, H; Guo-jian, Z; Tian-jiao, Z; Ji-xing, P; Dehai L.C.Q. Isolation and structure elucidation of lobophorin N from the South China Sea-derived *Streptomyces* sp. HDN1844000. *J. Antibiot.* **2021**, 40. doi:1002-3461(2021)05-009-07. [[Crossref](#)]
38. Nobutaka Takahashi; Satoshi Miyamoto, Rinpei Mori; Saburo Tamura. Isolation and Physiological Activities of Piericidin A, A Natural Insecticide Produced by *Streptomyces*. *Agric. Biol. Chem.* **1963**, 27, 576–582. doi:10.1080/00021369.1963.10858144. [[Crossref](#)]
39. Nobutaka Takahashi, Akinori Suzuki, Satoshi Miyamoto and Saburo Tamura. Structure of piericidin B and stereochemistry of piericidins. *Tetrahedron Lett.* **1967**, 21, 1961–1964. [[Crossref](#)]
40. Hiroshi Nishioka; Tsutomu Sawa; Kunio Isshiki, Yoshikazu Takahashi, Naganawa, H; Matsuda, N; Hattori, S; Hamada, M; Takeuchi K.T. Isolation and structure determination of a novel phosphatidylinositol turnover inhibitor, Piericidin B1 N-Oxide. *J. Antibiot.* **1991**, 44, 1283–1285. [[Crossref](#)] [[PubMed](#)]
41. Yoichi Hayakawa; Shingo Shirasaki; Takashi Kawasaki; Yoshihide Matsuo; Kyoko Adachi Y.S. Structures of New Cytotoxic Antibiotics, Piericidins C7 and C8. *J Antibiot.* **2007**, 60(3), 201–203. [[Crossref](#)] [[PubMed](#)]
42. Yoshida S; Nagao Y; Watanabe A; Takahashi N. Structure-activity relationship in piericidins inhibitors on the electron transport system in mitochondria. *Agric Biol Chem.* **1980**, 44(12), 2921–2924. doi:10.1080/00021369.1980.10864438. [[Crossref](#)]
43. Shaaban, K.A; Helmke, E; Kelter, G; Fiebig, H.H; Laatsch, H. Glucopiericidin C: A cytotoxic piericidin glucoside antibiotic produced by a marine *Streptomyces* isolate. *Journal of Antibiotics*. **2011**, 64(2), 205–209. doi:10.1038/ja.2010.125. [[Crossref](#)] [[PubMed](#)]
44. Jun-ya Ueda; Takushi Togashi; Susumu Matukura; Aya Nagai; Takuji Nakashima; Hisayuki Komaki; Kozue Anzai; Shigeaki Harayama; Takayuki Doi; Takashi Takahashi; Tohru Natsume; Yasutomo Kisu; Naoki Goshima; Nobuo Nomura; Motoki Takagi. A Novel Nuclear Export Inhibitor JBIR-02, a New Piericidin Discovered from *Streptomyces* sp. ML55. *J. Antibiot.* **2007**, 60(7) 459–462. [[Crossref](#)] [[PubMed](#)]
45. Wang, M; Carver, J.J; Phelan, V.V; Sanchez, L.M; Garg, N; Peng, Y., ...& Bandeira, N. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nature biotechnology*. **2016**, 34(8), 828–837. doi:10.1038/nbt.3597. [[Crossref](#)] [[PubMed](#)]

SUPPLEMENTARY DATA

Table S1. Complete 16S rRNA sequences of two *Streptomyces* spp.

Strain	No. of bp	BLASTn closest homolog (accession #) organism	Completeness /Identity (%)	Variation ratio
<i>Streptomyces</i> sp. B-81	1345	<i>Streptomyces olivaceus</i> NRRL B-3009 ^T	100.0/(99.93)	1/1345
		<i>Streptomyces pactum</i> NBRC 13433 ^T	99.6/(99.93)	1/1345
		<i>Streptomyces parvulus</i> NBRC 13193 ^T	99.2/(99.03)	13/1340
<i>Streptomyces</i> sp. 796.1	1370	<i>Streptomyces buecheerae</i> AC541 ^T	100/(99.70)	4/1352
		<i>Streptomyces youssoufiensis</i> X4 ^T	100/(99.48)	7/1352
		<i>Streptomyces zagrosensis</i> HM 1154 ^T	100/(99.48)	7/1352
		<i>Streptomyces philanthi triangulum</i> ^T	90.6/(99.45)	7/1265

Table S2. Complete 16S rRNA sequences of three *Streptomyces* spp.

>*Streptomyces* sp. B-81 Accession Number NCBI (MW562807)

CTGGCGGCGTGCTTAACACATGCAAGTCGAACGATGAACCACTTCGGTGGGGATTAGTGCGAACGGGTGAG 510
TAACACGTGGGCAATCTGCCCTGCACTCTGGGACAAGCCCTGGAAACGGGGTCTAATACCGGATATTGACCTT 511
CACGGGCATCTGTGAGGTTTCGAAAGCTCCGGCGGTGCAGGATGAGCCCGCGGCCTATCAGCTTGTGGTGAGG 512
TAATGGCTCACCAAGGCGACGACGGGTAGCCGGCCTGAGAGGGCGACCGGCCACACTGGGACTGAGACACGG 513
CCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGCGAAAGCCTGATGCAGCGACGCCGCG 514
TGAGGGATGACGGCCTTCGGGTTGTAAACCTCTTTCAGCAGGGAAGAAGCGAAAGTGACGGTACCTGCAGAA 515
GAAGCGCCGGCTAACTACGTGCCAGCAGCCGCGTAATACGTAGGGCGCAAGCGTTGTCCGGAATTATTGGG 516
CGTAAAGAGCTCGTAGGCGGCTTGTACGTCGGTGTGAAAGCCCGGGGCTTAACCCCGGGTCTGCAGTCGAT 517
ACGGGCAGGCTAGAGTTCGGTAGGGGAGATCGGAATTCCTGGTGTAGCGGTGAAATGCGCAGATATCAGGAG 518
GAACACCGGTGGCGAAGGCGGATCTCTGGGCCGATACTGACGCTGAGGAGCGAAAGCGTGGGGAGCGAACA 519
GGATTAGATACCCTGGTAGTCCACGCCGTAACGGTGGGCACTAGGTGTGGGCAACATTCCACGTTGTCCGTG 520
CCGCAGCTAACGCATTAAGTGCCCCGCTGGGGAGTACGGCCGCAAGGCTAAAACCTCAAAGGAATTGACGGG 521
GGCCCGCACAAGCGGCGGAGCATGTGGCTTAATTTCGACGCAACGCGAAGAACCTTACCAAGGCTTGACATAC 522
ACCGGAAACGGCCAGAGATGGTCGCCCCCTTGTGGTTCGGTGTACAGGTGGTGCATGGCTGTCGTCAGCTCGTG 523
TCGTGAGATGTTGGGTTAAGTCCCGCAACGAGCGCAACCCTTGTCCCGTGTGCCAGCAAGCCCTTCGGGGG 524
TGTTGGGGACTCACGGGAGACCGCCGGGGTCAACTCGGAGGAAGGTGGGGACGACGTCAAGTCATCATGCC 525
CTTATGTCTTGGGCTGCACACGTGCTACAATGGCCGGTACAATGAGCTGCGATACCGCAAGGTGGAGCGAATC 526
TCAAAAAGCCGGTCTCAGTTCGGATTGGGGTCTGCAACTCGACCCCATGAAGTCGGAGTCGCTAGTAATCGCA 527
GATCAGCATTGCTGCGGTGAATACGTTCCCGGGCCTTGTACA 528

>*Streptomyces* sp 796.1 Accession Number (MG654686)

GATTCCCTGGCTCAGGACGAACGCTGGCGGGCGTGCTTAACACATGCAGTCGAACGATGAGCCGCTTCGGTGG 530
TGGATTAGTGGCGAACGGGTGAGTAACACGTGGGCAATCTGCCCTGCACTCTGGGACAAGCCCTGGAAACGG 531
GGTCTAATACCGGATATGACTACCGATCGCATGGTTGGTGGTGGAAAGCTCCGGCGGTGCAGGATGAGCCCGC 532
GGCCTATCAGCTTGTGGTGGGGTGTATGGCCTACCAAGGCGACGACGGGTAGCCGGCCTGAGAGGGCGACCG 533
GCCACACTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGCGA 534
AAGCCTGATGCAGCGACGCCGCTGAGGGATGACGGCCTTCGGGTTGTAAACCTCTTTCAGCAGGGAAGAAG 535
CGCAAGTGACGGTACCTGCAGAAGAAGCGCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGCGC 536
GAGCGTTGTCCGGAATTATTGGGCGTAAAGAGCTCGTAGGCGGCTTGTTCGCGTCGGATGTGAAAGCCCGGGG 537
TTAACTCCGGGTCTGCATTTCGATACGGCAGGCTAGAGTTCGGTAGGGGAGATCGGAATTCCTGGTGTAGCGG 538
TGAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGATCTCTGGGCCGATACTGACGCTGAGGAG 539
CGAAAGCGTGGGGAGCGAACAGGATTAGATACCCTGGTAGTCCACGCCGTAACGTTGGGAACTAGGTGTGG 540
GCGACATTCCACGTCGTCCGTGCCGAGCTAACGCATTAAGTTCCCCGCTGGGGAGTACGGCCGCAAGGCTA 541
AAACTCAAAGGAATTGACGGGGGCCCGCACAAGCGGCGGAGCATGTGGCTTAATTCGACGCAACGCGAAGAA 542
CCTTACCAAGGCTTGACATACACCGGAAACGGCCAGAGATGGTCGCCCCCTTGTGGTTCGGTGTACAGGTGGTG 543
CATGGCTGTCGTCAGCTCGTGTGAGATGGTTGGGTTAAGTCCCGCAACGAGCGCAACCCTTGTCTGTGTT 544
GCCAGCATGCCTTTCGGGGTGTATGGGACTCACAGGAGACTGCCGGGGTCAACTCGGAGGAAGGTGGGGACG 545
ACGTCAAGTCATCATGCCCTTATGTCTTGGGCTGCACACGTGCTACAATGGCCGGTACAATGAGCTGCGATG 546
CCGTGAGGTGGAGCGAATCTCAAAAAGCCGGTCTCAGTTCGGATTGGGGTCTGCAACTCGACCCCATGAAGTC 547
GGAGTCGCTAGTAATCGCAGATCAGCATTGCTGCGGTGAATACGTTCCCGGGCCTTGTAAACACA 548

549

Top: mzspec:GNPS:TASK-b7a29b5a78864223bde01fb8af0c5eff-spectra/specs_ms.mgf:scan:5653

Precursor m/z: 1157.6379 Charge: 0

Bottom: mzspec:GNPS:GNPS-LIBRARY:accession:CCMSLIB00000223869

Precursor m/z: 1157.6400 Charge: 1

Cosine similarity = 0.9168




Fig. S1. MS/MS match between GNPS database (green) lobophorin A C61H93N2O19 (5653) compound (1) from *Streptomyces* sp. B-81 extract (black)

mzspec:GNPS:TASK-b7a29b5a78864223bde01fb8af0c5eff-spectra/specs_ms.mgf:scan:5178

Precursor m/z: 753.4320 Charge: 1




Fig. S2. Lobophorin H in the GNPS database or manually by accurately determining which ion precursor m/z 753.4320 showed mass errors below 0.10 ppm C42H60N2O10 (5178) from *Streptomyces* sp. B-81 extract.

mzspec:GNPS:TASK-b7a29b5a78864223bde01fb8af0c5eff-spectra/specs_ms.mgf:scan:4097
Precursor m/z: 554.2750 Charge: 1




Fig. S3. Divergolide A in the GNPS database or manually by accurate determination of which ion precursor m/z 554.2750 showed mass errors below 0.10 ppm C₃₁H₃₉NO₈ (4097) compound (3) from *Streptomyces* sp. B-81 extract.

587
588
589
590
591

mzspec:GNPS:TASK-b7a29b5a78864223bde01fb8af0c5eff-spectra/specs_ms.mgf:scan:3981
Precursor m/z: 536.2650 Charge: 1




Fig. S4. Divergolide B in the GNPS database or manually by accurate determination of which ion precursor m/z 536.2650 showed mass errors below 0.10 ppm C₃₁H₃₇NO₇ (3981) compound (4) from *Streptomyces* sp. B-81 extract.

592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608

mzspec:GNPS:TASK-b7a29b5a78864223bde01fb8af0c5eff-spectra/specs_ms.mgf:scan:4068
Precursor m/z : 550.2440 Charge: 1




Fig. S5. Divergolide C in the GNPS database or manually by accurate determination of which ion precursor m/z 550.2440 showed mass errors below 0.10 ppm C₃₁H₃₅NO₈ (4068) compound (5) from *Streptomyces* sp. B-81 extract.

Top: mzspec:GNPS:TASK-b7a29b5a78864223bde01fb8af0c5eff-spectra/specs_ms.mgf:scan:4214
Precursor m/z : 578.3330 Charge: 0

Bottom: mzspec:GNPS:GNPS-LIBRARY:accession:CCMSLIB00004678798
Precursor m/z : 578.3320 Charge: 1

Cosine similarity = 0.8682




Fig. S6. MS/MS match between GNPS database (green) glucopiericidin A C₃₁H₄₇NO₉ (4214) compound (6) from *Streptomyces* sp. 796.1 extract (black)

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

Top: mzspec:GNPS:TASK-b7a29b5a78864223bde01fb8af0c5eff-spectra/specs_ms.mgf:scan:3145

Precursor m/z : 414.2650 Charge: 1

Bottom: mzspec:GNPS:GNPS-LIBRARY:accession:CCMSLIB00004678734

Precursor m/z : 414.2640 Charge: 1

Cosine similarity = 0.9234




Fig. S7. MS/MS match between GNPS database (green) dehydropiericidin A1a C₂₅H₃₅NO₄ (3145) compound (7) from *Streptomyces* sp. 796.1 extract (black).