Large-scale duplication events underpin population-level flexibility
in tRNA gene copy number in *Pseudomonas fluorescens* SBW25

Zahra Khomarbaghi[†], Wing Y. Ngan[†], Gökçe B. Ayan, Sungbin Lim, Gunda Dechow-Seligmann,
Pabitra Nandy, Jenna Gallie[*]

Microbial Evolutionary Dynamics Research Group, Department of Theoretical Biology, Max Planck
Institute for Evolutionary Biology, Plön, 24306, Germany
8

9 [†] Joint authors: these authors contributed equally to this work.
10 [*] To whom correspondence should be addressed. Email: gallie@evolbio.mpg.de

12 Present Address: Zahra Khomarbaghi, Division of Evolution and Genomic Sciences, School of
13 Biological Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester,
14 M13 9PL, United Kingdom
15

ABSTRACT

The complement of tRNA genes within a genome is typically considered to be a (relatively) stable
characteristic of an organism. Here we demonstrate that bacterial tRNA gene set composition can be
more flexible than previously appreciated, particularly regarding tRNA gene copy number. We report
the high-rate occurrence of spontaneous, large-scale, tandem duplication events in laboratory
populations of the bacterium *Pseudomonas fluorescens* SBW25. The identified duplications are up to
~1 Mb in size (~15 % of the wildtype genome) and are predicted to change the copy number of up to
917 genes, including several tRNA genes. The observed duplications are inherently unstable: they
occur, and are subsequently lost, at extremely high rates. We propose that this unusually plastic type
of mutation provides a mechanism by which tRNA gene set diversity can be rapidly generated, while
simultaneously preserving the underlying tRNA gene set in the absence of continued selection. That
is, if a tRNA set variant provides no fitness advantage, then high-rate segregation of the duplication
ensures the maintenance of the original tRNA gene set. However, if a tRNA gene set variant is
beneficial, the underlying duplication fragment(s) may persist for longer and provide raw material for
further, more stable, evolutionary change.

INTRODUCTION

The complement of tRNA genes encoded by an organism (the 'tRNA gene set') is an important
determinant of translational efficiency (1, 2) and disease trajectories (3–5). In bacteria, tRNA gene set
composition varies considerably, yet the tRNA gene set of any given bacterial strain is typically
treated as a stable entity. For example, the tRNA gene sets of *Pseudomonas fluorescens* strains
SBW25 and Pf0-1 are reported to respectively contain 66 and 73 canonical tRNA genes, encoding 39

and 40 distinct species of mature tRNA (each with a unique anticodon-amino acid combination) (6–8). In this work we highlight that, at the population level, the tRNA gene set of a bacterial strain may be more flexible than previously appreciated, particularly with regard to the number of gene copies encoding each type of tRNA.

The composition of tRNA gene sets differs across the bacterial phylogeny; extant bacterial tRNA gene sets differ in both the tRNA species that they encode, and in the number of gene copies coding for each tRNA species (7, 9–15). Variations in tRNA species encoded are possible because of wobble base pairing, the process by which some tRNAs can – either naturally (16) or through post-transcriptional modifications (17, 18) – translate multiple codons. Ultimately, wobble base pairing allows all 61 sense codons to be translated by far fewer than the number of directly corresponding tRNA species, with most bacteria encoding between 25 and 46 different types of tRNA (7, 9, 19). In contrast to this apparently reductionist strategy, the tRNA species that are present within a bacterial genome may be encoded by multiple – and often, identical – gene copies (7, 10).

The content of a tRNA gene set has a strong influence on the composition of the mature tRNA pool, which in turn is a major determinant of the speed and accuracy of translation (1, 2, 20). The capacity of a tRNA set for efficient translation depends on a number of additional factors, many of which co-evolve with tRNA set composition. Prominent examples include (i) which post-transcriptional tRNA modification enzymes are present (affecting anticodon-codon matching patterns) (15, 18, 21, 22), (ii) codon usage (which influences the demand for specific tRNAs) (23–29), and (iii) environmental conditions that affect anticodon-codon base pairing properties (*e.g.*, pH, temperature) (30, 31). Evolutionary optimization of translational efficiency is expected to shape the evolution of each of these factors, particularly under conditions where efficient translation is most important (*e.g.*, during rapid bacterial growth) (32–35). For example, bacteria with faster growth rates tend to carry tRNA gene sets with fewer tRNA species, encoded by higher gene copy numbers (11). Empirically, multiple copies of several *Escherichia coli* tRNA genes have been demonstrated to be advantageous only under conditions that support rapid growth (and hence, rapid translation) (36).

The molecular mechanisms by which tRNA genes are gained and lost from tRNA gene sets have been inferred by phylogenetic analyses. Such computational studies have provided evidence for the evolution of extant bacterial tRNA gene sets via a number of routes, including: (i) mutation of tRNA genes to encode different tRNA species ('anticodon switching'), (ii) loss of tRNA genes by deletion, and (iii) acquisition of new tRNA genes, either by within-genome duplication events, or by horizontal gene transfer (10, 12–14, 37–39). In addition, we recently provided an empirical example of a bacterial tRNA gene set undergoing adaptive evolution; elimination of tRNA-Ser(CGA) from the mature tRNA pool of the fast-growing, model bacterium *P. fluorescens* SBW25 was compensated by within-genome duplication events that serve to increase the copy number of the tRNA gene *serTGA* (and corresponding levels of tRNA-Ser(UGA) in the mature tRNA pool) (6). Others have reported that translational challenges imposed by (i) novel codon usage patterns, or (ii) the fitness costs associated with antibiotic resistance mutations can be readily alleviated by similar duplication (and, sometimes, downstream amplification) events that affect the copy numbers of remaining tRNA genes (40, 41).

Together, these studies demonstrate a considerable degree of evolutionary flexibility in the

composition of bacterial tRNA gene sets.

In the above-mentioned study of the tRNA gene set of *P. fluorescens* SBW25 (6), the

adaptive increase in *serTGA* copy number was shown to be mediated through the tandem duplication

of large sections of the *P. fluorescens* SBW25 chromosome (ranging in size from ~45 kb to ~290 kb).

In addition to the 91 bp target gene (*serTGA*), these duplication events affect the dosage of up to a

further 278 genes, and generate considerable alterations in chromosome size and architecture.

Similar types of large-scale, tandem duplications are a pervasive phenomenon in bacterial genomes

(42–46) and – despite their large size and widespread gene dosage effects – they can provide a net

selective advantage in response to a range of selective pressures (47–53). Notably, these types of

mutations have been reported to arise spontaneously, and at extremely high rates; estimates range

from frequencies of 10^{-1} to 10^{-5} duplications per cell per generation (43, 54–57), several orders of

magnitude higher than those expected for single nucleotide polymorphisms (SNPs) (58). Given their

pervasiveness and ease of occurrence, such large-scale, tandem duplications challenge the

prevailing view of fixed, stable tRNA gene sets, particularly in large, rapidly growing bacterial

populations.

In this work, we firstly show that large-scale, tandem duplication events affecting the *P.*

fluorescens SBW25 tRNA gene content are not limited to the genomic region surrounding *serTGA*

(located at ~4.16 Mb on the SBW25 chromosome). Specifically, we demonstrate that a fitness defect

caused by deleting a locus containing four tRNA genes (two copies of each of *gluTTC* and *glyGCC*) is

repeatedly and rapidly compensated by large-scale, tandem duplication events encompassing the

remaining *glyGCC* gene (at genomic position ~2.38 Mb). Secondly, we provide empirical evidence

that these duplication fragments arise, and are lost, at very high frequencies; we show that in each

independently evolving mutant population, multiple distinct duplication fragments (and hence, tRNA

gene sets) rapidly arise and compete.

MATERIALS AND METHODS

Strains and growth conditions. A list of strains is provided in Supplementary Table S1; construction

details can be found in Supplementary Text S1. Unless otherwise stated, *P. fluorescens* SBW25 was

grown in King's Medium B (KB) (59) for ~16 hours at 28˚C with shaking. For growth curves, strains

were grown in either liquid KB, or minimal (M9) medium (60) supplemented with 4 % (w/v) glucose.

During engineering procedures, *E. coli* strains were grown in Lysogeny Broth (LB) for 16-18 hours at

37˚C with shaking.

Photography. Colonies were grown for the time indicated, at room temperature (~20-22˚C) or 28˚C,

on KB agar. Colonies were visualized under a Leica MS5 dissection microscope and photographed

with a VisiCam® 1.3 (VWR International). Photographs were cropped and, where noted in figure

legends, the exposure and/or brightness uniformly altered in Preview (v11.0) to improve visibility.

Genotype growth curves. Seven colonies per genotype were inoculated into individual wells of a 96-

well pre-plate (each containing 200 µL KB), and grown overnight (28˚C, 200 rpm). From each well, 2

µL of grown culture was transferred into 198 µL fresh KB, covered with a Breathe-Easy® membrane,

and grown at 28˚C in an Epoch 2 Microplate Spectrophotometer (Agilent BioTek). Absorbance at

600nm (OD_{600}) was measured at five-minute intervals, with 5 seconds of 3 mm orbital shaking

preceding each read. Gen5^{TM} Data Analysis Software (version 3.00.19) was used to calculate the

maximum growth rate (the calculated value of the mean slope, V_{max}) and lag time (time interval

between the line of maximum slope of the propagation phase and the absorbance baseline at time

zero) for each growth curve, with the following parameters: calculation window=1-8 hours (KB) or 1-10

hours (M9); sliding time interval per calculation ("*n*")=6 points). The data was then exported to Excel

for further processing and drawing of graphs (Figures 3 and 4). Where assumptions are satisfied,

parametric one-way ANOVA was used to test for differences in the mean maximum growth rate

between pairs of genotypes. Where assumptions are violated, non-parametric Dunn's tests, followed

by the Benjamini-Hochberg procedure to correct for multiple comparisons, were used to test for

differences in the median lag time between pairs of genotypes.

Competition assays. One-to-one competition experiments were used to determine the relative

fitness of 9 genotype pairs under conditions similar to those of a single growth cycle of the evolution

experiment (see below). Competitions were performed in two blocks, with between 5 and 8 replicates

per pair of genotypes. Single colonies of each genotype were grown on KB agar (48 hours, room

temperature) and, for each competition replicate, a single colony of each competitor was inoculated

into an overnight culture (4 mL KB, 28˚C, 200 rpm). Once grown, the two competitors for each

competition were mixed in a ~1:1 ratio. Ten µL of each mixture was used to inoculate 10 mL KB in a

50 ml Falcon tube, and grown for 24 hours (28˚C, 200 rpm). Cell samples taken from each

competition tube at the start (T_0) and end (T_{24}) of the incubation period were serially diluted and plated

on LB agar containing 60 µg mL^{-1} X-gal (48 hours, room temperature). The number of colonies per

competitor were recorded at T_0 and T_{24}, where differentiation between competing genotypes was

achieved through distinct colony morphologies and/or colours (in competitions including neutrally

marked SBW25-*lacZ* (61)). The change in the ratio of the competitors was calculated, and relative

fitness of the competitors determined (62). Parametric one-sample *t*-tests or, where normality

assumptions were violated, non-parametric Wilcoxon rank sum tests, were used to detect statistical

differences in the mean relative fitness of genotype pairs. The Bonferroni procedure was

subsequently used to correct for multiple comparisons.

Evolution experiment. SBW25 (wildtype) and ΔEGEG (quadruple tRNA gene deletion mutant) were

streaked from glycerol stocks onto KB agar and grown (48 hours, 28˚C). Five single colonies per

genotype were grown overnight (4 mL KB; 28˚C, 200 rpm). Grown cultures ("day 0") were vortexed

and 100 µL of each used to inoculate 9.9 mL KB in a 50 mL Falcon tube (28˚C, shaking, 24 hours).

Every 24 hours thereafter, 1 % (100 µL) of each culture was transferred to a fresh 9.9 mL KB, and a

sample of the population frozen in glycerol saline at -80˚C. The experiment was continued until day 28.

The later days (~day 21 onwards) of mutant lines M4 and M5 were discovered to contain external

contaminants (see Supplementary Figure S1), and so were removed from downstream population

analyses (*i.e.*, Figure 7 and Figure 8). Extensive checks for external and cross contamination events

were performed on the remaining 8 lines (including day 28 population-level whole genome re-

sequencing, population colony morphology checks, and population growth curve analyses).

Population growth curves. Forty populations of interest (10 evolving lines, at each of 4 time points)

were retrieved from glycerol stocks by re-suspending large scrapings in 200 µL sterile Ringer's

solution. Each re-suspension was used to inoculate 4 wells of 96-well pre-plates, and grown overnight

(*i.e.*, 4 technical replicates; 2 µL re-suspension into 198 µL KB per replicate). Subsequently, grown

pre-cultures were transferred to a new 96-well plate (2 µL pre-culture into 198 µL fresh KB), and the

protocol continued as described above for the genotype growth curves. This protocol was repeated

three times per population (*i.e.*, 3 biological replicates), giving a total of 12 replicates per population of

interest (4 technical replicates x 3 biological replicates). The final population growth curves are

presented in Supplementary Figure S1.

Genotype whole genome re-sequencing. Six single genotypes (W1-1, M1-1, M2-1, M3-1, M4-1,

M5-1) were purified from day 21 of the evolution experiment, including 1 genotype from each of the 5

mutant lines (M1-M5), and 1 from a representative wildtype control line (W1). Genomic DNA was

isolated from 1 mL of overnight culture of each genotype (DNeasy Blood & Tissue Kit; Qiagen).

Paired-end, 250-bp reads were generated with an Illumina MiSeq instrument at The Max Planck

Institute for Evolutionary Biology (Ploen, Germany), via standard procedures. Raw reads are available

at NCBI Sequence Read Archive (SRA accession number: PRJNA790786) (63). A minimum of 1.08

million raw reads per genotype were obtained and aligned to the SBW25 reference genome sequence

(NC_012660.1) (8) using *breseq* (64–66) and Geneious Prime (v2023.2.1). A minimum mean

coverage of 36.4 reads per bp was obtained. Per genotype, a list of the raw differences predicted by

breseq was manually curated to give a list of final mutation predictions (see Supplementary Table S2

for lists and further details). Each mutation prediction was confirmed by standard PCR or duplication

junction PCR (see below) and, where indicated, Sanger sequencing (see Supplementary Text S2).

Mature tRNA pool sequencing (YAMAT-seq). Three independent replicates of 8 genotypes (*i.e.*, 24

samples) were used for YAMAT-seq (67) according to the method described in (6). Briefly, (i) per

sample, a single colony was chosen from KB agar (in the case of genotypes carrying duplication

fragments, rapidly-growing colonies were chosen in an attempt to mitigate the inherent instability of

the duplication fragment; see stability test method below), grown in overnight KB culture, diluted 1/10

in fresh KB, and grown to mid-exponential phase, (ii) total RNA was isolated from each rapidly-

growing liquid culture using a TRIzol Max Bacterial RNA isolation kit, (iii) amino acids were removed

from mature tRNAs by deacylation treatment with 20 mM Tris-HCl, (iv) Y-shaped, DNA/RNA hybrid

adapters (Eurofins; see Supplementary Table S1) were annealed to the conserved, exposed 5'-

NCCA-3' and 3'-inorganic phosphate-5' ends of non-acylated tRNAs, and ligated T4 RNA ligase 2, (v)

ligation products were reverse-transcribed into cDNA using SuperScript™ III reverse transcriptase, (vi)

cDNA products were amplified, and a unique barcode added, with eleven cycles of PCR using

Phusion™, (vii) products were quantified using a Bioanalyzer and fluorescent Nanodrop, and mixed in

equimolar amounts, (viii) the mixed sample was run on a 5 % polyacrylamide gel, the bands between

~210 bp and ~280 bp (tRNAs plus adapters and barcodes) cut out and co-extracted overnight (by

soaking in deionized, ultrapure water (Illumina)), (ix) the extracted liquid sample was collected by

centrifugation through a filtered microcentrifuge tube (cellulose acetate membrane with pore size 0.45

µm; Merck CLS-8162), (x) the purified product was quantified and deep sequenced at the Max Planck

Institute for Evolutionary Biology (Ploen, Germany). Single-end, 150 bp reads were generated with a

NextSeq 550 Output v2.5 kit (Illumina) (NCBI Gene Expression Omnibus accession number

GSE196705) (68).

YAMAT-seq analysis. The raw YAMAT-seq sequencing reads were split into 24 samples by

extracting exact matches to each unique, 6-bp Illumina index. The resulting 24 set of raw reads, each

containing a minimum of 1,470,795 reads, were analysed using Geneious Prime (version 11.1.4).

Reads of the expected length (80-151 bp) were extracted (resulting in at least 99.8 % read retention

per sample). The extracted reads were aligned to a set of 42 reference tRNA sequences from SBW25

(see Supplementary Text S4) (Geneious settings: per read, up to 10 % mismatches, gaps of <3 bp,

up to five ambiguities; discard reads that align equally well to more multiple reference sequences).

The unused reads for each sample were *de novo* assembled, and the resulting contigs were checked

to ensure that none contained substantial numbers of tRNA reads. The within-sample proportion of

reads aligned to each tRNA species was calculated. Next, mean mature tRNA proportions were

calculated for each genotype across the 3 replicates (Supplementary Table S4). DESeq2 (69) was

used in R (version 3.6.0) to detect tRNA expression differences between pairs of genotypes

(Supplementary Table S5). DESeq2 corrects for multiple testing (Benjamini-Hochberg procedure) (70).

Stability test. Duplication fragment stability was estimated using colony morphology stability tests,

according to the method detailed in (71). Briefly, 10 genotypes (WT, rWT, ΔEGEG, ΔglyGCC, W1-1,

M1-1, M2-1, M3-1, M4-1, and M5-1) were streaked from glycerol stocks onto KB agar plates and

grown (48 hours, 28˚C). Five colonies per genotype were used to inoculate 4 mL KB cultures (28˚C,

200 rpm, overnight). For duplication-carrying genotypes, all chosen colonies were large (*i.e.*, were

expected to retain the duplication). Cultures were then dilution plated on KB agar (48 hours, room

temperature), and the number of large and small colonies per plate was enumerated. Between 41 and

295 colonies were recorded per replicate, with no more than ~200 colonies counted on any given

plate. For each replicate, the proportion of small colonies was calculated. A non-parametric Kruskal-

Wallis test was used to test for evidence of a difference in the median of at least 1 founder (*p*=6.25e-

07). To determine which genotypes showed a different median rate of small colony enumeration as

compared with SBW25 (wildtype), Dunn's test was used with Benjamini-Hochberg procedure to adjust

for multiple comparisons. A Welch non-parametric two-sample *t*-test was used to test for a difference

in the mean proportion in small colonies arising from the genotype carrying the small duplication

fragment (M3-1), and those carrying a large duplication fragment (M1-1, M2-1, M4-1, M5-1).

Duplication junction PCRs. Duplication junction PCRs were used in this work to (i) confirm the

computationally-predicted duplications in genotypes isolated from day-21 of the evolution experiment

(see Supplementary Text S2), and (ii) reveal the evolutionary trajectories of duplication fragments in

evolving populations (method described here). For each trajectory of interest, 3 independent rounds

of duplication junction PCR were performed. Per round, PCR templates were prepared as follows: a

large chunk of the relevant glycerol stock was melted at room temperature, and 10 µL used to

inoculate a 4 mL KB culture (20 h at 28˚C, 200 rpm). Genomic DNA was extracted from 1 mL of

culture (DNeasy Blood & Tissue Kit; Qiagen) and quantified on a Nanodrop. Each genomic DNA

sample was used as a template for the following PCRs, at a final concentration of 1 ng µL^{-1}: (PCRs 1

and 2) the duplication junction PCR(s) of interest, targeting ~1 Mb duplications (with primer pair

M1and2junct_f/M5_junct_r), and/or ~0.49 Mb duplications (with primer pair M3_junct_f/r); (PCR 3) a

standard, control PCR targeting a genomic region unaffected by the duplication events (at genomic

position ~0.01 Mb; primer pair *glyQ*_f_/r). PCRs were performed with GoTaq Flexi (Promega) for 30

cycles of: 30 sec at 95˚C, 30 sec at 58˚C or 59˚C, 72˚C for 1 min per kb of expected product. Five µL

of each PCR product was mixed with loading dye and run on a 1 % agarose gel (SYBR Safe; Life

Technologies) at 90 V for ~45 min, against 1 kb DNA ladder (Promega) and photographed under UV

illumination (see Supplementary Figures S3 and S4 for examples). PCR product intensities were

quantified using ImageJ2 (version 2.14.0/1.54f) (72), and duplication junction PCR intensities were

normalized using the corresponding control PCR (see Supplementary Table S6). Final evolutionary

trajectories are plotted in Figure 7 and Supplementary Figure S5A.

Population whole genome re-sequencing. Nine samples were used for population-level whole

genome re-sequencing: 8 day-28 populations (from lines M1-M3 and W1-W5), and the non-evolved

SBW25 wildtype genotype (which, as a control, was treated as a population sample in downstream

computational analyses). For each of the 8 day-28 samples, the corresponding day-27 glycerol stock

was grown for 1 cycle of the evolution experiment (in 9.9 mL KB: 180 µL thawed glycerol stock,

consisting of ~100 µL cell culture and ~80 µL glycerol; 28˚C, 200 rpm, 24 hours). For the control

sample, a single SBW25 colony picked from KB agar was grown under the same conditions. Genomic

DNA was extracted from 2 mL of each of the nine overnight cultures (DNeasy Blood & Tissue Kit;

Qiagen). Library preparation and sequencing was performed by Novogene Europe (Cambridge, UK),

using Illumina technology on a NovaSeq 6000 platform. The 150-bp, paired-end raw reads are

available at NCBI Sequence Read Archive (SRA accession number: PRJNA1035442) (63). Raw

reads were quality-trimmed using fastp (settings: --detect_adapter_for_pe, --cut_right, minimum read

length 50 bp) (73). The polymorphism setting in *breseq* (64–66) was used to align each set of trimmed

reads to the reference sequence (SBW25; NC_012660.1). This gave a minimum mean coverage of

154.9 reads per bp, and a set of predicted mutations in each sample (Supplementary Tables S2 and

S7). Geneious Prime (v2023.2.1) was used to (i) confirm and extend the set of the duplication

fragments predicted by *breseq* (Supplementary Text S3), and (ii) rigorously check for contamination in

evolving populations (none was detected).

RESULTS

Severe reduction in essential tRNA species' gene copy number slows growth

The *P. fluorescens* SBW25 genome is predicted to contain 66 canonical tRNA genes encoding 39

tRNA species, 14 of which are coded for by multiple (between two and five) gene copies (Figure 1A)

(6, 8, 74). This work initially focusses on two such multicopy tRNA species: tRNA-Gly(GCC) (encoded

by three identical *glyGCC* gene copies) and tRNA-Glu(UUC) (encoded by four identical *gluTTC* gene

copies). Two copies of each gene are found together at chromosomal position ~2.03 Mb (the *gluTTC-*

glyGCC-gluTTC-glyGCC locus, hereafter referred to as "*EGEG*") (Figure 1A-B). The third *glyGCC*

copy is a lone tRNA gene at genomic position ~2.38 Mb, while the final two *gluTTC* genes occur,

together with two copies of *alaGGC*, at ~4.65 Mb. Both of the focal tRNA species are essential: at

least one functional copy of each of *glyGCC* and *gluTTC* is expected to be required for translation and

growth. Specifically, tRNA-Gly(GCC) is predicted to be the sole decoder of codons 5'-GGC-3'

(cognate match) and 5'-GGU-3' (G_{34}:U_3 wobble match), while tRNA-Glu(UUC) is required to translate

5'-GAA-3' (cognate) and 5'-GAG-3' (G_{34}:U_3 wobble) (Figure 1C).

We began by constructing two tRNA gene deletion genotypes in the SBW25 background: (i)

ΔEGEG, in which the entire *EGEG* locus was removed from SBW25 (leaving two copies of *gluTTC*

and only a single copy of *glyGCC*), and (ii) Δ*glyGCC*, in which the lone *glyGCC* copy at position ~2.03

Mb was removed (leaving two copies of *glyGCC*, and all four copies of *gluTTC*) (Figure 1D;

Supplementary Text S1). The two *gluTTC* copies at position ~4.65 Mb were not initially targeted

because the tRNA genes with which they co-localize are expected to be essential for survival (*i.e.,*

they are the only two copies of *alaGGC*, encoding the essential tRNA-Ala(GGC)) (Figure 1A).

Deletion of *EGEG* results in an immediately obvious growth defect; compared with SBW25,

ΔEGEG forms smaller colonies on KB and M9 agar plates, and grows more slowly in liquid culture

(Figure 2). Further, wildtype-like growth is restored by the re-introduction of *EGEG* into ΔEGEG

(generating the reconstructed wildtype, rWT). The observed reduction in growth is presumably due to

a drop in levels of either tRNA-Glu(UUC), tRNA-Gly(GCC), or both. However, we note that there is at

least partial redundancy among the *glyGCC* gene copies; removal of the lone *glyGCC* gene has no

discernible effect on SBW25 growth under the conditions tested (Δ*glyGCC* in Figure 2). This

observation is consistent with earlier reports that multiple copies of various tRNA genes (including

glyGCC) show a degree of redundancy in *E. coli* (36).

The ΔEGEG growth defect is rapidly compensated by evolution

In order to determine whether, and how, the ΔEGEG growth defect can be genetically compensated,

a serial transfer evolution experiment was performed. Ten independent lines were passaged through

repeated cycles of growth and dilution in shaken rich medium (KB), for 28 days (a total of ~200

generations over 28 days; 1 % daily bottleneck). Five evolutionary lines were founded by ΔEGEG

(lines M1-M5), and five by the wildtype (control lines W1-W5). Within 21 days (~150 generations)

each of the five mutant lines showed visibly improved growth, while the wildtype lines did not (from

this point on, wildtype line W1 is used as a representative control line) (see Supplementary Figure S1).

Single genotypes purified from day 21 of each mutant line (namely, genotypes M1-1, M2-1, M3-1, M4-

1, and M5-1) showed improved growth on agar plates and in liquid culture, with the growth profile of

one genotype (M3-1) approximating that of the wildtype (Figure 3A-D). A control genotype, W1-1,

isolated from day 21 of the representative wildtype line showed a growth profile comparable to that of
SBW25 (Figure 3A-D).
Three of the evolved mutant genotypes – M1-1, M3-1, and M5-1, each representing a
different class of compensatory mutation (see next section) – were chosen for one-to-one competition
experiments with (i) ΔEGEG, and (ii) a neutrally marked wildtype genotype (SBW25-*lacZ*). Each of the
evolved mutant genotypes outperformed ΔEGEG (two-tailed one-sample *t*-tests $p<0.0006$), but were
themselves outperformed by SBW25-*lacZ* (two-tailed one-sample *t*-tests $p<0.05$) (Figure 3E).
Together, these results demonstrate that the growth defect caused by *EGEG* deletion can be readily
compensated, at least in part, by evolution.

Compensation of *EGEG* loss occurs via large-scale, tandem duplications spanning *glyGCC*

In order to identify the mutation(s) behind compensation, whole genome re-sequencing was
performed on the six genotypes isolated from day 21 of the evolution experiment (M1-1, M2-1, M3-1,
M4-1, M5-1, and W1-1). Per sample, over one million, 250-bp long Illumina sequencing reads were
aligned to the SBW25 reference genome (8), resulting in a minimum mean coverage of 36.4 reads
per genomic base. *Breseq* (64) and downstream analyses were used to predict the mutations in each
evolved genotype (Supplementary Table S2).
In each of the five genotypes isolated from a mutant line, a large, direct, tandem duplication
was identified at around ~2.38 Mb of the SBW25 chromosome (Figure 4A-C). No evidence of any
such duplication fragments was found in the representative wildtype control genotype, strongly
suggesting that these large duplications play an important role in the observed compensatory effect.
In one mutant genotype (M5-1), an additional, non-synonymous SNP was identified in *rplC* (located
outside of the duplicated region; see Figure 4B). While the *rplC* mutation affects the translational
apparatus (via amino acid change T43I in the 50S ribosomal protein L3), no evidence was found for a
role in the compensated phenotype of M5-1: in the absence of the accompanying duplication, the *rplC*
mutation did not improve the growth of ΔEGEG (see Supplementary Figure S2).
The duplicated region in each ΔEGEG-derived genotype was more precisely defined using
computational analyses and duplication junction PCRs (Table 1, Supplementary Text S2). While each
duplication fragment has distinct endpoints, the fragments fall into two broad classes based on size:
~1 Mb and ~0.49 Mb duplications. ~1 Mb duplications were identified in four evolved genotypes (M1-1,
M2-1, M4-1, and M5-1). This duplication type typically occurs between long stretches of repetitive
DNA at ~1.86 Mb and ~2.86 Mb of the SBW25 chromosome, and encompasses over 900 putative
genes (Supplementary Table S3). The ~0.49 Mb duplication was identified in one genotype, M3-1,
between positions ~2.30 Mb and ~2.79 Mb of the SBW25 chromosome. At approximately half the size
of the ~1 Mb duplication, the ~0.49 Mb duplication is predicted to contain only 426 genes
(Supplementary Table S3). Notably, both types of duplication contain distinct subsets of tRNA genes:
the ~1 Mb fragments encompass seven tRNA genes (resulting in 69 tRNA genes in the evolved
genome), while the ~0.49 Mb fragment contains one tRNA gene (giving 63 tRNA genes) (Figure 4D).
Both ~1 Mb and ~0.49 Mb duplications encompass many hundreds of genes, with each
constituent gene having a particular effect – positive, negative, or neutral – on growth. Since the net

effect of each duplication fragment is compensatory, at least one gene in each fragment is presumed

to have a sizeable positive effect on ΔEGEG growth. To gain insight into the gene(s) that may

contribute to this positive effect, we focused on the ~0.49 Mb fragment, for two reasons: (i) this region

is duplicated in all five evolved genotypes, and (ii) it affords the greatest growth advantage of all

observed duplication fragments. Of the 426 putative genes in the ~0.49 Mb fragment, only one has an

obvious role in translation: *glyGCC*, the sole remaining gene encoding tRNA-Gly(GCC) in ΔEGEG

(see Figure 1). Interestingly, none of duplicated genes present a clear link to the co-deleted *gluTTC*

genes. Hence, we hypothesize that changes in *glyGCC* gene copy number are the major determinant

of the observed fitness effects.

The observed tRNA gene set changes affect tRNA-Gly(GCC) levels

We sought to test whether the above *glyGCC* gene copy number changes are reflected in mature

tRNA pools, using YAMAT-seq (a deep-sequencing technique that quantifies the relative proportions

of tRNA species in the mature tRNA pool) (6, 67). YAMAT-seq was performed on 24 samples,

including three replicates of each of eight genotypes: wildtype SBW25, the two engineered genotypes

with reductions in *glyGCC* copy number (ΔEGEG, Δ*glyGCC*), three evolved mutant genotypes that

reflect the identified compensatory mutations (M1-1, M3-1, and M5-1), and two control genotypes

(engineering control rWT, evolution control W1-1). Illumina sequencing resulted in a minimum of 1.47

million raw tRNA reads per sample; these were aligned to a set of reference sequences, consisting of

all the unique tRNA gene sequences predicted for SBW25 (6, 7) (see Supplementary Text S4). Each

of the expected 39 tRNA species was detected in every sample, in varying proportions

(Supplementary Table S4). Notably, tRNA-Gly(GCC) – encoded by three *glyGCC* genes – is the most

highly represented of all tRNA species in the wildtype, accounting for 11.0 ± 1.0 % (mean $\pm$ SE) of the

mature tRNA pool (Figure 5A). Four tRNA species (tRNA-Ile(CAU), tRNA-Phe(GAA), tRNA-Thr(CGU),

and tRNA-Glu(UUC)) were detected only at very low levels (<0.1 % of the pool). It is probable that

various post-transcriptional modifications present in these tRNA species reduced the efficacy of the

YAMAT-seq protocol (specifically, the reverse transcription reaction; see Methods) (6, 67, 75). Due to

unreliably low read numbers, these four tRNA species were removed from downstream analyses.

DESeq2 analyses were used to detect differences in the relative proportions of all 39 tRNA

species between genotype pairs of interest (Figure 5B-D, Supplementary Table S5) (6, 36, 69).

Among the measurable tRNA species, the predominant – and in many cases, only statistically

significant – effect of the engineering and evolution experiments was on tRNA-Gly(GCC) (Figure 5B-

E). Deletion of either *glyGCC* or the *EGEG* locus saw a reduction in the relative contribution of

tRNA-Gly(GCC) to the mature tRNA pool. Specifically, compared with 11.0 ± 1.0 % of the wildtype

tRNA pool, tRNA-Gly(GCC) accounts for only 6.9 ± 0.43 % and 3.6 ± 0.34 % of the Δ*glyGCC* and

ΔEGEG mature tRNA pools, respectively. The severe reduction in tRNA-Gly(GCC) in ΔEGEG was

partially reversed by adaptive evolution; genotypes M1-1, M3-1, and M5-1 show tRNA-Gly(GCC)

levels of 5.1 ± 0.34 %, 6.3 ± 0.30 %, and 5.8 ± 0.42 %, respectively. No such changes were observed

in the tRNA pools of the engineering or evolution control genotypes (rWT, W1-1) (Figure 5B-E).

Unfortunately, due to the unreliably low read numbers obtained, we are unable to comment on the

effect of *EGEG* deletion, or subsequent evolution, on tRNA-Glu(UUC).

In addition to tRNA-Gly(GCC), the ~1 Mb duplication carried by genotypes M1-1 and M5-1

changes the number of genes coding for a further four tRNA species: tRNA-Ala(UGC) (from five to six

gene copies), tRNA-Asp(GUC) (from four to six gene copies), tRNA-Ile(GAU) (from five to six gene

copies), and tRNA-Val(UAC) (from three to five gene copies). While the ~1 Mb duplication fragment

increases the contribution of each of these tRNA species to the mature tRNA pool, these changes are

less pronounced than those observed for tRNA-Gly(GCC) (and very few are statistically significant)

(Figures 5C, 5F). One reason for these weaker effects could be that each of the additional four tRNA

species is encoded by multiple, dispersed tRNA gene copies, only some of which lie within the

duplicated region. The degree to which each of these dispersed gene copies contributes to the final

level of the mature tRNA species may vary considerably, and remains unknown (76, 77). Finally, we

note that in addition to incurring no detectable growth or fitness effects (see Supplementary Figure

S2), the *rplC* mutation had no major effects on the relative proportions of tRNA species in the mature

tRNA pool, under the tested conditions; no statistically significant differences were detected between

genotypes M1-1 (~1 Mb duplication) and M5-1 (~1 Mb duplication + *rplC* mutation) (Figure 5D).

The results in this section show that changes in *glyGCC* gene copy number are mirrored by

changes in the proportional contribution of tRNA-Gly(GCC) to the mature tRNA pool. Specifically,

reductions in *glyGCC* gene copy number result in progressively lower proportions of tRNA-Gly(GCC),

which we hypothesize eventually manifest as a reduction in growth rate. Growth rate is subsequently

restored by large-scale, tandem duplication events, one effect of which is to elevate *glyGCC* gene

copy number and (partially) reinstate tRNA-Gly(GCC) levels.

Duplication fragments are unstable, leading to genetically mixed populations

Large-scale, tandem duplications are typically highly unstable; they are known to be lost at high rates

via homologous recombination events during DNA replication (often leaving no trace in the segregant

genome) (55, 78, 79). The duplication fragments observed in this work are no exception. When grown

in overnight culture and plated on agar plates, each of the duplication-carrying genotypes repeatedly

gives rise to colonies of two distinct types: (i) a majority of larger, wildtype-like colonies that retain the

duplication fragment, and (ii) a sizeable minority of smaller, ΔEGEG-like colonies in which the initial

duplication fragment has been lost (Figure 6A-B; see also Supplementary Figure S2).

The degree of duplication fragment loss was quantified using a stability test, in which the

proportion of small colonies (*i.e.*, segregants) arising from independent cultures founded by each of

the five duplication-carrying, evolved mutant genotypes was determined. Between 3 % and 34 % of

colonies arising from duplication genotype cultures showed the small, ΔEGEG-like phenotype (Figure

6C). The proportion of small colonies observed was dependent on the size of the initial duplication

fragment, with small and large duplication fragments giving rise to respective averages of 5.0 % ±

1.0% and 18 % ± 1.5 % segregants (mean ± SE; non-parametric two-sample *t*-test for difference

between means p=3.5e-07***). The higher proportion of segregants in populations founded by

duplication genotypes carrying the larger, ~1 Mb fragments presumably results from a combination of

the higher number of recombination events occurring across a more extensive region of homology,

and the lesser fitness advantage afforded by the ~1 Mb duplications (see Figure 3).

The results in this section demonstrate that the adaptive duplication events are inherently

unstable; they, along with their compensatory tRNA gene copies, are lost at high rates. In growing

populations founded by duplication-carrying genotypes, this high rate of loss leads to the presence of

at least two genetically distinct sub-populations of cells: (i) those carrying the initial duplication

fragment, and (ii) those in which the fragment has been lost (*i.e.*, ΔEGEG) (Figure 6D).

Multiple duplication fragments rapidly arise and compete within evolving populations

The ease with which duplications arise and are lost suggests complex evolutionary dynamics within

the mutant lines. To begin investigating, we tracked ~1 Mb and ~0.49 Mb duplication fragments

through one of the evolutionary lines in which they were identified by the day 21 genotype sequencing

(M2 and M3, respectively). This was done by normalizing the intensity of a duplication junction PCR

band (which is expected to change as the population evolves) with that of a product amplified from a

genomic region unaffected by the duplication events (which is expected to remain constant) (see

Methods, Supplementary Figure S3, and Supplementary Table S6). Across an evolving population,

changes in the frequency of the target duplications are expected to be reflected by changes in the

normalized intensity values (we note that the normalized intensities are not intended to provide

precise frequency estimates, but to reveal patterns of change).

The normalized intensity profile of the ~1 Mb fragment across line M2 is shown in Figure 7A,

and that of the ~0.49 Mb fragment across line M3 is in Figure 7B. Both duplications were detected

early in their respective lines (~days 4-6), and spread rapidly to peak frequency (~days 12-18).

Thereafter, the ~1Mb fragment steadily declined to low levels, while the smaller (and hence fitter and

more stable) ~0.49 Mb fragment was maintained at high levels until the end of the experiment.

The above results hint at the presence of multiple adaptive genotypes in at least some

evolving populations. To further explore this possibility, normalized duplication junction PCRs were

used to test for the co-existence of ~1 Mb and ~0.49 Mb duplication fragments in evolving lines M1,

M2, M3, as well as a representative control line (W1). (Lines M4 and M5 were excluded due to the

discovery of external contaminants in the later stages of the experiment, see Supplementary Figure

S1.) The results presented in Figure 7C show that both the ~1 Mb and ~0.49 Mb duplications are

indeed detectable in all three evolving mutant lines (and not in W1). The co-existence of both

fragment types is most prominent during the early stages of the experiment (~days 7-21); in general,

the ~1 Mb fragment declines in the later stages of the experiment, while the ~0.49 Mb fragment

remains at relatively constant levels. Notably, distinct subtypes of the smaller duplication fragment –

each with subtly different endpoints and hence variably sized duplication junction PCR products –

were detected in lines M1-M3 (see Supplementary Figures S4 and S5A for more detail). Further,

since each duplication junction PCR tests only for a subset of hypothetically possible duplication

fragments, we note that there may be other duplications present that are not captured by these PCRs.

The results in this section provide evidence of complex evolutionary dynamics of duplications

in the evolving mutant populations; both ~1 Mb and ~0.49 Mb fragments, each typically occurring

between sets of long repeats, arise readily and repeatedly within evolving ΔEGEG populations. Over

time, the larger ~1 Mb fragments are outcompeted by their smaller (and fitter/more stable) ~0.49 Mb
counterparts.

Smaller duplication fragments arise and displace larger duplication fragments

To further investigate the evolutionary dynamics of duplications, we performed population-level whole
genome re-sequencing on the day 28 populations of mutant lines M1-M3 (day 28 wildtype populations
W1-W5, and an ancestral SBW25 sample were included as controls). For each of the nine samples,
150-bp paired-end Illumina reads were aligned to the SBW25 reference sequence (8) using *breseq*
(on polymorphism detection settings, see Methods) (64), giving a minimum mean coverage of 154
reads per genomic base. The raw *breseq* output, and a list of curated mutation predictions, are
provided in Supplementary Table S7.

From the *breseq* output and further computational analyses (see Supplementary Text S3),
eleven distinct duplication fragments were identified across the three day-28 ΔEGEG-derived
populations (Table 2). No such duplications were observed in any of the control samples. Ten of
these eleven duplication fragments are novel; the eleventh is identical to the ~0.49 Mb fragment
previously identified in the genotype M3-1 (isolated from the day-21, line M3 population). Each of the
three day-28 mutant populations was found to contain at least three distinct, co-existing duplication
fragments, including multiple (relatively) large fragments (between ~0.28 Mb and ~0.77 Mb in size,
often with endpoints in repetitive DNA regions), and one much smaller fragment (between ~2.2 kb and
~4.3 kb, with no obvious endpoint homology) (Figure 8A). Notably, each mutant population is
predicted to contain a ~0.49 Mb duplication fragment between genomic positions ~2.30 Mb and ~2.79
Mb (*i.e.*, is expected to give a PCR product with the duplication junction primers used in Figure 7C).

The smallest duplication fragment is predicted in line M3. It is just 2,170-bp long and contains
only three complete genes: *pgsA* (predicted to be involved in phospholipid biosynthesis)*, glyGCC*
(tRNA gene), and *pflu2192* (a putative pseudogene of phage origin) (Figure 8B). When all 15 of the
duplication fragments identified in this study are considered, only two genes are duplicated in every
fragment: *glyGCC* and *pflu2192* (Figure 8B, Supplementary Table S3). Taken together, the population
sequencing and duplication junction PCRs provide strong support for the hypothesis that the major
compensatory element of the duplication fragments is *glyGCC*; over time, progressively smaller
duplication fragments encompassing *glyGCC* (and pseudogene *pflu2192)* arise and spread in the
evolving, ΔEGEG-derived populations.

DISCUSSION

Various studies have reported adaptive changes in tRNA gene copy number in laboratory populations
of bacteria, in response to changing translational needs (6, 40, 41). In these studies, changes in gene
copy number are underpinned by large-scale, tandem duplications (6, 41) or amplifications (40). Such
mutations are both pervasive and, often, mechanistically reversible: large-scale duplications typically
arise, and are lost, extremely rapidly (43, 44, 78, 80, 81), in a manner that is reminiscent of
contingency loci in pathogenic bacteria (82, 83). Their transient nature means that their contributions
to population diversity, and downstream effects on evolution, can be underappreciated (44). In this

work, we (i) demonstrate the contribution of large-scale duplications to population-level diversity in
tRNA gene set content, and (ii) show that such duplications may provide the raw material for longer-
term evolutionary changes in basal tRNA gene set content.

Spontaneous duplications rapidly generate population-level tRNA set diversity

As outlined in the introduction, large-scale duplications typically arise spontaneously (during DNA
replication), at high rates, in a wide range of organisms (43, 46, 54–57). For instance, it has been
reported that – without invoking any growth advantage – around 10 % of cells in an overnight
Salmonella culture carry a large duplication of some region of the chromosome, with between
0.005 % and 3 % of cells carrying a duplication of a given locus (42, 43, 55). In short, a growing
population of initially isogenic cells is expected to rapidly consist of a mixture of genotypes that vary
with respect to the copy numbers of many genes (including tRNA genes).
We have demonstrated that similar large-scale, tandem duplications readily occur in at least
two regions of the *P. fluorescens* SBW25 chromosome: either side of the putative replication terminus,
centered around genomic positions ~2.38 Mb (this work) and ~4.16 Mb (6) (see Figure 4B). In this
work we have reported 15 distinct duplications at position ~2.38 Mb, including five duplications
identified in genotypes isolated from day 21 of the evolution experiment, and a further ten duplications
predicted in the day 28 population sequencing (see Tables 1-2 and Supplementary Table S3). These
15 duplications range from ~2.2 kb to ~1 Mb in size, and each one increases the initial tRNA gene set
(of ∆EGEG) from 62 tRNA genes to either 63 or 69 tRNA genes (see Figure 4D). In our earlier work,
we observed five duplications at position ~4.16 Mb (6). These five duplications are up to ~290 kb, and
each adds one tRNA gene copy to the initial genome (∆*serCGA*) (6). The tRNA gene set changes
incurred by large-scale duplications have phenotypic consequences: the additional tRNA gene copies
generally lead to higher proportional contributions of the corresponding tRNA species to the mature
tRNA pool (see Figure 5 and (6)). Taken together, these results show that large-scale duplications
readily occur at multiple locations in the SBW25 chromosome, thereby providing a mechanism for the
rapid generation of population-level diversity in mature tRNA pool content.
The duplications reported in this work are detected at unequal rates. For instance, examples
of similar ~1 Mb and ~0.49 Mb duplication fragments are found in the early stages of each evolving
mutant line (see Table 2, Figure 7). This observation is consistent with literature showing that some
chromosomal regions – particularly those delineated by long stretches of homologous DNA (*e.g., rrn*
operons) – give rise to duplication events at especially high rates (79). Indeed, the prevalent ~1 Mb
and ~0.49 Mb fragment classes are both delineated by long stretches of homologous DNA: the ~1 Mb
duplication fragments are typically flanked by two *rhs* genes at positions ~1.86 Mb and ~2.86 Mb (~3
kb of homologous sequence), while the ~0.49 Mb fragments are generally delineated by intergenic
repeat regions at ~2.30 Mb and ~2.79 Mb (~1.1 kb of homology).
In addition to the above mechanisms, tandem duplications can arise – via alternative
mechanisms, and at much lower rates – between regions with little or no DNA homology (79). Such
alternative mechanisms presumably contribute to the generation of the smaller duplication fragments
detected at the later stages of evolution in this work; these smaller fragments, many of which have no
obvious endpoint homology, could conceivably arise at low rates either directly from the founder

genotype (ΔEGEG), or from duplication-carrying genotypes (through the partial segregation of a

duplication fragment). Indeed, an example of the latter was recently reported in an experiment

examining the fate of an advantageous, large-scale (~1.66 Mb), tandem duplication fragment in

Salmonella; during 2,000 generations of positive selection, the initial duplication fragment repeatedly

shrank to ~0.66 Mb, via the loss of an internal ~1 Mb segment (84, 85).

Unstable but adaptive duplications as evolutionary stepping stones

Generally speaking, large-scale, tandem duplication fragments are inherently unstable; they are prone

to loss through RecA-mediated recombination during DNA replication (55, 79). The duplication

fragments described in this work are similarly precarious – the stability test showed that, despite the

net fitness advantage that they provide, duplications are lost by between 3 and 34 % of cells in

overnight culture (see Figure 6). The rate of segregation was higher for genotypes carrying the larger

(~1 Mb) versus the smaller (~0.49 Mb) duplications (median loss rates of 19 % and 5 %, respectively).

This difference presumably results from a combination of the larger duplications (i) carrying twice the

length of homologous DNA for recombination (*i.e.*, the duplication fragment), and (ii) conferring a

lesser fitness advantage (see Figure 3) (44).

The inherent instability of the large-scale duplications renders them unlikely to represent a

long-term evolutionary solution. That is, over time, inherently unstable duplication fragments are

expected to be displaced by a more stable type of mutation (*e.g.*, SNPs affecting tRNA gene

expression, or smaller/more stable duplication fragments). Indeed, we already see evidence of such

displacement over the course of our 28-day evolution experiment; the large, ~1 Mb fragments

detected at the early stages of each mutant line tested (M1-M3) falls to very low or undetectable

levels by the end of the experiment (see the population-level duplication junction PCRs in Figure 7).

Furthermore, in one mutant line (M3), the co-existing smaller/fitter ~0.49 Mb fragment also appears to

drop in frequency towards the end of the experiment, hinting at the presence of at least one additional,

higher-effect mutation in this population. Indeed, subsequent population-level genome re-sequencing

detected the presence of much smaller (and, presumably, fitter and mechanistically more stable)

duplication fragments in the later stages of all three mutant lines; each day 28 mutant population

contains at least one duplication fragment of <5 kb (see Table 2, Figure 8).

The small (<5 kb) duplications fragments provide considerable insight into the mechanism

behind the compensatory effects of the duplications. Together, the three smallest fragments reduce

the core, overlapping region of all 15 reported duplication fragments to just 1,064 bp, which

encompasses only two complete genes: tRNA gene *glyGCC*, and predicted pseudogene *pflu2192*

(see Figure 8B). This observation strongly supports the hypotheses that (i) the ΔEGEG growth defect

is primarily caused by a drop in tRNA-Gly(GCC), and (ii) this defect is readily compensated by

duplicating *glyGCC* and thus (partially) restoring tRNA-Gly(GCC) levels. We note that this does not

rule out a more minor role for tRNA-Glu(UUC), the second tRNA species targeted by the initial *EGEG*

deletion. For instance, it is conceivable that a secondary, lower-effect wave of adaptation could target

gluTTC. Alternatively, selection pressure on *gluTTC* may be stronger under different experimental

conditions.

The detection of progressively shorter duplication fragments raises the interesting question of what

would occur if the evolution experiment were continued. One possibility is that the duplications would

continue to reduce in size. Presumably, the smallest possible adaptive duplication is approximately

~150-200 bp long (encompassing merely the *glyGCC* gene and promoter). Mechanistically speaking,

such a short duplication fragment would be expected to be relatively stable, and hence a genotype

carrying such a duplication could be considered to encode a new, comparatively stable tRNA gene

set. The long-term evolutionary fate of the duplication fragments – and hence the new tRNA gene

copies – remains to be seen, and will provide insight into the extent to which within-genome

duplication events contribute to the evolution of extant bacterial tRNA gene sets.

DATA AVAILABILITY

Whole genome sequencing data has been deposited with NCBI Short Read Archive (SRA) under

accession numbers PRJNA790786 (isolate sequencing) and PRJNA1035442 (population sequencing).

YAMAT-seq data has been deposited with NCBI Gene Expression Omnibus (GEO) under accession

number GSE196705.

The remaining raw data relating to this project have been deposited on Zenodo (doi:

10.5281/zenodo.7276660).

SUPPLEMENTARY DATA

Supplementary Data are available at NAR online.

ACKNOWLEDGEMENTS

The authors thank Frederic Bertels for useful comments on the manuscript.

FUNDING

This work was supported by the German Research Foundation (DFG) [GA 2895/2-1 to JG], the Max

Planck Society [all authors], and the International Max Planck Research School for Evolutionary

Biology [WYN]. Funding for open access charge: The Max Planck Society.

AUTHOR CONTRIBUTIONS

Multi-author contributions are listed alphabetically. Conceived and designed the experiments: GBA,

JG, PN, SL, WYN, ZK. Performed the experiments: all authors. Analyzed the data: all authors. Wrote

the paper: JG. Commented on the paper: all authors. Conceived the research: JG.

CONFLICT OF INTEREST

The authors declare no conflict of interest exists.

REFERENCES

1. Gingold,H. and Pilpel,Y. (2011) Determinants of translation efficiency and accuracy. *Mol*
Syst Biol, **7**, 481.

2. Rak,R., Dahan,O. and Pilpel,Y. (2018) Repertoires of tRNAs: The couplers of genomics
and proteomics. *Annu Rev Cell Dev Biol*, **34**, 239–264.

3. Abbott,J.A., Francklyn,C.S. and Robey-Bond,S.M. (2014) Transfer RNA and human
disease. *Front Genet*, **5**, 158.

4. Lant,J.T., Berg,M.D., Heinemann,I.U., Brandl,C.J. and O'Donoghue,P. (2019) Pathways to
disease from natural variations in human cytoplasmic tRNAs. *J Biol Chem*, **294**,
5294–5308.

5. Knight,J.R.P., Garland,G., Poyry,T., Mead,E., Vlahov,N., Sfakianos,A., Grosso,S., De-
Lima-Hedayioglu,F., Mallucci,G.R., Von Der Haar,T., *et al.* (2020) Control of
translation elongation in health and disease. *Dis Model Mech*, **13**, dmm043208.

6. Ayan,G.B., Park,H.J. and Gallie,J. (2020) The birth of a bacterial tRNA gene by large-
scale, tandem duplication events. *eLife*, **9**, e57947.

7. Chan,P.P. and Lowe,T.M. (2016) GtRNAdb 2.0: An expanded database of transfer RNA
genes identified in complete and draft genomes. *Nucl Acids Res*, **44**, D184-189.

8. Silby,M.W., Cerdeño-Tárraga,A.M., Vernikos,G.S., Giddens,S.R., Jackson,R.W.,
Preston,G.M., Zhang,X.X., Moon,C.D., Gehrig,S.M., Godfrey,S.A.C., *et al.* (2009)
Genomic and genetic analyses of diversity and plant interactions of *Pseudomonas*
fluorescens. *Genome Biol*, **10**, R51.

9. Ikemura,T. and Ozeki,H. (1983) Codon usage and transfer RNA contents: Organism-
specific codon-choice patterns in reference to the isoacceptor contents. *Cold Spring*
Harb Symp Quant Biol, **47**, 1087–1097.

10. Marck,C. and Grosjean,H. (2002) tRNomics: Analysis of tRNA genes from 50 genomes
of Eukarya, Archaea, and Bacteria reveals anticodon-sparing strategies and domain-
specific features. *RNA*, **8**, 1189–1232.

11. Rocha,E.P.C. (2004) Codon usage bias from tRNA's point of view: Redundancy,
specialization, and efficient decoding for translation optimization. *Genome Res*, **14**,
2279–2286.

12. Withers,M., Wernisch,L. and Dos Reis,M. (2006) Archaeology and evolution of transfer
RNA genes in the *Escherichia coli* genome. *RNA*, **12**, 933–942.

13. Wald,N. and Margalit,H. (2014) Auxiliary tRNAs: Large-scale analysis of tRNA genes
reveals patterns of tRNA repertoire dynamics. *Nucl Acids Res*, **42**, 6552–6566.

14. Tremblay-Savard,O., Benzaid,B., Lang,B.F. and El-Mabrouk,N. (2015) Evolution of tRNA
repertoires in *Bacillus* inferred with OrthoAlign. *Mol Biol Evol*, **32**, 1643–1656.

15. Diwan,G.D. and Agashe,D. (2018) Wobbling forth and drifting back: The evolutionary history and impact of bacterial tRNA modifications. *Mol Biol Evol*, **35**, 2046–2059.

16. Crick,F.H.C. (1966) Codon-anticodon pairing: The wobble hypothesis. *J Mol Biol*, **19**, 548–555.

17. Shepherd,J. and Ibba,M. (2015) Bacterial transfer RNAs. *FEMS Microbiology Reviews*, **39**, 280–300.

18. Agris,P.F., Eruysal,E.R., Narendran,A., Väre,V.Y.P., Vangaveti,S. and Ranganathan,S.V. (2018) Celebrating wobble decoding: Half a century and still much is new. *RNA Biol*, **15**, 537–553.

19. Alkatib,S., Scharff,L.B., Rogalski,M., Fleischmann,T.T., Matthes,A., Seeger,S., Schöttler,M.A., Ruf,S. and Bock,R. (2012) The contributions of wobbling and superwobbling to the reading of the genetic code. *PLoS Genet*, **8**, e1003076.

20. Varenne,S., Buc,J., Lloubes,R. and Lazdunski,C. (1984) Translation is a non-uniform process. Effect of tRNA availability on the rate of elongation of nascent polypeptide chains. **180**, 549–576.

21. Ran,W. and Higgs,P.G. (2010) The influence of anticodon-codon interactions and modified bases on codon usage bias in bacteria. *Mol Biol Evol*, **27**, 2129–2140.

22. Novoa,E.M., Pavon-Eternod,M., Pan,T. and Ribas De Pouplana,L. (2012) A role for tRNA modifications in genome structure and codon usage. *Cell*, **149**, 202–213.

23. Ikemura,T. (1981) Correlation between the abundance of *Escherichia coli* transfer RNAs and the occurrence of the respective codons in its protein genes: A proposal for a synonymous codon choice that is optimal for the *E. coli* translational system. *J Mol Biol*, **151**, 389–409.

24. Ikemura,T. (1985) Codon usage and tRNA content in unicellular and multicellular organisms. *Mol Biol Evol*, **2**, 13–34.

25. Dong,H., Nilsson,L. and Kurland,C.G. (1996) Co-variation of tRNA abundance and codon usage in *Escherichia coli* at different growth rates. *J Mol Biol*, **260**, 649–663.

26. Sharp,P.M., Bailes,E., Grocock,R.J., Peden,J.F. and Sockett,R.E. (2005) Variation in the strength of selected codon usage bias among bacteria. *Nucl Acids Res*, **33**, 1141–1153.

27. Higgs,P.G. and Ran,W. (2008) Coevolution of codon usage and tRNA genes leads to alternative stable states of biased codon usage. *Mol Biol Evol*, **25**, 2279–2291.

28. Plotkin,J.B. and Kudla,G. (2011) Synonymous but not the same: The causes and consequences of codon bias. *Nat Rev Genet*, **12**, 32–42.

29. Novoa,E.M. and Ribas de Pouplana,L. (2012) Speeding with control: Codon usage, tRNAs, and ribosomes. *Trends Genet*, **28**, 574–581.

30. Serra,M.J., Baird,J.D., Dale,T., Fey,B.L., Retatagos,K. and Westhof,E. (2002) Effects of magnesium ions on the stabilization of RNA oligomers of defined structures. *RNA*, **8**, 307–323.

31. Nikolova,E.N. and Al-Hashimi,H.M. (2010) Thermodynamics of RNA melting, one base
pair at a time. *RNA*, **16**, 1687–1691.

32. Kurland,C.G. and Ehrenberg,M. (1987) Growth-optimizing accuracy of gene expression.
Ann Rev Biophys Chem, **16**, 291–317.

33. Emilsson,V. and Kurland,C.G. (1990) Growth rate dependence of transfer RNA
abundance in *Escherichia coli*. *EMBO J*, **9**, 4359–4366.

34. Kurland,C.G., Hughes,D. and Ehrenberg,M. (1996) Limitations of translational accuracy.
In Neidhardt,F.C. (ed), *Escherichia coli and Salmonella: Cellular and Molecular*
Biology. ASM Press, Washington, D.C, Vol. 1, pp. 979–1004.

35. Sharp,P.M., Emery,L.R. and Zeng,K. (2010) Forces that influence the evolution of codon
bias. *Philos Trans R Soc Lond B Biol Sci*, **365**, 1203–1212.

36. Raval,P.K., Ngan,W.Y., Gallie,J. and Agashe,D. (2022) The layered costs and benefits of
translational redundancy. 10.1101/2022.06.16.495782.

37. Saks,M.E., Sampson,J.R. and Abelson,J. (1998) Evolution of a transfer RNA gene
through a point mutation in the anticodon. *Science*, **279**, 1665–1670.

38. McDonald,M.J., Chou,C.H., Swamy,K.B.S., Huang,H.D. and Leu,J.Y. (2015) The
evolutionary dynamics of tRNA-gene copy number and codon-use in *E. coli*. *BMC*
Evol Biol, **15**, 163.

39. Callens,M., Scornavacca,C. and Bedhomme,S. (2021) Evolutionary responses to codon
usage of horizontally transferred genes in *Pseudomonas aeruginosa*: gene retention,
amelioration and compensatory evolution. *Microb Genom*, **7**.

40. Nilsson,A.I., Zorzet,A., Kanth,A., Dahlström,S., Berg,O.G. and Andersson,D.I. (2006)
Reducing the fitness cost of antibiotic resistance by amplification of initiator tRNA
genes. *Proc Natl Acad Sci USA*, **103**, 6976–6981.

41. Bedhomme,S., Amorós-Moya,D., Valero,L.M., Bonifaci,N., Pujana,M.-À. and Bravo,I.G.
(2019) Evolutionary changes after translational challenges imposed by horizontal
gene transfer. *Genome Biol Evol*, **11**, 814–831.

42. Anderson,R.P. and Roth,J.R. (1977) Tandem genetic duplications in phage and bacteria.
Annu Rev Microbiol, **31**, 473–505.

43. Anderson,P. and Roth,J. (1981) Spontaneous tandem genetic duplications in *Salmonella*
typhimurium arise by unequal recombination between rRNA (*rrn*) cistrons. *Proc Natl*
Acad Sci USA, **78**, 3113–3117.

44. Andersson,D.I. and Hughes,D. (2009) Gene amplification and adaptive evolution in
Bacteria. *Annu Rev Genet*, **43**, 167–195.

45. Elliott,K.T., Cuff,L.E. and Neidle,E.L. (2013) Copy number change: Evolving views on
gene amplification. *Future Microbiol*, **8**, 887–899.

46. Tomanek,I., Grah,R., Lagator,M., Andersson,A.M.C., Bollback,J.P., Tkačik,G. and
Guet,C.C. (2020) Gene amplification as a form of population-level gene expression
regulation. *Nat Ecol Evol*, **4**, 612–625.

47. Andersson,D.I., Susan Slechta,E. and Roth,J.R. (1998) Evidence that gene amplification
underlies adaptive mutability of the bacterial *lac* operon. *Science*, **282**, 1133–1135.

48. Riehle,M.M., Bennett,A.F. and Long,A.D. (2001) Genetic architecture of thermal
adaptation in Escherichia coli. **98**, 525–530.

49. Lind,P.A., Tobin,C., Berg,O.G., Kurland,C.G. and Andersson,D.I. (2010) Compensatory
gene amplification restores fitness after inter-species gene replacements. *Mol*
Microbiol, **75**, 1078–1089.

50. Paulander,W., Andersson,D.I. and Maisnier-Patin,S. (2010) Amplification of the gene for
isoleucyl-tRNA synthetase facilitates adaptation to the fitness cost of mupirocin
resistance in *Salmonella enterica*. *Genetics*, **185**, 305–312.

51. Wannarat,W., Wannarat,W., Motoyama,S., Masuda,K., Kawamura,F. and Inaoka,T.
(2014) Tetracycline tolerance mediated by gene amplification in Bacillus subtilis. **160**,
2474–2480.

52. Toussaint,J.P., Farrell-Sherman,A., Feldman,T.P., Smalley,N.E., Schaefer,A.L.,
Greenberg,E.P. and Dandekar,A.A. (2017) Gene duplication in Pseudomonas
aeruginosa improves growth on adenosine. *J Bacteriol*, **199**, e00261-17.

53. Blount,Z.D., Maddamsetti,R., Grant,N.A., Ahmed,S.T., Jagdish,T., Baxter,J.A.,
Sommerfeld,B.A., Tillman,A., Moore,J., Slonczewski,J.L., *et al.* (2020) Genomic and
phenotypic evolution of *Escherichia coli* in a novel citrate-only resource environment.
eLife, **9**, e55414.

54. Straus,D.S. and Straus,L.D.A. (1976) Large overlapping tandem genetic duplications in
Salmonella typhimurium. *J Mol Biol*, **103**, 143–153.

55. Reams,A.B., Kofoid,E., Savageau,M. and Roth,J.R. (2010) Duplication frequency in a
population of *Salmonella enterica* rapidly approaches steady state with or without
recombination. *Genetics*, **184**, 1077–1094.

56. Sun,S., Ke,R., Hughes,D., Nilsson,M. and Andersson,D.I. (2012) Genome-wide
detection of spontaneous chromosomal rearrangements in Bacteria. *PLoS ONE*, **7**,
e42639.

57. Reams,A.B., Kofoid,E., Duleba,N. and Roth,J.R. (2014) Recombination and annealing
pathways compete for substrates in making *rrn* duplications in *Salmonella enterica*.
Genetics, **196**, 119–135.

58. Westra,E.R., Sünderhauf,D., Landsberger,M. and Buckling,A. (2017) Mechanisms and
consequences of diversity-generating immune strategies. *Nat Rev Immunol*, **17**, 719–
728.

59. King,E.O., Ward,M.K. and Raney,D.E. (1954) Two simple media for the demonstration of
pyocyanin and fluorescin. *J Lab Clin Med*, **44**, 301–307.

60. Sambrook,J., Fritsch,E.F. and Maniatis,T. (1989) Molecular cloning: a laboratory manual
2nd. ed. Cold Spring Harbor laboratory press, Cold Spring Harbor.

61. Zhang,X.X. and Rainey,P.B. (2007) Construction and validation of a neutrally-marked
strain of *Pseudomonas fluorescens* SBW25. *J Microbiol Methods*, **71**, 78–81.

62. Lenski,R.E., Rose,M.R., Simpson,S.C. and Tadler,S.C. (1991) Long-term experimental evolution in *Escherichia coli*. I. Adaptation and divergence during 2000 generations. *Evolution*, **138**, 1315–1341.

63. Leinonen,R., Sugawara,H. and Shumway,M. (2011) The Sequence Read Archive. *Nucl Acids Res*, **39**, D19–D21.

64. Deatherage,D.E. and Barrick,J.E. (2014) Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using *breseq*. *Methods Mol Biol*, **1151**, 165–188.

65. Deatherage,D.E., Traverse,C.C., Wolf,L.N. and Barrick,J.E. (2015) Detecting rare structural variation in evolving microbial populations from new sequence junctions using *breseq*. *Front Genet*, **5**, 468.

66. Barrick,J.E., Colburn,G., Deatherage,D.E., Traverse,C.C., Strand,M.D., Borges,J.J., Knoester,D.B., Reba,A. and Meyer,A.G. (2014) Identifying structural variation in haploid microbial genomes from short-read resequencing data using *breseq*. *BMC Genomics*, **15**, 1039.

67. Shigematsu,M., Honda,S., Loher,P., Telonis,A.G., Rigoutsos,I. and Kirino,Y. (2017) YAMAT-seq: An efficient method for high-throughput sequencing of mature transfer RNAs. *Nucl Acids Res*, **45**, e70.

68. Edgar,R., Domrachev,M. and Lash,A.E. (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucl Acids Res*, **30**, 207–210.

69. Love,M.I., Huber,W. and Anders,S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*, **15**, 550.

70. Anders,S. and Huber,W. (2010) Differential expression analysis for sequence count data. *Genome Biol*, **11**, R106.

71. Mukherjee,A., Dechow-Seligmann,G. and Gallie,J. (2022) Evolutionary flexibility in routes to mat formation by *Pseudomonas*. *Mol Microbiol*, **117**, 394–410.

72. Schneider,C.A., Rasband,W.S. and Eliceiri,K.W. (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*, **9**, 671–675.

73. Chen,S., Zhou,Y., Chen,Y. and Gu,J. (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, **34**, i884–i890.

74. Chan,P.P., Lin,B.Y., Mak,A.J. and Lowe,T.M. (2021) tRNAscan-SE 2.0: improved detection and functional classification of transfer RNA genes. *Nucl Acids Res*, **49**, 9077–9096.

75. Warren,J.M., Salinas-Giegé,T., Hummel,G., Coots,N.L., Svendsen,J.M., Brown,K.C., Drouard,L. and Sloan,D.B. (2021) Combining tRNA sequencing methods to characterize plant tRNA expression and post-transcriptional modification. *RNA Biol*, **18**, 64–78.

76. Thornlow,B.P., Armstrong,J., Holmes,A.D., Howard,J.M., Corbett-Detig,R.B. and Lowe,T.M. (2020) Predicting transfer RNA gene activity from sequence and genome context. *Genome Res*, **30**, 85–94.

77. Scheepbouwer,C., Aparicio-Puerta,E., Gomez-Martin,C., Verschueren,H., Van
Eijndhoven,M., Wedekind,L.E., Giannoukakos,S., Hijmering,N., Gasparotto,L., Van
Der Galien,H.T., *et al.* (2023) ALL-tRNAseq enables robust tRNA profiling in tissue
samples. *Genes Dev*, **37**, 243–257.

78. Adler,M., Anjum,M., Berg,O.G., Andersson,D.I. and Sandegren,L. (2014) High fitness
costs and instability of gene duplications reduce rates of evolution of new genes by
duplication-divergence mechanisms. *Mol Biol Evol*, **31**, 1526–1535.

79. Reams,A.B. and Roth,J.R. (2015) Mechanisms of gene duplication and amplification.
Cold Spring Harb Perspect Biol, **7**, a016592.

80. Anderson,R.P. and Roth,J.R. (1979) Gene duplication in Bacteria: Alteration of gene
dosage by sister-chromosome exchanges. *Cold Spring Harb Symp Quant Biol*, **43**,
1083–1087.

81. Hastings,P.J., Lupski,J.R., Rosenberg,S.M. and Ira,G. (2009) Mechanisms of change in
gene copy number. *Nat Rev Genet*, **10**, 551–564.

82. Moxon,E.R. and Thaler,D.S. (1997) The tinkerer's evolving tool-box. *Nature*, **387**, 659–
662.

83. Moxon,E.R., Rainey,P.B., Nowak,M.A. and Lenski,R.E. (1994) Adaptive evolution of
highly mutable loci in pathogenic bacteria. *Curr Biol*, **4**, 24–33.

84. Brandis,G. and Hughes,D. (2020) The SNAP hypothesis: Chromosomal rearrangements
could emerge from positive selection during niche adaptation. *PLoS Genet*, **16**,
e1008615.

85. Cao,S., Brandis,G., Huseby,D.L. and Hughes,D. (2022) Positive selection during niche
adaptation results in large-scale and irreversible rearrangement of chromosomal
gene order in Bacteria. *Mol Biol Evol*, **39**, msac069.

 TABLES AND FIGURES

Geno type	Dup. size (bp)	# Dup. genes	# Dup. tRNA genes	Junct. name	Junction side 1		Junction side 2	
					Base	**Locus**	**Base**	**Locus**
M1-1	1,016,166	917	69	M1junct	2,864,810	*pflu2597 (rhs)*	1,848,645	Intergenic (no repeat)
M2-1	1,012,338	911	69	M2junct	2,865,243–2,865,277	*pflu2597 (rhs)*	1,852,904–1,853,090	*pflu1686 (rhs)*
M3-1	489,709	426	63	M3junct	2,786,207	*pflu2547* (no repeat)	2,296,499	Intergenic (repeat)
M4-1	~1,012,000	911	69	M4junct	~2,865,500	*pflu2597 (rhs)*	~1,853,600	*pflu1686 (rhs)*
M5-1	1,012,339	911	69	M5junct	2,866,737–2,866,785	*pflu2597 (rhs)*	1,854,339–1,854,447	*pflu1686 (rhs)*

843 **Table 1. Details of duplications identified in five compensated genotypes isolated from the five**

844 **evolving mutant populations on day 21 of the evolution experiment.** Base positions refer to the

845 SBW25 wildtype genome sequence (NCBI accession number: NC_012660.1). For a list of the genes

846 contained within each duplication fragment, see Supplementary Table S3.

Pop.	Frag.	Rank[a]	Dup. size (bp)	# Genes	Junction side 1		Junction side 2	
					Base	Locus	Base	Locus
M1 day 28	F1	3	767,013	670	3,142,292 – 3,142,294	*pflu2872*	2,375,280 – 2,375,282	*pgsA* (*pflu2191*)
	F2	4	490,582	427	2,786,879 – 2,786,885	IG repeat	2,296,298 – 2,296,304	IG repeat
	F3	2	444,617	403	2,712,554	*pflu2497*	2,267,938	IG (*pflu2087/8*)
	F4	1	4,273	7	2,379,967 – 2,379,968	*pflu2197*	2,375,695 – 2,375,696	*pgsA* (*pflu2191*)
M2 day 28	F1	2	490,215	427	2,787,135 – 2,787,168	IG repeat	2,296,921 – 2,296,954	IG repeat
	F2	4	318,344	278	2,666,045 – 2,666,046	*pflu2447*	2,347,702 – 2,347,703	IG repeat (*pflu2165/7*)
	F3	1	285,798	256	2,525,738	*soxB* (*pflu2319*)	2,239,941	IG *rrn* (23S/5S)
	F4	3	3,631	6	2,379,196	*pflu2196*	2,375,566	*pgsA* (*pflu2191*)
M3 day 28	F1*	1	489,709	426	2,786,206 – 2,786,207	*pflu2547*	2,296,498 – 2,296,499	IG repeat (*pflu2166/8*)
	F2	3	434,971	360	2,809,371	*pflu2553*	2,374,401	*uvrC* (*pflu2190*)
	F3	2	2,170	3	2,376,758	*ribD* (*pflu2193*)	2,374,589	*uvrC* (*pflu2190*)

Table 2. Details of the duplications computationally predicted in day 28 populations from

evolving mutant lines M1, M2, and M3. Base positions refer to the SBW25 wildtype genome

sequence (NCBI accession number: NC_012660.1). For a list of the genes contained within each

predicted duplication, see Supplementary Table S3. [a]Duplications are ranked within each population

in descending order of prevalence (according to the adjusted number of Illumina reads perfectly

matching the complete proposed duplication junction sequence; see Supplementary Text S3).

*Fragment M3d28-F1 is identical to the duplication identified in day 21 genotype M3-1 (see Table 1).

Figure 1. Translation of glycine (G) and glutamic acid (E) in *P. fluorescens* SBW25. (**A**) Cartoon

of the SBW25 chromosome depicting loci predicted by GtRNAdb2.0 to encode canonical tRNAs. Grey

lines=tRNA loci (each representing between one and four tRNA genes); black arrow=*gluTTC-glyGCC-*

gluTTC-glyGCC ("*EGEG*") tRNA locus deleted by genetic engineering; white arrow=*glyGCC* lone

tRNA locus (encoding the remaining copy of *glyGCC*); grey arrow=*gluTTC-alaGGC-gluTTC-alaGGC*

tRNA locus (encoding the remaining two copies of *gluTTC*). (**B**) Details of the *EGEG* tRNA locus

indicated by the black arrow in panel A. (**C**) Putative translational relationships between anticodons

and codons for glycine (top) and glutamic acid (bottom). Codons are listed with their cognate

anticodons (black=tRNA species encoded in genome; grey=not encoded). Predicted anticodon-codon

matching patterns are indicated by joining lines (black=cognate match; grey=wobble match).

Percentage contribution of each codon to (i) the relevant amino acid (% aa), and (ii) all amino acids, is

provided to 1 d.p. from GtRNAdb2.0 (tRNAscan-SE version 2.0.2; February 2019) (7). (**D**) Details of

the tRNA gene sets carried by engineered genotypes.

Figure 2. Deletion of the four gene tRNA locus *EGEG* reduces SBW25 growth in rich, undefined (KB) and poorer, defined (M9+glucose) media. (**A**-**B**) Colonies of SBW25, reconstructed wildtype (the engineering control, rWT), and the two engineered genotypes Δ*glyGCC* and ΔEGEG at room temperature for 30 hours on KB agar, or 72 hours on M9 agar. Scale bar applies to all four images in each panel. (**C**-**D**) Growth (measured as absorbance at 600 nm) of the same genotypes in liquid KB for 8 hours, or M9 for 16 hours. Lines show the mean of 7 replicates at 5-minute intervals; error bars are ± 1 SE. (**E**-**H**) Maximum growth rates (change in absorbance; mOD min^{-1}) and lag times of growth profiles in panel C (KB), and panel D (M9). In all four panels, red and blue lines are the medians of SBW25 and ΔEGEG, respectively. Non-parametric Dunn's tests, followed by the Benjamini-Hochberg procedure to correct for multiple comparisons, were used to test for a difference in medians between pairs of genotypes in each panel. Stars indicate statistically significant *p*-values for a difference from SBW25 (red stars) or ΔEGEG (blue stars).

Figure 3. The growth defect caused by deleting four-gene tRNA locus *gluTTC-glyGCC-gluTTC-

glyGCC (*EGEG*) undergoes repeated compensation during a serial transfer experiment.** (**A**)

Colonies of founders SBW25, ΔEGEG, and six genotypes from day 21 of the serial transfer evolution

experiment, grown on KB agar for 30 h at room temperature. Scale bar applies to all images. (**B**)

Growth in liquid KB culture. Lines show the mean of 7 replicates at 5-minute intervals; error bars are

± 1 SE. (**C,D**) Maximum growth rate (change in absorbance; mOD min^{-1}) and lag time for each growth

profile in panel B. In both panels, red and blue lines are the medians of SBW25 and ΔEGEG,

respectively. Non-parametric Dunn's tests, followed by the Benjamini-Hochberg procedure to correct

for multiple comparisons, were used to test for a difference in medians between pairs of genotypes.

Stars indicate statistically significant *p*-values for a difference from SBW25 (red stars) or ΔEGEG (blue

stars). The data for SBW25, rWT and ΔEGEG also appears in Figure 2. (**E**) Box plots of the relative

fitness of competitor 1 (*x*-axis) and competitor 2 (horizontal bars at top). Direct, one-to-one

competitions were performed in liquid KB for 24 hours (28˚C, shaking). Between 5 and 8 replicate

competitions were performed for each genotype pair. Relative fitness >1 means that the first

competitor wins, <1 means that the second competitor wins. Statistically significant deviations of

relative fitness from 1 were determined using either one-tailed one-sample *t*-tests, followed by a

Benjamini-Hochberg correction for multiple comparisons. Data points from all replicates are overlaid

on boxplots, and ***p<0.001, **p<0.01, *p<0.05.

Figure 4. Large-scale, tandem duplications compensate for deletion of *EGEG*. (A) Number of raw sequencing reads covering 1.8 - 3.0 Mb of the reference genome. A jump to two-fold coverage is evident in the genotypes isolated from mutant populations (M1-1 to M5-1), and not in the control genotype isolated from the representative wildtype population (W1-1). The ~0.49 Mb region of double coverage shared by all mutants is highlighted in green; tRNA genes are shown by dotted lines. **(B)** Cartoon depicting the positions of duplicated regions on the SBW25 genome. Black arc=~1 Mb duplication (in M1-1, M2-1, M4-1, M5-1); green arc=~0.49 Mb duplication (in M3-1); grey arc=distinct ~0.4 Mb region in which duplication fragments have previously been reported (6). Duplicated tRNA genes are marked by dotted lines; deleted tRNA genes are marked with a cross. Image drawn using BioRender.com. **(C)** Depiction of the duplication event in M3-1, resulting in an additional ~0.49 Mb of DNA and double copies of all genes in the duplicated region. Thick black line=duplication junction, which can be amplified using a duplication junction PCR (primer positions in black; see Supplementary Text S2). **(D)** Heatmap showing differences in the gene copy numbers coding for 39 tRNA species between wildtype (wt), engineered (eng.), and evolved (evol.) genotypes. Open/closed circles=tRNA species whose gene copy number changed by engineering/evolution.

Figure 5. *glyGCC* gene copy number changes are mirrored in the mature tRNA pool. (A) Proportions of the 39 tRNA species in the SBW25 mature tRNA pool (highest to lowest). Light grey=excluded tRNA species. Open/closed circles=tRNA species whose gene copy number (GCN) changes by engineering/evolution in downstream experiments. (**B-D**) Heatmaps of differences in tRNA species proportions (expressed as log2-fold change(strain1/strain2); see Supplementary Table S5) between genotype pairs, to detect the effect of engineering (panel B) and evolution (panels C-D). Box outlines indicate statistical significance (DESeq2 adjusted p-values; thin line *p*>0.01; thick grey line 0.01<*p*>0.001; thick black line *p*<0.001). (**E**) Bar graph of tRNA-Gly(GCC) proportion by genotype. Bars=means of 3 replicates ± 1 SE. DESeq2 adjusted *p*-values show tRNA-Gly(GCC) differences between genotypes pairs. (**F**) Scatter plot of mean proportions ± 1 SE of 7 tRNA species, by genotype. GCN for first 5 tRNA species varies by engineering and/or evolution (indicated by point size), while the final 2 are controls (constant GCN). *rrn*=tRNA species genes co-localize with rRNA operons.

Figure 6. The large-scale, tandem duplications are rapidly lost in overnight culture. (**A**) Two colony morphologies – large and small – are observed when plating from overnight cultures of duplication-carrying genotypes (colonies from duplication genotypes M3-1 and M5-1 are presented here). All images taken under the same magnification after ~48 hours' incubation on KB agar at room temperature (~21˚C). Scale bar applies to all images in panel. (**B**) Small colonies no longer amplify the unique junction introduced by the relevant duplication fragment (illustrated by the thick black lines in M3-1 cartoon of panel D). (**C**) Duplication fragment loss was quantified as the proportion of small colonies when plating from an overnight culture. Five replicates were included per genotype, and at least 41 colonies were counted per replicate. Boxplots summarize the data (medians), and grey dots are individual replicates. Statistical significance of a difference in the median proportion of small colonies compared to SBW25 (wildtype) was calculated using a Dunn test (with Benjamini-Hochberg correction for multiple comparisons). Significance levels: no stars (p>0.05), *0.05<p<0.001, **0.01<p<0.001, ***p<0.001. (**D**) Cartoon depicting the gain and loss of duplication fragments, leading to a mixed population of the duplication genotype (here, M3-1) and ΔEGEG.

Figure 7. The complex evolutionary dynamics of large-scale duplication fragments in evolving

mutant populations. Duplication junction PCRs were used to follow the dynamics of subsets of ~1

Mb and ~0.49 Mb duplications across evolving populations. The intensity of the PCR product obtained

across each emergent duplication junction was measured relative to a PCR product amplified from a

control region located outside of the genomic region affected by duplication events. Various control

samples – in which (almost) all or none of the population is expected to carry duplication fragments –

were included in each set of PCRs (see Methods for further details). The subsets of ~1 Mb (**A**; primer

pair M1and2_junct_f/M5_junct_r) and ~0.49 Mb (**B**; primer pair M3_junct_f/r) duplications show

different evolutionary trajectories in the lines in which they were originally identified; the ~1 Mb

fragment rises and then wanes in line M2, while the ~0.49 Mb fragment rises to, and is maintained for

longer at, a higher level in line M3. (**C**) Multiple duplication fragments arise and compete within

evolving populations. Each PCR was performed in triplicate, and the median is presented here (for full

data sets see Supplementary Figure S5A; Supplementary Table S6).

Figure 8. Eleven distinct duplication fragments are detected in mutant lines M1, M2, and M3 on

day 28 of the evolution experiment. (**A**) Number of raw sequencing reads covering 1.7 Mb – 3.2 Mb

of the reference genome, shown for the three mutant lines (M1, M2, M3) and one representative

wildtype control line (W1) on day 28. Large changes in coverage can be seen in all mutant

populations (but not the wildtype population). The black bars below each plot indicate approximate

positions of each computationally predicted duplication (see Table 2); grey bars show the duplication

identified in the corresponding day 21 isolate (see Table 1). (**B**) Structure of the ~7 kb genomic region

containing the three smallest duplication fragments. Grey highlighting indicates the 1,064 bp that is

duplicated in all three fragments. This region contains two complete genes: tRNA gene *glyGCC* and

predicted pseudogene *pflu2192* (see Supplementary Table S3).